Hominoid-specific transposable elements rewired the expression of neural crest

migration genes

Laura Deelen[1], Zoe H. Mitchell[1], Martina Demurtas[1], Andria Koulle[1], Beatriz Garcia Del Valle[1],

and Marco Trizzino[1,*]

1 Department of Life Sciences, Imperial College London, SW7 2AZ London, UK

*Corresponding author: E-mail: m.trizzino@imperial.ac.uk

Abstract

Craniofacial development is evolutionarily conserved, yet subtle changes in its regulatory
network drive species-specific traits. Transposable elements (TEs) contribute to genome
evolution, but their role in cranial neural crest cells (CNCCs) remains unclear. Here, we
investigate the domestication of hominoid-specific TEs (LTR5Hs and SVAs) as enhancers
during human CNCC specification, a process critical for vertebrate craniofacial development.
Using human iPSC-derived CNCCs, we identified ~550 hominoid-specific TEs functioning as
enhancers, including ~250 human-specific, predominantly LTR5Hs. These elements are
enriched for CNCC coordinator motifs, are bound by the CNCC signature factor TWIST1, and
their enhancer activity appears largely CNCC-specific. CRISPR-interference targeting ~75%
of these active TEs led to widespread transcriptional dysregulation of genes involved in neural
crest migration, and two orthogonal functional assays confirmed that CNCC migration is
impaired upon TE repression. Finally, genes near human-specific TEs showed higher
expression in human CNCCs compared to chimpanzee CNCCs, but TE repression restored
gene expression to chimpanzee levels. These findings highlight how young TEs were
domesticated to fine-tune CNCC regulatory networks, potentially contributing to lineage-
specific craniofacial evolution.
26
27

28 **Keywords**: Cell migration / coordinator motif / Cranial Neural Crest / LTR5Hs / SVAs

Introduction

Craniofacial development is a highly complex process that requires precise spatiotemporal regulation and involves contributions from all three germ layers, and in particular from neural crest cells (NCCs) (Gans & Northcutt, 1983; Ahlstrom & Erickson, 2009; Martik & Bronner, 2021; Theveneau & Mayor, 2012). NCCs emerge during early embryogenesis, between weeks 3-4 in humans, at the neural plate border between the neuroectoderm and non-neural ectoderm. As neurulation progresses, the neural plate invaginates and separates from the dorsal ectoderm, forming the neural tube. At this stage, NCCs undergo epithelial-to-mesenchymal transition (EMT), allowing them to delaminate and migrate to specific regions throughout the developing embryo (Ahlstrom & Erickson, 2009; Martik & Bronner, 2021; Theveneau & Mayor, 2012). Among the NCC subtypes, cranial neural crest cells (CNCCs) play a pivotal role in the formation of key craniofacial structures, including bones and cartilage (Bronner & LeDouarin, 2012; Cordero et al., 2011; Jheon & Schneider, 2009).

Although craniofacial development is an evolutionary conserved process, recent adaptations to the modern human craniofacial complex include changes in shape and function to accommodate the enlargement of the brain, the transition to bipedal posture, laryngeal extension for speech as well as adjustments for the evolvement of sensory organs (Lieberman, 1998; Sambataro et al., 2022; Spoor et al., 1994). The evolution of human-specific craniofacial traits has required precise modifications in gene expression and increasing evidence suggests that regulatory changes, rather than protein-coding mutations, have been key to shaping species-specific features (Carroll, 2005; King & Wilson, 1975; Wray, 2007). One major source of these regulatory innovations are the transposable elements (TEs). Comprising nearly half of the human genome, TEs are now recognized as key contributors to genomic evolution through their ability to integrate into the genome and act as cis-regulatory elements (Bourque et al., 2008; Chuong et al., 2013, 2016; Cosby et al., 2021; Goubert et al., 2020; Kunarso et al., 2010; Lynch et al., 2011; Pontis et al., 2019; Schmidt et al., 2012; Sundaram & Wysocka, 2020). Transposable element-mediated rewiring of gene regulatory networks has previously been implicated as a major driver of species-specific gene expression patterns (Chuong et al., 2013; Feschotte, 2008; Fueyo et al., 2022; Jacques et al., 2013; Patoori et al., 2022; Playfoot et al., 2021; Prescott et al., 2015; Sundaram et al., 2014; Trizzino et al., 2017).

However, much remains to be uncovered about the precise mechanisms through which transposable elements (TEs) have been co-opted as cis-regulatory elements in humans, particularly in the context of craniofacial development. In the human genome, SINE-VNTR-Alus (SVAs) and LTR5Hs have been previously linked to gene regulatory activity (Barnada et al., 2022; Chuong et al., 2016; Fuentes et al., 2018; Patoori et al., 2022; Pontis et al., 2019; Trizzino et al., 2017). SVAs are the youngest TE family and include over 3000 copies in the human genome. They are composed of a hexamer repeat, an Alu-like element, a variable

number of tandem repeats (VNTRs), a SINE element, and a poly-A tail (Fig. 1A). The SVAs

are hominoid-specific, comprise of six subfamilies (A–F), with SVA-E and SVA-F being found

exclusively in humans, with ~1700 copies in total (Quinn & Bubb, 2014; Wang et al., 2005).

LTR5Hs are also human-specific and are remnants of an ancestral endogenous retrovirus of

the HERV-K subtype.

Co-option of SVAs and LTR5Hs as active cis-regulatory elements has been observed in

different human tissues (Barnada et al., 2022; Chuong et al., 2013; Fuentes et al., 2018;

Ostertag et al., 2003; Patoori et al., 2022; Pontis et al., 2019; Trizzino et al., 2017). Since the

expansion of LTR5Hs and SVA-E/F subfamilies occurred around the time of the human–

chimpanzee split, we set out to investigate if these elements could have contributed to human-

specific craniofacial development.

Human cranial neural crest cell (CNCC) specification and migration can be effectively

modelled *in vitro* using human-induced pluripotent stem cells (hiPSCs). Therefore, in this

study, we employed an inducible CRISPR-interference (CRISPRi) hiPSC line to investigate

the impact of silencing human SVA and LTR5Hs elements to CNCC specification and

migration, which are essential processes for the development of the craniofacial structures,

including bones and cartilage. To achieve this, we employed previously published single-guide

RNAs (sgRNAs) targeting approximately 80% of these transposable elements (Pontis et al.,

2019).

With this approach, we identified approximately ~550 human SVAs and LTR5Hs that are

accessible and depleted of the repressive histone mark H3K9me3 in human CNCCs. We

found that the specific DNA sequence was the primary driver for the co-option of this set of

TEs. Importantly, silencing these retroelements attenuated the expression of hundreds of

genes involved in CNCC migration, which is a key process in craniofacial morphogenesis.

Functional assays confirmed that migration was disrupted, suggesting a potential role for the

transposons in species-specific craniofacial development. This was further supported by

comparisons with previously published chimpanzee CNCC data, where the corresponding

genes to those near accessible human-specific transposable elements showed expression

patterns similar to those in human CNCCs when these elements were silenced.

Results

An inducible CRISPR-interference iPSC system for the repression of SVAs and LTR5Hs

To investigate the role of hominoid-specific TEs (SVAs and LTR5Hs; Fig. 1a) in human cranial

neural crest cell (CNCC) development, we designed a stable human iPSC line with an

inducible CRISPR-interference (CRISPRi) system with single-guide RNAs (sgRNAs) targeting

~80% of all SVAs and LTR5Hs annotated in the human genome (Fig. 1b). The sgRNAs were

originally designed and validated in a recent seminal study (Pontis et al., 2019) and were
further used in subsequent studies (Barnada et al., 2022; Patoori et al., 2022). Briefly, we
cloned a stable iPSC line with a permanently integrated TET-inducible, catalytically dead,
Cas9 fused to a repressive KRAB domain (dCas9-KRAB), along with the two gRNAs targeting
LTR5Hs and SVAs (hereafter +gRNA line). The KRAB domain recruits the transcriptional
machinery necessary to deposit repressive histone methylation (H3K9me3) to the regions
targeted by dCas9. To account for potential off-target effects caused by exposure to
doxycycline (TET-ON) or by Cas9 expression, we cloned the same iPSC line with an identical
dCas9-KRAB construct but without any gRNAs (hereafter -gRNAs line). Treating the cells with
doxycycline for 24 hours was sufficient to activate dCas9 in both CRISPRi lines (i.e. with and
without gRNAs; Fig. 1c,d).

Next, we generated CNCCs from our CRISPRi-iPSC lines using an established 5-day protocol
(Fig. 1e,f; Leung et al., 2016). Since both SVAs and LTR5Hs have been shown to have
important roles in human embryonic stem cells and iPSCs (Barnada et al., 2022; Fuentes et
al., 2018; Pontis et al., 2019), doxycycline was only introduced 24h after differentiation (Fig.
1c). By day 5 of differentiation, both cell lines expressed markers typical of CNCC identity,
both at gene (*SOX9*, *SOX10*, *TWIST1*, *TFAP2A*) and protein (SOX9, AP2α) level. We
generally did not observe significant differences in CNCC marker expression between the two
lines, apart from *TFAP2A* (Fig. 1e,f), indicating that expression of the main genes essential for
CNCC identity was largely unaffected by the CRISPRi with the exception of *TFAP2A*.

Finally, using RNA-seq data, we explored the TE expression landscape in human CNCCs.
This revealed that many TE subfamilies are highly expressed (including SVAs and HERVK-
LTRs, with Alu (SINE) and L1 (LINE) elements dominating (Appendix Fig. S1a, Appendix File.
S1).

Hundreds of SVAs and LTR5Hs are accessible in human CNCCs

We first set out to determine whether any SVA and LTR5Hs exhibited chromatin accessibility
in human CNCCs, and whether these elements could be repressed using our CRISPRi
system. To this end, we performed ATAC-seq (paired-end, 150 bp reads) in hiPSC-derived
CNCCs generated with our CRISPRi-iPSC lines (- and +gRNAs). We started by examining
human-specific SVAs (i.e. SVA-E, SVA-F) and LTF5Hs. K-mer clustering of the ATAC-seq data
identified a total of 256 accessible LTR5Hs and human-specific SVAs (Fig. 2a; Appendix File
S2). Notably, 77% of these 256 TEs displayed decreased accessibility in the +gRNA CRISPRi
line relative to the -gRNA control (cluster 2, Fig. 2a). This is consistent with the assumption
that these gRNAs can target ~80% of all the human LTR5Hs and SVAs (Pontis et al. 2019).
LTR5Hs were significantly overrepresented among the 256 accessible human-specific TEs

(observed 87%, expected 29%; Fisher's Exact Test p < 2.2 x 10^{-16}; Fig. 2b), while the SVAs

were significantly underrepresented (13% observed vs 71% expected; Fisher's Exact Test p <

2.2 x 10^{-16}; Fig. 2b), highlighting a potential primary role for LTR5Hs in CNCCs.

We set out to confirm that the loss in chromatin accessibility observed in the +gRNA lines was

a direct consequence of the CRISPRi-mediated repression. To this end, we performed ChIP-

seq for H3K9me3 (paired-end 150 bp reads), which showed accumulation of this repressive

histone mark at approximately 75% of the 256 accessible LTR5Hs and human-specific SVAs

in the +gRNA samples (Fig. 2c). Notably, there was a remarkable overlap (92%) between the

TEs that lost chromatin accessibility (Fig. 2a) and those gaining H3K9me3 (Fig. 2c).

We further investigated the genomic locations of these accessible human-specific TEs.

Overall, 91% of the 256 accessible LTR5Hs and human-specific SVAs were located > 1 kb

from the nearest transcription start site (TSS), and the median distance was 8.9 kb (Appendix

Fig. S1b). This suggests that these accessible mobile elements could be putative CNCC

enhancers. To confirm this, we leveraged publicly available H3K27ac ChIP-seq data

previously generated by our group in hiPSC-derived CNCCs (Barnada et al., 2024; Data ref:

Barnada et al., 2024). This analysis revealed that the vast majority of the 256 accessible

human-specific TEs also displayed the H3K27ac modification, an established active enhancer

mark, supporting their role as bona-fide human-specific CNCC enhancers (Appendix Fig.

S1c).

To investigate whether these accessible human-specific TEs are actively transcribed in

CNCCs, we performed a H3K4me3 ChIP-seq (which is a marker for actively transcribing

promoters) in -gRNA CNCCs and +gRNA CNCCs. This experiment revealed that 39.5% of the

accessible LTR5Hs and human-specific SVAs were also decorated by H3K4me3, suggesting

that over a third of the accessible LTR5Hs and human-specific SVAs are likely engaged in

active transcription in CNCCs (Appendix Fig. S1d,e).

Next, we focused on the accessibility of all SVA subfamilies, including those that are also found

in other hominoid species (SVA_A, _B, _C, _D). To this end, we analysed the ATAC-seq signal

across all existing SVA subfamilies and LTR5Hs separately (Appendix Fig. S2a,b). This yielded

results consistent to the previous analysis, since we found that 5% (n = 258) of non-human-

specific SVAs (A–D subfamilies) were accessible in CNCCs, whilst 95% (n = 5716) were

inaccessible (Appendix Fig. S2a,b). Therefore, even when accounting for all SVA groups,

SVAs remained significantly underrepresented among accessible TEs.

In summary, these experiments identified approximately 550 hominoid-specific TEs, over half

of which human-specific, that exhibit signatures of active enhancers in human iPSC-derived

CNCCs, with LTR5Hs elements playing a prominent role. Notably, our CRISPR-based

approach enabled the inducible repression of ~75% of these putative TE-derived human-

specific CNCC enhancers.

To ensure that our findings were not biased by a specific differentiation protocol, we used an
alternative iPSC-to-CNCC differentiation method (Bajpai et al., 2010) and performed ATAC-
seq on the differentiated cells. With this protocol, migratory CNCCs are obtained in 2-3 weeks
(Bajpai et al., 2010; Barnada et al., 2024; Mitchell et al., 2025; Pagliaroli et al., 2021). Using
this approach, we identified 374 LTR5Hs and human-specific SVA elements that are
accessible in CNCCs, the majority of which exhibited reduced accessibility in the +gRNA line
compared to the -gRNA control (Appendix Fig. S2c). Moreover, comparable to our findings
obtained with the 5-day protocol, LTR5Hs were significantly overrepresented in the set of
accessible TEs (83% observed vs 29% expected; Fisher's Exact Test $p < 2.2 \times 10^{-16}$), while
SVAs were significantly underrepresented (17% observed vs 71% expected; Fisher's Exact
Test $p < 2.2 \times 10^{-16}$). Importantly, 91% of the 256 human-specific TEs identified as accessible
in the 5-day protocol were also found accessible in CNCCs derived using the alternative
protocol. This consistency suggests that our findings are robust and independent of the CNCC
differentiation method used.
189
Co-opted TEs are enriched with the CNCC *coordinator* motif

Next, we explored the genomic features driving the recruitment of the LTR5Hs and SVAs as
CNCC enhancers. First, we performed computational DNA motif analysis on the set of
accessible TEs using the non-accessible LTR5Hs and SVAs as background control for
differential enrichment. Recent studies have identified a specific DNA motif, known as
coordinator, which is enriched at enhancers critical for the regulation of CNCC identity (Kim et
al., 2024; Prescott et al., 2015). *Coordinator* is a composite motif which consists of a fusion
between a generic AT-rich homeobox motif (TTAATTA) and the binding motif of the CNCC
master regulator TWIST1, typically joined by a stretch of A nucleotides (Kim et al., 2024;
Prescott et al., 2015). Importantly, we found both components of the *coordinator motif* as highly
enriched in the set of accessible LTR5Hs and human-specific SVAs (Fig. 2d; Appendix File
S3). Specifically, the TWIST1 motif and the AT-rich homeobox motif were found in 83% and
62% of accessible human-specific TEs, respectively, with similar numbers observed when
incorporating non-human-specific SVAs in the analysis (Fig. 2d; Appendix Fig. 2d,e; Appendix
File S3). Moreover, we found that 29% of the accessible SVAs and LTR5Hs harboured the full
coordinator motif sequence, as opposed to only 14% of the non-accessible TEs. This suggests
that the human-specific TEs accessible in CNCCs are significantly more enriched for the
coordinator motif relative to the non-accessible ones (Fisher's Exact Test $p < 0.00001$).
In addition, the motifs for other CNCC signature transcription factors, such as AP2α (*TFAP2A*),
SOX9/10, SLUG and FOXD3 were also found as significantly enriched (Fig. 2d; Appendix Fig.
2d,e; Appendix File S3). Given the prevalence of the *coordinator* motif in accessible LTR5Hs

and SVAs, we investigated whether these TEs are directly bound by TWIST1 in human
CNCCs. To address this, we leveraged publicly available TWIST1 ChIP-seq data generated
in iPSC-derived CNCCs (Kim et al., 2024), which revealed TWIST1 ChIP-seq signal at over
half of the accessible LTR5Hs and SVA in human CNCCs (Fig. 2e).

Overall, these findings suggest that DNA sequence and transcription factor binding are a major
driver for TE co-option as active CNCC enhancers.

Finally, we examined whether the LTR5Hs and SVAs accessible in CNCCs integrated into
previously existing enhancers or transcriptionally inactive genomic regions. To investigate this,
we analyzed regions extending 250 bp upstream and downstream of these accessible TEs,
assessing their accessibility in chimpanzee iPSC-derived CNCCs using publicly available
paired-end ATAC-seq data (Prescott et al., 2015; Data ref: Prescott et al., 2015). Notably, we
observed no accessibility signal in any of these regions (Appendix Fig. S3), suggesting that
the hominoid-specific transposons likely established novel cis-regulatory elements rather than
integrating into pre-existing active enhancer regions.

Repression of LTR5Hs and SVAs impairs expression of CNCC-migration genes
Our experiments so far identified ~550 LTR5Hs and SVAs (half of which human-specific)
displaying active enhancer signature in human CNCCs. Since our CRISPRi system enables
the simultaneous repression of ~75% of these elements, we next investigated whether their
repression had significant effects on the CNCC transcriptome. To this end, we differentiated
our doxycycline-treated CRISPRi lines (+ and -gRNAs) into CNCCs and performed RNA-seq.
First, we analyzed gene expression in relation to the nearest transcription start sites (TSS) of
the 256 accessible human-specific SVAs and LTR5Hs. In total, we identified 107 genes
located near these elements that were actively expressed in CNCCs (median TPM >1 across
all -gRNA replicates). Overall, the expression of these genes was significantly decreased upon
TE repression (Wilcoxon's Rank Sum test p < 0.037; Appendix Fig. S4a), with 83/107 (=
77.5%) displaying lower expression in the +gRNA sample relative to the -gRNA counterpart
(Appendix File S4). Consistent with LTR5Hs playing a prominent role in CNCC, 71 of the 83
genes (= 86%) with decreased expression upon CRISPR-i were the nearest gene to an
accessible LTRH5s.

We leveraged publicly available RNA-seq data from CNCCs derived from chimpanzee iPSCs
(Prescott et al., 2015), which we reanalysed using our pipeline, to compare the expression of
these 107 genes between the two closely related ape species. This analysis revealed that
these genes are typically expressed at significantly higher levels in humans than in
chimpanzees (Fig. 3a; Appendix File S4). However, repressing the SVAs and LTR5Hs
abolished most of this expression difference between the two species (Fig. 3a), suggesting a
direct role for human-specific TEs in species-specific CNCC gene regulation. Consistent with

this, repressing the hominoid-specific TEs increased the correlation between human and
chimpanzee expression levels for these genes (Appendix Fig. S4b).
Next, we examined transcriptome-wide effects. Differential gene expression analysis identified
795 genes that were significantly differentially expressed between +gRNA and -gRNA CNCCs
(FDR < 0.05; FC < -1.5 or > 1.5; Fig. 3b; Appendix File S5). Of these, 501 genes were
downregulated, while 294 genes were upregulated in the +gRNA samples (Fig. 3b). Gene
ontology enrichment analysis of the downregulated genes revealed a significant enrichment
for cell migration-related processes, suggesting that many of these genes play key roles in
CNCC migration (Fig. 3c; Appendix File S6). In line with these findings, gene ontology
enrichment analysis performed on the 107 genes expressed genes located near accessible
LTR5Hs and human-specific SVAs also identified cell migration as one of the top-5 enriched
pathways, despite the small sample size (Appendix File S7). Conversely, upregulated genes
were primarily associated with mitochondrial and cell division processes (Appendix File S8).
It is worth noting that only 3 of the differentially expressed genes had an intronic SVA or
LTR5Hs, suggesting that the high number of differentially expressed genes is not an artifact
of CRISPRi mediated repression at gene bodies.
264
Silencing hominoid-specific TEs functionally affects CNCC migratory potential *in vitro*
Since cell migration was the predominant signature enriched in the downregulated genes, we
investigated whether repressing hominoid-specific SVAs and LTR5Hs could functionally affect
CNCC migration. To test this, we performed a transwell migration assay, in which hiPSC-
CNCCs were seeded on geltrex-coated transwell membranes overnight and subsequently
exposed to medium supplemented with the general chemoattractant fetal bovine serum (FBS)
in the lower chamber (Fig. 4a). After 24 hours of incubation, comparison of the –gRNA and
+gRNA lines revealed a significant reduction in CNCC migration across the membrane upon
repression of hominoid-specific TEs, suggesting impaired migratory capacity (Fig. 4b,c).
Additionally, we performed a wound healing assay which showed significantly impaired wound
closure of the +gRNA CNCCs compared to -gRNA CNCCs when exposed to medium+10%
FBS for 8h (Fig. 4d,e). Altogether, these findings are consistent with the RNA-seq data and
suggest that SVAs and LTR5Hs-driven regulatory activity contributes to regulate human CNCC
migration.
279
The LTR5Hs and SVAs active in CNCCs are silent in most human cell-types
Finally, we investigated whether the 256 human-specific SVAs and LTR5Hs with active
enhancer signature in CNCCs were also showing cis-regulatory signature in other human cell
types. To address this, we leveraged publicly available H3K27ac ChIP-seq data from 14 cell
types generated by the Roadmap Epigenomics Consortium, including distinct brain regions,

lung, aorta, adipose tissue, pancreas and spleen (Kundaje et al., 2015; Data ref: Kundaje et
al., 2015). This analysis revealed that, in addition to CNCCs, these 256 TE exhibit active
enhancer signature exclusively in endomesodermal cells, with a subset also active in iPSCs
(Fig. 5). In contrast, they remain completely silenced in all other cell types (Fig. 5). The
activation of these elements in endomesodermal cells is unsurprising, as a previous study
from our lab has shown that human-specific SVAs are highly enriched for the motif of the key
mesodermal regulator EOMES (= TBR2; Patoori et al., 2022). Similarly, the co-option of SVAs
and LTR5Hs as regulatory elements in human iPSCs and ESCs has been suggested by
previous research (Barnada et al., 2022; Fuentes et al., 2018; Pontis et al., 2019).

In summary, these findings support the notion that TE co-option as functional cis-regulatory
elements is largely a cell-type-specific process, with distinct subsets of human-specific TEs
contributing to regulatory networks in different developmental contexts. However, a more
comprehensive analysis of additional cell types should be performed to further support this
model.

300 Discussion

Transposable elements (TEs) have historically been viewed as genomic parasites with little
relevance beyond their self-propagation mechanisms. Yet, over the past two decades,
numerous studies have shown that TEs can substantially influence gene regulatory
architectures, driving lineage-specific developmental programs and morphological variation
(Chuong et al., 2017; Feschotte, 2008; Fueyo et al., 2022; Patoori et al., 2022; Sundaram &
Wysocka, 2020; Trizzino et al., 2017). In this regard, the vertebrate cranial neural crest cell
(CNCC) population, which plays a pivotal role in craniofacial morphogenesis, offers an
especially intriguing system for the investigation of TE co-option as a source of regulatory
novelty (Bronner & LeDouarin, 2012; Gokhman et al., 2021; Minoux & Rijli, 2010; Prescott et
al., 2015). Over the course of vertebrate evolution, craniofacial diversity has been shaped by
small but potent modifications in the gene regulatory circuits of the neural crest (Gokhman et
al., 2021; Prescott et al., 2015). In this context, our findings suggest that hominoid-specific
TEs contribute to these modifications by providing new enhancer platforms that alter CNCC
gene expression, pointing to a direct role for TEs in the emergence of human-specific
craniofacial features (Gokhman et al., 2021; Prescott et al., 2015).

CNCCs are a multipotent, migratory cell population that emerges from the border region of the
neural tube and subsequently disperses into developing craniofacial structures, the peripheral
nervous system, and other tissues (Simões-Costa et al., 2015; Theveneau & Mayor, 2012).
Understanding how species-specific genomic elements, such as hominoid-specific TEs,
modulate CNCC development can shed light on the evolutionary mechanisms underlying
morphological divergence, particularly in the craniofacial region (Capra et al., 2013; Prescott

et al., 2015). Specifically, in this study we demonstrate that LTR5Hs and SVAs, two families

of TEs unique to hominoids, function as enhancers within human CNCCs. Using integrative

approaches that combined chromatin accessibility profiling, transcriptomic analysis, and

functional perturbation via CRISPR-interference, we reveal that hundreds of these TEs display

active enhancer signatures and exert direct regulatory control over genes crucial for CNCC

migration. These findings are in line with previous studies showing that TEs can be repurposed

as developmental enhancers, thereby contributing to species-specific gene regulatory

landscapes in primates (Jacques et al., 2013; Patoori et al., 2022; Trizzino et al., 2017). Our

data also implicate these TE-derived enhancers in driving human-specific craniofacial

regulatory programs, as evidenced by the fact that repressing LTR5Hs and SVAs reduces the

expression of dozens of genes in human CNCCs to levels normally observed in chimpanzees.

Notably, LTR5Hs and SVAs have historically been linked to regulatory innovation in primates,

though their roles in neural crest biology have remained unexplored (Fuentes et al., 2018;

Imbeault et al., 2017; Patoori et al., 2022; Pontis et al., 2019).

One of the central questions in evolutionary developmental biology is how minor alterations in

deeply conserved developmental pathways can yield significant morphological divergence

(Carroll, 2005; King & Wilson, 1975). The CNCC population, which is indispensable for

craniofacial bone and cartilage formation, serves as an excellent proof of principle in this

regard. It has been suggested that relatively small changes in CNCC gene expression can

profoundly impact facial shape and size (Preprint: Khouri-Farah et al., 2025; Le Douarin &

Dupin, 2018; Minoux & Rijli, 2010; Preprint: Mitchell et al., 2025; Prescott et al., 2015). Our

study extends this framework by proposing that TEs represent a flexible reservoir of regulatory

motifs capable of integrating into pre-existing gene regulatory networks, fine-tuning CNCC

specification and migration. These findings are consistent with previous reports showing that

TEs have been exapted into tissue-specific enhancers in various lineages, including immune

cells, brain cell-types, embryonic stem cells, and endometrial tissue (Chuong et al., 2017;

Frost et al., 2023; Lynch et al., 2011; Trizzino et al., 2018; J. Wang et al., 2014).

Importantly, while most of the LTR5Hs elements are present a "solo LTRs" in the human

genome, a few of them, also targeted by our gRNAs, encode full-length proviral sequences

that express Rec, Env and Np9 subunits. Previous studies have shown that the Rec and Env

subunits can effectively influence cell migration, for instance in melanocytes in melanoma

(Singh et al., 2020). Therefore, we cannot rule out that the impairment of CNCC migration is

not only due to repression of the LTR5Hs-derived enhancer activity, but also to the repression

of the encoded proteins of the full-length proviral sequences.

While correlative evidence can implicate TEs as enhancers, functional validation is key to

establishing their regulatory impact. By employing a CRISPRi approach that targeted

approximately 75% of all the LTR5Hs and SVAs active in CNCCs, we observed pronounced

effects on gene expression with a strong enrichment for processes related to cell migration,

which is indispensable for CNCCs to navigate into the developing facial primordia (Theveneau

& Mayor, 2012). The accessible LTR5Hs and SVAs that remained unaffected by CRISPRi, as

they were not targeted by our gRNAs, comprised of only 68 elements. Analysis of the nearest

genes associated with these elements did not reveal enrichment for any specific biological

pathways. Nevertheless, motif enrichment analysis against scrambled backgrounds identified

significant enrichment for binding motifs of key CNCC transcription factors, including TFAP2A,

SOX9/10, and TWIST1. These findings suggest that a small subset of TEs potentially bound

by NCC transcription factors may escape effective silencing in our experimental system,

although they represent a minority of elements.

Overall, our results not only reinforce the notion that TEs influence CNCC gene networks but

also demonstrate that their regulatory input is significant enough to modulate key

developmental pathways. This model agrees with evidence from other contexts in which TEs

have driven evolutionary innovations. For instance, MER41 elements in the human genome

have been exapted as enhancers regulating immune responses (Chuong et al., 2017).

Similarly, in the endometrium and placenta, endogenous retroviral LTRs have been implicated

in the evolution of pregnancy-specific gene regulatory networks (Frost et al., 2023; Lynch et

al., 2011). Our study adds craniofacial development to the growing list of developmental

systems shaped by TE co-option, further highlighting the broad evolutionary relevance of

these mobile elements.

TE co-option relies on the inherent or mutated presence of transcription factor binding sites

within TE sequences (Barnada et al., 2022; Bejerano et al., 2006; Feschotte, 2008; Patoori et

al., 2022; Trizzino et al., 2017, 2018). In this study we show that LTR5Hs and SVAs

domesticated as enhancers in CNCCs harbour binding sites for the CNCC *coordinator* motif

(Kim et al., 2024; Prescott et al., 2015), as well as motifs for TFAP2A, SOX9/10 and other key

regulators of neural crest identity. We hypothesize that these factors bind cooperatively to

multi-part motifs within the TE loci, enabling the formation of enhancer complexes that drive

spatial and temporal gene expression programs in CNCCs. Notably, TFAP2A expression was

reduced, albeit modestly, upon TE repression. TFAP2A is critical for CNCC specification and

also migration, and it is possible that dysregulation of this factor could, at least in part, underpin

some of the aberrant migratory phenotypes.

Disruptions to CNCC gene regulatory networks can underlie congenital anomalies such as

cleft lip and palate, craniosynostosis, and other craniofacial malformations (Preprint: Khouri-

Farah et al., 2025; Trainor, 2010). Given that TE-derived enhancers are integral to the CNCC

transcriptional program, mutations or epigenetic dysregulations at these loci could contribute

to such disorders. Future studies investigating the links between TE variations and craniofacial

pathologies may reveal novel diagnostic markers or therapeutic targets.

In addition, comparative investigations in other hominoids, including gorillas and orangutans,
could help unravel whether other TEs have been similarly co-opted in the cranial neural crest.
Such research would clarify whether TE-driven craniofacial enhancer innovations are a
hallmark across all hominoids or largely unique to the human lineage, shedding light on how
small-scale genetic elements can produce macro-evolutionary changes in form and function.
Finally, our study has some limitations. Firstly, employing distinct gRNAs for SVAs and LTRHs
would be ideal to more precisely differentiate the roles of these two TE families in CNCC
development. Additionally, we assigned genes to TE-derived enhancers based on genomic
proximity, but enhancer-promoter contact maps would offer more precise enhancer-gene
associations. Unfortunately, genome-wide contact maps are currently unavailable for CNCCs.
Relatedly, the expression of a single gene is often regulated by multiple redundant enhancers,
complicating the precise determination of individual enhancer contributions without CNCC-
specific enhancer-promoter contact data. Consequently, the absence of a detectable gene
expression change upon CRISPRi does not necessarily exclude enhancer activity of a specific
TE, as compensatory enhancers might mask the loss-of-function phenotype.
Furthermore, while our Gene Ontology analysis robustly highlighted a "cell migration"
signature, functionally validated through transwell and wound healing assays, we could not
definitively identify the exact genes responsible for the altered migratory phenotype observed
upon repression of hominoid-specific TEs.
Another limitation is our reliance on uniquely mapping sequencing reads. Although this
approach ensures specificity, it likely omits biologically meaningful signals from multi-mapping
reads. Lastly, while we hypothesize that the observed phenotype arises from the co-option of
TEs into CNCC migration-associated enhancers, additional validation via techniques like
ChIP-STARR-seq combined with targeted genetic perturbations would be help substantiate
this model and confirm that disruption of these enhancers directly influences gene expression
and related cellular behaviors.
In conclusion, our findings provide further evidence for TEs as highly adaptive genomic
elements that actively contribute to shaping developmental and evolutionary trajectories.
Through combined roles in CNCC specification, migration, and potentially other lineage-
specific developmental programs, TEs demonstrate how evolutionary tinkering at the
regulatory level can yield morphological diversity.
427
428
429
430

Materials and Methods

Generation of doxycycline-inducible dCas9-KRAB hiPSC lines

A plasmid containing a tetracycline-inducible dCas9-KRAB expression cassette flanked by piggyBac recombination sites was obtained from the Wysocka Lab at Stanford University. This plasmid, referred to as 'p-dCas9-KRAB,' provides constitutive puromycin resistance, enabling selection of stable clones when co-expressed with the piggyBac transposase plasmid ('p-PB-Transposase,' Systems Bioscience). For the -gRNAs line, the C2277 hiPSC line was co-transfected with p-dCas9-KRAB and p-PB-Transposase by Applied StemCell Inc. (C2277-A, -gRNAs line). Next, a piggyBac transposon plasmid ('p-sgRNA,' Systems Bioscience) containing two sgRNAs targeting ~80% of annotated SVAs and LTR5Hs in humans was obtained based on a previously published study(Pontis et al., 2019). This plasmid confers constitutive dual sgRNA expression and geneticin resistance. For the +gRNAs line, the C2277 hiPSC line was co-transfected with p-dCas9-KRAB, p-PB-Transposase, and p-sgRNA (C2277-B, +gRNA line). Puromycin (0.125 µg/mL) only (-gRNAs line) or puromycin (0.125 µg/mL) and geneticin (100 µg/mL) (+gRNAs line) were introduced for multiple days to obtain purified colonies. Both cell lines were authenticated by STR profiling and screened for mycoplasma contamination prior to receival.

iPSC culture and NCC differentiation

Human iPSC lines C2277-A (-gRNAs) and C2277-B (+gRNAs) were cultured in feeder-free, serum-free 2D culture at 37°C and 5% CO_2. For the generation of NCCs, the differentiation method was derived from the protocol published by Leung et al. (2016). Briefly, hiPSCs were grown in mTeSR Plus (StemCell Technologies, 100-0276) on Geltrex-coated (Thermofisher, A1413302) 6-well plates until reaching ~80% confluency. Differentiation was initiated by plating 2×10^6 hiPSCs in Neural Crest Differentiation media (DMEM/F12 (Gibco, A4192002), 2% B-27 supplement (Fisher Scientific, 15717988), 3 µM CHIR99021 (Stratech, S1263-SEL-5mg), 0.5% bovine serum albumin (Fisher Scientific, 12881630), 1X glutaMAX supplement (Gibco, 35050061), 1% penicillin-streptomycin (Gibco, 15070063)) onto Geltrex-coated 6-well plates supplemented with 10 µM Y-27632 (Cambridge Bioscience, HY-119937-5MG). The following days daily media changes were performed and 2 µg/mL doxycycline was added to the culture media from day 2 onwards (i.e. 24h after the start of differentiation). NCC differentiation was assessed after 5 days.

Immunocytochemistry

HiPSCs-NCCs were harvested by incubation with StemPro Accutase (Gibco, A1110501) for 5-10 min followed by aspiration and detachment with DMEM/F12, and subsequently seeded

onto 16 mm coverslips in 12-well plates coated with Geltrex and left overnight to attach before

fixation with 4% paraformaldehyde/PBS for 15 min at 37°C. After washing with PBS, the fixed

cells were stored in PBS until staining. HiPSC-NCCs were subsequently permeabilised with

0.1% Triton-X-100/PBS (Merck, 648463) for 10 min at RT and blocked using 10% donkey

serum/PBS (Abcam, AB7475) for 1h at RT. Cells were stained with primary antibodies

overnight at 4°C in the dark, washed and stained with secondary antibodies for 30 min at 37°C.

The antibodies used can be found in Reagents and Tools Table. After a subsequent washing

step, cells were counterstained with 1 µg/mL DAPI (BioLegend, 422801). Washing steps were

performed with 0.1% Tween:PBS (Promega, H5152). Images were acquired using a Zeiss

Axio Observer and further processing was done using ImageJ. Normalised intensity was

calculated by dividing the signal intensity of each frame by the number of cells.

Western blot

Cells were lysed in radioimmunoprecipitation assay (RIPA) buffer (Fisher Scientific, 10230544)

supplemented with a Pierce protease inhibitor tablet (Fisher Scientific, 15614189). The lysates

were vortexed for 30 s and incubated on ice for 30 min. Samples were then centrifuged at

17,000 g for 10 min at 4°C, and the supernatant was collected. Protein concentration was

determined using a Pierce BCA Protein Assay Kit (Fisher Scientific, 23225) according to the

manufacturer's protocol. The plate was incubated at 37°C for 30 min and analyzed using the

SpectraMax M2 microplate reader (Molecular Devices) at 562 nm. 10 ug of protein were

denatured in Laemmli Buffer (Fisher Scientific, J61337.AC) at 95°C for 5 min. Proteins were

separated using a 4–12% Novex Tris-Glycine Mini Protein Gel (Invitrogen, XP04122BOX) in

Novex Tris-Glycine SDS Running Buffer (Fisher Scientific, 11559066) for 90 min. Proteins

were transferred onto a nitrocellulose membrane using the Mini Trans-Blot Cell system with

NuPAGE Transfer Buffer (Life Technologies, NP00061) for 1h. Membranes were blocked in

Li-Cor Intercept Blocking Buffer (Li-Cor, 927-70001) for 1h at RT, then incubated overnight at

4°C with primary antibodies (see Reagents and Tools Table) diluted in Intercept Blocking

Buffer with 0.2% Tween. The following day, washes were performed in 0.1% Tween/PBS and

membranes were incubated with secondary antibodies (see Reagents and Tools Table) for 30

min before imaging on a LiCor Odyssey XF imaging system.

Real-time qPCR

Total RNA was isolated using the Monarch Total RNA miniprep kit (New England BioLabs,

T2010) following the manufacturer's instructions. Briefly, harvested cells were lysed in RLT

buffer. The lysate was then passed through a gDNA column and centrifuged at 16,000 g for 1

min. Next, 70% ethanol was added to the lysate, which was transferred to an RNeasy silica

column and centrifuged at 16,000 g for 1 min. The column was washed followed by DNase I

treatment for 15 min at RT. Additional wash steps were performed followed by a final dry centrifugation at 12,000 rpm for 1 minute. RNA was eluted in RNase-free ddH$_2$O, and centrifuged at 16,000 g for 1 min. The eluted RNA was either used immediately or stored at -80°C. Reverse transcription into cDNA was performed using the Maxima First Strand cDNA Synthesis kit with the following thermal cycler settings: 10 min at 25°C, 15 min at 50°C, and 5 min at 85°C. Gene expression was quantified via real-time PCR PowerUp SYBR Green Supermix (Life Technologies, A46112). Primers for each gene are listed in the Reagents and Tools Table. GAPDH was used as housekeeping gene. RT-qPCR was performed using a CFX Connect Real-Time PCR Detection System (Bio-rad) with the following cycling parameters: 5 min at 95°C, 40 cycles of 15 s at 95°C, 30 s at 60°C, and 30 s at 72°C.

Transwell migration assay

24-well inserts (8 µM, Starstedt) were coated on both sides with ~22 µg/cm² Geltrex for 1 h at 37°C. After removing excess liquid, 1×10^5 hiPSC-NCCs were seeded in Neural Crest Differentiation media + 10 µM Y-27632 in the upper compartment and incubated overnight for attachment. The following day, the media was replaced and Neural Crest Differentiation media supplemented with 10% FBS was added to the lower compartment. Migration was assessed by fixing the membranes in 4% paraformaldehyde/PBS for 15 min at RT, followed by PBS washing and DAPI staining for 15 min at RT. To ensure visualization of migrated cells only, the upper surface of the insert was carefully swabbed with a cotton swab before fixation. Membranes were then excised and mounted on microscope slides for imaging. 10X images were captured from four different fields of view, and the average number of migrated cells was counted. Images were acquired using a Zeiss Axio Observer and migration was quantified using ImageJ.

Wound healing assay

HiPSC-NCCs were seeded into 24-well plates at a density of 2.5 x10^5 cells/cm^2 overnight to generate confluent monolayers. A scratch wound was made with a P200 tip and following a wash with PBS, hiPSC-NCCs were incubated with Neural Crest Differentiation media + 10% FBS. After 8h, migration was assessed and the wounds were imaged EVOS XL Core Cell Imaging System. Wound closure was measured automatically using an ImageJ plugin (Suarez-Arnedo et al., 2020).

Bulk RNA-seq library preparation

Total RNA was isolated using the Monarch Total RNA miniprep kit (New England BioLabs, T2010) following the manufacturer's instructions. RNA was quantified using a Nanodrop spectrophotometer and RNA integrity was checked on a Tapestation 2200 (Agilent). Only

samples with a RIN > 7.0 were used for transcriptome analysis. RNA libraries were prepared

using 300 ng of total RNA input using the NEBNext Poly(A) mRNA Magnetic Isolation

Module(E7490) and NEBNext® Ultra II Directional RNA Library Prep Kit for Illumina (E7760)

according to the manufacturer's instructions. PE150 sequencing was performed on a

NovaSeq X Plus instrument (Illumina).

RNA-seq analysis

Adapters were removed using TrimGalore!, and gene-level read counts were obtained with

Kallisto (Bray et al., 2016). Differential gene expression analysis was performed using DESeq2

(Love et al., 2014). GO enrichment analysis was performed using Webgestalt 2019 (Liao et

al., 2019). TE expression analysis was performed using TEtranscripts (Jin et al., 2015).

Additional statistical analysis was carried out using R (version 4.2.2) and GraphPad Prism 10.

ChIP-seq library preparation

10 million hiPSC-NCCs were crosslinked in 1% paraformaldehyde for 5 min at RT with mild

rotation. Crosslinking was quenched with 0.125M glycine by rotation for 5 min at RT. Cells

were washed twice with cold PBS and were snap-frozen for 15 min on dry ice. Samples were

stored at -80 °C until further processing. Cells were resuspended in 1 mL ChIP buffer (150

mM NaCl, 1% Triton X-100, 500 mM DTT, 10 mM Tris-HCl, 5 mM EDTA) supplemented with

Pierce protease inhibitor tablet and incubated on ice for 10 min. SDS was added to each

sample to a final concentration of 0.3%, and chromatin was sheared using the following

settings: 9 min, duty factor 5%, 6°C in a Covaris M220 Focused-Ultrasonicator. Chromatin

fragment size was assessed on a Tapestation 2200.

The chromatin lysates were then centrifuged at 13,000 g for 10 min at 4 °C and supernatant

was subsequently incubated with ChIP buffer supplemented with Pierce protease inhibitor and

Dynabeads Protein G magnetic beads (Invitrogen) along with 3 µg of H3K9me3 antibody

(Abcam, ab8898) and incubated overnight at 4 °C under mild rotation. The following day,

samples were placed in a magnetic rack and washed twice with Mixed Micelle Wash Buffer

(150 mM NaCl, 1% Triton X-100, 0.2% SDS, 20 mM Tris-HCl, 5 mM EDTA, 65% sucrose),

twice with Buffer 200 (200 mM NaCl, 1% Triton X-100, 0.1% sodium deoxycholate, 25 mM

HEPES, 10 mM Tris-HCl, 1 mM EDTA), twice with LiCl/detergent wash (250 mM LiCl, 0.5%

sodium deoxycholate, 0.5% NP-40, 10 mM Tris-HCl, 1 mM EDTA), once with cold TE. The

beads were resuspended in TE + 1% SDS and incubated at 65 °C for 10 min at 1200 rpm to

elute immunocomplexes. The elution was repeated twice and the samples were incubated

overnight at 65°C to reverse cross-linking. The following day, Proteinase K (0.5 mg/mL) was

added to digest samples at 65 °C for 1h. DNA was purified using the Zymo ChIP DNA Clean

and Concentrator kit (Zymo, D5205) and quantified with the Quantifluor ONE dsDNA system

(Promega, E4871). DNA libraries were prepared using the NEBNext Ultra II DNA library Prep
Kit (E7645L) for Illumina. PE150 sequencing was performed on a NovaSeq X Plus instrument
(Illumina).
582
ChIP-seq analysis
Adapters were removed with TrimGalore! and reads were aligned to hg38 using Burrows-
Wheeler Alignment tool, with the MEM algorithm (Li & Durbin, 2009). Uniquely mapping reads
were filtered (MAPQ>10) and PCR duplicates were removed and mitochondrial reads were
discarded. Peaks were called using HOMER with default parameters at 5% FDR (Heinz et al.,
2010). All statistical analyses were performed using BEDTools (Quinlan & Hall, 2010),
deepTools (Ramírez et al., 2014), R (version 4.2.2) and GraphPad Prism 10.
590
ATAC-seq library preparation
DNA libraries were prepared following the ATAC-Seq kit (Active Motif, 53150) according to the
manufacturer's instructions. 100,000 cells were tagmented per sample, DNA was purified
using SPRI beads and amplified with dual index primers. Libraries were assessed for size
distribution using the TapeStation 2200 and PE150 sequencing was performed on a NovaSeq
X Plus instrument (Illumina).
597
ATAC-seq analysis
Adapters were removed with TrimGalore! and reads were aligned to hg38 using Burrows-
Wheeler Alignment tool, with the MEM algorithm (Li & Durbin, 2009). With SAMTools (Li et al.,
2009), uniquely mapping reads were filtered (MAPQ>10), PCR duplicates were removed and
mitochondrial reads were discarded. Consensus peaks were determined in each cell line using
BEDTools (Quinlan & Hall, 2010). Motif analysis was performed using HOMER (Heinz et al.,
2010). All further downstream analysis was performed using BEDTools and deepTools
(Ramírez et al., 2014).
606
Statistics
RT-qPCR, transwell migration and image quantification data were analysed using Graphpad
Prism 10 software (Graphpad, San Diego, CA). Data is represented as mean ± standard error
of mean (SEM). Statistical significance was accepted at p <0.05. Sample sizes were not
determined using statistical methods but were based on previous experiments under similar
conditions. Blinding was not performed on experiments but data analysis was automated for
all experiments (e.g. intensity tresholds or automated plugins in FIJI). Sequencing data was
analysed using BEDTools, DeepTools, and R as indicated. Motif analysis was performed using
HOMER. GO enrichment analysis was performed using Webgestalt 2019.

Acknowledgements

The authors thank the Wysocka group at Stanford (and particularly Dr. Raquel Fueyo) for providing the dCAS9-KRAB vector, and Andrew Isopi and Dr. Samantha Barnada (Thomas Jefferson University), for designing the plasmid with the sg-RNAs. We thank the Facility for Imaging by Light Microscopy (FILM) at Imperial College London for providing access to instrumentation and technical support, which is part-supported by funding from the Wellcome Trust (grant 104931/Z/14/Z) and BBSRC (grant BB/L015129/1). For this work, MT was funded by Biotechnology and Biological Sciences Research Council (BBSRC, grant BB/Y000854/1).

Data availability

RNA-seq, ATAC-seq, and ChIP-seq data have been deposited in the Gene Expression Omnibus (GEO) under accession code GEO: GSE292478. (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE292478) and are publicly available as of the date of manuscript submission.

Disclosure and competing interest statement

The authors declare no competing interests.

Author contributions

MT and LD designed the project. LD performed most of the experiments. ZHM, MD and BGDV contributed to some of the experiments. LD, MT, and AK analyzed the data. LD and MT and wrote the manuscript. All the authors read and approved the manuscript.

References

Ahlstrom, J. D., & Erickson, C. A. (2009). The neural crest epithelial-mesenchymal transition in 4D: A 'tail' of multiple non-obligatory cellular mechanisms. *Development*, *136*(11), 1801–1812. https://doi.org/10.1242/dev.034785

Bajpai, R., Chen, D. A., Rada-Iglesias, A., Zhang, J., Xiong, Y., Helms, J., Chang, C. P., Zhao, Y., Swigut, T., & Wysocka, J. (2010). CHD7 cooperates with PBAF to control multipotent neural crest formation. *Nature*, *463*(7283), 958–962. https://doi.org/10.1038/nature08733

Barnada, S. M., Giner de Gracia, A., Morenilla-Palao, C., López-Cascales, M. T., Scopa, C., Waltrich, F. J., Mikkers, H. M. M., Cicardi, M. E., Karlin, J., Trotti, D., Peterson, K. A., Brugmann, S. A., Santen, G. W. E., McMahon, S. B., Herrera, E., & Trizzino, M. (2024). ARID1A-BAF coordinates ZIC2 genomic occupancy for epithelial-to-mesenchymal transition in cranial neural crest specification. *The American Journal of Human Genetics*. https://doi.org/10.1016/j.ajhg.2024.07.022

Barnada, S. M., Giner de Gracia, A., Morenilla-Palao, C., López-Cascales, M. T., Scopa, C., Waltrich, F. J., Mikkers, H. M. M., Cicardi, M. E., Karlin, J., Trotti, D., Peterson, K. A., Brugmann, S. A., Santen, G. W. E., McMahon, S. B., Herrera, E., & Trizzino, M. (2024). Gene Expression Omnibus GSE261846 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE261846). [DATASET]

Barnada, S. M., Isopi, A., Tejada-Martinez, D., Goubert, C., Patoori, S., Pagliaroli, L., Tracewell, M., & Trizzino, M. (2022). Genomic features underlie the co-option of SVA transposons as cis-regulatory elements in human pluripotent stem cells. *PLoS Genetics*, *18*(6), 1–19. https://doi.org/10.1371/journal.pgen.1010225

Bejerano, G., Lowe, C. B., Ahituv, N., King, B., Siepel, A., Salama, S. R., Rubin, E. M., Kent, W. J., & Haussler, D. (2006). A distal enhancer and an ultraconserved exon are derived from a novel retroposon. *Nature*, *441*(1), 87–90. https://doi.org/10.1038/nature04696

Bourque, G., Leong, B., Vega, V. B., Chen, X., Yen, L. L., Srinivasan, K. G., Chew, J. L., Ruan, Y., Wei, C. L., Huck, H. N., & Liu, E. T. (2008). Evolution of the mammalian transcription factor binding repertoire via transposable elements. *Genome Research*, *18*(11), 1752–1762. https://doi.org/10.1101/gr.080663.108

Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Near-optimal probabilistic RNA-seq quantification. *Nature Biotechnology*, *34*(5), 525–527. https://doi.org/10.1038/nbt.3519

Bronner, M. E., & LeDouarin, N. M. (2012). Development and evolution of the neural crest: An overview. In *Developmental Biology* (Vol. 366, Issue 1, pp. 2–9). Academic Press Inc. https://doi.org/10.1016/j.ydbio.2011.12.042

Capra, J. A., Hubisz, M. J., Kostka, D., Pollard, K. S., & Siepel, A. (2013). A Model-Based Analysis of GC-Biased Gene Conversion in the Human and Chimpanzee Genomes. *PLoS Genetics*, *9*(8). https://doi.org/10.1371/journal.pgen.1003684

Carroll, S. B. (2005). Evolution at two levels: On genes and form. In *PLoS Biology* (Vol. 3, Issue 7, pp. 1159–1166). Public Library of Science. https://doi.org/10.1371/journal.pbio.0030245

Chuong, E. B., Elde, N. C., & Feschotte, C. (2016). Regulatory evolution of innate immunity through co-option of endogenous retroviruses. *Science*, *351*(6277), 1083–1087. https://doi.org/10.1126/science.aad5497

Chuong, E. B., Elde, N. C., & Feschotte, C. (2017). Regulatory activities of transposable elements: From conflicts to benefits. In *Nature Reviews Genetics* (Vol. 18, Issue 2, pp. 71–86). Nature Publishing Group. https://doi.org/10.1038/nrg.2016.139

Chuong, E. B., Rumi, M. A. K., Soares, M. J., & Baker, J. C. (2013). Endogenous retroviruses function as species-specific enhancer elements in the placenta. *Nature Genetics*, *45*(3), 325–329. https://doi.org/10.1038/ng.2553

Cordero, D. R., Brugmann, S., Chu, Y., Bajpai, R., Jame, M., & Helms, J. A. (2011). Cranial neural crest cells on the move: Their roles in craniofacial development. *American Journal of Medical Genetics, Part A*, *155*(2), 270–279. https://doi.org/10.1002/ajmg.a.33702

Cosby, R. L., Judd, J., Zhang, R., Zhong, A., Garry, N., Pritham, E. J., & Feschotte, C. (2021). Recurrent evolution of vertebrate transcription factors by transposase capture. *Science*, *371*(6531). https://doi.org/10.1126/science.abc6405

Feschotte, C. (2008). Transposable elements and the evolution of regulatory networks. *Nature Genetics*, *9*, 397–405.

Frost, J. M., Amante, S. M., Okae, H., Jones, E. M., Ashley, B., Lewis, R. M., Cleal, J. K., Caley, M. P., Arima, T., Maffucci, T., & Branco, M. R. (2023). Regulation of

human trophoblast gene expression by endogenous retroviruses. *Nature Structural and Molecular Biology*, *30*(4), 527–538. https://doi.org/10.1038/s41594-023-00960-6

Fuentes, D., Swigut, T., & Wysocka, J. (2018). Systematic perturbation of retroviral LTRs reveals widespread long-range effects on human gene regulation. *ELife*. https://doi.org/10.7554/eLife.35989.001

Fueyo, R., Judd, J., Feschotte, C., & Wysocka, J. (2022). Roles of transposable elements in the regulation of mammalian transcription. *Nature Reviews Molecular Cell Biology*, *23*(7), 481–497. https://doi.org/10.1038/s41580-022-00457-y

Gans, C., & Northcutt, R. G. (1983). Neural Crest and the Origin of Vertebrates: A New Head. *Science*, *220*(4594), 268–273. https://doi.org/10.1126/science.220.4594.268

Gokhman, D., Agoglia, R. M., Kinnebrew, M., Gordon, W., Sun, D., Bajpai, V. K., Naqvi, S., Chen, C., Chan, A., Chen, C., Petrov, D. A., Ahituv, N., Zhang, H., Mishina, Y., Wysocka, J., Rohatgi, R., & Fraser, H. B. (2021). Human–chimpanzee fused cells reveal cis-regulatory divergence underlying skeletal evolution. *Nature Genetics*, *53*(4), 467–476. https://doi.org/10.1038/s41588-021-00804-3

Goubert, C., Zevallos, N. A., & Feschotte, C. (2020). Contribution of unfixed transposable element insertions to human regulatory variation. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *375*(1795). https://doi.org/10.1098/rstb.2019.0331

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., Cheng, J. X., Murre, C., Singh, H., & Glass, C. K. (2010). Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Molecular Cell*, *38*(4), 576–589. https://doi.org/10.1016/j.molcel.2010.05.004

Imbeault, M., Helleboid, P. Y., & Trono, D. (2017). KRAB zinc-finger proteins contribute to the evolution of gene regulatory networks. *Nature*, *543*(7646), 550–554. https://doi.org/10.1038/nature21683

Jacques, P. É., Jeyakani, J., & Bourque, G. (2013). The Majority of Primate-Specific Regulatory Sequences Are Derived from Transposable Elements. *PLoS Genetics*, *9*(5). https://doi.org/10.1371/journal.pgen.1003504

Jheon, A. H., & Schneider, R. A. (2009). The cells that fill the bill: Neural crest and the evolution of craniofacial development. In *Journal of Dental Research* (Vol. 88, Issue 1, pp. 12–21). https://doi.org/10.1177/0022034508327757

Jin Y, Tam OH, Paniagua E, Hammell M. (2015) TEtranscripts: a package for including transposable elements in differential expression analysis of RNA-seq datasets. *Bioinformatics*. 31(22):3593-9.

Khouri-Farah, N., Winchester, E. W., Schilder, B. M., Robinson, K., Curtis, S. W., Skene, N. G., Leslie-Clarkson, E. J., & Cotney, J. (2025). Gene expression patterns of the developing human face at single cell resolution reveal cell type contributions to normal facial variation and disease risk. *bioRxiv*. https://doi.org/10.1101/2025.01.18.633396

Kim, S., Morgunova, E., Naqvi, S., Goovaerts, S., Bader, M., Koska, M., Popov, A., Luong, C., Pogson, A., Swigut, T., Claes, P., Taipale, J., & Wysocka, J. (2024). DNA-guided transcription factor cooperativity shapes face and limb mesenchyme. *Cell*. https://doi.org/10.1016/j.cell.2023.12.032

Kim, S., Morgunova, E., Naqvi, S., Goovaerts, S., Bader, M., Koska, M., Popov, A.,
Luong, C., Pogson, A., Swigut, T., Claes, P., Taipale, J., & Wysocka, J. (2024).
Gene Expression Omnibus GSE230319
(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE230319) [DATASET]
King, M.-C., & Wilson, A. C. (1975). Evolution at Two Levels in Humans and
Chimpanzees. *Science*, *188*, 107–116. https://www.science.org
Kunarso, G., Chia, N. Y., Jeyakani, J., Hwang, C., Lu, X., Chan, Y. S., Ng, H. H., &
Bourque, G. (2010). Transposable elements have rewired the core regulatory
network of human embryonic stem cells. *Nature Genetics*, *42*(7), 631–634.
https://doi.org/10.1038/ng.600
Le Douarin, N. M., & Dupin, E. (2018). The "beginnings" of the neural crest. In
Developmental Biology (Vol. 444, pp. S3–S13). Elsevier Inc.
https://doi.org/10.1016/j.ydbio.2018.07.019
Leung, A. W., Leung, A. W., Murdoch, B., Salem, A. F., Prasad, M. S., Gomez, G. A.,
& García-Castro, M. I. (2016). WNT/β-catenin signaling mediates human neural
crest induction via a pre-neural border intermediate. *Development (Cambridge)*,
143(3), 398–410. https://doi.org/10.1242/dev.130849
Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-
Wheeler transform. *Bioinformatics*, *25*(14), 1754–1760.
https://doi.org/10.1093/bioinformatics/btp324
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., & Durbin, R. (2009). The Sequence Alignment/Map format and
SAMtools. *Bioinformatics*, *25*(16), 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352
Liao, Y., Wang, J., Jaehnig, E. J., Shi, Z., & Zhang, B. (2019). WebGestalt 2019:
gene set analysis toolkit with revamped UIs and APIs. *Nucleic Acids Research*,
47(W1), W199–W205. https://doi.org/10.1093/nar/gkz401
Lieberman, D. E. (1998). Sphenoid shortening and the evolution of modern human
cranial shape. *Nature*, *393*(6681), 158–162. https://doi.org/10.1038/30227
Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12).
https://doi.org/10.1186/s13059-014-0550-8
Lynch, V. J., Leclerc, R. D., May, G., & Wagner, G. P. (2011). Transposon-mediated
rewiring of gene regulatory networks contributed to the evolution of pregnancy in
mammals. *Nature Genetics*, *43*(11), 1154–1159. https://doi.org/10.1038/ng.917
Martik, M. L., & Bronner, M. E. (2021). Riding the crest to get a head: neural crest
evolution in vertebrates. In *Nature Reviews Neuroscience* (Vol. 22, Issue 10, pp.
616–626). Nature Research. https://doi.org/10.1038/s41583-021-00503-2
Minoux, M., & Rijli, F. M. (2010). Molecular mechanisms of cranial neural crest cell
migration and patterning in craniofacial development. In *Development* (Vol. 137,
Issue 16, pp. 2605–2621). https://doi.org/10.1242/dev.040048
Mitchell, Z. H., den Hoed, J., Claassen, W., Demurtas, M., Deelen, L., Campeau, P.
M., Liu, K., Fisher, S. E., & Trizzino, M. (2025). CHD3 regulates BMP signalling
response during cranial neural crest cell specification. *bioRxiv*.
https://doi.org/10.1101/2025.02.14.638260
Ostertag, E. M., Goodier, J. L., Zhang, Y., & Kazazian, H. H. (2003). SVA Elements
Are Nonautonomous Retrotransposons that Cause Disease in Humans. In *Am.
J. Hum. Genet* (Vol. 73).
Pagliaroli, L., Porazzi, P., Curtis, A. T., Scopa, C., Mikkers, H. M. M., Freund, C.,
Daxinger, L., Deliard, S., Welsh, S. A., Offley, S., Ott, C. A., Calabretta, B.,

Brugmann, S. A., Santen, G. W. E., & Trizzino, M. (2021). Inability to switch from
ARID1A-BAF to ARID1B-BAF impairs exit from pluripotency and commitment
towards neural crest formation in ARID1B-related neurodevelopmental
disorders. *Nature Communications*, *12*(1). https://doi.org/10.1038/s41467-021-
26810-x
Patoori, S., Barnada, S. M., Large, C., Murray, J. I., & Trizzino, M. (2022). Young
transposable elements rewired gene regulatory networks in human and
chimpanzee hippocampal intermediate progenitors. *Development (Cambridge)*,
49(23). https://doi.org/10.1242/dev.200413
Playfoot, C. J., Duc, J., Sheppard, S., Dind, S., Coudray, A., Planet, E., & Trono, D.
(2021). Transposable elements and their KZFP controllers are drivers of
transcriptional innovation in the developing human brain. *Genome Research*,
31(9), 1531–1545. https://doi.org/10.1101/gr.275133.120
Pontis, J., Planet, E., Offner, S., Turelli, P., Duc, J., Coudray, A., Theunissen, T. W.,
Jaenisch, R., & Trono, D. (2019). Hominoid-Specific Transposable Elements and
KZFPs Facilitate Human Embryonic Genome Activation and Control
Transcription in Naive Human ESCs. *Cell Stem Cell*, *24*(5), 724-735.e5.
https://doi.org/10.1016/j.stem.2019.03.012
Prescott, S. L., Srinivasan, R., Marchetto, M. C., Grishina, I., Narvaiza, I., Selleri, L.,
Gage, F. H., Swigut, T., & Wysocka, J. (2015). Enhancer Divergence and cis-
Regulatory Evolution in the Human and Chimp Neural Crest. *Cell*, *163*(1), 68–
83. https://doi.org/10.1016/j.cell.2015.08.036
Prescott, S. L., Srinivasan, R., Marchetto, M. C., Grishina, I., Narvaiza, I., Selleri, L.,
Gage, F. H., Swigut, T., & Wysocka, J. (2015). Gene Expression Omnibus
GSE70751 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE70751)
[DATASET]
Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for
comparing genomic features. *Bioinformatics*, *26*(6), 841–842.
https://doi.org/10.1093/bioinformatics/btq033
Quinn, J. P., & Bubb, V. J. (2014). SVA retrotransposons as modulators of gene
expression. *Mobile Genetic Elements*, *4*(4), e32102.
https://doi.org/10.4161/mge.32102
Ramírez, F., Dündar, F., Diehl, S., Grüning, B. A., & Manke, T. (2014). DeepTools: A
flexible platform for exploring deep-sequencing data. *Nucleic Acids Research*,
42(W1). https://doi.org/10.1093/nar/gku365
Roadmap Epigenomics Consortium, Kundaje, A., Meuleman, W., Ernst, J., Bilenky,
M., Yen, A., Heravi-Moussavi, A., Kheradpour, P., Zhang, Z., Wang, J., Ziller, M.
J., Amin, V., Whitaker, J. W., Schultz, M. D., Ward, L. D., Sarkar, A., Quon, G.,
Sandstrom, R. S., Eaton, M. L., … Kellis, M. (2015). Integrative analysis of 111
reference human epigenomes. *Nature*, *518*(7539), 317–329.
https://doi.org/10.1038/nature14248
Roadmap Epigenomics Consortium, Kundaje, A., Meuleman, W., Ernst, J., Bilenky,
M., Yen, A., Heravi-Moussavi, A., Kheradpour, P., Zhang, Z., Wang, J., Ziller, M.
J., Amin, V., Whitaker, J. W., Schultz, M. D., Ward, L. D., Sarkar, A., Quon, G.,
Sandstrom, R. S., Eaton, M. L., … Kellis, M. (2015). Gene Expression Omnibus
GSE16256 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE16256)
[DATASET]
Sambataro, S., Fiorillo, L., Bocchieri, S., Stumpo, C., Cervino, G., Herford, A. S., &
Cicciù, M. (2022). Craniofacial Evolution: From Australopithecus to Modern Man.

Journal of Craniofacial Surgery, *33*(1), 325–332.
https://doi.org/10.1097/SCS.0000000000007841

Schmidt, D., Schwalie, P. C., Wilson, M. D., Ballester, B., Gonalves, Â., Kutter, C., Brown, G. D., Marshall, A., Flicek, P., & Odom, D. T. (2012). Waves of retrotransposon expansion remodel genome organization and CTCF binding in multiple mammalian lineages. *Cell*, *148*(1–2), 335–348. https://doi.org/10.1016/j.cell.2011.11.058

Simões-Costa, M., Stone, M., & Bronner, M. E. (2015). Axud1 Integrates Wnt Signaling and Transcriptional Inputs to Drive Neural Crest Formation. *Developmental Cell*, *34*(5), 544–554. https://doi.org/10.1016/j.devcel.2015.06.024

Singh, M., Cai, H., Bunse, M., Feschotte, C., & Izsvák, Z. (2020). Human endogenous retrovirus k rec forms a regulatory loop with mitf that opposes the progression of melanoma to an invasive stage. *Viruses*, *12*(11). https://doi.org/10.3390/v12111303

Spoor, F., Wood, B., & Zonneveld, F. (1994). Implications of early hominid labyrinthine morphology for evolution of human bipedal locomotion. *Nature*, *369*(6482), 645–648. https://doi.org/10.1038/369645a0

Suarez-Arnedo, A., Figueroa, F. T., Clavijo, C., Arbeláez, P., Cruz, J. C., & Muñoz-Camargo, C. (2020). An image J plugin for the high throughput image analysis of in vitro scratch wound healing assays. *PLoS ONE*, *15*(7 July). https://doi.org/10.1371/journal.pone.0232565

Sundaram, V., Cheng, Y., Ma, Z., Li, D., Xing, X., Edge, P., Snyder, M. P., & Wang, T. (2014). Widespread contribution of transposable elements to the innovation of gene regulatory networks. *Genome Research*, *24*(12), 1963–1976. https://doi.org/10.1101/gr.168872.113

Sundaram, V., & Wysocka, J. (2020). Transposable elements as a potent source of diverse cis-regulatory sequences in mammalian genomes. In *Philosophical Transactions of the Royal Society B: Biological Sciences* (Vol. 375, Issue 1795). Royal Society Publishing. https://doi.org/10.1098/rstb.2019.0347

Theveneau, E., & Mayor, R. (2012). Neural crest migration: interplay between chemorepellents, chemoattractants, contact inhibition, epithelial–mesenchymal transition, and collective cell migration. *WIREs Developmental Biology*, *1*(3), 435–445. https://doi.org/10.1002/wdev.28

Trainor, P. A. (2010). Craniofacial birth defects: The role of neural crest cells in the etiology and pathogenesis of Treacher Collins syndrome and the potential for prevention. In *American Journal of Medical Genetics, Part A: Vol. 152 A* (Issue 12, pp. 2984–2994). https://doi.org/10.1002/ajmg.a.33454

Trizzino, M., Kapusta, A., & Brown, C. D. (2018). Transposable elements generate regulatory novelty in a tissue-specific fashion. *BMC Genomics*, *19*(1). https://doi.org/10.1186/s12864-018-4850-3

Trizzino, M., Park, Y. S., Holsbach-Beltrame, M., Aracena, K., Mika, K., Caliskan, M., Perry, G. H., Lynch, V. J., & Brown, C. D. (2017). Transposable elements are the primary source of novelty in primate gene regulation. *Genome Research*, *27*(10), 1623–1633. https://doi.org/10.1101/gr.218149.116

Wang, H., Xing, J., Grover, D., Hedges Kyudong Han, D. J., Walker, J. A., & Batzer, M. A. (2005). SVA elements: A hominid-specific retroposon family. *Journal of Molecular Biology*, *354*(4), 994–1007. https://doi.org/10.1016/j.jmb.2005.09.085

Wang, J., Xie, G., Singh, M., Ghanbarian, A. T., Raskó, T., Szvetnik, A., Cai, H., Besser, D., Prigione, A., Fuchs, N. V., Schumann, G. G., Chen, W., Lorincz, M.

903   C., Ivics, Z., Hurst, L. D., & Izsvák, Z. (2014). Primate-specific endogenous
904   retrovirus-driven transcription defines naive-like stem cells. *Nature*, *516*(7531),
905   405–409. https://doi.org/10.1038/nature13804
906   Wray, G. A. (2007). The evolutionary significance of cis-regulatory mutations. *Nature*
907   *Reviews Genetics*, *8*(3), 206–216. https://doi.org/10.1038/nrg2063

Figure legends

Figure 1. An inducible CRISPRi tool to repress human-specific SVAs and LR5Hs in iPSC-derived CNCCs – (a) Schematic overview of SVA and LTR5Hs transposable elements, illustrating their structure and phylogenetic distribution across Old World monkeys, apes, and humans. Human-specific SVA subfamilies (i.e., SVA_E/F/F1) and the LTR5Hs are highlighted, along with the approximate number of elements in each category. **(b)** CRISPR-dCas9-KRAB targeting strategy for SVAs and LTR5Hs. Two guide RNAs (gRNA1 and gRNA2) target ~80% of all SVAs and LTR5Hs, directing dCas9-KRAB to deposit repressive epigenetic marks at the TEs, thereby reducing their transcriptional activity. **(c)** Experimental timeline for doxycycline (dox)-induced expression of dCas9-KRAB. Dox is added to the media 24 hours after the start of differentiation into cranial neural crest cells, which spans 5 days. **(d)** Western blot confirming dox-inducible dCas9-KRAB expression. Lysates from cells cultured with or without TE-targetting gRNAs were probed with antibodies against dCas9 (top band, ~160 kDa) and GAPDH (bottom band, loading control). **(e)** Immunofluorescence displaying expression of CNCC signature markers AP2α (red) and SOX9 (green) in CNCCs differentiated using CRISPRi-iPSCs without (left) and with gRNAs (right). Signal intensity was quantified and normalised to the cell number per frame. This experiment was repeated at least once. **(f)** RT-qPCR of CNCC markers *SOX9*, *TWIST1*, *SOX10* and *TFAP2A*. n=6: each data point is derived from 6 independent rounds of NCC differentiation. Statistical analyses were performed using unpaired t-tests. Error bars indicate mean ± SEM. ***, p=0.0002.

Figure 2 – Hundreds of human-specific SVAs and LTR5Hs function as enhancers in human CNCCs – (a) ATAC-seq signal across human-specific SVAs and LTR5Hs. Heatmap rows represent individual transposable elements (TEs) aligned at their start or end (±500 bp), and the intensity reflects chromatin accessibility. Clusters were generated with deepTools using k-mer algorithm. Two independent replicates per group. **(b)** Pie charts showing the observed versus expected fractions of accessible TEs belonging to the LTR5Hs/SVA subfamilies. **(c)** H3K9me3 enrichment at the accessible human-specific SVAs and LTR5Hs. The top panel displays the average H3K9me3 signal (average profile) aligned to the TE start

or end (±3 kb), while the heatmap below shows individual TEs ranked by signal intensity. **(d)**

Transcription factor motif analysis of accessible human-specific SVAs and LTR5Hs.

Representative enriched motifs (TWIST1, HOMEOBOX, SOX9/10, AP2α, SLUG, and FOXD3)

in accessible human-specific TEs are shown with the percentage of TEs containing each motif.

(e) Heatmaps of TWIST1 ChIP-seq signal (left) and corresponding input control (right) centred

on the accessible human-specific SVAs and LTR5Hs (±2 kb). Colour intensity indicates

normalized enrichment, highlighting TWIST1 occupancy at these accessible elements.

Figure 3. Human-specific SVAs and LTR5Hs regulate the expression of CNCC migration

genes – (a) Violin plots displaying expression (log2 TPM) of genes located near accessible

human-specific SVAs and LTR5Hs. Three conditions are shown: human CNCCs generated

using CRISPRi iPSC line without guide RNAs (−gRNA), human CNCCs generated using

CRISPRi iPSC line with guide RNAs (+gRNA), and chimpanzee CNCCs. The decreased

expression in the +gRNA condition indicates potential regulatory contributions of these human-

specific TEs. **(b)** Volcano plot illustrating differentially expressed genes between the −gRNA

(n=3) and +gRNA (n=2) conditions in human CNCCs. The x-axis shows the log2 fold change,

while the y-axis represents the −log10 P-value. Significantly upregulated (green) and

downregulated (red) genes are highlighted, with total numbers indicated. Dashed lines mark

common significance thresholds. **(c)** Functional enrichment analysis of downregulated genes.

Bubble plots display top enriched pathways, with the x-axis indicating enrichment ratio (blue)

or the -log10 FDR (red) and the y-axis listing pathway terms. The size of each bubble reflects

statistical significance. Pathways involved in cell migration and motility are among the most

prominently enriched.

Figure 4. HiPSC-NCCs show inferior migration capability when human-specific TEs are

silenced. (a) Outline of the transwell migration assay. **(b)** 1 x 10^5 hiPSC-NCCs (+/- gRNAs)

were seeded overnight, followed by a media change in the upper chamber and incubation with

differentiation medium +10% FBS in the lower chamber for 24h. At 24h, non-migrated hiPSC-

NCCs on the top of the membrane were swabbed away and migrated hiPSC-CMs were fixed,

stained and imaged. DAPI staining was used to visualise nuclei. Scale bar: 50 µm. **(c)**

Quantification of transwell membranes. 10X images were taken from 4 different fields of view

and the average number of migrated cells was calculated. n=5. ***,p= 0.0008. **(d)** 5 x 10^4

hiPSC-NCCs (+/- gRNAs) were seeded overnight onto a 24-well plate. The next day, scratch

wounds were made and the hiPSC-NCCs were incubated with differentiation medium + 10%

FBS. After 8h, scratch wounds were imaged and wound closure was measured. n=12 (-

gRNAs), n=11 (+gRNAs). Scale bar: 200 µm. **(e)** Quantification of scratch wound closure was

performed with an automated plugin (Suarez-Arnedo et al., 2020). ****,p= 2.56 × 10^{-6}. Both

assays were independently performed twice. Statistical analyses were performed using an
unpaired t-test. Error bars indicate mean ± SEM.
977
Figure 5. Enhancer signature of SVAs and LTR5Hs is cell-type specific – Heatmaps of
normalized H3K27ac ChIP-seq signal (Roadmap Epigenomics Consortium) across 256
accessible human-specific SVAs and LTR5Hs in multiple tissues and cell types. Each row
corresponds to an individual transposable element (TE), aligned at its centre (±3 kb), while
columns represent different tissues/cell lines. The colour bar on the right indicates the relative
intensity of H3K27ac signal, with higher enrichment in red and lower enrichment in blue.

A

ATAC-seq

-gRNAs   -gRNAs   +gRNAs   +gRNAs

HUMAN-SPECIFIC SVAs and LTR5Hs

TE START/END +/- 500 bp

0   10   20   30   40   50

B

Observed
Accessible in CNCCs:

LTR5Hs
SVA-E/-F

Expected

29%

Expected

LTR5Hs
SVA-E/-F

Expected

LTR5Hs
SVA-E/-F

C

H3K9me3

ACCESSIBLE HUMAN-SPECIFIC SVAs and LTR5Hs (N = 256)

-gRNAs   +gRNAs

TE 0.5Kb

TE START/END +/- 3 KB

D

**Motifs enriched in accessible
human-specific SVAs and LTR5HS**

TWIST1	CCAGCTGTTC	83%
HOMEOBOX	TTAATTAAA	62%
SOX9/10	CCATTTGTTCG	88%
TFAP2A	ATGCCCTGAGGC	65%
SLUG	GGGCACCTGCTG	62%
FOXD3	TGTTTATTTAGC	32%

COORDINATOR
MOTIF

E

TWIST1   INPUT

ACCESSIBLE HUMAN-SPECIFIC SVAs and LTR5Hs (N = 256)

TE CENTER +/- 2 KB

A

Genes with decreased expression
near accessible human-specific TEs

B

501   294

C

Pathways enriched in the set of differentially expressed genes