The role of active mRNA-ribosome dynamics and closing constriction in daughter chromosome separation in *Escherichia coli*

Chathuddasie I. Amarasinghe[1], Mu-Hung Chang[1], Jaana Männik[1], Scott T. Retterer[2], Maxim O. Lavrentovich*[1,3], Jaan Männik*[1]

[1]Department of Physics and Astronomy, University of Tennessee, Knoxville, TN, USA,

[2]Center for Nanophase Materials Sciences, Oak Ridge National Laboratory, Oak Ridge, TN, USA,

[3]Department of Earth, Environment, and Physics, Worcester State University, Worcester, MA, USA.

Correspondence: mlavrent@utk.edu, jmannik@utk.edu

1   **Abstract**

The mechanisms by which two sister chromosomes separate and partition into daughter cells in bacteria
remain poorly understood. A recent theoretical model has proposed that out-of-equilibrium processes
associated with mRNA–ribosome (polysome) dynamics play a significant role in this process. Here we
investigate the role of ribosomal dynamics on nucleoid segregation and separation in *Escherichia coli* using
high-throughput fluorescence microscopy in microfluidic devices. We compare our experimental
observations with predictions from a reaction-diffusion model that includes the interactions among
ribosomal subunits, polysomes, and chromosomal DNA. Our results show that the non-equilibrium behavior
of mRNA and ribosomes causes them to aggregate at the midcell and this process contributes to the
separation of the two daughter chromosomes. However, this effect is considerably weaker than that
predicted by the model. Rather than relying solely on active mRNA–ribosome dynamics, our data suggest
that the closing division septum via steric interactions and potentially entropic forces between two DNA
strands coupled to cell elongation act as additional mechanisms to ensure faithful partitioning of the
nucleoids to two daughter cells.

15   **Significance**

The mitotic spindle separates chromosomes in eukaryotic cells, but bacteria lack this structure. It remains
unclear how bacterial chromosomes partition prior to cell division. It has been hypothesized that non-
equilibrium dynamics of polysomes, that is mRNA–ribosome complexes, actively drive the separation of
bacterial chromosomes. Using quantitative microscopy combined with computational modeling, we show
that polysome dynamics significantly contribute to chromosome segregation in *Escherichia coli* but this
process does not constitute the sole mechanism. Our findings suggest the closing division septum via steric
interactions and potentially entropic forces between two DNA strands act as additional mechanisms.

Introduction

Chromosome replication and segregation are essential processes in the life cycle of a cell (1-3). Segregation can be divided into two stages: the individualization of daughter chromosomes, and their further separation, which enables their partitioning into daughter cells during cytokinesis (4). Completion of both stages is required for the successful propagation of cells from one generation to the next.

In bacteria, the individualization stage, commonly referred to as DNA segregation, occurs for the most part concurrently with DNA replication (5). In *Escherichia coli*, the newly synthesized DNA loci near *oriC* begin to separate from each other with some delay after initiation of DNA replication (6). After this initial cohesion phase, the sister loci segregate from each other sequentially in the order in which they appear in the genome (7-9). Despite extensive previous research, the underlying molecular mechanism driving the segregation phase in bacteria remains poorly understood (2,3,10). Several active mechanisms for chromosome segregation have been investigated, including ParAB (ParA-ParB-*parS*) (2,10) and homologous Spo0J-Soj system (11), but these are missing in many bacterial species, including *E. coli* (12,13). In addition to the Par-systems, structural maintenance of chromosomes (SMC) proteins, encoded by MukBEF in *E. coli*, have also been implicated in the active segregation (14,15). SMC involvement in segregation is postulated to occur via a successive condensation of DNA around *oriC* via loop extrusion and repulsion of condensed DNA masses away from each other due to excluded volume interactions and chain entropy (14,15). While this is a plausible scenario, MukBEF is not essential for DNA segregation in *E. coli*, except possibly at the fastest growth rates when multifork replication is present (16).

Rather than relying on dedicated protein machinery for segregation, which would expend energy through ATP hydrolysis, it is plausible that the free energy needed for the segregation of chromosomal DNA is derived mostly from the replication process itself. Replication creates two newly formed proximal DNA strands, which, according to polymer physics, experience a repulsive entropic force in confined conditions (17,18). This scenario has been supported by numerous computer simulations (19-26) but it has yet to be experimentally validated. However, the driving force from this so-called entropic mechanism vanishes once the spatial overlap between the two nucleoids becomes zero (26). Therefore, if only the entropic mechanism were present, the two individualized and segregated nucleoids would remain in contact rather than fully separating from each other. The entropic mechanism can, thus, explain the segregation but not the separation phase of the two nucleoids.

The question then arises: what drives chromosomes apart during the separation stage? In eukaryotes the separation is carried out by the mitotic spindle, but no spindles are known to exist in bacteria. DNA translocase FtsK, which is a part of the bacterial cell division apparatus (divisome) (27-29) could be involved in this process (28). However, cells with inactivated FtsK (FtsK K997A) (30) appear to separate their nucleoids and partition them into two daughter cell compartments in the majority of cells (31).

Evidence for a different mechanism for separation was revealed by Wu *et al.* (32). These authors observed
that in filamentous cells (more than 10 μm) with two nonreplicating nucleoids, the nucleoids separated from
each other over large distances so that their centers stayed close to the cell's ¼ positions (32). Wu *et al.*
explained such positioning by osmotic pressure from the accumulation of macromolecular crowders, but
their explanation relied on the assumption that these crowders are produced in a specific way outside the
two nucleoids (32). An alternative explanation, which did not rely on this assumption, was proposed by
Miangolarra *et al.* (33). These authors developed a model that included reactions between chromosomal
DNA, mRNA, ribosomal subunits and polyribosomes (polysomes), along with their diffusion within the cell.
This reaction-diffusion model, which we refer to as the active mRNA-ribosome dynamics model, included
reactions for mRNA synthesis, ribosome assembly and disassembly on mRNA, and mRNA degradation.
The diffusion of these species was affected by the steric (excluded volume) interactions between these
molecules. The model predicted that due to these interactions, the 30S and 50S ribosomal subunits can
relatively unhindered diffuse into the nucleoid region, consistent with single-molecule tracking experiments
(34,35), while 70S ribosomes, which are further assembled into polyribosomes (polysomes), cannot. The
key insight from the model was a finding that an active mRNA synthesis within the nucleoid and the
assembly of 30S and 50S ribosomal subunits into polysomes around these mRNAs create instability within
the nucleoid. This instability drives the nucleoid apart into two distinct nucleoid lobes at some critical cell
length (33). The model also predicted that the two daughter nucleoid centers would position at the quarter
points of the cell once the transition occurred and stay in these locations even when cells filamented,
consistent with the experimental observations by Wu *et al*. (32).

Here, we investigate the mechanisms that individualize chromosomes in the late stages of segregation and
drive their separation in normal growth conditions. In particular, we test the active mRNA-ribosome
dynamics model by Miangolarra *et al.* in non-filamenting cells growing in slow-growth conditions with simple
DNA topology. To this end, we use high-throughput microscopy in microfluidic devices to determine the
distributions of ribosomes, nucleoids, and replication machinery during cell and replication cycles. We
quantitatively compare the acquired experimental data to our own reaction-diffusion model, which is
conceptually similar to the one proposed by Miangolarra *et al*. (33). We first find that the termination of DNA
replication does not have a significant effect on the nucleoid segregation dynamics. Instead, DNA expands
and daughter nucleoids separate in proportion to cell length increase during this stage of the cell cycle. Our
data, when compared to the model, supports the idea that active ribosome dynamics facilitates separation
of two nucleoids in normal growth conditions. However, in cells where active ribosome dynamics are halted,
we observe that the separation of daughter nucleoids still increases proportionally with cell length. In
addition to this effect, we identify closing constriction, as a new mechanism, that has a significant effect in
separating the two daughter nucleoids and partitioning them into two daughter cells. Based on this
evidence, we propose a multi-process model to explain the nucleoid segregation and separation dynamics
in *E. coli*.

Results

The midcell DNA density minimum evolves independent of replication termination

DNA segregation cannot occur without DNA being first replicated. We first hypothesized that the progression of segregation is directly driven by the replication stage. In previous works on the segregation of nucleoids using tracking of fluorescently labeled loci in the chromosome (5,6,9,36,37), following bulk segregation (25,38), or labeling the replication terminus by MatP fusion (25,39), the segregation process had not been directly tracked during the replication cycle at the single cell level. To do so now, we labeled the beta-clamp protein, DnaN, with fluorescent fusion (Ypet-DnaN) (40). To follow the dynamics of the whole chromosome, we used HupA-mCherry, which has been tested before and is known to bind uniformly throughout the chromosome (41). We chose to carry out measurements in slow growth conditions where, at most, only a single replication cycle is ongoing (42) and where we could accurately determine timings for the initiation and termination of the replication (SI Fig. S1). We imaged cells in mother machine microfluidic devices (43,44), obtaining about 30-50 time points per cell cycle from several hundred cells. We averaged these data over the cell population and produced kymographs (Fig. 1A-B).

To characterize the segregation process, we determined the total extent of all nucleoids in the cell, $L_{nuc,ext}$, and the sum of nucleoid lengths, $L_{nuc,sum}$ (Fig. 1C; for details, see Materials and Methods). These measurements showed that the outer edges of the nucleoids, as determined by $L_{nuc,ext}$, stayed at about a constant distance (0.5/2 μm ≈ 0.25 μm) from the cell ends throughout the replication cycle (Fig. 1C). Notably, the total extent and the sum of nucleoid lengths did not show any discontinuities at the termination of replication (Fig. 1C). To further verify this unexpected result, we aligned the curves from individual cells relative to replication termination time, Trt, but still found only smooth continuous changes (SI Fig. S2). We also found a comparable cell-cycle independent distance throughout the division cycle in both slow (SI Fig. S3) and moderately fast growth conditions (SI Fig. S4). The measured exclusion distance of nucleoids from poles (0.25 μm) is slightly less than the length of the hemispherical cap region (0.3-0.35 μm) (45) . Therefore, the chromosomal DNA is mostly excluded from the polar caps. The data furthermore show that instead of nucleoid elongation and separation being controlled by the replication cycle, the length of individual nucleoids, $L_{nuc,sum}$, and their total extent in the cell, $L_{nuc,ext}$, increases in proportion to cell length. This dependence has also been noticed in some earlier studies (25,32,46), although the replication period was not directly determined in these works.

In addition to $L_{nuc,sum}$ and $L_{nuc,ext}$, we also determined the DNA density at midcell during the replication cycle as a different measure of DNA segregation and separation (Fig. 1D). In the early stages of replication, there was a maximum DNA density at midcell (Fig. 1A). Upon completion of about 60% of the replication cycle, the DNA amount at the midcell started to decrease (Fig.1D). This decrease corresponded to the formation of a local DNA density minimum (Fig. 1E). As with our other nucleoid measurements, the midcell minimum continued to decline further after replication termination without any distinct changes. Altogether,

these results, combined with the nucleoid length measurements, indicate that while chromosome replication
is a prerequisite for segregation, nucleoid segregation is not strongly driven by the replication process itself
at late stages of the replication cycle. Instead, separation of two nucleoids follows overall cell growth.

Correlated and anti-correlated periods of DNA and ribosome density distributions during the cell
cycle

In balanced growth, cells size increases proportional to its ribosome number. We hypothesized that the
DNA segregation at late stages of the replication cycle could be driven by active ribosome dynamics that
involve mRNA synthesis and degradation and are associated with the assembly and disassembly of
polyribosomes (33). To test this hypothesis, we constructed a strain where the nucleoids were labeled with
HupA-mYpet fluorescent marker (25,41) and ribosomes with RpsB-mCherry (47,48) (for details, see
Materials and Methods). Both labels had been tested in previous studies (25,41,47,48). Since this strain
lacked a marker for replication, we followed nucleoid segregation and ribosome accumulations throughout
the division cycle (Fig. 2A-C), which we customarily refer to as the cell cycle.

In the first half of the cell cycle, the population-averaged data shows a unimodal distribution of DNA with a
maximum close to the cell center (Fig. 2D, Age 0.10, 0.30). Notably, the ribosome density distribution is
similar to the DNA density distribution from the same cells, showing a broad density maximum at the midcell.
The co-localization of the two distribution maxima appears not consistent with equilibrium thermodynamic
models (49,50), where ribosomes phase separate from DNA rich regions. Note that there are no local
maxima in the ribosome density at cell poles in distributions in Fig. 2D. The excess ribosome density at cell
poles is not expected to result in maxima in the extreme ends of 1D density distributions, which are
projections of the 3D density onto the long axis of the cell. The maxima are missing because of the rounded
shape of the poles and diffraction-limited imaging. As the cell cycle progresses, the nucleoid expands, and
the population-average DNA density distribution becomes bimodal (bilobed (6)) with a local minimum near
its center (Fig.2D, Age=0.50). This minimum is accompanied by the sharper ribosome density maximum at
the same location. In this case, the observed anti-correlated densities of ribosomes and chromosomal DNA
at the midcell are consistent with the phase separation of nucleoid and ribosomes, as expected from
equilibrium thermodynamics (49,50). This excess ribosome density increases as the cell cycle progresses,
and the DNA density at midcell decreases (Fig. 2D, Age=0.50, 0.70). At the end of the cell cycle, the
ribosome density, instead of further increasing, decreases while the DNA density continues also to decrease
(Fig. 2D, Age=0.90). This positively correlated behavior again cannot be explained by equilibrium
considerations of volume exclusion between ribosomes and chromosomes.

To determine the timing of these behaviors, we plot the Pearson correlation coefficient, R, between the DNA
and ribosome distributions along the long axis of the cell as a function of cell cycle age, expressed as the
fraction of completed cell cycle (Fig. 2E). At the beginning of the cell cycle, the two distributions show small
but positive R values, as expected based on the coincidence of their maxima. At about cell age 0.5, the

correlation coefficient starts to decrease and becomes negative. This decrease in R corresponds to a
concurrent decrease in DNA and an increase in ribosome amounts at midcell (Fig. 2F). The correlation
coefficient reaches its minimum at cell age ~0.75, then starts to increase and becomes positive by the end
of the cell cycle. At cell cycle age ~0.75, the midcell ribosome numbers reach their peak and start to
decrease thereafter, while the amount of DNA at midcell continues to decrease until the end of the cell cycle
(Fig. 2F). To understand the cause of the decrease in ribosome numbers at midcell at the end of the cell
cycle, we compare these data with the midcell phase contrast signal from the same cells (Fig. 2F, black
line). There is an upturn in the phase contrast signal at the time when the ribosome numbers peaked. The
increase in the phase contrast signal signifies the onset of constriction (51). The midcell ribosome numbers,
thus, start to decrease as the cell starts to constrict in this region.

Based on traces in Fig. 2D the excess fraction of ribosomes in the midcell appeared small throughout the
cell cycle. To quantify this fraction, we decompose the ribosomal density distribution into a midcell fraction
and a uniform background by fitting the data from individual cells (see Materials and Methods). We find that
the population average excess fraction of ribosomes is only 4% of the cell total at its peak at cell age ~0.75
(Fig. 2G). At the same time, the excess fraction before the DNA density minimum at midcell begins to form
is 2.5%. The equilibrium thermodynamics can qualitatively explain the increase in the excess fraction from
2.5% to 4%. However, it fails to explain the 2.5% fraction present in the early cell cycle.

To verify the validity of the above findings, we constructed a strain in which the fluorescent mCherry fusion
was added to the C-terminus of the 50S ribosomal protein RplI (RplI-mCherry) instead of labeling the 30S
subunit (SI Fig. S5A-C). This label has also been previously characterized and tested (51). The cells with
RplI-mCherry label showed the same behavior as the one with RpsB-mCherry (SI Fig. S5D-G), including
co-localization of ribosome and DNA density distributions at the midcell at early cell cycle (SI Fig. S5D,
Age=0.15) and comparably small excess fraction of ribosomes at midcell at its peak value (3.5%) (SI Fig.
S5G).

We also investigated whether the same behaviors would occur in faster growth rates in M9 glucose + CAS
medium (SI Fig. S6). Under this growth condition, the replication cycle initiates predominantly in the
previous cell cycle and completes at the early cell cycle (42). Accordingly, a minimum in the nucleoid density
distribution is already observed at the midcell at cell birth (SI Fig. S6C). Consequently, unlike in slow growth
conditions, there is no positive correlation between DNA and ribosome density distribution during the early
stages of the cell cycle (SI Fig. S6D). Aside from this difference, the other behaviors (SI Fig. S6E-F),
including a small excess ribosome fraction at the cell center at its peak value (5.5%), are consistent with
those observed under slower growth conditions.

The DNA and ribosome correlation patterns in the early and midcell cycle are qualitatively explained by the coupled DNA and ribosome dynamics model

To understand if the observed ribosome and nucleoid localization patterns can be explained by the active dynamics of mRNA and ribosomes, we developed our own reaction-diffusion model (see Methods, *The Coupled DNA and Ribosome Dynamics Model*). We then compared the calculated cell cycle-dependent DNA and ribosome densities from the model to the experimental data. Our 1D model is similar to the one developed by Miangalorra *et al*. (33), considering mRNA synthesis and degradation, as well as the assembly of polysomes (polyribosomes) from ribosomal subunits. Polysome and ribosomal subunits interact with each other and with DNA via excluded volume interactions. However, unlike the previously published model, ours uses a power-law term (50,52) instead of a virial expansion to describe DNA and polysome self-interactions. Additionally, our model does not explicitly simulate bare mRNA and polysomes of varying sizes; instead, it considers only an average-sized polysome. Unlike the previous model (33), ours also includes cell growth, assuming DNA and ribosome amounts increase with cell length (or volume).

To qualitatively compare the model to experimental data, we plotted the nucleoid and ribosome density distribution along the cell length at different cell cycle times (Fig. 2H, SI Fig. S7A-B). These times were selected to be comparable, though not identical to those shown in Fig. 2D. Notably, the ribosome distribution predicted by the model shows a small local maximum at the cell middle, coinciding with the maximum of the nucleoid distribution at the early stages of the cell cycle (Fig. 2H, Age 0.10, 0.30). This is consistent with experimental data (Fig. 2D, Age 0.15). In the model, the midcell maximum in ribosome distribution arises at the start of the cell cycle. This occurs because 30S and 50S ribosomal subunits can diffuse almost unhindered into the nucleoid region, where they assemble into polysomes. Since the rate of polysome formation is proportional to DNA density in the model, the maxima of ribosome and DNA densities coincide.

The model predicts that excess ribosome density at the midcell cleaves the initial single nucleoid into two nucleoid masses at some critical cell length due to excluded volume interactions (Fig 2H, Age=0.32). After this transition, ribosome and DNA density distributions become anti-correlated (Fig 2H, Age=0.72; SI Fig. S7C), which is consistent with experimental observations. Separation of the two DNA masses in the model occurs rapidly, within 5% (0.05) of the cell cycle. During this transition, the excess fraction of ribosomes at midcell reaches 48% and approximately maintains this value for the rest of the cycle (SI Fig. S7D). In contrast, the experiment shows that the fraction reaches only 4.5% (Fig. 2G). The model thus correctly predicts a local maximum in ribosome density at the cell middle in the early stages. However, it shows a significantly higher accumulation of ribosomes in the cell middle in the later stages of the cell cycle compared to the experimental data. We hypothesized that the substantially higher ribosome fractions observed in the model are due to its 1D nature. In a 3D cell, ribosomes could form an outer shell around the nucleoid, as has been proposed by earlier single-molecule tracking experiments of ribosomes (34). Since this shell could not be derived from our 1D model, we added its effect *ad hoc* to our model (SI Fig. S8A-B). Additionally, we assumed a uniform ribosome volume density in the shell that matches the density

predicted by the 1D model at the cell poles (see *The Concentric Shell Model* in Material and Methods for
details). With this addition, the experimental (Fig. 2D) and modeling densities (Fig. 2I) showed similar values
with the model yielding about 3% excess of ribosomes at midcell at its peak (SI Fig. S8E). However, to
reach this agreement, the inner radius of the added shell needed to be 50% of the cell radius while it has
been estimated based on single-molecule tracking experiments that the inner radius is about 80% of the
cell's radius (34). These comparisons suggest that the coupled DNA and ribosome dynamics model could
better explain the data if it were implemented in 3D.

Constriction pushes aside daughter nucleoids

In the late stages of the cell cycle, when a constriction forms at the midcell, the model and experimental
data do not agree with each other. The data reveals a decreasing ribosome fraction at the midcell (Fig. 2G),
whereas the model maintains almost a constant fraction (SI Fig. S7E). We hypothesized that the closing
constriction pushes aside the ribosomes and potentially also the nucleoids. To test this, we carried out
experiments in which constriction formation was inhibited by the beta-lactam antibiotic cephalexin. It is well
known that despite the inhibition of constriction, cephalexin-treated cells continue to grow in volume and
replicate DNA at the same rates as untreated cells (53). We then compared the changes in ribosome and
DNA densities between the cephalexin-treated and untreated cells. Since cephalexin prevents cell division,
cell age cannot be used to track the progression of DNA and ribosome densities, as done in the previous
measurements in Fig. 2. Instead, we determined the timing for the first appearance of the local DNA density
minimum at the cell center, T_{min}, following the procedure defined earlier (42) (for details see Materials and
Methods). We then aligned the time-dependent data relative to this time (Fig. 3A, B) processing the data
from untreated cells the same way (SI Fig. S9A-B). At T_{min}, both the treated and untreated cells show an
increase in midcell ribosome numbers (Fig. 3C) and a simultaneous decrease in the midcell DNA amount
(Fig. 3D), consistent with the excluded volume interactions between ribosomes and DNA. However, at later
times ($time - T_{ab} > 50$ min), the midcell ribosome numbers stay constant in treated cells, while they
decrease in untreated ones (Fig. 2C). This comparison confirms that the decrease in ribosome numbers in
untreated cells is caused by the closing constriction as we hypothesized.

The data also indicate that, in addition to reducing the number of ribosomes at midcell, the constriction
affects the separation of two daughter nucleoids. Comparison of DNA density at the midcell reveals a clear
decrease in DNA in untreated cells relative to treated ones for $time - T_{ab} > 50$ min (Fig. 3D). The effect of
constriction is also noticeable in the measured distance between two nucleoid lobes increasing more
significantly in untreated cells than in treated ones (Fig. 3E). Altogether, these data show that the closing
constriction separates two chromosomes in the late stages of the cell cycle. The effect is not caused by
FtsK, since FtsK is recruited to the divisome in both cephalexin-treated and untreated cells (54). The
constriction-driven separation appears, thus, to be a steric effect: the closing constriction physically pushes
aside the daughter nucleoids.

In untreated cells in slow growth conditions, the onset of constriction occurs effectively at the time of the
DNA density minimum, T_{min} (42). Therefore, the effects arising from the active ribosome dynamics and
constriction on the separation of two daughter nucleoids are overlayed for $time > T_{min}$. However, the data
from cephalexin-treated cells allows us to quantitatively compare the coupled dynamics of DNA and
ribosome separation with our model (for $time > T_{min}$). The DNA and ribosome density distributions
predicted by the model (Fig. 3F,G) are qualitatively similar to the experimental data from cephalexin-treated
cells (Fig. 3A,B). In quantitative comparisons, the model reproduced some trends in the experimental data
but missed others. The model quantitatively correctly predicts the distance between the peaks of two
nucleoid lobes (Fig. 3H), but fails to predict the time-dependent decrease in the amount of DNA at midcell
(Fig. 3I). In the experiments, the DNA amount changes only 10% over 60 min, while in the model it changes
50% within about 5 min. However, after this initial 5 min period, the model and experiment show comparable
rates of DNA density decrease at the midcell (Fig. 3I). The reason for the ten-fold slower segregation
dynamics observed in the experiments could be related to the choice of model parameters. Reducing the
diffusion coefficient of DNA 10-fold considerably slowed down the rate of the change in DNA amount at the
midcell, although the cell cycle time for the transition for DNA density from high to low values shifted to later
times (SI Fig. S10).

The experimental data show that after an initial increase at T_{min}, both the excess midcell ribosome fraction
(Fig. 3J) and the width of the excess ribosome distribution at midcell (Fig. 3K) remain nearly constant in
cephalexin-treated cells. The reasons behind this steady behavior are not intuitively clear, given that the
total ribosome number continues to rise as cells elongate. Notably, the model qualitatively reproduces these
observed trends but does not accurately predict their numerical values. The quantitative discrepancy for
the excess mid cell ribosome fraction (Fig. 3J) can potentially be resolved by including an *ad hoc* concentric
ribosome shell around the nucleoid, as previously suggested. Additionally, the experimentally measured
width of the ribosome distribution at midcell (Fig. 3K) is approximately three times greater than predicted
by the model. This discrepancy likely arises from uncertainties in the chosen Cahn–Hilliard parameters (κ),
for which no independent experimental measurements are yet available. Altogether, the model shows
qualitative agreement with the experimental data. Its quantitative accuracy could be potentially improved
by extending the model to three dimensions and refining its unknown parameters through additional
experiments and optimization.

The effect of disruption of polysome formation on nucleoid separation

To perturbatively test the effects of active ribosome dynamics on the separation of two nucleoids, we treated
E. coli cells with a lethal concentration of rifampicin (300 µg/ml), which inhibits the initiation of RNA synthesis
(55). As in the previous measurements, we also added cephalexin to the culture medium at the same time
as rifampicin to prevent cells from constricting. Since rifampicin inhibits RNA synthesis, no new polysomes
can form after the drug reaches the cells. The existing transcripts can still complete, but this occurs on the

time scale of tens of seconds for a single gene and about a minute for longer operons. These timescales

are much shorter than our measurement frame rate (4 min). On a slightly longer time scale (3-5 min (56)),

the existing polysomes disassemble as their associated mRNAs degrade, releasing ribosomal subunits.

To understand the effects arising from these processes, we compared data from cells treated

simultaneously with rifampicin and cephalexin (Fig. 4A–C) to cells treated only with cephalexin (Fig. 4D–

F). In this comparison, we align the time-dependent data to the antibiotic treatment start time, T_{ab}, instead

of T_{min} because most Rif-treated cells lack a detectable midcell DNA density minimum. Additionally, we

excluded from the analysis cells with visible constriction at the start of the treatment. As expected, the

localized accumulation of ribosomes at the midcell disappears following rifampicin treatment (Fig. 4B, C)

due to polysome dissociation and subsequent diffusion of free ribosomal subunits onto the nucleoid region.

Following rifampicin addition, the midcell DNA density minimum almost immediately ceases to deepen,

although it persists rather than disappearing (Fig. 4G). If steric interactions between DNA and polysomes

alone were responsible for creating this minimum, we would expect its complete disappearance on a

timescale comparable to polysome dissociation. In contrast, cephalexin-only treated cells continue to show

a gradual decrease in midcell DNA density (Fig. 4G). This difference between treatment conditions indicates

that active mRNA-ribosome dynamics contribute significantly to establishing the midcell DNA density

minimum but are not solely responsible.

We next examined nucleoid segregation by measuring the distance between nucleoid centers (D_{peak}) in

rifampicin-treated cells (Fig. 4H, SI Fig. S11A). This distance continues to increase for approximately 50

minutes, long after active ribosome dynamics ceases (within minutes of rifampicin addition) and polysomes

dissociate (3–5 min). Under normal growth conditions, nucleoids separate proportionally to cell elongation

(Fig. 1C). To determine if a similar relationship exists following rifampicin treatment, we compared nucleoid

center distances to cell length increases, normalizing both metrics at the time of antibiotic addition (Fig. 4H,

SI Fig. S11). Indeed, even without active ribosome dynamics, nucleoid centers continue separating

proportionally to cell elongation, demonstrating that cell growth alone is sufficient to drive nucleoid

segregation (Fig. 4I). However, rifampicin treatment also results in nucleoid expansion, causing increased

midcell DNA density when two nucleoids are present (Fig. 4G). Collectively, these findings show that

chromosome segregation progresses despite active ribosome dynamic and polysome dissociation while

the separation of daughter chromosomes is affected.

To compare these data to our model, we divide the rifampicin-treated cells into three age groups based on

their nucleoid segregation stages. This grouping was necessary because rifampicin treatment starts at

different cell cycle ages for the asynchronous cell population in Fig. 4. The first group, representing 12% of

cells, includes those cells where T_{min} was detected before treatment began ($T_{min} < T_{ab}$, $N = 92$) (SI Fig.

S12). The second group, comprising 17% of cells, includes cells in which nucleoid separation was detected

after treatment started ($T_{min} > T_{ab}$, $N = 138$). The largest group, accounting for 71% of cells ($N = 556$),

shows no detectable nucleoid separation either before or after treatment. Our following analysis focuses
on the first two groups, as the third group shows minimal changes in ribosome and nucleoid distributions
after the treatment. To model these two groups of cells, we set the polysome formation rate to zero at the
start of the treatment. We then evolve the model so that cell length changes the same way as in the
experiments (see Materials and Methods, *Rifampicin Treatment in the Model*, for details).

The group of cells that were late in their replication cycle at the onset of antibiotic treatment ($T_{min} < T_{ab}$),
show a local DNA density minimum (Fig. 5A) and a midcell ribosome accumulation at the start of rifampicin
treatment (Fig. 5B). The midcell ribosome accumulation vanishes within about 16 min from the start of the
treatment (Fig. 5B-C). The same occurs in the model, although at a faster rate (Fig. 5E-F). At the same
time, the model and the data qualitatively differ in how DNA distributions evolved. The experimental data
show that the midcell DNA density minimum persists, albeit becoming shallower over a 50-minute period,
before plateauing (Fig. 5A,C, SI Fig. S13A). In contrast, in the model, the two nucleoid masses merge so
that the local DNA density minimum at the midcell completely vanishes (Fig.5D,F, SI Fig. S13B-C). The
coupled DNA ribosome dynamics model is thus not able to explain the persistence of the experimentally
observed nucleoid minimum after treatment.

We next examine the second group of cells ($T_{min} > T_{ab}$) where the DNA density minimum develops after
the start of rifampicin treatment (Fig. 5G-I, SI Fig. S14A). Although the observed DNA density minimum is
small in this group, its presence also contradicts the active mRNA-ribosome dynamics model. In the model,
the midcell DNA density minimum does not form without active ribosome dynamics (Fig. 5J-L, SI Fig. S14B-
C). The lack of the minimum is a qualitative feature of the model, independent of parameter selection. This
results from the overdamped nature of diffusive processes. Once the driving force is removed, that is, the
formation of polysomes is halted, then the separation of two nucleoid lobes stops immediately (SI Fig.
S14B-C). The qualitative discrepancy between the model and the data for cells with $T_{min} > 0$ show that
active dynamics of mRNA and ribosomes is not responsible for DNA separation in this group of cells.

Discussion

Active mRNA-ribosome dynamics

This study provides both experimental and theoretical evidence supporting the role of active mRNA-
ribosome dynamics in the segregation and separation of chromosomal DNA in *E. coli* cells. A characteristic
feature of this mechanism, as also observed in our measurements (Fig. 2), is the accumulation of
polyribosomes (polysomes) at the nucleoid (and cell) center before it starts to split into two lobes. According
to our model, the accumulation of polyribosomes results from strong excluded volume interactions between
polysomes and chromosomal DNA (50), leading to the exclusion of DNA from the midcell region. These
qualitative features of the model are consistent with our data in steady-state growth conditions. Perturbative
measurements with rifampicin treatment furthermore show that the nucleoid split does not develop in the

majority of cells (77%) when polysome formation is inhibited due to the lack of mRNA substrates. However,
these measurements also show that in cells where nucleoid lobes have already formed at the start of the
treatment, the nucleoid centers continue to move apart as the cells continue to elongate (Fig. 4H, SI Fig.
S13C).

The latter finding indicates that the active mRNA-ribosome dynamics is not the sole process driving the late
segregation of nucleoids. Similar conclusions were recently reported by Papagiannakis *et al.,* although
based on different evidence (38). Notably, their measurements did not show further separation of nucleoid
lobes after the start of rifampicin treatment. This discrepancy likely arises from differences in growth
conditions between the two studies. Under faster growth rates, as used by Papagiannakis *et al.,* the cell
growth stops almost immediately after rifampicin treatment begins. In comparison, under the slower growth
conditions used in our study, cell growth continues for a considerable portion of time (more than 50 min in
our case). Accordingly, in faster growth conditions, further nucleoid lobe separation is not observed, given
that it is driven by continued cell elongation.

Entropic mechanism

What is the mechanism driving two nucleoid lobes apart in proportion to cell elongation, independent of
active mRNA-ribosome dynamics in the late stages of segregation? First, it can be ruled out that this
process is due to the passive macromolecular crowding, which remains present after rifampicin treatment.
Indeed, passive crowding without reactions could lead to aggregation of DNA at high crowding levels or
simply hinder DNA segregation and separation due to increased viscosity. Consistent with this view, a fusion
of nucleoids can be observed in *E. coli* cells treated with chloramphenicol (57), which stops protein
synthesis but not RNA synthesis. In this case, the high level of macromolecular crowding appears to result
from an increased level of RNA species (rRNA, tRNA, mRNA) in the cytosol (58). Alternatively, it has been
proposed that cell elongation could drive nucleoid segregation via transertion linkages (59). However, this
hypothesis can be ruled out by our rifampicin measurements because the linkages would disappear within
a few minutes from the start of the treatment, whereas the daughter nucleoids continue to move apart for a
much longer period (50 min).

At the same time, the cell elongation-dependent movement of nucleoid lobes apart can be consistent with
the entropic segregation mechanism (17,18,25). Polymer physics predicts that two daughter chromosomes,
under cylindrical confinement, will segregate from each other such that the center of each nucleoid remains
at the ¼ position of the cell, as long as two chromosomes remain in contact with each other (32). The
distance between the nucleoid centers, D_{peak} scales then proportional to cell length as observed in our
experiments. For the nucleoids to position exactly at the cell's ¼ positions the macromolecular crowding
effects need to be sufficiently weak so that each nucleoid fills half of the cell volume. This appears the case
in rifampicin treatment measurements. However, at stronger macromolecular crowding levels, the osmotic
pressure from crowders, can push two nucleoids together when there is no active mRNA-ribosome

dynamics in the end of the segregation phase. D_{peak} being proportional to L_{cell} then no longer holds. It is

unclear if the daughter nucleoids in normal growth conditions without rifampicin treatment experience weak

or strong crowding levels. We also emphasize that entropic mechanism is applicable only for the

segregation phase and it cannot explain any separation of two nucleoids.

Constriction effect

Our measurements also indicate that, in addition to active mRNA-ribosome dynamics a closing constriction

plays an important role in separating the two daughter nucleoids and partitioning them into daughter cell

compartments (Fig. 3). This likely occurs via steric interactions between the constriction and DNA. These

steric interactions could potentially be mediated by crowders, ensuring that the inner membrane and

chromosomal DNA do not come into direct contact. Interestingly, earlier research has shown that the onset

of constriction is blocked until DNA segregation has progressed to the point where there is a local DNA

density minimum at midcell /mid nucleoid (42). If the constriction were to start closing earlier, then it could

likely result in the trapping of incompletely replicated DNA under the constriction. Altogether, DNA

segregation to the point where there is a local DNA density minimum at the midcell appears to trigger the

onset of constriction, which completely separates one daughter chromosome from another.   The

segregation process can thus be self-amplifying through the formation of the constriction.

FtsK translocase

Constriction, in the majority of cases, provides a faithful mechanism for partitioning the two chromosomes

to the daughter cells. When it fails and some DNA becomes trapped by the closing septum, then DNA

translocase FtsK (60,61) provides a last-resort mechanism to correct the error (31). FtsK usually acts on

about 400 kb region near the replication termination region around the *dif*-site (28) but it is capable of

directional pumping of more than 3 Mb of chromosomal DNA across closing constriction (31). This activity

mostly occurs at the very end of the constriction when the pore separating two daughters has a diameter

of less than 250 nm (31). The pore does not close fully until all the DNA is pumped across it. Altogether,

FtsK effectively guarantees a fail-safe completion of DNA partitioning. However, in most cell divisions, FtsK

is dispensable, as mutant cells are viable even when FtsK translocase activity has been inhibited (31,62).

Integration

In summary, our measurements and modeling, combined with findings from previous research, bring out

the involvement of several partially overlapping processes in the segregation and separation of daughter

chromosomes in *E. coli*. Based on these findings, we propose the following sequence of events unfolding

during the cell cycle (Fig. 6): The entropic mechanism is likely involved during the late segregation stage

when two chromosomes overlap (17,18). When about half of the DNA is replicated in slow growth

conditions, active mRNA-ribosome dynamics via excluded volume interactions become a sufficiently strong

driving force to lead to a midcell DNA density minimum. At this point, the entropic mechanism of segregation

still remains important, and, in fact, the associated force increases as more DNA is replicated (25,26).

However, the entropic force vanishes once two DNA molecules no longer overlap. In parallel, the formation
of the midcell DNA density minimum via the active mRNA-ribosome dynamics triggers the onset of
constriction in slow growth conditions (42). The constriction further drives the two daughter chromosomes
apart and contributes to their separation. The final stages of DNA partitioning are further aided by FtsK if
DNA becomes trapped by the closing constriction.

The involvement of active mRNA-ribosome dynamics in the segregation of the two DNA molecules in the
process outlined above (Fig. 6) may appear to some degree redundant and may not be essential for
partitioning the DNA molecules into daughter cells. However, this mechanism is inherently present in
actively growing cells where gene expression takes place. Harnessing transcription and translation
processes to accelerate DNA segregation offers a clear advantage, especially in growth conditions where
DNA segregation is a limiting factor for the duration of the cell cycle (42,63).

Outlook and future directions

Further work is needed to validate the proposed processes outlined in Fig. 6. In particular, the existing
evidence for the entropic mechanism remains indirect. Using synthetic cell systems implemented in
microfluidic platforms, such as the genome-in-the-box (64) or minimal cells (65), will help further elucidate
the role of this mechanism. An open question is also whether additional molecular systems beyond those
listed in Fig. 6 are involved in the segregation and separation of chromosomal DNA in *E. coli*.

Significant effort is still required to develop models that can simulate these processes in sufficient detail,
ideally integrating them into a single platform. For the active mRNA-ribosome dynamics models, it remains
unclear to what degree the findings from 1D models apply in a 3D setting. Additionally, these reaction-
diffusion models lack chain connectivity for DNA, which may alter the results. We hope the experimental
evidence presented here motivates further computational efforts to address these gaps in our
understanding.

Materials and Methods

Details of bacterial growth conditions, microchip design, fluorescence microscopy setup, and image
analysis, and modeling are given in the SI Information, Materials and Methods. The list of all strains and
plasmids used in this study, including their sources, is given in the SI Information, Table S1.

Acknowledgments

The authors thank Rachel McCord and Steve Abel for their valuable comments. A part of this research was
conducted at the Center for Nanophase Materials Sciences, which is sponsored at Oak Ridge National
Laboratory by the Scientific User Facilities Division, Office of Basic Energy Sciences, U.S. Department of
Energy. This work was supported by the National Institutes of Health award GM127413 (JM) and NSF
MCB2313719 (JM).

References

1. Bloom K, Joglekar A (2010) Towards building a chromosome segregation machine. *Nature* **463**:446-456. doi:10.1038/nature08912.

2. Wang X, Llopis PM, Rudner DZ (2013) Organization and segregation of bacterial chromosomes. *Nat. Rev. Genet.* **14**:191-203. doi:10.1038/nrg3375.

3. Gogou C, Japaridze A, Dekker C (2021) Mechanisms for chromosome segregation in bacteria. *Frontiers in Microbiology* **12**:685687. doi:10.3389/fmicb.2021.685687.

4. Batty P, Gerlich DW (2019) Mitotic chromosome mechanics: how cells segregate their genome. *Trends in Cell Biology* **29**:717-726. doi:10.1016/j.tcb.2019.05.007.

5. Woldringh CL, Hansen FG, Vischer NOE, Atlung T (2015) Segregation of chromosome arms in growing and non-growing *Escherichia coli* cells. *Frontiers in Microbiology*. **6**. doi:10.3389/fmicb.2015.00448.

6. Bates D, Kleckner N (2005) Chromosome and replisome dynamics in *E. coli*: Loss of sister cohesion triggers global chromosome movement and mediates chromosome segregation. *Cell* **121**:899-911. doi:10.1016/j.cell.2005.04.013.

7. Viollier PH, Thanbichler M, McGrath PT, West L, Meewan M, McAdams HH, Shapiro L (2004) Rapid and sequential movement of individual chromosomal loci to specific subcellular locations during bacterial DNA replication. *Proc. Natl. Acad. Sci. U. S. A.* **101**:9257-9262. doi:10.1073/pnas.0402606101.

8. Nielsen HJ, Ottesen JR, Youngren B, Austin SJ, Hansen FG (2006) The *Escherichia coli* chromosome is organized with the left and right chromosome arms in separate cell halves. *Mol. Microbiol.* **62**:331-338. doi:10.1111/j.1365-2958.2006.05346.x.

9. Cass JA, Kuwada NJ, Traxler B, Wiggins PA (2016) *Escherichia coli* chromosomal loci segregate from midcell with universal dynamics. *Biophys J* **110**:2597-2609. doi:10.1016/j.bpj.2016.04.046.

10. Badrinarayanan A, Le TBK, Laub MT (2015) Bacterial chromosome organization and segregation. *Annu Rev Cell Dev Biol* **31**:171-199. doi:10.1146/annurev-cellbio-100814-125211.

11. Lin DC, Grossman AD (1998) Identification and characterization of a bacterial chromosome partitioning site. *Cell* **92**:675-685. doi:10.1016/s0092-8674(00)81135-6.

12. Livny J, Yamaichi Y, Waldor MK (2007) Distribution of centromere-like parS sites in bacteria: insights from comparative genomics. *J Bacteriol* **189**:8693-8703. doi:10.1128/JB.01239-07.

13. Cornet F, Blanchais C, Dusfour-Castan R, Meunier A, Quebre V, Sekkouri Alaoui H, . . . Bouet JY (2023) DNA segregation in enterobacteria. *EcoSal Plus* **11**:eesp00382020. doi:10.1128/ecosalplus.esp-0038-2020.

14. Brahmachari S, Marko JF (2019) Chromosome disentanglement driven via optimal compaction of loop-extruded brush structures. *Proc. Natl. Acad. Sci. U. S. A.* **116**:24956-24965. doi:10.1073/pnas.1906355116.

15. Harju J, van Teeseling MCF, Broedersz CP (2024) Loop-extruders alter bacterial chromosome topology to direct entropic forces for segregation. *Nat Commun* **15**:4618. doi:10.1038/s41467-024-49039-w.

16. Makela J, Sherratt D (2020) SMC complexes organize the bacterial chromosome by lengthwise compaction. *Curr Genet* **66**:895-899. doi:10.1007/s00294-020-01076-w.

17. Jun S, Mulder B (2006) Entropy-driven spatial organization of highly confined polymers: lessons for the bacterial chromosome. *Proc. Natl. Acad. Sci. U. S. A.* **103**:12388-12393. doi:10.1073/pnas.0605305103

18. Jun S, Wright A (2010) Entropy as the driver of chromosome segregation. *Nat. Rev. Microbiol.* **8**:600-607. doi:10.1038/nrmicro2391

19. Jun S, Arnold A, Ha BY (2007) Confined space and effective interactions of multiple self-avoiding chains. *Phys. Rev. Lett.* **98**:128303. doi:10.1103/PhysRevLett.98.128303.

20. Jung Y, Ha BY (2010) Overlapping two self-avoiding polymers in a closed cylindrical pore: implications for chromosome segregation in a bacterial cell. *Phys. Rev. E* **82**:051926. doi:10.1103/PhysRevE.82.051926.

21. Liu Y, Chakraborty B (2012) Segregation of polymers in confined spaces. *Phys. Biol.* **9**:066005. doi:10.1088/1478-3975/9/6/066005.

22. Dorier J, Stasiak A (2013) Modelling of crowded polymers elucidate effects of double-strand breaks in topological domains of bacterial chromosomes. *Nucleic Acids Res.* **41**:6808-6815. doi:10.1093/nar/gkt480.

23. Shin J, Cherstvy AG, Metzler R (2014) Mixing and segregation of ring polymers: spatial confinement and molecular crowding effects. *New Journal of Physics* **16**:053047. doi:10.1088/1367-2630/16/5/053047.

24. Chen YH, Yu WC, Wang JJ, Luo KF (2015) Polymer segregation under confinement: Influences of macromolecular crowding and the interaction between the polymer and crowders. *J. Chem. Phys.* **143**:134904. doi:10.1063/1.4932370.

25. Männik J, Castillo D, Yang D, Siopsis G, Männik J (2016) The role of MatP, ZapA, and ZapB in chromosomal organization and dynamics in *Escherichia coli*. *Nucleic Acids Res.* **44**:1216-1226. doi:10.1093/nar/gkv1484.

26. Polson JM, Kerry DRM (2018) Segregation of polymers under cylindrical confinement: effects of polymer topology and crowding. *Soft Matter* **14**:6360-6373. doi:10.1039/c8sm01062e.

27. Männik J, Bailey MW (2015) Spatial coordination between chromosomes and cell division proteins in *Escherichia coli*. *Frontiers in Microbiology* **6**:306. doi:10.3389/fmicb.2015.00306.

28. Stouf M, Meile J-C, Cornet F (2013) FtsK actively segregates sister chromosomes in *Escherichia coli*. *Proc. Natl. Acad. Sci. U. S. A.* **110**:11157-11162. doi:10.1073/pnas.1304080110.

29. Galli E, Midonet C, Paly E, Barre FX (2017) Fast growth conditions uncouple the final stages of chromosome segregation and cell division in *Escherichia coli*. *PLoS Genet.* **13**:1006702. doi:10.1371/journal.pgen.1006702.

30. Barre FX, Aroyo M, Colloms SD, Helfrich A, Cornet F, Sherratt DJ (2000) FtsK functions in the processing of a Holliday junction intermediate during bacterial chromosome segregation. *Genes Dev.* **14**:2976-2988. doi:10.1101/gad.188700.

31. Männik J, Bailey MW, O'Neill JC, Männik J (2017) Kinetics of large-scale chromosomal movement during asymmetric cell division in *Escherichia coli*. *PLoS Genet.* **13**:e1006638. doi:10.1371/journal.pgen.1006638.

32. Wu FB, Swain P, Kuijpers L, Zheng X, Felter K, Guurink M, . . . Dekker C (2019) Cell boundary confinement sets the size and position of the *E. coli* chromosome. *Curr. Biol.* **29**:2131-2144. doi:10.1016/j.cub.2019.05.015.

33. Miangolarra AM, Li SHJ, Joanny JF, Wingreen NS, Castellana M (2021) Steric interactions and out-of-equilibrium processes control the internal organization of bacteria. *Proc. Natl. Acad. Sci. U. S. A.* **118**: e2106014118. doi:10.1073/pnas.2106014118.

34. Sanamrad A, Persson F, Lundius EG, Fange D, Gynna AH, Elf J (2014) Single-particle tracking reveals that free ribosomal subunits are not excluded from the *Escherichia coli* nucleoid. *Proc. Natl. Acad. Sci. U. S. A.* **111**:11413-11418. doi:10.1073/pnas.1411558111.

35. Mohapatra S, Weisshaar JC (2018) Functional mapping of the *E. coli* translational machinery using single-molecule tracking. *Mol. Microbiol.* **110**:262-282. doi:10.1111/mmi.14103.

36. Youngren B, Nielsen HJ, Jun S, Austin S (2014) The multifork *Escherichia coli* chromosome is a self-duplicating and self-segregating thermodynamic ring polymer. *Genes Dev.* **28**:71-84. doi:10.1101/gad.231050.113.

37. Makela J, Uphoff S, Sherratt DJ (2021) Nonrandom segregation of sister chromosomes by *Escherichia coli* MukBEF. *Proc. Natl. Acad. Sci. U. S. A.* **118**:e2022078118. doi:10.1073/pnas.2022078118.

38. Papagiannakis A, Yu Q, Govers SK, Lin W-H, Wingreen NS, Jacobs-Wagner C (2025) DNA/polysome phase separation and cell width confinement couple nucleoid segregation to cell growth in *Escherichia coli*. doi:10.7554/elife.104276.1.

39. Sadhir I, Murray SM (2023) Mid-cell migration of the chromosomal terminus is coupled to origin segregation in Escherichia coli. *Nat Commun* **14**:7489. doi:10.1038/s41467-023-43351-7.

40. Reyes-Lamothe R, Sherratt DJ, Leake MC (2010) Stoichiometry and architecture of active DNA replication machinery in *Escherichia coli*. *Science* **328**:498-501.

41. Wery M, Woldringh CL, Rouviere-Yaniv J (2001) HU-GFP and DAPI co-localize on the *Escherichia coli* nucleoid. *Biochimie* **83**:193-200.

42. Tiruvadi-Krishnan S, Männik J, Kar P, Lin J, Amir A, Männik J (2022) Coupling between DNA replication, segregation, and the onset of constriction in *Escherichia coli*. *Cell Reports* **38**:110539. doi:10.1016/j.celrep.2022.110539.

43. Wang P, Robert L, Pelletier J, Dang WL, Taddei F, Wright A, Jun S (2010) Robust growth of *Escherichia coli*. *Curr. Biol.* **20**:1099-1103.

44. Yang D, Jennings AD, Borrego E, Retterer ST, Männik J (2018) Analysis of factors limiting bacterial growth in PDMS mother machine devices. *Frontiers in Microbiology* **9**:871. doi:10.3389/fmicb.2018.00871.

45. Yang D, Männik J, Retterer ST, Männik J (2020) The effects of polydisperse crowders on the compaction of the *Escherichia coli* nucleoid. *Mol Microbiol* **113**:1022-1037. doi:10.1111/mmi.14467.

46. Gray WT, Govers SK, Xiang YJ, Parry BR, Campos M, Kim S, Jacobs-Wagner C (2019) Nucleoid size scaling and intracellular organization of translation across bacteria. *Cell* **177**:1632-1648. doi:10.1016/j.cell.2019.05.017.

47. Bakshi S, Choi H, Mondal J, Weisshaar JC (2014) Time-dependent effects of transcription- and translation-halting drugs on the spatial distributions of the *Escherichia coli* chromosome and ribosomes. *Mol. Microbiol.* **94**:871-887. doi:10.1111/mmi.12805.

48. Bakshi S, Siryaporn A, Goulian M, Weisshaar JC (2012) Superresolution imaging of ribosomes and RNA polymerase in live *Escherichia coli* cells. *Mol. Microbiol.* **85**:21-38. doi:10.1111/j.1365-2958.2012.08081.x.

49. Mondal J, Bratton BP, Li YJ, Yethiraj A, Weisshaar JC (2011) Entropy-based mechanism of ribosome-nucleoid segregation in *E. coli* cells. *Biophys. J.* **100**:2605-2613. doi:10.1016/j.bpj.2011.04.030.

50. Chang MH, Lavrentovich MO, Mannik J (2024) Differentiating the roles of proteins and polysomes in nucleoid size homeostasis in *Escherichia coli*. *Biophys J* **123**:1. doi:10.1016/j.bpj.2023.11.010.

51. Chai Q, Singh B, Peisker K, Metzendorf N, Ge XL, Dasgupta S, Sanyal S (2014) Organization of ribosomes and nucleoids in *Escherichia coli* cells during growth and in quiescence. *Journal of Biological Chemistry* **289**:11342-11352. doi:10.1074/jbc.M114.557348.

52. Cunha S, Woldringh CL, Odijk T (2001) Polymer-mediated compaction and internal dynamics of isolated *Escherichia coli* nucleoids. *J. Struct. Biol.* **136**:53-66. doi:10.1006/jsbi.2001.4420.

53. Gelber I, Aranovich A, Feingold M, Fishov I (2021) Stochastic nucleoid segregation dynamics as a source of the phenotypic variability in *E. coli*. *Biophys. J.* **120**:5107-5123. doi:10.1016/j.bpj.2021.10.004.

54. Yu XC, Tran AH, Sun Q, Margolin W (1998) Localization of cell division protein FtsK to the Escherichia coli septum and identification of a potential N-terminal targeting domain. *J Bacteriol* **180**:1296-1304. doi:10.1128/JB.180.5.1296-1304.1998.

55. Wlodarski M, Raciti B, Kotar J, Cosentino Lagomarsino M, Fraser GM, Cicuta P (2017) Both genome and cytosol dynamics change in *E. coli* challenged with sublethal rifampicin. *Phys Biol* **14**:015005. doi:10.1088/1478-3975/aa5b71.

56. Bernstein JA, Khodursky AB, Lin PH, Lin-Chao S, Cohen SN (2002) Global analysis of mRNA decay and abundance in Escherichia coli at single-gene resolution using two-color fluorescent DNA microarrays. *Proc. Natl. Acad. Sci. U. S. A.* **99**:9697-9702. doi:10.1073/pnas.112318199.

57. Van Helvoort J, Huls PG, Vischer NOE, Woldringh CL (1998) Fused nucleoids resegregate faster than cell elongation in *Escherichia coli* pbpB(Ts) filaments after release from chloramphenicol inhibition. *Microbiology* **144**:1309-1317. doi:10.1099/00221287-144-5-1309.

58. Xiang YJ, Surovtsev IV, Chang YJ, Govers SK, Parry BR, Liu J, Jacobs-Wagner C (2021) Interconnecting solvent quality, transcription, and chromosome folding in *Escherichia coli*. *Cell* **184**:3626-3642. doi:10.1016/j.cell.2021.05.037.

59. Woldringh CL (2002) The role of co-transcriptional translation and protein translocation (transertion) in bacterial chromosome segregation. *Mol. Microbiol.* **45**:17-29. doi:10.1046/j.1365-2958.2002.02993.x.

60. Demarre G, Galli E, Barre FX (2013) The FtsK family of DNA pumps. *Adv Exp Med Biol* **767**:245-262. doi:10.1007/978-1-4614-5037-5_12.

61. Besprozvannaya M, Burton BM (2014) Do the same traffic rules apply? Directional chromosome segregation by SpoIIIE and FtsK. *Mol. Microbiol.* **93**:599-608. doi:10.1111/mmi.12708.

62. Kennedy SP, Chevalier F, Barre F-X (2008) Delayed activation of Xer recombination at *dif* by FtsK during septum assembly in *Escherichia coli*. *Mol. Microbiol.* **68**:1018-1028. doi:10.1111/j.1365-2958.2008.06212.x.

63. Micali G, Grilli J, Marchi J, Osella M, Lagomarsino MC (2018) Dissecting the control mechanisms for DNA replication and cell division in *E. coli*. *Cell Reports* **25**:761-771. doi:10.1016/j.celrep.2018.09.061.

64.   Birnie A, Dekker C (2021) Genome-in-a-box: building a chromosome from the bottom up. *ACS Nano* **15**:111-124. doi:10.1021/acsnano.0c07397.

65.   Gibson DG, Glass JI, Lartigue C, Noskov VN, Chuang RY, Algire MA, . . . Venter JC (2010) Creation of a bacterial cell controlled by a chemically synthesized genome. *Science* **329**:52.

Figures

Fig. 1. Segregation of nucleoids as a function of DNA replication time (age). (A) The population average

DNA density distribution along the long axis of the cell as a function of replication cycle age. Time zero

corresponds to replication initiation and one to termination (also indicated by a vertical dashed line). HupA-

mCherry labeling is used to infer DNA density distribution. Red corresponds to high- and blue to low-

intensity values. The dashed horizontal lines indicate a midcell region. (B) The same for the average

replisome density distribution. Ypet-DnaN labeling is used to infer this distribution. (C) Left: definition of

parameters characterizing the dimensions of nucleoids: $L_{nuc,sum}$ and $L_{nuc,ext}$. Right: change of cell

length L_{cell}, $L_{nuc,sum}$ and $L_{nuc,ext}$ as a function of replication cycle time. The dashed vertical line marks

replication termination. (D) Left: schematics showing the region where HupA-mCherry and Ypet-DnaN

midcell amount is measured (indicated also by the horizontal dashed lines in panels A and B). The midcell

amount is based on the integrated intensity of fluorescent labels in this band. Right: the normalized midcell

amount for nucleoid (red) and replisome (green) as a function of replication age. The curves for individual

cells are normalized by the maximum value during the whole measurement period and then averaged over

the cell population. (E) Left: schematics showing the DNA density distribution for a single cell at a single

time point and how the depth of the local minimum, A_{min}, is defined. Right: A_{min} as a replication cycle time.

Error bars in panels C-E are s.e.m. N=438

Figure 2. Cell-cycle dependent changes in DNA and ribosome density distributions and their comparison
to model predictions. (A) Population averaged kymograph of DNA density distribution (HupA-mYpet) along
the long axis of the cell as a function of cell age. Zero corresponds to cell birth and one to division. Red
corresponds to high- and blue to low-intensity values. The dashed horizontal lines indicate a midcell region.
(B) The same for ribosome density (RpsB-mCherry) distribution. (C) Phase contrast kymograph for the
same cell population. (D) The normalized population-averaged density of nucleoid (green) and ribosomes

(red) along the long axis of the cell at different cell ages. The curves are normalized by their peak values.

(E) Pearson correlation coefficient, R, between nucleoid and ribosome densities as a function of cell age.

The coefficient is averaged over the cell population at each cell age point. (F) The integrated intensity within

the 0.75 μm wide band around the cell middle is shown for the nucleoid (green), ribosome (red) and phase

contrast signals (black). The curves from individual cells are normalized by their peak values and then

averaged over the cell population. The left vertical dashed line corresponds to cell age of 0.40 and the right

one to 0.68. (G) The relative excess fraction of ribosomes in the midcell band as a function of cell cycle

time. The curve is derived from a fitting of the ribosome signal (see Methods). N=486 for panels A-G. The

error bars are s.e.m. for panels E-G. (H) The normalized DNA and ribosome densities along the long axis

of the cell at different cell ages, as predicted by the coupled DNA and ribosome dynamics model. The

curves are normalized by their peak values. Note that the DNA (left) and ribosome (right) signals have

different scales, and the cell ages are comparable but not identical as those in panel D. (I) The same curves

that are recalculated, assuming a concentric shell of ribosomes around the nucleoid (for details, see the

Concentric Shell Model in Methods). The shell's inner radius is $0.5 \times R_{cell}$.

Figure 3. Segregation of nucleoids with and without closing constriction. (A) A population-averaged kymograph of DNA density along the long axis of the cell as a function of absolute (unscaled) time from the formation of local DNA density minimum, T_{min}, for cephalexin-treated non-constricting cells. The vertical axis shows normalized cell length, which remains unchanged throughout the experiment. Note that cells filament in this measurement. (B) The same for ribosome density distribution in this cell population. (C) The

average number of ribosomes at midcell as a function of time from T_{min} for cephalexin-treated (red) and

untreated cells (violet). The curves are normalized so that the amounts are one at time T_{min}. (D) The

normalized DNA amount at midcell is presented the same way as in the previous panel. (E) The average

distance between nucleoid centers (peak distance) as a function of time from T_{min} for cephalexin-treated

(red) and untreated cells (violet). The distances are normalized at T_{min}. The schematic on the top shows

how the peak distance is defined. (F) Kymograph of nucleoid density from the coupled DNA and ribosome

dynamics model presented the same way as the experimental data in panel A. (G) Kymograph of ribosome

density from the model. (H) The peak distance as a function of time relative to T_{min} from cephalexin-treated

cells (red) and from the model (black). (I) Normalized midcell DNA amount for cephalexin-treated cells (red)

and model (black). The linear fits to both curves are shown by dashed lines. The model is fitted for $\text{time} -$

$T_{min} > 5 \text{ min}$. (J) The relative excess fraction of ribosomes at midcell for cephalexin-treated cells (red) and

model (black). The linear fits for both curves are shown as dashed blue lines. The fit coefficients for panels

I and J are listed in Table S3. (K) Left: Schematic showing how the width of the midcell accumulation of

ribosomes, W, is determined based on ribosome density profile (blue) and it's fit by generalized Gaussian

curves (red). Right: the time evolution of W for cephalexin-treated cells (red) and the model (black). All error

bars are s.e.m. N=235.

Figure 4. The effects from the disruption of active mRNA-polysome dynamics with rifampicin (Rif) on nucleoid and ribosome distributions. (A, D) The population-averaged kymographs of DNA density. The times are relative to the start of antibiotic treatment, T_{ab}, with Rif and Cef (cephalexin), and with only Cef, respectively. All analyzed cells in this figure are non-constricting. (B, E) The same for the average ribosome density. (C, F) The density distributions of DNA (green) and ribosome (red) along the long axis of the cell at different time points. In all panels, the curves are normalized relative to their maximum value at T_{ab}. (G) The depth of the local DNA density minimum at midcell as a function of time relative to the start of antibiotic treatment with Rif + Cef (blue) and Cef only (pink). (H) The average cell length (green) and the nucleoid peak distance (black) as a function of time from the start of the treatment for Rif + Cef treated cells. The curves from individual cells are normalized to one at T_{ab}. Error bars in panels G and H are s.e.m. (I) Schematic summarizing the expansion and movement of nucleoids and ribosomes during Rif treatment.

553 The top corresponds to a cell just before and bottom after Rif (+Cef) treatment. $N = 786$ for Rif+Cef and

554 $N = 310$ for only Cef treated cells.

Figure 5. Comparing DNA and ribosome dynamics in Rif+Cef treated cells to the coupled DNA and ribosome dynamics model. Cells from two age groups are compared. (A-F) Data and model for cells that have formed a local DNA density minimum before the antibiotic treatment, $T_{min} < T_{ab}$ ($N = 92$). (A, B) The kymographs of DNA and ribosome density, respectively. The times are relative to the start of antibiotic

treatment. (C) The average DNA (green) and ribosome (red) distributions along the long axis of the cell at

different time points of experiment. (D, E) The DNA and ribosome density kymographs, respectively, from

the model corresponding to group of cells with $T_{min} < T_{ab}$. The cell length in the beginning of the treatment

is 2.87 μm. Note shorter times in the model compared to experimental data. (F) The average DNA (green)

and ribosome (red) distributions along the long axis of the cell at different time points from the model. (G-

L) Data and model for cells that formed a local DNA density minimum after the start of the antibiotic

treatment, $T_{min} > T_{ab}$ ($N = 138$). (G,H) The kymographs of DNA and ribosome density, respectively. (I) The

average DNA (green) and ribosome (red) distributions along the long axis of the cell at different time points

of the experiment. (J, K) The DNA and ribosome density kymographs, respectively, from the model

corresponding to group of cells with $T_{min} > T_{ab}$. The cell length in the beginning of the treatment is 2.24 μm.

Note shorter times in the model compared to experimental data. (F) The average DNA (green) and

ribosome (red) distributions along the long axis of the cell at different time points from the model.

Figure 6. Progression of different mechanisms involved in segregation and separation of daughter
chromosomes during *E. coli* cell cycle. The schematic summarizes the results from this work and from
previously published studies. C-period marks the replication period, and D-period marks the duration from
the termination of DNA replication to the final division of two daughter cells. The drawn schematic
corresponds to slow growth conditions where the replication initiates after cell birth.