Connectome of a human foveal retina

Yeon Jin Kim[1], Orin Packer[1], Thomas Macrina[2], Andreas Pollreisz[3], Christine A. Curcio[4], Kisuk

Lee[2], Nico Kemnitz[2], Dodam Ih[2], Tri Nguyen[2], Ran Lu[2], Sergiy Popovych[2], Akhilesh Halageri[2],

J. Alexander Bae[2], Joe Strout[2], Stephan Gerhard[5], Robert G. Smith[6], Paul R. Martin[7], Ulrike

Grünert[7], Dennis M. Dacey[1,8,*]

[1]Department of Neurobiology & Biophysics, University of Washington, Seattle, WA, 98195

[2]Zetta AI LLC, Sherrill, NY, USA

[3]Department of Ophthalmology, Medical University of Vienna, Vienna, Austria

[4]Department of Ophthalmology and Vision Sciences, University of Alabama at Birmingham,

Alabama 35294

[5]Aware LLC, Rappenstrasse 19, 8307 Effretikon, Switzerland

[6]Department of Neuroscience, University of Pennsylvania, Philadelphia, PA 19104

[7]The University of Sydney, Save Sight Institute and Discipline of Ophthalmology, Faculty of

Medicine and Health, Sydney, NSW, 2000, Australia

[8]Washington National Primate Research Center, Seattle, WA 98195

*Corresponding author: Dennis M. Dacey (dmd@uw.edu)

This file includes: [5059 / 5000 words]

Main Text (Main text pages: 19, Total pages: 62)

Figures 1 to 10

Supplementary Information

Abstract

The foveal retina is a primate specialization which presents a feasible site for obtaining a

complete connectome of a human CNS structure. In the fovea, cells and circuits are

miniaturized and compressed to densely sample the visual image at high resolution, initiating

form, color, and motion perception. Here we present a sample of human foveal retina analyzed

by deep learning-based segmentation of all cells and synaptic connections. Cells in the volume

are provisionally classified into 51 morphological types. We discovered synaptic pathways that

are absent in non-human primates that may play a role in human color vision. We also

demonstrate a biophysical model based on electrical synapses among short-, middle-, and

long-wavelength sensitive cone photoreceptors. Segmentation of the ganglion cells, the output

cells of the retina, suggests only 11 visual pathways, with 5 high-density pathways accounting

for over 95% of the foveal output to the brain; this number is substantially lower than the 40+

types observed in mouse retina. The connectomic resource presented here reveals distinct

features of a human neural system and highlights the potential for deep learning-based

computational methods to advance understanding of the human brain.

The vertebrate retina is an accessible central nervous system (CNS) outpost where the visual

process begins. It has long been appreciated for its great complexity[1] and for linking neural

structure to visual function[2, 3, 4]. The retina was an early focus of synaptic-level connectomic

reconstruction[5, 6, 7, 8, 9, 10], and more recently, retinal cell populations have been linked to their

molecular profiles[11, 12, 13, 14] and the genetics of human disease[13, 15]. Yet even for this

intensively studied neural system, we lack a full wiring diagram that could explain how

vision—one of our most fundamental senses—is initiated.

The recent achievement of a complete connectome of the brain of the fruit fly, *Drosophila*

melanogaster[16, 17], and application of connectomic methods to mm-scale volumes of mouse

and human neocortex[18, 19, 20, 21], have made the connectome of a larger mammalian brain a

conceivable goal[18, 19, 22, 23]. In the present study, we take advantage of the distinctive neural

architecture of the foveal retina to provide a first draft connectome of this critical locus in the

human CNS.

The fovea is a specialized central region in the retina of humans and other day-active primates

where neural circuits are spatially compressed to facilitate high-resolution form, color and

motion perception. At barely 1 mm wide, the fovea encodes only a few degrees of visual space,

but its representation is magnified 100-fold across the surface of primary visual cortex[24, 25, 26, 27,

28]. Accordingly, the fovea is essential for normal vision, and failure of foveal function is a

hallmark of devastating blinding retinal diseases.

From a signal-processing viewpoint, the fovea is an input-to-output sensory structure, where

light transduction by photoreceptors is processed by diverse neural cell types across two

narrow cell and synaptic layers. Here we provide a catalog of human foveal cells and their

synaptic assignments to distinct functional circuits. We use this dataset to directly address the

long-standing question of the number and identity of visual pathways that link the fovea to the

brain. We find that only five visual pathways account for over 95% of foveal output to the

human brain: a striking reduction compared to non-primate mammals[29, 30, 31] and non-human

primates[32, 33]. In addition, we observed unexpected differences in synaptic connections

compared to non-human primates and provide evidence that neural coding for trichromatic

color vision begins with circuitry that is uniquely human.

Results

Acquisition of the foveal volume. Our goal was to obtain complete retinal circuitry near the peak of cone density (peak of visual acuity)[34, 35]. This is a difficult task because (among other reasons) foveal cones feed inner retinal circuits via very long axons (Henle fibers, Fig. 1a-1c)[36]. Our data sample was therefore centered at 500 μm retinal eccentricity, where cone axon length averages 300-400 μm[37, 38]; this location corresponds to cones located ~100-200 μm from the center of the foveal pit (~0.5-1 deg). We cut 3028 vertical sections at 50 nm thickness through a depth of ~150 μm and imaged an area 180 μm tall x 180 μm wide at 5 nm resolution (voxel size, 5 x 5 x 50 nm^3), extending from the Henle fiber layer (outer retina) to the nerve fiber layer (inner retina) (Fig. 1d). Advanced computational methods (convolutional neural networks)[39, 40] were used to align the images, detect membrane boundaries, segment neurons, glia, blood vessels, conventional (inhibitory) and ribbon (excitatory) synapses, and predict synaptic relationships in the outer and inner synaptic layers (Fig. 1d-1k; see Methods for procedures). We named this volume HFseg1 (Human Fovea, segmented 1). All annotated cell types and the draft catalog of synaptic connections are available for exploration using search tools in an openly accessible form (https://neuromaps.app/datasets, see Supplementary Table 3) which utilizes the Neuroglancer browser and CAVE infrastructure. In addition, we created a second site, accessed via NeuroMaps, that facilitates proofreading of cells and synapses by the user community.

Cell populations

To derive and understand any connectome, the identity and layout of all cell types must first be clearly understood. Here, we took advantage of the detailed anatomical picture of foveal cell types[41, 42] that has been linked to transcriptomically defined cell clusters in human and non-human primate retina[12, 13, 43]. Moreover, for many cell populations the presence of orderly spatial arrays could be used to estimate relative density. We were therefore able to use the

original segmentation followed by proofreading of cell morphology (~40,000 edits to date) to

annotate and classify all cells and to assess the quality of synapse prediction (Fig. 1f-1g; see

Supplementary Table 1a, 3008 total cells with cell bodies within the volume; ~304,800 total

synapses, including those related to processes extending into or through the volume but not

linked to cell bodies).

In brief, HFseg1 contained 313 cone (pedicles) and 24 rod (spherules) synaptic terminals (rod:

cone = 0.07, Supplementary Table 1c, 1d). 17 cone pedicles were identified as short

wavelength-sensitive (S) cones (Fig. 2a)[44]. The remaining 296 cones comprised long (L) and

medium (M) wavelength-sensitive cones, here referred to collectively as LM cones.

Postsynaptic to cones, we annotated 268 horizontal cells and 912 bipolar cells (Supplementary

Table 1e, 1f). It was possible to distinguish the H1 and H2 horizontal cell types (Fig. 3) and

most of the classically recognized bipolar cell types (Fig. 4) by distinctive morphology and

connectivity. Some exceptions will be considered further below.

We annotated 293 amacrine cells, including 40 cells displaced to the ganglion cell layer (Fig. 5;

Supplementary Table 1g; Supplementary Fig. 8), first dividing cells by dendritic field diameter

into small (diameter < 100 μm; n = 189), and large-field (diameter >100 μm) groups (n = 104).

We divided these groups in turn into 19 types (6 small-field and 13 large-field; Supplementary

Table 1g) based on morphology, stratification, mosaic interactions and nuclear morphology

(Supplementary Figs. 3, 6, 8). Figure 5 provides a partial view of this dataset by highlighting

four cell types (see also Supplementary Table 3); two additional amacrine populations are

illustrated in Supplementary Fig. 8 and others can be viewed at the NeuroMaps site.

We annotated 599 ganglion cells (Fig. 6; Supplementary Table 1h). As expected, the

classically recognized midget GCs constitute the great majority (536 cells, ~90%)[13, 45]; these

cells were readily divided into outer-OFF stratifying (n = 280 cells) and inner-ON stratifying (n =

256) types (Fig. 6a, 6e). A second well-defined cell group, the parasol cells (26 cells, 13 inner-

ON cells and 13 outer-OFF cells, ~4% of total GCs), was also easily distinguished (Fig. 6b, 6f).

Small bistratified GCs[46], found to show blue-yellow color opponency in macaque retina[47, 48]

were distinguished by their broad dendritic ramification (12 cells, ~2% of total GCs; Fig. 6c, 6g)

and extensive synaptic input from blue cone bipolar cells (BB cells)[49]. Together, the midget,

parasol and small bistratified cells accounted for 95.8% of the ganglion cells in HFseg1

(Supplementary Table 1h). The remaining 25 ganglion cells showed relatively large dendritic

fields (Figs. 6d, 6h–6n) and were divided into 5 provisional groups comprising 6 types

(Supplementary Table 1h; Supplementary Notes) pending a more detailed analysis of their

synaptic connections, giving a total of 11 ganglion cell populations in this volume–this number

is very close to the 12 populations identified for central human retina by molecular profiling[13].

Finally, glial cells were readily distinguished from all neurons by their distinctive ultrastructure

(Supplementary Figs. 3a, 3b, 3c, 3f). The Müller cells (555 cells, Fig. 1f,1g; Supplementary

Table 1b) were distinguished as radial glia with cell bodies in the inner nuclear layer

(Supplementary Fig. 1a). A second, morphologically distinct population of glial cells in the GCL

was closely associated with superficial blood vessels (Supplementary Figs. 1b, 3b) and

identified as astrocytes[50] (32 cells, Supplementary Table 1b; Supplementary Fig. 1b). We also

identified 15 microglial cells (Supplementary Table 1b) with cell bodies in the IPL or GCL

(Supplementary Figs. 1a, 1b, 3c, 3b). The density of microglial cells was low, consistent with

previous measurements in the macaque monkey foveal retina[51, 52]. In the current volume, we

have not yet attempted to characterize blood vessel and associated pericytes, though these

cellular elements are also segmented and are available to annotate and further characterize.

More detailed information on cell type identification is provided in the supplementary

information (Supplementary Notes on cell type identification).

Synapse detection

To assess the accuracy of ribbon segmentation and synapse prediction (Fig. 1h-1k), we

compared deep-learning-predicted synaptic assignments to manual (ground truth)

assignments. We focused on the cone-to-midget bipolar cell synapse in the OPL and the

midget bipolar cell synapses to amacrine and ganglion cells in the IPL. For ground truth, we

manually annotated all midget pathway synaptic connections for three LM cones (cones 21, 22,

and 58; Supplementary Fig. 9; Supplementary Table 2a-2c).

Figure 7 shows the predicted outgoing synapses for a pair of midget bipolar cell axon

terminals. We calculated precision (true synapses / [true synapses + false positives]; the

accuracy of positive predictions) and sensitivity (true synapses / [true synapses + false

negatives]; the proportion of true synapses identified) for ribbons and their synapses. The

segmentation of ribbons showed high precision and sensitivity near 100% (precision: 93.5% ±

2.7%; sensitivity: 99.7% ± 0.8%, n = 6 midget bipolar cells, 370 total ribbons; Fig. 7b inset).

Postsynaptic partner assignment for these ribbons was also excellent (precision: 89.0% ±

5.4%; sensitivity: 93.7% ± 4.6%, n = 6 midget bipolar cells, 637 total synapses; Fig. 7b inset).

These values demonstrate the high accuracy of convolutional nets for predicting and

segmenting ribbon synapses.

In contrast to the ribbon synapses made in the IPL, ribbon synapses made by the cone pedicle

in the OPL are extremely complex[53, 54] (Fig. 1h, 1i). Postsynaptic processes can either

invaginate into the face of the pedicle or occupy a basal position at the cone pedicle up to ~800

nm from the sites of transmitter release[54, 55]. We mapped all cone-to-bipolar synapses at the

dendritic tips for a sample of six midget bipolar cells. We found high precision and sensitivity for

ribbon detection (n = 51 ribbons) (precision: 92.3% ± 6.7%; sensitivity: 100%, n = 3 LM cones,

Supplementary Fig. 2a-2c) and for the postsynaptic processes (n = 153) deriving from the

invaginating (IMB) and flat (FMB) midget bipolar cells (precision: 98.7% ± 1.2%; sensitivity:

99.7% ± 0.6%, n = 153 total synaptic connections; see Supplementary Fig. 2c right). Diffuse

bipolar (DB) cells, which make sparser connections with multiple cones, were also sampled (n

= 2, DB4 and DB2; Supplementary Fig. 2d-2f), yielding similar precision and sensitivity values.

For inhibitory synapses, we focused on the AII amacrine cell (Figs. 5a, 7c, 9; Supplementary

Table 1g) where glycinergic synapses to OFF bipolar cell axon terminals are well established[56,

57, 58, 59]. We manually mapped all the outgoing synapses made by two neighboring AII amacrine

cells and compared our ground truth data to the predicted synaptic assignments (Fig. 7c-7g).

The prediction captured well the synaptic output from the outer stratifying lobular appendages

of the AII cell to outer-OFF bipolar cells (Fig. 7d-7g). Vesicle clouds were detected with high

sensitivity (87.5% ± 0.7%, n = 2 AII cells; false negatives were rare; Fig. 7c inset), but relative

to ribbon detection, there were more false positives, thereby reducing precision (76.0% ± 8.5%,

n = 2 AII cells). Despite this limitation, the results show that our dataset predicts synaptic

connectivity with good sensitivity and precision. However, we implemented methods within the

Neuromaps interface to proofead all synapses for accuracy.

Having established the accuracy of our connectomic dataset, we examined the cell types and

their synaptic connections to identify features that are distinctive in human compared to non-

human primate. Below we highlight examples where differences are dramatic and warrant

further study.

Modeling the connectome: S-LM cone electrical coupling alters spectral tuning.

Cone pedicles contact each other by short extensions (telodendria) or by distinct pedicle-to-

pedicle appositions at perforations in the glial sheath surrounding cone synapses (Fig. 2b).

These contacts harbor gap junctions, i.e., electrical synapses that mediate a conductance

between cones, including L and M cones[60]. In non-human primates, L to M cone contacts are

abundant, but S to LM contacts are greatly reduced or absent[44, 61]. In the macaque, electrical

coupling was absent between LM and S cones[60], consistent with a lack of contacts between

these cone types[61, 62]. The crosstalk between L and M cones serves to reduce noise in the

light-evoked response with a small cost to spectral separation[63, 64, 65]. The lack of gap junctions

between LM and S cones has been interpreted as reflecting the need to keep S and LM cone

signals separate to build color-coding receptive fields at later stages in color processing[66, 67].

Unlike in the macaque, S-LM cone contacts have been reported in human retina[44, 68]. We

therefore mapped the full cone-cone connectome to determine whether S-LM contacts are a

consistent feature of the human fovea. Cone-cone contacts (n = 1141) were characterized by a

thick, darkly stained region defining the contact area (see Fig. 2c)[69]. We found abundant S-LM

cone contacts (n = 17 S cones; 5.25 ± 2.11 contacts/S-cone pedicle), indistinguishable from

LM-LM cone contacts (n = 296 LM cones; 6.89 ± 3.08 contacts/pedicle). We then used this

cone-cone connectome dataset to estimate the effect of S-LM coupling on spectral tuning in a

simulated cone-cone network, using known parameters for gap junction number and single

channel conductance values as follows. We mapped the surface area of the contacts among all

cones in the volume (Fig. 2a), using automated methods to generate the "contactome" between

cone pedicles. Detected contacts were inspected and filtered by size to remove very small and

likely spurious contacts. The surface area of each contact was calculated (n = 1141 contacts;

mean surface area 0.56 ± 0.31 μm^2). In general, contacts were large and obvious with a

relatively thick, darkly stained region defining the contact area (see Fig. 2c), reflecting the

presence of large adherens junctions at the contact site[69]. We calculated the conductance for

each contact based on previously measured densities for gap junctions and the conductance of

single gap junction channels (9 gap junctions/μm^2 at 15 pS/junction; see Methods for details).

We used a fixed ratio of L to M cones of ~1.7:1, randomly assigned[70], and incorporated the

cone-cone coupling given by our contactome (Fig. 8a). Each cone was modeled

compartmentally as a soma, Henle fiber-axon, and synaptic terminal (see model parameters in

Supplementary Table 4). Our modeled average coupling conductances were low (LM-LM cone

pairs, 133 pS, n = 25 cones; S-LM cone pairs, 166 pS, n = 15) by comparison with

conductances measured directly between LM cone pairs in the macaque (> 500 pS) or cone

pairs in non-primate mammalian retina (~220-320 pS)[65, 67]. Nevertheless, we found that even

with this conservative estimate of coupling conductance, the S-cones showed an increased

sensitivity to light pulses at longer wavelengths in the model (Fig. 8b). We measured the effect

on spectral sensitivity by simulating light responses from L, M, and S cones at 5 nm

wavelength intervals across the visible spectrum. The results demonstrate a broad secondary

elevation in sensitivity for S cones at longer wavelengths that peaks at ~550 nm (Fig. 8c-8d) in

the coupled cone network. These data provide evidence that S-LM cone coupling can directly

influence the spectral tuning of the voltage response of S cones and raise the question of how

this network feature affects downstream processing.

Trichromatic horizontal cells

A second difference in synaptic connectivity that may impact color vision occurs at the synapse

between cones and horizontal cells. The horizontal cell network in the outer retina generates

lateral negative feedback that underlies the antagonistic receptive field surround of cones,

bipolar cells, and ganglion cells[71]. In the macaque monkey retina, the horizontal cell-mediated

surround is a key element in the circuitry that initiates red-green color vision[72, 73, 74]. In the

peripheral retina of the macaque and marmoset, H1 cells make dense contacts with LM cones

but largely or completely avoid contact with S cones, whereas H2 cells show some selectivity

for S cones[75, 76, 77]. While foveal H2 cells in HFseg1 maintained a clear preference for

synapsing with S cones (Fig. 3b), we found that H1 cells also received a major input from S

cones, comparable to H2 cells (H1, 43% ± 16%; H2, 57% ± 16%, n = 10 S cones). By contrast,

H2 cells formed less than 2% of the horizontal cell contacts with LM cones (H1, 98.1% ± 0.1%;

H2, 1.9% ± 0.1%, n = 2 LM cones) (also Supplementary Fig. 4). Consequently, human foveal

H1 cells are predicted to show a trichromatic receptive field with a small S cone contribution

(Fig. 3; Supplementary Fig. 4). Taken together, the S-LM cone coupling and the combined S

and LM cone connectivity of H1 horizontal cells provide new connectomic evidence for

differences between human and non-human primate circuits that initiate color vision[44].

Unexpected cell types and circuits in HFseg1

We also found differences in both bipolar and amacrine cell circuits compared to expectations

from the macaque monkey, whereby some bipolar cell types were either greatly reduced or

absent. For example, DB-OFF bipolar cells can be divided into 4 morphological types (DB 1, 2,

3a and 3b) in macaque monkey[78, 79] but we were only able to distinguish three populations by

mosaic arrangement and stratification in the IPL that we call DB1, 2 and 3 (see Supplementary

Notes). Bipolar cell morphologies not previously described in primates were also present. One

bipolar group showed amacrine-like morphology (outer-x and inner-x, Fig. 4a, 4b) but

contained abundant synaptic ribbons (20 cells, Supplementary Table 1f; Supplementary Fig. 6).

Similar cells were discovered recently in mouse retina[11, 80] and likely correspond to the OFF-x

bipolar cells identified by transcriptomics of human fovea[13]. While most of these cells stratified

in the outer half of the IPL (outer-x cells, n = 15, Supplementary Fig. 6a-6b), five cells stratified

in the inner half of the IPL (inner-x cells, Supplementary Fig. 6c-6g). Another previously

undescribed bipolar type showed the characteristic DB cell connection to multiple cones but

extended axonal branches in both outer and inner IPL where it provided synaptic output to

outer-OFF and inner-ON ganglion cell types (see also Supplementary Fig. 7c, 7e). We refer to

this cell group as DBbroad (18 cells, Fig. 4a; Supplementary Fig. 7; Supplementary Table 1f).

The HFseg1 volume offers a unique opportunity to further understand these novel bipolar cell

populations.

For the amacrine cells, the AII type (AC2) was present at the highest density of any amacrine

cell population (26%, 75 cells, Figs. 5a, 5b, 7c, 9; Supplementary Table 1g). This cell type was

recognized by its well-established morphology across species[56], including humans[41, 57, 81]. A

major function of AII cells is to transfer signals from rod bipolar to cone bipolar cells, serving as

conduits for rod signals to ganglion cells at scotopic light levels[82]. The presence of AII

amacrine cells at such a high density in the fovea, where rods are nearly absent, suggests,

however, a functional switch to a primary role in photopic spatial vision. Indeed, we observed

that AII cells are postsynaptic to all cone bipolar types, including IMB and FMB cells, across

the full extent of the dendritic tree (Fig. 9). These data support the general view that AII cells

"have a day job"[57] to provide an inhibitory pathway from ON to OFF bipolar cells under photopic

conditions[82, 83, 84] while at the same time suggesting specialized synaptic alterations for the AII

cells in the human fovea in relation to the midget circuit.

Visual pathway reduction: the vertical excitatory connectome

Despite its fundamental importance, the number of visual pathways linking the human fovea to

the brain is still unknown. Our classification of 11 ganglion cell types suggests that several

ganglion cell types observed in the monkey and human retinal periphery are absent from the

human fovea. This result also stands in sharp contrast to the retina of the mouse, where ~40

GC types have been distinguished independent of retinal location[29, 30, 31]. Our classification,

however, did not rule out the possibility that some ganglion cell types with somata outside

HFseg1 have dendritic trees which penetrate the volume and contribute substantially to the

synaptic connections of the inner retina. We therefore took a second approach to identifying

the visual pathways in HFseg1 by mapping what we called the "vertical excitatory

connectome".

We reasoned, as others have[85], that to generate parallel pathways to the brain, each cone

photoreceptor must be locally presynaptic to each cone bipolar type (For the S cones, there is

uncertainty regarding this point, due to their more specialized role in color vision[86, 87]). At the

next synaptic step, the combined output of all bipolar cell types should be distributed to all

ganglion cell types. In other words, in principle, *all visual pathways should be synaptically*

linked to the output of any given cone.

To test this hypothesis, we annotated the ribbons within five cones (three LM and two S) and

their postsynaptic bipolar cells. For each bipolar cell, all synaptic ribbons and their postsynaptic

GC partners were then identified (Fig. 10a-10b). Finally, we determined the relative synaptic

density from a given cone via its postsynaptic bipolar cells to each identified GC type (see

details in Supplementary Figs. 9 and 10; Supplementary Table 2).

We found that bipolar cells that were postsynaptic to the three LM cones provided their output

overwhelmingly to midget and parasol ganglion cells (Supplementary Fig. 9), accounting for

92.1% of the total bipolar synaptic output to ganglion cells (Fig. 10c, 10d; Supplementary Fig.

9; Supplementary Table 2a-2c). The remaining ~7.9% of bipolar synapses were directed to the

large-field ganglion cell types (LFGC1-LFGC5, Fig. 10d, 10h, and 10i-10n; Supplementary Fig.

9; Supplementary Table 2a-2c). Output to dendritic fragments of unidentified ganglion cells was

rare, accounting for only ~0.5% of the total bipolar output to ganglion cells (0.53 ± 0.76%, n = 3

LM cones, Supplementary Table 2a-2c). These data support the conclusion that, with the

possible exception of extremely large-field and low-density melanopsin-expressing ganglion

cells, we have accounted for all ganglion cell types present in our foveal sample.

Discussion

A human foveal connectome. Our aim was to provide a draft nanometer-scale connectome in the human CNS. Compared to ongoing studies of a larger volume of human neocortex[18], our foveal volume has the advantage that foveal circuitry is miniaturized and functionally complete – from receptors to ganglion cell outputs. Moreover, all cell types in the HFseg1 volume are identified and can be related to anatomical, physiological, and transcriptomic studies of human and non-human primate retina (Supplementary Table 1).

Cell type profiling. Comparable datasets to HFseg1 exist for primate central retina where cell types were clustered by gene expression profiles[12, 13, 15, 43, 88]. The HFseg1 volume permits a comparison of the two approaches and suggests generally good correspondence, though with some unexpected differences. For example, the ratio of H1 to H2 horizontal cell types identified by transcriptomic analysis was shown in the macaque and human central retina to be ~3:1 and 7:1, respectively. By contrast, the H1 to H2 ratio in HFseg1 is much higher (22:1), with H1 cells comprising 96% of the horizontal cells at this foveal location. Thus, the trend toward H1 cell dominance in the transcriptomic dataset is evident but more extreme in HFseg1. A similar trend holds for other clearly identified types: for example, midget ganglion cells make up ~90% of the ganglion cell population in our sample and 86% in the transcriptomic dataset; AII amacrine cells make up 26% of the amacrine cells but only 18% of the amacrines identified by transcriptomic analysis. One explanation for these differences may be that our smaller sample represents the cone-enriched foveal center, whereas the larger samples used for transcriptomics encompassed a much larger area (~1.5 mm, central 5 degrees). Cell type-specific density changes are extremely steep across this region, so that larger spatial samples would average out such changes.

Certain cell populations previously recognized were not evident in HFseg1. For example, the

recent functional, anatomical, and molecular distinction made between DB3a and DB3b in the

macaque monkey[79, 89] and human retina[13, 38] was not apparent in HFseg1. Further, we found

evidence for only 19 amacrine cell types, whereas molecular profiling in human retina

(including the retinal periphery) has proposed 73 amacrine cell types[15]. By comparison, 63

amacrine clusters have been identified in mouse retina[90, 91]. These differences suggest that the

relationship between cells defined by gene expression and anatomical types may not always

be straightforward. A further implication is that the extreme increase in density of the midget

circuit, required for high acuity and color vision, gives rise to an attendant density increase in

other foveal cell types tightly linked to the midget pathway (e.g., H1 horizontal cells).

The vertical excitatory connectome. In the non-human primates, fewer than 20 ganglion cell

types have been defined, with transcriptomic and anatomical identification in close

agreement[12, 32, 33]. In the human fovea, only 12 pathways have been suggested by

transcriptomic analysis[13, 15]. It was nevertheless striking that in the HFseg1 dataset

(Supplementary Table 1h), ~96% of the ganglion cells comprised just five anatomically distinct

pathways: the ON and OFF midget and parasol cells, and the small bistratified cells. When we

manually traced the vertical connectome from a single cone to its postsynaptic bipolar and

ganglion cells, we found that ~92% of synaptic output targeted these same five pathways (Fig.

10d), confirming their dominance. Thus, a fundamental, defining feature of foveal synaptic

organization may be a reduction in the number of visual pathways compared to the peripheral

retina.

Over 40 distinct ganglion cell populations have been recognized in the mouse retina, with

molecular profiling, physiology, and anatomy in agreement[29, 30, 31]. What explains this dramatic

reduction in ganglion cell types from the mouse retina to the primate fovea? The difference

suggests that humans and mice have evolved fundamentally different solutions to analyzing

the visual scene. Great emphasis has been placed on the behaviorally relevant feature-

detection properties of diverse mouse ganglion cell types[92, 93, 94]. By contrast, in primates,

considerable emphasis has been placed on the dual and more general role of the midget

pathway in both form and color perception[74, 95, 96].

Recent molecular orthotype analysis shows that the midget and parasol ganglion cell types

are likely homologous to sustained and transient alpha cell types, respectively, in the mouse[14].

What differs dramatically, however, is the extremely large dendritic tree and consequent low

spatial density (~2% of the total ganglion cells) of the mouse ortholog of the midget ganglion

cell. Thus, the mouse retina sacrifices spatial resolution[97], opting instead for an expansion of

feature detectors that rely on trigger features linked to motor output—with mice employing

feature-detection strategies analogous to those described in classical studies of amphibian

vision[98]. In primates, the midget circuit represents an evolutionary extreme where the spatial

resolution afforded by tightly packed photoreceptors is preserved through private-line

connectivity. This high-resolution sampling is behaviorally exploited through active vision

strategies. Humans not only explore a visual scene by making saccadic eye movements but

also make small fixational eye movements (microsaccades) to extract information at peak

resolution[99]. These microsaccades are dynamically modulated during highly precise

manipulation (e.g., threading a needle)[99, 100, 101]. Thus, the incessantly mobile fovea and

related retinal circuits are designed for inspection, manipulation, interpretation, and

understanding of an object—the converse of a feature detector.

The foveal connectome and human color vision.

S cone input to H1 cells. Most psychophysically based models of human color vision assume

the presence of two parallel color-opponent channels: red vs green and blue vs yellow[109]. The

red-green pathway shows opposing L vs M cone interaction and has been associated with the

circuitry that feeds midget ganglion cells[102]. However, the human psychophysically defined red-

green pathway includes a small S cone input[103]. The locus of this critical signal pathway

remains unclear, with recent data from the macaque monkey implicating both the retina[110] and

primary visual cortex[111]. The HFseg1 connectome shows that H1 horizontal cells, a major

source of the surround in the midget circuit[51], likely contribute antagonistic signals from all

three cone types (L, M, and S) to the midget pathway. In this scenario, the human foveal

midget circuit would derive its center response from either an L or M cone, while the surround

would provide cone opponency by non-selectively contacting all three cone types[73, 104, 105], with

the S cones making a small contribution to the chromatic receptive field, creating L+S vs M or

M+S vs L opponency.

S-LM cone coupling. As a first step in deriving the foveal connectome, we determined the

"contactome" among the tightly packed cone synaptic pedicles. Contacts among the cone

types harbor gap junctions and thus form electrical synaptic connections between cones[65, 69].

We confirmed and extended previous observations suggesting that, in contrast to non-human

primates and non-primate mammals, human foveal S cones show extensive telodendritic

contacts with neighboring LM cones[44, 68], and these contacts are identical in form to LM-LM

cone contacts.

A network model of this connectivity showed that, as a previous model suggested[66], that mixing

S and LM cone signals can alter the action spectrum of the S cone light response, with

increased sensitivity at longer wavelengths. This result raises a functional puzzle: the prevailing

view is that electrical coupling serves to increase signal relative to noise by reducing the

uncorrelated voltage difference (i.e., the noise component) among neighboring cones and

thereby increase contrast sensitivity[66]. However, any mixing of S and LM cone signals would

detract from the separation of cone signals required for color processing, in turn raising the

question of why such coupling occurs in human foveal retina, whereas it is absent in both the

macaque and marmoset retina[44].

The answer may be related, at least in part, to the connectivity of H1 horizontal cells with S

cones noted above. Coupling among the S, L, and M cones prior to this connection may serve

to preserve the smaller S cone signal within the H1 cell network. This in turn may be critical for

the unique requirements of human color vision whereby a small S cone signal contributes

critically to the "red-green" chromatic channel defined psychophysically[103]. This hypothesis

could be tested by adding the horizontal cell connectome to the current cone-cone network and

testing the effects of noise and coupling on the spectral tuning of the H1 cell response and

ultimately its contribution to cone opponency modeled in the midget ganglion cell.

Acknowledgements

This work was supported by NIH grants EY-028282 and RF1-MH129260 to D.M.D. and NIH grant P51 OD010425 to the Washington National Primate Research Center and EY01730 to the Vision Research Core at the University of Washington. We thank Sharm Knecht for tissue preparation and managing EM data acquisition.

Author contributions

Y.J.K. and D.M.D. conceived the project. Y.J.K. and D.M.D. generated the ground truth for segmentation and synapses, performed proofreading and created annotations. R.G.S constructed and ran the computational models. Y.J.K. and D.M.D. analyzed the data and generated figures. Y.J.K., D.M.D., P.R.M., and U.G., wrote the paper and consulted on cell type annotations and figure development. C.A.C. and R.G.S., edited the paper. A.P. acquired human tissue. D.M.D., Y.J.K., C.A.C. and A.P. acquired funding. Zetta AI performed EM data alignment, segmentation, synapse detection and cone-cone contactome detection and ingested all data into the CAVE infrastructure for proofreading. S.G. developed 3D visualizations and analysis tools at NeuroMaps.app.

Declaration of interests

K.L., N.K., D.I., T.N., R.L., S.P., A.H., J.A.B., J.S. and T.M. declare financial interests in Zetta AI. S.G. is owner of Aware LLC and developed the NeuroMaps.app. The remaining authors declare no competing interests.

Figures and figure legends (10 Figures)

Fig. 1. The HFseg1 volume. **a.** The fovea forms a pit in the retina ~3 mm temporal to the

optic disc (within the inset rectangle). **b**. Schematic foveal pit showing cone (red) to bipolar

cell (blue) to ganglion cell (green) circuit. ONL, outer nuclear layer; HFL, Henle fiber layer;

INL, inner nuclear layer; GCL, ganglion cell layer. **c**. Trimmed retinal block face and ROI

(yellow box) in temporal retina 400-500 μm from the foveal center. **d**. Single section from the

imaged volume (cropped laterally) to illustrate the retinal layers; HFL; OPL, outer plexiform

layer; INL; IPL, inner plexiform layer; GCL and NFL, nerve fiber layer. **e**. Volume segmented

into membrane-bound objects. **f**. Examples of segmented cells: photoreceptor axon terminal

(PC, white), horizontal cell (HC, orange), ON and OFF bipolar cells (BC, light and dark blue),

Müller glia cell (MC, yellow), amacrine cells (AC, green), ON and OFF ganglion cells (GC,

dark and light violet); astrocytes and microglia not shown here. **g**. Numbers and cell soma

locations of the cell types shown in **f**. **h.** Three segmented ribbons (varied colors) at synaptic

triads (red arrowheads) in a single cone pedicle. **i.** Synaptic assignments associated with each

ribbon (yellow dots and lines). An IMB (inner-ON midget bipolar; violet, white open arrowhead)

cell makes an invaginating contact and two FMB cells (outer-OFF midget bipolar; light and

dark blue, white open arrowheads) make a basal contact with the cone pedicle. **j.** Examples of

ribbon and synapse prediction in the inner plexiform layer (IPL) for an FMB cell axon terminal

at two locations (left and right panels). Segmented ribbons (varied colors, red arrowheads)

and synaptic predictions (blue dots, lines) are shown. **k.** Inhibitory vesicle cloud segmentation

(varied colors) and synaptic prediction (red dots and lines) in the IPL between an amacrine

cell and an FMB axon terminal are shown. Insets in **j** and **k** show zoomed-view of synapses

indicated by white boxes.

Fig. 2. Photoreceptors. a. Array of photoreceptor axon terminals (cone pedicles, rod

spherules) viewed from the synaptic face of the terminals (296 LM cones, gold; 17 S cones,

blue, 24 rods, brown) (see Supplementary Table 1a, 1c, 1d). Five cones marked with white

asterisks were used to generate the vertical connectome analysis shown in Supplementary

Figs. 9 and 10 (S cones 54 and 103; LM cones 21, 22, and 58). **b**. Six cone pedicles within the

white dotted box in **a**. These pedicles contact each other via telodendria (e.g., arrows) within

the circled areas. A pedicle identified as an S cone (white asterisk) contacts neighboring LM

cone 230. **c**. EM micrographs showing telodendritic contacts corresponding to the two solid red

circled areas in **b** (open white arrowheads). Left solid red circle shows contacts between LM

cones 231 and 224, and the right solid red circle shows S cone 14 contact with LM cone 230.

Cone-cone contact surface areas were calculated and used in the simulation shown in **Fig. 8**.

Fig. 3. H1 and H2 horizontal cells synapse with S cones. a. Six overlapping H1 cells are

shown (varied colors); each H1 cell forms lateral elements in S cone 15 (blue). The upper

inset is an inverted view that shows the cell bodies of the six H1 cells. Circular inset at lower

left shows EM view of H1 cell lateral elements at a synaptic invagination of S cone 15 (red

arrowhead). Circular inset at lower right shows a 3D view of S cone 15 (partial transparency)

contacting multiple H1 cells. H1 cells made 43% ± 16% of the total contacts with S cones (n =

489   10 S cones). **b**. Two H2 cells (tan and red) contact multiple S cones, including S cone 15.

490   Lower insets show H2 cell lateral elements at S cone 15 in single-layer EM view (left) and 3D

491   (right) view (conventions as in **a**). H2 cells made 57% ± 16% of the total contacts with S cones

492   (n = 10 S cones).

Fig. 4. Bipolar cell types. a. Outer stratifying types (FMB, flat midget bipolar; DB, diffuse

bipolar). Outer-x bipolar cells lacked a dendrite but made ribbon synapses in the IPL

(Supplementary Fig. 6a). DBbroad cells extend processes across the IPL depth

(Supplementary Fig. 7). **b.** Inner stratifying bipolar types: IMB, invaginating midget bipolar; DB,

diffuse bipolar types; BB, blue-cone bipolar type; Inner-x bipolar (Supplementary Fig. 6c); RB,

rod bipolar. **c**. Horizontal view of spatial mosaics of axon terminals for small groups of

neighboring cells of each bipolar type, as indicated.

Fig. 5. Amacrine cells. Amacrine cells in the INL (253 cells) and GCL (40 cells) were divided

into 19 types (AC1-19; Supplementary Table 1g). **a.** Small-field, AII amacrine cells (AC2), in

vertical view show outer, lobular dendrites (white arrowhead) and extensive, inner dendrites

(green arrowhead). **b.** Horizontal view of the entire AII population in HFseg1; lower panel,

vertical view. **c.** Starburst amacrine cells (AC7) were located primarily in the GCL (GCL, 14

cells; INL, 3 cells). White arrowheads indicate typical fasciculation of starburst dendrites. **d.**

Mosaic of starburst cells, with GCL cells (red-violet) and INL cells (yellow); lower panel,

vertical view. **e**. Small-field amacrine type (AC11) in horizontal view with densely branched

dendritic arbor. **f**. Horizontal view of the AC11 cell mosaic; lower panel, vertical view shows

dendrites extend broadly across the IPL. **g.** Large-field amacrine cells (AC8) with thick lobular

dendrites, narrowly stratified in the outer IPL. **h.** Horizontal view of the complete mosaic; lower

panel shows vertical view of outer IPL stratification. Scale bar in **c** applies to **e** and **g**. Scale

bar in lower panel of **h** applies to **b, d, f**.

Fig. 6. Ganglion cells. a. Cluster of inner-ON (red-violet) and outer-OFF (yellow-green)

midget ganglion cells; dendrites stratify broadly across inner (ON) and outer (OFF) half of the

IPL. **b**. Inner-ON (red-violet) and outer-OFF (yellow-green) parasol ganglion cells; dendrites

stratify broadly near the center of the IPL. **c**. Small bistratified ganglion cells extend dendrites

across the ON-OFF subdivision of the IPL. **d**. Large bistratified ganglion cells (LBGC1) show

large dendritic fields and broad stratification. White arrowhead shows branching processes

arising from the cell body of unknown significance. **e-h**, Mosaic of dendritic arbors in

horizontal view of the same cells shown in **a-d**. **e**. Outer-OFF (yellow-green, top) and inner-

ON (red-violet) midget GC. **f**. Outer-OFF parasol cell dendrites show minimal overlap (inner-

ON parasol mosaic not shown). **g**. Small bistratified cells show minimal dendritic tree overlap.

h. LFGC1 (Large bistratified cells) show a much larger and more sparsely branching irregular

tree. **i-k**. Morphology of three additional large field types. **i**. LFGC2 (Large diffuse cells) show

a densely branched tree broadly stratified across the IPL. **j**. LFGC3 (Recursive bistratified

cells) are narrowly bistratified near the center of the IPL and tend to fasciculate with starburst

amacrine cells. **k**. LFGC4 (Inner smooth monostratified cells) show large, monostratified

dendritic trees costratified with parasol cells. **l-n**. Horizontal views of the cells shown in **i-k.**

Fig. 7. Synapse detection. a. OFF midget bipolar (yellow) and ON midget bipolar (red)

presynaptic to an outer-OFF (blue) and inner-ON (green) midget ganglion cell. **b.** Outgoing

synapses detected for the midget bipolar cell axon terminals shown in **a** (ganglion cells

removed, semi-transparent view). Synaptic ribbons (red/violet dots) and postsynaptic profiles

were first identified manually (see Supplementary Fig. 9; cone 22; BC 308 (IMB) and BC 303

(FMB)). Predicted outgoing synapses (green dots and lines, IMB; blue dots and lines, FMB) are

shown. White arrowheads numbered 1-4 indicate the panels shown on the right in single-layer

EM view. Predicted and segmented synaptic ribbons (in varied colors, red arrowheads);

predicted postsynaptic segments are marked by the dots and lines linking the synaptic ribbon

to its postsynaptic structure (unlabeled postsynaptic profiles belong to amacrine cells). Inset

histogram shows precision and sensitivity of ribbon detection and synaptic assignment. **c.**

Inhibitory output synapses made by two neighboring AII amacrine cells (purple and orange,

shown in partial transparency). Yellow dots show synaptic output predictions; most synaptic

outputs arise in the outer half of the IPL and are presynaptic to OFF bipolar cells. Inset shows

precision and sensitivity for the synapses made by each AII cell. **d-e**. Single-layer images

showing segmented vesicle clouds (varied colors) in the purple AII cell (shown in **c**) and

synaptic connections predicted with an FMB cell (yellow dots and lines; ribbons in the FMB are

also segmented). **f-g**. Same conventions as in **d-e** but showing three segmented vesicle clouds

and synaptic predictions for the orange AII cell (shown in **c**), with another FMB cell.

Fig. 8. Cone-cone coupling alters S cone spectral tuning. a. Biophysical model of the cone-

cone connectome. Assignment of L/M cone type was random with a ~1.7:1 L:M ratio, based on

known proportions in the human fovea. Contacts between cones were determined from the

reconstructed connectome, and conductances were assigned based on the contact area, known

density gap junction density, and single connexin channel conductance. Cones were modeled

compartmentally as a soma that included the phototransduction cascade, connected to an axon,

in turn connected to the axon terminal (see model parameters in Supplementary Table 4). **b**.

Modeled voltage responses (averages from 5 different L, M and S cones in the network) to

monochromatic 10 msec light flashes at 420 nm and 550 nm. **Top:** the cone-cone network is

uncoupled (zero cone-cone conductance). **Bottom**: the network is coupled via the individual

cone-cone conductances given in the electrical model. Coupling among L, M, and S cones

produces a light-evoked response in S cones at 550 nm that is absent in the uncoupled

condition. **c**, **d**. Impact of cone-cone coupling on the action spectrum of the cone voltage

response. Average light responses were generated every 5 nm from 350 to 700 nm. In **c**

(uncoupled condition), the action spectrum is proportional to the spectral sensitivity of the L, M,

and S cone opsins. In **d** (coupled condition), the S cone voltage response shows a smaller

second peak of sensitivity at ~550 nm (teal arrowhead) which is not present when cones are

uncoupled.

Fig. 9. AII cells are postsynaptic to both ON and OFF cone bipolar cells at ribbon

synapses. a. Chemical (ribbon) synaptic inputs from multiple OFF and ON bipolar types. For

this cell, 70 synapses (red dots, presynaptic locus; blue, postsynaptic locus; most blue

postsynaptic dots are overlying the presynaptic position and are not visible in this zoomed-out

view) are dispersed throughout the dendritic tree. We found 300 synapses arising from all ON

and OFF cone bipolar types to an initial sample of 6 AII cells, with IMB and FMB cells

accounting for 19% and 24% of the synapses respectively. **b**. By contrast, this AII cell is

presynaptic only to OFF cone bipolar cells; many of these synapses are reciprocal. **c**. An

OFF-midget bipolar (FMB, violet) makes a ribbon dyad synapse (blue arrowhead points to

ribbon) with the AII cell (yellow) and a parasol ganglion cell. The AII cell makes a nearby

reciprocal synapse (red arrowhead) upon this FMB. **d.** Nonreciprocal AII synapse with a DB2

OFF bipolar type (teal). **e.** AII cell (yellow) postsynaptic to an ON-midget bipolar (IMB, green)

at two dyad synapses (blue arrowheads). **f**. 3D view of a reciprocal synapse between this AII

and a DB3 bipolar (blue); the AII inserts a finger-like projection into this bipolar axon terminal.

592   **g**. Single section through contact in **f,** showing two of three ribbon synapses (blue arrowhead

593   indicates one) and the reciprocal AII inhibitory synapse (red arrowhead).

Fig. 10. Vertical excitatory connectome. a. Vertical connectome for a single LM cone (cone

22, see details in Supplementary Fig. 9b). Cone pedicle and postsynaptic bipolar cells shown in

silhouette. Synaptic ribbons in the cone pedicle (top, OPL) and bipolar cell axon terminals

(lower, IPL) are shown as red balls. Inset shows all bipolar cells postsynaptic to this cone

pedicle (outer, yellow-gold; inner, blue-violet-red). **b**. The same views for an identified S-cone

(cone 103, see details in Supplementary Fig. 10b; synaptic ribbons, blue dots). **c**. Histogram

shows output synapse numbers (left axis), and percent (right axis) made via bipolar cells

connected to 3 LM and 2 S cones to postsynaptic GC types. Note that OFF-pathway-derived

synapses greatly outnumber ON pathway synapses. **d**. The great majority of synapses from

LM and S cones are distributed to midget, parasol, and small bistratified ganglion cells (Major

GCs), with all other large-field types receiving 7.9% (LM) and 6.4% (S) of the synaptic output

(see also Supplementary Table 2). Unidentified processes account for 0.5% of the total

synaptic output (LM) (see Supplementary Table 2a-2c).

Methods

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Human tissue acquisition and preparation

Human eyes (52-year-old, white male; brain dead organ donor) were acquired from the

Medical University of Vienna, Vienna, Austria at the time of death by surgical enucleation. A

small incision was made to give fixative access to the posterior chamber and the retina was in

this way immersion fixed in 4% glutaraldehyde in 0.1 M sodium cacodylate buffer, pH7.3-7.4 at

room temperature. Medical history confirmed no abnormalities of the visual system recorded

and no driving limitations; the ocular status was defined medically as unremarkable at time of

enucleation. Eyes were shipped in fixative to the University of Washington where the retinas

were dissected and prepared for electron microscopy.

METHOD DETAILS

Identification of foveal ROI, serial block-face SEM sample preparation and image

acquisition

Retinal tissue was dissected under a dissecting microscope to facilitate identification of the

optic disc and the center of the foveal pit as well as the major retinal meridians. We selected

an ROI that extends across absolute eccentricity of ~450-650 microns (temporal retina). This

location is linked to cones located ~100-300 microns from the foveal pit center due to the

lateral migration of retinal circuits from the cones at the foveal center. This location

encompasses much of the central 1 degree (~300 µm/deg) of visual angle (Fig. 1a-1c).

The dissected retina was then plastic embedded and prepared for electron microscopy as

previously described[1]. In brief, the tissue was incubated in a 1.5% potassium ferrocyanide and

2% osmium tetroxide (OsO4) solution for 1 h. After washing, the tissue was placed in a freshly

made thiocarbohydrazide solution for 20 min and then rinsed and incubated in 2% OsO4 for

30 min. Lastly, the tissue was stained en bloc in 1% uranyl acetate overnight and

subsequently stained with Walton's lead aspartate, dehydrated and plastic embedded

(Durcupan, 44610, Sigma Aldrich). After evaluation of the ROI by semithin sectioning the

tissue block was trimmed, gold-coated by standard methods, and mounted in an Apreo SEM

Volumescope (Thermo Fisher). We cut 3028 sections at 50 nm thick for a total depth of ~150

microns, imaging an ROI (180 µm tall x 180 µm wide; 5 nm pixels) that encompassed the full

vertical depth of the retina, from the Henle fiber layer (HFL) to the nerve fiber layer (Fig. 1d).

The overall ROI was imaged on the blockface after each section as 25 sequential tiles (8000 x

8000 pixels or 40 x 40 µm; 75,700 total image tiles) with 10% overlap. Images were stitched

within layer and aligned across layers using standard methods available in TrakEM2[2] (NIH

image plugin; Scale Invariant Feature Transform (SIFT) with affine translation) to create a

good quality volume. We first evaluated this TrakEM2-aligned volume by manual

reconstruction of a small number of selected cells and circuits[106]. To further develop this

volume as a connectomic resource we applied deep-learning based computational

approaches to realign, segment and reconstruct all cells and synaptic connections as

follows[20, 39].

Volume creation

Montaging. Tile offsets were determined manually, then refined by template matching at 40 nm

with a search window of 96 pixels. Tiles were cropped by 128 pixels to avoid edge distortions

and placed to minimize the sum of least squares from the new offsets. Outliers were inspected

and given manual placement. These cropped tiles were processed to create a pyramid of

encodings for resolutions between 20 nm and 640 nm[107], and then stitched to make the

montaged section in two steps. In the first step, only the two neighboring tiles in Y were

considered, with both neighbors being fixed. Using an online finetuner pyramid[107] with the

calculated placement as the starting point, a displacement field was generated that would

warp the encoded tiles to minimize the pixel difference with the fixed tiles. The encoded tiles

were then warped using this displacement field, eroded by a pixel to avoid partial pixels at the

edge, blended to the fixed tiles using the mean of the pixel values in the overlap, and then

passed through CLAHE (Contrast Limited Adaptive Histogram Equalization) enhancement to

mitigate the contrast loss from the warping and blending, resulting in strips of encoded tiles

where every other tile had been warped. In the second step, the strips of tiles from the first

step were stitched to each other, holding every other strip fixed in a similar process. An error

map was generated at both steps by computing the difference between the encodings in the

overlap at 40 nm and inspected. The non-encoded tiles were cropped by 394 pixels, warped

according to the composed displacement fields from the encodings, and combined to yield

montaged sections.

Alignment. The unaligned sections were first processed to create a pyramid of encodings[107]. A

rigid transformation for the entire dataset was estimated using SIFT features[108] extracted from

the encodings at 640 nm and correspondences established between nearest and next-nearest

section pairs. From the rigidly aligned encodings, displacement fields at 40 nm resolution were

generated between nearest and next-nearest section pairs using an online finetuner

pyramid[107].

A convolutional net trained to detect local misalignments and/or simple mismatches caused by

defects in the EM images was used to mask unreliable regions in each displacement field. In

addition, 25 sections with defects that presented with non-smooth deformations were masked

by hand. Pairwise displacement fields and transformation masks were then passed to a global

relaxation[107, 109]. This relaxation was performed first at 2560 nm across all 3028 sections of the

dataset, and subsequently in blocks of approximately 100 sections at a resolution of 640 nm,

using the previously generated drift-free fields as constraints for the start and end sections of

each of the 31 blocks. The resulting displacement fields were composed with the earlier rigid

transformation and the final aligned image stack warped and rendered at 10 nm.

Segmentation. We pretrained a convolutional net to predict affinities between neighboring

voxels at 16×16×40 nm³ using 1,368 µm³ of manually annotated data derived from three

mouse visual cortex datasets[110]. We used the residual symmetric U-Net architecture[111] with

modifications[112] and trained with data augmentation previously described [113]. The network was

trained for 2,000,000 iterations using four NVIDIA L4 GPUs with PyTorch data parallelism,

where each GPU processed individual patches with its own model replica.

To further fine-tune this network, we manually created training data consisting of 39 µm³ at a

resolution of 20×20×50 nm³. In contrast to the pretraining datasets, mitochondria were

intentionally over segmented to bias boundary detection toward a more conservative

approach, aiming primarily to reduce merge errors at the expense of increasing split errors.

We fine-tuned the network for 300,000 iterations.

We predicted an affinity map for the whole dataset. Dark voxels could sometimes span

multiple cells and cause segmentation errors. We therefore applied a binarized mask for dark

voxels with a threshold of 0.2, invalidating each dark voxel and its three incident affinities. The

resulting masked affinity map was then segmented with a mean affinity agglomeration

threshold of 0.25[113]. The segmentation was ingested into the Connectome Annotation

Versioning Engine proofreading platform, CAVE[114] for further analysis.

CAVE permitted proofreading to be performed by a user community accessing the web-based

neuroglancer interface[113, 115] with dynamic updating of segment ID and related synaptic

connectivity. The HFseg 1 segmentation was of high initial quality such that it was possible to

identify and annotate all major cell classes within the volume and to sort these cells into

populations of distinct types using the annotation tools available in neuroglancer. All cells are

accessible at NeuroMaps (https://neuromaps.app).

Cell types were identified provisionally based on segment 3D morphology (dendritic

morphology, stratification depth, and spatial regularity and interaction (mosaics) and in some

cases nuclear morphology). Four of the authors (Kim, Dacey, Grünert and Martin) conferred

and agreed on cell type identification. Proofreading is ongoing (~ 40,000 edits) and applied

thus far primarily to correct larger obvious mergers (primarily with glial cell processes) that

interfered with cell type identification and additional more detailed proofreading on circuits of

interest in the current survey of cone-bipolar-ganglion cell connections. Many cell populations

were unambiguously recognized based on previous studies of primate retina[41, 116, 117] (e.g., the

S cones, rods and LM cones, the H1 and H2 horizontal cells, the starburst and AII amacrine

cell types, the midget, diffuse, blue-cone bipolar, and rod bipolar types, or the midget, parasol

and small bistratified and large bistratified ganglion cell types; Müller glia, astrocytes and

microglia); some cell types (the low-density amacrine types) were provisionally designated

subject to further proofreading and connectivity analysis. Following FlyWire[40], we have

initiated an open community of retinal neurobiologists (currently 7 labs participating) to use

these tools to complete proofreading of the HFseg1 volume.

Synapse detection

Ribbon synapse detection in OPL and IPL. A symmetric residual U-Net architecture was

trained on data with a resolution of $10\times10\times50$ nm^3 using input patches of size

$128\times128\times20$ to generate a probability map that indicates the likelihood of each voxel

belonging to a synaptic ribbon [118]. To remove less robust predictions at the boundaries,

only the central $96\times96\times16$ voxels of the input patch was used as the output. The model

was trained on a total volume of 2,254 μm^3 of manually labeled cutouts sampled from

various regions across the entire dataset.

The ribbon segmentation was produced by (1) down sampling the probability map by 2×, (2)

thresholding for voxels above 0.15, and (3) using 26-connectivity to extract connected

components. Segments smaller than 5 voxels in a 20×20×50 nm^3 resolution were neglected to

eliminate spurious detections.

To achieve ribbon detection in the OPL segments smaller than 5 voxels (in a 20×20×50 nm^3

resolution volume) were first disregarded to eliminate spurious detections of extremely small

ribbon-like structures. For each detected ribbon within a cone pedicle, postsynaptic partners

were identified via the following steps: (1) All neighboring segments touching the pedicle

within 800 nm of the ribbon centroid were identified. (2) For each neighbor, a representative

point close to the ribbon was chosen as the postsynaptic site. These constitute the set of

putative synapses for each ribbon. (3) After cell identification, only putative synapses onto

bipolar or horizontal cells were kept; all others were filtered out (e.g. Müller glia, neighboring

cones, mitochondria). (4) Finally, for each distinct ribbon-postsynaptic cell pair, all but the

shortest putative synapse were filtered out. The result was that a single synaptic connection

was found onto any bipolar or horizontal cell contacting that cone pedicle within 800 nm of a

given cone ribbon.

For synaptic ribbons in the IPL, segments smaller than 5 voxels in a 20×20×50 nm^3 resolution

within the bounds of the IPL were also disregarded to eliminate spurious detections. For each

detected ribbon, postsynaptic partners were automatically identified via the following steps: (1)

Identify the two largest cells within a small (500 nm) window of the centroid of the ribbon

which are adjacent to the presynaptic cell and are also adjacent to each other. (2) If such a

pair is found, the ribbon is assumed to be presynaptic to both (i.e., a dyadic synapse). (3) If no

such pair is found, then within the window, the single largest profile adjacent to the

presynaptic cell is identified as the postsynaptic partner. (4) Putative synaptic connections

were filtered by cell type, requiring the presynaptic cell to be previously annotated as a bipolar

cell, and the postsynaptic partner to be previously annotated as amacrine, or ganglion cell.

Note we did not at this time try to use a convolutional net to detect a low density of possibly

ribbonless synapses.

Vesicle cloud detection and conventional synapse assignment in the IPL. A symmetric residual

U-Net architecture was trained on data with a resolution of 10×10×50 nm^3 using input patches

of size 128×128×20 to generate a probability map that indicates the likelihood of each voxel

belonging to a vesicle cloud[118]. To remove less robust predictions at the boundaries, only the

central 96×96×16 voxels of the input patch was used as the output. The model was trained on

a total volume of 2,254 μm^3 of manually labeled cutouts sampled from various regions across

the entire dataset (the same cutouts as ribbon synapse detection). Manual labels were

roughly annotated to mark approximate locations of vesicle clouds rather than precise

boundaries.

The vesicle cloud segmentation was produced by (1) downsampling the probability map by

2×, (2) thresholding for voxels above 0.15, and (3) using 26-connectivity to extract connected

components. Segments with fewer than 200 voxels were excluded, based on the range of

vesicle cloud sizes encountered.

IPL conventional synapse assignment. A symmetric residual U-Net architecture was trained on

data with a resolution of 20×20×50 nm^3 using input patches of size 24×24×8 to generate a

probability map that indicates the likelihood of each voxel belonging to the postsynaptic cell[118].

The model was trained on 682 μm^3 of manually annotated labeled cutouts.

The model was applied once for each vesicle cloud detected, in a window centered on the

centroid of the vesicle cloud. Using the initial automatic cell segmentation, the cell with the

highest mean probability within the window was selected as the putative postsynaptic

partner[118]. Further filtering by cell type was applied, requiring the presynaptic cell to be

previously annotated as an amacrine cell, and the postsynaptic partner to be previously

annotated as amacrine, bipolar, or ganglion cell.

Building and modeling the cone-cone contactome

Generating the cone-cone contactome. We identified gap junctions between cone pedicles

and between cone pedicles and rods as follows. Cones and rods were manually identified and

proofread to completion. Contacts were computed using a recently developed supervoxel

agglomeration process (abiss)[119] where a contact is considered to be a set of connected voxel

faces that are shared by the same pair of segments. For each contact, we counted the

number of voxel faces along each orientation (xy, xz, and yz), as well as the minimum

bounding box containing all of the faces. The surface area of a contact was estimated by

computing the sum of the area of all faces involved. Contacts were excluded from analysis if

(1) the surface area of the contact was $\leq$0.04 μm^2 (mean surface area of contact, 0.55 ± 0.3

μm^2, n = 1141 cone-cone contacts) or (2) the center of the bounding box was more than 15

μm from the approximate center of the cone pedicle (manually identified). All remaining

contacts were manually inspected with 1× coverage to determine if they corresponded to what

would be manually identified as a true cone-cone contacts harboring gap junctions.

Biophysical modeling of cone-cone contactome. To generate a biophysical model of the

electrically coupled photoreceptor network, we started with data from the X,Y locations of 310

cone and 24 rod synaptic terminals, and for the cones their M/L vs. S identity. We then

incorporated all reviewed and accepted cone-cone contacts and the surface areas of each

contact.

The cone morphology was modeled as a sphere (single compartment, dia. 2.5 μm) for the

outer and inner segments and soma, connected to an axon (dia. 1.6 μm, Ri 80 Ohm-cm, Rm

12,000 Ohm-cm^2 [7] ; V_{rev} -60 mV, compartment size 0.05 lambda, ~9 compartments/cone

axon) extending 300 μm to the terminal (diam 5 μm, Rm 12,000 Ohm-cm^2, V_{rev} -60mV). Rod

morphology was similar (OS/IS diam 2.5 μm, axon dia 0.45 μm, terminal dia. 3 μm). The

OS/IS/soma compartment contained a transduction mechanism (max conductance 880 pS)

with spectral sensitivity appropriate for the cone identity. The transduction mechanism of the

photoreceptor biochemical cascade generated a peak ~20 ms after the stimulus with a

damped recovery [8-15]. Spectral sensitivity for the L/M cones was taken from non-human

primate[9] and for the S cones from human cone data (peak sens. 420 nm[16] corrected for lens

spectral sensitivity[17]). In some models including both lens and macular filters, the S cone peak

sensitivity was shifted to ~440 nm.

The location of all the S-cones and rods in the reconstructed contactome was known but the

individual identity of L/M cones was unknown, so for the simulations the L/M identity was set

randomly. The target L/M ratio was 1.7 and the actual L/M ratio in the model shown was

1.697. The contacts between cones had not been resolved into gap junctions (groups of

connnexin molecules), but the area of each contact had been measured. The conductance for

each contact was calculated by taking the average area of the contacts (0.55 μm^2), assuming

5 connexins[18] (15 pS per connexin) per average contact[19,20] to produce an average

conductance of 75 pS, or 136 pS/μm^2. The conductance for each individual contact was then

calculated by multiplying by the contact area. Contacts were simulated with a linear gap

junction conductance (i.e. electrical synapse). The cones had on average 3.6 contacts with

neighboring cones. The average L/M total coupling conductance was ~270 pS, and for S-LM

cones was ~282 pS. There were a total of 715 coupled pairs of cones, 659 LM pairs, and 56

pairs with 1 S cone. The average coupling conductance between adjacent pairs of L/M cones

was 116 pS, and for L/M–S pairs was 123 pS. However, the average measured coupling

conductance between LM pairs was 133 pS (n = 25) and between S-LM pairs was 166 pS (n

= 14) due to the existence of multiple current paths between pairs through the gap junction

network. The average cone terminal input resistance was ~0.4 Gohm.

The simulated cones were stimulated with flashes of monochromatic light (10 ms, 1000 µm

spot, ~1e5 Ph/µm^2/sec) on a white background (1e4 Ph/µm^2/s, spectrally flat). The resting

potential V_{rest} in response to the background was ~ -40 mV but varied by ~0.3mV depending

on the cone identity and coupling. The flash response varied from 0 to -5 mV. To determine

the effect of the gap junctions, simulations were run with the gap junction conductance at 0%,

50%, 100%, and multiples of the nominal 136pS/µm^2 conductance. The model was

constructed in the Neuron-C simulation language [21] and comprised ~3000 compartments. The

model was run on AMD Opteron 3000 MHz CPUs under the Linux and Mosix operating

systems. Approximately 200 simulations were run of the 310-cone model to develop and

calibrate the model. Modeling code can be accessed at zenodo.org upload of Neuron-C at:

https://doi.org/10.5281/zenodo.17792059.

Proofreading, synaptic connectivity and updating with the Connectome Annotation

Versioning Engine (CAVE) and NeuroMaps.

As noted above we ingested synapse predictions and cell type annotations into CAVE. A

critical functionality given by CAVE is that is allows collaborative users to programmatically

query the up-to-date proofread connectivity graph (synaptic connectivity is updated during

proofreading of each segment-cell type) as well as any changes made to the metadata (cell

type IDs, descriptions, or other annotations). We then adapted NeuroMaps to combine this

metadata with synaptic connectivity for each segment ID and to permit synaptic proofreading

(addition or deletion of specific synaptic predictions).

DATA AND SOFTWARE AVAILABILITY

The HFseg1 dataset (volume, segmentation, synaptic detection and full annotation) are

currently hosted by Zetta.ai (https://zetta.ai/) and all data is accessible for community

proofreading by permission at NeuroMaps. The latest version of the HFseg1 dataset is also

available for interrogation at NeuroMaps (https://neuromaps.app). Modeling code can be

accessed at zenodo.org upload of Neuron-C at: https://doi.org/10.5281/zenodo.17792059

References

1.   Ramon y Cajal S. La rètine des vertèbrès. *La Cellule* **9**, 119-257 (1892).

2.   Masland RH, Raviola E. Confronting complexity: strategies for understanding the
microcircuitry of the retina. *Annual review of neuroscience* **23**, 249-284 (2000).

3.   Dowling JE. *The Retina: An approachable part of the brain*, 2nd edn. Harvard
University Press (2012).

4.   Rodieck RW. *The First Steps in Seeing*. Sinauer Associates, Inc. (1998).

5.   Helmstaedter M, Briggman KL, Turaga SC, Jain V, Seung HS, Denk W. Connectomic
reconstruction of the inner plexiform layer in the mouse retina. *Nature* **500**, 168-174
(2013).

6.   Kim JS*, et al.* Space-time wiring specificity supports direction selectivity in the retina.
Nature **509**, 331-336 (2014).

7.   Briggman KL, Helmstaedter M, Denk W. Wiring specificity in the direction-selectivity
circuit of the retina. *Nature* **471**, 183-188 (2011).

8.   Anderson JR*, et al.* Exploring the retinal connectome. *Molecular vision* **17**, 355-379
(2011).

9.   Calkins DJ, Schein SJ, Tsukamoto Y, Sterling P. M and L cones in macaque fovea
connect to midget ganglion cells by different numbers of excitatory synapses. *Nature*
371, 70-72 (1994).

10.   Calkins DJ, Sterling P. Absence of spectrally specific lateral inputs to midget ganglion
cells in primate retina. *Nature* **381**, 613-615 (1996).

11.   Shekhar K, *et al.* Comprehensive Classification of Retinal Bipolar Neurons by Single-
897   Cell Transcriptomics. *Cell* **166**, 1308-1323 e1330 (2016).

899   12.   Peng YR, *et al.* Molecular Classification and Comparative Taxonomics of Foveal and
900   Peripheral Cells in Primate Retina. *Cell* **176**, 1222-1237 (2019).

902   13.   Yan W, *et al.* Cell Atlas of The Human Fovea and Peripheral Retina. *Sci Rep* **10**, 9802
903   (2020).

905   14.   Hahn J, *et al.* Evolution of neuronal cell classes and types in the vertebrate retina.
906   *Nature* **624**, 415-424 (2023).

908   15.   Li J, *et al.* Integrated multi-omics single cell atlas of the human retina. *Res Sq*, (2023).

910   16.   Dorkenwald S, *et al.* Neuronal wiring diagram of an adult brain. *Nature* **634**, 124-138
911   (2024).

913   17.   Azevedo A, *et al.* Connectomic reconstruction of a female Drosophila ventral nerve
914   cord. *Nature* **631**, 360-368 (2024).

916   18.   Shapson-Coe A, *et al.* A petavoxel fragment of human cerebral cortex reconstructed at
917   nanoscale resolution. *Science* **384**, eadk4858 (2024).

919   19.   Loomba S, *et al.* Connectomic comparison of mouse and human cortex. *Science* **377**,
920   eabo0924 (2022).

922   20.   Turner NL, *et al.* Reconstruction of neocortex: Organelles, compartments, cells,
923   circuits, and activity. *Cell* **185**, 1082-1100 e1024 (2022).

925   21.   Ding Z, *et al.* Functional connectomics reveals general wiring rule in mouse visual
926   cortex. *Nature* **640**, 459-469 (2025).

927
928   22.   Abbott LF, *et al.* The Mind of a Mouse. *Cell* **182**, 1372-1376 (2020).

929
930   23.   Lichtman JW, Sanes JR. Ome sweet ome: what can the genome tell us about the
931   connectome? *Current opinion in neurobiology* **18**, 346-353 (2008).

932
933   24.   Wandell BA, Dumoulin SO, Brewer AA. Visual field maps in human cortex. *Neuron* **56**,
934   366-383 (2007).

935
936   25.   Sereno MI, Sood MR, Huang RS. Topological Maps and Brain Computations From
937   Low to High. *Frontiers in systems neuroscience* **16**, 787737 (2022).

938
939   26.   Harvey BM, Dumoulin SO. The relationship between cortical magnification factor and
940   population receptive field size in human visual cortex: constancies in cortical
941   architecture. *J Neurosci* **31**, 13604-13612 (2011).

942
943   27.   Sereno MI, *et al.* Borders of multiple visual areas in humans revealed by functional
944   magnetic resonance imaging. *Science* **268**, 889-893 (1995).

945
946   28.   Schira MM, Tyler CW, Breakspear M, Spehar B. The foveal confluence in human visual
947   cortex. *J Neurosci* **29**, 9050-9058 (2009).

948
949   29.   Baden T, Berens P, Franke K, Roman Roson M, Bethge M, Euler T. The functional
950   diversity of retinal ganglion cells in the mouse. *Nature* **529**, 345-350 (2016).

951
952   30.   Bae JA, *et al.* Digital Museum of Retinal Ganglion Cells with Dense Anatomy and
953   Physiology. *Cell* **173**, 1293-1306 e1219 (2018).

31. Goetz J*, et al.* Unified classification of mouse retinal ganglion cells using function,
morphology, and gene expression. *Cell reports* **40**, 111040 (2022).

32. Masri RA, Percival KA, Koizumi A, Martin PR, Grünert U. Survey of retinal ganglion cell
morphology in marmoset. *J Comp Neurol* **527**, 236-258 (2019).

33. Kim YJ*, et al.* Origins of direction selectivity in the primate retina. *Nature*
communications **13**, 2862 (2022).

34. Curcio CA, Sloan KR, Kalina RE, Hendrickson AE. Human photoreceptor topography.
JCompNeurol **292**, 497-523 (1990).

35. Curcio CA, Kar D, Owsley C, Sloan KR, Ach T. Age-Related Macular Degeneration, a
Mathematically Tractable Disease. *Investigative ophthalmology & visual science* **65**, 4
(2024).

36. Schein SJ. Anatomy of macaque fovea and spatial densities of neurons in foveal
representation. *J Comp Neurol* **269**, 479-505 (1988).

37. Drasdo N, Millican CL, Katholi CR, Curcio CA. The length of Henle fibers in the human
retina and a model of ganglion receptive field density in the visual field. *Vision Res* **47**,
2901-2911 (2007).

38. Masri RA, Grünert U, Martin PR. Analysis of Parvocellular and Magnocellular Visual
Pathways in Human Retina. *J Neurosci* **40**, 8132-8148 (2020).

39. Lee K, Turner N, Macrina T, Wu J, Lu R, Seung HS. Convolutional nets for
reconstructing neural circuits from brain images acquired by serial section electron
microscopy. *Current opinion in neurobiology* **55**, 188-198 (2019).

40.   Dorkenwald S*, et al.* FlyWire: online community for whole-brain connectomics. *Nature*
986   *methods* **19**, 119-128 (2022).

988   41.   Grünert U, Martin PR. Cell types and cell circuits in human and non-human primate
989   retina. *Progress in retinal and eye research*, 100844 (2020).

991   42.   Grünert U, Martin PR. Morphology, Molecular Characterization, and Connections of
992   Ganglion Cells in Primate Retina. *Annu Rev Vis Sci* **7**, 73-103 (2021).

994   43.   Zhang L*, et al.* Evolutionary and developmental specialization of foveal cell types in the
995   marmoset. *Proceedings of the National Academy of Sciences of the United States of*
996   *America* **121**, e2313820121 (2024).

998   44.   Kim YJ, Packer O, Pollreisz A, Martin PR, Grünert U, Dacey DM. Comparative
999   connectomics reveals noncanonical wiring for color vision in human foveal retina.
1000   *Proceedings of the National Academy of Sciences of the United States of America* **120**,
1001   e2300545120 (2023).

1002

1003   45.   Dacey DM. The mosaic of midget ganglion cells in the human retina. *JNeurosci* **13**,
1004   5334-5355 (1993).

1005

1006   46.   Dacey DM. Morphology of a small-field bistratified ganglion cell type in the macaque
1007   and human retina. *Visual neuroscience* **10**, 1081-1098 (1993).

1008

1009   47.   Dacey DM, Lee BB. The blue-ON opponent pathway in primate retina originates from a
1010   distinct bistratified ganglion cell type. *Nature* **367**, 731-735 (1994).

1011

1012   48.   Chichilnisky EJ, Baylor DA. Receptive-field microstructure of blue-yellow ganglion cells
1013   in primate retina. *NatNeurosci* **2**, 889-893 (1999).

1014

1015 49. Calkins DJ, Tsukamoto Y, Sterling P. Microcircuitry and mosaic of a blue-yellow
1016   ganglion cell in the primate retina. *JNeurosci* **18**, 3373-3385 (1998).

1017

1018 50. Masri RA, Greferath U, Fletcher EL, Martin PR, Grünert U. Immunohistochemistry and
1019   spatial density of Müller cells in the human fovea. *Investigative ophthalmology & visual*
1020   *science* **in press**, (2025).

1021

1022 51. Singaravelu J, Zhao L, Fariss RN, Nork TM, Wong WT. Microglia in the primate
1023   macula: specializations in microglial distribution and morphology with retinal position
1024   and with aging. *Brain Struct Funct* **222**, 2759-2771 (2017).

1025

1026 52. Goyal M, Bordt AS, Neitz J, Marshak DW. Trogocytosis of neurons and glial cells by
1027   microglia in a healthy adult macaque retina. *Sci Rep* **13**, 633 (2023).

1028

1029 53. Haverkamp S, Grünert U, Wassle H. The cone pedicle, a complex synapse in the
1030   retina. *Neuron* **27**, 85-95 (2000).

1031

1032 54. Grabner CP*, et al.* Mechanisms of simultaneous linear and nonlinear computations at
1033   the mammalian cone photoreceptor synapse. *Nature communications* **14**, 3486 (2023).

1034

1035 55. Herr S, Ngo IT, Huang TM, Klug K, Sterling P, Schein S. Cone synapses in macaque
1036   fovea: II. Dendrites of OFF midget bipolar cells exhibit Inner Densities similar to their
1037   Outer synaptic Densities in basal contacts with cone terminals. *Visual neuroscience*
1038   **28**, 17-28 (2011).

1039

1040 56. Strettoi E, Raviola E, Dacheux RF. Synaptic connections of the narrow-field, bistratified
1041   rod amacrine cell (AII) in the rabbit retina. *JCompNeurol* **325**, 152-168 (1992).

1042

1043 57. Strettoi E, Masri RA, Grünert U. AII amacrine cells in the primate fovea contribute to
1044   photopic vision. *Sci Rep* **8**, 16429 (2018).

1045

1046   58.   Wässle H, Grünert U, Chun MH, Boycott BB. The rod pathway of the macaque monkey
1047   retina: Identification of AII-amacrine cells with antibodies against calretinin.
1048   *JCompNeurol* **361**, 537-551 (1995).

1049

1050   59.   Grünert U. Anatomical evidence for rod input to the parvocellular pathway in the visual
1051   system of the primate. *EurJNeurosci* **9**, 617-621 (1997).

1052

1053   60.   Hornstein EP, Verweij J, Schnapf JL. Electrical coupling between red and green cones
1054   in primate retina. *Nature neuroscience* **7**, 745-750 (2004).

1055

1056   61.   O'Brien JJ, Chen X, Macleish PR, O'Brien J, Massey SC. Photoreceptor coupling
1057   mediated by connexin36 in the primate retina. *J Neurosci* **32**, 4675-4687 (2012).

1058

1059   62.   Wool LE, Packer OS, Zaidi Q, Dacey DM. Connectomic identification and three-
1060   dimensional color tuning of S-OFF midget ganglion cells in the primate retina. *J*
1061   *Neurosci* **39**, 7893-7909 (2019).

1062

1063   63.   Lamb TD, Simon EJ. The relation between intercellular coupling and electrical noise in
1064   turtle photoreceptors. *J Physiol* **263**, 257-286 (1976).

1065

1066   64.   Tessier-Lavigne M, Attwell D. The effect of photoreceptor coupling and synapse
1067   nonlinearity on signal:noise ratio in early visual processing. *Proc R Soc Lond B Biol Sci*
1068   **234**, 171-197 (1988).

1069

1070   65.   DeVries SH, Qi X, Smith R, Makous W, Sterling P. Electrical coupling between
1071   mammalian cones. *Current biology : CB* **12**, 1900-1907 (2002).

1072

1073   66.   Hsu A, Smith RG, Buchsbaum G, Sterling P. Cost of cone coupling to trichromacy in
1074   primate fovea. *Journal of the Optical Society of America A, Optics, image science, and*
1075   *vision* **17**, 635-640 (2000).

1076

1077   67.   Li W, DeVries SH. Separate blue and green cone networks in the mammalian retina.
1078   *Nature neuroscience* **7**, 751-756 (2004).

1079

1080   68.   Kolb H, Goede P, Roberts S, McDermott R, Gouras P. Uniqueness of the S-cone
1081   pedicle in the human retina and consequences for color processing. *JCompNeurol*
1082   **386**, 443-460 (1997).

1083

1084   69.   Tsukamoto Y, Masarachia P, Schein SJ, Sterling P. Gap junctions between the pedicles
1085   of macaque foveal cones. *Vision Res* **32**, 1809-1815 (1992).

1086

1087   70.   Hofer H, Carroll J, Neitz J, Neitz M, Williams D. Organization of the human trichromatic
1088   cone mosaic. *J Neurosci* **25**, 9669-9679 (2005).

1089

1090   71.   Thoreson WB, Mangel SC. Lateral interactions in the outer retina. *Progress in retinal*
1091   *and eye research* **31**, 407-441 (2012).

1092

1093   72.   Crook JD, Manookin MB, Packer OS, Dacey DM. Horizontal cell feedback without
1094   cone type-selective inhibition mediates "red-green" color opponency in midget
1095   ganglion cells of the primate retina. *J Neurosci* **31**, 1762-1772 (2011).

1096

1097   73.   Wool LE, Crook JD, Troy JB, Packer OS, Zaidi Q, Dacey DM. Nonselective Wiring
1098   Accounts for Red-Green Opponency in Midget Ganglion Cells of the Primate Retina. *J*
1099   *Neurosci* **38**, 1520-1540 (2018).

1100

1101   74.   Martin PR, Blessing EM, Buzas P, Szmajda BA, Forte JD. Transmission of colour and
1102   acuity signals by parvocellular cells in marmoset monkeys. *J Physiol* **589**, 2795-2812
1103   (2011).

1104

1105   75.   Goodchild AK, Chan TL, Grünert U. Horizontal cell connections with short-wavelength-
1106   sensitive cones in macaque monkey retina. *Vis Neurosci* **13**, 833-845 (1996).

1107

1108   76.   Dacey DM, Lee BB, Stafford DK, Pokorny J, Smith VC. Horizontal cells of the primate

1109   retina: cone specificity without spectral opponency. *Science* **271**, 656-659 (1996).

1110

1111   77.   Chan TL, Grünert U. Horizontal cell connections with short wavelength-sensitive cones

1112   in the retina: A comparison between New World and Old World primates.

1113   *JCompNeurol* **393**, 196-209 (1998).

1114

1115   78.   Tsukamoto Y, Omi N. Some OFF bipolar cell types make contact with both rods and

1116   cones in macaque and mouse retinas. *Frontiers in neuroanatomy* **8**, 105 (2014).

1117

1118   79.   Tsukamoto Y, Omi N. OFF bipolar cells in macaque retina: type-specific connectivity in

1119   the outer and inner synaptic layers. *Frontiers in neuroanatomy* **9**, 122 (2015).

1120

1121   80.   Della Santina L*, et al.* Glutamatergic Monopolar Interneurons Provide a Novel Pathway

1122   of Excitation in the Mouse Retina. *Current biology : CB* **26**, 2070-2077 (2016).

1123

1124   81.   Lee SC, Weltzien F, Madigan MC, Martin PR, Grünert U. Identification of AII amacrine,

1125   displaced amacrine, and bistratified ganglion cell types in human retina with antibodies

1126   against calretinin. *J Comp Neurol* **524**, 39-53 (2016).

1127

1128   82.   Demb JB, Singer JH. Intrinsic properties and functional circuitry of the AII amacrine

1129   cell. *Visual neuroscience* **29**, 51-60 (2012).

1130

1131   83.   Diamond JS. Inhibitory Interneurons in the Retina: Types, Circuitry, and Function. *Annu

1132   Rev Vis Sci* **3**, 1-24 (2017).

1133

1134   84.   Manookin MB, Beaudoin DL, Ernst ZR, Flagel LJ, Demb JB. Disinhibition combines

1135   with excitation to extend the operating range of the OFF visual pathway in daylight. *J

1136   Neurosci* **28**, 4136-4150 (2008).

1137

1138   85.   Wassle H, Puller C, Muller F, Haverkamp S. Cone contacts, mosaics, and territories of

1139   bipolar cells in the mouse retina. *J Neurosci* **29**, 106-117 (2009).

1140

1141   86.   Dacey DM, Crook JD, Packer OS. Distinct synaptic mechanisms create parallel S-ON

1142   and S-OFF color opponent pathways in the primate retina. *Visual neuroscience* **31**,

1143   139-151 (2014).

1144

1145   87.   Miyagishima KJ, Grünert U, Li W. Processing of S-cone signals in the inner plexiform

1146   layer of the mammalian retina. *Visual neuroscience* **31**, 153-163 (2014).

1147

1148   88.   Cowan CS*, et al.* Cell Types of the Human Retina and Its Organoids at Single-Cell

1149   Resolution. *Cell* **182**, 1623-1640 e1634 (2020).

1150

1151   89.   Puthussery T, Venkataramani S, Gayet-Primo J, Smith RG, Taylor WR. NaV1.1

1152   channels in axon initial segments of bipolar cells augment input to magnocellular visual

1153   pathways in the primate retina. *J Neurosci* **33**, 16045-16059 (2013).

1154

1155   90.   Li J*, et al.* Comprehensive single-cell atlas of the mouse retina. *iScience* **27**, 109916

1156   (2024).

1157

1158   91.   Yan W, Laboulaye MA, Tran NM, Whitney IE, Benhar I, Sanes JR. Mouse Retinal Cell

1159   Atlas: Molecular Identification of over Sixty Amacrine Cell Types. *J Neurosci* **40**, 5177-

1160   5195 (2020).

1161

1162   92.   Riccitelli S, Vlasits AL, Franke K. Behavior-Specific Computations in the Vertebrate

1163   Retina. *Annu Rev Vis Sci* **11**, 149-173 (2025).

1164

1165   93.   Kerschensteiner D. Feature Detection by Retinal Ganglion Cells. *Annu Rev Vis Sci* **8**,

1166   135-169 (2022).

1167

1168   94.   Hsiang JC, Shen N, Soto F, Kerschensteiner D. Distributed feature representations of

1169   natural stimuli across parallel retinal pathways. *Nature communications* **15**, 1920

1170   (2024).

1171

1172   95.   Lennie P, Movshon JA. Coding of color and form in the geniculostriate visual pathway

1173   (invited review). *Journal of the Optical Society of America A, Optics, image science,*

1174   *and vision* **22**, 2013-2033 (2005).

1175

1176   96.   Crook JD, Packer OS, Troy JB, Dacey DM. Synaptic mechanisms of color and

1177   luminance coding: rediscovering the X-Y dichotomy in primate retinal ganglion cells. In:

1178   *The New Visual Neurosciences* (eds Chalupa LM, Werner JS). MIT press (2014).

1179

1180   97.   Prusky GT, Douglas RM. Characterization of mouse cortical spatial vision. *Vision Res*

1181   **44**, 3411-3418 (2004).

1182

1183   98.   Lettvin JY, Maturana HR, McCulloch W, Pitts WH. What the frog's eye tells the frog's

1184   brain. *Proc IRE* **47**, 1940-1951 (1959).

1185

1186   99.   Rucci M, Poletti M. Control and Functions of Fixational Eye Movements. *Annu Rev Vis*

1187   *Sci* **1**, 499-518 (2015).

1188

1189   100.   Ko HK, Poletti M, Rucci M. Microsaccades precisely relocate gaze in a high visual

1190   acuity task. *Nature neuroscience* **13**, 1549-1553 (2010).

1191

1192   101.   Poletti M, Rucci M. A compact field guide to the study of microsaccades: Challenges

1193   and functions. *Vision Res* **118**, 83-97 (2016).

1194

1195   102.   Shevell SK, Martin PR. Color opponency: tutorial. *Journal of the Optical Society of*

1196   *America A, Optics, image science, and vision* **34**, 1099-1108 (2017).

1197

1198   103.   Stockman A, Brainard DH. Color Vision Mechanisms. In: *OSA Handbook of Optics* (ed
1199   Bass M). 3rd edn. McGraw-Hill (2010).

1200

1201   104.   Lennie P, Haake PW, Williams DR. The design of chromatically opponent receptive
1202   fields. In: *Computational models of visual processing* (eds Landy MS, Movshon JA).
1203   The MIT Press (1991).

1204

1205   105.   Field GD*, et al.* Functional connectivity in the retina at the resolution of photoreceptors.
1206   *Nature* **467**, 673-677 (2010).

1207

1208   106.   Kim YJ, Packer O, Dacey DM. A circuit motif for color in the human foveal retina.
1209   *Proceedings of the National Academy of Sciences of the United States of America* **121**,
1210   e2405138121 (2024).

1211

1212   107.   Popovych S*, et al.* Petascale pipeline for precise alignment of images from serial
1213   section electron microscopy. *Nature communications* **15**, 289 (2024).

1214

1215   108.   Lowe DG. Object recognition from local scale-invariant features. In: *Proceedings of the*
1216   *Seventh IEEE International Conference on Computer Vision*). IEEE (1999).

1217

1218   109.   Saalfeld S, Fetter R, Cardona A, Tomancak P. Elastic volume reconstruction from
1219   series of ultra-thin microscopy sections. *Nature methods* **9**, 717-720 (2012).

1220

1221   110.   Consortium TM*, et al.* Functional connectomics spanning multiple areas of mouse
1222   visual cortex. *bioRxiv*, 2021.2007.2028.454025 (2023).

1223

1224   111.   Lee K, Zung J, Li PH, Jain V, Seung HS. Superhuman accuracy on the SNEMI3D
1225   connectomics challenge. *arXiv* **preprint arXiv:1706.00120**, (2017).

1226

1227   112.   Lee K, Lu R, Luther K, Seung HS. Learning and Segmenting Dense Voxel
1228   Embeddings for 3D Neuron Reconstruction. *IEEE Trans Med Imaging* **40**, 3801-3811
1229   (2021).

1230

1231   113.   Macrina T*, et al.* Petascale neural circuit reconstruction: automated methods. *bioRxiv*,
1232   2021.2008.2004.455162 (2021).

1233

1234   114.   Dorkenwald S*, et al.* CAVE: Connectome Annotation Versioning Engine. *Nature*
1235   *methods*, (2025).

1236

1237   115.   Maitin-Shepard J*, et al.* google/neuroglancer.) (2021).

1238

1239   116.   Boycott BB, Wässle H. Morphological classification of bipolar cells of the primate
1240   retina. *EurJNeurosci* **3**, 1069-1088 (1991).

1241

1242   117.   Wässle H, Boycott BB. Functional architecture of the mammalian retina. *PhysiolRev*
1243   **71**, 447-480 (1991).

1244

1245   118.   Turner NL, Lee K, Lu R, Wu J, Ih D, Seung HS. Synaptic Partner Assignment Using
1246   Attentional Voxel Association Networks. In: *IEEE 17th International Symposium on*
1247   *Biomedical Imaging (ISBI)*) (2020).

1248

1249   119.   Lu R, Zlateski A, Seung HS. Large-scale image segmentation based on distributed
1250   clustering algorithms. *arXiv* **arXiv. http://arxiv.org/abs/2106.10795 [cs.CV]**, 1-8
1251   (2021).

1252

1253