Title:

Evolution of the olfactory system during the radiation of Heliconiini

butterflies

Authors:

Yi Peng Toh[1], Francesco Cicconardi[2], Giorgio Bianchini[3], Richard M Merrill[1]* & Stephen H.

Montgomery[2]*

Institutional affiliations:

[1] Division of Evolutionary Biology, Faculty of Biology, LMU Munich, Germany

[2] School of Biological Science, University of Bristol, Bristol, United Kingdom

[3] School of Geographical Sciences, University of Bristol, Bristol, United Kingdom

Corresponding authors:

YPT: toh@bio.lmu.de

RMM: merrill@bio.lmu.de

SHM: s.montgomery@bristol.ac.uk

Key words: olfactory system evolution, comparative neuroanatomy, Heliconiini,

developmental plasticity, antennal lobe morphology

Abstract

Sensory system evolution plays a crucial role in shaping species' interactions with their environment, yet the extent to which olfactory system diversity reflects ecological and evolutionary pressures at a macroevolutionary scale remains unclear. Here, we investigate the evolution of the olfactory system across the Heliconiini butterfly tribe, an ecologically diverse but closely related group. Using a comparative approach, we examined variation in antennal lobe morphology and its constituent structures, the glomeruli and antennal lobe hub, as well as olfactory receptor repertoires across species. We found that antennal lobe size variation is driven by independent shifts in glomerular and antennal lobe hub volumes, with species-specific differences occurring against a backdrop of broader phylogenetic stability. While no direct associations with ecological traits were observed, certain species showed large expansions in total glomerular volume and olfactory receptor numbers, warranting further investigation into unmeasured ecological or behavioural factors. Additionally, comparisons between wild-caught and insectary-reared individuals revealed a surprising pattern of developmental plasticity, with antennal lobe hub volumes increasing and glomeruli volumes decreasing in captivity, highlighting the influence of environmental conditions on neural development. These findings suggest that olfactory evolution in Heliconiini is shaped by both evolutionary divergence and developmental plasticity, emphasizing the need to integrate phylogenetic, ecological, and developmental perspectives to fully understand sensory system adaptation.

Introduction

When confronted with new or changing environments, populations must adapt to different selective pressures (Darwin, 1859; Coyne & Orr, 2004; Rundle & Nosil, 2005; Schluter, 2009). These adaptations are often behavioural, which ultimately depend on changes in sensory and/or neural systems (Endler, 1992; Boughman, 2002; Montgomery et al., 2021). As such, sensory system evolution allows organisms to meet challenges imposed by their environment by increasing their ability to detect and respond to environmental cues (Dell'Aglio et al., 2024a). For example, cichlids living in depth-differentiated environments display shifts in sensitivity towards long and short-wavelength light, which are associated with habitat-dependent survival (Wright et al.,2019, Seehausen et al., 2008; Maan et al., 2006), and in New World warblers, differences in opsin gene expression are similarly associated with habitat (Bloch, 2015). Beyond the visual system, evidence also exists for adaptive divergence of the mechanoreception, electroreception and olfactory systems (Morris et al., 2021; Arnegard et al., 2010; Wark & Peichel, 2010).

The olfactory system plays a crucial role in most animals, as odour detection is often critical for both mate-finding and foraging (Togunov et al., 2017; Zjacic & Scholz, 2022). In insects, the primary olfactory processing centre is comprised of paired sphere-shaped structures within the midbrain called the antennal lobes, analogous to the olfactory bulbs in mammals (Boeckh et al., 1990) (Fig. 1). Each antennal lobe is made up of glomeruli positioned around a central axonal network of mainly olfactory projection neurons and local interneurons, called the antennal lobe hub (Malnic et al., 1999; Montgomery et al., 2016; Kymre et al., 2021). Each glomerulus houses the terminal projections of olfactory receptor neurons and the dendritic arbors of local interneurons, which help to arbitrate complex interactions across the antennal lobe (Fusca & Kloppenburg, 2021). Different odours are recognized by different combinations of olfactory receptors, and this translates to different glomeruli being activated, such that olfactory information is represented by a combinatorial coding strategy (Hansson et al., 1992; Malnic et al., 1999; Hallem & Carlson, 2006; Bisch-Knaden et al., 2018). This activation pattern is then relayed via projection neurons, which also synapse in the glomeruli, to integrative processing centres in the brain, the mushroom bodies and the lateral horn (Homberg et al., 1989) (Fig 1).

Individual olfactory receptor neurons originate from peripheral structures, predominantly the antennae, but also the maxillary palps and legs, and terminate within one of the antennal lobe glomeruli (Briscoe et al., 2013; Hansson & Anton, 2000; Arnold et al., 1985; Boeckh et al., 1977; Homberg et al., 1989). Two general principles of olfactory system architecture are that each olfactory receptor neuron only expresses a single olfactory receptor, and that each glomerulus receives input from only one class of olfactory receptor neurons. This essentially results in a one-receptor-one-glomerulus relationship (Hildebrand

& Shepherd, 1997). Although this paradigm has been recently challenged by evidence of co-
expression of multiple receptors in single receptor neurons in *Aedes aegypti* and *Drosophila*
melanogaster (Herre et al., 2022; Task et al., 2022), it remains unclear how taxonomically
widespread this exception is.

Insect olfactory receptors function as heteromeric tetramers, consisting of a single
ligand-binding olfactory receptor and a coreceptor, *Orco* (Störtkuhl and Kettler, 2001;
Dobritsa et al., 2003; Vosshall et al., 2000; Larsson et al., 2004). Except for *Orco*, the
conservation of close orthologs and subfamily structures are not detected across insect
orders (Hansson & Stensmyr, 2011). Studies looking at the number of olfactory receptors in
invertebrates range widely from around 10 in Phthiraptera (i.e., lice (Kirkness et al., 2010)) to
~200–400 in Hymenoptera (i.e., bees, ants, and wasps (Robertson et al., 2010, Wurm et al.,
2011)), suggesting olfactory receptor repertoires are both highly variable across species and
intimately linked to species' ecology. Among closely related species, olfactory receptors
show a high rate of birth and death, even when overall receptor numbers remain relatively
constant (Cicconardi et al., 2024; Nei & Rooney, 2005) and are often associated with
signatures of positive selection (Rooney et al., 2002; Piontkivska et al., 2002; Cicconardi et
al., 2017). This rapid turnover suggests that olfactory receptor repertoires can evolve quickly
to meet immediate ecological demands. For example, comparisons amongst close relatives
in *Drosophila* showed an increased expression and upregulation of odorant receptor and
odorant-binding proteins in ecological specialists compared to more generalist species
(Kopp et al., 2008). As another example, in closely related species of *Heliconius* butterflies
living in divergent environments, widespread gene expression divergence was also found,
consistent with evolutionary divergence in the sensory tissues, and this was argued to have
facilitated speciation in the early stages (Wu et al., 2022). Thus, together with the broader
loss of conserved subfamilies across insect orders indicating deep evolutionary divergence
over millions of years, these examples suggest dynamic patterns of evolution in olfactory
receptor repertoires across both long and short evolutionary timescales (Yohe et al., 2020).

While most research on sensory system evolution has focused on changes in these
olfactory receptor gene repertoires (e.g. Carlsson et al., 2013; Maan et al., 2006; Bloch,
2015; Wark & Peichel, 2010), variation in downstream neural processing also plays a
significant role in olfactory evolution. Comparative studies often reveal volumetric differences
in regions of the brain involved in sensory processing, which likely reflects changes in cell
number or size (Gronenberg & Hölldobler, 1999; Kondoh et al., 2003; Streinzer et al., 2013;
O'Donnell et al., 2013; Montgomery & Ott, 2015). For example, detailed studies in
Drosophila have not only revealed evolution in the number of olfactory receptor genes and
the types of olfactory sensilla, but also volumetric changes in the antennal lobe and specific
glomeruli (Auer et al., 2020; Prieto-Godino et al., 2020, 2021; Linz et al., 2013). This

provides evidence that olfactory evolution can occur at multiple points along the olfactory

pathway and illustrate the advantages of studying olfactory system evolution in the context of

an adaptive radiation, where closely related species are ecologically distinct (Depetris-

Chauvin et al., 2023; Auer et al., 2020; Prieto-Godino et al., 2016, 2017).

Studies of Lepidoptera are proving informative as case studies linking ecological

divergence, behaviour and sensory evolution (Carlsson et al., 2013; Montgomery & Merrill,

2017; Montgomery et al., 2021; Hebberecht et al., 2023, Borrero et al., 2024). For example,

differences in the overall volume of the antennal lobe have been found between nocturnal

and diurnal moth species, with these volumetric differences corresponding to differences in

sensory weighting during foraging experiments (Stöckl et al., 2016), lending support to the

prevailing theory that expansions of certain brain regions correspond to enhanced

information processing (Barton, 1998; Barton et al., 1995). Classic comparative studies in

moths have also revealed differences in the arrangement and volume of a region of the

glomeruli termed the macroglomerular complex (Boeckh and Boeckh, 1979; Hansson et al.,

1991; Namiki et al., 2014; Berg et al., 2014; Liu et al., 2021). The macroglomerular complex

is enlarged in males (or is male specific) and is involved in receiving and processing

information about female sex pheromones (Koontz and Schneider, 1987). In contrast to

moths, however, most butterflies lack a macroglomerular complex (Rospars, 1983; Heinze

and Reppert, 2012; Carlsson et al., 2013), although a similar cluster of morphologically

distinct glomeruli has re-emerged at least once in butterflies, in the tribe Ithomini, associated

with increased expression of a small number of olfactory receptors (Montgomery & Ott,

2015; Morris et al., 2021; Cicconardi et al., 2024).

The Nymphalid butterfly tribe, Heliconiini, includes the diverse genus *Heliconius*,

which has radiated across the Neotropics to now include ~50 species, having diverged from

its sister genus *Eueides* as recently as 18 million years ago (Kozak et al., 2015, Cicconardi

et al., 2023). Although *Heliconius* research has largely focused on visual cues and

behaviour (e.g. Dell'Aglio et al., 2016; Rossi et al., 2024, Merrill et al., 2015, Hausmann et

al., 2023), chemoreception plays an important role in many aspects of *Heliconius* behaviour.

Heliconius males produces sex and species-specific compounds which alter female mating

decisions (Darragh et al., 2017; Mérot et al., 2015, González-Rojas et al.,2020), and are also

known to generate anti-aphrodisiac pheromones, which are transferred to females during

copulation, and repel subsequent males (Schulz et al., 2008; Gilbert, 1976; Estrada et al.,

2011). Differences in olfactory receptor expression have also been documented between

the sympatric species, *H. cydno* and *H. melpomene*, which have distinct mating and host

plant preferences, highlighting the potential for olfactory evolution in maintaining species

barriers in this group (Darragh et al., 2017; van Schooten et al., 2020). The antennal lobe

also appears to show frequent differences in size, possibly associated with habitat shifts,

between closely related *Heliconius* species (Montgomery & Merrill, 2017; Montgomery et al.,
2021; Hebberecht et al., 2023). However, while these studies have provided valuable
insights, they have largely been confined to comparisons between individual taxon pairs,
leaving us without a comprehensive understanding of chemosensory evolution across the
Heliconiini.
To further investigate what drives changes in the olfactory system across the
Heliconiini, we expanded neuroanatomical and molecular comparisons across the whole
Heliconiini tribe. Specifically, we explored how the volume of the antennal lobe, as well as its
constituent units, the antennal lobe hub and glomeruli, in addition to the number of olfactory
receptors, vary across the Heliconiini. We tested whether these components of the olfactory
system are associated with ecological variation. Finally, we made use of data from
individuals reared under common garden conditions to determine if any of the morphological
variation we observed across wild populations can be attributed to developmentally induced
plasticity.
183
184   **Methods**
185
Sampling across Heliconiini
We took advantage of an extensive collection of brain samples previously used to examine
mushroom body evolution across the Heliconiini (Couto et al., 2023)*.* These include 318 wild
butterflies from 41 species (average 8 individuals per species; Fig. 2) belonging to the
Heliconiini tribe sampled across sites in Costa Rica, Panama, French Guiana, Ecuador and
Peru. We also made use of total antennal lobe volume segmented in prior studies (Couto et
al., 2023; Hebberecht et al., 2023*)* but extended these data by re-segmenting these samples
to include total antennal lobe, total glomeruli and antennal lobe hub volume.
To determine whether any variation we observed was due to environmentally induced
plasticity, we additionally examined neuropil volumes for five *Heliconius* species raised under
common garden conditions, as described in Hebberecht *et al.* (2023). In brief, stocks of *H.*
erato, *H. melpomene*, *H. cydno*, *H. hecale* and *H. ismenius* were established from wild
individuals caught in Gamboa and the nearby Soberanía National Park, Panama, and
maintained in insectaries (c. 2 × 2 × 2 m) at the Smithsonian Tropical Research Institute. All
butterflies were provided with a ~10% sugar solution, and either *Lantana* spp*.* or *Psiguria*
spp*.* flowers as a source of pollen. *Passiflora* spp. were provided for oviposition, and
caterpillars were raised on their native host plant (Merrill et al., 2013). Adults were
maintained for 2-3 weeks to ensure maturity before sampling, and their age recorded.
All brains had been dissected, imaged, and partially segmented for previous studies
(Couto et al. 2023; Hebberecht et al., 2023). This followed established protocols that stain

against a synaptic protein, illuminating synapse-dense regions of the brain referred to as
neuropils (Ott, 2008). We additionally scanned one side of the antennal lobe, chosen at
random, of two individuals of *E. lybia,* and three individuals of *E. aliphera* for imaging at
higher resolution with an x-y resolution of 1024 × 1024 pixels using a Leica SP8, 10X 0.4NA
lens, and the same settings as previously collected data. These higher-resolution antennal
lobe stacks, together with original confocal scans of *Dryadula phaetusa* in the resolution of
512 x 512 pixels, were used to estimate the number of glomeruli present in these species by
segmenting all observable glomeruli to complement prior estimations of glomeruli numbers
for *H. hecale* and *H. erato* (Montgomery et al., 2016). Measurement error due to
compression of neighbouring glomeruli blurring boundaries between individual glomeruli
creates uncertainty in the precise count of the total glomeruli for an individual, leading to
possible slight over, or under, estimation.
218
Image segmentations
We reconstructed the antennal lobe into the antennal lobe hub and glomeruli in a total of 318
wild-caught individuals, representing 41 species in the Heliconiini tribe, as well as an
additional 44 *Heliconius* insectary reared individuals. To do so, we manually segmented
each neuropil in the whole brain based on the boundaries that we see using the *labelfield*
module of AMIRA using the *magic wand* function. We then obtained the volume of each
neuropil using the *material statistics module* in AMIRA. We reconstructed these neuropils on
one side of the brain, chosen at random, and multiplied them by two to get the total volume
across the brain. We used the volume of the rest of the central brain (rCBR) as an allometric
control which was obtained by previous reconstructions of the constituent brain regions
(Couto et al., 2023; Hebberecht et al., 2022).
230
Ecological data
We obtained data for three ecological traits for each Heliconiini species, in addition to a
number of climatic traits. The ecological traits were the presence/absence of pollen feeding,
the number of host plant species exploited, and the degree of social roosting. Pollen feeding
was categorised as a binary trait (present/absent) following Couto et al. (2023), host plant
use number was obtained from McLellan and Montgomery (2024), and roosting behaviour
was classified as either solitary, loose group, small group, or large group, which was factored
respectively into levels 1-4, following Brown (1981). These ecological traits were chosen as
possible factors that would affect the evolution of sensory systems in Heliconiini: first, pollen
feeding, the active collection of pollen grains as an adult source of amino acids, may involve
processing of floral cues (Gilbert, 1972; Young & Montgomery, 2020); second, host plant
generalism potentially requires recognition of more olfactory cues compared to species that

utilise only a small number of host plant species (Wang et al., 2022; Burger et al., 2013), as
recognition of each specific host plant involves the detection of its unique blend of volatile
organic compounds (Bruce & Pickett, 2011); third, some *Heliconius* species are known to
roost together in large groups at night, potentially representing a site of high-density social
behaviours (Mallet, 1986).
To obtain climatic data, we followed a previous protocol by Rivas-Sánchez et al.
(2023). In brief, we first obtained geo-referenced occurrence data and associated climatic
data for all 41 species from Rosser et al. (2012). Outliers above the elevational range of
2100m were excluded from the dataset. We took into account sampling biases by spatially
thinning occurrence records with the thin function in *spThin* (Aiello-Lammens et al., 2015;
distance parameter = 5 km). For each occurrence data point, we then extracted the climatic
variables BIO1 (Annual Mean Temperature) and BIO12 (Annual Precipitation) (30-arc sec
resolution [~1 km2]) from WorldClim (Fick & Hijmans, 2017) as well as the variables, relative
humidity and average surface wind speed (30-arc sec resolution [~1 km2]) in CHELSEA
(Karger et al., 2017). We took the mean values of each variable for each species in the
thinned data set and used them as the environmental data for our individuals, meaning that
individuals from the same species have the same value for the same environmental factor.
We selected these traits as temperature, precipitation, relative humidity and wind speed
have all been shown to affect the concentration of odourants in the air (Hill & Petrucci, 1999;
Martin et al., 2011; Dalton, 2000; Kuehn et al., 2008; Nagappan et al., 2017). At higher
temperatures, vapor pressure is increased thereby increasing the rate of diffusion of
odourant molecules (Hill & Petrucci, 1999). Both temperature and high odour concentrations
can impact neuronal activity (Martin et al., 2011; Dalton, 2000; Devaud et al., 2001; Devaud,
2003; Störtkuhl et al., 1999). Increased humidity may increase the capacity of the air to carry
odourants and possibly make detection easier (Kuehn et al., 2008). An increase in
precipitation would also increase humidity and so we included it in our analysis. Finally, wind
speeds have also been reported as a potential environmental factor impacting odour
reception and locating (Nagappan et al., 2017).
271
Odorant receptor annotation, phylogeny, and orthology assignment
To obtain the olfactory receptor repertoire across Heliconiini, annotated olfactory receptor
coordinates and protein sequences obtained from previous studies (Cicconardi et al., 2023,
2024) were used to extend the list of annotated genes for Heliconinae. To do so, we
implemented a modification of the pipeline used in Cicconardi et al. (2023), which used a
combination of manual and automatic procedures. In brief, we utilized Miniprot (Li, 2023) to
map protein sequences onto the genomes of 72 species: 44 *Heliconius* species, 6 *Eueides*
species, 8 species sister to *Heliconius* and *Eueides* within the Heliconiini tribe, 2 *Argynnini*

species, and 12 outgroup species from the Nymphalid family (including those most distantly
related to the Heliconinae). This comprehensive mapping aimed to infer orthologous
relationships across the Nymphalid family. Coding sequences (CDS) were extracted from all
mapping hits, and a number of features were computed: the conserved PFAM v33.1
domains, using HMMscan (Eddy 2011) and DAMA (Bernardes et al. 2015), the presence of
the peptide signal, using SignalP v5.0b (Emanuelsson et al. 2007), and the R
implementation of TMHMM (Krogh et al. 2001) to predict the number of transmembrane
helices (TMHs). At each locus, the best annotation was therefore selected based on the
optimal protein length, conserved domain length and score, presence of P-signal and best
number of TMHs. The amino acid sequences were then aligned using CLUSTALW v2.1
(settings: dnamatrix = IUB; gapopen = 10; gapext = 0.1; gapdist = 10; iteration = TREE;
numiter = 1000; clustering = NJ), and the phylogeny was inferred using approximate
Maximum Likelihood search as implemented in FastTree v2.1.11 SSE3, using Le-Gascuel
(2008) model with pseudocounts and the slow exhaustive search algorithm to search for
neighbor-joining (Price et al. 2010). To root the tree, ORco was used as an outgroup, and
gene orthology was subsequently assigned based on the phylogenetic tree or reference
genes in Cicconardi et al. (2024). For the graphical representation of the phylogenetic tree,
we adopted TreeViewer (Bianchini & Sánchez-Baracaldo 2024). Due to the large number of
sequences, we collapsed all monophyletic groups of orthologous genes, showing their
relationships, while the full phylogenetic tree is available in the supplementary materials
(Figure S1).
301
Statistics
All neuropil volumes were $\log_{10}$-transformed prior to analysis, and all analyses were run in R
(v4.4.1; R Core Team, 2024). Using a recent phylogenetic tree of the Heliconiini (Couto et
al., 2023) to control for phylogenetic effects, we ran a series of phylogenetic generalized
linear mixed models (GLMM) with Gaussian distributions using the R package MCMCglmm v
2.32 (Hadfield, 2010). We assigned species as a random effect and tested whether the
scaling relationship between the i) the total antennal lobe volume, ii) antennal lobe hub
volume, and iii) the total volume of the glomeruli, and the volume of the rest of the central
brain (rCBR) changes across broad phylogenetic groups within the Heliconiini. Specifically,
we tested for differences between a) *Heliconius* and non-*Heliconius* Heliconiini, b) between
Heliconiini genera, and c) between sub-clades within *Heliconius* (excluding outgroups),
including the effects of sex. We also ran a fourth model to test for shifts in the scaling
relationship between the glomeruli and the antennal lobe hub across the three different
phylogenetic groupings noted above. To determine to what extent the volumetric differences
observed can be accounted for by phylogeny, we additionally calculated the posterior mean

of Pagel's lambda (λ), after running a null GLMM model, with no other variables other than

the allometric control rCBR or antennal lobe hub. We subsequently conducted Likelihood

ratio testing of a model with the observed lambda and a model that assumes phylogenetic

independence, as described in Münkemüller et al. (2012).

To test for species-specific differences, we also conducted a pairwise comparison of

the 29 species with more than 5 individuals using the posterior distribution of the random

effects. Here we adjusted the significant threshold to 0.99 to account for false positives when

conducting multiple pairwise comparisons. Finally, we explored associations between the

antennal lobe, antennal lobe hub, and glomeruli (after correcting for allometry), as well as

the scaling relationship between antennal lobe hub and glomeruli volume, with

environmental and life history variables. Associations between the neuropil with

environmental and ecological factors were tested separately by taking the best fitting of

these MCMCglmm models and then including these factors individually. Throughout, model

checking for convergence was done using the *gelman.diag* function and for auto-correlation

using the *autocorr* function in MCMCglmm, in addition to visually inspecting the trace plots.

All models were run for 500,000 iterations, with a burn-in of 10,000 and a thinning factor of

500.

We repeated the above analyses to i) test for an association between the number of

olfactory receptors annotated for each Heliconiini species and the same three neuropils

(antennal lobe, antennal lobe hub and glomeruli), and ii) test for association between the

number of olfactory receptors and environmental and ecological factors. We referenced the

olfactory receptor data from a previous study (Cicconardi et al. 2023) and controlled for the

quality of the annotation by including the N50 value. We also repeated the analyses with an

alternative control, specifically the Benchmarking Universal Single-Copy Orthologues

(BUSCO) score (Simão et al., 2015). When using the BUSCO as the alternative control, we

included only species with a BUSCO completeness score of more than 90%. All N50 values,

BUSCO scores and BUSCO completeness score were similarly referenced from Cicconardi

et al. (2023).

To test for potential developmental plasticity, we compared the volume of the total

antennal lobe, the antennal lobe hub, and glomeruli, between wild-caught and insectary-

reared individuals for 5 *Heliconius* species (*H. erato cyrbia, H. melpomene, H. ismenius, H.*

cydno and *H. hecale*). We first tested for differences using a series of nested linear mixed

models implemented in the R package lme4 (Bates et al., 2015). In each case, $\log_{10}$

transformed values were used as the dependent variable. We included the rest of the central

brain (rCBR), species and origin (wild-caught vs insectary-reared) as fixed factors. We also

included an interaction term between species and origin in our full model. Sex was included

as a random factor in all models. To acquire the minimum adequate model (MAM), and to

determine the significance of individual terms, we used Likelihood Ratio Tests. All p-values

were subsequently adjusted with Bonferroni correction to account for multiple testing.

We next conducted standardized major axis regression analyses using the SMATR

package in R (Warton et al., 2012) to identify specific pairwise species differences in the

volume of the antennal lobe, the antennal lobe hub and glomeruli, as well as differences the

scaling relationship between antennal lobe hub and glomeruli volume in wild-caught and

insectary-reared individuals. Following the allometric relationship $y=\beta \log x + \alpha,$ we first

tested for significant shift in slope (β), which would suggest a change in the allometric

scaling exponent between the groups (Kruska, 2005). If a common slope was found, we

subsequently tested for a shift in elevation (α), or a major-axis shift. Elevation shifts (or

"grade-shifts") suggest non-allometric divergence with a change in the intercept (Kruska,

2005), while a major-axis shift suggests that the groups change along a shared allometry

which underlies changes in neuropil volume in concert with the rest of the brain (Barton &

Harvey, 2000; Montgomery et al., 2016). Correction for multiple comparisons was done

using a Sidak-correction, available as an option in the *sma* function of the SMATR package.

3. Results

Variation in antennal lobe investment is driven by species-level differences rather

than broad phylogenetic trends.

Across all individuals, we saw considerable variation in the raw volumes of the antennal lobe

(AL), which varied 6.2-fold across Heliconiini (from ~$3.8 \times 10^6 \mu m^3$ to ~$23.5 \times 10^6 \mu m^3$). This was

reflected in variation in both the total glomeruli volume (GL) (6.7-fold from ~$3.2 \times 10^6 \mu m^3$ to

~$21.5 \times 10^6 \mu m^3$), and especially in the volume of the antennal lobe hub (ALH) (17-fold from

~$3.2 \times 10^5 \mu m^3$ to ~$5.6 \times 10^6 \mu m^3$). However, we did not find evidence for significant variation

between *Heliconius* and non-*Heliconius,* in any of the four models tested (pMCMC>0.05)

(Fig. 2), or between narrower phylogenetic groupings (i.e. outgroup genus or within

Heliconius subclades) (pMCMC>0.05). Our analysis of phylogenetic signal suggests that the

variation that we see for the three neuropils of interest (AL, $\lambda=0.710$; ALH, $\lambda=0.671$ and GL,

$\lambda=0.698$), as well as the relationship between glomeruli and antennal lobe hub volumes ($\lambda=$

0.605), are only moderately influenced by phylogeny. Likelihood ratio testing comparing a

model with and without phylogeny for all three neuropils and the fourth model showed that

only the antennal lobe volume and the glomeruli volume have phylogenetic signal present

(p<0.05). This suggest that the antennal lobe and glomeruli volumes are more strongly

constrained by evolutionary history, whereas the antennal lobe hub volume and the

relationship between glomeruli and antennal lobe hub volumes appear to be more influenced

by other factors, such as ecological or functional constraints. In general, there were no

interactions with sex detected in any of the four models (pMCMC>0.05), suggesting that the
antennal lobe and antennal lob hub volumes of males and females are largely similar.
However, we did find a steeper relationship between glomeruli volume and the antennal lobe
hub volume in males than females (pMCMC=0.018).
Despite the lack of broad phylogenetic trends, pairwise comparisons revealed a
number of differences between species (Table S1A-D). Due to sample size, these analyses
were largely restricted to comparisons within *Heliconius* and some *Eueides* species for all 4
comparisons (i.e. AL, ALH, GL, and GL to ALH volume; Table S1A-D). Comparing the
posterior distribution of each species' random effects with each other, we noted that a few
species constantly showed high probability that their posterior distribution is significantly
different than the other species. This means that if the random effect for species A is greater,
species A will have a higher baseline value for the trait bring measured compared to others.
For example, *E. lybia* emerged as a clear outlier in the pairwise comparisons of individual
antennal lobe structures, and in the scaling relationship between glomeruli and antennal
lobe hub volumes (Fig. 2). We also noted that *E. lybia* had a greater total glomeruli volume
than many other Heliconiini species. To test whether this increase in volume is associated
with an increase in the number of glomeruli, we took a subset of five Heliconiini species, two
Heliconius, two *Euiedes* and one *Dryadula* to examine if *E. lybia* had pronounced increases
in glomeruli number. The number of glomeruli ranges from 65 to 87 (Table S2). Of the five
species, *Dryadula phaetusa* had the greatest number of glomeruli (85-87), followed by *E.*
lybia (82-83). However, *E. lybia* has the largest glomeruli volume (from $\sim 13 \times 10^6 \mu m^3$ to
$\sim 18.\times 10^6 \mu m^3$), and *Dryadula phaetusa* (from $\sim 10.8 \times 10^6 \mu m^3$ to $\sim 12.6 \times 10^6 \mu m^3$) had similar
glomeruli volume to *H. erato* (from $\sim 11.5 \times 10^6 \mu m^3$ to $\sim 12.4 \times 10^6 \mu m^3$), despite the increased
number of glomeruli (Table S2). *H. hecale* had the smallest number of glomeruli (68) and the
smallest glomeruli volume (from $\sim 7.5 \times 10^6 \mu m^3$ to $\sim 9.5.\times 10^6 \mu m^3$) (Table S2).
416
Diversity of olfactory receptors across Heliconiini
As the number of glomeruli is thought to be directly related to the number of functional
olfactory receptors, we fully annotated olfactory receptor genes (ORs) in 58 Heliconiini and
14 other Nymphalids genomes. In total we were able to annotate 4,658 genes (Table S3),
which clustered into ~70 orthologous groups (OGs) (Fig. 3), 21 of them specific to different
subclades within Nymphalids. We recorded 10 distinct orthologous group duplications at the
stem of Heliconinae (51, 40, 35a, 35b, 05, 04a, 04b, 04c, 56a and 56b), and five at the stem
of Heliconiini (24, 24b, 05a, 05b and 56c) (Fig. 3, 4). Interestingly, OR05, belonging to the
pheromone receptor clade, was duplicated once at the stem of Heliconinae and twice more
at the stem of Heliconiini, while OR56, possibly linked to plant volatile compounds (Bastin-
Héline et al. 2019), was found to be duplicated twice in Heliconinae and again,

independently, in Heliconiini. As previously described (Cicconardi et al. 2023), we did not
detect any duplications at the stem of *Heliconius*, also supporting previous findings that the
great majority of duplications happened at base the of the tribe and subtribe radiation
(Cicconardi et al. 2023). We also confirm three putative orthologous groups deletions of
OR15, OR27, and OR35.

Overall, *Heliconius* species have on average ~65 ORs per species (excluding
species with low quality assemblies), broadly within the range of other Heliconiini (and
average 68 ORs) including *Agraulis* and *Dione* genera species (62 ORs). *H. aoedes*, a non-
pollen feeding *Heliconius* species with 66 ORs, was also within range of other *Heliconius*.
Despite this, within more closely related clades there are distinct outliers with higher
numbers of ORs. For example, *H. pardalinus butleri* and *H. timareta* had 89 and 90 ORs,
respectively, while *Eueides* species with the best quality genomes (*E. isabella* and *E. tales*)
have 73 and 81 ORs, respectively. Of particular note is *Dryadula phaetusa*, which has 95
ORs, the largest repertoire within the whole dataset, which is in line with their high number of
glomeruli. The OR expansion in *Dryadula* is driven largely by a large expansion of a few
orthologous groups; OR04 (25 copies), OR41 (7 copies), and OR56 (from the pheromone
clade; 11 copies).

Our MCMCglmm analyses, including the N50 value to account for genome
completeness, did not show any significant relationships between OR number and the
volumes of the three different neuropils (AL/ALH/GL) (Table S4). We obtained the same
results when repeating the analyses with BUSCO completeness scores, instead of N50.
Next, to explore whether the number of glomeruli is related to the number of olfactory
receptors, we looked at the subset of data of four Heliconiini species in which all the
individual glomeruli were segmented and counted. We observed that in three of the five
species, the number of olfactory receptors was smaller than the number of glomeruli.
However, two of these species (*E.lybia and E.aliphera*) had a BUSCO completeness score
of less than 90%, which could have led to an underestimation of the number of olfactory
receptors. Disregarding these 2 species, there was a disparity of ±5-10 between the
olfactory receptor numbers and glomeruli numbers in the remaining 3 species (Table S2).
457

Major environmental and ecological factors do not explain variation in antennal lobe
morphology across the tribe
We found no significant association between any of the olfactory neuropils tested and the
number of host plants used, the degree of social roosting, or the presence of pollen-feeding
behaviours (Table S5). We also tested for associations with environmental factors that have
been argued to affect the odorant concentration in the air, using datasets of annual
temperature and annual precipitation. Once again, we did not find any significant

associations between these variables and antennal lobe, antennal lobe hub, and total

glomeruli volumes, or with the scaling relationship between the antennal lobe hub and total

glomeruli volumes (Table S5). We further repeated these analyses with number of olfactory

receptors as the independent variable, and again our results showed no significant

associations (Table S5).

Rearing environment influences relative volumes of the antennal lobe hub and

glomeruli, but not overall antennal lobe volume

Our broad phylogenetic comparison of antennal lobe morphology took advantage of wild

caught individuals. Although the differences observed might reflect heritable variation, they

might also be the result of plastic responses to the environment. To test for potential effects

of the local environment, we used a subset of five species from our wild dataset for which

imaging was also available for insectary-reared individuals. We found that the interaction

between species and origin (i.e. insectary-reared vs wild) was not significant in our

comparison of antennal lobe volumes between wild and insectary individuals ($\chi2_4$=4.127,

P_{adj}=1.00), suggesting that the volumes between these two groups are similar across

species (Table S6). There was no overall effect of origin on antennal volume ($\chi2_1$ =4.643,

P_{adj}=0.094), and only species ($\chi2_4$=12.341, P_{adj}<0.05) and rCBR (as an allometric control)

($\chi2_1$ =14.628, P_{adj}<0.05) were retained in our minimum adequate model (MAM). Subsequent

analyses using SMATR on species-specific data showed no difference in the overall

antennal lobe volume between wild and insectary individuals, with the exception of *H.*

melpomene (significant slope shift; LR_1=4.999, P_{adj}<0.05), and *H. hecale* (significant grade

shift; Wald $\chi2_1$ =4.505, P_{adj}<0.05).

We next examined variation in the two constituent parts of the antennal lobe, the

antennal lobe hub and total glomeruli volumes, and once again the interaction between

species and origin was not significant (ALH: $\chi2_4$=3.453, P_{adj}=1; GL: $\chi2_4$ =2.425, P_{adj}=1.000).

In both cases, species was not retained in the MAM, but there was a significant effect of

origin (ALH: $\chi2_1$ =68.033, P_{adj}<0.001; GL: $\chi2_1$ =41.297, P_{adj}<0.001). Subsequent major axis

regressions focused on the antennal lobe hub revealed that all differences that were

observed among wild-caught and insectary-reared individuals in the five species tested were

a result of significant grade shifts (α) or significant major-axis shifts, with the slopes being

conserved (Table S7; Fig. S2). This pattern was similarly seen in the total glomeruli volume,

with the exception of two species, *H. ismenius and H. hecale* (Table S7; Fig. S2).

Surprisingly, although we observed volumetric differences between the wild and

insectary-reared individuals for both the antennal lobe hub and total glomeruli volume, these

were not in the same direction. Specifically, although the antennal lobe hub was larger in

insectary-reared individuals, the total glomeruli volume was smaller (Fig. 5). To investigate

this relationship further we directly compared these two components of the antennal lobe. In
our model of total glomeruli volume as a function of antennal lobe hub volume, neither the
interaction between species and origin or species was significant (species*origin: $\chi2_4$=2.084,
P_{adj}=1.000, species: $\chi2_4$=11.435, P_{adj}=0.066), suggesting that different species do not
respond differently to captivity. However, there was an overall effect of origin ($\chi2_1$ =33.037,
P_{adj}<0.001). When looking at the scaling relationship between the total glomeruli volume
and antennal lobe hub volume, we again observed a common slope, but an overall
significant grade shifts across all five species (Fig. 5).
510
511 **Discussion**
512
Groups of closely related, but ecologically diverse species provide an opportunity to test
hypotheses about divergence in sensory and neural systems (Montgomery et al., 2021;
Montgomery & Merrill, 2017; Wainwright & Montgomery, 2022). We examined variation in
different components of the olfactory system across the Heliconiini tribe of Neotropical
butterflies. Unlike previous studies in this group that have focused on a limited number of
species pairs within this group (Montgomery et al., 2021; Montgomery & Merrill, 2017;
Hebberecht et al., 2023), this allowed us to assess relationships between components of the
olfactory system, and how these may correlate with ecological or environmental traits at a
macroevolutionary scale. Our results reveal a mix of conserved and dynamic patterns of
olfactory evolution, with species-specific shifts in antennal lobe morphology and olfactory
receptor diversity occurring against a backdrop of broader phylogenetic stability. Notably, the
variation in antennal lobe size across species appears to be driven by independent shifts in
its two major components, the glomeruli and the antennal lobe hub, which likely reflect
different selective pressures acting on these structures, and some independent variation
between total glomeruli volume and OR number. We also uncovered a surprising pattern of
developmental plasticity, with insectary-reared individuals exhibiting larger antennal lobe
hubs but smaller glomeruli volumes compared to their wild-caught counterparts, suggesting
complex environmental influences on the maturation of the olfactory system.
Although our results suggest that the architecture of the olfactory system is broadly
conserved across the Heliconiini, we did observe substantial shifts in antennal lobe size for
certain species. For example, *Eueides lybia* frequently emerged as an outlier in our
analyses, displaying an overall increase in the antennal lobe, as well as significantly larger
volumes of both its components, the glomeruli and antennal lobe hub volume. These
findings might suggest enhanced olfactory discrimination capabilities linked to ecological
demands, but current knowledge about *Eueides* provides little insight into the specific factors
driving this difference. Similarly, for most species across the Heliconiini, we also see that

investment into the glomeruli and the antennal lobe hub are mostly equal, again except for a

few outlier species (e.g *Eueides lybia*), which suggest a general correlation of growth for the

glomeruli and antennal lobe hub.

In addition to changes in the antennal lobe, we also found species-specific shifts in

olfactory receptor number (e.g. *E. tales, E. isabella*, *H. timareta* etc.). *Dryadula phaetusa*

exhibited the largest olfactory receptor repertoire in our dataset, driven by expansions in

specific olfactory receptor subgroups, including those potentially linked to pheromone

detection and plant volatile cues. Although further investigation of this interesting pattern is

outside the scope of this study, we do also note that we distinguished ammino acid

sequences having, lacking or partially lacking the PFAM domain for the 7-TMHs (7tm_6).

Dryadula has 62 ORs with a complete conserved domain, in line with other Heliconiini, but

has 28 loci with a shorter domain at the N-terminus. But only ~10 of them are actually

shorter than ~300aa and have less than 5 TMHs; further analyses could evaluate the extent

of which this lack of domain conservation is a limitation of the PFAM HMM or a signature of

pseudogenisation.

Elsewhere, as with changes in antennal lobe volume, shifts in olfactory receptor

numbers are difficult to ascribe to specific ecological events. The expansion of olfactory

receptors in *H. timareta*, however, is intriguing given what we know of its ecology. Within a

sympatric community, closely related *Heliconius* tend to differ in warning pattern, and it is

well established that males often use colour to identify suitable mates (Merrill et al., 2011,

2015). This is certainly true for *H. cydno* (white or yellow) and its close relative *H.*

melpomene (red), which coexist to the west of the Eastern Cordillera in the Andes (Merrill et

al, 2019). However, on the eastern slopes of the Andes *H. cydno* is replaced by its sister

species *H. timareta*, which shares the red patterns of the local *H. melpomene.* Interestingly,

while neither species can discriminate based on colour alone, *H. timareta* (but not *H.*

melpomene) males preferentially court *H. timareta* over *H. melpomene* females, presumably

as a result of olfactory cues (Mérot et al., 2015; Rossi et al., 2024). As such, it is tempting to

hypothesise that the expansion of olfactory receptors in *H. timareta* may have arisen as a

result of a reinforcement-like process, to aid mate discrimination and avoid the production of

unfit hybrids.

Overall, our analyses revealed little evidence to illuminate broader ecological

correlates of species-specific expansions and reductions of the antennal lobe (and/or its

constituent structures), or olfactory receptors. Broader trends in antennal lobe size evolution

reported across the Lepidoptera have often revealed correlations with major shifts in activity

pattern, and in particular diurnal versus nocturnal lifestyles (Stöckl et al., 2016; O'Donnell et

al., 2013; Montgomery & Ott, 2015). In contrast, all Heliconiini species are diurnal, and

Heliconius butterflies, in particular, are reported to predominantly forage for nectar and

pollen in the early morning (Gilbert, 1975; Barp et al., 2011; Dell'Aglio et al., 2024b).

Although we cannot discount undiscovered differences in diurnal activity across populations

and species, evidence for this has not been found in captive populations (Dell'Aglio et al.,

2024b). Although light intensity in closed forests will vary throughout the day, it will never be

as low as at night, where light levels drop by eight orders of magnitude (Warrant, 2017). As

such, it is perhaps unlikely that this difference necessitates significant investment in olfactory

processing. Similar logic may also explain our failure to detect broad correlates between

evolution in olfactory structures and ecology. Indeed, while shifts in antennal lobe volume

have been reported among closely related *Heliconius* species inhabiting different forest

types (Montgomery & Merrill, 2017), there is little evidence that this association is repeated

across different species pairs, in contrast to the changes in the optic lobe (Montgomery et

al., 2021, Rivas-Sánchez et al.; 2024). Indeed, across more diverse communities of

ithomiine butterflies, variation in the visual system is more extensive and more closely tied to

variation in habitat use than the olfactory system (Wainwright et al., 2024).

The species-specific variation we see in antennal lobe volume could stem from

changes in the antennal lobe hub, or from changes in glomeruli volume. Overall, our

analyses suggest that variation in antennal lobe size across species can result from both

differential investment in glomeruli volume (6.7-fold absolute change) and antennal lobe hub

volume (17-fold absolute change), however the range of variation in glomeruli volume is

more similar to that for overall brain size (7.7-fold absolute change). While in some cases

changes in the antennal lobe hub and glomeruli are correlated, this was not always the case,

suggesting that these two components can evolve semi-independently. For example, *E. lybia*

and *E. isabella* stood out with large glomeruli volumes, while other species like *H. erato*

cyrbia and *H. himera* show disproportionately larger antennal lob hub volumes relative to

glomeruli.

An increase in overall glomeruli volume can result from two factors, larger individual

glomeruli or an increase in glomeruli number. Larger individual glomeruli would suggest an

increase in olfactory receptor neurons housed within each glomerulus. Because studies

have linked glomerular volume to olfactory receptor neuron number, this expansion may

enhance sensitivity to olfactory cues (Takagi et al., 2024; Linz et al., 2013; Meisami, 1989),

and *Drosophila* species with more but same types of olfactory receptor neurons have been

shown to track host odors more efficiently and persistently (Takagi et al., 2024). Alternatively,

glomeruli volume may also arise from an increase in glomeruli number, as suggested by our

data on a subset of *Heliconiini* species (see below; Table S2). Since each glomerulus is

expected to process input from a single class of olfactory receptor neuron (Hildebrand &

Shepherd, 1997), an increased number of glomeruli could enhance odor discrimination by

expanding the combinatorial possibilities for olfactory recognition. This has been observed in

hornets, where the region of the antennal lobe with the most extensive variation (T_B Cluster)
also had the highest glomeruli count, likely due to the greater demand for olfactory
discrimination in high-density environments (Couto et al., 2021).
Changes in antennal lobe volume may also be driven by variation in antennal lobe
hub volume. For instance, previous research found that differences in antennal lobe size
between the glasswing butterfly (*Godyris zavaleta*) and two *Heliconius* species (*H. erato* and
H. hecale) were largely due to expansion of the antennal lobe hub in *Godyris* rather than
changes in glomeruli size (Montgomery et al., 2016). Since the antennal lobe hub contains
axons of interneurons that integrate and relay olfactory information to integrative centres, its
expansion could reflect differences in how olfactory signals are integrated across glomeruli
and processed (Montgomery et al., 2016). In particular, a larger antennal lobe hub may
enhance signal integration or prioritize specific odor types, potentially influencing behavioral
responses.
The variation in glomeruli number in our dataset contrasts with the relatively constant
glomeruli count observed across Lepidoptera, which typically ranges between 60 and 70
(Boeckh & Boeckh, 1979; Rospars, 1983; Berg et al., 2002; Kazawa et al., 2009; Carlsson et
al., 2013), albeit with some exceptions, such as the oligophagous moth genus *Cydia*, which
only has around 50 glomeruli (Varela et al., 2009; Couton et al., 2009; Trona et al., 2010). In
contrast to this apparent consistency, we found that glomeruli number ranged from 65 to 87
in the five sampled Heliconiini species (Table S2). Although measurement errors and the
lack of clear boundaries between individual glomeruli may have led to slight under- or
overestimation, the number of glomeruli in our dataset aligns relatively well with the range of
odor receptors identified in a much larger sample of Heliconiini. While the higher number of
odor receptors compared to glomeruli in *Dryadula phaetusa* could be attributed to the
expansion of certain odor receptor groups, we also acknowledge the possibility of functional
redundancy or divergence in these butterflies. This could result in a deviation from the
traditional 1:1 model of olfactory coding observed in other insects (Couto et al., 2005).
However, future studies could expand the glomeruli counts to more species in the Heliconiini
to get a more robust association between olfactory receptor number and glomeruli number,
and single cell sequencing methods could provide insights into possible redundancy and co-
expression of ORs in single sensory neurons.
Our analyses also reveal intriguing patterns of developmental plasticity. Insectary-
reared individuals displayed larger antennal lobe hub volumes, but reduced glomeruli
volumes compared to their wild-caught counterparts. This leads to a significant grade shift in
the relationship between glomeruli and antennal lobe hub volumes between wild and
insectary bred individuals. This divergent pattern suggests that the absence of natural
environmental stimuli in captivity may differentially influence the development of these two

structures. While the biological implications of these changes remain unclear, they raise

questions about the potential trade-offs in sensory investment under constrained

environments. It is possible that some of the interspecific variation we observe in our larger

sampling of wild individuals reflect environmentally induced effects. However, the lack of

current ecological correlates makes it unlikely that this would be driven by consistent

environmental cues, although we cannot reject the possibility. However, examining olfactory

receptor numbers in three species that differ greatly in terms of their antennal lobe hub (*H.*

himera), glomeruli (*H. sara*) and glomeruli to antennal lobe hub volume (*H. aoede*) from

other species — and accounting for genome completeness scores — we found that their

olfactory receptor counts (*H. himera*: 64, *H. sara*: 70, *H. aoede*: 66) were within or close to

the average for Heliconiini (68 ORs). This suggests that, despite their neuroanatomical

differences, these species likely have similar capacities to process olfactory information,

reducing the likelihood that their divergence is driven by evolutionary plasticity, although

olfactory receptors turnover may result in different olfactory tuning. Expanding comparative

studies of wild versus insectary-reared individuals to include species we identify as outliers

would help eliminate the possibility of evolutionary plasticity. Additionally, experimental

studies measuring olfactory sensitivity and behavioural outcomes in both wild and captive

derived individuals could provide valuable insights into the functional consequences of the

plastic responses in the glomeruli and antennal lobe hub.

Overall, our findings have important implications for future studies that use antennal

lobe size as a proxy for olfactory capabilities across species. Our study demonstrates that

glomeruli and antennal lobe hub volumes can evolve independently, with the environment

acting on them in unexpected and even opposing ways. Consequently, total antennal lobe

size might not be a straightforward indicator of olfactory abilities, underscoring the need to

examine the functional contributions of glomeruli and antennal lobe hub separately in

evolutionary and ecological contexts. In the Heliconiini, olfactory evolution is shaped by

species-specific adaptations and developmental plasticity. Modular adjustments within the

antennal lobe—for example, differential investment in glomeruli versus antennal lobe hubs—

provide a flexible framework for rapid adaptation to diverse ecological niches. Although we

did not observe direct associations with specific ecological traits, species-level differences,

such as the large glomerular volume in *Eueides lybia* and the expansion of olfactory

receptors in species like *H. timareta*, suggest that diverse evolutionary pressures are at play,

warranting further investigation into unmeasured ecological or behavioural factors. Notably,

our identification of a curious pattern of developmental plasticity—where insectary-reared

individuals exhibited larger antennal lobe hubs but smaller glomeruli volumes compared to

their wild-caught counterparts—highlights the influence of environmental conditions on

olfactory system development and raises questions about how such plasticity may affect

sensory processing and behaviour. Together, these results emphasize the importance of
integrating evolutionary, ecological, and developmental perspectives when investigating
olfactory evolution.
690

691 **Data availability**
692 The underlying data and R-scripts supporting the findings of this study are available as part
693 of the supplementary data "Heliconiini_olfactory_evolution_supplementary_tables.xlsx" and
694 on github (https://github.com/SpeciationBehaviour/olfactory_system_evolution_heliconiini)
695 respectively.
696

Acknowledgements
We thank the environmental ministries of Costa Rica, Panama, French Guiana, Ecuador and
Peru for permission to collect and export the sample in previous studies. We are grateful to
the Wolfson Bioimaging Centre, University of Bristol, and the Center for Advanced Light
Microscopy at the University of Munich Biocenter for imaging assistance. This work was
supported by NERC Independent Research Fellowship NE/N014936/1 (SHM), and ERC
Starter Grants 758508 (SHM) and 851040 (RMM).
704

705 **Conflict of interest statement**
706 The authors declare no conflict of interest.
707

708 **Role of authors**
709 Conceptualisation: SHM, RMM; Data curation: YP, FC, SHM; Investigation and Methodology:
710 YP, FC, SHM; Writing – original draft: YP, FC; Writing – review and editing; YP, FC, GB,
711 RMM, SHM. Visualisation: YP, FC, GB; Funding acquisition, Project Administration and
712 Supervision: SHM, RMM.
713

References
Aiello-Lammens, M. E., Boria, R. A., Radosavljevic, A., Vilela, B. and Anderson, R. P. (2015).
spThin: an R package for spatial thinning of species occurrence records for use in
ecological niche models. *Ecography* 38, 541–545.
https://doi.org/10.1111/ecog.01132.
Arnegard, M. E., McIntyre, P. B., Harmon, L. J., Zelditch, M. L., Crampton, W. G., Davis, J.
K., Sullivan, J. P., Lavoué, S. and Hopkins, C. D. (2010) 'Sexual signal evolution
outpaces ecological divergence during electric fish species radiation', *Am. Nat.*,
176(3), pp. 335–356. Available at: https://doi.org/10.1086/655221.

Arnold, G., Masson, C. and Budharugsa, S. (1985). Comparative study of the antennal lobes

and their afferent pathway in the worker bee and the drone (*Apis mellifera*). *Cell*

Tissue Res. 242, 593–605. https://doi.org/10.1007/BF00225425.

Auer, T. O., Khallaf, M. A., Silbering, A. F., Zappia, G., Ellis, K., Álvarez-Ocaña, R., Arguello,

J. R., Hansson, B. S., Jefferis, G. S. X. E., Caron, S. J. C. et al. (2020). Olfactory

receptor and circuit evolution promote host specialization. *Nature* 579, 402–408.

https://doi.org/10.1038/s41586-020-2073-7.

Barp, E. A., Soares, G. L. G., Giani, E. J. M., Rodrigues, D. and Moreira, G. R. P. (2011).

Variation in nectar and pollen availability, sucrose preference, and daily response in

the use of flowers by *Heliconius erato phyllis*. *J. Insect Behav.* 24, 200–219.

https://doi.org/10.1007/s10905-010-9248-2.

Barton, R. A. (1998). Visual specialization and brain evolution in primates. *Proc. R. Soc.*

Lond. B Biol. Sci. 265, 1933–1937. https://doi.org/10.1098/rspb.1998.0523.

Barton, R. A. and Harvey, P. H. (2000). Mosaic evolution of brain structure in mammals.

Nature 405, 1055–1058. https://doi.org/10.1038/35016580.

Barton, R. A., Purvis, A. and Harvey, P. H. (1995). Evolutionary radiation of visual and

olfactory brain systems in primates, bats and insectivores. *Philos. Trans. R. Soc.*

Lond. B Biol. Sci. 348, 381–392. https://doi.org/10.1098/rstb.1995.0076.

Bates, D., Mächler, M., Bolker, B. and Walker, S. (2015). Fitting linear mixed-effects models

using lme4. *J. Stat. Softw.* 67, 1–48. https://doi.org/10.18637/jss.v067.i01.

Beltrán, M., Jiggins, C. D., Brower, A. V. Z., Bermingham, E. and Mallet, J. (2007). Do pollen

feeding, pupal-mating and larval gregariousness have a single origin in *Heliconius*

butterflies? Inferences from multilocus DNA sequence data. *Biol. J. Linn. Soc.* 92,

221–239. https://doi.org/10.1111/j.1095-8312.2007.00830.x.

Berg, B. G., Galizia, C. G., Brandt, R. and Mustaparta, H. (2002). Digital atlases of the

antennal lobe in two species of tobacco budworm moths, the oriental *Helicoverpa*

assulta (male) and the American *Heliothis virescens* (male and female). *J. Comp.*

Neurol. 446, 123–134. https://doi.org/10.1002/cne.10180.

Berg, B. G., Zhao, X.-C. and Wang, G. (2014). Processing of pheromone information in

related species of heliothine moths. *Insects* 5, 742–761.

https://doi.org/10.3390/insects5040742.

Bisch-Knaden, S., Dahake, A., Sachse, S., Knaden, M. and Hansson, B. S. (2018). Spatial

representation of feeding and oviposition odors in the brain of a hawkmoth. *Cell Rep.*

22, 2482–2492. https://doi.org/10.1016/j.celrep.2018.01.082.

Bloch, N. I. (2015). Evolution of opsin expression in birds driven by sexual selection and

habitat. *Proc. R. Soc. B Biol. Sci.* 282, 20142321.

https://doi.org/10.1098/rspb.2014.2321.

Boeckh, J. and Boeckh, V. (1979). Threshold and odor specificity of pheromone-sensitive
neurons in the deutocerebrum of *Antheraea pernyi* and *A. polyphemus* (Saturnidae).
J. Comp. Physiol. 132, 235–242. https://doi.org/10.1007/BF00614495.

Boeckh, J., Boeckh, V. and Kühn, A. (1977) 'Further data on the topography and physiology
of central olfactory neurons in insects', *Olfact. Taste*, 6, pp. 315–321.

Boeckh, J., Distler, P., Ernst, K. D., Hösl, M. and Malun, D. (1990). Olfactory bulb and
antennal lobe, in: Schild, D. (Ed.), Chemosensory Information Processing. Springer
Berlin Heidelberg, Berlin, Heidelberg, pp. 201–227.

Borrero, J., Mogollon, P.E., Wright, D.S., Lozano-Urrego, D., Rueda-Muñoz, G., Pardo-Diaz,
C., Salazar, C., Montgomery, S.H., Merrill, R.M. (2024). Weighting of sensory cues
reflect changing patterns of visual investment during ecological divergence in
Heliconius butterflies. *Biol. Lett*. 20, 20240377. http://doi.org/10.1098/rsbl.2024.0377

Boughman, J. W. (2002). How sensory drive can promote speciation. *Trends Ecol. Evol.* 17,
571–577. https://doi.org/10.1016/S0169-5347(02)02595-8.

Briscoe, A. D., Macias-Muñoz, A., Kozak, K. M., Walters, J. R., Yuan, F., Jamie, G. A.,
Martin, S. H., Dasmahapatra, K. K., Ferguson, L. C., Mallet, J., et al. (2013). Female
behaviour drives expression and evolution of gustatory receptors in butterflies. *PLoS*
Genet. 9, e1003620. https://doi.org/10.1371/journal.pgen.1003620.

Brown, K. S. (1981). The biology of *Heliconius* and related genera. *Annu. Rev. Entomol.* 26,
427–457. https://doi.org/10.1146/annurev.en.26.010181.002235.

Bruce, T. J. A. and Pickett, J. A. (2011). Perception of plant volatile blends by herbivorous
insects – finding the right mix. *Phytochemistry, Plant-Insect Interactions* 72, 1605–
1611. https://doi.org/10.1016/j.phytochem.2011.04.011.

Burger, H., Ayasse, M., Dötterl, S., Kreissl, S. and Galizia, C. G. (2013). Perception of floral
volatiles involved in host-plant finding behaviour: comparison of a bee specialist and
generalist. *J. Comp. Physiol. A* 199, 751–761. https://doi.org/10.1007/s00359-013-
0835-5.

Carlsson, M., Schäpers, A., Nässel, D. and Janz, N. (2013). Organization of the olfactory
system of Nymphalidae butterflies. *Chem. Senses* 38.
https://doi.org/10.1093/chemse/bjt008.

Cicconardi, F., Di Marino, D., Olimpieri, P. P., Arthofer, W., Schlick-Steiner, B. C., & Steiner,
F. M. (2017). Chemosensory adaptations of the mountain fly Drosophila nigrosparsa
(Insecta: Diptera) through genomics' and structural biology's lenses. *Sci. Rep.* 7,
43770. https://doi.org/10.1038/srep43770

Cicconardi, F., Milanetti, E., Pinheiro de Castro, E. C., Mazo-Vargas, A., Van Belleghem, S.
M., Ruggieri, A. A., Rastas, P., Hanly, J., Evans, E., Jiggins, C. D., et al. (2023).
Evolutionary dynamics of genome size and content during the adaptive radiation of

Heliconiini butterflies. *Nat. Commun.* 14, 5620. https://doi.org/10.1038/s41467-023-
41412-5.

Cicconardi, F., Morris, B. J., Martelossi, J., Ray, D. A. and Montgomery, S. H. (2024). Novel
sex-specific genes and diverse interspecific expression in the antennal
transcriptomes of *Ithomiine* butterflies. *Genome Biol. Evol.* 16, evae218.
https://doi.org/10.1093/gbe/evae218.

Couto, A., Alenius, M., & Dickson, B. J. (2005). Molecular, Anatomical, and Functional
Organization of the Drosophila Olfactory System. *Curr. Biol*. 15, 1535–1547.
https://doi.org/10.1016/j.cub.2005.07.034

Couto, A., Arnold, G., Ai, H. et al. (2021). Interspecific variation of antennal lobe composition
among four hornet species. *Sci Rep.* 11, 20883 https://doi.org/10.1038/s41598-021-
00280-z

Couto, A., Young, F. J., Atzeni, D., Marty, S., Melo-Flórez, L., Hebberecht, L., Monllor, M.,
Neal, C., Cicconardi, F., McMillan, W. O., et al. (2023). Rapid expansion and visual
specialisation of learning and memory centres in the brains of *Heliconiini* butterflies.
Nat. Commun. 14, 4024. https://doi.org/10.1038/s41467-023-39618-8.

Couton, L., Minoli, S., Kiêu, K., Anton, S., Rospars, JP. (2009). Constancy and variability of
identified glomeruli in antennal lobes: computational approach in Spodoptera
littoralis. *Cell Tissue Res*. 337,491–511.

Coyne, J. A. and Orr, H. A. (2004). *Speciation*. Oxford University Press, Incorporated.

Dalton, P. (2000). Psychophysical and behavioral characteristics of olfactory adaptation.
Chem. Senses 25, 487–492. https://doi.org/10.1093/chemse/25.4.487.

Darragh, K., Vanjari, S., Mann, F., Gonzalez-Rojas, M. F., Morrison, C. R., Salazar, C.,
Pardo-Diaz, C., Merrill, R. M., McMillan, W. O., Schulz, S., et al. (2017). Male sex
pheromone components in *Heliconius* butterflies released by the androconia affect
female choice. *PeerJ* 5, e3953. https://doi.org/10.7717/peerj.3953.

Darwin, C. (1859) *On the origin of species by means of natural selection, or the preservation*
of favoured races in the struggle for life. London, UK: Murray.

Dell'Aglio, D. D., Stevens, M. and Jiggins, C. D. (2016). Avoidance of an aposematically
coloured butterfly by wild birds in a tropical forest. *Ecol. Entomol.* 41, 627–632.
https://doi.org/10.1111/een.12335.

Dell'Aglio, D. D., Rivas-Sánchez, D. F., Wright, D. S., Merrill, R. M. and Montgomery, S. H.
(2024a) 'The sensory ecology of speciation', *Cold Spring Harb. Perspect. Biol.*, 16(1),
a041428. Available at: https://doi.org/10.1101/cshperspect.a041428.

Dell'Aglio, D. D., McMillan, O. W. and Montgomery, S. (2024b). Using motion-detection
cameras to monitor foraging behaviour of individual butterflies. *Ecol. Evol.* 14,
e70032. https://doi.org/10.1002/ece3.70032.

Depetris-Chauvin, A., Galagovsky, D., Keesey, I. W., Hansson, B. S., Sachse, S. and
Knaden, M. (2023). Evolution at multiple processing levels underlies odor-guided
behavior in the genus *Drosophila*. *Curr. Biol.* 33, 4771–4785.e7.
https://doi.org/10.1016/j.cub.2023.09.039.

Devaud, J.-M. (2003). Experimental studies of adult *Drosophila* chemosensory behaviour.
Behav. Process. Drosophila behaviour: concepts and tools 64, 177–196.
https://doi.org/10.1016/S0376-6357(03)00134-7.

Devaud, J.-M., Acebes, A. and Ferrús, A. (2001). Odor exposure causes central adaptation
and morphological changes in selected olfactory glomeruli in *Drosophila*. *J. Neurosci.*
21, 6274–6282. https://doi.org/10.1523/JNEUROSCI.21-16-06274.2001.

Dobritsa, A. A., van der Goes van Naters, W., Warr, C. G., Steinbrecht, R. A. and Carlson, J.
R. (2003). Integrating the molecular and cellular basis of odor coding in the
Drosophila antenna. *Neuron* 37, 827–841. https://doi.org/10.1016/s0896-
6273(03)00094-1.

Endler, J. A. (1992). Signals, signal conditions, and the direction of evolution. *Am. Nat.* 139,
S125–S153.

Estrada, C., Schulz, S., Yildizhan, S. and Gilbert, L. E. (2011). Sexual selection drives the
evolution of antiaphrodisiac pheromones in butterflies. *Evolution* 65, 2843–2854.
https://doi.org/10.1111/j.1558-5646.2011.01352.x.

Fick, S. E. and Hijmans, R. J. (2017). WorldClim 2: new 1-km spatial resolution climate
surfaces for global land areas. *Int. J. Climatol.* 37, 4302–4315.
https://doi.org/10.1002/joc.5086.

Fusca, D. and Kloppenburg, P. (2021). Task-specific roles of local interneurons for inter- and
intraglomerular signaling in the insect antennal lobe. *eLife* 10, e65217.
https://doi.org/10.7554/eLife.65217.

Gilbert, L. E. (1972). Pollen feeding and reproductive biology of *Heliconius* butterflies. *Proc.*
Natl. Acad. Sci. U.S.A. 69, 1403–1407. https://doi.org/10.1073/pnas.69.6.1403.

Gilbert, L. E. (1975). Coevolution of animals and plants, in: *Gilbert, L. E., Raven, P. H.*
(Eds.), Symposium V, First International Congress of Systematic and Evolutionary
Biology, 1973. University of Texas Press, pp. 210–240.
https://doi.org/10.7560/710313-011.

Gilbert, L. E. (1976). Postmating female odor in *Heliconius* butterflies: A male-contributed
antiaphrodisiac? *Science* 193, 419–420. https://doi.org/10.1126/science.935877.

González-Rojas, M. F., Darragh, K., Robles, J., Linares, M., Schulz, S., McMillan, W. O.,
Jiggins, C. D., Pardo-Diaz, C. and Salazar, C. (2020). Chemical signals act as the
main reproductive barrier between sister and mimetic *Heliconius* butterflies. *Proc. R.*
Soc. B Biol. Sci. 287, 20200587. https://doi.org/10.1098/rspb.2020.0587.

Gronenberg, W. and Hölldobler, B. (1999). Morphologic representation of visual and
antennal information in the ant brain. *J. Comp. Neurol.* 412, 229–240.
https://doi.org/10.1002/(SICI)1096-9861(19990920)412:2

Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models:
The MCMCglmm R package. *J. Stat. Softw.* 33, 1–22.
https://doi.org/10.18637/jss.v033.i02.

Hallem, E. A. and Carlson, J. R. (2006). Coding of odors by a receptor repertoire. *Cell* 125,
143–160. https://doi.org/10.1016/j.cell.2006.01.050.

Hansson, B. S. and Anton, S. (2000). Function and morphology of the antennal lobe: New
developments. *Annu. Rev. Entomol.* 45, 203–231.
https://doi.org/10.1146/annurev.ento.45.1.203.

Hansson, B. S., Christensen, T. A. and Hildebrand, J. G. (1991). Functionally distinct
subdivisions of the macroglomerular complex in the antennal lobe of the male sphinx
moth *Manduca sexta*. *J. Comp. Neurol.* 312, 264–278.
https://doi.org/10.1002/cne.903120209.

Hansson, B. S., Ljungberg, H., Hallberg, E. and Löfstedt, C. (1992). Functional specialization
of olfactory glomeruli in a moth. *Science* 256, 1313–1315.
https://doi.org/10.1126/science.1598574.

Hansson, B. S. and Stensmyr, M. C. (2011). Evolution of insect olfaction. *Neuron* 72, 698–
711. https://doi.org/10.1016/j.neuron.2011.11.003.

Hausmann, A. E., Freire, M., Alfthan, S. A., Kuo, C.-Y., Linares, M., McMillan, O., Pardo-
Diaz, C., Salazar, C. and Merrill, R. M. (2023). Does sexual conflict contribute to the
evolution of novel warning patterns? *J. Evol. Biol.* 36, 563–578.
https://doi.org/10.1111/jeb.14151.

Hebberecht, L., Wainwright, J. B., Thompson, C., Kershenbaum, S., McMillan, W. O. and
Montgomery, S. H. (2023). Plasticity and genetic effects contribute to different axes of
neural divergence in a community of mimetic *Heliconius* butterflies. *J. Evol. Biol.* 36,
1116–1132. https://doi.org/10.1111/jeb.14188.

Heinze, S., Reppert, S.M. (2012). Anatomical basis of sun compass navigation I: the general
layout of the monarch butterfly brain. *J. Comp. Neurol.* 520, 1599–1628.
https://doi.org/10.1002/cne.23054.

Herre, M., Goldman, O. V., Lu, T. C., Caballero-Vidal, G., Qi, Y., Gilbert, Z. N., Gong, Z.,
Morita, T., Rahiel, S., Ghaninia, M., et al. (2022). Non-canonical odor coding in the
mosquito. *Cell*, 185, 3104–3123.e28. https://doi.org/10.1016/j.cell.2022.07.024

Hildebrand, J.G., Shepherd, G.M. (1997). Mechanisms of olfactory discrimination:
converging evidence for common principles across phyla. *Annu. Rev. Neurosci.* 20,
595–631. https://doi.org/10.1146/annurev.neuro.20.1.595.

Hill, J.W., Petrucci, R.H. (1999). *General Chemistry: An Integrated Approach*. Prentice Hall.

Homberg, U., Christensen, T.A., Hildebrand, J.G. (1989). Structure and function of the
deutocerebrum in insects. *Annu. Rev. Entomol.* 34, 477–501.
https://doi.org/10.1146/annurev.en.34.010189.002401.

Ian, E., Chu, X., Berg, B.G. (2022). Brain investigation on sexual dimorphism in a
gynandromorph moth. *Insects* 13, 284. https://doi.org/10.3390/insects13030284.

Karger, D.N., Conrad, O., Böhner, J., Kawohl, T., Kreft, H., Soria-Auza, R.W., Zimmermann,
N.E., Linder, H.P., Kessler, M., (2017). Climatologies at high resolution for the earth's
land surface areas. *Sci Data.* 4, 170122. https://doi.org/10.1038/sdata.2017.122

Kazawa, T., Namiki, S., Fukushima, R., Terada, M., Soo, K., Kanzaki, R. (2009). Constancy
and variability of glomerular organization in the antennal lobe of the silkmoth. *Cell*
Tissue Res. 336:119–136.

Kirkness, E.F., Haas, B.J., Sun, W., Braig, H.R., Perotti, M.A., Clark, J.M., Lee, S.H.,
Robertson, H.M., Kennedy, R.C., Elhaik, E., et al. (2010). Genome sequences of the
human body louse and its primary endosymbiont provide insights into the permanent
parasitic lifestyle. *Proc. Natl. Acad. Sci. U.S.A.* 107, 12168–12173.
https://doi.org/10.1073/pnas.1003379107

Kondoh, Y., Kaneshiro, K.Y., Kimura, K., Yamamoto, D. (2003). Evolution of sexual
dimorphism in the olfactory brain of Hawaiian Drosophila. *Proc. R. Soc. Lond. B* 270,
1005–1013. https://doi.org/10.1098/rspb.2003.2331.

Koontz, M.A., Schneider, D. (1987). Sexual dimorphism in neuronal projections from the
antennae of silk moths (Bombyx mori, Antheraea polyphemus) and the gypsy moth
(Lymantria dispar). *Cell Tissue Res.* 249, 39–50.
https://doi.org/10.1007/BF00215416.

Kopp, A., Barmina, O., Hamilton, A. M., Higgins, L., McIntyre, L. M., & Jones, C. D. (2008).
Evolution of gene expression in the Drosophila olfactory system. *MBE*. 25, 1081–
1092. https://doi.org/10.1093/molbev/msn055

Kozak, K.M., Wahlberg, N., Neild, A.F.E., Dasmahapatra, K.K., Mallet, J., Jiggins, C.D.
(2015). Multilocus species trees show the recent adaptive radiation of the mimetic
Heliconius butterflies. *Systematic Biology* 64, 505–524.
https://doi.org/10.1093/sysbio/syv007.

Kruska, D.C.T. (2005). On the evolutionary significance of encephalization in some eutherian
mammals: effects of adaptive radiation, domestication, and feralization. *Brain Behav.*
Evol. 65, 73–108. https://doi.org/10.1159/000082979.

Kuehn, M., Welsch, H., Zahnert, T., Hummel, T. (2008). Changes of pressure and humidity
affect olfactory function. *Eur. Arch. Otorhinolaryngol.* 265, 299–302.
https://doi.org/10.1007/s00405-007-0446-2.

Kymre, J.H., Berge, C.N., Chu, X., Ian, E., Berg, B.G. (2021). Antennal-lobe neurons in the
moth *Helicoverpa armigera*: Morphological features of projection neurons, local
interneurons, and centrifugal neurons. *J. Comp. Neurol.* 529, 1516–1540.
https://doi.org/10.1002/cne.25034.

Larsson, M.C., Domingos, A.I., Jones, W.D., Chiappe, M.E., Amrein, H., Vosshall, L.B.
(2004). Or83b encodes a broadly expressed odorant receptor essential for
Drosophila olfaction. *Neuron* 43, 703–714.
https://doi.org/10.1016/j.neuron.2004.08.019.

Linz, J., Baschwitz, A., Strutz, A., Dweck, H.K.M., Sachse, S., Hansson, B.S., Stensmyr,
M.C. (2013). Host plant-driven sensory specialization in *Drosophila erecta*. *Proc. Biol.*
Sci. 280, 20130626. https://doi.org/10.1098/rspb.2013.0626.

Liu, J., He, K., Luo, Z., Cai, X., Bian, L., Li, Z., Chen, Z. (2021). Anatomical comparison of
antennal lobes in two sibling *Ectropis* moths: Emphasis on the macroglomerular
complex. *Front. Physiol.* 12. https://doi.org/10.3389/fphys.2021.685012.

Løfaldli, B.B., Kvello, P., Mustaparta, H. (2010). Integration of the antennal lobe glomeruli
and three projection neurons in the standard brain atlas of the moth *Heliothis*
virescens. *Front. Syst. Neurosci.* 4, 5. https://doi.org/10.3389/neuro.06.005.2010.

Ma, B., Chen, Q., Chu, X., Zuo, Y., Wang, J., Yang, Y., Wang, G., Ren, B. (2024). Glomerular
organization in the antennal lobe of the oriental armyworm *Mythimna separata*.
Journal of Integrative Agriculture 23, 3812–3829.
https://doi.org/10.1016/j.jia.2024.04.030.

Maan, M.E., Hofker, K.D., van Alphen, J.J.M., Seehausen, O. (2006). Sensory drive in cichlid
speciation. *Am. Nat.* 167, 947–954. https://doi.org/10.1086/503532.

Mallet, J. (1986). Gregarious roosting and home range in *Heliconius* butterflies. *National*
Geographic Research 2, 198–215.

Malnic, B., Hirono, J., Sato, T., Buck, L.B. (1999). Combinatorial receptor codes for odors.
Cell 96, 713–723. https://doi.org/10.1016/s0092-8674(00)80581-4.

Martin, F., Riveron, J., Alcorta, E. (2011). Environmental temperature modulates olfactory
reception in *Drosophila melanogaster*. *J. Insect Physiol.* 57, 1631–1642.
https://doi.org/10.1016/j.jinsphys.2011.08.016.

McLellan, C. F., & Montgomery, S. H. (2024). Evolution of larval gregariousness is
associated with host plant specialisation, but not host morphology, in Heliconiini
butterflies. *Ecol. Evol.* 14, e11002. https://doi.org/10.1002/ece3.11002

Meisami, E. (1989). A proposed relationship between increases in the number of olfactory
receptor neurons, convergence ratio and sensitivity in the developing rat. *Dev. Brain*
Res. 46, 9–19.

981 Mérot, C., Frérot, B., Leppik, E., Joron, M. (2015). Beyond magic traits: Multimodal mating
982   cues in *Heliconius* butterflies. *Evolution* 69, 2891–2904.
983   https://doi.org/10.1111/evo.12789.

984 Merrill, R., Dasmahapatra, K.K., Davey, J.W., Dell'Aglio, D.D., Hanly, J.J., Huber, B., Jiggins,
985   C.D., Joron, M., Kozak, K.M., Llaurens, V., et al. (2015). The diversification of
986   *Heliconius* butterflies: What have we learned in 150 years? *J. Evol. Biol.* 28, 1417–
987   1438. https://doi.org/10.1111/jeb.12672.

988 Merrill, R.M., Gompert, Z., Dembeck, L.M., Kronforst, M.R., McMillan, W.O., Jiggins, C.D.
989   (2011). Mate preference across the speciation continuum in a clade of mimetic
990   butterflies. *Evolution* 65, 1489–1500. https://doi.org/10.1111/j.1558-
991   5646.2010.01216.x.

992 Merrill, R.M., Naisbit, R.E., Mallet, J., Jiggins, C.D. (2013). Ecological and genetic factors
993   influencing the transition between host-use strategies in sympatric *Heliconius*
994   butterflies. *J. Evol. Biol.* 26, 1959–1967. https://doi.org/10.1111/jeb.12194.

995 Montgomery, S.H., Merrill, R. (2017). Divergence in brain composition during the early
996   stages of ecological specialization in *Heliconius* butterflies. *J. Evol. Biol.* 30, 571–
997   582. https://doi.org/10.1111/jeb.13027.

998 Montgomery, S.H., Merrill, R.M., Ott, S.R. (2016). Brain composition in *Heliconius* butterflies,
999   posteclosion growth and experience-dependent neuropil plasticity. *J. Comp. Neurol.*
1000   524, 1747–1769. https://doi.org/10.1002/cne.23993.

1001 Montgomery, S.H., Ott, S.R. (2015). Brain composition in *Godyris zavaleta*, a diurnal
1002   butterfly, reflects an increased reliance on olfactory information. *J. Comp. Neurol.*
1003   523, 869–891. https://doi.org/10.1002/cne.23711.

1004 Montgomery, S.H., Rossi, M., McMillan, W.O., Merrill, R.M. (2021). Neural divergence and
1005   hybrid disruption between ecologically isolated *Heliconius* butterflies. *Proc. Natl.*
1006   *Acad. Sci.* 118. https://doi.org/10.1073/pnas.2015102118.

1007 Morris, B.J., Couto, A., Aydin, A., Montgomery, S.H. (2021). Re-emergence and
1008   diversification of a specialized antennal lobe morphology in ithomiine butterflies.
1009   *Evolution* 75, 3191–3202. https://doi.org/10.1111/evo.14324.

1010 Münkemüller, T., Lavergne, S., Bzeznik, B., Dray, S., Jombart, T., Schiffers, K. and Thuiller,
1011   W. (2012). How to measure and test phylogenetic signal. Methods Ecol. Evol. 3, 743-
1012   756. https://doi.org/10.1111/j.2041-210X.2012.00196.x

1013 Nagappan, P.G., Subramaniam, S., Wang, D.-Y. (2017). Olfaction as a soldier—A review of
1014   the physiology and its present and future use in the military. *Mil. Med. Res.* 4, 9.
1015   https://doi.org/10.1186/s40779-017-0119-4.

Namiki, S., Daimon, T., Iwatsuki, C., Shimada, T., Kanzaki, R. (2014). Antennal lobe
organization and pheromone usage in bombycid moths. *Biol. Lett.* 10, 20140096.
https://doi.org/10.1098/rsbl.2014.0096.

Nei, M., Rooney, A.P. (2005). Concerted and birth-and-death evolution of multigene families.
Annu. Rev. Genet. 39, 121–152.
https://doi.org/10.1146/annurev.genet.39.073003.112240.

O'Donnell, S., Clifford, M.R., DeLeon, S., Papa, C., Zahedi, N., Bulova, S.J. (2013). Brain
size and visual environment predict species differences in paper wasp sensory
processing brain regions (Hymenoptera: Vespidae, Polistinae). *Brain Behav. Evol.*
82, 177–184. https://doi.org/10.1159/000354968.

Ott, S.R. (2008). Confocal microscopy in large insect brains: Zinc–formaldehyde fixation
improves synapsin immunostaining and preservation of morphology in whole-mounts.
J. Neurosci. Methods 172, 220–230. https://doi.org/10.1016/j.jneumeth.2008.04.031.

Piontkivska, H., Rooney, A.P., Nei, M. (2002). Purifying selection and birth-and-death
evolution in the histone H4 gene family. *Mol. Biol. Evol.* 19, 689–697.
https://doi.org/10.1093/oxfordjournals.molbev.a004127.

Prieto-Godino, L. L., Rytz, R., Bargeton, B., Abuin, L., Arguello, J. R., Peraro, M. D. and
Benton, R. (2016) 'Olfactory receptor pseudo-pseudogenes', *Nature*, 539(7627), pp.
93–97. Available at: https://doi.org/10.1038/nature19824.

Prieto-Godino, L.L., Rytz, R., Cruchet, S., Bargeton, B., Abuin, L., Silbering, A.F., Ruta, V.,
Peraro, M.D., Benton, R. (2017). Evolution of acid-sensing olfactory circuits in
Drosophilids. *Neuron* 93, 661–676.e6. https://doi.org/10.1016/j.neuron.2016.12.024.

Prieto-Godino, L.L., Silbering, A.F., Khallaf, M.A., Cruchet, S., Bojkowska, K., Pradervand,
S., Hansson, B.S., Knaden, M., Benton, R. (2020). Functional integration of "undead"
neurons in the olfactory system. *Science Adv.* 6, eaaz7238.
https://doi.org/10.1126/sciadv.aaz7238.

Prieto-Godino, L.L., Schmidt, H.R., Benton, R. (2021). Molecular reconstruction of recurrent
evolutionary switching in olfactory receptor specificity. *Elife* 10, e69732.
https://doi.org/10.7554/eLife.69732.

Rivas-Sánchez, D.F., Melo-Flórez, L., Aragón, A., Pardo-Díaz, C., Salazar, C., Montgomery,
S.H., Merrill, R.M. (2023). Parallel evolution of behavior, physiology, and life history
associated with altitudinal shifts in forest type in Heliconius butterflies. *Evolution* 77,
1458–1467. https://doi.org/10.1093/evolut/qpad062.

R Core Team (2024) *R: A language and environment for statistical computing.* Vienna,
Austria: R Foundation for Statistical Computing. Available at: http://www.R-
project.org.

Rivas-Sánchez, D.F., Salazar, C., Pardo-Díaz, C., Merrill, R.M., Montgomery, S.H. (2024).
Repeated evolution of reduced visual investment at the onset of ecological speciation
in high-altitude Heliconius butterflies. https://doi.org/10.1101/2024.06.12.598660.

Robertson, H.M., Gadau, J., Wanner, K.W. (2010). The insect chemoreceptor superfamily of
the parasitoid jewel wasp *Nasonia vitripennis*. *Insect Mol. Biol.* 19 Suppl 1, 121–136.
https://doi.org/10.1111/j.1365-2583.2009.00979.x.

Rooney, A.P., Piontkivska, H., Nei, M. (2002). Molecular evolution of the nontandemly
repeated genes of the histone 3 multigene family. *Mol. Biol. Evol.* 19, 68–75.
https://doi.org/10.1093/oxfordjournals.molbev.a003983.

Rospars, J.P. (1983). Invariance and sex-specific variations of the glomerular organization in
the antennal lobes of a moth, *Mamestra brassicae*, and a butterfly, *Pieris brassicae*.
J. Comp. Neurol. 220, 80–96. https://doi.org/10.1002/cne.902200108.

Rosser, N., Phillimore, A.B., Huertas, B., Willmott, K.R., Mallet, J. (2012). Testing historical
explanations for gradients in species richness in heliconiine butterflies of tropical
America. *Biol. J. Linn. Soc.* 105, 479–497. https://doi.org/10.1111/j.1095-
8312.2011.01814.x.

Rossi, M., Hausmann, A.E., Alcami, P., Moest, M., Roussou, R., Van Belleghem, S.M.,
Wright, D.S., Kuo, C.-Y., Lozano-Urrego, D., Maulana, A., et al. (2024). Adaptive
introgression of a visual preference gene. *Science* 383, 1368–1373.
https://doi.org/10.1126/science.adj9201.

Rundle, H.D., Nosil, P. (2005). Ecological speciation. *Ecol. Lett.* 8, 336–352.
https://doi.org/10.1111/j.1461-0248.2004.00715.x.

Schluter, D. (2009) 'Evidence for ecological speciation and its alternative', Science,
323(5915), pp. 737–741. Available at: https://doi.org/10.1126/science.1160006.

Schulz, S., Estrada, C., Yildizhan, S., Boppré, M., Gilbert, L.E. (2008). An antiaphrodisiac in
Heliconius melpomene butterflies. *J. Chem. Ecol.* 34, 82–93.
https://doi.org/10.1007/s10886-007-9393-z.

Seehausen, O., Terai, Y., Magalhaes, I.S., Carleton, K.L., Mrosso, H.D.J., Miyagi, R., van der
Sluijs, I., Schneider, M.V., Maan, M.E., Tachida, H., et al. (2008). Speciation through
sensory drive in cichlid fish. *Nature* 455, 620–626.
https://doi.org/10.1038/nature07285.

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E.V., Zdobnov, E.M. (2015).
BUSCO: assessing genome assembly and annotation completeness with single-copy
orthologs. *Bioinformatics* 31, 3210–3212.
https://doi.org/10.1093/bioinformatics/btv351.

Skiri, H.T., Rø, H., Berg, B.G., Mustaparta, H. (2005). Consistent organization of glomeruli in
the antennal lobes of related species of heliothine moths. *J. Comp. Neurol.* 491, 367–
380. https://doi.org/10.1002/cne.20692.

Stöckl, A., Heinze, S., Charalabidis, A., el Jundi, B., Warrant, E., Kelber, A. (2016).
Differential investment in visual and olfactory brain areas reflects behavioural choices
in hawk moths. *Sci. Rep.* 6, 26041. https://doi.org/10.1038/srep26041.

Störtkuhl, K.F., Hovemann, B.T., Carlson, J.R. (1999). Olfactory adaptation depends on the
Trp Ca2+ channel in *Drosophila. J. Neurosci.* 19, 4839–4846.
https://doi.org/10.1523/JNEUROSCI.19-12-04839.1999.

Störtkuhl, K.F., Kettler, R. (2001). Functional analysis of an olfactory receptor in *Drosophila*
melanogaster. *Proc. Natl. Acad. Sci. U. S. A.* 98, 9381–9385.
https://doi.org/10.1073/pnas.151105698.

Streinzer, M., Brockmann, A., Nagaraja, N., Spaethe, J. (2013). Sex and caste-specific
variation in compound eye morphology of five honeybee species. *PLOS ONE* 8,
e57702. https://doi.org/10.1371/journal.pone.0057702.

Talekar, N.S. and Shelton, A.M. (1993) 'Biology, Ecology, and Management of the
Diamondback Moth', *Annual Review of Entomology*. Annual Reviews. Available at:
https://doi.org/10.1146/annurev.en.38.010193.001423.

Task, D., Lin, C.-C., Vulpe, A., Afify, A., Ballou, S., Brbic, M., Schlegel, P., Raji, J., Jefferis,
G.S., Li, H., et al. (2022). Chemoreceptor co-expression in *Drosophila melanogaster*
olfactory neurons. *eLife* 11, e72599. https://doi.org/10.7554/eLife.72599.

Togunov, R.R., Derocher, A.E., Lunn, N.J. (2017). Windscapes and olfactory foraging in a
large carnivore. *Sci. Rep.* 7, 46332. https://doi.org/10.1038/srep46332.

Trona F, Anfora G, Bengtsson M, Witzgall P, Ignell R. (2010). Coding and interaction of sex
pheromone and plant volatile signals in the antennal lobe of the codling moth Cydia
pomonella. *J Exp Biol*. 213:4291–4303.

van Schooten, B., Meléndez-Rosa, J., Van Belleghem, S.M., Jiggins, C.D., Tan, J.D.,
McMillan, W.O., Papa, R. (2020). Divergence of chemosensing during the early
stages of speciation. *Proc. Natl. Acad. Sci.* 117, 16438–16447.
https://doi.org/10.1073/pnas.1921318117.

Varela N, Couton L, Gemeno C, Avilla J, Rospars J-P, Anton S. (2009). Three-dimensional
antennal lobe atlas of the oriental fruit moth, Cydia molesta (Busck) (Lepidoptera:
Tortricidae): comparison of male and female glomerular organization. *Cell Tissue*
Res. 337:513–526.

Vosshall, L.B., Wong, A.M., Axel, R. (2000). An olfactory sensory map in the fly brain. *Cell*
102, 147–159. https://doi.org/10.1016/S0092-8674(00)00021-0.

Wainwright, J.B., Loupasaki, T., Ramírez, F., Penry Williams, I.L., England, S.J., Barker, A., Meier, J.I., How, M.J., Roberts, N.W., Troscianko, J., et al. (2024). Mutualisms within light microhabitats drive sensory convergence in a mimetic butterfly community. *bioRxiv* p. 2024.08.16.607924. https://doi.org/10.1101/2024.08.16.607924

Wainwright, J.B., Montgomery, S.H. (2022). Neuroanatomical shifts mirror patterns of ecological divergence in three diverse clades of mimetic butterflies. *Evolution* 76, 1806–1820. https://doi.org/10.1111/evo.14547.

Wang, Y., Fang, G., Xu, P., Gao, B., Liu, X., Qi, X., Zhang, G., Cao, S., Li, Z., Ren, X., et al. (2022). Behavioural and genomic divergence between a generalist and a specialist fly. *Cell Reports* 41, 111654. https://doi.org/10.1016/j.celrep.2022.111654.

Wark, A.R., Peichel, C.L. (2010). Lateral line diversity among ecologically divergent threespine stickleback populations. *J. Exp. Biol.* 213, 108–117. https://doi.org/10.1242/jeb.031625.

Warrant, E.J. (2017). The remarkable visual capacities of nocturnal insects: vision at the limits with small eyes and tiny brains. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 372, 20160063. https://doi.org/10.1098/rstb.2016.0063.

Warton, D.I., Duursma, R.A., Falster, D.S., Taskinen, S. (2012). smatr 3 – an R package for estimation and inference about allometric lines. *Methods Ecol. Evol.* 3, 257–259. https://doi.org/10.1111/j.2041-210X.2011.00153.x.

Wright, D.S., van Eijk, R., Schuart, L., Seehausen, O., Groothuis, T.G.G., Maan, M.E. (2019). Testing sensory drive speciation in cichlid fish: Linking light conditions to opsin expression, opsin genotype, and female mate preference. *J. Evol. Biol.* 33, 422–434. https://doi.org/10.1111/jeb.13577.

Wu, N., Evans, E., van Schooten, B., Meléndez-Rosa, J., Ortiz, Y., Planas Soto-Navarro, S. M., Van Belleghem, S. M., Counterman, B. A., Papa, R., & Zhang, W. (2022). Widespread Gene Expression Divergence in Butterfly Sensory Tissues Plays a Fundamental Role During Reproductive Isolation and Speciation. *MBE*. 39, msac225. https://doi.org/10.1093/molbev/msac225

Wurm, Y., Wang, J., Riba-Grognuz, O., Corona, M., Nygaard, S., Hunt, B.G., Ingram, K.K., Falquet, L., Nipitwattanaphon, M., Gotzek, D., et al. (2011). The genome of the fire ant *Solenopsis invicta*. *Proc. Natl. Acad. Sci.* 108, 5679–5684. https://doi.org/10.1073/pnas.1009690108.

Yan, X., Wang, Z., Xie, J., Deng, C., Sun, X., Hao, C. (2019). Glomerular organization of the antennal lobes of the diamondback moth, *Plutella xylostella* L. *Front. Neuroanat.* 13. https://doi.org/10.3389/fnana.2019.00004.

1158   Yohe, L.R., Fabbri, M., Hanson, M., Bhullar, B.-A.S. (2020). Olfactory receptor gene
1159   evolution is unusually rapid across Tetrapoda and outpaces chemosensory
1160   phenotypic change. *Curr. Zool.* 66, 505–514. https://doi.org/10.1093/cz/zoaa051.
1161   Young, F.J., Montgomery, S.H. (2020). Pollen feeding in *Heliconius* butterflies: the singular
1162   evolution of an adaptive suite. *Proc. R. Soc. B: Biol. Sci.* 287, 20201304.
1163   https://doi.org/10.1098/rspb.2020.1304.
1164   Zjacic, N., Scholz, M. (2022). The role of food odour in invertebrate foraging. *Genes Brain*
1165   *Behav.* 21, e12793. https://doi.org/10.1111/gbb.12793.

1195 **Figures**

1196
1197

Figure 1: Olfactory pathway in invertebrates, including an overview of the antennal lobes and
their constituent components in the brain. A) Odours are detected by the olfactory receptor
neurons (ORNs) located in the maxillary palps and antennae in invertebrates. Sensory information is
then passed on by the ORNs to the first processing centre in the brain, the antennal lobe (AL). This
centre is made up of glomeruli (GL) and antennal lobe hub (ALH) which houses several different types
of neurons, ORNs, local interneurons (LNs) and projection neurons (PNs). LNs help to arbitrate
complex interactions in the antennal lobe. The signal is then forwarded to higher order processing
centres by the PNs. The butterfly used in the figure is *H. hecale*. Figure concept adapted from
Norwegian University of Science and Technology; https://shorturl.at/JAiDS. Scale bar: rendered
volume of whole brain: 500µm; *H. hecale*: 25mm **B)** Top: view of the antennal lobes in the X-Y plane
of a 3D image confocal stack of a Heliconiini brain; Bottom: zoomed-in view of the glomeruli and
antennal lobe hub, the constituent units of the antennal lobe. Scale bar: Top: 500µm, Bottom: 50µm.

1210

1211

Figure 2: Variation in the size of olfactory neuropils and associated ecological and
environmental traits in the Heliconiini. **A)** Phylogeny showing the Heliconiini species sampled and
the major phylogenetic groupings that were also used in the comparative analyses. Species that
displayed the greatest number of pairwise differences in terms of AL, ALH and GL volume, as well as
the change in GL~ALH volume are in bold. The associated environmental and ecological traits for each
species are plotted at the tips of the tree. **B)** Variation in the AL volume in the Heliconiini. **C)** Variation
in the ALH volume in the Heliconiini. **D)** Variation in the GL volume in the Heliconiini. **E)** Variation in the
GL~ALH volume in the Heliconiini. The volumes were allometrically controlled by plotting them against
the volume of the rCBR. The top 3 species that had the greatest number of pairwise differences in each
comparison are circled and labelled in the plots (note there are 4 species labelled in D) as 2 species
had a similar number of pairwise differences. Point size based on the number of individuals for each
species.

1224

1225

1226

Figure 3: Approximate Maximum likelihood phylogeny reconstruction of olfactory receptor gene family across Lepidoptera using non-Papilionoidea as reference (dashed lines). All monophyletic orthologous groups (OGs) are collapsed into a single node to show the entire phylogenetic tree, which contains 4658 butterfly genes and 428 reference genes. OGs containing butterfly sequences are listed in bold, together with its phylogenetic attribute. OGs without phylogenetic attribute are monophyletic at Lepidoptera level (e.g. ORco). The circle colours on nodes indicates bootstrap values.

1234

Figure 4: Copy number variation across Heliconinae and other Nymphalid species. On the left panel the ultrametric tree of all Nymphalid species generated with the chronos function (ape library), taking into account topologies and data phylogeny from (Cicconardi et al. 2023, 2024). In the central panel a bubble plot showing the expansions/contractions for each orthologous group (OG) (lower panel) for each nymphalid species. In the lower panel all butterfly OGs are listed with topological information for OG that are duplicated in butterflies. The right side of the panel shows a stacked histogram of percentage and absolute values of the number of OGs with complete or incomplete PFAM (7tm_6) domain. The upper panel shows instead boxplots of the distributions of fraction of protein sequence occupied by the 7tm_6 domain in each OGs. Broad distributions are a proxy of higher sequence variability.

1249

1250

Figure 5: Comparisons of wild (dark orange) and insectary-reared (light orange) olfactory

neuropil volumes for 5 species of *Heliconius*. A) phylogenetic relationship between *H. erato*, *H.*

melpomene, *H. cydno*, *H. ismenius*, and *H. hecale*. **B), C)** and **D)** boxplots of the volume of AL, ALH,

and GL respectively in insectary and wild individuals. **E)** Scaling relationship between GL and ALH

volume in insectary and wild individuals across species. The common logarithm ($\log_{10}$) of all volumes

were taken prior to analysis.

1257