The increase in cell volume and nuclear number of the koji-fungus *Aspergillus oryzae* contributes

to its high enzyme productivity

Ayaka Itani[1], Haruto Motomura[1], Ken Oda[2], Hideyuki Yamashita[3], Kanae Sakai[4], Ken-ichi Kusumoto[4],

Shinsuke Shigeto[5], Takehiko Ichikawa[6], Hosain Mohammad Mubarak[6], Takeshi Fukuma[6], Takuya

Katayama[7,8], Jun-ichi Maruyama[7,8], Shunsuke Masuo[1], Naoki Takaya[1], Norio Takeshita[1*]

[1] Microbiology Research Center for Sustainability (MiCS), Faculty of Life and Environmental

Sciences, University of Tsukuba, Tsukuba, 305-8572, Japan

[2] National Research Institute of Brewing, 3-7-1 Kagamiyama, Higashi-hiroshima, Hiroshima, 739-

0046, Japan

[3] Higuchi Matsunosuke Shoten Co., Ltd., 1-14-2 Harima-cho, Abeno-ku, Osaka 545-0022, Japan

[4] Department of Biotechnology, Graduate School of Engineering, Osaka University, 2-1 Yamadaoka,

Suita, Osaka 565-0871, Japan

[5] Department of Chemistry, School of Science, Kwansei Gakuin University, Sanda, Hyogo, 669-1330,

Japan

[6] Nano Life Science Institute (WPI-NanoLSI), Kanazawa University, Kakuma-machi, Kanazawa 920-

1192, Japan

[7] Department of Biotechnology, The University of Tokyo, Tokyo 113-8657, Japan

[8] Collaborative Research Institute for Innovative Microbiology, The University of Tokyo, Tokyo 113-

8657, Japan

* Correspondence to: Norio Takeshita, takeshita.norio.gf@u.tsukuba.ac.jp

Abstract

The regulation of cell volume and nuclear number in multi-nucleate organisms remains largely unknown. The koji-fungus *Aspergillus oryzae* has been used in traditional Japanese brewing and fermentation. We discovered that cell volume of hyphae increases tenfold, and the number of nuclei in the hyphal cells also increases tenfold, exceeding 200. The increase in cell volume and nuclear number is unique to *A. oryzae* and correlates with its high enzyme production capabilities. Through transcriptome analysis of thick hyphae isolated via laser microdissection and SNP analysis of practical strains from koji manufacturers, we identified a calcium channel and glycosyl transferase involved in cell volume expansion. Additionally, we indicated that specific amino acids and activation of the TOR pathway are essential for increasing nuclear number. This study not only deepens our understanding of the evolutionary processes that promote high enzyme productivity through fungal breeding, but also provides valuable insights into the molecular mechanisms regulating cell volume and nuclear number in multi-nucleate organisms.

Introduction

In unicellular organisms, cell volume is tightly regulated in coordination with nuclear division timing (1-3). Similarly, in multinucleated filamentous fungi, a correlation exists between cell volume and nuclear number (4, 5). However, the number of nuclei varies across species (6), the factors governing nuclear number and cell volume remain largely unknown. Filamentous fungi are industrially significant microorganisms, and the strains selected through breeding provide ideal research models for studying these evolutionary processes.

Filamentous fungi secrete a variety of hydrolytic enzymes to support their growth, and humans have long harnessed this enzyme secretion ability for the production of a wide range of fermented foods and beverages (7). *Aspergillus oryzae has been used in Japan for over 1,000 years to produce traditional fermented foods like sake, soy sauce, and miso* (8, 9). A distinctive feature of the fermentation techniques is solid-state cultivation using substrates like rice, soybeans, and wheat bran. The inoculum of *A. oryzae* used in fermentation, known as "koji," has been commercially produced for around 700 years (10). Koji is created by cultivating koji-fungus on grains, and koji starter manufacturers have selected and bred strains to improve the flavor and color of various fermented products. The *A. oryzae* strains currently in use are mainly preserved and managed by koji manufacturers and research institutions. A notable characteristic of *A. oryzae* is its high ability to secrete starch-degrading enzymes and proteases, which has led to extensive research into its genome and gene expression regulation (11-17).

Although *A. oryzae* shares 99.5% genomic similarity with *Aspergillus flavus*, which produces the carcinogenic aflatoxin (18), there have been no reports of *A. oryzae* producing aflatoxins, and its safety has been confirmed at the molecular level (19, 20). Phylogenomic analyses suggest that *A. oryzae* and *A. flavus* diverged approximately 50,000 to 189,000 years ago (21, 22). Over the following millennia, human use of *A. oryzae* is thought to have driven significant genomic

recombination, resulting in its evolution into a "cell factory" specifically optimized for the breakdown

of sugars and proteins (23).

Additionally, modern biotechnology utilizes filamentous fungi as cell factories to produce

organic acids, enzymes, and pharmaceuticals (24-26). *A. oryzae* also shows high production capacity

as a host for both homologous and heterologous protein production in modern biotechnology (27, 28).

Recently, its application has expanded to include the production of pharmaceutical proteins and

secondary metabolites (29, 30). Furthermore, filamentous fungi involved in fermentation play a crucial

role in creating a more sustainable food system, such as by upcycling agricultural by-products into

food through fungal fermentation and using A. oryzae mycelium for alternative meats (31, 32).

While basic and applied research on *A. oryzae* continues, fundamental questions remain

unanswered, such as why *A. oryzae* exhibits high enzyme production capacity and why it excels as a

host for heterologous expression. This study shows that the key lies in the increase in cell volume and

nuclear number, and analyzes the molecular mechanisms.

Result

Increase in nuclear number and cell volume in *Aspergillus oryzae*

In the model strain *A. oryzae* RIB40, hyphae on day 1 of cultivation contain 10-20 nuclei per cell, but

by day 2 of cultivation, apical cells with more than 200 nuclei appear (33). We found that the

proportion of hyphae with a higher number of nuclei (represented by class III, over 200 nuclei in the

cell, see methods) increased as the culture period increased from 1 to 3 days (Fig. 1A, B) (33). This

phenotype was not observed in the model fungus *Aspergillus nidulans* or the closely related species *A.*

flavus (Fig. 1A, B) (33). As far as we know, such phenotypes have not been observed in other strains

of *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus fumigatus*, or other strains commonly used in

research. Among other *Aspergillus* species used in fermentation (9, 10), the proportion of hyphae with

a higher number of nuclei increased in *Aspergillus sojae*, which is used for miso and soy sauce

production. This phenotype was not observed in *Aspergillus luchuensis* nor in its albino mutant

Aspergillus luchuensis mut. kawachii, which are used for distilled spirits such as awamori and shochu.

Among several strains of *A. oryzae* used in industrial applications, the RIB915 strain used for soy

sauce fermentation does not show an increase in the number of nuclei, while the RIB128 and RIB430

strains used in sake brewing exhibit a significant increase in the number of nuclei, suggesting that

different phenotypes were selected through distinct breeding processes.

The distribution of nuclei in the hyphae of *A. oryzae* RIB40 was visualized through 3D

reconstruction (Fig. 1C, Movie 1). In the class I hyphae, the nuclei were spaced relatively evenly,

whereas in the class III hyphae with increased nuclei, the nuclei were densely packed and arranged in

a disordered manner. Furthermore, the nuclear size in the class III hyphae was significantly smaller

than that in the class I hyphae (Fig. 1D, E).

Strains without an increase in nuclear number had hyphae with diameters ranging from 2 to

7 μm, while strains with an increase in nuclear number had some hyphae of similar thickness but often

exhibited thicker hyphae exceeding 10 μm (Fig. 1F). The number of nuclei and cell volume in hyphae

at 100 μm from the tips were compared among *A. oryzae* strains (RIB40, RIB128, RIB915) and *A.*

flavus (Fig. 1G). A strong positive correlation was observed between number of nuclei and cell volume.

This is consistent with previously reports that the number of nuclei per hyphal volume remains

constant (4, 5).

Thick hyphae with increased nuclei emerge by branching

Time-lapse imaging using the *A. oryzae* RIB40 expressing H2B-GFP indicated that thick hyphae with

increased nuclei emerged by branching from hyphae without increased nuclei (Fig. 2A, Movie 2). In

the emerged thick hyphae, rapid nuclear division occurred at the sites of branching (Fig. 2B, Movie

3). Immediately after branching, the thick hyphae showed a significantly faster rate of nuclear

proliferation, increasing 4–5 times within 8 hours compared to the thin hyphae immediately after

branching (Fig. 2C). Branches from hyphae with increased nuclei produced both hyphae with and

without increased nuclei (Fig. 2D).

Time-lapse imaging was conducted over a wider area, and image processing was applied to

color-code the nuclei: green for those in hyphae with increased nuclei and white for those in hyphae

without increased nuclei (Fig. 2E, Movie 4, Fig.S1A, see methods). Up to 24 hours after inoculation,

most hyphae were thin with no increase in nuclei number, but thick hyphae with increased nuclei

began to appear as time progressed (Fig. 2E, F). Since thick hyphae elongated faster than thin hyphae

(Fig. 2G), thick hyphae outgrew and overtook thin hyphae, eventually dominating the colony

perimeter (Fig. 2E, F).

Transmission electron microscopy confirmed basic nuclear structures, such as the nuclear

membrane and nucleolus, in both thick and thin hyphae (Fig. S1B). As hyphal diameter increased, cell

walls became more uneven and thicker (Fig. 2H, I). This finding corresponds with Raman

spectroscopy results showing pronounced peaks corresponding to cell wall polysaccharides in thick

hyphae (Fig. S1C)(34). Additionally, peaks associated with active mitochondria were prominent in

thick hyphae (Fig. S1C)(35), which was further confirmed by fluorescence staining (Fig. S1D). The

surface structure and mechanical properties of the cell wall were analyzed using atomic force

microscopy. Thick hyphae exhibited greater surface roughness and higher elasticity compared to thin

hyphae (Fig. 2J, K, Fig. S1E, F).

Correlation between nuclear number and enzyme secretion

We compared the amounts of secreted proteins in *A. oryzae* RIB40, RIB128, RIB915, *A. nidulans*, and

A. flavus (Fig. 3A). The strains that show an increase in nuclear number had significantly higher levels

of protein secretion than the strains that do not. In this condition, α-amylases are the dominant proteins

secreted by *A. oryzae* (15). The amylase enzyme activity was compared between in *A. oryzae* RIB40,

which shows an increase in nuclear number, and *A. oryzae* RIB915, which does not show an increase.

In RIB40, starting from day 2, when hyphae with increased nuclear numbers appeared, the enzyme

activity per fungal biomass significantly increased more than RIB915 (Fig. 3B).

To compare the amylase activity between hyphae with a higher and a lower number of nuclei,

amylase activity was monitored under conditions where a single hypha grew in a microfluidic channel

with a fluorescent substrate (Fig. 3C, S2A, Movie 5)(36). After 12 hours, the fluorescence intensity in

the channel with thick hyphae was about three times higher than that in the channel with thin hyphae,

indicating that thicker hyphae secreted more amylase than thinner hyphae (Fig. 3D).

We found that addition of yeast extract increased in both hyphal width and nuclear number

in the RIB915 strain (Fig. 3E, Fig. S2B), whereas adding yeast extract to *A. nidulans* did not result in

an increase in hyphal width or nuclear number. In *A. oryzae* RIB40, the addition of yeast extract did

not make the already thick hyphae any thicker but increased the proportion of thicker hyphae. Addition

of yeast extract increased protein secretion approximately 1.8 times in RIB40 and 8.5 times in RIB915

(Fig. 3F).

The addition of nucleic acids, peptone, or casamino acids to the minimal medium increased

the number of nuclei in *A. oryzae* RIB915, while vitamin B did not cause a significant change in

nuclear number (Fig. S2C). Although the addition of individual amino acids did not show the same

effect as yeast extract, the addition of asparagine, proline, glutamine, and branched-chain amino acids

increased the proportion of hyphae with increased nuclei by 40-50% (Fig. 3G, Fig. S2D). The amino

acids that induced nuclear number increase also led to an increase in protein secretion, indicating a

positive correlation between the ratio of class III hyphae and the increase in secreted proteins (Fig. 3H,

Fig. S2E). Since the amino acids inducing an increase in nuclear number may activate the TOR

pathway (37), *A. oryzae* RIB40 was grown with low concentration of rapamycin, a TOR pathway

inhibitor, which minimally affects colony size. Under the conditions, RIB40 did not produce hyphae

with an increased number of nuclei (Fig. 3I-K, Fig. S2F).

Transcriptome analyses in hyphae with increased nuclei

To investigate gene expression changes in hyphae with increased nuclei, transcriptome analysis was

performed under six conditions: In minimal medium; *A. oryzae* RIB40, which shows increased nuclei,

A. oryzae RIB915 and *A. nidulans*, which do not show increased nuclei. In minimal medium with yeast

extract, *A. oryzae* RIB40 and RIB915, which shows increased nuclei, and *A. nidulans,* which do not

show increased nuclei (Fig. 4A, Fig. S2B). In RIB915, supplementation with yeast extract led to a

more than four-fold increase in the expression of 660 genes, among which 449 genes did not show

such increase in RIB40 or *A. nidulans* (Fig. 4B, Table S1). GO analysis of these genes revealed that

processes related to cell wall synthesis and divalent metal ion transport were significantly enriched

(Fig. 4C).

To capture gene expression changes associated with increased nuclei with higher resolution,

thick and thin hyphae from RIB40 were dissected using laser microdissection, collected separately,

and subjected to transcriptome analysis (Fig. 4D, Movie 6). Clear differences in gene expression

pattern were observed between thick and thin hyphae (Fig. 4E). 558 genes that were more than four-

fold upregulated in thick hyphae but not in thin hyphae (Table S2). GO analysis of the upregulated

genes indicated an enrichment of processes related to biogenesis of ribosome and cellular component

(Fig. 4F). Among them, 21 matched those from Table S1 (Fig. 4G, Table S3). Of these, 13 had GO

annotations, which were related to conidiation, cell wall synthesis, Ca^{2+} transport and rRNA

processing (Fig. 3H). KEGG map visualization indicates that overall ribosomal gene expression

increased in the thick hyphae than the thin hyphae (Fig. S3A).

Among the gene list, we focused on *msy1* and *msy2*, which are involved in Ca^{2+} transport

and maintaining cell volume homeostasis in *Schizosaccharomyces pombe* (38, 39). Disruption of the

orthologue genes, *msyA* and *msyB,* in *A. oryzae* RIB40 did not have a significant impact on colony

growth, hyphal morphology and number of nuclei in minimal medium (Fig. S3B). Under hypoosmotic

shock conditions, the Δ*msyA* often indicated cell lysis near the hyphal tips (Fig. 4I, J). Under

hypoosmotic conditions, water influx into the cells causes hyphal cell expansion, but the volume and

turgor pressure are regulated to prevent cell lysis. The *msyA* disruption, however, impairs the

regulation of the hyphal cell expansion, leading to hyphal lysis (Fig. S3C-E).

SNP analysis among industrial strains with distinct phenotypes

The fact that currently used *A. oryzae* strains are managed and preserved by koji manufacturers and

research institutions is a notable advantage of *A. oryzae* research. Whole-genome sequencing and

comparative genome analysis of 82 *A. oryzae* strains with different applications have shown that these

strains cluster into several clades (24). We investigated whether nuclear numbers increase in 20 strains

selected from six different clades (Fig. 5A). In the minimal media supplemented with yeast extract, all

strains exhibited an increase in nuclear number (Fig. S4A). In the minimal media without yeast extract,

three strains from Clade C showed an increase in nuclear number, whereas strains from Clades A, B,

and E did not (Fig. 5A, B, Fig. S4A). Clades F and G included both strains with increased nuclear

numbers and strains without. SNP analysis of ORFs in Clades F and G revealed 108 mutations in

Clade F between TK-32 and TK-38, and 23 mutations in Clade G between TK-41 and TK-47 (Table

S4 and S5).

Among these, AO090038000626; *rseA*, predicted glycosyl transferase, was the only

common gene. This gene has been identified as a mutation in a *A. sojae* mutant with high extracellular

enzyme production (40). It is also shown that deletion of *rseA* in *A. nidulans* leads to cell wall defects

and promotes enzyme secretion (40). Comparison of mutations in *rseA* among strains in Clade C, F

and G indicates a few amino acid substitutions within the ORF, but no common mutations

corresponded to the phenotype (Fig. 5C, Fig. S4B). Structural prediction suggests that the mutations

are located inside the enzyme domain (Fig. S4C).

The disruption of the *rseA* in *A. oryzae* RIB40 resulted in significant growth delay, increased

branching, and the absence of hyphae with increased nuclei (Fig. 5D, Fig. S4D). The *rseA* gene DNA

fragments from the strains with increased nuclei (RIB40, TK-47) or from the strain without increased

nuclei (TK-41) were amplified by PCR and introduced ectopically into RIB915 (Fig. 5E). The strains

expressing *rseA* from RIB40 or TK-47 exhibited an increase in hyphal width and a corresponding

increase in the proportion of hyphae with more nuclei, whereas the strain expressing *rseA* from TK41

showed no increase in hyphal width or nuclear number (Fig. 5E, F, Fig. S4E).

220 Industrial breeding strains from other genera

To investigate whether the phenomenon of increased nuclear numbers also applies to other industrial

fungi, we compared the bred strain of *Trichoderma reesei* (QM9414) for cellulase production and its

control strain (QM6a)(41), as well as the bred strain of *Penicillium chrysogenum* (IFO4688) for

penicillin production and wild type strain (42). In minimal medium, no significant differences in

hyphal width or nuclear number were observed between the bred and control strains, whereas with the

addition of yeast extract, a notable increase in nuclear number and hyphal width was observed only in

the bred strains (Fig. 5G, H, Fig. S5A-C). The addition of yeast extract significantly increased cellulase

and protease activity only in the bred strain of *T. reesei* and *P. chrysogenum,* respectively (Fig. 5I, J).

These results suggest that even in modern biotechnology, where high enzyme or compound producing

strains are bred, strains with increased nuclear numbers were selected, like the traditional breeding

practices of koji-fungi.

Discussion

In *A. oryzae* some strains, hyphae grow about three times thicker and contain roughly ten times more

nuclei than the initial hyphae (Fig. 1). These thicker hyphae emerge from existing branches, where

nuclear division is promoted, resulting in the formation of hyphae with an increased number of nuclei

(Fig. 2). A clear correlation was observed between hyphal cell volume and nuclear number (Fig. 1G).

This consistency aligns with findings in other organisms, from unicellular yeast to multicellular plants

and animals, where the nuclear-to-cytoplasmic ratio is tightly regulated, allowing cell cycle

progression only upon reaching a critical cell size (1-3). Both the G1/S and G2/M transitions are

controled by size checkpoints, with specific mechanisms differing between species (44, 45).

In *A. nidulans*, it is known that the duration of the G1 and G2 phases is regulated by

temperature (46). While the mechanisms governing the G1/S phase transition remain unclear (47), the

G2/M phase transition has been shown to involve the following: During the interphase of *A. nidulans*,

NimX (CDK1) binds to NimE (cyclin B), and its activity is regulated through phosphorylation by

AnkA (Wee1 kinase) and NimT (Cdc25 phosphatase)(48-50). The activated NimX-NimE complex

subsequently dephosphorylates by NimT and activates NimA (Never-in-Mitosis A) protein kinase (51,

52). NimA is essential for the transition from G2 to mitosis, and its expression is tightly regulated

during the cell cycle (52). The mRNA level of *nimA* and *nimT* begin to increase in G2, reach a plateau

during late G2 and M phases, and are sharply degraded at the end of mitosis upon re-entry into

interphase (52, 53), meaning high expression levels of *nimA* and *nimT* result in a longer G2 phase. In

thick hyphae, *nimA* and *nimT* mRNA was hardly detected, suggesting that thick hyphae sense cell

volume in some way and subsequently shorten the G2 phase (Fig. S6A).

Branching emerges as a result of cell wall loosening at the branching site. SNP analysis

between industrial strains with different phenotypes identified RseA, a predicted glycosyltransferase

(Fig. 5). Deletion of *rseA* caused defects in cell wall synthesis in *A. nidulans*, leading to increased

enzyme secretion (40, 54). Although the targets of RseA are unknown, RseA could be involved in the

cell wall integrity by modifying glycosyl chains on enzymes responsible for cell wall synthesis and

degradation. Mutations in *rseA* may impair the regulation of cell wall loosening at the branching site,

resulting in the formation of thicker hyphae. In fact, in the thick hyphae of RIB40, localized

imbalances in cell wall components have been demonstrated by TEM and AFM.

Thicker hyphae can only form if high turgor pressure is maintained within the hyphae, in

addition to changes in cell wall remodeling during branching. *msyA* and *msyB*, whose orthologues are

calcium channels involved in the regulation of cell volume, were identified among the genes that show

a significant increase in expression in thicker hyphae (Fig. 4)(38, 39). Analysis of the knockout strains

demonstrated the function of cell volume regulation adapted to increased turgor pressure in

hypoosmotic conditions. The function of these calcium channels is likely crucial for maintaining the

balance of turgor pressure and cell volume.

The Target of Rapamycin (TOR) pathway is a central regulator of cellular growth,

responding to a variety of nutrients, such as amino acids, and carbon sources, as well as environmental

factors like oxygen levels, osmotic pressure, pH, and temperature stress (55, 56). Upon sensing these

inputs, the TOR pathway influences translation activity, metabolic flow, and cell size (56, 57). In both

yeast and mammalian models, amino acids like branched-chain amino acids (BCAAs), glutamine, and

alanine are crucial for nutrient sensing, forming the basis for TOR signaling activation (57-59). These

amino acids are like those contributing to nuclear increase in the RIB915 strain (Fig. 3G). Furthermore,

TOR inhibition by rapamycin in the RIB40 strain suppressed the formation of hyphae with increased

nuclei (Fig. 3I-K). Consistent with the activation of translation by the TOR pathway, the expression

of ribosome-related genes was elevated in thicker hyphae (Fig. S3A). The TOR pathway directly

regulates mitochondrial activity (60). Thick hyphae exhibited heightened mitochondrial density and

activity (Fig. S1C, D), suggesting that the TOR pathway support the high ATP demand associated with

rapid biosynthesis.

282   Cell volume and nuclear number mutually regulate each other, meaning that thicker hyphae

always possess more nuclei. For the formation of thicker hyphae with numerous nuclei, an increase in

cell volume and a corresponding increase in nuclei must occur simultaneously (Fig. S6A). In our

model, mutations in *rseA* loosen the cell wall during branching, and the regulation of turgor pressure

and cell volume by MsyA represents the necessary condition to increase cell volume and maintain

thick hyphae. At the same time, enhanced translational capacity involving TOR activation and

mechanisms that accelerate the cell cycle constitute another essential condition enabling an increase

in nuclear number.

290   *A. nidulans* and *A. flavus* did not exhibit an increase in nuclear number. *A. sojae* displayed

increased nuclear numbers, whereas *A. kawachii* and *A. luchuensis* did not (Fig. 1). The 20 strains of

A. oryzae selected from different clades showed an increase in nuclear number in the minimal medium

supplemented with yeast extract. In the medium without yeast extract, there were strains that showed

an increase in nuclear number and others that did not. These differences might be due to variations in

their applications or the selection of strains with diverse characteristics, such as not only enzyme

activity but also their impact on the flavor, aroma, and color of the final products. Additionally, there

might be dominant mutations that activate TOR pathway even in the minimal medium. Multiple gene

mutations were identified as contributing to the phenotype of increased hyphal volume and nuclear

number, suggesting that the phenotype was not the result of a single mutation but rather the gradual

accumulation of mutations through breeding. The koji-fungus strains used to produce high-quality

sake and soy sauce have been selectively bred over many years, long before the advent of modern

biotechnology, to enhance their saccharolytic and proteolytic activities. Modern biotechnological bred

strains such as *T. reesei* QM9414 and *P. chrysogenum* IFO4688 also exhibit a phenotype of thicker

hyphae with increased nuclear numbers under nutrient-rich conditions (Fig. 5). This consistency

suggests that by breeding naturally multinucleate filamentous fungi under nutrient-rich conditions, it

is possible to obtain strains with increased nuclear numbers and enhanced enzyme or compound

production.

The number of nuclei in filamentous fungi varies greatly among species (61). For example,

Neurospora crassa can exceed 100 nuclei, which correlates with its exceptionally rapid growth rate

and high protein secretion capacity (62). An increase in the number of nuclei per cell is expected to

enhance transcription and translation per cell, thereby improving enzyme production and secretion

capacity (Fig. S6B). However, maintaining a high nuclear number requires a substantial amount of

energy, and in conditions where external nutrient supply is insufficient, cells rapidly die. Thus, the

tendency to maintain a high nuclear number could be a selective pressure unlikely to occur in nature.

Koji-kin strains were bred specifically for high enzyme production in nutrient-rich environments, and

we propose that some of them acquired the specialized trait of increased cell volume and nuclear

number through selection. This study elucidates the evolutionary processes driven by breeding

selection in *A. oryzae* and the underlying mechanisms contributing to high enzyme productivity. These

findings not only offer new approaches for optimizing filamentous fungi for bioindustry applications

but also lead to the elucidation of the molecular mechanisms regulating cell volume and nuclear

number in multinucleated organisms.

Acknowledgements.

This work was funded by MEXT KAKENHI grant number 21H02095, 21K19062 and Scientific

Research on Innovative Areas "Post-Koch Ecology" grant number 22H04878 to NT. Ohsumi Frontier

Science Foundation, Noda Institute for Scientific Research Grant, and Japan Science and Technology

Agency (JST) ERATO grant number JPMJER1502 to NT. AFM measurement was supported by the

World Premier International Research Center Initiative (WPI).

Methods

Strains and Culture Conditions. The filamentous fungal strains used in this study are listed in Table S6. *A. oryzae* RIB40 expressing H2B-GFP, UtH2BG stain, is described previously (33). Minimal medium for *Aspergillus* species was shown in Table S7. When peptone, casamino acids, or yeast extract was added, the final concentration was 1%. Amino acids were added at a final concentration of 0.1%, B vitamins (biotin, pyridoxine hydrochloride, thiamin hydrochloride, riboflavin, PABA, and nicotinic acid) were added at 100 μM, and nucleotides (inosine and uridine) were added at 10 mM.

Genetic Manipulation. To delete target genes, genome editing plasmids and circular donor DNA plasmids were constructed as described previously (63). The sgRNA cassette was designed under the U6 promoter with the following target sequences: GTGACGATCTACGGTAACGC for the *msyA* gene, GATCAGTTGAGCGTCCCCTA for the *msyB* gene, and AGCTAGCGCAGCCGGTCTGC for the *rseA* gene. The genome editing plasmid was constructed by introducing the sgRNA expression cassette into the *Sma*I restriction site of the pRGE-gRT6 plasmid (63). For the donor plasmid, 1 kb fragments upstream and downstream of the target genes were amplified, ligated, and inserted into the linearized pUC19 vector (Takara Bio). Both plasmids were applied for transformation using the *A. oryzae* RIB40 strain. To amplify the *RseA* region, spanning 1 kb upstream to 1 kb downstream, the templates used were *A. oryzae* RIB40, TK-42, and TK-47 strains. The amplified fragments were inserted into the *Hind*III site of the pPTRI vector (TaKaRa) to construct the plasmid. Transformations were performed using the protoplast-PEG method according to the manufacturer's protocol for pPTRI. Transformants were selected on the minimal medium plates containing 0.1 mg/l pyrithiamine. Integration of constructs was confirmed by PCR analysis.

Microscopy. Images were captured using an Axio Observer Z1 inverted epifluorescence microscope (Carl Zeiss) equipped with Plan-Apochromat objectives, an AxioCam 506 mono camera, and a Colibri.2 LED light source. The stage temperature was maintained at 30°C with thermoplate (Tokai Hit). Nuclei were stained with SYBR Green (Takara), mitochondrial ware stained with Rhodamine 123 (Invitrogen), cell membrane were stained with FM4-64 (Invitrogen). Images were collected using Zen (Carl Zeiss) and ImageJ software.

Image Analysis. The distribution pattern of nuclei was classified into the following three types; Class I: nuclei distribute at a constant interval without overlapping; Class II: nuclei align but sometimes overlap; Class III: nuclei scattered throughout hyphae but not aligned. Nuclei, hyphal size, and fluorescence intensity were quantified using Zen and ImageJ software. Z-stack 3D reconstruction and nuclear color coding were performed with Imaris (Oxford Instruments). Hyphal elongation distances were analyzed using the MTrackJ plugin in ImageJ. For Fig. 2E, nuclei were binarized, and those in non-increased hyphae were displayed in white based on size and circularity.

Transmission Electron Microscopy (TEM). The culture samples were embedded in 1% water agar

and fixed with 2.5% glutaraldehyde in 0.1 M phosphate buffer (pH 7.0) at room temperature for 2

hours. Sample preparation and sectioning were performed at the Institute of Medicine, University of

Tsukuba. Observations were conducted using a JEM-1400 transmission electron microscope (JEOL).

Imaging with Microfluidic Devices. Microfluidic devices were fabricated as described previously

(33). Two types of devices were constructed: a 2D observation device to visualize hyphal expansion

(Fig. S1A) and a single-hypha isolation device for enzymatic activity measurement (Fig. S2A).

Conidia suspensions were loaded into a 10 ml plastic syringe (SS-10ESZ, Terumo) and connected to

polyethylene tubing (inner diameter 0.38 mm, outer diameter 1.09 mm, BD Intramedic). Air was

expelled from the syringe, and the suspension was injected into the device inlet. The number of conidia

flowing through the microfluidic channel was adjusted under a microscope. The syringe was then

replaced with another containing medium, and the medium was delivered at the minimum flow rate

using a syringe pump (YSP-101, YMC). The device was incubated at 30°C on a thermal stage (TOKAI

HIT) and imaged using an Axio Observer Z1 microscope (Carl Zeiss). α-amylase activity in the device

was measured using the EnzChek Ultra amylase assay kit (Invitrogen).

Enzyme Activity Assays. Fungal conidia 10^4 were inoculated onto 25 ml of medium in a petri dish

and statically cultured at 30 °C for 4 days. Protein concentration and enzymatic activities in the culture

supernatant were measured. Protein quantification was performed using the Bradford assay (Protein

Assay Kit, BIO-RAD), with bovine gamma globulin as the standard. Carboxymethyl cellulase

(CMCase) activity was determined by measuring reducing sugars using the 3,5-dinitrosalicylic acid

method. CMCase activity was assayed at 50 °C for 15 min in a reaction mixture containing 50 mM

sodium acetate buffer (pH 5.0) and 1% carboxymethyl cellulose. One unit of enzyme activity was

defined as the amount of enzyme required to release 1 μmol of reducing sugar (as glucose equivalent)

per minute. α-amylase activity was measured using an α-Amylase Assay Kit (Kikkoman Biochemifa)

according to the manufacturer's instructions. Acid carboxypeptidase (ACP) activity was determined

using the Acid Carboxypeptidase Assay Kit (Kikkoman Biochemifa) following the provided protocol.

RNA-seq Analysis. Sterilized cellophane was placed on agar plates, and conidia were inoculated.

After culturing at 30 °C for 3 days, samples were frozen, homogenized with a mortar and pestle, and

total RNA was extracted using the RNeasy Mini Kit (QIAGEN). Library preparation, sequencing, and

partial data analysis were performed by the Faculty of Medicine, University of Tsukuba. Sequencing

was conducted using 36 bp paired end reads on the Illumina NextSeq 500, and reads were mapped to

the *A. oryzae* RIB40 reference genome using the CLC Genomic Workbench (QIAGEN). For laser

microdissection, sterilized membrane slides (Carl Zeiss) were immersed in MM liquid, and conidia

were inoculated onto the membrane. After static culturing at 30 °C for 3 days, sections were collected

using a PALM MicroBeam laser microdissection system (Carl Zeiss). RNA was extracted from 500-

700 sections respectively using the RNeasy Mini Kit. Library preparation and sequencing were

performed by Takara Bio Inc., using 150 bp paired end reads on the Illumina NovaSeq 6000. Reads

were mapped to the *A. oryzae* RIB40 reference genome using the CLC Genomic Workbench. Genes

showing a ≥4-fold change in expression were considered significant, and gene ontology (GO)

enrichment analysis of biological processes was performed using ShinyGO 0.81 (64). Heatmaps were

visualized using the Python Seaborn package (65). Gene annotations were obtained from FungiDB.

Low Osmotic Pressure Shock Analysis. Conidia were spot inoculated onto the minimal medium

containing 1 M NaCl and cultured at 30 ℃ for 3 days. Colony tips were treated with distilled water

for 3 min, and the number of tip ruptures was counted. Hyphal tip swelling was assessed by staining

hyphae cultured in the minimal medium liquid for 3 days with FM4-64, mounting them on a coverslip

with a small volume of medium, and adding an equal volume of distilled water.

Raman Spectroscopy. The conidia were inoculated into minimal medium liquid in a glass-bottom

dish and cultured at 30 ℃ for 3 days. Raman spectra were acquired using an XploRA PLUS Raman

microscope, as described previously (32). Hyphae attached to the dish bottom were irradiated with a

532 nm laser at 0.93 mW for 30 seconds. Spike artifacts caused by cosmic rays were manually removed.

Ten spectra were averaged, and background signals attributed to the medium and glass were subtracted.

Protein Structure Analysis. RseA protein structure was predicted using AlphaFold2 and visualized

with PyMOL (v1.20). Domains were annotated using InterPro.

Atomic Force Microscopy (AFM). The conidia were inoculated into minimal medium liquid on a

Poly-L-Lysine-coated (Sigma Aldrich) glass-bottom dish (ibidi) and cultured at 30 ℃ for 3 days. We

used a JPK Nanowizard 4 (Bruker) equipped with an inverted fluorescence microscope (Eclipse Ti2,

Nikon). The temperature was maintained at 30 ℃ using a dish heater integrated into the stage. BL-

AC40TS-C2 cantilevers (Olympus, spring constant approximately 0.1 N/m) were used for AFM

imaging under the following conditions: QI mode, 1.5 × 1.5 μm or 0.5 × 0.5 μm scan size, 128 × 128

pixels, Z-length 1 μm, setpoint 0.1 nN, and Z speed 166 μm/s. The Young's modulus was calculated

using JPKSPM Data Processing software (Bruker). After averaging and baseline subtraction, Young's

modulus was calculated using a Hertz model with a triangular pyramid half-cone angle of 35°.

SNP analysis. The genome data of *A. oryzae* required for SNP identification was obtained from the

National Center for Biotechnology Information (NCBI). and mapped to *A. oryzae* RIB40 reference

sequence using BWA-MEM2 (64). The SNPs were listed by FreeBayes (67), and detailed comparative

analysis of genomes was performed on Integrative Genome Viewer (IGV) manually.

RNA seq. The RNA seq data have been deposited in DDBJ as BioProject PRJDB19992.

References

1. Amodeo, A. A. & Skotheim, J. M. Cell-Size Control. *Cold Spring Harb Perspect Biol* **8**, a019083 (2016).

2. Nurse, P. Genetic control of cell size at cell division in yeast. *Nature* **256**, 547–551 (1975).

3. Jorgensen, P. *et al.* The Size of the Nucleus Increases as Yeast Cells Grow. *MBoC* **18**, 3523–3532 (2007).

4. Dynesen J, Nielsen J. Branching is coordinated with mitosis in growing hyphae of *Aspergillus nidulans*. *Fungal Genet Biol* 40, 15-24 (2003).

5. Dundon SE, Chang SS, Kumar A, Occhipinti P, Shroff H, Roper M, Gladfelter AS. Clustered nuclei maintain autonomy and nucleocytoplasmic ratio control in a syncytium. *Mol Biol Cell* **27**, 2000-7(2016).

6. Roper, M., Ellison, C., Taylor J. W. & Glass N. L. Nuclear and Genome Dynamics in Multinucleate Ascomycete Fungi. *Curr Biol* **21**, R786-793 (2011).

7. Hurst, C. J. Microbial Fermentations in Nature and as Designed Processes. John Wiley & Sons Ltd. (2023)

8. Kitamoto, K. Cell biology of the *Koji* mold *Aspergillus oryzae*. *Biosci, Biotechnol, Biochem* **79**, 863–869 (2015).

9. Ichishima, E. Development of enzyme technology for *Aspergillus oryzae*, *A. sojae*, and *A. luchuensis,* the national microorganisms of Japan. *Bioscience, Biotechnology, and Biochemistry* **80**, 1681–1692 (2016).

10. Yamashita, H. Koji Starter and Koji World in Japan. *JoF* **7**, 569 (2021).

11. Machida, M. *et al.* Genome sequencing and analysis of *Aspergillus oryzae*. *Nature* **438**, 1157–1161 (2005).

12. Machida, M., Yamada, O. & Gomi, K. Genomics of Aspergillus oryzae: Learning from the History of Koji Mold and Exploration of Its Future. *DNA Research* **15**, 173–183 (2008).

13. Gomi, K. Regulatory mechanisms for amylolytic gene expression in the koji mold *Aspergillus oryzae*. *Bioscience, Biotechnology, and Biochemistry* **83**, 1385–1401 (2019).

14. Tanaka, M. & Gomi, K. Induction and Repression of Hydrolase Genes in *Aspergillus oryzae*. *Front. Microbiol.* **12**, 677603 (2021).

15. Wang, B. *et al.* Survey of the transcriptome of *Aspergillus oryzae* via massively parallel mRNA sequencing. *Nucleic Acids Research* **38**, 5075–5087 (2010).

16. Ishida, H. *et al.* Identification of functional elements that regulate the glucoamylase-encoding gene (*glaB*) expressed in solid-state culture of *Aspergillus oryzae*. *Current Genetics* **37**, 373–379 (2000).

17. Kitano, H., Kataoka, K., Furukawa, K. & Hara, S. Specific expression and temperature-

dependent expression of the acid protease-encoding gene (*pepA*) in *Aspergillus oryzae* in solid-

state culture (Rice-Koji). *J Biosci Bioengineering* **93**, 563–567 (2002).

18. Rokas, A. *et al.* What can comparative genomics tell us about species concepts in the genus

Aspergillus? *Studies in Mycology* **59**, 11–17 (2007).

19. Kusumoto, K.-I., Nogata, Y. & Ohta, H. Directed deletions in the aflatoxin biosynthesis gene

homolog cluster of *Aspergillus oryzae*. *Current Genetics* **37**, 104–111 (2000).

20. Kobayashi, T. *et al.* Genomics of *Aspergillus oryzae*. *Biosci, Biotechnol, Biochem* **71**, 646–670

(2007).

21. Watarai, N., Yamamoto, N., Sawada, K. & Yamada, T. Evolution of *Aspergillus oryzae* before

and after domestication inferred by large-scale comparative genomic analysis. *DNA Res* **26**,

465–472 (2019).

22. Chang, P.-K. & Ehrlich, K. C. What does genetic diversity of *Aspergillus flavus* tell us about

Aspergillus oryzae? *Internat J Food Microbiol* **138**, 189–199 (2010).

23. Gibbons, J. G. *et al.* The Evolutionary Imprint of Domestication on Genome Variation and

Function of the Filamentous Fungus *Aspergillus oryzae*. *Current Biol* **22**, 1403–1409 (2012).

24. Meyer, V. *et al.* Current challenges of research on filamentous fungi in relation to human

welfare and a sustainable bio-economy: a white paper. *Fungal Biol Biotechnol* **3**, 6, s40694-

016-0024–8 (2016).

25. Liu, D., Garrigues, S. & de Vries, R. P. Heterologous protein production in filamentous

fungi. *Appl Microbiol Biotechnol* **107**, 5019–5033 (2023).

26. Meyer, V. *et al.* Growing a circular economy with fungal biotechnology: a white paper. *Fungal*

Biol Biotechnol **7**, 5 (2020).

27. Christensen, T. *et al.* High Level Expression of Recombinant Genes in *Aspergillus oryzae*. *Nat*

Biotechnol **6**, 1419–1422 (1988).

28. Saito, N., Katayama, T., Minami, A., Oikawa, H. & Maruyama, J. Versatile filamentous fungal

host highly-producing heterologous natural products developed by genome editing-mediated

engineering of multiple metabolic pathways. *Commun Biol* **7**, 1263 (2024).

29. Huynh, H. H. *et al.* Functional production of human antibody by the filamentous fungus

Aspergillus oryzae. *Fungal Biol Biotechnol* **7**, 7 (2020).

30. Itoh, T., Tokunaga, K., Matsuda, Y. *et al.* Reconstitution of a fungal meroterpenoid biosynthesis

reveals the involvement of a novel family of terpene cyclases. *Nature Chem* **2**, 858–864 (2010).

31. Maini Rekdal, V., Villalobos-Escobedo, J.M., Rodriguez-Valeron, N. *et al.* Neurospora

intermedia from a traditional fermented food enables waste-to-food conversion. *Nat*

Microbiol **9**, 2666–2683 (2024).

32. Maini Rekdal, V., van der Luijt, C.R.B., Chen, Y. *et al.* Edible mycelium bioengineered for

enhanced nutritional value and sensory appeal using a modular synthetic biology toolkit. *Nat*

Commun **15**, 2099 (2024).

33. Yasui M, Oda K, Masuo S, Hosoda S, Katayama T, Maruyama JI, Takaya N, Takeshita N.

Invasive growth of *Aspergillus oryzae* in rice koji and increase of nuclear number. *Fungal Biol*

Biotechnol. 7, 8 (2020).

34. Hossain IM, Mazumder N, Kaino T, Kawamukai M, Noothalapati H, Yamamoto T. Direct

Visualization of Structurally Similar Polysaccharides in Single Yeast Cells in Vivo by

Multivariate Analysis Assisted Raman Microspectroscopy. *J Phys Chem B* 127, 5249-5256

(2023).

35. Yasuda, M., Takeshita, N. & Shigeto, S. Inhomogeneous Molecular Distributions and

Cytochrome Types and Redox States in Fungal Cells Revealed by Raman Hyperspectral

Imaging Using Multivariate Curve Resolution–Alternating Least Squares. *Anal. Chem.* **91**,

12501–12508 (2019).

36. Itani, A., Shida, Y. & Ogasawara, W. A microfluidic device for simultaneous detection of

enzyme secretion and elongation of a single hypha. *Front. Microbiol.* **14**, 1125760 (2023).

37. Wouters, B. G. & Koritzinsky, M. Hypoxia signalling through mTOR and the unfolded protein

response in cancer. *Nat Rev Cancer* **8**, 851–864 (2008).

38. Nakayama, Y., Yoshimura, K. & Iida, H. Organellar mechanosensitive channels in fission yeast

regulate the hypo-osmotic shock response. *Nat Commun* **3**, 1020 (2012).

39. Nakayama, Y., Hirata, A. & Iida, H. Mechanosensitive channels Msy1 and Msy2 are required

for maintaining organelle integrity upon hypoosmotic shock in *Schizosaccharomyces*

pombe. *FEMS Yeast Res* **14**, 992–994 (2014).

40. Ogawa, M. *et al.* Deletion of *Aspergillus nidulans cpsA/rseA* induces increased extracellular

hydrolase production in solid-state culture partly through the high osmolarity glycerol

pathway. *J Biosci Bioeng* **131**, 589–598 (2021).

41. Vitikainen, M. *et al.* Array comparative genomic hybridization analysis of Trichoderma reesei

strains with enhanced cellulase production properties. *BMC Genomics* **11**, 441 (2010).

42. Barreiro, C., Martín, J. F. & García-Estrada, C. Proteomics Shows New Faces for the Old

Penicillin Producer *Penicillium chrysogenum*. *J Biomed Biotechnol* **2012**, 1–15 (2012).

43. Murphy, C. M. & Michael, W. M. Control of DNA Replication by the Nucleus/Cytoplasm Ratio

in Xenopus. *J Biol Chem* **288**, 29382–29393 (2013).

44. Futcher, B. & Kellogg, D. R. SnapShot: Cell size control. *Cell* **187**, 2896-2896.e1 (2024).

45. Fantes, P. & Nurse, P. Control of cell size at division in fission yeast by a growth-modulated

size control over nuclear division. *Experiment Cell Research* **107**, 377–386 (1977).

46. Bergen, L. G. & Morris, N. R. Kinetics of the nuclear division cycle of *Aspergillus nidulans*. *J*

Bacteriol **156**, 155–160 (1983).

47. Dörter, I. & Momany, M. Fungal Cell Cycle: A Unicellular versus Multicellular

Comparison. *Microbiol Spectr* **4**, 4.6.28 (2016).

48. Gould, K. L. & Nurse, P. Tyrosine phosphorylation of the fission yeast cdc2+ protein kinase

regulates entry into mitosis. *Nature* **342**, 39–45 (1989).

49. Wu, L., Osmani, S. A. & Mirabito, P. M. A Role for NIMA in the Nuclear Localization of

Cyclin B in *Aspergillus nidulans*. *J Cell Biol* **141**, 1575–1587 (1998).

50. Ye, X. S., Fincher, R. R., Tang, A. & Osmani, S. A. The G2/M DNA damage checkpoint

inhibits mitosis through Tyr15 phosphorylation of p34cdc2 in *Aspergillus nidulans*. *EMBO*

J **16**, 182–192 (1997).

51. Shen, K. F. & Osmani, S. A. Regulation of mitosis by the NIMA kinase involves TINA and its

newly discovered partner, An-WDR8, at spindle pole bodies. *MBoC* **24**, 3842–3856 (2013).

52. Osmani, S. A., May, G. S. & Morris, N. R. Regulation of the mRNA levels of *nimA*, a gene

required for the G2-M transition in *Aspergillus nidulans*. *J Cell Biol* **104**, 1495–1504 (1987).

53. Osmani, A. H., O'Donnell, K., Pu, R. T. & Osmani, S. A. Activation of the *nimA* protein kinase

plays a unique role during mitosis that cannot be bypassed by absence of the *bimE* checkpoint.

EMBO J **10**, 2669–2679 (1991).

54. Feng, X. *et al. cpsA* regulates mycotoxin production, morphogenesis and cell wall biosynthesis

in the fungus *Aspergillus nidulans*. *Mol Microbiol* **105**, 1–24 (2017).

55. Urban, J. *et al.* Sch9 Is a Major Target of TORC1 in *Saccharomyces cerevisiae*. *Molecular*

Cell **26**, 663–674 (2007).

56. De Virgilio, C. & Loewith, R. The TOR signalling network from yeast to man. *The Internat J*

Biochem Cell Biol **38**, 1476–1481 (2006).

57. Kim, J. & Guan, K.-L. mTOR as a central hub of nutrient signalling and cell growth. *Nat Cell*

Biol **21**, 63–71 (2019).

58. Saxton, R. A. & Sabatini, D. M. mTOR Signaling in Growth, Metabolism, and

Disease. *Cell* **169**, 361–371 (2017).

59. Meng, D. *et al.* Glutamine and asparagine activate mTORC1 independently of Rag GTPases. *J*

Biol Chem **295**, 2890–2899 (2020).

60. Düvel, K. *et al.* Activation of a Metabolic Gene Regulatory Network Downstream of mTOR

Complex 1. *Mol Cell* **39**, 171–183 (2010).

61. Roper, M., Ellison, C., Taylor J. W. & Glass N. L. Nuclear and Genome Dynamics in

Multinucleate Ascomycete Fungi. *Curr Biol* **21**, R786-793 (2011).

62. Havlik, D., Brandt, U., Bohle, K. & Fleißner, A. Establishment of *Neurospora crassa* as a host

for heterologous protein production using a human antibody fragment as a model product.

Microbial Cell Factories **16**, 128 (2017).

63. Katayama, T. *et al.* Forced Recycling of an AMA1-Based Genome-Editing Plasmid Allows for

Efficient Multiple Gene Deletion/Integration in the Industrial Filamentous Fungus *Aspergillus*

oryzae. Appl Environ Microbiol **85**, e01896-18 (2019).

64. Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for animals and

plants. *Bioinformatics* **36**, 2628–2629 (2020).

65. Waskom, M. seaborn: statistical data visualization. *JOSS* **6**, 3021 (2021).

66. Vasimuddin, M., Misra, S., Li, H. & Aluru, S. Efficient Architecture-Aware Acceleration of

BWA-MEM for Multicore Systems. *IEEE Parallel and Distributed Processing Symposium*

(IPDPS), 314-324 (2019)

67. Garrison, E. & Marth, G. Haplotype-based variant detection from short-read sequencing.

arXiv:1207.3907 [q-bio.GN] https://doi.org/10.48550/arXiv.1207.3907

Figure legends.

Figure 1. Increase in number of nuclei and cell volume in *Aspergillus* species. (A) The nuclear

distribution in the tip cells was categorized into Classes I–III. The ratio of hyphae in each class was

measured in *Aspergillus* species at 24, 48, and 72 hours of growth (n=50). Data for *A. oryzae* RIB40

and *A. nidulans* are reproduced from a previous study (33). (B) Colony morphology after 3 days of

culture on the minimal medium. The nuclear distribution in the hyphae at the colony periphery stained

with SYBR Green. Scale bars: hyphae, 20 μm; colonies, 1 cm. (C) 3D images of hyphae without

increased nuclei (left) and with increased nuclei (right) in *A. oryzae* RIB40 from Movie 1. Each

nucleus is indicated with different colors by Imaris soft. Scale bar: 5 μm. (D) Comparison of nuclear

size in Class I hyphae (2.4 μm) and Class III hyphae (1.6 μm), indicated by white lines. (E) Box plots

of nucleus diameters in hyphae without increased nuclei and those with increased nuclei (n=35, ** p

< 0.01, t-test). (F) Box plots of hyphal width at the colony periphery grown for 72 hours (n=10–14, **

p< 0.01, * p< 0.05, t-test). Strains with increased nuclei are marked in red, and those without are

marked in blue. (G) Correlation between nuclear number and cell volume of hyphae at 100 μm from

the tips in *A. oryzae* RIB40, RIB128, RIB915, and *A. flavus*.

Figure 2. Thick hyphae with increased nuclei emerge by branching. (A) Time-lapse image sequence

of *A. oryzae* RIB40 expressing H2B-GFP showing the emergence of thick hyphae with increased

nuclei by branching from Movie 2. Scale bar: 20 μm. Elapsed time is indicated in min. (B) Image

sequence of successive nuclear division within the newly emerged branched hypha from Movie 3.

Scale bar: 10 μm. Elapsed time is indicated in min. (C) Time course of nuclear number per hypha.

Nuclear numbers were counted with the start time of branching set as 0, every hour for 5 to 10 h.

Thick; hyphal width >7 μm, Thin; hyphal width <5 μm. (D) Branching from hyphae with increased

nuclei generates both thick hyphae with increased nuclei (left) and thin hyphae without increased

nuclei (right). Scale bars: 20 μm. (E) Image sequence of mycelial growth showing hyphae with

increased nuclei (green) and without increased nuclei (white) from Movie 4. Scale bars: 200 μm.

Elapsed time is indicated in h. (F) Line histogram of hyphal width at the colony periphery at 20, 40,

and 60 h, calculated from Movie 4 (n=40). (G) Correlation between hyphal width and maximum

elongation rate calculated from Movie 4 (n=45). The maximum elongation rate is determined from

elongation rates measured every hour over 10 hours period. (H) TEM images of *A. oryzae* RIB40

hyphae. Hyphal diameter and cell wall thickness are indicated white and black text respectively. Scale

bar: 1 μm. (I) Correlation plots of hyphal width and cell wall thickness based on TEM images. Thick

hyphae (>7 μm) are marked in red, and thin hyphae (<7 μm) are marked in blue. (J) 3D surface images

of *A. oryzae* RIB40 hyphal tips in the thick hypha (upper) and the thin hypha (lower) constructed using

AFM. (K) Surface roughness of cell walls in thick (red) and thin (blue) hyphae along the arrows in J.

Figure 3. Correlation between number of nuclei and enzyme secretion. (A) Secreted protein per

biomass in strains with increased nuclei (red) and without increased nuclei (blue) after 4 days of culture

in minimal medium with 1% maltose (mean ± S.E., n=3, ** $p < 0.01$, t-test). (B) Time course of α-

amylase activity per biomass in *A. oryzae* RIB40 and RIB915 cultured in minimal medium with 1%

maltose (mean ± S.E., n=3). (C) Measurement of α-amylase activity in a single hypha using a

microfluidic device. A fluorescent substrate increases in fluorescence upon hydrolysis by α-amylase.

ROIs for thick hypha are marked in red, and thin hypha in blue. Scale bar: 10 μm. (D) Temporal

changes in fluorescence intensity measured in individual flow channels, as described in C. (E) Box

plots of hyphal width in colonies grown in minimal medium with or without 1% yeast extract (n=10–

18, ** $p < 0.01$, * $p < 0.05$, t-test). Conditions with increased nuclei are marked in red, and those

without in blue. (F) Secreted protein per biomass in minimal medium with or without 1% yeast extract

(mean ± S.E., n=3, ** $p < 0.01$, t-test). (G) Ratio of class I-III hyphae in *A. oryzae* RIB915 colonies

grown in minimal medium supplemented with 1% yeast extract or 0.1% individual amino acids (n=50).

(H) Correlation between the ratio of Class III hyphae and secreted protein per biomass in the 0.1%

amino acid-supplemented medium. (I) Images of hyphae and nuclear distribution in *A. oryzae* RIB40

grown on minimal medium with or without 0.5 ng/ml rapamycin. Scale bars: 10 μm. (J) Ratio of class

I-III hyphae in *A. oryzae* RIB40 cultured on minimal medium containing 0, 0.5, 5, or 100 ng/ml

rapamycin (n=50). (K) Box plots of hyphal width in the colonies cultured under the conditions in J

(n=11-18, ** $p < 0.01$, t-test).

Figure 4. Transcriptome analyses in hyphae with increased nuclei. (A) Heatmap of gene expression in

A. oryzae RIB40, RIB915, and *A. nidulans* grown in the minimal medium with or without 1% yeast

extract. The top 500 genes with the largest variation in expression are shown. (B) Venn diagram of

genes upregulated more than four-fold in the medium with yeast extract. (C) GO process analysis of

genes uniquely upregulated in *A. oryzae* RIB915 grown in the medium with yeast extract. (D) Image

sequence of cutting and collecting the targeted hypha by using laser microdissection. Scale bar: 100

μm. (E) Heatmap of gene expression in thick (n=3) and thin (n=2) hyphae dissected from *A. oryzae*

RIB40. The top 300 genes with the largest expression differences are shown. (F) GO process analysis

of genes upregulated more than four-fold in thick hyphae compared to thin hyphae. (G) Venn diagram

of 449 genes upregulated in RIB915 with yeast extract and 558 genes upregulated in thick hyphae of

RIB40. (H) GO annotations of 13 genes from the 21 shared genes in G. Genes related to cell wall

processes are shown in green, cell membrane in blue, Ca^{2+} transport in yellow, and rRNA processing

in orange. (I) Images of hyphae of RIB40 and Δ*msyA* strains after 3 min of low osmotic stress. Scale

bar: 200 μm. (J) Ratio of hyphal tip lysis under the hypoosmotic shock (mean ± S.E., n=50, ** $p <$

0.01, t-test).

Figure 5. Comparison of industrial bred strains in *A. oryzae, T. reesei* and *P. chrysogenum*. (A)

Phylogenetic clade A-F of *A. oryzae* TK strains. The strains with increased nuclei are indicated in red.

(B) Colonies and hyphae stained with SYBR Green of strains in clades C, F, and G grown for 3 days

on the minimal medium. Scale bar: 20 μm. (C) Sequence differences in *rseA* gene among strains in

clades C, F, and G. UTR regions are marked in pink, exons in purple, and nucleotide differences in

white. (D) Colonies and hyphae stained with SYBR Green of RIB40 and the Δ*rseA* strain grown for 3

days on the minimal medium. Scale bar: 10 μm. (E) Colonies and hyphae stained with SYBR Green

of RIB915 expressing *rseA* gene from RIB40 and TK-47 strains with increased nuclei, and TK-41

strain without increased nuclei. Scale bar: 20 μm. (F) Hyphal width of RIB915 and strains in E (n=14–

23). Strains with *rseA* from strains with increased nuclei are marked in red, and those without are

marked in blue. (G) Hyphal images stained with SYBR Green of *T. reesei* (QM9414) and *P.*

chrysogenum (IFO4688) grown for 3 days on the minimal medium with or without 1% yeast extract.

Scale bar: 20 μm. (H) Ratio of class I-III hyphae in the *T. reesei* and *P. chrysogenum* control and bred

strains cultured on the minimal medium with or without 1% yeast extract for 3 days (n=50). (I)

CMCase activity per biomass of *T. reesei* industrial and control strains under the conditions in G (mean

± S.E., n=3, ** $p < 0.01$, t-test). (J) Protease activity per biomass of *P. chrysogenum* industrial and

control strains under the conditions in G (mean ± S.E., n=3, ** $p < 0.01$, t-test).

680
681
682
683

Fig. 1

Fig. 2

Fig. 3

Fig. 4

H

Gene ID	GO annotation	Symbol
AO090003000583	Conidium formation, Establishment or maintenance of actin cytoskeleton polarity	*rho2*
AO090026000532	Asexual sporulation, Conidium formation	*flbA*
AO090701000589	Chitin biosynthetic process, Conidium formation	*chsB*
AO090012000084	Chitin synthase	*chsA*
AO090120000132	UDP-N-acetylglucosamine biosynthetic process	*gna1*
AO090009000257	Cellular sphingolipid homeostasis	*orm1/orm2*
AO090011000706	Cell volume homeostasis, Cellular calcium ion homeostasis	*msy2*
AO090012000418	Cell volume homeostasis, Calcium ion transmembrane transport	*msy1*
AO090038000531	Calcium ion import across plasma membrane	*trp1322*
AO090026000636	rRNA processing	*efg1*
AO090005001611	rRNA processing	*erb1*
AO090701000108	Mitochondrial tRNA wobble uridine modification, tRNA processing	*mto1*
AO090001000552	Cellular morphogenesis protein	*bud22*

Fig. 5