Intranasal Leukemia Inhibitory Factor as a late-stage treatment for delayed white matter damage in concussive head injury

Veera D'Mello[1], Jelena Mihailovic[2], Sidra Ali[1], Basavaraju G. Sanganahalli[2], Daniel Coman[2] Fahmeed Hyder[2], Merisha Fernando[1], Anita Mampilly[1], Sridhar S. Kannurpatti[1] and Steven W. Levison[1].

[1] Rutgers-New Jersey Medical School, Newark, NJ, 07103 USA
[2] Yale University, Magnetic Resonance Research Center, New Haven, CT, 06520 USA

Running Title: Delayed Intranasal LIF prevents neurodegeneration after mTBI

Keywords: traumatic brain injury, astrogliosis, axonal damage, MRI, DTI, corpus callosum, prefrontal cortex.

Corresponding Author

Steven W. Levison, PhD
Dept of Pharmacology Physiology and Neuroscience
Rutgers New Jersey Medical School
205 S. Orange Ave.
Newark, NJ 07101
E-mail: levisosw@rutgers.edu

Abstract

Leukemia Inhibitory Factor (LIF) is an injury-induced cytokine that peaks 48 hours after a traumatic brain injury (TBI). Juvenile LIF haplodeficient mice exhibit desynchronized glial responses, increased neurodegeneration, decreased axonal conductivity and behavioral deficits after a concussive head injury. Given the necessity of LIF during the acute recovery phase after injury, we hypothesized that intranasal (IN) LIF treatment would prevent neurodegeneration when administered during the chronic recovery period from a mild TBI (mTBI). Young adult male CD1 mice were subjected to a midline, closed-head frontal cortex injury using a flat metal impactor with a 3mm tip to induce a mTBI. In the 6-8 weeks post-mTBI, known to precede axonal atrophy in this mTBI model, two doses of 40 ng and 100 ng of LIF were administered twice daily, 5 days/week for two consecutive weeks. Sensorimotor functions were assessed at 4 and 8 weeks post mTBI, followed by ex-vivo brain magnetic resonance imaging at 9.4T and histopathology. mTBI mice showed sensorimotor deficits at 4 weeks, which worsened by 8 weeks post-injury. IN-LIF treatment prevented the progressive sensorimotor loss seen in the vehicle-treated controls. Increased mean diffusivity (MD) and decreased fractional anisotropy (FA) were observed in the corpus callosum and prefrontal cortex of mTBI brains. In a dose-dependent manner, IN-LIF prevented the mTBI-induced MD increase and FA decrease. Histologically, there was significantly less astrogliosis, microgliosis and axonal injury in the IN-LIF treated mice vs. controls. These results support the therapeutic potential of IN-LIF to reduce delayed neurodegeneration and improve neurological outcomes after mTBIs.

Introduction

Almost 80% of all traumatic brain injuries (TBIs) are classified as mild (mTBI) based on clinical severity. Whereas it had been thought that a single concussive hit to the head did not produce long-term effects, approximately 1/5 of mTBI patients report poor memory, reduced processing speed, fatigue, impulsivity, anxiety or depression that may be due to delayed neurodegeneration (1, 2). Recently, CTE pathology was found in young (~23-year-old) deceased athletes, underscoring the fact that unaddressed TBI can progress to more serious neurodegenerative pathologies (3). Analogously, several studies have shown persistent behavioral deficits in single-hit adult mouse mTBI models across various behavioral tasks, including spatial memory (4), olfactory discrimination (5), object recognition (6), passive avoidance (7), the forced swim test (7) , and analyses of circadian rhythms (8). Progress is being made in developing neuroprotective strategies to reduce acute brain injury, and several FDA-approved drugs are being investigated (9); however, clinical trials are still in the early stages. By contrast, drugs are lacking that can be administered during the chronic stage of recovery. Indeed, in a clinical trial to evaluate the efficacy of minocycline, microglial activation appeared to be inhibited. However, this did not prevent tertiary neurodegeneration as revealed by increased plasma neurofilament light chain (10) levels.

Histopathological and electrophysiological analyses in a single hit closed head injury of mTBI have provided insights into the chronic phase of neurodegeneration(11). Notably, with this injury, the subcortical white matter (WM) appears normal up to 6 weeks after the mTBI. However, tertiary neurodegeneration becomes apparent after 6 weeks of recovery, where the compound action potential becomes weak, accompanied by Wallerian degeneration (12). Moreover, the enzyme Sterile alpha and TIR motif containing 1 (SARM1), which is the executor for Wallerian axonal degeneration, is necessary for this delayed axonal injury, as the axonal degeneration does not occur in SARM1 null mice after a single hit TBI (13).

Intranasal delivery of drugs into the brain has great clinical potential due to (1) simplicity of administration, (2) noninvasiveness, (3) relatively rapid CNS delivery, (4) ability to repeat dosing easily,

(5) non-necessity of drug modification, (6) minimal systemic exposure and (7) avoidance of the liver and gut first-pass metabolism. Neuroprotective cytokines similar in size to LIF (20 kDa), including BDNF (14 KDa), CNTF (23 kDa), NGF (13 kDa) VEGF (38 kDa) and HB-EGF (22 kDA) have been successfully introduced into the brain via IN administration (14). Other studies have demonstrated that IN delivery of IGF-1 and EGF promote the proliferation and migration of SVZ progenitors in models of demyelination (15, 16). While most of these studies have been performed using rodents, studies have demonstrated the efficacy of intranasal delivery in non-human primates (17, 18) and humans(14, 19, 20).

LIF is the most pleiotropic member of the interleukin-6 family of cytokines, whose expression is induced by central nervous system (CNS) injury (21, 22). LIF haplodeficient mice exhibit desynchronized glial responses, increased neurodegeneration, decreased corpus callosum (CC) conductivity and motor deficits after a mTBI (23). LIF has been shown to peak in the acute/subacute phase (at 48 hours) after a mTBI (23), suggesting its possible involvement in the repair and regenerative processes within the injured brain. Given the beneficial role of LIF during the acute/subacute recovery phase after CNS injury and its pleiotropic effects in the brain (24), we hypothesized that IN-LIF treatment would prevent neurodegeneration when administered during the chronic period of recovery from an mTBI. We initiated these studies to test that hypothesis.

Methods

All experiments were performed in accordance with research guidelines set forth by the Institutional Animal Care and Use Committee of Rutgers New Jersey Medical School (Protocol #999900840) and were in accordance with the National Institute of Health Guide for the Care and Use of Laboratory Animals (publication No. 80-23) revised in 1996 and the ARRIVE guidelines.

Male CD1 mice were randomly assigned to sham or injured groups. The mTBI model that we used entails administering a single closed head injury. This model was developed in Dr. Regina Armstrong's laboratory, as described in Mierzwa et al., 2015 (25). Briefly, 2-month-old mice were administered 0.5

mg/kg Buprenorphine HCL sustained release prophylactically and then anesthetized with 2.5% isoflurane. Once they were confirmed to be deeply anesthetized, the mice were administered Bupivacaine (2.0 µg/g) subcutaneously on the scalp and then a 1 cm midline incision was made to expose the skull. After deflecting the periosteum the animals were placed in a mouse stereotactic device with ear cuffs. The cortical impact device (Custom Design & Fabrication, Richmond, VA), outfitted with a 3-mm-diameter flat metal impactor tip was zeroed on the sagittal suture just in front of Bregma. Twenty seconds after removing the anesthesia, the impactor was driven perpendicularly onto the exposed sagittal suture at a velocity of 4.0m/sec to a depth of 1.3 mm farther than the zero point with a dwell of 100 ms. The mouse was removed from the stereotactic device and the righting time and apnea times were recorded. The animals were re-anesthetized so that the scalp incision could be closed. The mice were then administered 0.9% saline (3% of body weight), i.p. Sham-injured mice were subjected to the same procedures without receiving an impact. All animals were allowed to recover on a heating pad set at 37°C, and upon becoming ambulatory were returned to their home cages.

Mice were lightly anesthetized with isofluorane so that they would lie on their backs. Mice were administered either 20 µL of recombinant murine LIF (rmLIF) (EMD Millipore, cat #LIF2010) or water using a fine plastic pipette in 2 ul doses. These substances were administered to alternate nostrils within 10 minutes. Two doses of rmLIF were administered, either 40 ng or 100 ng, administered twice daily and 5 days/week for two consecutive weeks during the 6-8 weeks following mTBI.

Sensorimotor behavior

Mice were evaluated using the Horizontal Ladder with irregular rungs which we have found to be a very sensitive test of sensorimotor function (26). They were assessed at 4 and 8 weeks post-mTBI. Mice were trained to ambulate across a horizontal ladder with irregularly spaced rungs for 3 days prior to testing **(Fig 1A)**. The task was conducted 3 times non-consecutively per session for each mouse. Video recordings

were made and scored by an individual blinded to the group identity of the subject. The mean number of missteps and the average latency to cross the length of the apparatus were determined as the dependent measures for sensorimotor function. After testing, the mice were perfused with PBS pH 7.4 containing 4% paraformaldehyde (PFA). The brains were removed and stored in 4% PFA at 4^oC overnight and then changed to PBS.

Magnetic Resonance Imaging

Two weeks prior to the magnetic resonance imaging (MRI) scans brain samples were washed and stored in fresh PBS containing 0.05% sodium azide, which was changed every week to wash out the fixative to prevent interference with MRI relaxation times (27). One hour before the scans, brains were washed three times for 10 min in 10 mL PBS to remove the PFA solution. Then, the brains were placed into a custom-built MRI-compatible tube filled with Fluorinert, an MRI susceptibility-matching fluid that generates no MRI signal (Sigma-Aldrich, Inc., St. Louis, MO). Microstructural integrity was assessed using ex-vivo diffusion tensor imaging (DTI) of the brain samples using a horizontal bore 9.4T scanner (Bruker BioSpin, Billerica, MA). . Proper alignment of each brain sample was performed by keeping the distance from the opening of the tube consistent across all specimens during positioning in the birdcage [1]H radio frequency coil (15 mm diameter positioning and power optimization were optimized through Bruker defined gradient echo (GE) and fast spin echo (FSE) sequences. Shimming was done by second-order shim with cubic voxel to bring the H20 line width to less than 30Hz. A fast low-angle shot (FLASH 3D) spin echo sequence was used for the T1 images with an FOV of 12x14mm and matrix 120x40, containing 80 slices of thickness of 0.1 mm, TR/TE=50/6 ms. The rapid acquisition with relaxation enhancement (RARE) sequence was used for T2 weighted imaging with a FOV of 12x8 mm and matrix 102x64, containing 64 slices of thickness 0.24 mm, TR/TE=3354/24 ms, and number of echoes=8. DTI acquisitions were performed using a Stejskal-Tanner spin-echo diffusion-weighted with a diffusion

gradient duration of 4ms and a delay between the two diffusion gradients of 9.3ms. 64 contiguous slices of 240μm thickness were acquired with a field of view of 12 mm×8 mm with a matrix of 102×67, resulting in an in-plane resolution of 120μm × 120μm. TR/TE was 2000/20m with 30 diffusion weighted images and 5 images with no diffusion weighting were acquired for each slice. The diffusion gradient directions for the diffusion weighted images were non-collinear and had the same b-value, 1000 s/mm^2. Eight averages were acquired with a total experiment time of 10 hours and 24 minutes.

Histological Analyses

Immunofluorescence was performed as previously described (23). No signal above background was obtained when the primary antibodies were replaced with pre-immune sera. The antibodies used for these studies are listed in Table 1. Images were collected using a Q-imaging Retiga-2000R CCD camera (Surrey, BC, Canada) on an Olympus AX70 microscope (Center Valley, PA, USA). Images were analyzed using FiJi software in automated batch mode using a custom script. Briefly, to measure intensity, RGB channels were split, the background was subtracted by applying a rolling ball radius, and the individual channel intensity was read. For % Area and particle count analyses, images were further thresholded, converted to mask, and outlined. Then the "Analyze particles" command was run defining size (0-infinity) and circularity (0-1) that produced a summary file with particle count and % Area values. Images were randomly checked in all 3 groups to ensure the set parameters did not result in false positives or negatives. Images quantified for Iba1 positive microglia and Iba1/CD68 double positive microglia were coded and manually counted. Cells were classified as CD68+ if they had 3 or more bright CD68+ puncta inside of them. White matter astrocytes were classified as ADK+/GFAP+ if they had a stellate morphology and appeared yellow on the overlay of the green fluorescence for ADK and red fluorescence for GFAP. Images quantified for ADK+/GFAP+ were also coded and manually counted.

Statistical analyses

All sham T2 weighted images were registered to the mouse Franklin and Paxinos MRI atlas and averaged to obtain a study anatomical template (28). Subsequently all anatomical T1, T2 and DTI images were spatially registered to the anatomical study brain template using non-linear registration in Bioimage Suite (29). The resulting brain matrix volume data were used to calculate the DTI eigenvalues (λ_1, λ_2, λ_3) and vectors (e_1, e_2, e_3). At each voxel, the eigenvalues represented the magnitude of diffusion and eigenvectors represented maximal and minimal diffusion directions respectively. The fractional anisotropy (FA) and mean diffusion (MD) were estimated from the eigen values (30, 31) using the BioImage Suite software (29).

Statistical parametric mapping of the whole brain was performed in a voxel-wise manner using a two sample t-test (two-tailed) using AFNI (32) . Statistical differences between experimental groups were analyzed with a corrected threshold probability of $P<0.05$, determining voxels with a significant difference (active voxels) using a family wise error control of a minimum cluster of 30 contiguous voxels to correct for multiple comparisons. MRI intensity differences obtained from the t-test means were depicted in color, with hot colors indicating a positive mean difference, and cold colors depicting a negative mean difference, which were overlayed on the mean anatomical MRI of the control group animals. Regions of interest (ROI) analysis was manually performed by drawing regions using the Franklin Paxinos mouse stereotactic atlas (28). Average DTI parameters across each ROI were determined across each animal subject in the respective experimental groups. Group-level statistical comparisons were performed using a one-way ANOVA followed by a post-hoc Tukey's HSD test with a significance of $p<0.05$ required for significance.

Results

Delayed LIF treatment prevents sensorimotor behavior debilitation after mTBI

Horizontal ladder tests were performed at 4 and 8 weeks post-mTBI in naïve, sham, and mTBI mice with vehicle or LIF. Between 4 weeks and 8 weeks after a mTBI the motor performance of the mice that were administered the vehicle showed significant functional deterioration on the horizontal ladder as evidenced by increased numbers of foot slips (**Fig 1B** and **1C**; blue). While only 10% of animals showed an average of >2 foot slips at 4 weeks post-mTBI (**Fig 1B;** blue), 50% of animals showed >2 foot slips at 8 weeks post-injury (**Fig 1C;** blue), indicating delayed sensorimotor deterioration. Remarkably, all of the mTBI mice treated with LIF showed <2 foot slips at 8 weeks post-injury, a level of motor performance that was no worse than the sham injured mice and significantly better ($p < 0.01$) than the vehicle administered mTBI mice (**Fig 1C**; red). Catwalk gait analysis showed no differences in any of the evaluated parameters (**Fig 1E-F**). For example, there was no difference in mean body speed or regularity index between groups at 8 weeks post-injury (**Fig 1D** and **1E**).

We considered using a repeated mild TBI model using the parameters described by Hall et al., 2016 (33). In initial studies using their repeated mild TBI model we documented that the mice subjected to a repeated mild TBI had impaired motor performance 3 days after the last concussive injury (**Fig S1**) and that these mice had both increased astrogliosis in both the gray matter and white matter as determined by increased GFAP (**Fig S2)** and decreased adenosine kinase (**Fig S3**), along with increased microgliosis as determined by increased percentage of CD68+/Iba1+ cells in the injured neocortex (**Fig S4**). However, when their motor performance was evaluated over 6 weeks of recovery, they showed almost full recovery (**Fig S5**). Since the goal of the current studies was to evaluate the neurorestorative properties of intranasal LIF during the chronic period of recovery from mTBI, we did not pursue studies using the repeated mild TBI model further as a single hit model was sufficient.

Evidence of LIF neuroprotection from T1 and T2 contrast MRI changes after mTBI

After completing the behavioral tests, the mice were perfused with paraformaldehyde, and their brains were removed. They were washed in PBS for 2 weeks to remove the paraformaldehyde before ex-vivo MRI measures. Conventional MRI T1 and T2 weighted acquisitions were obtained using the FLASH and RARE sequences, respectively (**Fig S6**). Comparing the T1 contrast changes between the mTBI+vehicle treated versus sham group, significant clusters of T1 decrease were observed across the neocortex proximal to the TBI epicenter along with diffuse clusters of T1 decreases within the striatum, basal brain areas and the corpus callosum (**Fig 2A**). Relatively fewer T2 intensity changes were detected after mTBI, with T2 decreases observed in the neocortex adjacent to the mTBI epicenter compared to sham (**Fig 2D**). LIF treatment significantly reduced the mTBI-induced T1 intensity decreases across most frontal regions of the brain and adjacent to the mTBI epicenter, in addition to increasing the T1 intensity across the basal forebrain, somatosensory and entorhinal cortices (**Fig 2B**). T2 intensity decreases observed after mTBI were absent with LIF treatment, and increased T2 was observed across the frontal corpus callosum (**Fig 2E**). The areas affected by LIF were determined by comparing mTBI groups treated with vehicle versus LIF. As shown in **Fig 2C**, the basal forebrain, somatosensory cortex, and lateral ventricles showed significantly reduced T1 intensity in mTBI+vehicle treated mice compared to mTBI+LIF mice. T2 was relatively less sensitive than T1, and the LIF effects were only evident in the lateral ventricular space in the mTBI+LIF treated vs. the mTBI+vehicle treated mice (**Fig 2F**). Sparse T2 decreases were observed across the corpus callosum in the mTBI+vehicle mice compared to mTBI+LIF mice (**Fig 2F**), signifying the protective effect of LIF over the corpus callosal area.

LIF partially restores callosal DTI changes after mTBI

Following the conventional T1 and T2 acquisitions, diffusion-sensitive acquisition was performed, and DTI parameters such as the fractional anisotropy (FA) and mean diffusivity (MD) were estimated across the whole brain (**Fig 3**). Comparing group differences between mTBI mice treated with vehicle and sham

mice, FA decreased predominantly in the corpus callosum and other subcortical white matter areas (**Fig 3A**; vehicle vs sham). MD increased between vehicle-treated mTBI mice and sham mice and was prominent across the neocortex adjoining the impact epicenter and deeper areas such as the corpus callosum, subventricular zone, and the striatum (**Fig 3B**; vehicle vs sham). LIF treatment inhibited the mTBI-induced FA decrease in a dose-dependent manner (**Fig 3A**; LIF-40 vs sham and LIF-100 vs sham). However, sparse increases in FA were observed in certain deeper areas of the brain such as the internal capsule (IC) and anterior commissure pars posterior (ACP), possibly owing to the proximity of the intranasal pathways and higher LIF dose (**Fig 3A**; LIF-100 vs sham). LIF treatment inhibited the mTBI-induced MD increases in a dose-dependent manner (**Fig 3B**; LIF-40 vs. sham and LIF-100 vs. sham). However, higher LIF dose increased MD in the subventricular zone and deeper brain regions (**Fig 3B; LIF-100 vs sham**).

A region of interest (ROI) analysis involving 8 ROIs was performed over the different animal groups. Significant FA decreases were observed across the corpus callosum (CC) and prefrontal cortex (PFC) in the mTBI vehicle group (**Fig 4A,B,** $p < 0.001$). Significant increases in MD also were observed in the CC and the PFC (**Fig 4C,D,** $p < 0.01$). No significant changes in FA or MD occurred across 6 other ROIs tested (**Figs S6** and **Fig S7**). However, an increasing trend in MD was observed in the motor cortex, primary somatosensory cortex, striatum, hippocampus, insula, and thalamus (**Fig S7**). LIF treatment, in a dose-dependent manner, reversed the mTBI-induced FA decrease across the corpus callosum and the PFC, respectively (**Fig 4A,B,** $p < 0.001$). However, LIF at the lower dose of 40 ng restored the mTBI-induced MD increase observed across the corpus callosum and in the PFC, with no further improvements with the 100 ng dose (**Fig 4C,D,** $p < 0.01$). When FA intensity was plotted against the number of foot slips on the horizontal ladder for Vehicle Rx TBI mice vs. LIF Rx TBI mice a Pearson's correlation analysis showed that while the slopes were not different ($p = 0.266$), the elevations were statistically different ($p = 0.0013$) (**Fig 4E**).

Compared to the sham mice, TBI mice showed decreased FA across the corpus callosum and other white matter tracts and increased FA across neocortical and subcortical gray matter regions **(Fig 5A,B)**. When the mTBI mice were treated with 40 ng LIF, the FA decrease across the corpus callosum was inhibited and the neocortical and subcortical gray matter FA increases were also prevented **(Fig 5C)**. To ascertain the underlying cellular changes representing these FA increases in the neocortex, the brains after the ex-vivo DTI studies were stained for GFAP as an index of astroglial reactivity. Astroglial scars were evident at the dorsal edges of the site of impact from the mTBI, which was mitigated by 40 ng LIF treatment **(Fig 5D)**, linking glial reactivity to the observed neocortical FA increases around the mTBI epicenter and subcortical gray matter areas.

IN-LIF protects against delayed axonal degeneration following mTBI

Axonal transport is compromised in damaged axons leading to the accumulation of amyloid precursor protein (APP) in axonal swellings along the length of the axon. Additionally, axons can be transected during TBI, producing end-bulbs (34). Therefore, we evaluated the extent of axonal damage following mTBI by immunostaining coronal sections using the Y188 antibody that is highly specific for APP (35). Negligible APP staining was seen in the CC in the sham group **(Fig 6A)**. By contrast, APP+ puncta accumulated within CC fibers of the IN-vehicle group as distinct swellings or end bulbs **(Fig 6B)**. 40 ng IN-LIF Rx decreased the number of APP swellings and endbulbs **(Fig 6C)**. A particle analysis showed significantly increased numbers of APP+ particles in the CC of the vehicle group over sham, which was almost entirely reversed by IN-LIF **(Fig 6D**, *$p<0.05$). To determine whether IN-LIF would also improve the myelin integrity, sections were stained for MBP and the thickness of the corpus callosum was measured **(Fig 6E-H)**. This analysis revealed a 30% decrease in the CC thickness in the TBI-vehicle group vs. the sham group ($p <0.001$) with a 17% decrease for the IN-LIF Rx mice that was not statistically different from the sham group **(Fig 6H)**.

Delayed IN-LIF attenuates astrocyte activation in the injured subcortical WM after mTBI.

GFAP has long been used as a marker for reactive gliosis, and GFAP expression was significantly increased in the subcortical WM of the vehicle treated group **(Fig 7B,D,** $p < 0.001$), but GFAP is a structural protein that does not provide information about the functional state of the astrocytes. Studies using the controlled cortical impact model have shown that TBI causes the overexpression of ADK in reactive astrocytes, a major adenosine metabolizing enzyme. Increases in ADK will result in chronic adenosine deficiency which can decrease the threshold for seizures (36). Therefore, we stained sections for ADK and evaluated the effect of 40 ng IN-LIF Rx on its expression. The subcortical white matter of the sham group showed low ADK immunofluorescence **(Figs 7A,E.** By contrast, ADK was strongly expressed in the cytoplasm of hypertrophic GFAP+ astrocytes in the vehicle Rx mTBI group **(Fig 7B,E)**. IN-LIF decreased levels of ADK to that seen in the sham injured group which was significantly reduced from those seen in the vehicle-treated group **(Fig 7C,E,** $p < 0.01$). Not only did IN-LIF Rx decrease the overall level of ADK, but it reduced the percentage of astrocytes that were ADK+ to that seen in the sham-injured mice **(Fig 7I,** $p < 0.01$ vs. vehicle).

Delayed IN-LIF attenuates macrophage/microglial activation in the injured white matter after mTBI

In our studies of mTBI in LIF heterozygous juvenile mice, we observed delayed and amplified microgliosis after injury (23). Therefore, we hypothesized that IN LIF would repress microgliosis. To test this hypothesis, we immunostained coronal sections for Iba1 and CD68. The sham injured white matter was populated by ramified IBA1+ cells with few CD68+ puncta **(Fig 7F)**. As expected, the subcortical WM of the vehicle-treated mTBI group was populated by IBA1+ cells that also contained multiple lysosomes that labeled for CD68 **(Fig 7G)**. By contrast, most of the microglia in the 40 ng IN-LIF Rx brains had a ramified morphology and they contained few CD68+ lysosomes **(Fig 7H)**. When we

quantified the percentage of CD68+/IBA1+ cells in the subcortical WM, on average $30 \pm 9.7\%$ of the microglia were double positive in the vehicle-treated mTBI group. In contrast, less than $2.3 \pm 2.6\%$ were double positive in the sham injured group and there were $7.8 \pm 3.6\%$ double positive in the LIF Rx group ($p<0.05$ comparing vehicle vs IN-LIF) **(Fig 7J)**.

Discussion

Histopathological analyses after mTBI have revealed both acute as well as chronic phases of neurodegeneration. To date the majority of pre-clinical studies have focused on strategies to reduce acute brain injury (9) with a paucity of studies evaluating the mechanisms of delayed neurodegeneration and its treatment. In the single hit model that we have employed, the subcortical white matter appears normal for up to 6 weeks after the injury. Still, tertiary neurodegeneration becomes apparent by 8 weeks of recovery. Accompanying the Wallerian degeneration that ensues the compound action potential becomes weak and electron microscopic analyses support the conclusion that the axons are undergoing pathological changes that precede any changes in myelin maintenance (12). Pediatric, juvenile and adult TBI studies have all revealed delayed white matter injury. There are several hypotheses for this late-onset neurodegeneration. One hypothesis is that the white matter becomes hypoxic which triggers the delayed white matter loss (37). Alternatively, the Wallerian degeneration could result from SARM1 activation as delayed axonal degeneration does not occur in SARM1 null mice after either mTBI or repeated head injury (8, 13, 38). While the trigger for the delayed neurodegeneration seen in this model remains elusive, our studies have reproduced the delayed white matter neurodegeneration that is seen after single hit concussive injury as revealed by high-resolution DTI, histopathology for axonal swellings and progressive sensorimotor deficits. Notably, all of these aspects of tertiary neurodegeneration and the accompanying astrogliosis and microgliosis subsequent to the mTBI could be largely prevented (or reversed) by administering LIF intranasally twice daily starting at 6 weeks of recovery.

LIF broadly affected the injured brain, and its effects were especially prominent at the impact epicenter and in specific deeper brain structures as visualized by DTI. Moreover, LIF treatment inhibited the mTBI-induced FA decrease, the MD increase, and there was a strong trend for it to reduce the sensorimotor deficits in a dose dependent manner. While the LIF had a widespread neuroprotective effect, it appeared to more strongly affect basal brain areas **(Fig 2B)**, likely because those regions have more robust structural connections with the olfactory bulb, which may have resulted in greater bioavailability of LIF in these areas, especially with exposure to the higher dose of LIF (**Fig 3B**); further studies are required to ascertain this interpretation.

A novel finding in the present study was that FA and MD changed across the prefrontal cortex 8 weeks after an mTBI. Furthermore, MD showed a trend for increasing across several gray matter brain regions, indicating milder effects across a large area of gray matter. This increasing trend in MD was significantly reversed by LIF treatment across all of these regions. The prefrontal cortex plays a vital role in social cognition in humans and across animal models (39) . Changes in social cognition have been observed after TBIs (40) stemming from post-injury frontal lobe changes and prefrontal cortical reorganization (41). While our behavioral paradigms did not test for differences in social cognition after mTBI, the current DTI findings, indicating a significant impact on the prefrontal cortex, warrant behavioral tests of social cognition during the delayed phase of recovery from a mTBI.

An additional finding was the formation of glial scars across the cortical gray matter adjoining the mTBI epicenter (ascertained by GFAP staining; **Fig 5D**), which manifested as areas of increased FA at the margins of the TBI epicenter **(Fig 5B;** arrows**)**. The mTBI-induced glial scar formation was prevented after treatment with 40 ng LIF **(Fig 5C)**. Within the white matter, the IN-LIF Rx mice also showed reduced astrogliosis as illustrated by fewer GFAP+/ADK+ stellate astrocytes. As ADK overexpression will reduce levels of adenosine, which lowers the threshold for epileptic seizures through insufficient

activation of adenosine A_1 receptors (36), our data show that not only is LIF reducing the formation of the glial scar but also enhancing a key function of the astrocytes that should reduce seizure incidence.

Microglia become activated rapidly after brain injuries and produce chemokines that attract macrophages, contributing to secondary and tertiary brain injury (42, 43). Therefore, we assessed the functional state of the microglia/macrophages using antibodies to Iba-1 and CD-68. CD-68 is a lysosomal protein that participates in antigen processing in late endosomes and the degradation of proteins in lysosomes. Therefore, it is a useful marker for highly phagocytic, activated microglia. As the microglia were significantly less reactive using delayed 40 ng IN LIF treatment, a reasonable conclusion is that the production and release of cytokines and chemokines from the microglia has been suppressed and, therefore, they are no longer stoking an inflammatory response. Indeed, if LIF has shifted the microglia towards an M2-like phenotype, then they are likely releasing growth factors and trophic factors that are facilitating repair. Reminiscent of our results, systemically administered LIF reduces the numbers of activated macrophage/microglia in the ischemic rat brain and LIF has been shown to reduce macrophage IL-12 levels in vitro (44).

Conclusions

Using T1/T2 MRI and DTI imaging we have obtained clinically relevant markers of gray matter and white matter changes across the brain in a mouse model of mTBI. The deterioration of sensorimotor function seen between 6 and 8 weeks of recovery correlated with white matter loss across the corpus callosum and in the PFC. Combining behavioral analyses, histology and translational imaging, our studies document the restorative effects of intranasally administered LIF. The dose-dependent effects of LIF in improving white matter integrity suggest that strategies that can increase LIF bioavailability (e.g., nanoparticle encapsulation) may improve its translational and clinical trial prospects as a treatment for preventing tertiary neurodegeneration.

Transparency, Rigor and Reproducibility Summary

Sample size 6-8 mice per group, were based on our preliminary studies with observed differences between means (>1.5 fold), observed standard deviations (40% of lowest mean) which indicated >80% power to detect both an overall significant effect by ANOVA and a post-hoc difference between the experimental groups vs. control with $p<0.05$ after correction for multiple comparisons. Thirty mice were subjected to the single hit mTBI model and 14 mice were subjected to the repeated mTBI protocol. None were excluded for technical reasons and 2 mice died immediately after the mTBI and one mouse died after the rmTBI. TBI mice were randomly assigned to vehicle or LIF Rx groups. The investigators who performed the behavioral tests were blinded to the group identity as were the investigators performing the MRI analyses. The normalality of the data behavioral and histological outcome data were evaluated using Shapiro-Wilk tests. The preprocessed MRI and DTI data sets cannot be shared in public due to ongoing therapeutic development of LIF, but data will be made available to readers upon receipt of a request by the corresponding author.

Contributors: Conceptualization: SWL, SSK and FH; funding acquisition: SWL, SSK and FH; project supervision: SWL, SSK and FH; experimental data acquisition: VD, AM, SA, MF, JM, BGS, DC and YW; behavioral data analysis: VD, AM and SWL, Immunofluorescence analyses: VD, MF and SWL MR image processing and analysis, JM, BGS and SSK, writing original manuscript draft: VD, SWL and SSK. Reviewing and editing revised versions: SWL, SSK and FH.

Author's Disclosure: The Authors declare that they have no conflicts of interest.

Funding: Supported by R21 NS125201, which was awarded to SWL, SK, and FH, and Rutgers Busch Biomedical Grant IRES 21-002946 to SWL and SK and CBIR25IRG016 awarded to SWL from the New Jersey Commission on Brain Injury Research.

References

1.   Prigatano GP, Gale SD. The current status of postconcussion syndrome. Curr Opin Psychiatry. 2011;24(3):243-50.
2.   Elder GA. Update on TBI and Cognitive Impairment in Military Veterans. Curr Neurol Neurosci Rep. 2015;15(10):68.
3.   McKee AC, Mez J, Abdolmohammadi B, Butler M, Huber BR, Uretsky M, et al. Neuropathologic and Clinical Findings in Young Contact Sport Athletes Exposed to Repetitive Head Impacts. JAMA Neurology. 2023.
4.   Zohar O, Rubovitch V, Milman A, Schreiber S, Pick CG. Behavioral consequences of minimal traumatic brain injury in mice. Acta Neurobiol Exp (Wars). 2011;71(1):36-45.
5.   Siopi E, Calabria S, Plotkine M, Marchand-Leroux C, Jafarian-Tehrani M. Minocycline restores olfactory bulb volume and olfactory behavior after traumatic brain injury in mice. J Neurotrauma. 2012;29(2):354-61.
6.   Siopi E, Llufriu-Dabén G, Fanucchi F, Plotkine M, Marchand-Leroux C, Jafarian-Tehrani M. Evaluation of late cognitive impairment and anxiety states following traumatic brain injury in mice: the effect of minocycline. Neurosci Lett. 2012;511(2):110-5.
7.   Milman A, Zohar O, Maayan R, Weizman R, Pick CG. DHEAS repeated treatment improves cognitive and behavioral deficits after mild traumatic brain injury. Eur Neuropsychopharmacol. 2008;18(3):181-7.
8.   Bradshaw DV, Jr., Knutsen AK, Korotcov A, Sullivan GM, Radomski KL, Dardzinski BJ, et al. Genetic inactivation of SARM1 axon degeneration pathway improves outcome trajectory after experimental traumatic brain injury based on pathological, radiological, and functional measures. Acta neuropathologica communications. 2021;9(1):89.
9.   Huang TQ, Song JN, Zheng FW, Pang HG, Zhao YL, Gu H, et al. Protection of FK506 against neuronal apoptosis and axonal injury following experimental diffuse axonal injury. Mol Med Rep. 2017;15(5):3001-10.
10.   Scott G, Zetterberg H, Jolly A, Cole JH, De Simoni S, Jenkins PO, et al. Minocycline reduces chronic microglial activation after brain trauma but increases neurodegeneration. Brain. 2018;141(2):459-71.
11.   Armstrong RC, Sullivan GM, Perl DP, Rosarda JD, Radomski KL. White matter damage and degeneration in traumatic brain injury. Trends Neurosci. 2024;47(9):677-92.
12.   Marion CM, Radomski KL, Cramer NP, Galdzicki Z, Armstrong RC. Experimental Traumatic Brain Injury Identifies Distinct Early and Late Phase Axonal Conduction Deficits of White Matter Pathophysiology, and Reveals Intervening Recovery. J Neurosci. 2018;38(41):8723-36.
13.   Marion CM, McDaniel DP, Armstrong RC. Sarm1 deletion reduces axon damage, demyelination, and white matter atrophy after experimental traumatic brain injury. Exp Neurol. 2019;321:113040.
14.   Meredith ME, Salameh TS, Banks WA. Intranasal Delivery of Proteins and Peptides in the Treatment of Neurodegenerative Diseases. AAPS J. 2015;17(4):780-7.
15.   Cantarella C, Cayre M, Magalon K, Durbec P. Intranasal HB-EGF administration favors adult SVZ cell mobilization to demyelinated lesions in mouse corpus callosum. Developmental neurobiology. 2008;68(2):223-36.
16.   Scafidi J, Hammond TR, Scafidi S, Ritter J, Jablonska B, Roncal M, et al. Intranasal epidermal growth factor treatment rescues neonatal brain injury. Nature. 2014;506(7487):230-4.
17.   Putnam PT, Roman JM, Zimmerman PE, Gothard KM. Oxytocin enhances gaze-following responses to videos of natural social behavior in adult male rhesus monkeys. Psychoneuroendocrinology. 2016;72:47-53.
18.   Parris BA, Dienes Z, Bate S, Gothard S. Oxytocin impedes the effect of the word blindness post-hypnotic suggestion on Stroop task performance. Soc Cogn Affect Neurosci. 2014;9(7):895-9.
19.   Mistry A, Stolnik S, Illum L. Nanoparticles for direct nose-to-brain delivery of drugs. International journal of pharmaceutics. 2009;379(1):146-57.
20.   Craft S, Baker LD, Montine TJ, Minoshima S, Watson GS, Claxton A, et al. Intranasal insulin therapy for Alzheimer disease and amnestic mild cognitive impairment: a pilot clinical trial. Arch Neurol. 2012;69(1):29-38.
21.   Nicola NA, Babon JJ. Leukemia inhibitory factor (LIF). Cytokine & growth factor reviews. 2015;26(5):533-44.
22.   Banner LR, Moayeri NN, Patterson PH. Leukemia inhibitory factor is expressed in astrocytes following cortical brain injury. Exp Neurol. 1997;147(1):1-9.
23.   Goodus MT, Kerr NA, Talwar R, Buziashvili D, Fragale JE, Pang KC, et al. Leukemia Inhibitory Factor Haplodeficiency Desynchronizes Glial Reactivity and Exacerbates Damage and Functional Deficits after a Concussive Brain Injury. J Neurotrauma. 2016;33(16):1522-34.

24.   Bauer S, Patterson PH. Leukemia inhibitory factor promotes neural stem cell self-renewal in the adult brain. J Neurosci. 2006;26(46):12089-99.

25.   Mierzwa AJ, Marion CM, Sullivan GM, McDaniel DP, Armstrong RC. Components of myelin damage and repair in the progression of white matter pathology after mild traumatic brain injury. J Neuropathol Exp Neurol. 2015;74(3):218-32.

26.   D'Mello V, Subramaniam M, Bhalla AP, Saavedra S, Leiba O, Levison SW. Intranasal Leukemia Inhibitory Factor Attenuates Gliosis and Axonal Injury and Improves Sensorimotor Function After a Mild Pediatric Traumatic Brain Injury. Neurotrauma Rep. 2023;4(1):236-50.

27.   Thelwall PE, Shepherd TM, Stanisz GJ, Blackband SJ. Effects of temperature and aldehyde fixation on tissue water diffusion properties, studied in an erythrocyte ghost tissue model. Magn Reson Med. 2006;56(2):282-9.

28.   Franklin KBJ, Paxinos G. Paxinos and Franklin's the Mouse Brain in Stereotaxic Coordinates, Compact: The Coronal Plates and Diagrams: Elsevier Science; 2019.

29.   Papademetris X, Jackowski MP, Rajeevan N, DiStasio M, Okuda H, Constable RT, et al. BioImage Suite: An integrated medical image analysis suite: An update. Insight J. 2006;2006:209.

30.   Basser PJ, Jones DK. Diffusion-tensor MRI: theory, experimental design and data analysis - a technical review. Nmr in Biomedicine. 2002;15(7-8):456-67.

31.   Basser PJ, Pierpaoli C. Microstructural and physiological features of tissues elucidated by quantitative-diffusion-tensor MRI. J Magn Reson Ser B. 1996;111(3):209-19.

32.   Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance neuroimages. Comput Biomed Res. 1996;29(3):162-73.

33.   Bolton Hall AN, Joseph B, Brelsfoard JM, Saatman KE. Repeated Closed Head Injury in Mice Results in Sustained Motor and Memory Deficits and Chronic Cellular Changes. PLoS One. 2016;11(7):e0159442.

34.   Tang-Schomer MD, Johnson VE, Baas PW, Stewart W, Smith DH. Partial interruption of axonal transport due to microtubule breakage accounts for the formation of periodic varicosities after traumatic axonal injury. Exp Neurol. 2012;233(1):364-72.

35.   Guo Q, Li H, Gaddam SS, Justice NJ, Robertson CS, Zheng H. Amyloid precursor protein revisited: neuron-specific expression and highly stable nature of soluble derivatives. J Biol Chem. 2012;287(4):2437-45.

36.   Aronica E, Sandau US, Iyer A, Boison D. Glial adenosine kinase--a neuropathological marker of the epileptic brain. Neurochem Int. 2013;63(7):688-95.

37.   Parikh U, Williams M, Jacobs A, Pineda JA, Brody DL, Friess SH. Delayed Hypoxemia Following Traumatic Brain Injury Exacerbates White Matter Injury. J Neuropathol Exp Neurol. 2016;75(8):731-47.

38.   Maynard ME, Redell JB, Zhao J, Hood KN, Vita SM, Kobori N, et al. Sarm1 loss reduces axonal damage and improves cognitive outcome after repetitive mild closed head injury. Exp Neurol. 2020;327:113207.

39.   Bicks LK, Koike H, Akbarian S, Morishita H. Prefrontal Cortex and Social Cognition in Mouse and Man. Front Psychol. 2015;6:1805.

40.   Milders M. Relationship between social cognition and social behaviour following traumatic brain injury. Brain Inj. 2019;33(1):62-8.

41.   Dall'Acqua P, Johannes S, Mica L, Simmen HP, Glaab R, Fandino J, et al. Prefrontal Cortical Thickening after Mild Traumatic Brain Injury: A One-Year Magnetic Resonance Imaging Study. J Neurotrauma. 2017;34(23):3270-9.

42.   Babcock AA, Kuziel WA, Rivest S, Owens T. Chemokine expression by glial cells directs leukocytes to sites of axonal injury in the CNS. J Neurosci. 2003;23(21):7922-30.

43.   Abe N, Choudhury ME, Watanabe M, Kawasaki S, Nishihara T, Yano H, et al. Comparison of the detrimental features of microglia and infiltrated macrophages in traumatic brain injury: A study using a hypnotic bromovalerylurea. Glia. 2018;66(10):2158-73.

44.   Davis SM, Collier LA, Winford ED, Leonardo CC, Ajmo CT, Jr., Foran EA, et al. Leukemia inhibitory factor modulates the peripheral immune response in a rat model of emergent large vessel occlusion. J Neuroinflammation. 2018;15(1):288.

Antibody	Host	Application and dilution	Source	Catalog #
Glial fibrillary acidic protein (GFAP)	Rabbit	Immunofluorescence, 1:200; Western blot, 1:1000	Dako, Santa Clara, CA	Z0334
Amyloid precursor protein (APP) Y188	Rabbit	Immunofluorescence, 1:200	Abcam, Cambridge, MA	ab32136
Ionized calcium binding adaptor molecule 1 (IBA1)	Rabbit	Immunofluorescence, 1:200	Wako Pure Chemical Corporation, Wako, TX	019-19741
CD-68	Rat	Immunofluorescence, 1:200; Western blot, 1:1000	Bio-Rad, San Diego, CA	MCA1957GA
Adenosine Kinase	Rabbit	Immunofluorescence, 1:500	Fisher Scientific, Pittsburgh PA	501567320
RedX-conjugated anti-mouse IgG	Donkey	Immunofluorescence, 1:300	Jackson ImmunoResearch, West Grove, PA	715-295-150
Alexa 488-conjugated anti-rabbit IgG	Donkey	Immunofluorescence, 1:300	Jackson ImmunoResearch, West Grove, PA	711-545-152
Alexa 594-conjugated anti-rabbit	Donkey	Immunofluorescence, 1:300	Jackson ImmunoResearch, West Grove, PA	711-585-152
Alexa 488-conjugated donkey anti-mouse IgG	Donkey	Immunofluorescence, 1:300	Jackson ImmunoResearch, West Grove, PA	715-545-150

Table 1. List of antibodies used for immunofluorescence.

Figures

Figure 1. A. IN-LIF Rx improves motor behavior in mTBI mice. The pattern of the irregular horizontal ladder used to quantify fine motor function. Mice were trained for 3 days on the horizontal ladder before performing the test. **B, C.** Average number of foot slips after 3 non-consecutive runs on the horizontal ladder 4 weeks after the mTBI but before treatment and 8 weeks after the mTBI following 2 weeks of vehicle or 40ng LIF treatments between 6-8 weeks post-TBI. (*p<0.05, n=4-8) by one-way ANOVA followed by Tukey's multiple comparison test. **D, E.** Catwalk gait analysis measurements showed no differences in mean body speed or regularity index at 8 weeks post-mTBI between groups; one-way ANOVA followed by Tukey's multiple comparison test, n=4-8 per group.

Fig 2. IN-LIF Rx modifies the mTBI-induced MRI-T1 and T2 intensity changes. Group differences in T1 (FLASH) and T2 (RARE) signal intensities estimated using a two-sample t-test on a voxel-by-voxel basis across the whole brain. A,D mTBI mice treated with vehicle vs sham mice, B,E. LIF treated mTBI mice vs sham mice and C,F. vehicle treated mTBI mice vs LIF treated mTBI mice. Activated voxels have a threshold of P<0.05, corrected for multiple comparisons using a family wise error control of 20-voxel contiguous cluster volume. Hot colors depict larger T1 or T2 intensity across the first group vs second group. Sham mice (n=7), mTBI mice treated with vehicle (n=7) and mTBI mice treated with LIF (n=8). Arrows indicates the mTBI epicenter.

Figure 3. IN-LIF Rx reduces mTBI-induced DTI-FA and MD changes, indicating axonal protection. Group differences between sham, vehicle-treated, and LIF-treated mTBI mice. **A.** fractional anisotropy (FA) changes **B.** mean diffusivity (MD) changes. The TBI epicenter is located at +0.5 mm AP from the bregma, as indicated by red arrows. Activated voxels represent a threshold of $p<0.05$ after a 2-tailed t-test corrected for multiple comparisons using a contiguous cluster volume of 30 voxels. Color map indicates FA or MD change magnitude.

Figure 4. IN-LIF Rx reduces mTBI-induced DTI-FA changes in a dose-dependent manner.
Quantitative DTI parameter changes across the corpus callosum (CC) and prefrontal cortex (PFC) regions of interest in sham (blue; n=7), mTBI+vehicle (orange; n=8), mTBI+40ng LIF (gray; n=8) and mTBI+100ng LIF (yellow; n=9). **A,B.** FA and **C,D.** MD. One-way ANOVA followed by a post-hoc Tukey's HSD test with p<0.05 was required for significance between groups. **A.** F=13.63, **B.** F=12.8, **C.** F=6.74 and **D.** F=5.47. **E.** FA intensity was plotted against the number of foot slips on the horizontal ladder for Vehicle Rx TBI mice vs. LIF Rx TBI mice. A Pearson's correlation analysis showed that while the slopes were not different (F = 1.37; p =0.27), the elevations were statistically different (F=17.4; p =0.001).

Figure 5. DTI-FA captures glial reactivity after mTBI, which IN-LIF Rx mitigates. Pseudocolor images of the FA maps from **A.** sham, **B.** mTBI+vehicle and **C.** mTBI+LIF groups. Colors represent FA from group averages, sham (n=7), mTBI+vehicle (n=8) and mTBI+LIF (n=8). Increased FA was observed across cortical and subcortical gray matter areas in the mTBI mice treated with vehicle (red arrows).

Fig 6. IN-LIF Rx decreases delayed axonal injury in corpus callosum after mTBI. Mice were injured at P60 and received IN-LIF for 2 weeks as described in Fig. 2. Images of APP staining in the CC in sham, IN-vehicle treated and IN-LIF Rx mice. Graph depicts particle analysis of APP staining. Sections were stained for MBP and the thickness of the corpus callosum was measured. Scale bar represents 50 µm for panels A-C and 200 µm for panels E-G. Data were analyzed by one-way ANOVA followed by Tukey's multiple comparison test. Statistical significance is indicated as follows: $p < 0.05$ (*), $p < 0.01$. n=6-8/group.

Fig 7. IN-LIF Rx decreases astroglial and microglial reactivity after mTBI. Mice were injured at P60 and received IN LIF for 2 weeks as described in Fig 2. After behavioral analyses, the mice were perfused, the brains sectioned and stained for GFAP, ADK, IBA-1 and CD-68. Scale. bar represents 100 µm. Data were analyzed by one-way ANOVA followed by Tukey's multiple comparison test for GFAP immunofluorescence (D) ADK immunofluorescence (E), % CD68/IBA-1+ (I0 and for CD68 immunofluorescence (J) . * p < 0.5, ** p<0.01, ** p<0.005, *** p < 0.001. n=6-8/group.

Supplementary Figure S1. rmTBI mice take significantly more time than the sham to exit an enclosure. A) Quantification of the time that each mouse took to exit the circle. n=4/group, *p<0.05 by Student's t-test. B) A schematic of the exit circle apparatus.

Supplementary Figure 2. **Activated astrocytes are increased at 72 h after r**mTBI. Immunofluorescence for GFAP (red) in the cerebral cortex (A,B) and the corpus callosum (C,D). Data were analyzed by one-way ANOVA followed by Tukey's multiple comparison test. Statistical significance is indicated as follows: $p < 0.05$ (*), $p < 0.01$ (), $p < 0.001$ (***) n=4/group. Scale bar represents 50 μm.

Supplementary Figure 3. **Adenosine kinase (ADK) is reduced in rmTBI mice vs. controls**.

Immunofluorescence for ADK (green) and GFAP (red) in the cerebral cortex (A,B) and the corpus callosum (C,D). Data were analyzed by one-way ANOVA followed by Tukey's multiple comparison test. Statistical significance is indicated as follows: $p < 0.05$ (*), $p < 0.01$ (), $p < 0.001$ (***) n=4/group. Scale bar represents 50 μm.

Supplementary Figure 4. **Activated microglia are increased at 72 h after rm**TBI. Immunofluorescence for CD68 (green) and IBA1 (red) in the cerebral cortex (A,B) and the corpus callosum (C,D). Data were analyzed by one-way ANOVA followed by Tukey's multiple comparison test. Statistical significance is indicated as follows: $p < 0.05$ (*), $p < 0.01$ (), $p < 0.001$ (***) n=4/group.

Supplementary Figure 5. Mice recover from sensorimotor deficits subsequent to rmTBI. A) Numbers of foot-slips recorded for mice on the horizontal ladder tested administered every 2 weeks after rmTBI. Data were analyzed by one-way ANOVA followed by Tukey's multiple comparison test. Statistical significance is indicated as follows: $p < 0.05$ (*), $p < 0.01$ (), $p < 0.001$ (***) n=3 mice/group. B). Photograph of the horizontal ladder.

Fig S6. Group differences in T1 (FLASH) and T2 (RARE) signal intensities estimated using a two-sample t-test on a voxel-by-voxel basis across the whole brain. A,D mTBI mice treated with vehicle vs sham mice, B,E. LIF treated mTBI mice vs sham mice and C,F. vehicle treated mTBI mice vs LIF treated mTBI mice. Activated voxels have a threshold of P<0.05, corrected for multiple comparisons using a family wise error control of 20-voxel contiguous cluster volume. Hot colors depict larger T1 or T2 intensity across the first group vs second group. Sham mice (n=7), mTBI mice treated with vehicle (n=7) and mTBI mice treated with LIF (n=8). Arrow indicates the mTBI epicenter.

Supplementary Figure 7. Six remaining regions of interest which did not show significant FA change between mTBI treated with vehicle (blue) versus sham (green) or mTBI treated with both doses of LIF versus sham (brown and red).

Supplementary Figure 8. Six remaining regions of interest which did not show significant MD change between mTBI treated with vehicle (blue) versus sham (green) or mTBI treated with both doses of LIF versus sham (brown and red).

A Sham

B mTBI+vehicle

C mTBI+LIF

Sham-epicenter

mTBI+veh epicenter

mTBI+LIF epicenter

D GFAP Vehicle Rx

GFAP LIF Rx

0.8

T1

A — veh vs sham

B — LIF vs sham

C — veh vs LIF

T2

D — veh vs sham

E — LIF vs sham

F — veh vs LIF

-10