Universal and quantitative detection of double-stranded RNAs as a signature of

pan-virus infections using a luciferase-based biosensor

Authors: Michihito Sasaki[1,2]*, Eri Fujii[1], Satoko Sasaki[1], Takuma Ariizumi[1], Kei

Konishi[3,4], Akihiko Sato[2,3,4], William W. Hall[5,6,7], Hirofumi Sawa[2,3,5,7] and Yasuko

Orba[1,2,3,5]*

Affiliations: [1]Division of Molecular Pathobiology, International Institute for Zoonosis

Control, Hokkaido University, Sapporo, Japan

[2]Institute for Vaccine Research and Development (IVReD), Hokkaido University, Sapporo,

Japan

[3]Division of Anti-Virus Drug Research, International Institute for Zoonosis Control,

Hokkaido University, Sapporo, Japan

[4]Laboratory for Drug Discovery & Disease Research, Shionogi & Co., Ltd., Osaka, Japan

[5]International Collaboration Unit, International Institute for Zoonosis Control, Hokkaido

University, Sapporo, Japan

[6]National Virus Reference Laboratory, School of Medicine, University College of Dublin,

Ireland

[7]Global Virus Network, Baltimore, Maryland, USA

*Correspondence: m-sasaki@czc.hokudai.ac.jp (M.S.), orbay@czc.hokudai.ac.jp (Y.O.)

Running title: dsRNA biosensor for quantifying viral replication

Abstract

Viral infections produce double-stranded RNA (dsRNA) during replication, which trigger

host innate immune responses. Immunoassays using anti-dsRNA antibodies have been

widely employed to detect viral dsRNA. In this study, we used a luciferase-based dsRNA

biosensor which consists of protein kinase R (PKR)-derived dsRNA binding domains fused

to split luciferase subunits. Here, we demonstrate the use of the dsRNA biosensor to

measure viral dsRNA in RNA specimens extracted from cells infected with Japanese

encephalitis virus (JEV). Moreover, the biosensor reacts to a broad-spectrum of dsRNAs

from infection with representatives of various viral families including positive- and

negative-sense single-stranded RNA (ssRNA) viruses, dsRNA viruses, and DNA viruses.

We validated the specific interaction between the dsRNA biosensor and viral RNA

including subgenomic flavivirus RNA (sfRNA) through RNA immunoprecipitation.

Additionally, we observed luminescence signals directly from lysates of JEV-infected cells

after cell lysis and phase separation with Triton X-114. Finally, we used the biosensor to

assess the activity of antiviral compounds. In summary, our results demonstrate that the

luciferase-based dsRNA biosensor offers a simple, homogeneous, and high-throughput

platform for quantifying viral replication, presenting a promising alternative to

antibody-based dsRNA detection methods.

Keywords: dsRNA, bioluminescence, RNA virus, DNA virus, flavivirus, PKR, RNA

binding protein, RNA structure

Introduction

The genomes of viruses include encapsided DNA, single-stranded RNA (ssRNA), or double-stranded RNA (dsRNA) and viruses are classified based on their genome types and replication strategies (Baltimore classification)(1). dsRNA comprises the nucleic acid of dsRNA viruses; however, this type of nucleic acid is also produced by ssRNA viruses as a replication intermediate and as a byproduct during viral replication in cells(2). As positive-sense ssRNA viruses replicate, they generate both viral genome–antigenome hybrid dsRNA (replicative form) and partial dsRNA that has multiple transcribing genomic RNAs (replicative intermediate) (3, 4). Copy-back defective viral genomic RNA is a partial viral genomic RNA characterized by reverse complementary 5′ and 3′ ends forming a dsRNA stem structure and is generated during the replication of various non-segmented negative-sense ssRNA viruses(5). In the case of DNA virus infection, converging bidirectional transcription of the viral genome produces complementary mRNA transcripts that can form dsRNA(6, 7). Collectively, the aberrant accumulation of dsRNA serves as a biomarker for viral infection.

Host cells counter viral infection through several dsRNA sensor proteins such as RIG-I-like receptors (RLRs), protein kinase R (PKR), and oligoadenylate synthases (OASes)(2). In particular, PKR is a dsRNA-dependent Ser/Thr protein kinase consisting of two tandem dsRNA binding domains (dsRBD) and a kinase domain. PKR recognizes dsRNA in a sequence-independent manner and requires a minimum of 30 to 33 bp of dsRNA for its activation(8, 9). When PKR interacts with dsRNA, the former is autophosphorylated, followed by the phosphorylation of eukaryotic initiation factor 2 alpha

(eIF2α) which results in translational shutoff and viral proliferation(10, 11). PKR can sense

various viral infections, and it mediates a universal antiviral effect in host cells. In response,

some viruses have evolved strategies to antagonize the innate immune responses mediated

by the dsRNA-PKR-eIF2α axis(10, 11).

Anti-dsRNA monoclonal antibodies specifically recognize dsRNA longer than 40 bp in

a sequence-independent manner(12). Clone J2, the most commonly employed anti-dsRNA

monoclonal antibody, has been used as a tool to detect a wide range of viral infections in

various immunoassays including immunofluorescence(3, 13, 14),

immunohistochemistry(15, 16), immunoprecipitation(17), and immunochromatographic

assays(18). J2 has also been used to determine the subcellular localization and

accumulation of dsRNA during the process of viral genome replication; furthermore, it has

been used to study the mechanisms of antiviral innate immune responses in various viral

infections(3, 14). Virological studies have predominantly relied on immunoassays using

anti-dsRNA antibodies to detect and quantify dsRNA; however, these assays have

limitations in sensitivity, quantitation and throughput. Therefore, it is desirable to develop

other more sophisticated dsRNA detection methods.

One alternative dsRNA detection tool is currently available as a luciferase assay

system using a dsRNA biosensor (Lumit dsRNA Detection Assay, Promega). This system

is based on a split luciferase complementation using small and large subunits of NanoLuc

luciferase, referred to as smBiT and LgBiT, respectively, by the manufacturer(19). The

biosensor consisting of PKR-derived dsRBD fused with smBiT or LgBiT (dsRBD-smBiT

or dsRBD-LgBiT, respectively) binds to dsRNA and reconstitutes the dsRBD-tagged

smBiT and LgBiT subunits as a functional NanoLuc luciferase(20, 21). The performance of

this luciferase assay system has been evaluated by quantifying artificially synthesized

dsRNA and detecting dsRNA contamination as a byproduct in *in vitro* transcription

products(20). The affinity of PKR for virus-induced dsRNA facilitates the use of this

luciferase-based dsRNA biosensor as a new research tool in the field of virology. In this

study, we have characterized the interaction between this biosensor and viral RNA and

examined its utility in various types of viral infections.

Results

Detection of dsRNA accumulation produced by flavivirus infection using the

luciferase-based dsRNA biosensor

We firstly used Japanese encephalitis virus (JEV), a single-stranded positive-sense

RNA virus in the family *Flaviviridae*, to examine whether the dsRNA biosensor can detect

viral replication in cells. Previous studies have used immunostaining with anti-dsRNA

monoclonal antibodies to show the accumulation of dsRNA in cells infected with

flaviviruses(22, 23). Consistent with previous reports, our immunofluorescence analysis

with anti-dsRNA antibody J2 demonstrated the accumulation of dsRNA in BHK-21 cells

infected with JEV (Fig. 1A). Total RNA was extracted from JEV-infected cells at different

time points, and quantitative RT-PCR (qRT-PCR) analysis revealed increases in

intracellular viral RNA levels over time (Fig. 1B). We also observed a time-dependent

increase of dsRNA in Quantitative ELISA with J2 antibody, while no clear dsRNA signal

was detected at 24 hpi (Fig. 1C). To examine the reactivity of dsRBD-LgBiT and

dsRBD-smBiT with dsRNA produced during viral replication, RNA samples from

JEV-infected cells were incubated with dsRBD-LgBiT and dsRBD-smBiT, and the

resulting luminescence was measured (Fig. 1D). The results showed higher luminescence

signals in the RNA samples at 24 (p=0.053), 48 ($p<0.0001$) and 72 hpi ($p<0.0001$) than that

in RNA of uninfected controls (Fig. 1E). Next, we performed the luciferase assay and

qRT-PCR using RNA samples mixed with RNAs from JEV-infected and uninfected cells in

different ratios. The luminescence and qRT-PCR signals were closely correlated (R^2 =

0.9829, Fig. 1F), although we observed background luminescence in the assay using RNA

from uninfected cells, which affected the limit of detection of the assay. The luminescence

signals from JEV-infected Vero and N2a cells indicated that the assay is independent of cell

type (Fig. 1G and 1H). Overall, these results indicate the applicability of the

luciferase-based dsRNA biosensor for detecting dsRNA accumulation during JEV

replication.

Broad-spectrum reactivity of the luciferase-based dsRNA biosensor to different viral

infections

To examine whether the dsRNA biosensor can detect the replication of various virus

types, we tested this in infections caused by the following viruses: positive-sense ssRNA

chikungunya virus (CHIKV, family *Togaviridae*) and Severe acute respiratory coronavirus

2 (SARS-CoV-2, family *Coronaviridae*); negative-sense ssRNA rabies virus (RABV,

family *Rhabdoviridae*) and La Crosse virus (LACV, family *Peribunyaviridae*); dsRNA

rotavirus A (RVA, family *Sedoreoviridae*) and Nelson Bay orthoreovirus (NBV, family

Spinareoviridae); and dsDNA mpox virus (MPXV, family *Poxviridae*) and herpes simplex

virus 1 (HSV-1, family *Orthoherpesviridae*). Immunofluorescence analysis with

anti-dsRNA and anti-viral protein antibodies demonstrated dsRNA accumulation in viral

antigen-positive Vero cells infected with CHIKV, LACV, NBV and MPXV,

Vero-TMPRSS2 cells infected with SARS-CoV-2, BHK-21 cells infected with RABV, and

MA104-T2T11D cells infected with RVA (Fig. 2A). dsRNA luciferase assay of total RNA

extracted from cells showed 9.5-fold to 22.3-fold increases in luminescence signal in RNA

samples from cells infected with CHIKV, SARS-CoV-2, RABV, LACV, RVA, NBV and

MPXV compared with uninfected cells (Fig. 2B). In contrast, we failed to detect

immunofluorescence signal of dsRNA and observed limited (2.0-fold) increase in

luminescence signal in Vero cells infected with HSV-1 (Fig. 2A and 2B), consistent with

previous reports that virion host shutoff protein of HSV-1 destabilizes dsRNA and limits its

accumulation in cells(6, 7). These results suggest that the dsRNA biosensor detects the

infection with a broad range of virus families through the accumulation of dsRNA in the

infected cells.

Specificity of the interaction of luciferase-based dsRNA biosensor with flavivirus

genomic and subgenomic RNAs

We investigated the target of the dsRNA biosensor in RNA samples from JEV infected

cells. First, we compared the reactivity of the dsRNA biosensor to RNA samples with or

without heat denaturation, which resolves double-stranded structure of dsRNA and

secondary and tertiary structures of ssRNA. Heat denaturation of sample RNA reduced the

luminescence signal of JEV-infected cells to a baseline level, similar to that of uninfected

cells (Fig. 3A), indicating that the viral infection-specific increase of the luminescence

signal depends on the RNA structure. We next examined the nature of the interaction

between the dsRNA biosensor and JEV RNA by RNA immunoprecipitation (RIP) with

RNA from JEV-infected cells using dsRBD-LgBiT, anti-LgBiT antibody, and protein G

beads. The LgBiT protein lacking the dsRBD domain was used as a control in the RIP

assay. Immunoblotting results showed the successful capture of dsRBD-LgBiT and LgBiT

on the Protein G beads (Fig. 3B). Viral genome RNA, viral mRNA, and subgenomic

flavivirus RNA (sfRNA) derived from the 3′-UTR are synthesized during active viral

replication in cells(24). Quantification of the amount of viral RNA in the

immunoprecipitate by qRT-PCR targeting the NS2A region and 3′-UTR of the JEV genome

revealed the specific interaction between the viral RNA and dsRBD-LgBiT (Fig. 3C and

3D). To determine the type of viral RNA in the immunoprecipitate, we performed northern

blot analysis with an RNA probe targeting the 3′-UTR of the JEV genome. The denaturing

gel electrophoresis followed by northern blot analysis identified the JEV sfRNA and

genomic RNA including replicative intermediate and replicative form in the RNA samples

extracted from JEV virions and JEV-infected cells (Fig. 3E). Furthermore, the analysis of

the immunoprecipitants revealed that dsRBD-LgBiT binds to not only the JEV RNA

genome but also sfRNA (Fig. 3E). To validate the specific interaction between

dsRBD-LgBiT and sfRNA, we performed additional RIP assay with *in vitro* transcribed

sfRNA (525 nt in length). Sequence scrambled sfRNA (525 nt in length), 525 and 3000 nt

ssRNAs of partial coding sequence (CDS) regions of JEV genome [CDS(525) and

CDS(3000)], and 717 nt ssRNA of AcGFP complete CDS were also synthesized by *in vitro*

transcription and used as ssRNA controls for the RIP assay. The synthesized sfRNA, but

not control ssRNAs, was enriched in the immunoprecipitate with anti-LgBiT and

dsRBD-LgBiT (Fig. 3F). Collectively, these results suggest that the dsRNA biosensor

specifically recognizes viral RNA including sfRNA synthesized in the cells infected with

JEV.

Quantitative detection of dsRNA in a homogeneous cell lysates using the

luciferase-based dsRNA biosensor

All the above experiments for dsRNA detection were performed with RNA extracted

from cells. We aimed to simplify the protocol by skipping the RNA extraction process and

detecting viral dsRNA directly in cell lysates using the dsRNA luciferase assay. First, we

assessed whether detergent contaminating the cell lysate could affect the dsRNA luciferase

assay. Using an extracted RNA sample containing 1% of conventional detergents, we found

that contamination by all tested detergents decreased the luminescence signal from the

dsRNA biosensor (Fig. 4A). Next, we suspended JEV-infected cells in lysis buffer with

different concentrations of digitonin and added the dsRNA biosensor to the mixture. The

luminescence signal from the dsRNA biosensor was accentuated by adding high

concentrations of digitonin in the cell lysis buffer, but the signal peaked at 0.25% of

digitonin and decreased at 1% and 4% of digitonin, probably due to the carry-over of

digitonin from the input cell lysate containing high concentrations of the detergent to the

dsRNA luciferase assay mixture (Fig. 4B). To reduce detergent carry-over after sufficient

cell lysis, we employed a phase separation approach using Triton X-114. Triton X-114 is a

nonionic detergent that forms a homogeneous solution on ice, but the solution separates

into aqueous and detergent-enriched phases above 20□(25). We suspended JEV-infected

cells in the lysis buffer with Triton X-114 on ice and then incubated the cell lysate at 37□,

followed by brief centrifugation to separate the phases (Fig. 4C). The dsRNA luciferase

assay detected a high luminescence signal in the aqueous phase of the cell lysate even with

high concentrations of Triton X-114. Notably, the phase separation procedure significantly

increased the luminescence signals from cell lysates containing 0.25%, 1% and 4% of

Triton X-114 (Fig. 4D). Overall, these results demonstrate that phase separation of cell

lysates with Triton X-114 enable us to obtain the RNA fraction with low detergent

concentrations and is a simple and reproducible way to directly detect dsRNA in cells.

We next examined whether our assay can measure the activity of antiviral compounds.

Ribavirin and NITD008 are nucleotide analogs that act as antiviral agents against

flaviviruses, including JEV(26-28). Both immunofluorescence staining and qRT-PCR

assays demonstrated the anti-JEV activities of ribavirin and NITD008 (Fig. 5A-5D).

Consistent with the results of conventional assays, the cell-directed dsRNA luciferase

assays showed that JEV infection decreased in a dose-dependent manner in response to

these antivirals (Fig. 5E and 5F). These results suggest that the cell-directed dsRNA

luciferase assay is a simple and fast method to estimate the magnitude of viral replication

and the activity of antiviral compounds.

Discussion

Replication of RNA and DNA viruses in cells produces viral genome-derived RNA

duplexes that stimulate the host antiviral immune responses through dsRNA sensors such

as RLRs, TLR-3, and PKR. In this study, we used the dsRNA biosensor dsRBD-LgBiT and

dsRBD-smBiT that employ the NanoLuc luciferase complementation process to detect viral

dsRNA in infected cells. We could demonstrate that luminescence signal from the

biosensor increases in a time-dependent manner and correlates with the viral RNA load in

JEV-infected cells. The biosensor could also be used to detect different viral infections

including those caused by ssRNA viruses, dsRNA viruses, and DNA viruses, without the

need for antibodies and oligonucleotide primers targeting specific viral proteins and

genomes. These results indicate that the luciferase-based dsRNA biosensor is capable of

estimating the magnitude of various types of viral infections including emerging and as yet

undiscovered viruses.

Anti-dsRNA monoclonal antibodies have been widely used to detect viral dsRNA in

various immunoassays as the protocol has broad applications and involves a simple

workflow. However, antibody-based immunoassays can sometimes be associated with

nonspecific reactions and variable signal intensity(29). The dsRNA luciferase assay is not

an immunoassay; rather, the dsRNA biosensor carries the dsRBD from PKR. Both the

dsRNA antibody and PKR recognize dsRNA in a sequence-independent manner. Using a

crosslinking and immunoprecipitation sequencing (CLIP-seq) analysis, Kim *et al*. revealed

that PKR as well as the anti-dsRNA antibody J2 can interact with various types of

dsRNA(30). Compared with the conventional immunoassays, the dsRNA luciferase assay

with the dsRNA biosensor offers a simple, rapid and high-throughput workflow to detect

dsRNA in RNA samples. However, this assay has been developed for dsRNA detection in

the extracted RNA samples and is susceptible to the presence of high amount of ssRNA and

detergent in the specimens. The results from our studies suggest that the dsRNA luciferase

assay is a versatile method for viral dsRNA detection with quantification and would be

applicable to sequence-independent detection of known and unknown viral infections in

cells.

PKR displays sequence-nonspecific interaction and autoactivates by not only dsRNA

but also ssRNA with an imperfect short stem flanked by single-stranded tails(31, 32). The

dsRNA biosensor recognizes dsRNA larger than 30 bp regardless of the nucleotide

sequence composition(20). Although we were able to demonstrate the interaction between

the biosensor and the JEV genome and sfRNA, the details on the target regions within the

viral RNAs remain elusive. There are multiple stem-loop structures in the 5′- and 3′-UTR

of the flavivirus genome and sfRNA(33, 34). Notably, a previous study showed that viral

sfRNA induces the activation of host PKR in cells(35). In addition, the long dsRNA

structure of flavivirus RNA genome consisting of a hybrid of the 5′-UTR and 3′-UTR was

formed during genomic RNA cyclization (a process that occurs during flavivirus

replication)(33, 34). We thus speculate that the stem-loops in JEV genomic RNA and

sfRNA are putative binding targets of dsRNA biosensor.

In this study, we showed the applicability of the dsRNA biosensor in a homogeneous

cell lysate. Although the interaction between the dsRNA biosensor and dsRNA was

attenuated by common detergents used for cell lysis, implementing a phase separation step

following cell lysis with Triton X-114 can effectively mitigate detergent interference while

maintaining the assay's sensitivity. Our cell-directed dsRNA luciferase assay was

applicable to measure the antiviral activity of compounds from cell lysates, and the results

correlated well with those of qRT-PCR. Implementing the dsRNA luciferase assay directly

on cells is a simple and fast way to evaluate the magnitude of viral infections. We believe

that the assay will also be useful for high-throughput screening of antiviral compounds and

host factors.

Although our study demonstrated the broad reactivity of the dsRNA biosensor in the

detection of various viral infections, it has the following limitations. First, the target

dsRNAs in each viral infection, except for JEV, remain elusive. The dsRBD in the

biosensor is derived from PKR; thus, identifying the targets of the biosensor will contribute

to our knowledge on the process of PKR activation in response to viral infection. Second,

the presence of endogenous dsRNA needs to be considered. The cellular levels of

host-derived dsRNAs such as inverted Alu repeats, bidirectional transcription of

mitochondrial DNA, and cellular mRNA with an extended 3′-UTR increase in response to

pathophysiological conditions(30, 36). In addition to virus-induced dsRNA, host-derived

dsRNAs are also sensed by endogenous dsRNA sensors. In neurons, mRNA with an

extended 3′-UTR acts as an endogenous dsRNA that triggers constitutive type I IFN

production, conferring an intrinsic antiviral state(37). However, we assumed that the

endogenous dsRNA had little effect in this study based on the result of heat denaturation

treatment of total RNA from uninfected cells, which showed limited impact on the

luciferase-based dsRNA assay. Finally, we observed background luminescence even in

total RNA samples from uninfected cells. This indicates that it may be necessary to further

optimize the luciferase assay protocol to ensure its wide dynamic range and robustness in

future applications.

<u>Materials and methods</u>

Cell lines

BHK-21 (RCB1423, RIKEN BRC) and C6/36 cells (CRL-1660, ATCC) were

maintained in Eagle's Minimum Essential Medium (MEM) containing 10% fetal bovine

serum (FBS). N2a cells (IFO50081, JCRB) were maintained in MEM containing 10% FBS

and nonessential amino acids (Gibco). Vero-TMPRSS2 cells stably expressing human

TMPRSS2 were established in our laboratory as described previously(38, 39).

Vero-TMPRSS2, Vero (CRL-1586, ATCC), and 293T cells (632180, Takara Bio) were

maintained in Dulbecco's Modified Eagle's Medium (DMEM) containing 10% FBS.

MA104-T2T11D cells expressing human TMPRSS2 and TMPRSS11D were established in

our laboratory as described previously(40) and maintained in MEM containing 10% FBS

and 10% tryptose phosphate broth (BD Difco).

Viruses

JEV (Mie/41/2002 strain)(41), CHIKV (SL10571 strain)(42), SARS-CoV-2 (WK-521

strain)(43), RABV (HEP strain)(44), NBV (MB/07 strain)(45) and MPXV (Zr-599

strain)(46) were provided by the National Institute of Infectious Diseases, Japan. HSV-1 (F

strain) was provided by the Institute of Medical Science, University of Tokyo, Japan. RVA

(SA11 strain) and LACV (VR-1834) were obtained from the ATCC. Working virus stocks

were prepared by passaging through C6/36 cells for JEV; BHK-21 cells for RABV; Vero

cells for CHIKV, NBV, MPXV, LACV and HSV-1; Vero-TMPRSS2 cells for

SARS-CoV-2; and MA104-T2T11D cells for RVA. The working virus stocks were titrated

by standard plaque assay (JEV, CHIKV, NBV, MPXV, LACV, HSV-1 and SARS-CoV-2)

or focus assay (RABV and RVA)(40, 47).

Immunofluorescence staining

BHK-21 cells on 24 well plate (Corning) were inoculated with 100 μl of inoculum

containing JEV or RABV at a multiplicity of infection (MOI) of 0.1 for 1h and then

cultured in MEM containing 2% FBS for 24 h. Vero cells on 24 well plate were

respectively inoculated with 100 μl of inoculum containing CHIKV, LACV, NBV, or

HSV-1 at an MOI of 0.1 and then cultured in DMEM containing 2% FBS for 24 h. Other

subset of Vero cells on 24 well plate were inoculated with 100 μl of inoculum containing

MPXV at an MOI of 0.01 for 1h and then cultured in DMEM containing 2% FBS for 48 h.

Vero-TMPRSS2 cells on 24 well plate were inoculated with 100 μl of inoculum containing

SARS-CoV-2 at an MOI of 0.1 for 1h and then cultured in DMEM containing 2% FBS for

24 h. MA104-T2T11D cells on 24 well plate were inoculated with 100 μl of inoculum

containing RVA at an MOI of 0.1 for 1h and then cultured in MEM containing 2% FBS

and 10% tryptose phosphate broth for 24 h. Cells were fixed at 24 h post-infection (hpi)

except for MPXV, which was fixed at 48 hpi with 3.7% buffered formaldehyde, followed

by permeabilization with 0.5% Triton X-100 in PBS for 5 min and double–staining with

anti-dsRNA mouse IgG2a monoclonal antibody (clone J2; 10010200, SCICONS) and

virus-specific primary antibody in 25% Block Ace (KAC) in PBS for 1 h. We used

anti-CHIKV E1 mouse IgG1 monoclonal (MAB12424, Abnova), anti-SARS-CoV-2

nucleocapsid rabbit monoclonal (GTX635679, GeneTex), anti-RABV nucleoprotein mouse

IgG2b monoclonal (3R7-4G4, HyTest), anti-LACV mouse IgG2b monoclonal

(MA1-10801, Thermo Fisher Scientific), anti-RVA goat polyclonal (AB1129, Merck),

anti-Vaccinia virus rabbit polyclonal (cross reactive with MPXV; PA17258, Thermo Fisher

Scientific), anti-HSV-1 rabbit polyclonal (B0114, DAKO), anti-JEV rabbit polyclonal

(serum from JEV-immunized rabbit in our laboratory), and anti-NBV rabbit polyclonal

(serum from NBV-immunized guinea pig in our laboratory) antibodies(48) for the viral

antigen staining. The cells were subsequently stained for 30 min with the following

appropriate secondary antibodies: anti-mouse IgG2a Alexa Fluor 488-conjugated goat

polyclonal (A21131, Thermo Fisher Scientific), anti-mouse IgG1 Alexa Fluor

594-conjugated goat polyclonal (A21125, Thermo Fisher Scientific), anti-mouse IgG2b

Alexa Fluor 594-conjugated goat polyclonal (A21145, Thermo Fisher Scientific),

anti-rabbit IgG Alexa Fluor 594-conjugated goat polyclonal (A11037, Thermo Fisher

Scientific), anti-guinea pig IgG Alexa Fluor 647-conjugated goat polyclonal (A21450,

Thermo Fisher Scientific), anti-mouse IgG Alexa Fluor 488-conjugated donkey polyclonal

(A21202, Thermo Fisher Scientific), and anti-goat IgG Alexa Fluor 568-conjugated donkey

polyclonal (A11057, Thermo Fisher Scientific). Cell nuclei were stained with Hoechst

33342 (Thermo Fisher Scientific). Fluorescent images were captured using an IX73

fluorescence microscope (Olympus).

RNA measurement by qRT-PCR

Virus growth kinetics were determined with BHK-21 cells infected with JEV at an MOI of 0.01. Total RNA was extracted from the cells at different time points with the Direct-zol RNA MiniPrep Kit (Zymo Research) and analyzed by qRT-PCR with the EXPRESS One-Step Superscript qRT-PCR Kit (Thermo Fisher Scientific) and a primer/probe set targeting JEV *NS2A*(49). The hamster *Actb* gene was used as an internal control for normalization(50). For the RIP assay, *in vitro* transcribed (IVT) ssRNAs and the 3′-UTR of the JEV genome in the immunoprecipitates were quantified, by qRT-PCR on the extracted RNA using a One-Step TB Green PrimeScript PLUS RT-PCR Kit (Takara Bio). JEV RNA levels were calculated by the standard curve, comparative ΔCt or $\Delta\Delta$Ct methods. The probe and primers used in this study are summarized in Table S1 (Supplementary materials).

Detection of dsRNA in RNA extracts

BHK-21 cells on 24 well plate were infected with JEV or RABV at an MOI of 0.01. Vero cells on 24 well plate were individually infected with CHIKV, LACV, NBV, MPXV, and HSV-1 at an MOI of 0.01. Vero-TMPRSS2 cells and MA104-T2T11D cells on 24 well plate were infected with SARS-CoV-2 and RVA, respectively, at an MOI of 0.01. At 1 and 48 hpi, the cells were lysed with 300 μl of RNAiso Plus (Takara Bio). Total RNA was extracted from the cell lysates with Direct-zol RNA MiniPrep Kit , with mock-infected cells included as controls. The extracted RNA samples were subjected to dsRNA luciferase assay using the Lumit dsRNA biosensor (dsRBD-LgBiT and dsRBD-smBiT) with dsRNA

assay buffer and luminescence detection substrate B provided in the Lumit dsRNA

Detection Assay kit (W2041, Promega) according to the manufacturer's instructions with

slight modifications. Briefly, a total of 200 to 500 ng RNA in 5 μl water was mixed and

incubated with dsRBD-LgBiT and dsRBD-smBiT in 95 μl of dsRNA assay buffer for 1 h at

room temperature. The mixture of RNA and biosensor was then added to 25 μl of

luminescence detection substrate B diluted in dsRNA assay buffer on 96-well half area

black microplates (3694, Corning). After 15-min of incubation, the luminescence was

measured with a GloMax Discover Microplate Reader (Promega). In parallel with the

dsRNA luciferase assay, dsRNA amount in RNA samples was also measured by ELISA

with dsRNA ELISA kit, J2-based (Exalpha Biologicals) following the manufacture's

instruction. 500 ng of RNA samples were loaded in each well and the values of optical

density at 450 nm were measured with iMark microplate reader (Bio-Rad).

Heat denaturation treatment of dsRNA was conducted by heating RNA samples for 5

min at 65□, then chilling on ice for 3 min, heating for 2 min at 95□, and finally chilling on

ice for 3 min. The heat denatured RNA samples was subjected to the dsRNA luciferase

assay as described above.

dsRNA detection in the cell lysates

BHK-21 cells on a 24-well plate were infected with JEV at an MOI of 0.01. At 48 hpi,

cells were lysed in 100 μl/well of cell lysis buffer (25 mM Tris-HCl [pH 7.5], 150 mM

NaCl, 5 mM EDTA) supplemented with different concentrations of digitonin (0.001 to 4%),

Triton X-114 (0.001 to 4%), NP-40 (1%), Brij 98 (1%) (all from Sigma), and Triton X-100

(1%) (Nacalai Tesque) for 5 min on ice. The phases in the Triton X-114-containing cell

lysates were separated by first incubating the mixture for 1 min at 37□ and then

centrifuging at $9000 \times g$ for 30 s. The amount of dsRNA was measured in the upper

detergent-free phase. We analyzed cell lysates that had both undergone and not undergone

phase separation by mixing 5 μl of the lysates with dsRBD-LgBiT and dsRBD-smBiT in 95

μl of dsRNA assay buffer and analyzing the solution using the Lumit dsRNA Detection

Assay kit as described above.

RIP assay with cellular RNA samples

Fifty microliters of protein G-coupled magnetic beads (Dynabeads protein G; Veritas)

were conjugated with 10 μg of anti-LgBiT monoclonal antibody (N7100, Promega) for 15

min and then incubated with 22.8 pmol of control LgBiT (Promega) or dsRBD-LgBiT in

PBS with 0.002% Tween 20 (PBST) for 30 min. Then 10 μg of total RNA extracted from

BHK-21 cells at 48 hpi with JEV was incubated with the magnetic beads in 500 μl of

dsRNA assay buffer supplemented with 60 units of a recombinant RNase inhibitor (Takara

Bio) overnight at 4□ with rotation. The mixture was then washed 3 times with PBST, and

10% of the beads were suspended in SDS sample buffer [4% SDS, 100 mM Tris-HCl (pH

6.8), 20% glycerol, 10% 2-mercaptoethanol] for immunoblotting. The remaining 90% of

the beads were suspended in RNAiso plus and the suspension was subjected to RNA

extraction with Direct-zol-96 MagBead RNA Kit (Zymo Research). The RNA extract was

analyzed by qRT-PCR for JEV RNA levels in the immunoprecipitate as described above

and by Northern blotting for JEV RNA size in the immunoprecipitate.

Our immunoblotting protocol began by resolving input and immunoprecipitated

proteins by SDS-PAGE, followed by blotting of the proteins onto an Immobilon-PSQ

PVDF membrane (Merck). The membrane was incubated with anti-LgBiT monoclonal

antibody (N7100) overnight at 4□ and then incubated with HRP-conjugated anti-mouse

IgG polyclonal antibody (AMI3404, BioSource) for 30 min at room temperature. Immune

complexes were detected using Immobilon Western Chemiluminescent Substrate (Merck)

on an Amersham ImageQuant 800 (Cytiva).

Northern blotting was performed using the DIG Northern Starter Kit (Roche)

according to the manufacturer's instructions with slight modifications. Briefly, we resolved

the following on 1.2% SeaKem GTG agarose-2.2 M formaldehyde gels : RNA samples

eluted from the magnetic beads, RNA extracted from BHK-21 infected with JEV, control

RNAs from uninfected BHK, and JEV virions in culture supernatants. The RNAs were

transferred onto a Nytran SuPerCharge nylon membrane using a Turboblotter (Cytiva) and

UV crosslinked at 120 mJ/cm^2 in an HL-2000 HybriLinker (UVP). For the digoxigenin

(DIG)–labeled RNA probes, template cDNA of the 3′-UTR of the JEV genome fused with

T7 promoter sequence was amplified by RT-PCR and applied to *in vitro* transcription with

T7 RNA polymerase. The UV-crosslinked membrane was hybridized with DIG–labeled

RNA probe in ULTRAhyb Ultrasensitive Hybridization Buffer (Thermo Scientific).

Hybridization complexes were detected with alkaline phosphatase-conjugated anti-DIG

antibody and CDP-Star (all from Roche) on Amersham ImageQuant 800.

RIP assay with IVT ssRNA samples

For *in vitro* transcription, DNA templates at nucleotide positions 10441 to 10965 for of

sfRNA, positions 9871 to 10395 for CDS (525) and positions 7396 to 10395 for CDS

(3000) in JEV genome (GenBank accession no. AB241119) were prepared and fused with

T7 promoter sequence by RT-PCR with PrimeScript II High Fidelity One Step RT-PCR Kit

(Takara Bio). DNA template of sequence-scrambled sfRNA with T7 promoter was

synthesized through gBlocks gene fragment synthesis (IDT). DNA template of AcGFP

CDS with T7 promoter was prepared by PCR with PrimeSTAR Max DNA polymerase and

pAcGFP1-C1 vector (Takara Bio). Sequence information of scrambled sfRNA and primers

for PCR/RT-PCR was summarized in Table S2 and S3 (Supplementary materials). IVT

ssRNAs were prepared by *in vitro* transcription with Takara IVTpro mRNA Synthesis Kit

(low dsRNA) (Takara Bio) and then purified with NucleoSpin RNA Clean-up XS kit

(Macherey Nagel). For RIP assay, 3.5 μl of protein G-coupled magnetic beads were

conjugated with 0.5 μg of anti-LgBiT monoclonal antibody (N7100) for 15 min and then

incubated with 1.4 pmol of control LgBiT or dsRBD-LgBiT in PBST for 30 min. The

magnetic beads were incubated with 10 ng of IVT ssRNA samples in 300 μl of dsRNA

assay buffer supplemented with 10 units of a recombinant RNase inhibitor and 3000 ng of

total RNA from uninfected BHK-21 cells for 2h at 4□ with rotation. The mixtures were

then washed 3 times with PBST and suspended in RNAiso plus. The RNA was extracted

with Direct-zol-96 MagBead RNA Kit and analyzed by qRT-PCR for each ssRNA levels in

the immunoprecipitates as described above.

Evaluation of antiviral efficacy

465   BHK-21 cells on 24-well plates were infected with JEV at an MOI of 0.01 in the

466 presence of different concentrations of ribavirin (Wako) or NITD008 (MedChemExpress).

467 At 48 hpi, a subset of cells were fixed with 3.7% buffered formaldehyde and stained with

468 anti-JEV rabbit polyclonal antibody. Another subset of cells on different 24-well plates

469 were lysed in 100 μl/well of cell lysis buffer supplemented with 1% Triton X-114 for 5 min

470 on ice. After phase separation, 5 μl of the upper detergent-free phase was analyzed using

471 the Lumit dsRNA Detection Assay kit to quantify dsRNA as described above. The

472 remaining lysate was subjected to RNA extraction and qRT-PCR as described above.

474 **Statistical analysis and reproducibility**

475  Statistical significance was determined by one-way analysis of variance (ANOVA)

476 with Dunnett's test (Figs. 1B, 1C, 1E, 1G, 1H, 2B, 4A and 5C-5F), one-way ANOVA with

477 Tukey's test (Fig. 3A and 3F), two-tailed Welch's t-test (Fig. 3C and 3D) and two-tailed

478 Student's t-test (Fig. 4D). All statistical tests were carried out using Prism version 9.5.1

479 (GraphPad software). All experiments were repeated independently 2 or 3 times with

480 similar results. Most results are presented as mean± SD of triplicate samples as specified in

481 the figure legends.

483 **Data availability**

484 All data associated with this work are included in this article.

486 **Acknowledgments**

We thank Dr. Saijo and Dr. Ebihara of the National Institute of Infectious Diseases, Japan

for providing JEV, CHIKV, SARS-CoV-2, RABV, NBV and MPXV, Dr. Kawaguchi of

Tokyo University for providing HSV-1, and Dr. Ito of Gifu University for support in the

establishment of MA104-T2T11D cells. We also thank Dr. Inagawa of Promega Japan for

technical support in the Lumit dsRNA detection assays.

Funding information

This work was supported in part by the Japan Agency for Medical Research and

Development (AMED) under Grant numbers JP24fk0108637 [Y.O.], JP223fa627005 [A.S.,

H.S.], JP24wm0125008 [H.S., Y.O.], JP24wm0325073 [Y.O.]; Japan Science and

Technology Agency (JST) Moonshot R&D under Grant number JPMJMS2025 [Y.O.];

Japan Society for the Promotion of Science (JSPS) KAKENHI under Grant number

JP23H02376 [M.S.].

Conflicts of interests

The authors K.K. and A.S. are employees and shareholders of Shionogi & Co., Ltd. The

remaining authors declare no competing interests.

References

1.   Koonin, E. V., Krupovic, M., and Agol, V. I. (2021) The Baltimore Classification of
Viruses 50 Years Later: How Does It Stand in the Light of Virus Evolution?
Microbiol Mol Biol Rev **85**, e0005321

2.   Chen, Y. G., and Hur, S. (2022) Cellular origins of dsRNA, their recognition and
consequences *Nat Rev Mol Cell Biol* **23**, 286-301

3.   Triantafilou, K., Vakakis, E., Kar, S., Richer, E., Evans, G. L., and Triantafilou, M.
(2012) Visualisation of direct interaction of MDA5 and the dsRNA replicative
intermediate form of positive strand RNA viruses *J Cell Sci* **125**, 4761-4769

4.   Son, K. N., Liang, Z., and Lipton, H. L. (2015) Double-Stranded RNA Is Detected
by Immunofluorescence Analysis in RNA and DNA Virus Infections, Including
Those by Negative-Stranded RNA Viruses *J Virol* **89**, 9383-9392

5.   Brennan, J. W., and Sun, Y. (2024) Defective viral genomes: advances in
understanding their generation, function, and impact on infection outcomes *mBio* **15**,
e0069224

6.   Burgess, H. M., and Mohr, I. (2018) Defining the Role of Stress Granules in Innate
Immune Suppression by the Herpes Simplex Virus 1 Endoribonuclease VHS *J Virol*
92,

7.   Dauber, B., Saffran, H. A., and Smiley, J. R. (2019) The herpes simplex virus host
shutoff (vhs) RNase limits accumulation of double stranded RNA in infected cells:
Evidence for accelerated decay of duplex RNA *PLoS Pathog* **15**, e1008111

8.   Manche, L., Green, S. R., Schmedt, C., and Mathews, M. B. (1992) Interactions
between double-stranded RNA regulators and the protein kinase DAI *Mol Cell Biol*

12, 5238-5248

9.   Husain, B., Mukerji, I., and Cole, J. L. (2012) Analysis of high-affinity binding of

protein kinase R to double-stranded RNA *Biochemistry* **51**, 8764-8770

10.   Cesaro, T., and Michiels, T. (2021) Inhibition of PKR by Viruses *Front Microbiol*

12, 757238

11.   Zhang, R., and Karijolich, J. (2024) RNA recognition by PKR during DNA virus

infection *J Med Virol* **96**, e29424

12.   Schönborn, J., Oberstrass, J., Breyel, E., Tittgen, J., Schumacher, J., and Lukacs, N.

(1991) Monoclonal antibodies to double-stranded RNA as probes of RNA structure

in crude nucleic acid extracts *Nucleic Acids Res* **19**, 2993-3000

13.   Weber, F., Wagner, V., Rasmussen, S. B., Hartmann, R., and Paludan, S. R. (2006)

Double-stranded RNA is produced by positive-strand RNA viruses and DNA

viruses but not in detectable amounts by negative-strand RNA viruses *J Virol* **80**,

5059-5064

14.   Deng, X., Hackbart, M., Mettelman, R. C., O'Brien, A., Mielech, A. M., Yi, G. *et al.*

(2017) Coronavirus nonstructural protein 15 mediates evasion of dsRNA sensors

and limits apoptosis in macrophages *Proc Natl Acad Sci U S A* **114**, E4251-E4260

15.   Richardson, S. J., Willcox, A., Hilton, D. A., Tauriainen, S., Hyoty, H., Bone, A. J.

et al. (2010) Use of antisera directed against dsRNA to detect viral infections in

formalin-fixed paraffin-embedded tissue *J Clin Virol* **49**, 180-185

16.   Piantadosi, A., Shariatzadeh, N., Bombin, A., Arkun, K., Alexandrescu, S.,

Kleinschmidt-DeMasters, B. K. *et al.* (2023) Double-stranded RNA

immunohistochemistry as a screening tool for viral encephalitis *Am J Clin Pathol*

160, 210-219

17. Decker, C. J., Steiner, H. R., Hoon-Hanks, L. L., Morrison, J. H., Haist, K. C.,
Stabell, A. C. *et al.* (2019) dsRNA-Seq: Identification of Viral Infection by
Purifying and Sequencing dsRNA *Viruses* **11**,

18. Ku, J., Kim, S., Park, J., Kim, T. S., Kharbash, R., Shin, E. C. *et al.* (2020) Reactive
Polymer Targeting dsRNA as Universal Virus Detection Platform with Enhanced
Sensitivity *Biomacromolecules* **21**, 2440-2454

19. Dixon, A. S., Schwinn, M. K., Hall, M. P., Zimmerman, K., Otto, P., Lubben, T. H.
et al. (2016) NanoLuc Complementation Reporter Optimized for Accurate
Measurement of Protein Interactions in Cells *ACS Chem Biol* **11**, 400-408

20. Moravec, R., Wang, J., Hartnett, J., Cong, M., and Grailer, J. (2024) Development
of novel luminescent assays for sensitive and specific quantitation of
double-stranded RNA in mRNA vaccines *Cancer Research* **84**, 3254-3254

21. Silas, D. S., Juneja, B., Kaur, K., Narayanareddy Gari, M., You, Y., Moon, Y. *et al.*
(2024) Development of Biolayer Interferometry (BLI)-Based Double-Stranded
RNA Detection Method with Application in mRNA-Based Therapeutics and
Vaccines *Pharmaceutics* **16**,

22. Miorin, L., Romero-Brey, I., Maiuri, P., Hoppe, S., Krijnse-Locker, J.,
Bartenschlager, R. *et al.* (2013) Three-dimensional architecture of tick-borne
encephalitis virus replication sites and trafficking of the replicated RNA *J Virol* **87**,
6469-6481

23. Tabata, K., Itakura, Y., Ariizumi, T., Igarashi, M., Kobayashi, H., Intaruck, K. *et al.*
(2023) Development of flavivirus subviral particles with low cross-reactivity by
mutations of a distinct antigenic domain *Appl Microbiol Biotechnol* **107**, 7515-7529

24. Slonchak, A., and Khromykh, A. A. (2018) Subgenomic flaviviral RNAs: What do

we know after the first decade of research *Antiviral Res* **159**, 13-25

25.   Bordier, C. (1981) Phase separation of integral membrane proteins in Triton X-114

580   solution *J Biol Chem* **256**, 1604-1607

581   26.   Tsao, C. H., Su, H. L., Lin, Y. L., Yu, H. P., Kuo, S. M., Shen, C. I. *et al.* (2008)

582   Japanese encephalitis virus infection activates caspase-8 and -9 in a

583   FADD-independent and mitochondrion-dependent manner *J Gen Virol* **89**,

584   1930-1941

585   27.   Zhang, Z. R., Zhang, H. Q., Li, X. D., Deng, C. L., Wang, Z., Li, J. Q. *et al.* (2020)

586   Generation and characterization of Japanese encephalitis virus expressing GFP

587   reporter gene for high throughput drug screening *Antiviral Res* **182**, 104884

588   28.   Li, C., Chen, X., Hu, J., Jiang, D., Cai, D., and Li, Y. (2022) A Recombinant

589   Genotype I Japanese Encephalitis Virus Expressing a *Int J Mol Sci* **23**,

590   29.   Thomsen, C., Røge, R., Fred, Å., and Wanders, A. (2023) Immunohistochemical

591   detection of double-stranded RNA in formalin-fixed paraffin-embedded tissue

592   *APMIS* **131**, 197-205

593   30.   Kim, Y., Park, J., Kim, S., Kim, M., Kang, M. G., Kwak, C. *et al.* (2018) PKR

594   Senses Nuclear and Mitochondrial Signals by Interacting with Endogenous

595   Double-Stranded RNAs *Mol Cell* **71**, 1051-1063.e1056

596   31.   Zheng, X., and Bevilacqua, P. C. (2004) Activation of the protein kinase PKR by

597   short double-stranded RNAs with single-stranded tails *RNA* **10**, 1934-1945

598   32.   Nallagatla, S. R., Hwang, J., Toroney, R., Zheng, X., Cameron, C. E., and

599   Bevilacqua, P. C. (2007) 5'-triphosphate-dependent activation of PKR by RNAs

600   with short stem-loops *Science* **318**, 1455-1458

33.   Villordo, S. M., and Gamarnik, A. V. (2009) Genome cyclization as strategy for

flavivirus RNA replication *Virus Res* **139**, 230-239

34. Saeedi, B. J., and Geiss, B. J. (2013) Regulation of flavivirus RNA synthesis and

capping *Wiley Interdiscip Rev RNA* **4**, 723-735

35. Pallarés, H. M., González López Ledesma, M. M., Oviedo-Rouco, S., Castellano, L.

A., Costa Navarro, G. S., Fernández-Alvarez, A. J. *et al.* (2024) Zika virus

non-coding RNAs antagonize antiviral responses by PKR-mediated translational

arrest *Nucleic Acids Res* **52**, 11128-11147

36. Dhir, A., Dhir, S., Borowski, L. S., Jimenez, L., Teitell, M., Rötig, A. *et al.* (2018)

Mitochondrial double-stranded RNA triggers antiviral signalling in humans *Nature*

560, 238-242

37. Dorrity, T. J., Shin, H., Wiegand, K. A., Aruda, J., Closser, M., Jung, E. *et al.* (2023)

Long 3'UTRs predispose neurons to inflammation by promoting

immunostimulatory double-stranded RNA formation *Sci Immunol* **8**, eadg2979

38. Sasaki, M., Uemura, K., Sato, A., Toba, S., Sanaki, T., Maenaka, K. *et al.* (2021)

SARS-CoV-2 variants with mutations at the S1/S2 cleavage site are generated in

vitro during propagation in TMPRSS2-deficient cells *PLoS Pathog* **17**, e1009233

39. Uemura, K., Sasaki, M., Sanaki, T., Toba, S., Takahashi, Y., Orba, Y. *et al.* (2021)

MRC5 cells engineered to express ACE2 serve as a model system for the discovery

of antivirals targeting SARS-CoV-2 *Sci Rep* **11**, 5376

40. Sasaki, M., Itakura, Y., Kishimoto, M., Tabata, K., Uemura, K., Ito, N. *et al.* (2021)

Host serine proteases TMPRSS2 and TMPRSS11D mediate proteolytic activation

and trypsin-independent infection in group A rotaviruses *J Virol* **95**,

41. Tajima, S., Taniguchi, S., Nakayama, E., Maeki, T., Inagaki, T., Lim, C. K. *et al.*

(2020) Amino Acid at Position 166 of NS2A in Japanese Encephalitis Virus (JEV) is

Associated with In Vitro Growth Characteristics of JEV *Viruses* **12**,

42.   Lim, C. K., Nishibori, T., Watanabe, K., Ito, M., Kotaki, A., Tanaka, K. *et al.* (2009)

Chikungunya virus isolated from a returnee to Japan from Sri Lanka: isolation of

two sub-strains with different characteristics *Am J Trop Med Hyg* **81**, 865-868

43.   Matsuyama, S., Nao, N., Shirato, K., Kawase, M., Saito, S., Takayama, I. *et al.*

(2020) Enhanced isolation of SARS-CoV-2 by TMPRSS2-expressing cells *Proc*

Natl Acad Sci U S A **117**, 7001-7003

44.   Horiya, M., Posadas-Herrera, G., Takayama-Ito, M., Yamaguchi, Y., Iizuka-Shiota,

I., Kato, H. *et al.* (2022) Genetic Characterization of Human Rabies Vaccine Strain

in Japan and Rabies Viruses Related to Vaccine Development from 1940s to 1980s

Viruses **14**,

45.   Yamanaka, A., Iwakiri, A., Yoshikawa, T., Sakai, K., Singh, H., Himeji, D. *et al.*

(2014) Imported case of acute respiratory tract infection associated with a member

of species nelson bay orthoreovirus *PLoS One* **9**, e92777

46.   Saijo, M., Ami, Y., Suzaki, Y., Nagata, N., Iwata, N., Hasegawa, H. *et al.* (2006)

LC16m8, a highly attenuated vaccinia virus vaccine lacking expression of the

membrane protein B5R, protects monkeys from monkeypox *J Virol* **80**, 5179-5188

47.   Anindita, P. D., Sasaki, M., Nobori, H., Sato, A., Carr, M., Ito, N. *et al.* (2016)

Generation of recombinant rabies viruses encoding NanoLuc luciferase for antiviral

activity assays *Virus Res* **215**, 121-128

48.   Harima, H., Sasaki, M., Orba, Y., Okuya, K., Qiu, Y., Wastika, C. E. *et al.* (2021)

Attenuated infection by a Pteropine orthoreovirus isolated from an Egyptian fruit

bat in Zambia *PLoS Negl Trop Dis* **15**, e0009768

49.   Bharucha, T., Sengvilaipaseuth, O., Vongsouvath, M., Davong, V., Panyanouvong,

P., Piorkowski, G. *et al.* (2018) Development of an improved RT-qPCR Assay for

detection of Japanese encephalitis virus (JEV) RNA including a systematic review

and comprehensive comparison with published methods *PLoS One* **13**, e0194412

50. Zivcec, M., Safronetz, D., Haddock, E., Feldmann, H., and Ebihara, H. (2011)

Validation of assays to monitor immune responses in the Syrian golden hamster

(Mesocricetus auratus) *J Immunol Methods* **368**, 24-35

 Figure Legends

Figure 1. Quantitative detection of dsRNA in JEV infection by luciferase assays with

the dsRNA biosensor

(A) Immunofluorescence staining of BHK-21 cells infected with JEV or mock-infected

cells at 24 hpi. Cells were stained with anti-JEV (red), anti-dsRNA (green), and Hoechst

33342 (blue). **(B)** Viral RNA loads in cells at the indicated time points of JEV infection

were analyzed by qRT-PCR. qRT-PCR data were normalized to endogenous *Actb*. **(C)**

dsRNA loads in cells at the indicated time points of JEV infection were analyzed by J2

antibody-based ELISA. **(D)** Schematic representation of dsRNA detection in RNA sample

by split nanoluciferase complementation assay using the dsRNA biosensor (dsRBD-smBiT

and dsRBD-LgBiT). **(E)** Total RNA was extracted from BHK-21cells infected with JEV

and analyzed by luciferase assay with the dsRNA biosensor. **(F)** Correlation of the

qRT-PCR targeting JEV RNA and luciferase assays with the dsRNA biosensor. RNA

samples from JEV-infected and uninfected BHK-21 cells were mixed at different ratios.

The RNA mixtures were analyzed by qRT-PCR and luciferase assay to measure the levels

of JEV RNA and dsRNA, respectively. The horizontal dashed line indicates the

luminescence value of uninfected cells in the assay (the detection limit of dsRNA related to

JEV infection). R^2 indicates the linear correlation coefficients between qRT-PCR and

dsRNA detection assays. **(G, H)** Total RNA was extracted from Vero (G) and N2a (H) cells

infected with JEV and analyzed by luciferase assays with the dsRNA biosensor. The values

shown are mean ± standard deviation (SD) of triplicate (B, C, E, G, H) or duplicate (F)

samples. ****$p < 0.0001$ by one-way ANOVA with Dunnett's test.

Figure 2. Application of the dsRNA biosensor to different viral infections

Positive-sense ssRNA viruses [Chikungunya virus (CHIKV), SARS-CoV-2],

negative-sense ssRNA viruses [Rabies virus (RABV), La Crosse virus (LACV)], dsRNA

viruses [Rotavirus A (RVA), Nelson Bay orthoreovirus (NBV)], DNA viruses [Mpox virus

(MPXV), and Herpes simplex virus 1 (HSV-1)] were inoculated onto monolayers of Vero

(CHIKV, LACV, NBV, MPXV and HSV-1), Vero-TMPRSS2 (SARS-CoV-2), BHK-21

(RABV), and MA104-T2T11D cells (RVA). **(A)** Immunofluorescence staining of the

infected cells with Hoechst 33342 nuclear dye (blue), antibodies targeting dsRNA (green)

and virus proteins (red). Scale bars, 50 μm. **(B)** Luciferase assay with the dsRNA biosensor

was used to measure dsRNA in RNA extracts from mock-infected and virus-infected cells.

The values shown are mean ± SD of triplicate samples. **$p < 0.01$, ****$p < 0.0001$ by

one-way ANOVA with Dunnett's test.

Figure 3. Target specificity of the dsRNA biosensor

(A) Effect of heat denaturation on dsRNA sensing by the dsRNA biosensor. Heat-denatured

and non-denatured RNA samples were analyzed by luciferase assays with the dsRNA

biosensor. **(B-D)** RNA immunoprecipitation (RIP) assays with total RNA extracted from

BHK-21 cells at 48 hpi with JEV using dsRNA biosensor dsRBD-LgBiT and anti-LgBiT

antibody. LgBiT protein without dsRBD was used as a control in the RIP assay.

Immunoprecipitates were analyzed by immunoblotting with anti-LgBiT (B) and qRT-PCR

targeting the JEV NS2A gene (C) and the 3′-UTR of the JEV genome (D). **(E)** Northern

blotting of the immunoprecipitates using digoxigenin DIG–labeled RNA probe targeting

the 3′-UTR of the JEV genomic RNA including replicative form and replicative

intermediate. RNA extracted from the culture supernatant containing JEV virions,

JEV-infected and uninfected BHK-21 cells were included in the blotting to estimate the

size of the JEV genome and subgenomic flavivirus RNA (sfRNA). **(F)** RIP assays with v

ssRNAs of JEV sfRNA, sequence-scrambled sfRNA (scrambled), partial coding sequences

of JEV genome [CDS(525) and CDS(3000)], and complete coding sequence of AcGFP.

Relative enrichment of each ssRNA was calculated by dividing the ssRNA amounts of

immunoprecipitates using dsRBD-LgBiT with those of immunoprecipitates using LgBiT

protein control. The values shown are mean ± SD of triplicate samples. ****$p < 0.0001$ by

one-way ANOVA with Tukey's test (A, F) or two-tailed Welch's t-test (C, D).

Figure 4. Detection of dsRNA in cell lysates using the dsRNA biosensor

(A) RNA samples extracted from JEV BHK-21 cells at 48 hpi with JEV were mixed with

the indicated detergents at a final concentration of 1% and then analyzed by the luciferase

assay for dsRNA detection with the dsRNA biosensor. **(B)** Cells at 48 hpi following JEV

infection were suspended in lysis buffer containing different concentrations of digitonin

and analyzed by the luciferase assay with the dsRNA biosensor. **(C)** Schematic

representations of cell lysis and phase separation with Triton X-114. The aqueous phase of

the cell lysate was used for the luciferase assay with the dsRNA biosensor. **(D)** Cells at 48

hpi with JEV were suspended in lysis buffer containing different concentrations of Triton

X-114. Cell suspensions containing Triton-X114 [phase separation (-)] and the aqueous

phase of the cell lysate after centrifugation [phase separation (+)] were analyzed by the

luciferase assay with the dsRNA biosensor. The values shown are mean ± SD of triplicate

samples. $*p < 0.05$, $**p < 0.01$, $****p < 0.0001$ by one-way ANOVA with Dunnett's test

(A) or two-tailed Student's *t*-test (D).

Figure 5. Application of the dsRNA biosensor to antiviral assays

(A, B) Cells were infected with JEV in the presence of different concentrations of ribavirin

(A) or NITD008 (B). Immunofluorescence staining of the cells at 48 hpi using anti-JEV

(green) and Hoechst 33342 (blue). Scale bars, 50 μm. **(C, D)** Viral RNA levels were

measured in the cells at 48 hpi by qRT-PCR. Relative viral RNA amounts in the presence

of ribavirin (C) and NITD008 (D) were compared with those in the absence of the antivirals.

Data were normalized to endogenous *Actb*. **(E, F)** Cells at 48 hpi with JEV were lysed in

Triton X-114-containing lysis buffer. The aqueous phase of the lysates was obtained after

phase separation and used for dsRNA quantification with the luciferase assays with the

dsRNA biosensor. Relative luminescence signals in the presence of ribavirin (E) and

NITD008 (F) were compared with those in the absence of the antivirals. The values shown

are mean ± SD of triplicate samples. $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$ by one-way

ANOVA with Dunnett's test (C-F).