Simple and quantitative detection of double-stranded RNA as a signature of pan-virus

infections using a luciferase-based biosensor

Authors: Michihito Sasaki[1,2,9]*, Eri Fujii[1], Satoko Sasaki[1], Takuma Ariizumi[1], Kei

Konishi[3,4], Akihiko Sato[2,3,4], William W. Hall[5,6,7], Hirofumi Sawa[2,3,5,7,8] and Yasuko

Orba[1,2,3,5,8]*

Affiliations: [1]Division of Molecular Pathobiology, International Institute for Zoonosis

Control, Hokkaido University, Sapporo, Japan

[2]Institute for Vaccine Research and Development (IVReD), Hokkaido University, Sapporo,

Japan

[3]Division of Anti-Virus Drug Research, International Institute for Zoonosis Control,

Hokkaido University, Sapporo, Japan

[4]Laboratory for Drug Discovery & Disease Research, Shionogi & Co., Ltd., Osaka, Japan

[5]International Collaboration Unit, International Institute for Zoonosis Control, Hokkaido

University, Sapporo, Japan

[6]National Virus Reference Laboratory, School of Medicine, University College of Dublin,

Ireland

[7]Global Virus Network, Baltimore, Maryland, USA

[8]One Health Research Center, Hokkaido University, Sapporo, Japan

[9]Lead contact

*Correspondence: m-sasaki@czc.hokudai.ac.jp (M.S.), orbay@czc.hokudai.ac.jp (Y.O.)

<u>Summary</u>

Viral infections produce double-stranded RNA (dsRNA) during replication, which trigger

host innate immune responses. Immunoassays using anti-dsRNA antibodies have been

widely employed to detect viral dsRNA. In this study, we used a luciferase-based dsRNA

biosensor which consists of protein kinase R (PKR)-derived dsRNA binding domains fused

to split luciferase subunits. Here, we demonstrate the use of the dsRNA biosensor to

measure viral dsRNA in RNA specimens extracted from cells infected with Japanese

encephalitis virus (JEV). Moreover, the biosensor reacts to a broad-spectrum of dsRNAs

from infection with representatives of various viral families including positive- and

negative-sense ssRNA viruses, dsRNA viruses, and DNA viruses. We validated the specific

interaction between the dsRNA biosensor and viral RNA through RNA

immunoprecipitation. Additionally, we observed luminescence signals directly from lysates

of JEV-infected cells after cell lysis and phase separation with Triton X-114. Finally, we

used the biosensor to assess the activity of antiviral compounds. Collectively, our results

demonstrate that the luciferase-based dsRNA biosensor provides a simple, homogeneous,

and high-throughput platform for the quantification of viral replication, offering a

promising alternative to antibody-based dsRNA detection methods.

<u>Keywords</u>

dsRNA, virus infection, biosensor, RNA virus, DNA virus, luciferase complementation,

cell-based assay

Introduction

The genomes of viruses include encapsided DNA, single-stranded RNA (ssRNA), or double-stranded RNA (dsRNA) and viruses are classified based on their genome types and replication strategies (Baltimore classification)[1]. dsRNA comprises the nucleic acid of dsRNA viruses; however, this type of nucleic acid is also produced by ssRNA and DNA viruses as a replication intermediate and as a byproduct during viral replication in cells[2]. As positive-sense ssRNA viruses replicate, they generate both viral genome–antigenome hybrid dsRNA (replicative form) and partial dsRNA that has multiple transcribing genomic RNAs (replicative intermediate) [3,4]. Copy-back defective viral genomic RNA is a partial viral genomic RNA characterized by reverse complementary 5′ and 3′ ends forming a dsRNA stem structure and is generated during the replication of various non-segmented negative-sense ssRNA viruses[5]. In the case of DNA virus infection, converging bidirectional transcription of the viral genome produces complementary mRNA transcripts that can form dsRNA[6,7]. Collectively, the aberrant accumulation of dsRNA serves as a biomarker for viral infection.

Host cells counter viral infection through several dsRNA sensor proteins such as RIG-I-like receptors (RLRs), protein kinase R (PKR), and oligoadenylate synthases (OASes)[2]. In particular, PKR is a dsRNA-dependent Ser/Thr protein kinase consisting of two tandem dsRNA binding domains (dsRBD) and a kinase domain. PKR recognizes dsRNA in a sequence-independent manner and requires a minimum of 30 to 33 bp of dsRNA for its activation[8,9]. When PKR interacts with dsRNA, the former is autophosphorylated, followed by the phosphorylation of eukaryotic initiation factor 2 alpha

(eIF2α) which results in translational shutoff and viral proliferation[10,11]. PKR can sense

various viral infections, and it mediates a universal antiviral effect in host cells. In response,

some viruses have evolved strategies to antagonize the innate immune responses mediated

by the dsRNA-PKR-eIF2α axis[10,11].

Anti-dsRNA monoclonal antibodies specifically recognize dsRNA longer than 40 bp in

a sequence-independent manner[12]. Clone J2, the most commonly employed anti-dsRNA

monoclonal antibody, has been used as a tool to detect a wide range of viral infections in

various immunoassays including immunofluorescence[3,13,14], immunohistochemistry[15,16],

immunoprecipitation[17], and immunochromatographic assays[18]. J2 has also been used to

determine the subcellular localization and accumulation of dsRNA during the process of

viral genome replication; furthermore, it has been used to study the mechanisms of antiviral

innate immune responses in various viral infections[3,14]. Virological studies have relied

solely on immunoassays using anti-dsRNA antibodies to detect and quantify dsRNA, and

these are limited in their sensitivity, quantitation and throughput. Therefore, it is desirable

to develop other more sophisticated dsRNA detection methods.

One alternative dsRNA detection tool is currently available as a luciferase assay

system using a dsRNA biosensor (Lumit dsRNA Detection Assay, Promega). This system

is based on a split luciferase complementation using small and large subunits of NanoLuc

luciferase, referred to as smBiT and LgBiT, respectively, by the manufacturer[19]. The

biosensor consisting of PKR-derived dsRBD fused with smBiT or LgBiT (dsRBD-smBiT

or dsRBD-LgBiT, respectively) binds to dsRNA and reconstitutes the dsRBD-tagged

smBiT and LgBiT subunits as a functional NanoLuc luciferase[20,21]. The performance of this

luciferase assay system has been evaluated by quantifying artificially synthesized dsRNA

and detecting dsRNA contamination as a byproduct in *in vitro* transcription products[20]. The

affinity of PKR for virus-induced dsRNA facilitates the use of this luciferase-based dsRNA

biosensor as a new research tool in the field of virology. In this study, we have

characterized the interaction between this biosensor and viral RNA and examined its utility

in various types of viral infections.

Results

Detection of dsRNA accumulation produced by flavivirus infection using the

luciferase-based dsRNA biosensor

Japanese encephalitis virus (JEV) is a single-stranded positive-sense RNA virus in the

family *Flaviviridae*. Previous studies have used immunostaining with anti-dsRNA

monoclonal antibodies to show the accumulation of dsRNA in cells infected with

flaviviruses[22,23]. Consistent with previous reports, our immunofluorescence analysis

demonstrated the accumulation of dsRNA in BHK-21 cells infected with JEV (Figure 1A).

Total RNA was extracted from JEV-infected cells at different time points, and quantitative

RT-PCR (qRT-PCR) analysis revealed increases in intracellular viral RNA levels over time

(Figure 1B). To examine the reactivity of dsRBD-LgBiT and dsRBD-smBiT with dsRNA

produced during viral replication, RNA samples from JEV-infected cells were incubated

with dsRBD-LgBiT and dsRBD-smBiT, and the resulting luminescence was measured

(Figure 1C). The results showed significant higher signals in the RNA samples at 48 and 72

hpi than that in RNA of uninfected controls (Figure. 1D). Next, we performed the luciferase

assay and qRT-PCR using RNA samples mixed with RNAs from JEV-infected and

uninfected cells in different ratios. The luminescence and qRT-PCR signals were closely

correlated (R^2 = 0.9829, Figure 1E), although we observed background luminescence in the

assay using RNA from uninfected cells, which affected the limit of detection of the assay.

The luminescence signals from JEV-infected Vero and N2a cells indicated that the assay is

independent of cell type (Figures 1F and 1G). Overall, these results indicate the

applicability of the luciferase-based dsRNA biosensor for detecting dsRNA accumulation

during JEV replication.

Broad-spectrum reactivity of the luciferase-based dsRNA biosensor to different viral

infections

To examine whether the dsRNA biosensor can detect the replication of various virus

types, we tested this in infections caused by the following viruses: positive-sense ssRNA

chikungunya virus (CHIKV, family *Togaviridae*) and SARS-CoV-2 (family *Coronaviridae*

family); negative-sense ssRNA rabies virus (RABV, family *Rhabdoviridae*) and La Crosse

virus (LACV, family *Peribunyaviridae*); dsRNA rotavirus A (RVA, family *Sedoreoviridae*)

and Nelson Bay orthoreovirus (NBV, family *Spinareoviridae*); and dsDNA mpox virus

(MPXV, family *Poxviridae*) and herpes simplex virus 1 (HSV-1, family

Orthoherpesviridae). Immunofluorescence analysis with anti-dsRNA antibody

demonstrated dsRNA accumulation in Vero (CHIKV, LACV, NBV and MPXV),

Vero-TMPRSS2 (SARS-CoV-2), BHK-21 (RABV), and MA104-T2T11D cells (RVA)

infected with all the viruses, except Vero cells infected with HSV-1 (Figure 2A). It has

been reported that virion host shutoff RNase (vhs) of HSV-1 destabilizes dsRNA and limits

its accumulation in cells[6,7]. dsRNA luciferase assay of total RNA extracted from cells

infected with each virus showed a significant increase in luminescence signal in RNA

samples at 24 or 48 hpi with all the tested viruses (Figure 2B). This confirms that the

dsRNA biosensor reacts with a broad range of virus families.

Specificity of the interaction of luciferase-based dsRNA biosensor with flavivirus

genomic and subgenomic RNAs

We investigated the target specificity of the dsRNA biosensor. Heat denaturation of

sample RNA reduced the luminescence signal of JEV-infected cells to a baseline level,

similar to that of uninfected cells (Figure 3A), indicating that the viral infection-specific

increase of the luminescence signal depends on the RNA structure. We examined the nature

of the interaction between the dsRNA biosensor and JEV RNA by RNA

immunoprecipitation (RIP) with JEV-infected cells using dsRBD-LgBiT, anti-LgBiT

antibody, and protein G beads. The LgBiT protein lacking the dsRBD domain was used as

a control in the RIP assay. Immunoblotting results showed the successful capture of

dsRBD-LgBiT and LgBiT on the Protein G beads (Figure 3B). Viral genome RNA, viral

mRNA, and subgenomic flavivirus RNA (sfRNA) are synthesized during active viral

replication in cells[24]. Quantification of the amount of viral RNA in the immunoprecipitate

by qRT-PCR targeting the NS2A region and 3′-UTR of the JEV genome revealed the

specific interaction between the viral RNA and dsRBD-LgBiT (Figures 3C and 3D).

Northern blot analysis with an RNA probe targeting the 3′-UTR of the JEV genome

identified the JEV RNA genome, viral replicative form RNA, and sfRNA in the RNA

extracted from JEV-infected cells (Figure 3E). Furthermore, the analysis of the

immunoprecipitants revealed that dsRBD-LgBiT binds to the JEV RNA genome and

sfRNA (Figure 3E). Due to the limited availability of dsRBD-LgBiT, we failed to scale-up

the RNA yield in the RIP assay. Thus, we were unable to detect the band of JEV replicative

form RNA in the immunoprecipitants in the Northern blot analysis. Collectively, these

results suggest that the dsRNA biosensor specifically recognizes viral RNA synthesized in

the cells infected with JEV.

Quantitative detection of dsRNA in a homogeneous cell lysates using the

luciferase-based dsRNA biosensor

All the above experiments for dsRNA detection were performed with RNA extracted

from cells. We aimed to simplify the protocol by skipping the RNA extraction process and

detecting viral dsRNA directly in cell lysates using the dsRNA luciferase assay. First, we

assessed whether detergent contaminating the cell lysate could affect the dsRNA luciferase

assay. Using an extracted RNA sample containing 1% of conventional detergents, we found

that contamination by all tested detergents decreased the luminescence signal from the

dsRNA biosensor (Figure 4A). Next, we suspended JEV-infected cells in lysis buffer with

different concentrations of digitonin and added the dsRNA biosensor to the mixture. As

expected, a high concentration of digitonin was necessary for sufficient cell lysis, and the

carry-over of digitonin from the input cell lysate with high concentrations of the detergent

attenuated the luminescence signal in the dsRNA luciferase assay (Figure 4B). To reduce

detergent carry-over after sufficient cell lysis, we employed a phase separation approach

using Triton X-114, a nonionic detergent that forms a homogeneous solution on ice.

However, the solution separates into aqueous and detergent-enriched phases above 20□[25].

We suspended JEV-infected cells in the lysis buffer with Triton X-114 on ice and then

incubated the cell lysate at 37□, followed by brief centrifugation to separate the phases

(Figure 4C). The dsRNA luciferase assay detected a high luminescence signal in the

aqueous phase of the cell lysate even with high concentrations of Triton X-114. Without

phase separation, the detergent contamination reduced the luminescence signals in the cell

lysates (Figure 4D). Overall, these results demonstrate that phase separation of cell lysates

with Triton X-114 is a simple and reproducible way to directly detect dsRNA in cells.

We next examined whether our assay can measure the activity of antiviral compounds.

Ribavirin and NITD008 are nucleotide analogs that act as antiviral agents against

flaviviruses, including JEV[26-28]. Both immunofluorescence staining and qRT-PCR assays

demonstrated the anti-JEV activities of ribavirin and NITD008 (Figures 4E-4H). Consistent

with the results of conventional assays, the cell-directed dsRNA luciferase assays showed

that JEV infection decreased in a dose-dependent manner in response to these antivirals

(Figures 4I, 4J). These results suggest that the cell-directed dsRNA luciferase assay is a

simple and fast method to estimate the magnitude of viral replication and the activity of

antiviral compounds.

Discussion

Replication of RNA and DNA viruses in cells produces viral genome-derived RNA

duplexes that stimulate the host antiviral immune responses through dsRNA sensors such

as RLRs, TLR-3, and PKR. In this study, we used the dsRNA biosensor dsRBD-LgBiT and

dsRBD-smBiT that employ the NanoLuc luciferase complementation process to detect viral

dsRNA in infected cells. We could demonstrate that luminescence signal from the

biosensor increases in a time-dependent manner and correlates with the viral RNA load in

JEV-infected cells. The biosensor could also be used to detect different viral infections

including those caused by ssRNA viruses, dsRNA viruses, and DNA viruses, without the

need for antibodies and oligonucleotide primers targeting specific viral proteins and

genomes. These results indicate that the luciferase-based dsRNA biosensor is capable of

estimating the magnitude of various types of viral infections including emerging and as yet

undiscovered viruses.

Anti-dsRNA monoclonal antibodies have been widely used to detect viral dsRNA in

various immunoassays as the protocol has broad applications and involves a simple

workflow. However, antibody-based immunoassays can sometimes be associated with

nonspecific reactions and variable signal intensity[29]. The dsRNA luciferase assay is not an

immunoassay; rather, the dsRNA biosensor carries the dsRBD from PKR. Both the dsRNA

antibody and PKR recognize dsRNA in a sequence-independent manner. Using a

crosslinking and immunoprecipitation sequencing (CLIP-seq) analysis, Kim *et al*. revealed

that PKR as well as the anti-dsRNA antibody J2 can interact with various types of dsRNA[30].

The results from our studies suggest that the luciferase-based dsRNA biosensor is a

versatile method for viral dsRNA detection with /quantification.

PKR displays sequence-nonspecific interaction and autoactivates by not only dsRNA

but also ssRNA with an imperfect short stem flanked by single-stranded tails[31,32]. The

dsRNA biosensor recognizes dsRNA larger than 30 bp regardless of the nucleotide

sequence composition[20]. Although we were able to demonstrate the interaction between the

biosensor and the JEV genome and sfRNA, the details on the target regions within the viral

RNAs remain elusive. There are multiple stem-loop structures in the 5′- and 3′-UTR of the

flavivirus genome and sfRNA[33,34]. Notably, a previous study showed that viral sfRNA

induces the activation of host PKR in cells[35]. In addition, the long dsRNA structure of

flavivirus RNA genome consisting of a hybrid of the 5′-UTR and 3′-UTR was formed

during genomic RNA cyclization (a process that occurs during flavivirus replication)[33,34].

We thus speculate that the stem-loops in JEV genomic RNA and sfRNA are putative

binding targets of dsRNA biosensor.

In this study, we showed the applicability of the dsRNA biosensor in a homogeneous

cell lysate. Although the interaction between the dsRNA biosensor and dsRNA was

attenuated by common detergents used for cell lysis, implementing a phase separation step

following cell lysis with Triton X-114 can effectively mitigate detergent interference while

maintaining the assay's sensitivity. Our cell-directed dsRNA luciferase assay was

applicable to measure the antiviral activity of compounds from cell lysates, and the results

correlated well with those of qRT-PCR. Implementing the dsRNA luciferase assay directly

on cells is a simple and fast way to evaluate the magnitude of viral infections. We believe

that the assay will also be useful for high-throughput screening of antiviral compounds and

host factors.

Limitations of the study

Although our study demonstrated the broad reactivity of the dsRNA biosensor in the

detection of various viral infections, it has the following limitations. First, the target

dsRNAs in each viral infection, except for JEV, remain elusive. The dsRBD in the

biosensor is derived from PKR; thus, identifying the targets of the biosensor will contribute

to our knowledge on the process of PKR activation in response to viral infection. Second,

the presence of endogenous dsRNA needs to be considered. The cellular levels of

host-derived dsRNAs such as inverted Alu repeats, bidirectional transcription of

mitochondrial DNA, and cellular mRNA with an extended 3′-UTR increase in response to

pathophysiological conditions[30,36]. In addition to virus-induced dsRNA, host-derived

dsRNAs are also sensed by endogenous dsRNA sensors. In neurons, mRNA with an

extended 3′-UTR acts as an endogenous dsRNA that triggers constitutive type I IFN

production, conferring an intrinsic antiviral state[37]. However, we assumed that the

endogenous dsRNA had little effect in this study based on the result of heat denaturation

treatment of total RNA from uninfected cells, which showed limited impact on the

luciferase-based dsRNA assay. Finally, we observed background luminescence even in

total RNA samples from uninfected cells. This indicates that it may be necessary to further

optimize the luciferase assay protocol to ensure its wide dynamic range and robustness in

future applications.

Acknowledgments

We thank Dr. Saijo and Dr. Ebihara of the National Institute of Infectious Diseases, Japan

for providing JEV, CHIKV, SARS-CoV-2, RABV, NBV and MPXV, and Dr. Kawaguchi

of Tokyo University for providing HSV-1. We also thank Dr. Inagawa of Promega Japan

for technical support in the Lumit dsRNA detection assays. This work was supported by the

Japan Agency for Medical Research and Development (AMED) under Grant numbers

JP24fk0108637, JP243fa627005, JP24wm0125008, JP24wm0325073; Japan Science and

Technology Agency (JST) Moonshot R&D under Grant number JPMJMS2025.

Author Contributions

Conceptualization, M.S., and Y.O.; Investigation, M.S., E.F., S.S. and T.A.; Formal

analysis, M.S.; Methodology, K.K., A.S.; Resources, H.S., Y.O.; Funding acquisition, A.S.,

H.S., and Y.O.; Writing – original draft, H.S., W.W.H., H.S., and Y.O.; Writing – review &

editing, all authors.

Declaration of interests

The authors K.K. and A.S. are employees and shareholders of Shionogi & Co., Ltd. The

remaining authors declare no competing interests.

Materials and Methods

Cell lines

BHK-21 (RCB1423, RIKEN BRC) and C6/36 cells (CRL-1660, ATCC) were

maintained in Eagle's Minimum Essential Medium (MEM) containing 10% fetal bovine

serum (FBS). N2a cells (IFO50081, JCRB) were maintained in MEM containing 10% FBS

and nonessential amino acids (Gibco). Vero-TMPRSS2 cells stably expressing human

TMPRSS2 were established in our laboratory as described previously[38,39]. Vero-TMPRSS2,

Vero (CRL-1586, ATCC), and 293T cells (632180, Takara Bio) were maintained in

Dulbecco's Modified Eagle's Medium (DMEM) containing 10% FBS. MA104-T2T11D

cells expressing human TMPRSS2 and TMPRSS11D were established in our laboratory as

described previously[40] and maintained in MEM containing 10% FBS and 10% tryptose

phosphate broth (BD Difco).

Viruses

JEV (Mie/41/2002 strain)[41], CHIKV (SL10571 strain)[42], SARS-CoV-2 (WK-521

strain)[43], RABV (HEP strain)[44], NBV (MB/07 strain)[45] and MPXV (Zr-599 strain)[46] were

provided by the National Institute of Infectious Diseases, Japan. HSV-1 (F strain) was

provided by the Institute of Medical Science, University of Tokyo, Japan. RVA (SA11

strain) and LACV (VR-1834) were obtained from the ATCC. Working virus stocks were

prepared by passaging through C6/36 cells for JEV; BHK-21 cells for RABV; Vero cells

for CHIKV, NBV, MPXV, LACV and HSV-1; Vero-TMPRSS2 cells for SARS-CoV-2;

and MA104-T2T11D cells for RVA.

Immunofluorescence staining

BHK-21 cells were infected with JEV or RABV at a multiplicity of infection (MOI) of

0.1. Vero cells were infected individually with CHIKV, LACV, NBV, and HSV-1 at an

MOI of 0.1; and with MPXV at an MOI of 0.01. Vero-TMPRSS2 cells and

MA104-T2T11D cells were respectively infected with SARS-CoV-2 and RVA at an MOI

of 0.1. Cells were fixed at 24 h post-infection (hpi) except for MPXV, which was fixed at

48 hpi with 3.7% buffered formaldehyde, followed by permeabilization with 0.5% Triton

X-100 in PBS for 5 min and double–staining with anti-dsRNA mouse IgG2a monoclonal

antibody (clone J2; 10010200, SCICONS) and virus-specific primary antibody in 25%

Block Ace (KAC) in PBS for 1 h. We used anti-CHIKV E1 mouse IgG1 monoclonal

(MAB12424, Abnova), anti-SARS-CoV-2 nucleocapsid rabbit monoclonal (GTX635679,

GeneTex), anti-RABV nucleoprotein mouse IgG2b monoclonal (3R7-4G4), anti-LACV

mouse IgG2b monoclonal (MA1-10801, Thermo Fisher Scientific), anti-RVA goat

polyclonal (AB1129, Merck), anti-Vaccinia virus rabbit polyclonal (cross reactive with

MPXV; PA17258, Thermo Fisher Scientific), anti-HSV-1 rabbit polyclonal (B0114,

DAKO), anti-JEV rabbit polyclonal (serum from JEV-immunized rabbit in our laboratory),

and anti-NBV rabbit polyclonal (serum from NBV-immunized guinea pig in our

laboratory) antibodies[47] for the viral antigen staining. The cells were subsequently stained

for 30 min with the following appropriate secondary antibodies: anti-mouse IgG2a Alexa

Fluor 488-conjugated goat polyclonal (A21131, Thermo Fisher Scientific), anti-mouse

IgG1 Alexa Fluor 594-conjugated goat polyclonal (A21125, Thermo Fisher Scientific),

anti-mouse IgG2b Alexa Fluor 594-conjugated goat polyclonal (A21145, Thermo Fisher

Scientific), anti-rabbit IgG Alexa Fluor 594-conjugated goat polyclonal (A11037, Thermo

Fisher Scientific), anti-guinea pig IgG Alexa Fluor 647-conjugated goat polyclonal

(A21450, Thermo Fisher Scientific), anti-mouse IgG Alexa Fluor 488-conjugated donkey

polyclonal (A21202, Thermo Fisher Scientific), and anti-goat IgG Alexa Fluor

568-conjugated donkey polyclonal (A11057, Thermo Fisher Scientific). Cell nuclei were

stained with Hoechst 33342 (Thermo Fisher Scientific). Fluorescent images were captured

using an IX73 fluorescence microscope (Olympus).

Viral RNA measurement by qRT-PCR

Virus growth kinetics were determined with BHK-21 cells infected with JEV at an

MOI of 0.01. Total RNA was extracted from the cells at different time points with the

Direct-zol RNA MiniPrep Kit (Zymo Research) and analyzed by qRT-PCR with the

EXPRESS One-Step Superscript qRT-PCR Kit (Thermo Fisher Scientific) and a

primer/probe set targeting JEV *NS2A*[48]. The hamster *Actb* gene was used as an internal

control for normalization[49]. For the RIP assay, the 3′-UTR of the JEV genome was

quantified, by qRT-PCR on the extracted RNA using a One-Step TB Green PrimeScript

PLUS RT-PCR Kit (Takara Bio). JEV RNA levels were calculated by the standard curve,

comparative ΔCt or ΔΔCt methods.

Detection of dsRNA in RNA extracts

BHK-21 cells were infected with JEV or RABV at an MOI of 0.01. Vero cells were

individually infected with CHIKV, LACV, NBV, MPXV, and HSV-1 at an MOI of 0.01.

Vero-TMPRSS2 cells and MA104-T2T11D cells were infected with SARS-CoV-2 and

RVA, respectively, at an MOI of 0.01. At 1 and 48 hpi, total RNA was extracted from the

cells with RNAiso Plus and the Direct-zol RNA MiniPrep Kit, with mock-infected cells

included as controls. We quantified dsRNA using the Lumit dsRNA biosensor

(dsRBD-LgBiT and dsRBD-smBiT) with dsRNA assay buffer and luminescence detection

substrate B provided in the Lumit dsRNA Detection Assay kit (W2041, Promega)

according to the manufacturer's instructions with slight modifications. Briefly, a total of

200 to 500 ng RNA in 5 μl water was mixed and incubated with dsRDB-LgBiT and

dsRBD-smBiT in 95 μl of dsRNA assay buffer for 1 h at room temperature. The mixture of

RNA and biosensor was then added to 25 μl of luminescence detection substrate B diluted

in dsRNA assay buffer on 96-well half area black microplates (3694, Corning). After

15-min of incubation, the luminescence was measured with a GloMax Discover Microplate

Reader (Promega).

dsRNA was denatured by heating RNA samples for 5 min at 65□, then chilling on ice

for 3 min, heating for 2 min at 95□, and finally chilling on ice for 3 min. The amount of

dsRNA in the denatured RNA samples was quantified as described above.

dsRNA detection in the cell lysates

BHK-21 cells on a 24-well plate were infected with JEV at an MOI of 0.01. At 48 hpi,

cells were lysed in 100 μl/well of cell lysis buffer (25 mM Tris-HCl [pH 7.5], 150 mM

NaCl, 5 mM EDTA) supplemented with different concentrations of digitonin (0.001 to 4%),

Triton X-114 (0.001 to 4%), NP-40 (1%), Brij 98 (1%) (all from Sigma), and Triton X-100

(1%) (Nacalai Tesque) for 5 min on ice. The phases in the Triton X-114-containing cell

lysates were separated by first incubating the mixture for 1 min at 37□ and then

centrifuging at $9000 \times g$ for 30 s. The amount of dsRNA was measured in the upper

detergent-free phase. We analyzed cell lysates that had both undergone and not undergone

phase separation by mixing 5 μl of the lysates with dsRDB-LgBiT and dsRBD-smBiT in 95

μl of dsRNA assay buffer and analyzing the solution using the Lumit dsRNA Detection

Assay kit as described above.

RIP assay

Fifty microliters of protein G-coupled magnetic beads (Dynabeads protein G; Veritas)

were conjugated with 10 μg of anti-LgBiT monoclonal antibody (N7100, Promega) for 15

min and then incubated with 0.3 μM concentrations of control LgBiT (Promega) or

dsRBD-LgBiT in PBS with 0.002% Tween 20 (PBST) for 30 min. Then 10 μg of total

RNA extracted from BHK-21 cells at 48 hpi with JEV was incubated with the magnetic

beads in 500 μl of dsRNA assay buffer supplemented with 60 units of a recombinant RNase

inhibitor (Takara Bio) overnight at 4□ with rotation. The mixture was then washed 3 times

with PBST, and 10% of the beads were suspended in SDS sample buffer [4% SDS, 100

mM Tris-HCl (pH 6.8), 20% glycerol, 10% 2-mercaptoethanol] for immunoblotting. The

remaining 90% of the beads were suspended in RNAiso plus (Takara Bio) and the

suspension was subjected to RNA extraction. The RNA extract was analyzed by qRT-PCR

for JEV RNA levels in the immunoprecipitate as described above and by Northern blotting

for JEV RNA size in the immunoprecipitate.

Our immunoblotting protocol began by resolving input and immunoprecipitated

proteins by SDS-PAGE, followed by blotting of the proteins onto an Immobilon-PSQ

PVDF membrane (Merck). The membrane was incubated with anti-LgBiT monoclonal

antibody (N7100) overnight at 4□ and then incubated with HRP-conjugated anti-mouse

IgG polyclonal antibody (AMI3404, BioSource) for 30 min at room temperature. Immune

complexes were detected using Immobilon Western Chemiluminescent Substrate (Merck)

on an Amersham ImageQuant 800 (Cytiva).

Northern blotting was performed using the DIG Northern Starter Kit (Roche)

according to the manufacturer's instructions with slight modifications. Briefly, we resolved

the following on 1.2% SeaKem GTG agarose-2.2 M formaldehyde gels : RNA samples

eluted from the magnetic beads, RNA extracted from BHK-21 infected with JEV, control

RNAs from uninfected BHK, and JEV virions in culture supernatants. The RNAs were

transferred onto a Nytran SuPerCharge nylon membrane using a Turboblotter (Cytiva) and

UV crosslinked at 120 mJ/cm^2 in an HL-2000 HybriLinker (UVP). For the digoxigenin

(DIG)–labeled RNA probes, template cDNA of the 3′-UTR of the JEV genome fused with

T7 promoter sequence was amplified by RT-PCR and applied to *in vitro* transcription with

T7 RNA polymerase. The UV-crosslinked membrane was hybridized with DIG–labeled

RNA probe in ULTRAhyb Ultrasensitive Hybridization Buffer (Thermo Scientific).

Hybridization complexes were detected with alkaline phosphatase-conjugated anti-DIG

antibody and CDP-Star (all from Roche) on Amersham ImageQuant 800.

Evaluation of antiviral efficacy

BHK-21 cells on 24-well plates were infected with JEV at an MOI of 0.01 in the

presence of different concentrations of ribavirin (Wako) or NITD008 (MedChemExpress).

At 48 hpi, a subset of cells were fixed with 3.7% buffered formaldehyde and stained with

anti-JEV rabbit polyclonal antibody. Another subset of cells on different 24-well plates

were lysed in 100 μl/well of cell lysis buffer supplemented with 1% Triton X-114 for 5 min

on ice. After phase separation, 5 μl of the upper detergent-free phase was analyzed using

the Lumit dsRNA Detection Assay kit to quantify dsRNA as described above. The

remaining lysate was subjected to RNA extraction and qRT-PCR as described above.

Statistical analysis and reproducibility

Statistical significance was determined by one-way analysis of variance (ANOVA)

with Dunnett's test (Figs. 1B, 1D, 1F, 1G, 2B, 4A and 4G-4J), one-way ANOVA with

Tukey's test (Figure 3A), two-tailed Welch's t-test (Figures 3C and 3D) and two-tailed

Student's t-test (Figure 4D). All statistical tests were carried out using Prism version 9.5.1

(GraphPad software). All experiments were repeated independently 2 or 3 times with

similar results. Most results are presented as mean± SD of triplicate samples as specified in

the figure legends.

References

1. Koonin, E.V., Krupovic, M., and Agol, V.I. (2021). The Baltimore Classification of Viruses 50 Years Later: How Does It Stand in the Light of Virus Evolution? Microbiol Mol Biol Rev *85*, e0005321. 10.1128/MMBR.00053-21.

2. Chen, Y.G., and Hur, S. (2022). Cellular origins of dsRNA, their recognition and consequences. Nat Rev Mol Cell Biol *23*, 286-301. 10.1038/s41580-021-00430-1.

3. Triantafilou, K., Vakakis, E., Kar, S., Richer, E., Evans, G.L., and Triantafilou, M. (2012). Visualisation of direct interaction of MDA5 and the dsRNA replicative intermediate form of positive strand RNA viruses. J Cell Sci *125*, 4761-4769. 10.1242/jcs.103887.

4. Son, K.N., Liang, Z., and Lipton, H.L. (2015). Double-Stranded RNA Is Detected by Immunofluorescence Analysis in RNA and DNA Virus Infections, Including Those by Negative-Stranded RNA Viruses. J Virol *89*, 9383-9392. 10.1128/JVI.01299-15.

5. Brennan, J.W., and Sun, Y. (2024). Defective viral genomes: advances in understanding their generation, function, and impact on infection outcomes. mBio *15*, e0069224. 10.1128/mbio.00692-24.

6. Burgess, H.M., and Mohr, I. (2018). Defining the Role of Stress Granules in Innate Immune Suppression by the Herpes Simplex Virus 1 Endoribonuclease VHS. J Virol *92*. 10.1128/JVI.00829-18.

7. Dauber, B., Saffran, H.A., and Smiley, J.R. (2019). The herpes simplex virus host shutoff (vhs) RNase limits accumulation of double stranded RNA in infected cells: Evidence for accelerated decay of duplex RNA. PLoS Pathog *15*, e1008111.

10.1371/journal.ppat.1008111.

8.   Manche, L., Green, S.R., Schmedt, C., and Mathews, M.B. (1992). Interactions

between double-stranded RNA regulators and the protein kinase DAI. Mol Cell Biol

12, 5238-5248. 10.1128/mcb.12.11.5238-5248.1992.

9.   Husain, B., Mukerji, I., and Cole, J.L. (2012). Analysis of high-affinity binding of

protein kinase R to double-stranded RNA. Biochemistry *51*, 8764-8770.

10.1021/bi301226h.

10.   Cesaro, T., and Michiels, T. (2021). Inhibition of PKR by Viruses. Front Microbiol

12, 757238. 10.3389/fmicb.2021.757238.

11.   Zhang, R., and Karijolich, J. (2024). RNA recognition by PKR during DNA virus

infection. J Med Virol *96*, e29424. 10.1002/jmv.29424.

12.   Schönborn, J., Oberstrass, J., Breyel, E., Tittgen, J., Schumacher, J., and Lukacs, N.

(1991). Monoclonal antibodies to double-stranded RNA as probes of RNA structure

in crude nucleic acid extracts. Nucleic Acids Res *19*, 2993-3000.

10.1093/nar/19.11.2993.

13.   Weber, F., Wagner, V., Rasmussen, S.B., Hartmann, R., and Paludan, S.R. (2006).

Double-stranded RNA is produced by positive-strand RNA viruses and DNA

viruses but not in detectable amounts by negative-strand RNA viruses. J Virol *80*,

5059-5064. 10.1128/JVI.80.10.5059-5064.2006.

14.   Deng, X., Hackbart, M., Mettelman, R.C., O'Brien, A., Mielech, A.M., Yi, G., Kao,

C.C., and Baker, S.C. (2017). Coronavirus nonstructural protein 15 mediates

evasion of dsRNA sensors and limits apoptosis in macrophages. Proc Natl Acad Sci

U S A *114*, E4251-E4260. 10.1073/pnas.1618310114.

15.   Richardson, S.J., Willcox, A., Hilton, D.A., Tauriainen, S., Hyoty, H., Bone, A.J.,

Foulis, A.K., and Morgan, N.G. (2010). Use of antisera directed against dsRNA to

detect viral infections in formalin-fixed paraffin-embedded tissue. J Clin Virol *49*,

180-185. 10.1016/j.jcv.2010.07.015.

16. Piantadosi, A., Shariatzadeh, N., Bombin, A., Arkun, K., Alexandrescu, S.,

Kleinschmidt-DeMasters, B.K., and Solomon, I.H. (2023). Double-stranded RNA

immunohistochemistry as a screening tool for viral encephalitis. Am J Clin Pathol

160, 210-219. 10.1093/ajcp/aqad039.

17. Decker, C.J., Steiner, H.R., Hoon-Hanks, L.L., Morrison, J.H., Haist, K.C., Stabell,

A.C., Poeschla, E.M., Morrison, T.E., Stenglein, M.D., Sawyer, S.L., and Parker, R.

(2019). dsRNA-Seq: Identification of Viral Infection by Purifying and Sequencing

dsRNA. Viruses *11*. 10.3390/v11100943.

18. Ku, J., Kim, S., Park, J., Kim, T.S., Kharbash, R., Shin, E.C., Char, K., Kim, Y., and

Li, S. (2020). Reactive Polymer Targeting dsRNA as Universal Virus Detection

Platform with Enhanced Sensitivity. Biomacromolecules *21*, 2440-2454.

10.1021/acs.biomac.0c00379.

19. Dixon, A.S., Schwinn, M.K., Hall, M.P., Zimmerman, K., Otto, P., Lubben, T.H.,

Butler, B.L., Binkowski, B.F., Machleidt, T., Kirkland, T.A., et al. (2016). NanoLuc

Complementation Reporter Optimized for Accurate Measurement of Protein

Interactions in Cells. ACS Chem Biol *11*, 400-408. 10.1021/acschembio.5b00753.

20. Moravec, R., Wang, J., Hartnett, J., Cong, M., and Grailer, J. (2024). Development

of novel luminescent assays for sensitive and specific quantitation of

double-stranded RNA in mRNA vaccines. Cancer Research *84*, 3254-3254.

21. Silas, D.S., Juneja, B., Kaur, K., Narayanareddy Gari, M., You, Y., Moon, Y., Chen,

Y., Arora, S., Hansen, J., Muthusamy, K., et al. (2024). Development of Biolayer

Interferometry (BLI)-Based Double-Stranded RNA Detection Method with

Application in mRNA-Based Therapeutics and Vaccines. Pharmaceutics *16*.

10.3390/pharmaceutics16091227.

22.   Miorin, L., Romero-Brey, I., Maiuri, P., Hoppe, S., Krijnse-Locker, J.,

Bartenschlager, R., and Marcello, A. (2013). Three-dimensional architecture of

tick-borne encephalitis virus replication sites and trafficking of the replicated RNA.

J Virol *87*, 6469-6481. 10.1128/JVI.03456-12.

23.   Tabata, K., Itakura, Y., Ariizumi, T., Igarashi, M., Kobayashi, H., Intaruck, K.,

Kishimoto, M., Kobayashi, S., Hall, W.W., Sasaki, M., et al. (2023). Development

of flavivirus subviral particles with low cross-reactivity by mutations of a distinct

antigenic domain. Appl Microbiol Biotechnol *107*, 7515-7529.

10.1007/s00253-023-12817-5.

24.   Slonchak, A., and Khromykh, A.A. (2018). Subgenomic flaviviral RNAs: What do

we know after the first decade of research. Antiviral Res *159*, 13-25.

10.1016/j.antiviral.2018.09.006.

25.   Bordier, C. (1981). Phase separation of integral membrane proteins in Triton X-114

solution. J Biol Chem *256*, 1604-1607.

26.   Tsao, C.H., Su, H.L., Lin, Y.L., Yu, H.P., Kuo, S.M., Shen, C.I., Chen, C.W., and

Liao, C.L. (2008). Japanese encephalitis virus infection activates caspase-8 and -9

in a FADD-independent and mitochondrion-dependent manner. J Gen Virol *89*,

1930-1941. 10.1099/vir.0.2008/000182-0.

27.   Zhang, Z.R., Zhang, H.Q., Li, X.D., Deng, C.L., Wang, Z., Li, J.Q., Li, N., Zhang,

Q.Y., Zhang, H.L., Zhang, B., and Ye, H.Q. (2020). Generation and characterization

of Japanese encephalitis virus expressing GFP reporter gene for high throughput

drug screening. Antiviral Res *182*, 104884. 10.1016/j.antiviral.2020.104884.

28.   Li, C., Chen, X., Hu, J., Jiang, D., Cai, D., and Li, Y. (2022). A Recombinant

Genotype I Japanese Encephalitis Virus Expressing a. Int J Mol Sci *23*.

10.3390/ijms232415548.

29.   Thomsen, C., Røge, R., Fred, Å., and Wanders, A. (2023). Immunohistochemical

detection of double-stranded RNA in formalin-fixed paraffin-embedded tissue.

APMIS *131*, 197-205. 10.1111/apm.13300.

30.   Kim, Y., Park, J., Kim, S., Kim, M., Kang, M.G., Kwak, C., Kang, M., Kim, B.,

Rhee, H.W., and Kim, V.N. (2018). PKR Senses Nuclear and Mitochondrial Signals

by Interacting with Endogenous Double-Stranded RNAs. Mol Cell *71*,

1051-1063.e1056. 10.1016/j.molcel.2018.07.029.

31.   Zheng, X., and Bevilacqua, P.C. (2004). Activation of the protein kinase PKR by

short double-stranded RNAs with single-stranded tails. RNA *10*, 1934-1945.

10.1261/rna.7150804.

32.   Nallagatla, S.R., Hwang, J., Toroney, R., Zheng, X., Cameron, C.E., and Bevilacqua,

P.C. (2007). 5'-triphosphate-dependent activation of PKR by RNAs with short

stem-loops. Science *318*, 1455-1458. 10.1126/science.1147347.

33.   Villordo, S.M., and Gamarnik, A.V. (2009). Genome cyclization as strategy for

flavivirus RNA replication. Virus Res *139*, 230-239.

10.1016/j.virusres.2008.07.016.

34.   Saeedi, B.J., and Geiss, B.J. (2013). Regulation of flavivirus RNA synthesis and

capping. Wiley Interdiscip Rev RNA *4*, 723-735. 10.1002/wrna.1191.

35.   Pallarés, H.M., González López Ledesma, M.M., Oviedo-Rouco, S., Castellano,

L.A., Costa Navarro, G.S., Fernández-Alvarez, A.J., D'Andreiz, M.J., Aldas-Bulos,

V.D., Alvarez, D.E., Bazzini, A.A., and Gamarnik, A.V. (2024). Zika virus
non-coding RNAs antagonize antiviral responses by PKR-mediated translational
arrest. Nucleic Acids Res *52*, 11128-11147. 10.1093/nar/gkae507.

36. Dhir, A., Dhir, S., Borowski, L.S., Jimenez, L., Teitell, M., Rötig, A., Crow, Y.J.,
Rice, G.I., Duffy, D., Tamby, C., et al. (2018). Mitochondrial double-stranded RNA
triggers antiviral signalling in humans. Nature *560*, 238-242.
10.1038/s41586-018-0363-0.

37. Dorrity, T.J., Shin, H., Wiegand, K.A., Aruda, J., Closser, M., Jung, E., Gertie, J.A.,
Leone, A., Polfer, R., Culbertson, B., et al. (2023). Long 3'UTRs predispose neurons
to inflammation by promoting immunostimulatory double-stranded RNA formation.
Sci Immunol *8*, eadg2979. 10.1126/sciimmunol.adg2979.

38. Sasaki, M., Uemura, K., Sato, A., Toba, S., Sanaki, T., Maenaka, K., Hall, W.W.,
Orba, Y., and Sawa, H. (2021). SARS-CoV-2 variants with mutations at the S1/S2
cleavage site are generated in vitro during propagation in TMPRSS2-deficient cells.
PLoS Pathog *17*, e1009233. 10.1371/journal.ppat.1009233.

39. Uemura, K., Sasaki, M., Sanaki, T., Toba, S., Takahashi, Y., Orba, Y., Hall, W.W.,
Maenaka, K., Sawa, H., and Sato, A. (2021). MRC5 cells engineered to express
ACE2 serve as a model system for the discovery of antivirals targeting
SARS-CoV-2. Sci Rep *11*, 5376. 10.1038/s41598-021-84882-7.

40. Sasaki, M., Itakura, Y., Kishimoto, M., Tabata, K., Uemura, K., Ito, N., Sugiyama,
M., Wastika, C.E., Orba, Y., and Sawa, H. (2021). Host serine proteases TMPRSS2
and TMPRSS11D mediate proteolytic activation and trypsin-independent infection
in group A rotaviruses. J Virol *95*. 10.1128/JVI.00398-21.

41. Tajima, S., Taniguchi, S., Nakayama, E., Maeki, T., Inagaki, T., Lim, C.K., and

Saijo, M. (2020). Amino Acid at Position 166 of NS2A in Japanese Encephalitis

Virus (JEV) is Associated with In Vitro Growth Characteristics of JEV. Viruses *12*.

10.3390/v12070709.

42. Lim, C.K., Nishibori, T., Watanabe, K., Ito, M., Kotaki, A., Tanaka, K., Kurane, I.,

and Takasaki, T. (2009). Chikungunya virus isolated from a returnee to Japan from

Sri Lanka: isolation of two sub-strains with different characteristics. Am J Trop Med

Hyg *81*, 865-868. 10.4269/ajtmh.2009.09-0009.

43. Matsuyama, S., Nao, N., Shirato, K., Kawase, M., Saito, S., Takayama, I., Nagata,

N., Sekizuka, T., Katoh, H., Kato, F., et al. (2020). Enhanced isolation of

SARS-CoV-2 by TMPRSS2-expressing cells. Proc Natl Acad Sci U S A *117*,

7001-7003. 10.1073/pnas.2002589117.

44. Horiya, M., Posadas-Herrera, G., Takayama-Ito, M., Yamaguchi, Y., Iizuka-Shiota,

I., Kato, H., Okamoto, A., Saijo, M., and Lim, C.K. (2022). Genetic

Characterization of Human Rabies Vaccine Strain in Japan and Rabies Viruses

Related to Vaccine Development from 1940s to 1980s. Viruses *14*.

10.3390/v14102152.

45. Yamanaka, A., Iwakiri, A., Yoshikawa, T., Sakai, K., Singh, H., Himeji, D., Kikuchi,

I., Ueda, A., Yamamoto, S., Miura, M., et al. (2014). Imported case of acute

respiratory tract infection associated with a member of species nelson bay

orthoreovirus. PLoS One *9*, e92777. 10.1371/journal.pone.0092777.

46. Saijo, M., Ami, Y., Suzaki, Y., Nagata, N., Iwata, N., Hasegawa, H., Ogata, M.,

Fukushi, S., Mizutani, T., Sata, T., et al. (2006). LC16m8, a highly attenuated

vaccinia virus vaccine lacking expression of the membrane protein B5R, protects

monkeys from monkeypox. J Virol *80*, 5179-5188. 10.1128/JVI.02642-05.

47.   Harima, H., Sasaki, M., Orba, Y., Okuya, K., Qiu, Y., Wastika, C.E., Changula, K.,

Kajihara, M., Simulundu, E., Yamaguchi, T., et al. (2021). Attenuated infection by a

Pteropine orthoreovirus isolated from an Egyptian fruit bat in Zambia. PLoS Negl

Trop Dis *15*, e0009768. 10.1371/journal.pntd.0009768.

48.   Bharucha, T., Sengvilaipaseuth, O., Vongsouvath, M., Davong, V., Panyanouvong,

P., Piorkowski, G., Garson, J.A., Newton, P.N., de Lamballerie, X., and Dubot-Pérès,

A. (2018). Development of an improved RT-qPCR Assay for detection of Japanese

encephalitis virus (JEV) RNA including a systematic review and comprehensive

comparison with published methods. PLoS One *13*, e0194412.

10.1371/journal.pone.0194412.

49.   Zivcec, M., Safronetz, D., Haddock, E., Feldmann, H., and Ebihara, H. (2011).

Validation of assays to monitor immune responses in the Syrian golden hamster

(Mesocricetus auratus). J Immunol Methods *368*, 24-35. 10.1016/j.jim.2011.02.004.

Figure Legends

Figure 1. Quantitative detection of dsRNA in JEV infection by luciferase assays with

the dsRNA biosensor

(A) Immunofluorescence staining of BHK-21 cells infected with JEV or mock-infected

cells at 24 hpi. Cells were stained with anti-JEV (red), anti-dsRNA (green), and Hoechst

33342 (blue). **(B)** Viral RNA loads in cells at the indicated time points of JEV infection

were analyzed by qRT-PCR. qRT-PCR data were normalized to endogenous *Actb*. **(C)**

Schematic representation of dsRNA detection in RNA sample by split nanoluciferase

complementation assay using the dsRNA biosensor (dsRBD-smBiT and dsRBD-LgBiT).

(D) Total RNA was extracted from BHK-21cells infected with JEV and analyzed by

luciferase assay with the dsRNA biosensor. **(E)** Correlation of the qRT-PCR targeting JEV

RNA and luciferase assays with the dsRNA biosensor. RNA samples from JEV-infected

and uninfected BHK-21 cells were mixed at different ratios. The RNA mixtures were

analyzed by qRT-PCR and luciferase assay to measure the levels of JEV RNA and dsRNA,

respectively. The horizontal dashed line indicates the luminescence value of uninfected

cells in the assay (the detection limit of dsRNA related to JEV infection). R^2 indicates the

linear correlation coefficients between qRT-PCR and dsRNA detection assays. **(F, G)** Total

RNA was extracted from Vero (F) and N2a (G) cells infected with JEV and analyzed by

luciferase assays with the dsRNA biosensor. The values shown are mean ± standard

deviation (SD) of triplicate (B, D, F, G) or duplicate (E) samples. ****$p < 0.0001$ by

one-way ANOVA with Dunnett's test.

Figure 2. Application of the dsRNA biosensor to different viral infections

Positive-sense ssRNA viruses [Chikungunya virus (CHIKV), SARS-CoV-2],

negative-sense ssRNA viruses [Rabies virus (RABV), La Crosse virus (LACV)], dsRNA

viruses [Rotavirus A (RVA), Nelson Bay orthoreovirus (NBV)], DNA viruses [Mpox virus

(MPXV), and Herpes simplex virus 1 (HSV-1)] were inoculated onto monolayers of Vero

(CHIKV, LACV, NBV, MPXV and HSV-1), Vero-TMPRSS2 (SARS-CoV-2), BHK-21

(RABV), and MA104-T2T11D cells (RVA). **(A)** Immunofluorescence staining of the

infected cells with Hoechst 33342 nuclear dye (blue), antibodies targeting dsRNA (green)

and virus proteins (red). Scale bars, 50 μm. **(B)** Luciferase assay with the dsRNA biosensor

was used to measure dsRNA in RNA extracts from mock-infected and virus-infected cells.

The values shown are mean ± SD of triplicate samples. **$p < 0.01$, ****$p < 0.0001$ by

one-way ANOVA with Dunnett's test.

Figure 3. Target specificity of the dsRNA biosensor

(A) Effect of heat denaturation on dsRNA sensing by the dsRNA biosensor. Heat-denatured

and non-denatured RNA samples were analyzed by luciferase assays with the dsRNA

biosensor. **(B-D)** RNA immunoprecipitation (RIP) assays with total RNA extracted from

BHK-21 cells at 48 hpi with JEV using dsRNA biosensor dsRBD-LgBiT and anti-LgBiT

antibody. LgBiT protein without dsRBD was used as a control in the RIP assay.

Immunoprecipitates were analyzed by immunoblotting with anti-LgBiT (B) and qRT-PCR

targeting the JEV NS2A gene (C) and the 3′-UTR of the JEV genome (D). **(E)** Northern

blotting of the immunoprecipitates using digoxigenin DIG–labeled RNA probe targeting

the 3′-UTR of the JEV genome. RNA extracted from the culture supernatant containing

JEV virions, JEV-infected and uninfected BHK-21 cells were included in the blotting to

estimate the size of the JEV genome and subgenomic flavivirus RNA (sfRNA). The values

shown are mean ± SD of triplicate samples. ****$p < 0.0001$ by one-way ANOVA with

Tukey's test (A) or two-tailed Welch's t-test (C, D).

Figure 4. Detection of dsRNA in cell lysates using the luciferase-based dsRNA

biosensor

(A) RNA samples extracted from JEV BHK-21 cells at 48 hpi with JEV were mixed with

the indicated detergents at a final concentration of 1% and then analyzed by the luciferase

assay for dsRNA detection with the dsRNA biosensor. **(B)** Cells at 48 hpi following JEV

infection were suspended in lysis buffer containing different concentrations of digitonin

and analyzed by the luciferase assay with the dsRNA biosensor. **(C)** Schematic

representations of cell lysis and phase separation with Triton X-114. The aqueous phase of

the cell lysate was used for the luciferase assay with the dsRNA biosensor. **(D)** Cells at 48

hpi with JEV were suspended in lysis buffer containing different concentrations of Triton

X-114. Cell suspensions containing Triton-X114 [phase separation (-)] and the aqueous

phase of the cell lysate after centrifugation [phase separation (+)] were analyzed by the

luciferase assay with the dsRNA biosensor. **(E, F)** Cells were infected with JEV in the

presence of different concentrations of ribavirin (E) or NITD008 (F). Immunofluorescence

staining of the cells at 48 hpi using anti-JEV (green) and Hoechst 33342 (blue). Scale bars,

50 μm. **(G, H)** Viral RNA levels were measured in the cells at 48 hpi by qRT-PCR.

Relative viral RNA amounts in the presence of ribavirin (G) and NITD008 (H) were

compared with those in the absence of the antivirals. Data were normalized to endogenous

Actb. **(I, J)** Cells at 48 hpi with JEV were lysed in Triton X-114-containing lysis buffer.

The aqueous phase of the lysates was obtained after phase separation and used for dsRNA

quantification with the luciferase assays with the dsRNA biosensor. Relative luminescence

signals in the presence of ribavirin (I) and NITD008 (J) were compared with those in the

absence of the antivirals. The values shown are mean ± SD of triplicate samples. *$p < 0.05$,

**$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$ by one-way ANOVA with Dunnett's test (A,

G-J) or two-tailed Student's *t*-test (D).