Dorsoventral-mediated *Shh* induction is required for axolotl limb regeneration

Sakiya Yamamoto[1], Saya Furukawa[1], Ayaka Ohashi[1], Mayuko Hamada [1, 2], and Akira Satoh[1]

Affiliations:

[1] Okayama University, Graduate School of Environmental, Life, Natural Science and Technology, Okayama, Japan

[2] Ushimado Marine Institute (UMI), Okayama University, Okayama, Japan

*Corresponding author

Akira Satoh, Ph.D.,

Okayama University, Okayama University, Graduate School of Environmental, Life, Natural Science and Technology

Email: satoha@cc.okayama-u.ac.jp

Tel: +81-86-251-8421

Abstract

Axolotls (*Ambystoma mexicanum*) exhibit a remarkable ability to regenerate limbs after amputation.

Classical experiments have suggested that contact between cells derived from distinct orientations—

dorsal, ventral, anterior, and posterior—within the regenerating blastema is necessary for accurate

limb pattern formation. However, the molecular basis for this requirement has remained largely

unknown. Here, we demonstrate that both dorsal and ventral tissues are required for limb formation

via induction of *Shh* expression, which plays a crucial role in limb patterning. Using the accessory

limb model (ALM), we induced position-specific blastemas lacking cells derived from a single

orientation (anterior, posterior, dorsal, or ventral). Patterned limb formation occurred only in

blastemas containing both dorsal- and ventral-derived cells. We further observed that *Shh* expression

requires dorsoventral contact within a blastema, highlighting the necessity of dorsoventral contact

for inducing *Shh* expression. In addition, we identified WNT10B and FGF2 as dorsal- and ventral-

mediated signals, respectively, that create the inductive environment for *Shh* expression. Our

findings clarify the role of dorsal and ventral cells in inducing *Shh*, a mechanism that has rarely been

studied in the context of limb regeneration and pattern formation. This model provides new insights

into how cells with different positional identities drive the regeneration process.

Introduction

Axolotls (*Ambystoma mexicanum*) possess remarkable regenerative abilities, enabling the regeneration of entire limbs after amputation. Among tetrapods (four-limbed vertebrates), only urodele amphibians retain the lifelong ability to regenerate fully developed limbs. Investigating the molecular mechanisms underlying axolotl limb regeneration could provide valuable insights for advancing regenerative medicine in humans, potentially leading to new therapies for tissue repair and organ regeneration.

The limb regeneration process can be divided into two stages: the blastema (regenerating limb primordium) induction process and the limb patterning process. The induction of a blastema, which contains highly proliferative multipotent and unipotent cells, depends on the presence of nerves at the injured region; when a denervated limb is amputated, a blastema is not induced (Todd, 1823). This blastema induction process is followed by the limb patterning process. The limb patterning process has been considered to be mainly a recapitulation of limb development. Actually, activation of many developmental genes can be found during this phase. Such sequential events accomplish axolotl limb regeneration.

To accomplish limb patterning, cells need to know its address within a limb and exert its position specific role. Such address of cells has been referred as positional identities. In limb development/regeneration, positional identities are thought to be established along three-dimensional axes—anteroposterior, dorsoventral, and proximodistal. In amniote limb development, WNT7A, secreted from dorsal ectoderm, induces *Lmx1b* expression in the underlying mesenchyme, thereby specifying dorsal identity (Riddle et al., 1995; Vogel et al., 1995; Cygan et al., 1997; Chen et al., 1998; Chen, 2002). On the ventral side, the ventral identity is specified indirectly by *En1*, which is expressed in the ventral ectoderm, restricts *Wnt7a* expression to the dorsal ectoderm and thereby prevents induction of *Lmx1b* in the ventral mesenchyme (Loomis et al., 1996; Logan et al., 1997; Chen and Johnson, 2002). Similarly, SHH is known to regulate anteroposterior patterning (Riddle et al., 1993). In axolotl limb regeneration, the establishment of the dorsoventral positional identities is still largely unknown, except for dorsal-specific *Lmx1b* expression (Shimokawa et al., 2013). Regarding the anteroposterior axis, *Shh* expression is highly conserved and restricted to the posterior margin. Though the process of the establishment of the positional identities is still veiled in axolotl limb regeneration, it is apparent that the regeneration blastema possesses regional specificity.

For successful limb patterning in limb regeneration, interactions between anterior, posterior, dorsal, and ventral cells within a blastema have been considered essential. The importance of these interpositional interactions was investigated in creating double-half limbs. Amputating the double-half limbs can provide a blastema in a "one-positional-identity-missing" state. For example,

when a double-dorsal limb—a chimeric limb surgically generated by excising the ventral half and

replacing it with a dorsal half from the contralateral limb while preserving anteroposterior

orientation—is amputated, the resulting blastema lacks ventral positional identity. Such blastemas

lacking one positional identity can form hypomorphic, spike-like structures or fail to regenerate; in

other cases, they regenerate limbs with complete anteroposterior axes accompanied by symmetric

dorsoventral duplications (Bryant, 1976; Bryant and Baca, 1978; Burton et al., 1986). It is

noteworthy that, in the non-regenerating cases, structural patterns along the anteroposterior axis

appear to be lost even though both anterior and posterior cells should, in principle, be present in a

blastema induced from a double-half limb. These observations are consistent with the idea that,

during axolotl limb regeneration, signals mediated by dorsal cells must act on ventral cells, and

signals mediated by ventral cells must act on dorsal cells, during the limb patterning process. In

contrast, in amniote limb development, *Wnt7a/Lmx1b* or *En1* mutants show that limbs can exhibit

anteroposterior patterning even when tissues are dorsalized or ventralized—i.e., in the relative

absence of ventral or dorsal cells, respectively (Riddle et al., 1995; Chen et al., 1998; Loomis et al.,

1996). Taken together, these findings indicate that the mechanism underlying limb patterning during

axolotl regeneration differs from that operating during amniote limb development. Additional

experiments involving ectopic contact between opposite regions (e.g., anterior-posterior or dorsal-

ventral) of a blastema further support the critical role of interpositional interactions in the limb

patterning process, as such interactions can lead to ectopic limb formation (Iten and Bryant, 1975;

Bryant and Iten, 1976; Maden, 1980; Stocum, 1982). These findings suggest that signals mediated

by each positional identity are required for successful limb patterning, such that the absence of any

one of them leads to failure. To further understand limb regeneration, the molecular basis of these

positional identity-mediated signals in limb patterning process should be elucidated.

The accessory limb model (ALM), a non-amputation experimental system, provides a

valuable approach for studying axolotl limb regeneration, particularly the role of interpositional

interactions (Endo et al., 2015). In this model, a blastema is induced through skin wounding

combined with nerve deviation (Endo et al., 2004, Satoh et al., 2007). When ALM surgery is

performed on the anterior side, the induced blastema (ALM blastema) lacks posterior-derived cells

and is unable to form a properly patterned limb. This limitation can be overcome by supplying

posterior-derived cells via a skin graft from the posterior side, enabling the successful regeneration

of a patterned limb. Similarly, an ALM surgery can be performed on the posterior, dorsal, or ventral

side of a limb; in each case, the induced ALM blastema is expected to lack cells from the

corresponding opposite orientation (anterior, ventral, or dorsal, respectively). This condition allows

the investigation of blastema induction and limb patterning by selectively removing cells from

specific orientations of the limb. Analyzing the characteristics and developmental limitations of

these ALM blastemas provides valuable insights into the positional identity-mediated signals that

regulate axolotl limb regeneration.

The molecular basis of the interpositional interactions within a blastema in the limb

patterning process has been partially elucidated (Nacu et al., 2016). In a normal blastema, *Fgf8* and

Shh are expressed at the anterior and the posterior regions, respectively. A previous study

demonstrated that FGF8 and SHH proteins can substitute for anterior and posterior tissues,

respectively, to form a patterned limb. Moreover, these proteins function as mutually inductive

signals, maintaining each other's expression. These findings suggest that FGF8 and SHH serve as

the anterior- and posterior-mediated signals, respectively. In contrast, the molecular basis of dorsal-

and ventral-mediated signals remains poorly understood. Previous studies have shown that retinoic

acid supplementation can convert the dorsal identity to the ventral, enabling patterned limb

formation even in the absence of ventral tissues (McCusker et al., 2014; Ludolph et al 1990; Vieira et

al., 2019). However, the ventral-mediated signal remains uncertain, as retinoic acid induces ventral

positional identities rather than acting as the ventral-mediated signal itself, suggesting the existence

of another, downstream ventral-mediated signal. Consequently, the mechanisms underlying dorsal-

and ventral-mediated signals, as well as how they relate to the anterior- and posterior-mediated

signals (FGF8 and SHH, respectively), remain unclear.

In the present study, we investigate the roles of dorsal- and ventral-mediated signals in

limb patterning. We confirmed the necessity of dorsoventral tissue contact for limb patterning using

the ALM. We found that the induction of *Shh* expression was dependent on the co-existence of

dorsal and ventral cells in axolotl limb regeneration. Furthermore, we identified WNT10B and FGF2

as dorsal- and ventral-mediated signals, respectively, by RNA-seq analysis, and confirmed that these

factors mediate *Shh* expression in axolotl blastemas. Our results suggest that the crucial role of the

dorsal- and ventral-mediated signals in the limb patterning process is to induce *Shh* expression,

thereby enabling anteroposterior interaction. These results contribute to understanding how the

integration of four positional identities—dorsal, ventral, anterior, and posterior—drives proper limb

patterning during axolotl limb regeneration.

Results

Dorsoventral tissue contact is required for limb regeneration in the ALM

We performed the ALM experiment on *Ambystoma mexicanum* to directly test whether dorsoventral

tissue contact, as well as anteroposterior tissue contact, is required for limb patterning (Fig. 1). The

skin was removed from one side (anterior, posterior, dorsal, or ventral) of a limb, and the large nerve

fibers running along the center of a limb (*Nervus medianus* and *Nervus ulnaris*, Fig. 1A) were

dissected and rerouted to the wounded region. We supplied the cells from the contralateral side of a

limb as a skin graft in experimental conditions to ensure the presence of sufficient cells for patterned

limb formation. We conducted eight types of ALM blastema inductions, which can be categorized

into two groups: basic ALM blastemas and skin-grafted ALM blastemas. The basic ALM blastemas

included the anteriorly induced ALM blastema (AntBL), the posteriorly induced ALM blastema

(PostBL), the dorsally induced ALM blastema (DorBL), and the ventrally induced ALM blastema

(VentBL), all of which should lack cells from the contralateral side. The skin-grafted ALM blastemas

included AntBL with posterior skin grafting (AntBL+P), PostBL with anterior skin grafting

(PostBL+A), DorBL with ventral skin grafting (DorBL+V), and VentBL with dorsal skin grafting

(VentBL+D); these ALM blastemas should contain cells with all anteroposterior and dorsoventral

orientations. For the ALM experiment, we defined the two large blood vessels running on the

anterior and posterior sides at the stylopod level (*Vena cephalica* and *Vena. basilica*, Fig. 1A) as the

anterior and posterior dorsoventral borders, respectively. At 10 days post-surgery (dps), an ALM

blastema was observed in all experimental groups (Fig. 1B–I). Consistent with previous studies

(Endo et al., 2004; Nacu et al., 2016), limbs with multiple digits were formed from AntBL+P and

PostBL+A (Table 1, Fig. 1F, G), whereas AntBL and PostBL either regressed or formed bump

structures (Table 1, Fig. 1B, C). Similarly, DorBL+V and VentBL+D formed limbs (Table 1, Fig. 1H,

I), whereas most DorBL and VentBL either regressed or formed bump structures (Table 1, Fig. 1D,

E). Notably, only 1 out of 12 DorBL formed a limb, which may have resulted from ventral tissue

contamination, as the rerouted nerves originally ran along the ventral side of the humerus. In

contrast, none of the 22 VentBL formed multi-digit limbs. These results suggest that both dorsal and

ventral tissues, as well as anterior and posterior tissues, are required for successful limb formation.

Gene expression patterns in ALM blastemas

To elucidate the characteristics of the ALM blastemas that typically fail to form limbs (AntBL,

PostBL, DorBL, and VentBL), we analyzed gene expression patterns at 10 dps (Fig. 2, Fig. S1).

Lmx1b, a gene necessary and sufficient for establishing the dorsal identity in mesenchymal cells in

developing limb buds (Vogel et al., 1995; Chen et al., 1998; Chen, 2002), was used as a dorsal

marker because we previously reported that *Lmx1b* is exclusively expressed in dorsal-derived cells

and can be activated in the ALM experiment (Iwata et al., 2020; Yamamoto et al., 2022). The

expression patterns of *Fgf8*, *Shh*, and *Lmx1b* were investigated using *in situ* hybridization (ISH). In

AntBL and PostBL, *Lmx1b* expression was restricted to the dorsal half of the blastemas (Fig. 2B, G,

Fig. S1A, D, n=5/5 for both), suggesting the presence of both dorsal and ventral tissues. These

expression patterns correspond to the anatomically defined dorsoventral borders (Fig. 1A). In

contrast, *Lmx1b* was expressed throughout the entire region of DorBL (Fig. 2L, Fig. S1G, n=6/6),

whereas no *Lmx1b* expression was detected in VentBL (Fig. 2Q, Fig. S1J, n=6/6), suggesting the

absence of ventral tissue in DorBL and dorsal tissue in VentBL. Consistent with previous studies,

Fgf8 expression was observed in AntBL (Fig. 2C, Fig. S1B, n=4/5), but not in PostBL (Fig. 2H, Fig.

S1E, n=5/5, Nacu et al., 2016). Conversely, *Shh* expression was detected in PostBL (Fig. 2I, Fig.

S1F, n=5/5), but not in AntBL (Fig. 2D, Fig. S1C, n=5/5). In DorBL and VentBL, *Shh* expression

was largely absent (Fig. 2N, Fig. S1I, n = 5/6; Fig. 2S, Fig. S1L, n = 6/6), whereas *Fgf8* expression

was present in both (Fig. 2M, Fig. S1H, n=4/6; Fig. 2R, Fig. S1K, n=4/6). In DorBL, *Shh* expression

was observed in only 1 out of 6 samples, possibly due to ventral tissue contamination, as described

above. These results suggest that *Fgf8* expression is independent of the co-existence of dorsal and

ventral tissues, whereas *Shh* expression appears to require their presence.

Induction of Shh expression requires both dorsal and ventral cells

The absence of *Shh* expression in most DorBL and VentBL, combined with its consistent expression

in all PostBL—which contain both *Lmx1b*-positive and *Lmx1b*-negative regions—suggests that the

induction of *Shh* expression depends on the presence of both dorsal and ventral cells (Fig. 2). To test

this hypothesis, we performed cell-tracing experiments using GFP-expressing skin from transgenic

animals (Fig. 3). We induced VentBL in leucistic animals and then grafted the posterior half of the

dorsal skin (VentBL+PD$_{gfp}$) or the posterior half of the ventral skin (VentBL+PV$_{gfp}$) from GFP

transgenic animals (Fig. 3A). Similarly, DorBL was induced in wild-type animals, and the posterior

half of the dorsal skin (DorBL+PD$_{gfp}$) or the posterior half of the ventral skin (DorBL+PV$_{gfp}$) from

GFP transgenic animals was grafted (Fig. 3A). GFP-positive mesenchymal cells derived from the

posterior skin were expected to express *Shh* if provided with a suitable environment, because

posteriorly derived cells, not anteriorly derived cells are known to have the competency to express

Shh in a blastema—i.e., whether a cell is capable of expressing *Shh* depends on its original positional

identity (Iwata et al., 2020), but whether it actually expresses *Shh* should depend on the environment

in which the cell is placed. If the induction of *Shh* expression depends on the co-existence of dorsal

and ventral cells, *Shh* expression in the GFP-positive cells should be observed in VentBL+PD$_{gfp}$ and

DorBL+PV$_{gfp}$ but not in DorBL+PD$_{gfp}$ or VentBL+PV$_{gfp}$. Samples were collected at 10 dps, and the

expression of *Shh* and *Lmx1b* was analyzed using ISH. In all four experimental groups, *Lmx1b*

expression patterns in the GFP-negative regions were consistent with those in DorBL and VentBL

(Fig. 3D, F, H, J). In DorBL+PD$_{gfp}$ and VentBL+PD$_{gfp}$, *Lmx1b* expression was observed in GFP-

positive mesenchymal cells, whereas GFP-positive cells in DorBL+PV$_{gfp}$ and VentBL+PV$_{gfp}$ were

Lmx1b-negative (Fig. 3C–F). These *Lmx1b* expression patterns suggest that dorsoventral tissue

contact was present in VentBL+PD$_{gfp}$ and DorBL+PV$_{gfp}$ but absent in DorBL+PD$_{gfp}$ and

VentBL+PV$_{gfp}$. In the GFP-positive cells derived from PD$_{gfp}$ skin, *Shh* expression was observed in

VentBL+PD$_{gfp}$ (Fig. 3C, n=4/6) but not in DorBL+PD$_{gfp}$ (Fig. 3E, n=6/6). Similarly, in the GFP-

positive cells derived from PV$_{gfp}$ skin, *Shh* expression was observed in DorBL+PV$_{gfp}$ (Fig. 3I, n=3/6)

but not in VentBL+PV$_{gfp}$ (Fig. 3G, n=7/7). In addition, *Shh* expression was observed in some GFP-

negative mesenchymal cells in samples where *Shh* expression was observed in GFP-positive cells.

These results suggest that the induction of *Shh* expression in posteriorly derived cells requires the

co-existence of dorsal and ventral cells.

Limb formation in the absence of dorsoventral tissue contact

DorBL and VentBL failed to form limbs (Fig. 1D, E). *Shh* expression was absent in these blastemas,

whereas *Fgf8* was expressed (Fig. 2). Results from the cell-tracing experiments suggest that *Shh*

expression requires the co-existence of dorsal and ventral cells (Fig. 3). Based on these findings, we

hypothesized that while the co-existence of dorsal and ventral cells is necessary to induce *Shh*

expression, it is not directly required for limb patterning if SHH protein is externally provided. To

test this hypothesis, we overexpressed *Shh* in DorBL and VentBL (Fig. 4A, B). As a positive control,

Shh was overexpressed in AntBL, where *Fgf8* was expressed but *Shh* expression was absent (Fig.

2C, D), because a previous study has shown that *Shh* overexpression to AntBL can induce limb

patterning (Nacu et al., 2016). As a result, *Shh*-electroporated AntBL, DorBL, and VentBL formed

limbs (n=10/19, 6/18, and 8/14, respectively, Fig. 4C–G). In contrast, GFP-electroporated DorBL

and VentBL (negative controls) failed to form limbs (n=7/7 and 6/6, respectively). We further

analyzed the anatomy of the induced limbs (Fig. 4H–K, Fig. S2). Because ALM-induced limbs

frequently exhibit abnormal and highly variable morphologies, which makes it difficult to use

consistent anatomical landmarks such as particular digits or muscle groups, we focused our analysis

on morphological symmetry. We found that limbs formed from DorBL and VentBL exhibited

symmetric structures along the dorsoventral axis, suggesting double-dorsal and double-ventral

structures, respectively (Fig. 4J, K, Fig. S2C, D). In contrast, limbs formed from AntBL, which

contained both *Lmx1b*-positive and *Lmx1b*-negative regions (Fig. 2B), exhibited asymmetric

structures similar to those of normal limbs (Fig. 4H, I, Fig. S2A, B). To evaluate the symmetry of the

limbs, we applied a machine learning-based method using ilastik because staining intensity varied

among samples, such that a region identified as "muscle" in one sample could be assigned

differently in another if classification were based solely on color. The machine-learning classifier

trained separately for each sample allowed us to group the same tissues consistently within that

sample irrespective of intensity differences. To minimize the effects of curvature or fixation-induced

distortion, the boxes with a width of 400 μm were selected, and the angle was adjusted so that the

outer contour (epidermal surface) was aligned symmetrically; this procedure was applied uniformly

across all conditions to avoid bias. Each pixel in the images was classified into five classes (Classes

1 to 5, Fig. 4L). In this process, we annotated regions of background (Class 1), cartilage (Class 2),

muscle (Class 3), other connective tissue (Class 4), and epidermis (Class 5) as training data for

classification. Using these annotated regions as references, pixels were automatically classified into

the five classes. Each class was assumed to primarily represent these tissues and regions with similar

characteristics. Symmetry scores were then calculated for each class individually and for the

combined set of all classes (Fig. 4M–R, see Materials and Methods). The analysis revealed that

symmetry scores for Classes 2 and 3, which encompass cartilage and muscle, and scores for the

combined set of all classes were significantly higher in limbs formed from DorBL and VentBL

compared to intact limbs (Fig. 4N, O, R). In contrast, the differences in symmetry scores for Classes

1, 4, and 5 in these limbs were relatively small (Fig. 4M, P, Q), likely because the axis of symmetry

was manually set to maximize the external shape as symmetrically as possible. No significant

differences in symmetry scores across all classes were observed between intact limbs and limbs

formed from AntBL. These results suggest that limbs formed from DorBL and VentBL exhibit

symmetric internal structures compared to normal limbs. Therefore, *Shh* overexpression appears to

compensate for the lack of co-existence of dorsal and ventral cells in limb patterning, without

inducing new dorsal or ventral identities. Thus, we conclude that the co-existence of both dorsal and

ventral cells is critical for inducing *Shh* expression, which in turn is essential for limb patterning.

The molecular basis of the dorsal-mediated and ventral-mediated signals

To investigate the molecular basis of dorsal- and ventral-mediated signals, we performed RNA-seq

analysis on DorBL and VentBL and identified differentially expressed genes (DEGs) between the

two groups ($p < 0.05$, Fig. 5A). Among the DEGs, we specifically focused on genes annotated as

"intercellular signaling molecules" in PANTHER and identified 21 genes (Fig. 5B). In this analysis,

we found that 5 genes were expressed at higher levels in DorBL and 16 genes were expressed at

higher levels in VentBL (Fig. 5B). Notably, *Wnt4*, *Wnt10b*, *Fgf2*, *Fgf7*, and *Tgfb2*, which encode

secreted proteins, belong to the WNT, FGF, and TGFB families, which play key roles in major

signaling pathways regulating limb developmental processes and we focused on these 5 genes. To

examine whether these genes regulate the induction of *Shh* expression in ALM blastemas, we

overexpressed each gene in either DorBL or VentBL. As a result, *Shh* expression was observed in

Wnt10b-electroporated VentBL (n=4/5) and *Fgf2*-electroporated DorBL (n=5/7, Fig. 5C). In contrast,

Shh expression was not detected in *Wnt10b*-electroporated DorBL (n=6/6) or *Fgf2*-electroporated

VentBL (n=5/5). Similarly, *Shh* expression was not detected in *Fgf7*- or *Tgfb2*-electroporated DorBL

(n=5/5 for both) or in *Wnt4*-electroporated VentBL (n=6/6). To confirm the ISH data, we performed

RT-qPCR on *Fgf2*- or GFP (control)-electroporated DorBL and *Wnt10b*- or GFP (control)-

electroporated VentBL and observed significant upregulation in *Shh* expression (Fig. 5D, E). In

Wnt10b-electroporated VentBL, *Axin2*, a downstream transcriptional target of canonical WNT

signaling, and *Lef1*, a canonical WNT pathway effector expressed in axolotl limb mesenchyme

(Glotzer et al., 2022), were also upregulated (Fig. 5F, G). Next, to confirm *Shh* induction by WNT

signaling in VentBL, we treated VentBL with 1 μM 6-bromoindirubin-3-oxime (GSK-3 Inhibitor,

BIO). As a result, *Shh* expression was relatively higher in BIO-treated VentBL compared to DMSO-

treated control VentBL (Fig. S3B, D, E). Additionally, symmetric limbs were formed in *Wnt10b*-

electroporated VentBL (n=3/12), BIO-treated VentBL (n=3/8), and in *Fgf2*-electroporated DorBL

(n=5/10), consistent with the results of *Shh* overexpression (Fig. 5H–J, Fig. S3A, C, F–H, Fig. S4A,

B). These findings suggest that WNT10B, expressed highly in dorsal blastema cells, and FGF2,

expressed highly in ventral blastema cells, function as the dorsal- and ventral-mediated signals,

respectively, to induce *Shh* expression, which subsequently supports limb patterning.

To test whether our model applies to normal regeneration, we analyzed *Wnt10b* and *Fgf2*

expression in amputation-induced blastemas (Fig. S5). We first performed *in situ* hybridization on

blastemas at several stages (early bud [EB], middle bud [MB], and late bud [LB]), but the signals

were weak and inconsistent, and we could not reliably detect clear expression domains (Fig. S5A).

We then performed RT-qPCR on manually microdissected dorsal and ventral halves of MB

blastemas (Fig. S5B). We found that *Wnt10b* and *Fgf2* were expressed at significantly higher levels

in the dorsal and ventral halves, respectively, compared to the opposite half. This dorsoventral-

biased expression of *Wnt10b* and *Fgf2* is consistent with our RNA-seq data from ALM blastemas.

We next quantified *Wnt10b*, *Fgf2*, and *Shh* expression across stages (intact, EB, MB, LB, and early

digit [ED]) and found that *Wnt10b* and *Fgf2* expression peaked at the MB stage, whereas *Shh*

expression peaked later, at the LB stage (Fig. S5C). This temporal offset in *Shh* upregulation relative

to *Wnt10b* and *Fgf2* supports a model in which WNT10B and FGF2 act upstream to induce *Shh*

expression.

To identify the cell populations expressing *Wnt10b* and *Fgf2* during normal regeneration,

we reanalyzed published single-cell RNA-seq data from a 7 dpa (MB) blastema (Li et al., 2021, Fig.

S6). The dataset was reclustered, and clusters were assigned using known markers (*Prrx1* for

mesenchyme and *Krt17* for epithelium, Fig. S6A, B). As expected, *Lmx1b*, *Fgf8*, and *Shh* were

detected in the mesenchymal cluster (Fig. S6C, F, G). *Fgf2* was also expressed in the mesenchymal

cluster (Fig. S6E). In contrast, *Wnt10b* expression was detected in both mesenchymal and epithelial

clusters (Fig. S6D). Both *Wnt10b* and *Fgf2* were expressed in only a few cells, consistent with the *in*

situ hybridization data (Fig. S5A). We then examined the relationships between these genes. *Fgf8*

and *Shh* were expressed in both *Lmx1b*-positive and *Lmx1b*-negative cells (Fig. S6H, I), but *Fgf8*

and *Shh* themselves were mutually exclusive (Fig. S6M). These expression patterns of *Fgf8*, *Shh*,

and *Lmx1b* in the normal blastema are consistent with those observed in ALM blastemas (Fig. 2).

For *Wnt10b* and *Fgf2*, their expression did not follow *Lmx1b* expression (Fig. S6J, K), and *Wnt10b*

and *Fgf2* themselves were not exclusive (Fig. S6L). Together with the RT-qPCR data (Fig. S5B),

these results suggest that *Wnt10b* and *Fgf2* are not exclusively confined to purely dorsal or ventral

cells at the single-cell level, even though they show dorsoventral bias when assessed in bulk tissue.

Limb formation without nerve deviation and ventral skin grafting

A previous study demonstrated that a cocktail of BMP2, FGF2, and FGF8 can substitute for nerve

deviation in the ALM experiment on the anterior region, suggesting that nerves contribute to

blastema induction by supplying these proteins (Makanae et al., 2014). We found that FGF2 can also

substitute for ventral tissue in DorBL (Fig. 5), suggesting that FGF2 serves not only as part of the

nerve factors in blastema induction but also as the ventral-mediated signal in limb patterning. To

investigate whether supplementation with BMP2, FGF2, and FGF8 to the dorsal region could

substitute for both nerve deviation and ventral skin grafting, we performed experiments involving

gelatin beads soaked in these proteins (Fig. 6). The dorsal or ventral skin was removed at the

zeugopod level, and a BMP2+FGF2+FGF8-soaked gelatin bead was grafted at 3 dps without nerve

deviation. Blastema induction was observed in both experimental groups (Fig. 6A, B, upper panels).

We investigated gene expression patterns in the induced blastemas and found that both *Fgf8* and *Shh*

were expressed in blastemas induced at the dorsal site (n=8/8), whereas *Shh* expression was absent

in ventral blastemas (n=5/5), although *Fgf8* expression was detected in most cases (n=4/5, Fig. 6C,

D, F, G). *Lmx1b* expression patterns corresponded to those observed in DorBL and VentBL (Fig. 6E,

H, n=8/8 and 5/5, respectively). Furthermore, limb patterning was observed in the dorsal groups

(n=12/20) but not in the ventral groups (n=17/17, Fig. 6A, B, lower panels). These findings

demonstrate that a straightforward procedure involving dorsal skin wounding and supplementation

with BMP2, FGF2, and FGF8 is sufficient to induce accessory limb formation. This method enabled

the induction of an ectopic limb without surgical nerve deviation, but a previous study has shown

that fine nerve ingrowth can still occur when blastemas are induced by a BMP2+FGF2+FGF8-

soaked bead (Makanae et al., 2014). These recruited nerves may be functional after blastema

induction. Remarkably, this method also enabled the induction of multiple limbs within the same

limb. Grafting BMP2+FGF2+FGF8-soaked beads to multiple dorsal sites resulted in the formation

of multiple limbs along the proximodistal axis (n=16/35, Fig. 6I–L). Despite careful implantation

designed to avoid injuring deep tissues, one sample displayed a fusion of the stylopod with the host

humerus—a phenotype associated with deep wounding (Satoh et al., 2010; Makanae et al., 2014).

This suggests that contributions from a broader cellular population cannot be excluded. However,

because this fusion was not consistently observed, and because ectopic limbs induced at the forearm

(zeugopod) level did not exhibit such fusion (n=1/6 for stylopod-level inductions; n=0/10 for

zeugopod-level inductions, Fig. 6L1, L2), the data suggest that most ectopic limbs would have

developed without substantial ventral-cell contribution. These results refine our understanding of the

role of FGF2 in the ALM, particularly its critical function on the dorsal side.

Discussion

Dorsal- and ventral-mediated signals are required for the induction of Shh expression

In axolotl limb regeneration, cells derived from all four positional origins—anterior, posterior,

dorsal, and ventral—are required for an induced blastema to form a limb. This was confirmed

through the ALM experiments (Fig. 1). The gene expression patterns in these ALM blastemas

revealed distinct molecular characteristics for each group (Fig. 2). *Fgf8* expression in AntBL and *Shh*

expression in PostBL are consistent with a previous report (Nacu et al., 2016). The absence of SHH

in AntBL and of FGF8 in PostBL likely accounts for their failure to form limbs. *Lmx1b* expression in

the dorsal half of both AntBL and PostBL suggests that both dorsal and ventral tissues co-existed in

these blastemas (Fig. 2B, G, Fig. S1A, D). In contrast, DorBL and VentBL exhibited distinct *Lmx1b*

expression patterns, suggesting that most cells in DorBL and VentBL are derived from dorsal or

ventral origins, respectively (Fig. 2L, Q, Fig. S1G, J). A possible explanation for the lack of *Shh*

expression in DorBL and VentBL is that the induction of *Shh* expression may depend on the co-

existence of dorsal and ventral cells, although *Fgf8* can be expressed independently of the co-

existence of dorsal and ventral cells (Fig. 2M, N, R, S, U, V, Fig. S1H, I, K, L). This idea is

supported by the cell-tracing experiment (Fig. 3). In these assays, the posteriorly derived cells

expressed *Shh* only when both dorsal and ventral cells were present. These results strongly suggest

that the co-existence of dorsal and ventral cells is necessary for inducing *Shh* expression.

Consequently, one of the essential roles of the dorsal and ventral cells in limb patterning is to

mediate signals required for *Shh* induction in the posterior cells, which facilitates the anteroposterior

interaction.

Our findings clarify the hierarchical relationship between dorsal- and ventral-mediated

signals and anteroposterior interaction. Our data demonstrate that dorsal and ventral cells are

essential for inducing *Shh* expression in a regenerating blastema. Furthermore, we showed that

DorBL and VentBL, which typically fail to form a limb, could form patterned limbs with ectopic *Shh*

expression, even in the absence of dorsal and ventral cell co-existence (Fig. 4). Furthermore, such

limbs exhibited dorsally or ventrally symmetric structures (Fig. 4F, G, J–R, Fig. S2C, D),

highlighting that dorsoventral patterning depends on the cell origin. Thus, we concluded that dorsal-

and ventral-mediated signals are essential for patterned limb formation via inducing *Shh* expression.

WNT10B and FGF2 as dorsal- and ventral-mediated signals

We identified *Wnt10b* and *Fgf2* as candidate genes encoding the dorsal- and ventral-mediated signals

required to induce *Shh* expression, respectively (Fig. 5). Our DEG analysis between DorBL and

VentBL revealed higher *Wnt10b* expression in DorBL and higher *Fgf2* expression in VentBL (Fig.

5A, B). *Wnt10b*-electroporation in VentBL and *Fgf2*-electroporation in DorBL induced *Shh*

expression and subsequent limb patterning (Fig. 5C–E). *Wnt4* expression was also elevated in

DorBL, while *Fgf7* and *Tgfb2* were upregulated in VentBL. However, we did not observe *Shh*

induction in the *Fgf7*-electroporated or *Tgfb2*-electroporated DorBL or in the *Wnt4*-electroporated

VentBL. FGF7 is known to be capable of inducing the apical ectodermal ridge (AER) in chick limb

development (Yonei-Tamura et al., 1999), but little is known about its role in limb regeneration. In

newt limb regeneration, KGFR, which acts as an FGF7 receptor, is expressed in the basal layer of

the wound epithelium, while FGFR1, which acts as an FGF2 receptor, is primarily expressed in the

mesenchyme (Poulin et al., 1993). This differential expression pattern suggests that FGF2 and FGF7

target distinct cell populations, potentially explaining the differences in their ability to induce *Shh*

expression. The difference in the ability of *Wnt10b* and *Wnt4* to induce *Shh* expression in VentBL

may reflect differences in how these ligands activate downstream WNT signaling programs.

WNT10B is a potent activator of the canonical WNT signaling pathway (Bennett et al., 2005),

although WNT10B has also been reported to trigger a β-catenin–independent pathway (Lin et al.,

2021). Similarly, WNT4 can activate both the canonical WNT signaling pathways and a non-

canonical, β-catenin–independent pathway (Li et al., 2013; Li et al., 2019). We also observed *Shh*

induction in BIO-treated VentBL, indicating that the canonical WNT signaling pathway regulates

Shh expression. However, it is uncertain why *Shh* expression was observed in *Wnt10b*-electroporated

VentBL, but not in *Wnt4*-electroporated VentBL. One possible explanation is that different WNT

ligands can engage the same receptors (Frizzled/LRP6) yet elicit distinct downstream programs,

suggesting that such ligand-specific outputs may vary depending on cell context (Voss et al., 2025).

Regarding *Tgfb2*, TGF-β signaling has been implicated in blastema induction during salamander

limb regeneration (Sader and Roy, 2022; Lévesque et al., 2007). However, little is known about the

relationship between TGF-β signaling and *Shh* induction in limb development and regeneration,

suggesting that TGF-β signaling does not play a primary role in *Shh* induction. Consistent with this,

we did not observe *Shh* induction in the *Tgfb2*-electroporated DorBL, as described above. This

suggests that TGFB2 signaling may be involved in dorsoventral regulation independent of *Shh*

induction. Taken together, among the candidate genes we identified, WNT10B via the canonical

WNT pathway and FGF2 via FGFR1 appear to regulate *Shh* induction and the subsequent patterning

process.

WNT signaling may be the key for the dorsal properties in limb formation. It has been

reported that canonical WNT/β-catenin signaling plays essential roles in limb regeneration among

vertebrates, including axolotls (Kawakami et al., 2006; Lovely et al., 2022). In the present study, we

demonstrated that introducing *Wnt10b* or BIO treatment could induce *Shh* expression in VentBL,

facilitating limb patterning. In mouse limb development, *Wnt10b* is expressed in the apical

ectodermal ridge (AER, Witte et al., 2009), and mutations in *Wnt10b* are associated with Split-

Hand/Foot Malformation (Ugur and Tolun, 2008; Al Ghamdi et al., 2020; Bilal et al., 2023).

However, there is no evidence suggesting that *Wnt10b* contributes to dorsal specificity during limb

development. This raises questions about whether the function of *Wnt10b* is unique to axolotls or

whether its orthologs in other species might share functional redundancy with other WNT genes.

Further studies are needed to clarify this point. Notably, WNT10B utilizes the canonical Wnt

signaling pathway, similar to WNT7A, a well-known WNT family gene critical for dorsoventral

limb patterning in amniotes. In amniotes, *Wnt7a* is expressed in the dorsal ectoderm of developing

limb buds and induces *Lmx1b* expression in dorsal mesenchyme, thereby establishing dorsal

characteristics (Riddle et al., 1995; Cygan et al., 1997; Chen, 2002). In axolotls, dorsal-specific

Wnt7a expression has not been confirmed. Our RNA-seq analysis showed that there was almost no

significant difference in *Wnt7a* expression levels between DorBL and VentBL ($\log_2$ (normalized

counts) = 5.97, $\log_2$ (fold change) = -0.810, p = 0.401), consistent with previous studies (Shimokawa

et al., 2013). Similarly, there was no significant difference in *En1* expression ($\log_2$ (normalized

counts) = 0.909, $\log_2$ (fold change) = -1.812, p = 0.264). In amniote limb development, *En1* is

expressed in the ventral ectoderm, where it restricts *Wnt7a* expression to the dorsal ectoderm and

thereby prevents induction of *Lmx1b* in the ventral mesenchyme (Loomis et al., 1996; Logan et al.,

1997; Chen and Johnson, 2002). These results suggest that *Wnt7a* does not have a dorsal-specific

function, at least in axolotl limb regeneration. While WNT10B could function as a dorsal-mediated

signal, WNT10B is unlikely to induce dorsal identity, as ectopic *Lmx1b* expression was not observed

in *Wnt10b*-introduced VentBL, which formed double-ventral limbs (Fig. 5F–H). This indicates that

Wnt10b in axolotl limb regeneration does not simply replace the function of *Wnt7a* in amniote limb

development. Nevertheless, the involvement of canonical WNT signaling in dorsal function and limb

morphogenesis remains an important area for further investigation.

We identified FGF2 as the ventral-mediated signal, which plays a crucial role in the limb

patterning process during axolotl limb regeneration. This aligns with previous studies showing that

Fgf2 expression correlates with limb regeneration in salamanders (Giampaoli et al., 2003). *Fgf2*

expression was relatively high in VentBL, and *Fgf2*-introduced DorBL formed patterned limbs (Fig.

5). These results indicate that *Fgf2* overexpression can substitute for the presence of ventral cells in

the limb patterning process. It is noteworthy that FGF2 application does not appear to induce ventral

identity, as the limbs formed from *Fgf2*-introduced DorBL exhibited dorsally symmetric limb

structures. The use of FGF signaling as a ventral-mediated output downstream of dorsoventral

identity has not been documented in other species examined to date. Whether this represents an

axolotl-specific regulatory mechanism or a broader function of FGF signaling in limb

morphogenesis remains to be determined.

Our findings suggest that although WNT10B and FGF2 act as dorsal- and ventral-

mediated signals, they do not alter dorsal or ventral identity itself. In amniote limb development,

WNT7A and EN1 regulate dorsoventral identity through *Lmx1b* expression. In contrast, in axolotls,

Wnt10b-electroporation in VentBL or *Fgf2*-electroporation in DorBL did not affect the expression

patterns of *Lmx1b* (Fig. 5C). Moreover, the limbs formed from such DorBL and VentBL exhibited

dorsally or ventrally symmetric structures (Fig. 5H–J, Fig. S4). These findings suggest that

dorsoventral identities in axolotls are not affected by *Wnt10b* or *Fgf2* overexpression. We previously

reported that the expression patterns of *Lmx1b* in axolotl limb regeneration are likely to depend on

the positional origins of cells (Iwata et al., 2020; Yamamoto et al., 2022). The identities of cells

along the dorsoventral axis may be controlled by their positional memory and determined before the

initiation of limb regeneration.

The present study revealed that the dorsal- and ventral-mediated signals WNT10B and

FGF2 regulate *Shh* expression during limb patterning in axolotls. These findings highlight a degree

of conservation between axolotl limb regeneration and amniote limb development. In amniote limb

development, FGF and WNT signaling pathways are key regulators of *Shh* expression. In developing

amniote limb buds, *Fgf2* is expressed in the ectoderm, including the AER, and in adjacent

mesoderm, and promotes distal outgrowth. FGFs, including FGF2, supplied from the AER are

known to maintain *Shh* expression (Laufer et al., 1994; Niswander et al., 1994; Yang and Niswander,

1995; Li et al., 1996). Similarly, WNT7A regulates *Shh* expression in amniote limb development,

and loss of WNT7A function results in reduced *Shh* expression in the zone of polarizing activity

(ZPA) and the deletion of posterior structures (Parr and McMahon, 1995). Thus, our findings provide

new insight into both conserved and divergent aspects of dorsal- and ventral-mediated signaling in

the regulation of *Shh* expression, thereby furthering our understanding of limb morphogenesis.

In this study, RNA-seq analysis of ALM blastemas induced on the dorsal or ventral side

listed *Wnt10b* and *Fgf2* as genes that are more highly expressed in DorBL and VentBL, respectively.

In the ALM context, *Wnt10b* or *Fgf2* overexpression was sufficient to substitute for dorsal or ventral

tissues, respectively, and to drive *Shh* induction and subsequent limb patterning even in the absence

of those tissues (Fig. 5). However, in amputation-induced blastemas during normal regeneration, *in*

situ hybridization did not reveal clear expression patterns for *Wnt10b* or *Fgf2* (Fig. S5A), and

reanalysis of single-cell RNA-seq from the regenerating blastema (Li et al., 2021) showed that their

expression did not strictly follow *Lmx1b* expression (Fig. S6J, K). By contrast, and consistent with

our bulk RNA-seq results, RT-qPCR of manually microdissected dorsal and ventral halves of

regenerating blastemas showed that *Wnt10b* and *Fgf2* were expressed at significantly higher levels in

the dorsal and ventral halves, respectively (Fig. S5B). These results suggest that *Wnt10b* expression

and *Fgf2* expression are mediated by dorsal and ventral cells, respectively, but their expression is not

restricted to dorsal or ventral cells. Our results on *Wnt10b*-electroporated VentBL and *Fgf2*-

electroporated DorBL suggest that both activation of WNT10B or FGF2 is required for *Shh*

induction and proceed with limb patterning. To fully understand axolotl limb regeneration, it will be

important to determine how *Wnt10b* and *Fgf2* expression is regulated, how their downstream

programs are deployed, and how dorsal and ventral cells respond to these signals in future studies.

The dorsoventral-mediated Shh induction mechanism

In the present study, we found the importance of the co-presence of cells carrying the

anterior, posterior, dorsal, and ventral identity for successful axolotl limb patterning (Fig. 7). In our

proposed model, following amputation, nerves first trigger blastema induction by secreting nerve

factors, such as BMPs and FGFs (Makanae et al., 2014, Satoh et al., 2016). These nerve factors

stimulate connective tissue cells, including dermal cells, to generate multipotent mesenchymal

blastemal cells (Muneoka et al., 1986; Kragl et al., 2009; Hirata et al., 2010; Gerber et al., 2018).

Within the induced blastema, *Wnt10b* and *Fgf2* expression are mediated by the dorsal and ventral

cells, respectively. In the next phase, the co-existence of WNT10B and FGF2 signaling induces *Shh*

expression in the posterior region of the blastema. During this phase, *Fgf8* is expressed in the

anterior mesenchyme independently of the co-existence of dorsal and ventral cells (Fig. 7A). These

expression patterns of *Wnt10b*, *Fgf2*, *Shh*, and *Fgf8* may be mediated by the positional memory of

cells (Otsuki et al., 2021). In the subsequent phase, the anteroposterior interaction, mediated by

FGF8 and SHH, supports distal outgrowth to form a limb. This model explains previous

observations in studies on double-half limbs and ALM blastemas (AntBL, PostBL, DorBL, and

VentBL), which typically fail to form a limb. In blastemas induced from double-anterior and double-

posterior limbs, or AntBL (Fig. 7B) and PostBL (Fig. 7C), SHH or FGF8 proteins are likely absent

due to the lack of posteriorly or anteriorly derived cells, respectively. Similarly, in blastemas induced

by amputating double-dorsal and double-ventral limbs, or DorBL (Fig. 7D) and VentBL (Fig. 7E),

FGF2 or WNT10B proteins are likely absent or insufficient because of the lack of ventrally or

dorsally derived cells, respectively. This results in the absence of SHH, even if posteriorly derived

cells are present. In all these cases, the absence of either FGF8 or SHH disrupts the anteroposterior

interaction, leading to failure in limb patterning. We conclude that the requirement for cells derived

from all four positional origins is underpinned by this model. In this model, dorsal- and ventral-

mediated signals activate the posterior SHH, enabling mutual interaction with the anterior FGF8.

This interplay ensures proper anteroposterior interaction and complete limb patterning. Our findings

contribute to understanding how the integration of four positional identities—dorsal, ventral,

anterior, and posterior—drives proper limb patterning during axolotl limb regeneration.

Materials and Methods

Animal procedures

Axolotls between 4 and 15 cm from snout to tail tip were housed in tap water at 22°C. Both forelimbs and hind limbs were used for surgical procedures without distinction. Transgenic axolotls were obtained from the Ambystoma Genetic Stock Center (http://www.ambystoma.org/genetic-stock-center). Animals were anesthetized in 0.1% MS222 (Sigma-Aldrich) solution before surgical procedures.

In the ALM experiments, skin was peeled from the anterior, posterior, dorsal, or ventral side of a limb at the stylopod level, so that the skin of the opposite side was not injured. Thus, the size of the injured area depended on the limb size. Then, thick bundles of nerve trunks running the center of a limb was deviated the injured region. For the cell-tracing experiment, a piece of posterior half of the dorsal or ventral skin was obtained from GFP-expressing transgenic animals and grafted to the ALM region.

Protein-soaked beads for grafting were prepared as previously described (Makanae et al., 2014; Kashimoto et al., 2023). In brief, air-dried gelatin beads were allowed to swell in stock solution (1 μg/μL) prepared following the manufacturer's instructions. Equal amounts of proteins were used when formulating the combination protein mixture. Beads were soaked in the mixture of proteins (Bmp2 [mouse], Fgf2 [mouse], and Fgf8 [human/mouse]; R&D Systems) overnight at 4°C. For control experiments, gelatin beads were soaked in DDW. Before grafting, dorsal or ventral skin was peeled because full-thickness skin inhibits limb regeneration (Thornton, 1962; Mescher, 1976; Tsai, 2020). The beads were grafted under the wounded epidermis at 3 dps. The limbs were fixed 10 days after grafting. The details of grafting procedures are as previously described (Makanae et al., 2014; Kashimoto et al., 2023).

To analyze skeletal patterns, induced limbs were stained with alcian blue and alizarin red. Samples were fixed in 100% ethanol for 1 day at room temperature, then stained with Alcian blue solution (Wako, pH 2.0) in 20% acetic acid (Nacalai Tesque) with 80% ethanol solution at 37°C overnight. Then, samples were washed in tap water several times and fixed in 10% Formaldehyde Neutral Buffer Solution (Nacalai Tesque) for 1 day. Samples were then stained with Alizarin red S (Nacalai Tesque) in the solution (4% KOH:10% Formaldehyde Neutral Buffer Solution = 2:3) at room temperature for 1 day. Finally, samples were placed in graded glycerol with 4% KOH for clearing.

For BIO treatment, 10 mM stock solution of 6-bromoindirubin-3-oxime (BIO, Selleck, S7198) dissolved in DMSO was stored in the dark at 4°C. Axolotls soon after surgery were raised in tap water with BIO solution (experimental) or with the same amount of solvent, DMSO (control), until 14 dps for RT-qPCR and 30 dps for phenotype analysis. The containers including water and axolotls were kept in the dark during BIO or DMSO treatment. For BIO treatment, axolotls between

568 4 and 5 cm from snout to tail tip were used.

Sectioning and histological staining

571 Samples were fixed in 4% PFA/PBS overnight at room temperature before sectioning. The fixed

572 samples were embedded in O.C.T. compound (Sakura) following 30% sucrose/PBS treatment for

573 approximately 12 h at 4℃. Frozen sections of 14 μm thickness were prepared using Leica CM1850

574 (Nussloch). The sections were well dried under an air dryer and kept at -80℃ until use.

575   Standard hematoxylin and eosin (HE) staining and Masson's trichrome staining were used

576 for histological analysis. To visualize cartilage formation, Alcian blue staining was performed before

577 HE staining. In brief, sections were washed in tap water several times to remove the O.C.T.

578 compound. Then, the sections were stained with Alcian blue (Wako, pH 2.0), and then HE staining

579 was performed. Trichrome stain (Masson) kit (Sigma, HT15-1KT) was used for trichrome staining.

580 The stained sections were mounted using Softmount (Wako).

In situ hybridization

583 *In situ* hybridization was performed as described previously (Yamamoto et al., 2022). For probe

584 synthesis, target genes cloned on pTAC-2 plasmid (BioDynamics Lab Inc.) were amplified by PCR

585 using M13 primers. PCR fragments were purified and used as an RNA probe template. RNA probe

586 synthesis was performed with Sp6 or T7 RNA polymerase (Takara) for 3 h, and RNA was

587 hydrolyzed, depending on the length of targets. The sections were washed in PBT to remove the

588 O.C.T. compound, treated with proteinase K (10 μg/mL) (Invitrogen)/PBT for 20 min at room

589 temperature, washed in PBT, treated with 4% PFA/PBS for 20 min at room temperature, washed in

590 PBT, and then probes were hybridized at 62.5℃ for approximately 18 h. The sections were washed

591 in wash buffer 1 (formamide:H_2O:20× SSC [3M NaCl:0.3M sodium citrate, pH 5.0] = 2:1:1), and

592 then in wash buffer 2 (formamide:H_2O:20× SSC = 5:1:4). The samples were then incubated with

593 anti-digoxigenin-AP Fab fragments (Sigma-Aldrich, 1/1000) for 2 h at room temperature. Samples

594 were stained with BCIP (Nacalai Tesque) and NBT (Nacalai Tesque) in alkaline phosphatase buffer

595 (0.1M NaCl, 0.1M Tris-HCl [pH9.5], 0.1% Tween20) for 18 h at room temperature after washing in

596 TBST.

Immunofluorescence

599 Immunofluorescence on sections was carried out based on a previous report (Yamamoto et al.,

600 2022). The antibodies were as follows: anti-GFP (MBL, #598, 1/500), anti-acetylated alpha tubulin

601 (Santa Cruz, #sc-23950, 1/1000), anti-mouse IgG Alexa 488 (Invitrogen, #A11017, 1/1000), anti-

602 rabbit IgG Alexa 488 (Invitrogen, #A21206, 1/500). Nuclei were stained with Hoechst (Nacalai

Tesque), and images were captured using an Olympus BX51 system.

RNA-Seq analysis

Total RNA was extracted from DorBL and VentBL at 10 dps using TriPure reagent (Roche),

following the manufacturer's instructions. Three biological replicates were prepared for both

samples. 150 bp paired-end RNA-seq reads were obtained under contract to Rhelixa (Tokyo, Japan),

using Illumina Nova Seq 6000 and SMART-Seq HT Plus Kit (#R400449). RNA-seq data were

analyzed on Galaxy (https://usegalaxy.eu/root) as follows. Sequence reads were trimmed and the

quality was filtered by Trimmomatic v0.39 (Bolger et al., 2014) with the following parameters

(LEADING:20, TRAILING:20, SLIDINGWINDOW:4:15, MINLEN:36). Axolotl genome data

(AmexG_v6.0-DD) and annotation data (AmexT_v47-AmexG_v6.0-DD.gtf) were obtained from

AXOLOTL-OMICS (https://www.axolotl-omics.org/) (Schloissnig et al., 2021). Mapping to the

Axolotl genome (AmexG_v6.0-DD) was performed with HISAT2 v2.2.1 (Kim et al., 2015). Count

data were obtained with featureCounts v2.0.8 (Liao et al., 2013) on the basis of the Axolotl gene

model (AmexT_v47-AmexG_v6.0-DD.gtf). DEGs were identified with DESeq2 v2.11.40.8 (Love et

al., 2014) ($p < 0.05$). Among identified DEGs (DorBL > VentBL; 762 genes, VentBL > DorBL; 513

genes), genes annotated as "intercellular signaling molecule" were explored on PANTHER

(https://www.pantherdb.org/) (Thomas et al., 2003), and 21 genes (DorBL > VentBL; 5 genes,

VentBL > DorBL; 16 genes) were identified as candidate genes of dorsal- and ventral-mediated

signals, as shown below:

DorBL > VentBL:

WNT4 (AMEX60DD052091), WNT10B (AMEX60DD029981), SEMA3A (AMEX60DD023165),

LECT2 (AMEX60DD041044), ANGPTL5 (AMEX60DD049512).

VentBL > DorBL:

FCN1 (AMEX60DD050926), CXCL14 (AMEX60DD028973), DNER (AMEX60DD002010),

FGF7 (AMEX60DD003767), CXCL12 (AMEX60DD052412), CXCL8 (AMEX60DD044133),

SEMA6A (AMEX60DD042813), ANGPTL2 (AMEX60DD050490), ANGPTL6

(AMEX60DD031854), FGL2 (AMEX60DD006387), VWDE (AMEX60DD022491), TGFB2

(AMEX60DD036126), FGF2 (AMEX60DD044865), EFNA4 (AMEX60DD014882), EFNA3

(AMEX60DD014867), ANGPTL1 (AMEX60DD018261).

Cloning candidate genes and preparation of construct vectors for overexpression

Among the identified candidate genes, we focused on *Wnt10b*, *Wnt4*, *Fgf2*, *Fgf7*, and *Tgfb2*. We

cloned these genes in pTAC-2 plasmids from cDNA obtained from blastemas. The following primers

were used:

Wnt10b Fwd:ATGGCCCACAGCTCACCCTCCGACACC

Wnt10b Rev:TCACTTGCACACATTCACCCATTCTGTG

Wnt4 Fwd:ATGGATGCTCACGAAAGCAGCGTATATC

Wnt4 Rev:TCACCGGCAGGTGTGCATTTCTACCAC

Fgf2 Fwd:ATGGCGGCGGGGAGCATCACCACCTTGC

Fgf2 Rev:TCAACTCTTGGCCGACATGGGAAGGAAAG

Fgf7 Fwd:ATGCGCAGATGGGTGCTAGCTTGGATC

Fgf7 Rev:TCATGTGTTATTGGATATACGCATTGGA

Tgfb2 Fwd:ATGAGATTACAATTACTGAGAAAAAAAATG

Tgfb2 Rev:TTAGCTGCACTTGCAAGATTTTACAATCA

We then inserted these genes to p2a-AcGFP-pCS2 vectors with In-Fusion® HD Cloning Kit

(Clontech). The following primers were used for PCR before In-Fusion reaction:

pCS2 inverse Fwd:GCTACTAACTTCAGCCTGCTGAAGCAGG

pCS2 inverse Rev:CATCGATGGGATCCTGCAAAAAGAACAAGTAGCTT

Wnt10b Fwd:AGGATCCCATCGATGGCCCACAGCTCACCCTCCGA

Wnt10b Rev:GCTGAAGTTAGTAGCCTTGCACACATTCACCCA

Wnt4 Fwd:AGGATCCCATCGATGGATGCTCACGAAAGCAGCG

Wnt4 Rev:GCTGAAGTTAGTAGCCCGGCAGGTGTGCATTTCTA

Fgf2 Fwd:AGGATCCCATCGATGGCGGCGGGGAGCATCACCA

Fgf2 Rev:GCTGAAGTTAGTAGCACTCTTGGCCGACATGGGAA

Fgf7 Fwd:AGGATCCCATCGATGCGCAGATGGGTGCTAGCTTG

Fgf7 Rev:GCTGAAGTTAGTAGCTGTGTTATTGGATATACGCA

Tgfb2 Fwd:AGGATCCCATCGATGAGATTACAATTACTGAGAAA

Tgfb2 Rev:GCTGAAGTTAGTAGCGCTGCACTTGCAAGATTTTA

Finally, the following DNA constructs were obtained; *Wnt10b*-p2a-GFP (pCS2), *Wnt4*-p2a-GFP

(pCS2), *Fgf2*-p2a-GFP (pCS2), *Fgf7*-p2a-GFP (pCS2), and *Tgfb2*-p2a-GFP (pCS2).

Electroporation

Each DNA construct was injected directly into the target region. Immediately after injection, electric

pulses were applied (20 V, 50 ms pulse length, 950 ms interval, 10 times) with NEPA21(Nepa gene).

The injected DNA constructs were as follows: pCS2-AcGFP, pCS2-*Shh*-p2a-AcGFP, *Wnt10b*-p2a-

GFP (pCS2), *Wnt4*-p2a-GFP (pCS2), *Fgf2*-p2a-GFP (pCS2), *Fgf7*-p2a-GFP (pCS2), and *Tgfb2*-p2a-

GFP (pCS2). All plasmids were purified using a Genopure Maxi kit (Roche). Electroporation was

performed at -3, 4, and 7 dps and samples were fixed at 10 dps for *in situ* hybridization. To analyze

the phenotype of 90 dps samples, electroporation was performed at -3, 4, 7, 10, 15, 20, 25, and 30

dps.

qRT-PCR

The procedures of qRT-PCR were previously described (Yamamoto et al., 2022). In brief, RT was

performed using Prime Script II (Takara), and RT-qPCR was performed using KAPA SYBR FAST

qPCR Master Mix (Kapa Biosystems) and StepOne (ThermoFisher Scientific). Primers were as

follows:

Ef-1α Fwd: AACATCGTGGTCATCGGCCAT

Ef-1α Rev: GGAGGTGCCAGTGATCATGTT

Shh Fwd: GCTCTGTGAAAGCAGAGAACTCG

Shh Rev: CGCTCCGTCTCTATCACGTAGAA

Axin2 Fwd: GGCACTGACTTATCCCCAGG

Axin2 Rev: GCATCATTGGCTGTCAACGG

Lef1 Fwd: CTACACCGAGATCAGCCACC

Lef1 Rev: GCTGTGGTAGGAGTTGTGGG

Lmx1b Fwd: CTGGTCCATGGCTACGATCT

Lmx1b Rev: TTAGCAGCAGAAACGGGACT

Wnt10b Fwd: CAGAAGAGACCCAGGTGCAG

Wnt10b Rev: CGAAGGCCCAAGATGTCTGT

Fgf2 Fwd: TCTTCCTTCGCATCAACCCC

Fgf2 Rev: TTTCATTGCCATCAACCGCC

RNA for Fig. S3E was prepared from 1μM BIO- or DMSO-treated VentBL at 14 dps. *Ef-1α* was

used as the internal control.

Pixel classification and calculating symmetry scores

A machine learning-based method was applied for pixel classification using ilastik software

(downloaded on https://www.ilastik.org/). The details and workflows were previously described

(Berg et al., 2019). Each pixel in the images was classified into five classes. The regions of the

background (Class 1), cartilage (Class 2), muscle (Class 3), other connective tissue (Class 4), and

epidermis (Class 5) were annotated for training data. Using these annotated regions as references,

pixels were automatically classified into the respective classes. In this process, the same training data

were used for images obtained from the same limb. Then, symmetry scores were calculated for each

class individually and for the combined set of all classes. In this process, the external shape of the

section, which could bend randomly during sample fixation process, would affect the symmetry

scores if the entire region were used. To focus on the symmetry of the anatomical patterns, images

with 400 μm width were prepared (Fig. 4L, 5E, F). The symmetry scores of pixel-classified images

were calculated using python. First, the center of the dorsal end and ventral end was set as the axis of

symmetry. Then, one side of the image was flipped. Next, color masks were generated for both sides

of the image. These masks identified pixels that matched the specified color. Pixels were considered

to match if their RGB values were within the given tolerance range for all three channels. In pixel

comparison, the masks for one side and the other side were compared to calculate the following

pixels:

Matching pixels: The number of pixels that match in both sides for the specified color.

Total pixels: The total number of pixels matching the color in either side.

The symmetry scores were computed as (Number of Matching Pixels)/(Total Pixels in Both Sides).

The scores were obtained from 12 areas of each group. For statistical analysis, a two-tailed a two-

tailed Welch's *t*-test was used. In this analysis, each group was compared to the intact limb group

because the intact limb should be set as a typical asymmetrical structure.

Reanalyzing single-cell data

We reanalyzed published single-cell RNA-seq data from a 7 dpa (MB) blastema (Li et al., 2021, Fig.

S6). We constructed a transcriptome index for axolotl from the available genome assembly and gene

annotation (AmexG_v6.0-DD genome and AmexT_v47-AmexG_v6.0-DD annotation, Schloissnig et

al., 2021). The transcript set was used to build a salmon index with default k-mer length (31), and

without decoy sequences, and gene-level UMI counts were generated using the Alevin module of

salmon, which performs lightweight (quasi-)mapping of reads directly to the transcriptome index

(Patro et al., 2017; Srivastava et al., 2019). The mapping was done on Galaxy (https://usegalaxy.eu).

This procedure yielded a whitelist of 8,925 barcodes, corresponding to putative cells, and produced a

per-cell by per-gene UMI count matrix. According to the Alevin summary output, 77.4% of reads

aligned to the indexed axolotl transcriptome after barcode correction and UMI deduplication. The

resulting count matrix was exported in Matrix Exchange (MEX) format (matrix.mtx, barcodes.tsv,

features.tsv).

The data were then imported into R (RStudio environment) as a Seurat object, a data
structure for scRNA seq data (Hao et al., 2021). Cells were filtered based on standard quality-control
metrics, excluding droplets with extremely low gene complexity, extremely high total UMI counts,
or unusually high mitochondrial or ribosomal RNA content. For normalization and integration of
cells into a shared space, counts were log-normalized using Seurat's NormalizeData (default
LogNormalize method with a scale factor of 10,000), and highly variable genes (HVGs) were
identified using FindVariableFeatures (vst method; 2,000 features). The data were scaled with
ScaleData, and principal component analysis (PCA) was performed on the HVGs (RunPCA, 50
components). The first 30 principal components were used to construct a nearest-neighbor graph
(FindNeighbors) and to perform community detection–based clustering (FindClusters, resolution =
0.3). The same set of PCs (dims 1:30) was also used for nonlinear dimensionality reduction by
UMAP (RunUMAP). Gene expression patterns were visualized using FeaturePlot with order =
TRUE so that high-expressing cells are drawn on top of low- or non-expressing cells. We also
generated two-gene "co-expression" maps by classifying cells as expressing gene A only, gene B
only, both, or neither, and overlaying these classes on the UMAP using DimPlot.

Author contributions
Conceptualization: S.Y., A.S.; Methodology: S.Y., M.H., A.S; Validation: S.Y.; Formal analysis: S.Y.;
Investigation: S.Y.; Resources: S.Y., S.F., A.O., A.S.; Data curation: S.Y.; Writing - original draft:
S.Y.; Writing - review & editing: S.Y., S.F., A.O., M.H., A.S.; Visualization: S.Y., A.S.; Supervision:
A.S.; Project administration: A.S.; Funding acquisition: A.S., S.Y.

Data availability
RNA-seq FASTQ files have been deposited in the DNA Data Bank of Japan (DDBJ;
https://www.ddbj.nig.ac.jp/) under BioProject accession PRJDB38065.

Acknowledgments
We are grateful to R. Iwata and T. Satoh for supporting office work and animal housing. Animals
were obtained through Hiroshima University Amphibian Research Center. This work is supported
by the Japan Society for the Promotion of Science KAKENHI grant-in-aid for scientific research (B)
(20H03264 and 24K02034 to A.S.) and by a Grant-in-Aid for Japan Society for the Promotion of
Science fellows (24KJ1718 to S.Y.).

Figure Legends

Figure 1. ALM experiments at the four orientations.

(A) Schematic image of anatomy at the stylopod level of axolotl limb. HE staining (bright field) and acetylated alpha tubulin, visualized by immunofluorescence (green, dark field), are shown in the right panels. Black and white arrows, respectively, indicate major blood vessels and nerves. H: *humerus*, AHL: *Anconaeus humeralis lateralis*, HAB: *Humeroantebrachialis*, CBL: *Coracobrachialis longus*. (B–I) Blastemas induced at the anterior, posterior, dorsal, or ventral region by skin wounding plus nerve deviation without (B–E) or with (F–I) skin grafting from the opposite side of the limb. (F–I) Limb patterning was observed (n=8/9 for F, 4/7 for G, 7/14 for H, and 7/11 for I, see Table 1 for more detail). Images were captured at 10 and 60 dps. Scale bar = 3 mm.

Figure 2. Gene expression patterns of the ALM-induced blastemas.

Sections of anteriorly (A–D), posteriorly (F–I), dorsally (K–N), or ventrally (P–S) induced blastemas at 10 dps. Acetylated alpha tubulin (A–P) was visualized by immunofluorescence. Expression of *Lmx1b* (B–Q), *Fgf8* (C–R), and *Shh* (D–S) in the regions indicated by white boxes in (A–P) was visualized by *in situ* hybridization. Images of the entire blastema are provided in Fig. S1. *Fgf8* expression was observed in AntBL (C), DorBL (M), and VentBL (R) (n=4/5, 4/6, and 4/6, respectively. *Shh* expression was observed in PostBL (I) (n=5/5). In each case, these expression patterns of *Fgf8* and *Shh* were focal and only a few cells expressed *Fgf8* or *Shh*. Black and white arrowheads indicate the signals of *Fgf8* and *Shh* expression, respectively. The dotted line indicates the epithelial-mesenchyme border. (E–T) Schematic images of gene expression patterns. (U, V) Quantitative analysis of *Fgf8* and *Shh* expression in ALM blastemas (n=5 for all groups). n.s.; no significant difference, *$p<0.05$, **$p<0.005$ (two-tailed Welch's t-test). Scale bar in (A) = 700 μm.

Figure 3. Co-existence of dorsal and ventral cells induces Shh expression.

(A) Experimental scheme. Posterior half of dorsal (PD_{gfp}) or ventral (PV_{gfp}) GFP-expressing skin was grafted on VentBL (VentBL+PD_{gfp}; C, D / VentBL+PV_{gfp}; G, H), or DorBL (DorBL+PD_{gfp}; E, F / DorBL+PV_{gfp}; I, J) region. (B) Induced blastemas at 10 dps; images of bright and dark fields are merged. (C–J) Dark and bright fields of the same sections of induced blastemas at 10 dps. Red boxes in (C–I) indicate the corresponding regions of lower images. Expression of *Shh* and *Lmx1b* were visualized by *in situ* hybridization. GFP signals were visualized by immunofluorescence. Arrowheads indicate the cells expressing *Shh*. *Shh* expression was observed in VentBL+PD_{gfp} (C) and DorBL+PV_{gfp} (I) (n=4/6 and 3/6, respectively), but not in DorBL+PD_{gfp} (E) and VentBL+PV_{gfp} (G) (n=6/6 and 7/7, respectively). The dotted line indicates the epithelial-mesenchyme border. For all samples, we collected serial sections spanning the entire blastema. For blastemas in which *Shh* expression was observed, we present representative sections showing the signal. For blastemas

without detectable *Shh* expression, we present a section from the central region that contains GFP-

positive cells. Scale bar = 3 mm (B) and 700 μm (C).

Figure 4. Limb formation without the co-existence of dorsal and ventral cells by Shh

overexpresion.

(A) Experimental scheme. *Shh*-p2A-GFP- or GFP-containing pCS2 vector was electroporated (EP)

into AntBL, DorBL, and VentBL. (B) Induced blastemas at 14 dps; images of bright and dark

fields are merged. (C–G) Phenotypes at 90 dps. Limb patterning was observed in (E–G) (n=10/19,

6/18, and 8/14, respectively). (H–K) Histological analysis of intact and induced limbs. Standard

Masson's trichrome staining was performed on the transverse sections. The dotted boxes indicate the

regions shown in (L). (L) Upper panels: analyzed regions for calculating symmetry scores. Lower

panels: images after pixel classification by machine learning. (M–R) Symmetry scores of each class.

Scores obtained from the same limb are plotted at the same x-coordinates. n.s.; no significant

difference, *$p<0.05$, **$p<0.005$ (two-tailed Welch's *t*-test). Scale bar = 2 mm (B), 4 mm (C–G), and

1 mm (H–K).

Figure 5. Identification of candidate molecules of the dorsal- and ventral-mediated signals.

RNA-Seq was performed on DorBL and VentBL at 10 dps. (A) MA plot of the result. (B) List of

DEGs annotated as "intercellular signaling molecules." (C) Bright and dark fields of sections of

DorBL and VentBL at 10 dps with candidate genes introduced. Expression of *Shh* and *Lmx1b* was

visualized by *in situ* hybridization, and GFP signals were visualized by immunofluorescence.

Arrowheads indicate the cells expressing *Shh*. The white boxes indicate the regions of the lower

panels. Images of dark and bright fields of *Shh* are obtained from the same section. For samples with

detectable *Shh* expression, the window was placed in the region where the signal was observed, and

for conditions without detectable *Shh* expression, the window was positioned in a comparable region

containing GFP-positive cells. *Shh* expression was observed in *Wnt10b*-electroporated VentBL

(n=4/5) and *Fgf2*-electroporated DorBL (n=5/7), but not in *Wnt10b*-electroporated DorBL (n=6/6),

Fgf2-electroporated VentBL (n=5/5), *Fgf7*-electroporated DorBL (n=5/5), *Tgfb2*-electroporated

DorBL (n=5/5), or in *Wnt4*-electroporated VentBL (n=6/6). (D, E) Quantitative analysis of *Shh*

expression in *Fgf2*- or GFP-electroporated DorBL and *Wnt10b*- or GFP-electroporated VentBL (n=7

for both). In each case, *Fgf2* or *Wnt10b* was electroporated into the DorBL or VentBL induced in the

left limb, and GFP was electroporated into the contralateral right limb of the same animal. (F, G)

Quantitative analysis of *Axin2* and *Lef1* expression in *Wnt10b*- or GFP-electroporated VentBL (n=7

for both). (H, I) The limbs formed from VentBL with *Wnt10b* and DorBL with *Fgf2* at 90 dps.

Histological analysis and pixel classification was performed in the same way as in Fig. 4. The dotted

boxes indicate the regions shown in the right panels. (J) Symmetry scores of each class. Scores

obtained from the same limb are plotted at the same x-coordinates. The plots of "int" are the same

plots as in Fig. 4. n.s.; no significant difference, $*p<0.05$, $**p<0.005$ (two-tailed paired t-test for D–

G, and two-tailed Welch's t-test for J). Scale bar = 700 μm (C), 4 mm (H, I, upper panels), 1 mm (H,

I, lower panels).

Figure 6. Limb formation at the dorsal region by BMP2+FGF2+FGF8 supplementation.

(A, B) 14 and 60 dps phenotypes of BMP2+FGF2+FGF8 supplementation by bead grafting at the

dorsal (A) or ventral (B) region. Neither nerve deviation nor skin grafting was performed. Limb

patterning was observed in the dorsa group (A) (n=12/20) but not in the ventral group (B) (n = 0/17).

(C–H) Expression patterns of *Fgf8*, *Shh* and *Lmx1b* of induced blastemas at 10 dps. Gene expression

was visualized by *in situ* hybridization. The dotted line indicates the external shape of the blastema.

Fgf8 expression was detected in both dorsal and ventral groups (n=8/8 for C and 5/5 for F), whereas

Shh expression was detected only in the dorsal group (D) (n=8/8) and not in the ventral group (G)

(n= 0/5). (I–J) Phenotype obtained by BMP2+FGF2+FGF8 supplementation to two dorsal regions of

an identical limb. (K, L) Phenotype obtained by BMP2+FGF2+FGF8 supplementation to five dorsal

regions of an identical limb (n=16/35). Scale bar = 3 mm (B), 700 μm (H), 2 cm (J), 1 cm (K, L).

Figure 7. The dorsoventral-mediated Shh induction mechanism.

Schematic images of a normal blastema (A), AntBL (B), PostBL (C), DorBL (D), and VentBL (E).

Green, red, blue, and yellow boxes within the blastema represent cells derived from anterior,

posterior, dorsal, and ventral regions, respectively. Colored arrows indicate the presence of the

corresponding signals mediated by these cells. In this model, limb patterning requires both FGF8 and

SHH, and *Shh* expression in posteriorly derived cells is induced by the co-existence of WNT10B and

FGF2.

Figure S1. Gene expression patterns across the entire ALM-induced blastemas.

Sections of ALM blastemas induced at the anterior (A–C), posterior (D–F), dorsal (G–I), or ventral

(J–L) region at 10 dps. *Lmx1b* (A, D, G, J), *Fgf8* (B, E, H, K), and *Shh* (C, F, I, L) expression was

detected by *in situ* hybridization. These panels show the full blastema regions corresponding to the

higher-magnification views in Fig. 2. Scale bar = 700 μm (A).

Figure S2. Histological analysis of intact limbs and limbs induced by Shh electroporation.

(A–D) Transverse sections taken at multiple levels along the proximodistal axis from an intact limb

and from the limbs shown in Fig. 4. Sections from limbs induced from DorBL and VentBL exhibit

dorsoventrally symmetric internal structures, whereas intact limbs and limbs induced from AntBL do

not. Scale bar = 1 mm.

Figure S3. Limb patterning from BIO-treated VentBL.

(A, C) VentBL treated with 1μM BIO (C) or DMSO (A) at 90 dps. (E) Quantitative analysis of *Shh*

expression in VentBL treated with 1μM BIO and with DMSO. RNA was prepared from 4 identical

VentBL for both. Technological replicates are plotted in the same color. (C–D) Histological analysis

and pixel classification was performed in the same way as in Fig. 4. The dotted boxes indicate the

regions shown in (D). (E) Symmetry scores of each class. Scores obtained from the same limb are

plotted at the same x-coordinates. The plots of "int" are the same plots as in Fig. 4. n.s.; no

significant difference, $*p<0.05$, $**p<0.005$. Scale bar = 4 mm (B), 1 cm (C).

Figure S4. Histological analysis of limbs induced by Fgf2 or Wnt10b electroporation.

(A, B) Transverse sections taken at multiple levels along the proximodistal axis from the limbs

shown in Fig. 5. The induced limbs exhibit dorsoventrally symmetric internal structures. Scale bar =

1 mm.

Figure S5. Gene expression patterns during normal limb regeneration.

(A) Sections of amputation-induced blastemas at the early bud (EB), middle bud (MB), and late bud

(LB) stages. *Lmx1b*, *Wnt10b*, and *Fgf2* expression was visualized by *in situ* hybridization. (B)

Quantitative analysis of *Lmx1b*, *Wnt10b*, and *Fgf2* expression in MB-stage blastemas during normal

regeneration (n=13). Gene expression was quantified by RT-qPCR on manually microdissected

dorsal and ventral halves of each blastema. $*p<0.05$, $**p<0.005$ (two-tailed paired *t*-test). (C)

Quantitative analysis of *Wnt10b*, *Fgf2*, and *Shh* expression across stages (intact, EB, MB, LB, and

early digit [ED]; n = 5 for all stages). *Wnt10b* and *Fgf2* expression peaked at the MB stage, whereas

Shh expression peaked later, at the LB stage. One-way ANOVA detected a stage-dependent

difference (p < 0.05 for *Wnt10b* and *Fgf2*; p < 0.001 for *Shh*), followed by Dunnett's test. $*p<0.05$,

$**p<0.005$.

Figure S6. Gene expression patterns in a normal blastema assessed by reanalysis of axolotl

single-cell RNA-seq data.

Reanalysis of single-cell RNA-seq data from a middle bud (MB) stage blastema (Li et al., 2021).

(A–G) FeaturePlot visualizations of *Prrx1* (mesenchyme marker), *Krt17* (epithelium marker),

Lmx1b, *Wnt10b*, *Fgf2*, *Fgf8*, and *Shh*. *Lmx1b*, *Fgf2*, *Fgf8*, and *Shh* expression are detected in the

mesenchymal cluster, whereas *Wnt10b* expression is detected in both mesenchymal and epithelial

clusters. (H–M) "Co-expression" maps generated by classifying cells as expressing gene A only,

gene B only, both genes, or neither gene, and overlaying these classes on the UMAP using DimPlot.

Table 1. The induction rate of bump/limb formation in ALM.

References

Al Ghamdi, M. A., Al-Qattan, M. M., Hadadi, A., Alabdulrahman, A., Almuzzaini, B., Alatwi, N., & AlBalwi, M. A. (2020). A classification system for split-hand/ foot malformation (SHFM): A proposal based on 3 pedigrees with WNT10B mutations. European Journal of Medical Genetics, 63(3), 103738. https://doi.org/https://doi.org/10.1016/j.ejmg.2019.103738

Bennett, C. N., Longo, K. A., Wright, W. S., Suva, L. J., Lane, T. F., Hankenson, K. D., & MacDougald, O. A. (2005). Regulation of osteoblastogenesis and bone mass by Wnt10b. Proceedings of the National Academy of Sciences, 102(9), 3324–3329. https://doi.org/10.1073/pnas.0408742102

Berg, S., Kutra, D., Kroeger, T., Straehle, C. N., Kausler, B. X., Haubold, C., Schiegg, M., Ales, J., Beier, T., Rudy, M., Eren, K., Cervantes, J. I., Xu, B., Beuttenmueller, F., Wolny, A., Zhang, C., Koethe, U., Hamprecht, F. A., & Kreshuk, A. (2019). ilastik: interactive machine learning for (bio)image analysis. Nature Methods, 16(12), 1226–1232. https://doi.org/10.1038/s41592-019-0582-9

Bilal, M., Haack, T. B., Buchert, R., Peralta, S., Ahmad, I., Faisal, Abbasi, S., & Ahmad, W. (2023). Sequence Variants in the WNT10B Underlying Non-Syndromic Split-Hand/Foot Malformation. Molecular Syndromology, 14(6), 469–476. https://doi.org/10.1159/000531069

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics, 30(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

BRYANT, S. V. (1976). Regenerative failure of double half limbs in Notophthalmus viridescens. Nature, 263(5579), 676–679. https://doi.org/10.1038/263676a0

Bryant, S. V, & Baca, B. A. (1978). Regenerative ability of double-half and half upper arms in the newt, Notophthalmus viridescens. Journal of Experimental Zoology, 204(3), 307–323. https://doi.org/https://doi.org/10.1002/jez.1402040302

Bryant, S. V, & Iten, L. E. (1976). Supernumerary limbs in amphibians: Experimental production in Notophthalmus viridescens and a new interpretation of their formation. Developmental Biology, 50(1), 212–234. https://doi.org/https://doi.org/10.1016/0012-1606(76)90079-8

Burton, R., Holder, N., & Jesani, P. (1986). The regeneration of double dorsal and double ventral limbs in the axolotl. Development, 94(1), 29–46. https://doi.org/10.1242/dev.94.1.29

Chen H, J. R. (2002). Interactions between dorsal-ventral patterning genes lmx1b, engrailed-1 and wnt-7a in the vertebrate limb. Int J Dev Biol, 46(7), 937–941.

Chen, H., Lun, Y., Ovchinnikov, D., Kokubo, H., Oberg, K. C., Pepicelli, C. V, Can, L., Lee, B., & Johnson, R. L. (1998). Limb and kidney defects in Lmx1b mutant mice suggest an involvement of LMX1B in human nail patella syndrome. Nature Genetics, 19(1), 51–55. https://doi.org/10.1038/ng0598-51

Cygan, J. A., Johnson, R. L., & McMahon, A. P. (1997). Novel regulatory interactions revealed by studies of murine limb pattern in Wnt-7a and En-1 mutants. Development, 124(24), 5021–5032.

https://doi.org/10.1242/dev.124.24.5021

Endo, T., Bryant, S. V, & Gardiner, D. M. (2004). A stepwise model system for limb regeneration.

Developmental Biology, 270(1), 135–145.

https://doi.org/https://doi.org/10.1016/j.ydbio.2004.02.016

Endo, T., Gardiner, D. M., Makanae, A., & Satoh, A. (2015). The Accessory Limb Model: An

Alternative Experimental System of Limb Regeneration. In A. Kumar & A. Simon (Eds.),

Salamanders in Regeneration Research: Methods and Protocols (pp. 101–113). Springer New

York. https://doi.org/10.1007/978-1-4939-2495-0_8

Gerber, T., Murawala, P., Knapp, D., Masselink, W., Schuez, M., Hermann, S., Gac-Santel, M.,

Nowoshilow, S., Kageyama, J., Khattak, S., Currie, J. D., Camp, J. G., Tanaka, E. M., & Treutlein,

B. (2018). Single-cell analysis uncovers convergence of cell identities during axolotl limb

regeneration. Science, 362(6413), eaaq0681. https://doi.org/10.1126/science.aaq0681

Giampaoli, S., Bucci, S., Ragghianti, M., Mancino, G., Zhang, F., & Ferretti, P. (2003). Expression of

FGF2 in the limb blastema of two Salamandridae correlates with their regenerative capability.

Proceedings of the Royal Society of London. Series B: Biological Sciences, 270(1530), 2197–

2205. https://doi.org/10.1098/rspb.2003.2439

Glotzer, G. L., Tardivo, P., & Tanaka, E. M. (2022). Canonical Wnt signaling and the regulation of

divergent mesenchymal Fgf8 expression in axolotl limb development and regeneration. ELife,

11, e79762. https://doi.org/10.7554/eLife.79762

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., Zheng, S., Butler, A., Lee, M. J., Wilk, A. J.,

Darby, C., Zager, M., Hoffman, P., Stoeckius, M., Papalexi, E., Mimitou, E. P., Jain, J., Srivastava,

A., Stuart, T., Fleming, L. M., Yeung, B., … Satija, R. (2021). Integrated analysis of multimodal

single-cell data. Cell, 184(13), 3573-3587.e29.

https://doi.org/https://doi.org/10.1016/j.cell.2021.04.048

Hirata, A., Gardiner, D. M., & Satoh, A. (2010). Dermal fibroblasts contribute to multiple tissues in

the accessory limb model. DGD, 52(4), 343–350. https://doi.org/https://doi.org/10.1111/j.1440-

169X.2009.01165.x

Iten, L. E., & Bryant, S. V. (1975). The interaction between the blastema and stump in the

establishment of the anterior-posterior and proximal-distal organization of the limb regenerate.

Developmental Biology, 44(1), 119–147. https://doi.org/https://doi.org/10.1016/0012-

1606(75)90381-4

Iwata, R., Makanae, A., & Satoh, A. (2020). Stability and plasticity of positional memory during limb

regeneration in Ambystoma mexicanum. Developmental Dynamics, 249(3), 342–353.

https://doi.org/https://doi.org/10.1002/dvdy.96

Kashimoto, R., Furukawa, S., Yamamoto, S., & Satoh, A. (2023). Bead Implantation and Delivery of

Exogenous Growth Factors. In A. W. Seifert & J. D. Currie (Eds.), Salamanders: Methods and

Protocols (pp. 209–216). Springer US. https://doi.org/10.1007/978-1-0716-2659-7_14

Kawakami, Y., Rodriguez Esteban, C., Raya, M., Kawakami, H., Martí, M., Dubova, I., & Izpisúa
Belmonte, J. C. (2006). Wnt/β-catenin signaling regulates vertebrate limb regeneration. Genes
& Development, 20(23), 3232–3237. https://doi.org/10.1101/gad.1475106

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: a fast spliced aligner with low memory
requirements. Nature Methods, 12(4), 357–360. https://doi.org/10.1038/nmeth.3317

Kragl, M., Knapp, D., Nacu, E., Khattak, S., Maden, M., Epperlein, H. H., & Tanaka, E. M. (2009).
Cells keep a memory of their tissue origin during axolotl limb regeneration. Nature, 460(7251),
60–65. https://doi.org/10.1038/nature08152

Laufer, E., Nelson, C. E., Johnson, R. L., Morgan, B. A., & Tabin, C. (1994). Sonic hedgehog and Fgf-
4 act through a signaling cascade and feedback loop to integrate growth and patterning of the
developing limb bud. Cell, 79(6), 993–1003. https://doi.org/https://doi.org/10.1016/0092-
8674(94)90030-2

Lévesque, M., Gatien, S., Finnson, K., Desmeules, S., Villiard, É., Pilote, M., Philip, A., & Roy, S.
(2007). Transforming Growth Factor: β Signaling Is Essential for Limb Regeneration in Axolotls.
PLOS ONE, 2(11), e1227-. https://doi.org/10.1371/journal.pone.0001227

Li, H., Wei, X., Zhou, L., Zhang, W., Wang, C., Guo, Y., Li, D., Chen, J., Liu, T., Zhang, Y., Ma, S.,
Wang, C., Tan, F., Xu, J., Liu, Y., Yuan, Y., Chen, L., Wang, Q., Qu, J., … Xu, X. (2021). Dynamic
cell transition and immune response landscapes of axolotl limb regeneration revealed by single-
cell analysis. Protein & Cell, 12(1), 57–66. https://doi.org/10.1007/s13238-020-00763-1

Li, Q., Kannan, A., Das, A., DeMayo, F. J., Hornsby, P. J., Young, S. L., Taylor, R. N., Bagchi, M. K.,
& Bagchi, I. C. (2013). WNT4 Acts Downstream of BMP2 and Functions via β-Catenin
Signaling Pathway to Regulate Human Endometrial Stromal Cell Differentiation. Endocrinology,
154(1), 446–457. https://doi.org/10.1210/en.2012-1585

Li, S., Anderson, R., Reginelli, A. D., & Muneoka, K. (1996). FGF-2 influences cell movements and
gene expression during limb development. Journal of Experimental Zoology, 274(4), 234–247.
https://doi.org/https://doi.org/10.1002/(SICI)1097-010X(19960301)274:4<234::AID-
JEZ4>3.0.CO;2-Q

Li, X., Li, Z., Wang, J., Li, Z., Cui, H., Dai, G., Chen, S., Zhang, M., Zheng, Z., Zhan, Z., & Liu, H.
(2019). Wnt4 signaling mediates protective effects of melatonin on new bone formation in an
inflammatory environment. The FASEB Journal, 33(9), 10126–10139.
https://doi.org/https://doi.org/10.1096/fj.201900093RR

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinformatics, 30(7), 923–930.
https://doi.org/10.1093/bioinformatics/btt656

Lin, Y.-C., Haas, A., Bufe, A., Parbin, S., Hennecke, M., Voloshanenko, O., Gross, J., Boutros, M.,

Acebron, S. P., & Bastians, H. (2021). Wnt10b-GSK3β–dependent Wnt/STOP signaling prevents aneuploidy in human somatic cells. Life Science Alliance, 4(1), e202000855. https://doi.org/10.26508/lsa.202000855

Logan, C., Hornbruch, A., Campbell, I., & Lumsden, A. (1997). The role of Engrailed in establishing the dorsoventral axis of the chick limb. Development, 124(12), 2317–2324. https://doi.org/10.1242/dev.124.12.2317

Loomis, C. A., Harris, E., Michaud, J., Wurst, W., Hanks, M., & Joyner, A. L. (1996). The mouse Engrailed-1 gene and ventral limb patterning. Nature, 382(6589), 360–363. https://doi.org/10.1038/382360a0

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology, 15(12), 550. https://doi.org/10.1186/s13059-014-0550-8

Lovely, A. M., Duerr, T. J., Qiu, Q., Galvan, S., Voss, S. R., & Monaghan, J. R. (2022). Wnt Signaling Coordinates the Expression of Limb Patterning Genes During Axolotl Forelimb Development and Regeneration. Frontiers in Cell and Developmental Biology, 10. https://www.frontiersin.org/journals/cell-and-developmental-biology/articles/10.3389/fcell.2022.814250

Ludolph, D. C., Cameron, J. A., & Stocum, D. L. (1990). The effect of retinoic acid on positional memory in the dorsoventral axis of regenerating axolotl limbs. Developmental Biology, 140(1), 41–52. https://doi.org/https://doi.org/10.1016/0012-1606(90)90051-J

Maden, M. (1980). Structure of supernumerary limbs. Nature, 286(5775), 803–805. https://doi.org/10.1038/286803a0

Makanae, A., Mitogawa, K., & Satoh, A. (2014a). Co-operative Bmp- and Fgf-signaling inputs convert skin wound healing to limb formation in urodele amphibians. Developmental Biology, 396(1), 57–66. https://doi.org/https://doi.org/10.1016/j.ydbio.2014.09.021

Makanae, A., Mitogawa, K., & Satoh, A. (2014b). Implication of two different regeneration systems in limb regeneration. Regeneration, 1(3), 1–9. https://doi.org/https://doi.org/10.1002/reg2.16

McCusker, C., Lehrberg, J., & Gardiner, D. (2014). Position-specific induction of ectopic limbs in non-regenerating blastemas on axolotl forelimbs. Regeneration, 1(1), 27–34. https://doi.org/https://doi.org/10.1002/reg2.10

Mescher, A. L. (1976). Effects on adult newt limb regeneration of partial and complete skin flaps over the amputation surface. Journal of Experimental Zoology, 195(1), 117–127. https://doi.org/https://doi.org/10.1002/jez.1401950111

Muneoka, K., Fox, W. F., & Bryant, S. V. (1986). Cellular contribution from dermis and cartilage to the regenerating limb blastema in axolotls. Developmental Biology, 116(1), 256–260. https://doi.org/https://doi.org/10.1016/0012-1606(86)90062-X

Nacu, E., Gromberg, E., Oliveira, C. R., Drechsel, D., & Tanaka, E. M. (2016). FGF8 and SHH
substitute for anterior–posterior tissue interactions to induce limb regeneration. Nature,
533(7603), 407–410. https://doi.org/10.1038/nature17972

Niswander, L., Jeffrey, S., Martin, G. R., & Tickle, C. (1994). A positive feedback loop coordinates
growth and patterning in the vertebrate limb. Nature, 371(6498), 609–612.
https://doi.org/10.1038/371609a0

Otsuki, L., & Tanaka, E. (2021). Positional Memory in Vertebrate Regeneration: A Century's Insights
from the Salamander Limb. Cold Spring Harbor Perspectives in Biology, 14, a040899.
https://doi.org/10.1101/cshperspect.a040899

Parr, B. A., & McMahon, A. P. (1995). Dorsalizing signal Wnt-7a required for normal polarity of D–
V and A–P axes of mouse limb. Nature, 374(6520), 350–353. https://doi.org/10.1038/374350a0

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides fast and
bias-aware quantification of transcript expression. Nature Methods, 14(4), 417–419.
https://doi.org/10.1038/nmeth.4197

Poulin, M. L., Patrie, K. M., Botelho, M. J., Tassava, R. A., & Chiu, I.-M. (1993). Heterogeneity in
the expression of fibroblast growth factor receptors during limb regeneration in newts
(Notophthalmus viridescens). Development, 119(2), 353–361.
https://doi.org/10.1242/dev.119.2.353

Riddle, R. D., Ensini, M., Nelson, C., Tsuchida, T., Jessell, T. M., & Tabin, C. (1995). Induction of the
LIM homeobox gene Lmx1 by WNT6a establishes dorsoventral pattern in the vertebrate limb.
Cell, 83(4), 631–640. https://doi.org/https://doi.org/10.1016/0092-8674(95)90103-5

Riddle, R. D., Johnson, R. L., Laufer, E., & Tabin, C. (1993). Sonic hedgehog mediates the polarizing
activity of the ZPA. Cell, 75(7), 1401–1416. https://doi.org/https://doi.org/10.1016/0092-
8674(93)90626-2

Sader, F., & Roy, S. (2022). Tgf-β superfamily and limb regeneration: Tgf-β to start and Bmp to end.
Developmental Dynamics, 251(6), 973–987. https://doi.org/https://doi.org/10.1002/dvdy.379

Satoh, A., Cummings, G. M. C., Bryant, S. V, & Gardiner, D. M. (2010). Regulation of proximal-distal
intercalation during limb regeneration in the axolotl (Ambystoma mexicanum). Development,
Growth & Differentiation, 52(9), 785–798. https://doi.org/https://doi.org/10.1111/j.1440-
169X.2010.01214.x

Satoh, A., Gardiner, D. M., Bryant, S. V, & Endo, T. (2007). Nerve-induced ectopic limb blastemas in
the axolotl are equivalent to amputation-induced blastemas. Developmental Biology, 312(1),
231–244. https://doi.org/https://doi.org/10.1016/j.ydbio.2007.09.021

Satoh, A., Makanae, A., Nishimoto, Y., & Mitogawa, K. (2016). FGF and BMP derived from dorsal
root ganglia regulate blastema induction in limb regeneration in Ambystoma mexicanum.
Developmental Biology, 417(1), 114–125.

https://doi.org/https://doi.org/10.1016/j.ydbio.2016.07.005

Schloissnig, S., Kawaguchi, A., Nowoshilow, S., Falcon, F., Otsuki, L., Tardivo, P., Timoshevskaya,
N., Keinath, M. C., Smith, J. J., Voss, S. R., & Tanaka, E. M. (2021). The giant axolotl genome
uncovers the evolution, scaling, and transcriptional control of complex gene loci. Proceedings
of the National Academy of Sciences, 118(15), e2017176118.
https://doi.org/10.1073/pnas.2017176118

SHIMOKAWA, T., YASUTAKA, S., KOMINAMI, R., & SHINOHARA, H. (2013). Lmx-1b and
Wnt-7a expression in axolotl limb during development and regeneration. Okajimas Folia
Anatomica Japonica, 89(4), 119–124. https://doi.org/10.2535/ofaj.89.119

Srivastava, A., Malik, L., Smith, T., Sudbery, I., & Patro, R. (2019). Alevin efficiently estimates
accurate gene abundances from dscRNA-seq data. Genome Biology, 20(1), 65.
https://doi.org/10.1186/s13059-019-1670-y

Stocum, D. L. (1982). Determination of axial polarity in the urodele limb regeneration blastema.
Development, 71(1), 193–214. https://doi.org/10.1242/dev.71.1.193

Thomas, P. D., Kejariwal, A., Campbell, M. J., Mi, H., Diemer, K., Guo, N., Ladunga, I., Ulitsky-
Lazareva, B., Muruganujan, A., Rabkin, S., Vandergriff, J. A., & Doremieux, O. (2003).
PANTHER: a browsable database of gene products organized by biological function, using
curated protein family and subfamily classification. Nucleic Acids Research, 31(1), 334–341.
https://doi.org/10.1093/nar/gkg115

Thornton, C. S. (1962). Influence of head skin on limb regeneration in urodele amphibians. Journal of
Experimental Zoology, 150(1), 5–15. https://doi.org/https://doi.org/10.1002/jez.1401500103

Todd, T. (1823). On the process of reproduction of the members of the aquatic salamander. Quart J Sci
Lit Arts Lib, 16, 84–86.

Tsai, S. (2020). Inhibition of Wound Epidermis Formation via Full Skin Flap Surgery During Axolotl
Limb Regeneration. JoVE, 160, e61522. https://doi.org/doi:10.3791/61522

Ugur, S. A., & Tolun, A. (2008). Homozygous WNT10b mutation and complex inheritance in Split-
Hand/Foot Malformation. Human Molecular Genetics, 17(17), 2644–2653.
https://doi.org/10.1093/hmg/ddn164

Vieira, W. A., Wells, K. M., Raymond, M. J., De Souza, L., Garcia, E., & McCusker, C. D. (2019).
FGF, BMP, and RA signaling are sufficient for the induction of complete limb regeneration from
non-regenerating wounds on Ambystoma mexicanum limbs. Developmental Biology, 451(2),
146–157. https://doi.org/https://doi.org/10.1016/j.ydbio.2019.04.008

Vogel, A., Rodriguez, C., Warnken, W., & Belmonte, J. C. I. (1995). Dorsal cell fate specified by chick
Lmxl during vertebrate limb development. Nature, 378(6558), 716–720.
https://doi.org/10.1038/378716a0

Voss, J. H., Koszegi, Z., Yan, Y., Shorter, E., Grätz, L., Lanner, J. T., Calebiro, D., & Schulte, G. (2025).

WNT-induced association of Frizzled and LRP6 is not sufficient for the initiation of WNT/β-catenin signaling. Nature Communications, 16(1), 4848. https://doi.org/10.1038/s41467-025-60096-7

Witte, F., Dokas, J., Neuendorf, F., Mundlos, S., & Stricker, S. (2009). Comprehensive expression analysis of all Wnt genes and their major secreted antagonists during mouse limb development and cartilage differentiation. Gene Expression Patterns, 9(4), 215–223. https://doi.org/https://doi.org/10.1016/j.gep.2008.12.009

Yamamoto, S., Kashimoto, R., Furukawa, S., Ohashi, A., & Satoh, A. (2022). Lmx1b activation in axolotl limb regeneration. Developmental Dynamics, 251(9), 1509–1523. https://doi.org/https://doi.org/10.1002/dvdy.476

Yang, Y., & Niswander, L. (1995). Interaction between the signaling molecules WNT7a and SHH during vertebrate limb development: Dorsal signals regulate anteroposterior patterning. Cell, 80(6), 939–947. https://doi.org/https://doi.org/10.1016/0092-8674(95)90297-X

Yonei-Tamura, S., Endo, T., Yajima, H., Ohuchi, H., Ide, H., & Tamura, K. (1999). FGF7 and FGF10 Directly Induce the Apical Ectodermal Ridge in Chick Embryos. Developmental Biology, 211(1), 133–143. https://doi.org/https://doi.org/10.1006/dbio.1999.9290

Yamamoto et al., 2025, Figure 1

Yamamoto et al., 2025, Figure 2

Yamamoto et al., 2025, Figure 3

Yamamoto et al., 2025, Figure 4

Yamamoto et al., 2025, Figure 5

Dorsal + bead

Ventral + bead

Dorsal + bead

Ventral + bead

Yamamoto et al., 2025, Figure 6

Yamamoto et al., 2025, Figure 7

Yamamoto et al., 2025, Supplemental Figure 1

Yamamoto et al., 2025, Supplemental Figure 2

Yamamoto et al., 2025, Supplemental Figure 3

A DorBL + *Fgf2*

B VentBL + *Wnt10b*

distal ↕ proximal

distal ↕ proximal

Yamamoto et al., 2025, Supplemental Figure 4

Yamamoto et al., 2025, Supplemental Figure 5

Yamamoto et al., 2025, Supplemental Figure 6

	N	Regress		Bump		Limb	
AntBL	8	3	(37.5 %)	5	(62.5 %)	0	(0.0 %)
AntBL+P	9	0	(0.0 %)	1	(11.1 %)	8	(88.9 %)
PostBL	8	6	(75.0 %)	2	(25.0 %)	0	(0.0 %)
PostBL+A	7	2	(28.6 %)	1	(14.3 %)	4	(57.1 %)
DorBL	12	8	(66.7 %)	3	(25.0 %)	1	(8.3 %)
DorBL+V	14	5	(35.7 %)	2	(14.3 %)	7	(50.0 %)
VentBL	22	15	(68.2 %)	7	(31.8 %)	0	(0.0 %)
VentBL+D	11	2	(18.2 %)	2	(18.2 %)	7	(63.6 %)