Title: Glial *betaPix* is essential for blood vessel integrity in the zebrafish brain

ShihChing Chiu[1], Qinchao Zhou[1], Chenglu Xiao[2], Linlu Bai[1], Xiaojun Zhu[1,*], Wanqiu

Ding[1,*], and Jing-Wei Xiong[1,2,*,#]

[1]Beijing Key Laboratory of Cardiometabolic Molecular Medicine, Institute of

Molecular Medicine, and College of Future Technology, Peking University, Beijing

100871, China; [2]School of Basic Medical Sciences, The Second Affiliated Hospital,

Institute of Biomedical Innovation, The MOE Basic Research and Innovation Center

for the Targeted Therapeutics of Solid Tumors, Jiangxi Medical College, Nanchang

University, Nanchang 330031, China

*Corresponding authors: WD (dingwq@pku.edu.cn), XZ (zhuxiaojun@pku.edu.cn),

and JWX (jingwei_xiong@pku.edu.cn)

#Lead Contact: Jing-Wei Xiong, Ph.D. (jingwei_xiong@pku.edu.cn)

Abstract

The formation of blood-brain barrier and vascular integrity depends on the coordinative development of different cell types in the brain. Previous studies have shown that zebrafish *bubblehead* (*bbh*) mutant, which has mutation in the *betaPix* locus, develops spontaneous intracerebral hemorrhage during early development. However, it remains unclear in which brain cells *betaPix* may function. Here, we established a highly efficient conditional knockout method in zebrafish by using homology-directed repair (HDR)-mediated knockin and knockout technology, and generated *betaPix* conditional trap (*betaPix^{ct}*) allele in zebrafish. We found that *betaPix* in glia, but neither neurons, endothelial cells, nor pericytes, was critical for glial and vascular development and integrity, thus contributing to the formation of blood-brain barrier. Single-cell transcriptome profiling revealed that microtubule aggregation signaling stathmins and pro-angiogenic transcription factors *Zfhx3/4* were down-regulated in glial and neuronal progenitors, and further genetic analysis suggested that betaPix acted upstream on the PAK1-Stathmin and Zfhx3/4-Vegfaa signaling to regulate glia migration and vascular integrity. Therefore, this work reveals that glial *betaPix* plays an important role in brain vascular integrity in zebrafish embryos and possibly human cells.

Keywords: *betaPix*; hemorrhage; glial migration; knock-in; stathmin

Introduction

Intracerebral hemorrhage (ICH) is a life-threatening stroke type with the worst prognosis and few proven clinical treatments. Most patients who survive intracerebral hemorrhage end up with disabilities and are at risk for recurrency, cognitive decline, and systemic vascular issues, making this condition particularly significant among neurological disorders [1,2]. Several rodent models have been used for modeling ICH, including autologous blood injection, collagenase injection, thrombin injection, and micro-balloon inflation techniques [3]. Zebrafish (Danio rerio) exhibit a closed circulatory system and regulatory pathways that are highly conserved among vertebrates [4] with relatively cheap and easy to maintain disease models. Over the past decades, varieties of zebrafish mutants that spontaneously developed brain hemorrhage have been generated in mutagenesis screens, such as *redhead* [5], *reddish* [6,7] and *bubblehead* (*bbh*) [8,9]. Elucidating mutated gene function in these zebrafish models enables us to gain insights into the genetic basis and pathophysiological mechanisms of ICH development. The *bbh* mutant was identified independently in two large-scale ethylnitrosourea-induced mutagenesis screens [8,9] and positional cloning revealed that p21-activated kinase (Pak)-interacting exchange factor beta (*betaPix*) gene was mutated in *bbh* mutants [10]. *betaPix* contains SH3 domain that binds group I PAKs, and Dbl homology (DH) and pleckstrin homology (PH) domains that function as a RhoGEF to interact with Rac/Cdc42 small GTPases [11], thus participating in multiple cellular pathways to regulate cell polarity, adhesion and migration [12].

Various studies have reported that brain vessels are highly supported by neurons and

glial cells during development and regeneration [13,14]. Glial-specific *betaPix* have been

proposed to interact with *αvβ8* integrin and the Band 4.1s in glia, which link to

multiple intracellular signaling effectors such as *Wnt7a* and *Wnt7b* [15], but this

hypothesis was not experimentally addressed [16]. To this end, *betaPix* has been shown

to interact with *αvβ8* integrins and mediate focal adhesion formation and thus

modulate cerebral vascular stability in *bbh* zebrafish mutant [17]. Integrins are the main

molecular link between cells and the extracellular matrix (ECM) which serve as

scaffolds in the perivascular space. Based on a chemical suppressor screening of *bbh*

hemorrhages, we have previously reported that a small molecule called miconazole

downregulated the pErk-matrix metalloproteinase 9 (Mmp9) signaling to reduce ECM

degradation, thus improving vascular integrity [18]. However, how *betaPix* cell-type

specifically contributes to vascular integrity and maturation remains unclear. Here, we

generated *betaPix* conditional trap (*betaPix^ct^*) alleles by using a CRISPR/HDR-based

Zwitch method, and found that betaPix acted mainly in glia to regulate glial and

vascular integrity *via* Stathmin and Zfhx3/4 signaling. Therefore, this work provides

the first experimental evidence that betaPix acts in glia for vascular integrity

development, and its functional conservation between zebrafish and human cells may

further guide us to decipher the genetic basis of ICH in the future.

Results

Generating *betaPix* conditional trap (*betaPix^{ct}*) allele by an HDR-mediated knockin and knockout method

To study *betaPix* function in cell-specific manner, we utilized a donor vector with an invertible gene trap cassette *via* HDR (Zwitch) [19-21] for generating *betaPix* conditional trap (*betaPix^{ct}*) alleles. Zwitch consists of a splice acceptor conjugated to a red fluorescent protein *via* a 2A self-cleaving peptide, flanked by two LoxP sites and two Lox 5171 sites in reciprocal orientations. Outside this conditional trap cassette, we included a lens-specific enhanced green fluorescence protein α-crystallin:EGFP for screening knockout founders, as well as inserted both left and right arms for targeting homologous sequences on the genome. We modified the Zwitch vector by adding a glycine-serine-glycine (GSG) spacer in front of the 2A peptide to enhance cleavage efficiency, and adding universal guide RNA (UgRNA)[22] target sequences outside the left and right homologous arms for linearizing donor vector (Figures 1A and S1A). We chose CRISPR/Cas9 system to generate double-strand breaks (DSBs) on intron 5 of the *betaPix* locus, and chose the most efficient guide RNAs based on their efficiency to induce CRISPR indels. We used either long arms such as ~1000 bp or short arms such as 24 bp that located upstream and downstream the guide RNA sites. We then co-injected the donor vectors with targeting guide RNA, universal guide RNA, and Cas9 mRNA into one-cell-stage embryos. At 4 days post-fertilization (dpf), we selected α-crystallin-EGFP-positive embryos (F_0) for raising to adulthood, and pre-selected F_0 founders by examining inheritable EGFP reporter expression of F_1 embryos. To confirm

the correct homologous recombination in the *betaPix* locus, we identified potential

founders by genomic PCR to examine Zwitch insertion in the *betaPix* locus. Precise

knock-in genotypes were further verified using Sanger sequencing (Figures 1B, 1C,

S1B). Among 184 adult F_0 founders with *α*-crystallin-EGFP expression, we found 84

founders had EGFP-positive F_1 embryos, and 7 founders had expected PCR fragments

around both 5' and 3' arms, which two founders had correct insertions in the *betaPix*

locus by Sanger sequencing (Figure S1C), thus achieving ~1% efficiency on generating

conditional alleles. Briefly, we established an efficient method for generating

conditional knockin/knockout mutants in general, and generated a conditional gene trap

line Ki(*betaPix*:Zwitch) in the *betaPix* locus in particular, which is referred to as

betaPix^{ct} hereafter.

A previous work has shown *bubblehead* (*bbh^{m292}*) mutant with a hypomorphic mutation

in *betaPix*, and *bbh^{fn40a}* mutant with reduced *betaPix* transcripts without finding

mutations in the coding region and the splicing sites [10]. *bbh^{fn40a}* mutants developed

cerebral hemorrhages phenotype between 36 to 52 hpf (Figure S1D and S1E). To

characterize the *betaPix^{ct}* allele, we injected one-cell-stage *betaPix^{ct/ct}* embryos with

Cre mRNA to induce global inversion of Zwitch and deletion of *betaPix*. As expected,

we found Cre-induced precise inversion in *betaPix^{m/m}* as confirmed by Sanger

sequencing (Figure 1C) and severe brain hemorrhages developed in the time window

similar to *bbh^{fn40a}* mutants (Figures 1D and 1E). Consistently, qPCR analysis showed

that *betaPix* decreased and *RFP* increased in heterozygous *betaPix^{m/+}* and homozygous

betaPix^{m/m} mutants compared with *betaPix^{ct/ct}* wild-type siblings, while *alphaPix*

expression was not affected (Figure 1F). Of note, *TagRFP* expression in *betaPix^{m/+}*

(after Cre-mediated recombination in *betaPix^{ct/+}* embryos) reflected endogenous

betaPix expression in the boundary of midbrains and hindbrains as well as the

hindbrains (Figure S1F and S1G). Together, these results suggest that the conditional

trap *betaPix^{ct/ct}* zebrafish is functional for visualizing endogenous *betaPix* expression

(knockin) and loss-of-function study (conditional knockout).

betaPix^{m/m} mutant has brain hemorrhages, central arteries defects and abnormal

glial structure that is partially rescued by Pak1 inhibitor IPA3 treatment

The main phenotypes in *bbh* mutants consist of brain hemorrhage and hydrocephalus

as early as 36 hpf, as well as poor endothelial–mesenchymal contacts and defective

central arteries (CtAs) sprouting in the hindbrain [10,17]. To demonstrate the practicability

of *betaPix* conditional trap lines, we assessed vasculature and glia in the hindbrain by

using light sheet fluorescence microscopy. Global *betaPix* inactivation in *betaPix^{m/m}*

mutants resulted in severe cerebral hemorrhages and defective angiogenesis in

hindbrain central arteries (Figures 1D, 1E, S2A and S2B), which is consistent with

previous studies [10,18].

It has been suggested that endothelial *betaPix* might not contribute to the occurrence of

brain hemorrhage [10]. During brain development, neurons and glia are important

perivascular cells and they orchestrate with endothelial cells in a temporal and spatial

pattern. Thus, we studied hindbrain radial glia development and found that radial glial

structure had disoriented arrangement and shorten process length in *betaPix^{m/m}* embryos

compared to siblings (Figures 2A, 2B and S2C). As a result, neuronal and glial

precursors marker *nestin* increased while differentiated neuronal marker *pax2a*

decreased in *betaPix^{m/m}* embryos, suggesting delayed neuronal development and

differentiation after *betaPix* inactivation (Figure 2C). By using global *betaPix* knockout

with multiple guide-RNA simultaneously [23], we found that CRISPR-induced *betaPix*

mutant F_0 embryos had almost identical phenotypes to those in *betaPix^{fn40a}* mutants

(Figure S3A-S3B), such as cerebral hemorrhages, central arteries defects, and abnormal

hindbrain glia and neuronal precursor development (Figures S3C to S3G), further

confirming the loss-of-function phenotypes in *betaPix^{m/m}* mutants.

Zebrafish central arteries vascularization in the hindbrains starts at 29 hpf. Tip cells

start to form in the dorsal side of the primordial hindbrain channels (PHBC), sprout and

migrate into the hindbrain tissue, moving towards the center around 36 hpf to establish

connections with the basilar artery. A characteristic arch architecture forms at 48 hpf as

boundaries dividing the hindbrain into rhombomeres[24]. In *betaPix* knockouts, CtA

sproutings were restricted to the bottom of glial processes, thus not being able to

migrate upwards to form an arched structure (Figure 2D). The transcription factor SRY

box-2 (Sox2) is another marker gene for neuron and glial precursors. *Sox2*-positive

precursor cells tightly wrapped around the central artery in control siblings, but Sox2

precursor cells had loose contacts with the central arteries, with larger perivascular

distances in *betaPix* mutants (Figure 2E). These results implicate the impaired

interactions between endothelial cells and neuronal/glial cells during development after

betaPix deletion.

p21-activated kinase (Pak) is a binding partner for *betaPix*. *Pak2a* has been shown

mediating downstream signaling in *bbh^{m292}* mutants. Group I PAK allosteric

inhibitor IPA-3 covalently modifies and stabilizes the autoinhibitory N-terminal

region of PAK1, PAK2 and PAK3 [25]. As expected, IPA-3 treatment significantly

decreased incidence and intensity of cerebral hemorrhages in *betaPix$^{m/m}$* mutant

embryos (Figure 2F and 2G), while IPA-3 treatment had no effect on hematopoiesis in

betaPix$^{ct/ct}$ control siblings. In addition, central arterial defects were partially rescued

in *betaPix$^{m/m}$* mutant embryos after IPA-3 treatment, showing increased endothelial

protruding into the hindbrain and more arch structure formation (Figure 2H and 2I).

IPA-3 treatment also decreased abnormal glial process arrangements, which the

lengths of glial processes statistically reached the levels of the control siblings (Figure

2J). Thus, these data suggest that the betaPix-Pak1/2 signaling regulates brain

vascular integrity during development.

Glial-specific *betaPix* knockouts recapture its global knockout phenotypes

To investigate *betaPix* function in different types of perivascular cells, we generated

glial- or neuronal-specific transgenic lines using either glial fibrillary acidic protein

(*gfap*) [26] or *huC* [27] promoters to drive *EGFP-Cre* fusion gene expression under

betaPix^{ct/ct} background, respectively (Figure S4A). EGFP signals reported *Cre*

expression while RFP signals efficiently reported Cre-induced gene trap cassette

inversion and disruption of *betaPix* expressions (Figure S4B to S4F). Interestingly,

glial-specific *betaPix* knockouts developed severe cerebral hemorrhages and abnormal

central arteries vascularization, which were partially rescued by IPA-3 treatment

(Figure 3A to 3C), highlighting betaPix function in glia *via* Pak1/2 signaling. On the

other hand, neither vascular-, neuronal-, nor mural-specific deletion of *betaPix* had

evident *betaPix* mutant phenotypes (Figure S5A to S5D). These results suggest that

glial *betaPix* plays crucial roles in zebrafish embryonic vascular integrity and glial

development *via* regulating *Pak* activities.

Single-cell transcriptome profiling reveals that stathmins in *gfap*-positive

progenitors were affected in *betaPix* knockouts

To investigate the interplays between glial and hemorrhagic pathology caused by

betaPix loss-of-function, we profiled the cranial tissues of CRISPR-edited zebrafish at

1 and 2 dpf using the 10X Chromium single-cell RNA sequencing (scRNA-seq)

platform (Figure 4A). Multi-guide targeting was able to generate almost 100% F_0 null

mutants, allowing us to select mutant embryos before brain hemorrhages from 36 hpf.

Low-quality cells were excluded based on the numbers of genes detected and

percentages of reads mapped to mitochondrial genes per sample. A total of 38,670 cells

passed quality control and were used for subsequent analyses. Uniform manifold

approximation and projection (UMAP) identified 71 cell clusters, which represented 24

zebrafish cranial cell types based on known marker gene expression profiles (Figure 4B

and S6A). Enriched gene markers were compared with previously annotated gene

markers in the ZFIN database and literatures[28,29]. By comparing the proportion of cells

in each sample, we found that most neuronal clusters had increased trends, while the

numbers of progenitors, endothelial cells, erythrocytes, neural crests, muscles,

cartilages, retinas (photoreceptor precursor cells), olfactory bulbs, or epidermis and

pharyneal arches were reduced in *betaPix* knockout heads at 2 dpf (Figure 4C and S6B).

This is consistent with the open public databases of single-cell transcriptome atlas

where *betaPix* weakly expressed in a broad range of cells including glia, neurons and

neuronal precursors, retinas, endothelial cells, pharynx, exocrine pancreas, olfactory

cells and heart cells [30]. Given that *betaPix* is critical in glia during brain vascular

integrity development (Figures 3 and S5), we examined the *gfap* expression among

each cluster, and found relatively high *gfap* expression in clusters including progenitors,

hindbrain, ventral diencephalon, ventral midbrain and floor plate (Figure S6A, arrow

indicates). We next focused on the progenitor cluster, as of its enriched *gfap* expression

and relatively more decreased cell numbers in *betaPix* mutants. Furthermore, this

progenitor cluster exhibited high-level expression of cell proliferation and cell cycle

related genes (*mki67*, *pcna*, *ccnd1*, *rrm1* and *rrm2*) as well as key glial-associated genes

(*gfap*, *fabp7a*, *her4*.1, *cx43*, *id1*, *fgfbp3*, *atp1a1b* and *mdka*) (Table S1). From

differentially expressed genes (DEGs) in this progenitor cluster between controls and

betaPix knockouts at 2 dpf, enriched gene ontology (GO) terms revealed three major

categories: epigenetic remodeling including chromatin silencing, negative regulation of

transcription, chromatin packaging and nucleosome positioning; microtubule organizations including microtubule depolymerization, regulation of microtubule polymerization and microtubule cytoskeleton organization; and neurotransmitter secretion/transportation (Figure 4D). Of these signaling genes, we were particularly interested in microtubule genes for further studies.

Stathmin acts downstream of *betaPix* in glial migration *via* regulating tubulin polymerization

Microtubules are essential cytoskeletal elements composed of α/β-tubulin heterodimers. The regulation of microtubule polymerization affects important cellular functions such as mitosis, motor transport, and migration [31]. Based on the above scRNA-seq data (Figures 4E and S6C), we then examined if microtubule polymerization genes stathmins were affected in *betaPix* mutants. We found that stathmin1a (*stmn1a*), stathmin1b (*stmn1b*), and stathmin4l (*stmn4l*) were relatively abundant in both the whole brain and the progenitor cluster, and decreased specifically at 2dpf in *betaPix* knockout progenitor cells (Figure 4E). Subsequent qRT-PCR revealed that these stathmins expression decreased in both global *betaPix* knockouts (*betaPix^{m/m}*) and glial-specific betaPix knockout glia (*gfap*:EGFP-Cre;*betaPix^{ct/ct}*) (Figure 4F and 4G), which was further verified by RNA *in situ* hybridization analysis (Figure 4H). We next investigated whether *Stathmins* act downstream to *betaPix* signaling on regulating vascular integrity development. To this end, we found that *Pak1/2* inhibition enabled to rescue down-regulated expressions of *stmn1a*, *1b* and *4l* in either global or glial-specific

betaPix knockout zebrafish (Figures 5A, 5B and S7A), supporting previous findings

that *stathmin-1* acts downstream to *betaPix-Pak* signaling in neurite outgrowth in

mice[32].

To test whether *Stathmin* family genes are important for vascular stability, we utilized

CRISPR-induced F_0 knockout system simultaneously targeting the three stathmin genes.

While singular stathmin gene knockout led to no evident phenotypes (data not shown),

we found that simultaneously inactivating *stmn1a*, *stmn1b* and *stmn4l* caused brain

hemorrhages and hydrocephalus in the brain, similar to that from *betaPix^{fn40a}* and

betaPix^{m/m} mutants (Figures 5C, 5D, S7B to S7E). By labeling both glia and vasculature

with Tg(*gfap*:GFP; *kdrl*:mCherrry) transgenic embryos, we found that triple stathmin

gene knockouts had impaired CtAs, abnormal glia and neuronal development in the

hindbrain region (Figure 5E, 5F, S7F to S7H). Furthermore, overexpressing *stmn1b* by

the *gfap* promoter partially rescued brain hemorrhages and delayed neuronal

development in *bbh^{fn40a}* mutants (Figures 5G, 5H, S7I and S7J), of which glial processes

elongation also improved, but CtAs defects failed to restore (Figures 5I, 5J and S7K).

Thus, these data support the critical role of stathmins in glia.

To examine the direct role of *betaPix-stathmin* in glia, we turned to use human

glioblastoma cell lines and manipulated the *betaPix* levels by siRNA knockdown or

transgenic overexpression. We found that *betaPix* knockdown attenuated glial

migration (Figures 5K, 5L, S7L to S7N), which is consistent with previous studies on

betaPix function in the endoderm [33], epithelial cells [34], neurons [35] and multiple

carcinoma cell lines including glioblastoma [36-38]. Either *betaPix* or *Stmn1*

overexpression led to a partial restoration of the impaired glial migration, suggesting

Stathmins as downstream effectors of *betaPix*. Furthermore, immunofluorescence

analysis revealed that *betaPix* knockdown led to abnormal accumulation of

microtubules at the cell periphery, whereas ectopic expression of *betaPix* and *Stathmin*

partially rescued these microtubule defects (Figures 5M and 5N). Together, these

results suggest that *betaPix* depletion disrupts microtubule dynamics and leads to

impaired glial development acting upstream to the *Pak-Stathmin* signaling,

subsequently disrupting vascular integrity and resulting in brain hemorrhages. While

the brain hemorrhages were only partially rescued and abnormal CtAs defects failed to

restore, indicating that additional pathways may act downstream of *betaPix*.

Zfhx3/4 acts downstream of *betaPix* in regulating vascular integrity development.

Previous studies have identified multiple signaling pathways important for zebrafish

hindbrain angiogenic sprouting, such as VEGF, Notch, Wnt, Integrin β1,

angiopoietin/TIE, and insulin-like growth factor signaling [39-42]. To investigate whether

angiogenic signal is disrupted by *betaPix* depletion, we examined several angiogenic

gene expression pattern by performing RNA *in situ* hybridization of embryos at 36 hpf.

The expression level of *Vegfaa* in sprouting CtAs decreased in *bbh*[fn40a] mutants (Figure

6A). Consistently, *betaPix* knockdown significantly reduced *VEGFA* expression in

cultured glial cells, which was rescued by ectopic expression of *betaPix* (Figure 6B;

right panel). To explore how *betaPix* regulate *Vegf* signaling in glia, we also found a

significant enrichment of epigenetic and transcriptional regulation in progenitor cluster

indicated by the GO analysis of single-cell sequencing data (Figure 4D). Among these

candidate genes, transcription factor Zinc Finger Homeobox 3 (ZFHX3) is known to

mediate hypoxia-induced angiogenesis in hepatocellular carcinoma *via* transcriptional

activation of VEGFA [43]. Furthermore, ZFHX3 has been associated with stroke in

multiple genome-wide association studies [44-46]. Interestingly, we found that

transcription factors *Zfhx3* and *Zfhx4* significantly downregulated after *betaPix*

inactivation in cultured glial cells (Figure 6B), CRISPR-induced *betaPix* mutants at 2

dpf (Figure S8A), and glia-specific *betaPix* mutants (Figure S8B), thus suggesting that

betaPix acts upstream to Zfhx3/4-VEGFA signaling in regulating angiogenesis.

To determine whether *Zfhx3/4* is critical for vascular integrity by interacting with

betaPix in zebrafish, we depleted *Zfhx3* and *Zfhx4* simultaneously *via* CRISPR-

mediated F_0 knockouts (Figure S8C to S8E). We found that *zfhx3/4* knockouts

decreased *Vegfaa* expression at sprouting CtAs at 36 hpf (Figure 6C). *Zfhx3/4*

knockouts at 48 hpf had under-developed CtAs and reduced penetration of endothelial

cells into the hindbrain (Figure 6F and 6G), developed cerebral hemorrhages during 36

to 52 hpf, which are comparable with *betaPix* mutant phenotypes (Figure 6D-6G).

Despite up-regulated expression level of *nestin*, glial arrangement and processes

elongation remained statistically unaffected after *Zfhx3/4* inactivation (Figures 6F, 6G

and S8F). We next investigated whether increasing expression level of *Vegfaa* in

betaPix mutants was able to rescue impaired vascular integrity. By using *Zfhx4* mRNA

microinjection into one-cell stage embryos in either *bbh*^{fn40a} mutants or CRISPR-

induced *betaPix* knockouts, we found that ectopic expression of *Zfhx4* increased *Vegfaa*

expression level (Figure 6L), thus leading to a drastic reduction of both percentages and

volumes of brain hemorrhage shown in these mutants (Figures 6H, 6I, S8G to S8I), as

well as improved CtAs and glial development (Figures 6J and 6K). In addition, scRNA-

seq data showed that CRISPR-induced *betaPix* knockouts had down-regulated

angiogenic gene expression in a cluster of endothelial cells (Figure 6M). Together, these

data support the critical role of *Zfhx3/4* in vascular integrity and glial development

probably acting upstream to VEGFA and downstream to betaPix.

Discussion

Conditional knockout technology in mice has been widely used to modify and

determine targeted gene function in a cell-specific manner. Spatio-temporal specific

modification is beneficial for studying embryonic lethality genes or investigating the

function of targeted genes in specific cell populations. Despite being broadly utilized

in mice, conditional knockouts in other animal models are limited due to technical

difficulties. To this end, multiple research groups have designed different strategies

for establishing floxed-mutant alleles, gene traps, and inducible Cas9 in zebrafish [47].

In particular, the Zwitch gene trap-based approach has been established successfully

in zebrafish, enabling to simultaneously produce knockin reporter driven by the

endogenous gene promoter and disrupting endogenous gene function [19-21]. We adapted

this strategy with the aid of CRISPR/Cas9 technology, and added a GSG spacer to

enhance cleavage efficiency and universal guide RNA target sites on both

homologous arms for producing linearized targeting vector. Moreover, we also found

that utilizing either short or long homologous arms enables to achieve precise targeted

integrations with this donor vector. Our short arms-based method can be broadly

adapted because of less restrictions on arm sequences and easily engineering the

donor vector. From the *betaPix* locus and other five unpublished loci that we already

generated Zwitch alleles with ~1% correct insertions based on homologous

recombination, we achieved Cre-induced, cell-specific deletion and functional

analysis of targeted genes in both embryos and adult hearts. Therefore, our CRISPR-

based Zwitch method can be widely applied for generating conditional mutant

zebrafish in particular and possibly extended for other animal mutants in general.

Development of the vasculature in the central nervous system can be classified into

three phases: vasculogenesis, angiogenesis and blood-brain barrier formation. The

hallmark of CNS vasculature maturation is acquisition of BBB properties,

establishing a stable environment crucial for brain homeostasis in response to

extrinsic factors and physiological changes. The core features of the BBB include

specialized tight junctions, highly selective transporters and limited immune cell

trafficking [48,49]. In zebrafish, BBB can be functionally characterized from 3 dpf [50,51].

While *bubblehead* mutants acquire vessel rapture phenotype early before BBB

maturation, presenting a good hemorrhagic model for studying early BBB

development. Multiple types of cells contribute to diverse aspects of BBB

development and maintenance, leading to proper vasculature function across

developmental timeline. Pericytes are perivascular cells embedded within the same

basement membrane with endothelial cells [52]. Pericyte-deficient mice with disrupting

PDGF-B signaling displayed BBB leakiness, reflecting altered pericyte coverage and

endothelial gene expressions [53-55]. Pericyte-secreted signals promoting the endothelial

survival, proliferation and sprout formation *via* multiple pathways such as

transforming growth factor-β (TGF-β) signaling [56-59], VEGF-VEGFR signaling [60,61],

Angiopoietin-Tie2 signaling [62-64], Sphingosine-1-phosphate (S1P) signaling [65] and

Notch signaling [66,67]. Apart from endothelial cells, pericytes also regulate

development of other members of the neurovascular unit. Pericytes regulate astrocyte

aquaporin-4 polarization and guide end-foot processes toward endothelial tubes [53,68].

In addition, pericytes secrete ECM molecules supporting the basement membrane and

regulation of the angioarchitecture [69-72].

Endothelial cells form the continuous inner layer of blood vessels with junctional

protein and selective transporters, which are the main subject of permeability

regulation in BBB. It is well accepted that endothelial cells do not show a

predetermined role and that the brain environment is sufficient to induce endothelial

barrier [73]. Accumulative evidences also point to critical role of endothelial signals in

regulating several aspects of BBB development. In particular, endothelial cell-derived

PDGF-B directly binds to surface receptor PDGFRβ on pericytes that are critical for

pericyte recruitment and function [74,75]. Although their barrier properties are non-

assessed, angiocrine signals such as leukaemia inhibitory factor (LIF) and TGF-β

induce both astrocyte [76] and oligodendrocyte differentiation [77], respectively. Wnt7a/b

ligands secreted by neural progenitor cells bind to Frizzled receptors on endothelial

cells, activating canonical Wnt signaling and downstream genes. WNT ligands

knockouts lead to angiogenic defects and BBB leakages specifically in the CNS [78].

Endothelial GPR124, as a coactivator of canonical Wnt signaling, has important

function in CNS-specific angiogenesis and BBB establishment [79-81]. Other signaling

ligands such as angiopoietins and Ephrin family ligands orchestrate with Tie or EphB

receptors on endothelial cells, respectively[82]. Briefly, complex interactions between

BBB cell types orchestrate proper angiogenesis and CNS development in a spatio-

temporal manner.

Compared with pericytes and endothelial cell signaling, we have less knowledge on

glia signaling in the BBB development and maturation. Astrocytes ensheathe

capillaries through polarized end feet that are enriched with aquaporin-4 (Aqp4)

proteins, colocalized with inwardly rectified K^+ channels[83,84]. Functional coupling of

ion and water fluxes plays a critical role in regulating local osmotic

equilibrium. During development, radial glia is a neuroepithelial origin with

heterogeneous populations that are able to generate neurons, astrocytes, and

oligodendrocytes[85-88]. In zebrafish, radial glia has long been considered serving an

astrocytic role. Radial glia in zebrafish enriches with several key molecular markers

for astrocytes and tight junctions while Aqp4[+] radial glial processes rarely contact the

vasculature[89]. A more recent article has reported that zebrafish *Glast*-expressing radial

glia transform into astrocyte-like cells, displaying dense cellular processes, tiling

behavior and proximity to synapses [90]. Whether end feet of these astrocytes enwrap

capillary blood vessels and resemble mammalian astrocytes warranty future

investigations. In agreement with mammalian astrocytes, two independent groups

have shown that ablation of pan-glia results in progressive brain hemorrhage[91,92] or

deficiency of spinal cord arteries development in zebrafish [93], demonstrating

important role of zebrafish glia in BBB formation. Multiple glia/astrocyte-derived

signals have been shown to contribute to endothelial barrier properties development.

Astrocyte-expressed sonic hedgehog (Shh) bind to hedgehog (Hh) receptors on

endothelial cells and contribute to the BBB functions by promoting junctional protein

expression and the quiescence of the immune system [94-97]. Before BBB formation,

notochord-derived Shh activity promote arterial cell fate on developing endothelial

cells [98]. Shh stimulates the production of angiogenic cytokines, angiopoietins and

interleukins, *via* downstream transcription factor *Gli* or non-canonical pathways

crosstalks with iNOS/Netrin-1/PKC, RhoA/Rock, ERK/MAPK, PI3K/Akt, Wnt/β-

catenin, and Notch signaling pathways [99]. VEGF is the main pro-angiogenic growth

factor produced by various cell types that controls multiple vascular development

steps, such as proliferation, migration, permeability and survival *via* interacting with

VEGF receptors on endothelial cells. Inactivation of VEGFA in astrocytes leads to

BBB disruption in multiple sclerosis models [100,101]. In early development of avascular

neural tube, VEGF derived from neuroectoderm serve as a central role in inducing tip

cells sprouting from the perineural vascular plexus (PVNP) and formation of

vasculature lumens from stem cells [102,103]. Ablating neuroglia reduces *vegfab* signals

and leads to ectopic intersegmental vessels sprouting and vertebral arteries in

zebrafish [93]. VEGF signaling orchestrates elaborately with other signaling pathways

such as delta-like 4 (Dll4)/Notch signaling [104] and activates downstream components

including the Ras/Raf/MEK, PI3K/AKT, and p38/MAPK/HSP27 pathways [105]. Our

data suggest glial *betaPix* as an important factor in vascular integrity development.

Together with previous studies, *betaPix-Pak1-stathmin* signaling regulates glial

growth and differentiation[32] and betaPix regulates *Vegfaa* secretion *via* downstream

transcription factors *Zfhx3/4*. Thus, this work presents a novel glial betaPix signaling

in regulating zebrafish BBB development.

Basement membrane is a non-cellular component consisting of several extracellular

matrix (ECM) proteins, which laminins are the most abundant ones, while collagen

IV, fibronectin, and perlecan serve as hub for structural supports, anchoring and

intercellular communication. Components of basement membrane are synthesized in

surrounding cells with diverse expression patterns [106]. For example, mice with

deletion of Collagen IV in either brain microvascular endothelial cells (BMEC)- or

pericytes lead to fully penetrant intracerebral haemorrhage, while COL4A1 deletion

in astrocytes results in mild haemorrhage phenotypes [69]. Differential cell-specific

expression pattern of laminin isoforms contribute differently to BBB formation [107]. As

astrocytic laminins mediate pericyte differentiation and regulate capillary

permeability [108], intercellular interactions with the ECM occur by cell surface integrin

receptors. Integrin β1 on endothelial cells are critical during angiogenesis[109]. On the

other hand, integrin αvβ8 has glial-specific enrichment which mediate focal adhesion

to modulate vascular integrity[17,110,111]. Notably, degradation of ECM and focal

adhesions at the perivascular space is responsible for the hemorrhagic phenotypes of

bbh mutants [17,18]. It would be of great interest to determine whether glial *betaPix*

account for ECM breakdown in *bbh* mutants in future studies.

Acknowledgements

The authors thank Drs. Bo Zhang, Jiulin Du, Liangyi Chen, Kazu Kikuchi, and Jian

Chen for providing fish lines, plasmid clones, and human glial cell lines; the members

of Dr. Jing-Wei Xiong's laboratory for helpful discussions and technical assistance;

and the National Center for Protein Sciences at Peking University, particularly Dr.

Liying Du at the Flow Cytometry Core for technical help on the Beckman Coulter

MoFlo XDP; and Dr. Hua Liang at National Biomedical Imaging Center, Peking

University for assistance with the Luxendo Multi-View Selective-Plane Illumination

Microscopy (Bruker). This work is supported by grants from the National Key R&D

Program of China (2019YFA0801602 and 2023YFA1800600); the National Natural

Science Foundation of China (32230032 and 31730061).

Declaration of Interest

The authors declare no competing financial interests.

Materials and Methods

Fish maintenance

Zebrafish were raised and handled in accordance with the guidelines of the Peking University Animal Care and Use Committee accredited by the AAALAC.

The bbh^{fn40a} mutant was isolated from a large-scale mutagenesis screen of the zebrafish genome at Massachusetts General Hospital, Boston[9]. Tg(*acta2:GFP-Cre*) was purchased from Xinjia Pharmaceutical Technology Co., Ltd (Nanjing, China); Tg(*neurod:EGFP*) were kindly provided by Dr. Bo Zhang (Peking University, China); Tg(*huC:GFP*) were kindly provided by Dr. Liangyi Chen (Peking University, China); Tg(*kdrl:GFP*) [112] and Tg(*kdrl:mCherry*) [113] were described previously.

Cell culture and transfection

U251 and A172 human glioblastoma cell lines, kindly provided by Dr. Jian Chen at Chinese Institute for Brain Research, Beijing, were cultured in high glucose DMEM medium (Hyclone) supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin, at 37°C in 5% CO_2 humidified incubator. The *betaPix* siRNAs and *STMN1* siRNAs were purchased from GenePharma (Shanghai, China). The sequences of the siRNAs are listed in **Table S2**. Transfection was conducted with Lipofectamine 3000 (Invitrogen) at a final siRNA concentration of 30 nM.

mRNA/gRNA synthesis and microinjections

mRNA and gRNA were synthesized as described previously [114]. In brief, linearized

pT3TS-nls-zCas9-nls [115], pXT7-*Cre*, and pXT7-*Zfhx4* plasmid DNA were purified as

templates using TIANquick Mini Purification Kit (TIANGEN, DP203, China). Next,

in vitro transcription reactions were performed using the mMESSAGE mMACHINE

T3 (for pT3TS-nls-zCas9-nls) or T7 (for pXT7-*Cre* and pXT7-*Zfhx4*) kits (Life

Technologies) according to manufacturer's instructions. Templates for gRNA were

generated by complementary annealing and elongation of two oligos. Forward oligo

contained a T7 promoter and gRNA target sequence, and reverse oligo contained the

universal gRNA scaffold [23]. The resulting double-stranded DNA served as the templates

for *in vitro* transcription using HiScribe® T7 High Yield RNA Synthesis Kit (NEB).

The sequences of the gRNAs are listed in **Table S3**. For global *betaPix* inactivation,

Cre mRNA with a working concentration of 200 ng/µL were injected into the *betaPix*$^{ct/ct}$

embryos at one-cell stage. For CRISPR-mediated knockouts, 300 ng/µL Cas9 mRNA

and 20-50 ng/µL gRNAs were injected into zebrafish embryos at one-cell stage. For

Zfhx4 rescue experiments, *Zfhx4* mRNA with a working concentration of 400 ng/µL

were injected into the *bbh*^{fn40a} mutants or in combination with CRISPR-mediated

betaPix knockout system at one-cell stage. Injected embryos were scored for mutant

phenotypes at 36 or 48 hpf.

Construction of *betaPix* conditional knockout zebrafish

The pZwitch plasmid clone was kindly provided by Dr. Kazu Kikuchi (National

Cerebral and Cardiovascular Center Research Institute, Suita, Japan). GSG spacer was

added to pZwitch+3 between P2A and TagRFP coding sequences by overlapping primer

pairs. Next, a highly efficient gRNA was found from the fifth intron of *betaPix*. Primer

pairs were designed to amplify the left and right homologous arms from the gRNA site

for 1000 bp and 24 bp with following modifications: 1) The 5' end of the forward primer

of the left homologous arm were added with an *Nhe*1 site and a universal gRNA site

which provided an *in vivo* linearization cleavage site; a *Mlu*1 site to the 5' end of the

reverse primer of the left arm. 2) Added *EcoR*1 site to the 5' end of the forward primer

of the right homologous arm; added *Xho1* site and universal gRNA site to the 5' end of

the reverse primer of the right homologous arm. Then we cloned the inverted

homologous arms into polyclonal sites of the modified pZwitch+3, and purified plasmid

DNA with EndoFree Mini Plasmid Kit (Tiangen, DP118). *In vivo* knock-in system was

composed of 300 ng/μL Cas9 mRNA, 50 ng/μL *betaPix*-intron5 gRNA, 50 ng/μL

Universal gRNA, and 20 ng/μL donor vector. Microinjected *in vivo* knock-in system

into wildtype embryos at one-cell stage. Fish founders were screened for *α*-crystallin

reporter expression, raised to adulthood, and re-screened for germline transmission and

precise knock-in by Sanger sequencing.

Construction of transgenic reporters

The glial reporter plasmid *gfap*-EGFP [26] was kindly provided by Dr Jiulin Du (Shanghai

Institute for Biological Sciences, Shanghai, China). The *gfap* promoter was used to

establish Tg(*gfap:GFP-Cre*) plasmid. The *stmn1b* coding sequences were amplified

from wild-type zebrafish cDNA library, which were used to establish Tg(*gfap:GFP-*

stmn1b) plasmid.

Transgenic reporters were generated by using *Tol2*-based transgenesis [116]. In brief, 100

ng/µL *Tol2* transposase mRNA in combination with 20 ng/µL donor plasmid DNA were

co-injected into the embryos at one-cell stage. Transgenic founders were screened

for specific transgenic GFP expression, raised to adulthood, and re-screened for

germline transmission.

Inhibitor treatment

IPA-3 (Proteintech, CM05727) were dissolved in DMSO to form a 10 mM stock, and

then diluted with E3 medium to 3 µM in 6-well plates. Zebrafish embryos were treated

from 24 to 48 hpf, and then washed with E3 medium for phenotype analysis.

***o*-dianisidine staining**

o-dianisidine staining was performed as described previously [18]. In brief, embryos were

dechorionated, anesthetized and incubated in fresh staining solution (0.6 mg/mL *o*-

dianisidine, 0.01 M sodium acetate pH4.5, 0.65% H_2O_2 and 40% Ethanol) for 20

minutes in dark. Washed three times with methanol, and performed benzyl

alcohol/benzyl benzoate (BABB) tissue clearing before imaging with stereo

microscope (Leica, 160F).

Whole-mount *in situ* hybridization (WISH)

Whole-mount *in situ* RNA hybridization was performed as described previously [117].

Antisense probes were synthesized using a digoxigenin RNA labeling kit (Roche,

11277073910). Primer sequences for all WISH probes used in this paper are provided

in **Table S4**.

Quantitative real-time RT-PCR

Total RNA was isolated from embryos using Trizol (invitrogen) and cDNA was

generated using HiScript® III RT SuperMix (Vazyme) according to the manufacturer's

instruction. Quantitative real-time PCR was performed using Lightcycler (Roche) and

ChamQ SYBR qPCR Master Mix (Vazyme). Primer sequences are listed in

Supplementary Materials **Table S5**. Gene expression level was normalized against

GAPDH level.

Light-sheet fluorescence microscopy imaging

Light-sheet microscopy imaging was performed as described previously [118]. In brief,

transgenic zebrafish embryos were collected and maintained at 28.5°C. At the required

developmental stages, the embryos were carefully dechorionated, paralyzed with

tricaine, and transferred into 1% ultrapure low melting point agarose (16520-050;

Invitrogen). The embryo was then drawn into a glass tube in top-down position using a

1 mL syringe with an 18G blunt needle. After agarose coagulation, a wire was inserted

from bottom to push the agarose with embryo upwards, removed excess agar until the

head area is exposed from the top of the glass tube. Finally, the glass tube was fixed in

a sample holder for subsequent imaging.

Imaging was carried out with Luxendo Multi-View Selective-Plane Illumination

Microscopy (Bruker). Optical calibrations were adjusted according to the instruction

manual.

The optical plates are emitted from two Nikon CFI Plan Fluor 10x W 0.3 NA immersion

objectives in opposite directions. The detection is completed by two Olympus 20x 1.0

NA immersion lenses. The parameters are set as follows: the green fluorescence channel

uses a laser of 488 nm coupled with BP497-554 filter, the red fluorescent channel uses

laser 561 nm coupled with BP580-627 filter, with an exposure time of 100 ms, a delay

of 11 ms, a line mode of 50 px, and z-Stack interval of 3 μm. The raw data were stored

in H5 file format, processed into TIFF format and merged Multi-View dataset using

MATLAB software. After pre-processing, multi-fluorescence channels were merged in

ImageJ, and 3D visualization and measurement were performed by Imaris.

Single-cell RNA sequencing

For knockout samples, 300 ng/μL Cas9 mRNA and a mixture of number 1 to 4 *betaPix*

gRNAs with 50 ng/μL each were co-injected into wildtype embryos at one-cell stage.

Injection with equivalent concentration of Cas9 mRNA with PBS served as siblings.

After microinjection, embryos were collected and maintained at 28.5°C. At 24 or 48

hpf, zebrafish were dechorionated, paralyzed and transferred onto agarose plate. Heads

were harvested with dissecting scissors in cold sterile 1X PBS with pooling 200 heads

for each group. Heads were then dissociated in 900 μL Accutase cell detachment

solution (Sigma-Aldrich, A6964) at 28.5°C for 3 hours and re-suspended by pipetting

every 30 min. Once digestion was complete, 100 μL FBS was added to cell suspension

and centrifuged at 4℃, 500 g for 3 min. The cells were gently re-suspended in cold 500

μL 2.5% FBS in 1X PBS and filtered through a 40 μm strainer. PI and Hoechst33342

were stained for distinguishing living cells. The single, living cells were sorted by Aria

SORP (BD biosciences) into 1.5 ml tubes. The cell counts and vitality were verified by

AOPI staining coupled with an Automated Cell Counter (Countstar BioTech). Single

cells were barcoded with Chromium Next GEM Single Cell 3 'Reagent Kits V3.1 kit

(10x Genomics, 1000269) in 10× Chromium Controller (10× Genomics). After

qualified by peak shapes, fragment size and tailing with Fragment Analyzer System kit

(Agilent Technologies, DNF-915), single-cell transcriptome libraries were sequenced

via Illumina High Throughput Sequencing PE150 (Novogene, Beijing). The sequencing

data were analyzed using the CellRanger-6.1.1 (10x Genomics) and mapped to

reference genome GRCz11-GRCz11.103.

Cells were excluded from subsequent analyses under the following conditions: when

the number of expressed genes was less than 500, when there were abnormally high

counts of UMIs or genes (outliers of a normal distribution), or when the mitochondrial

content exceeded 9%. Unsupervised clustering was performed using Seurat (version:

4.0.2) with a resolution of 2.5, resulted in 71 cell populations and further annotated into

24 zebrafish major cranial cell types. Differential expression analysis in Seurat v4 was

used to identify cluster/cell type markers by Wilcoxon rank sum test.

Scratch assay

U251 cells were planted into culture plate and performed transfection at the desired

density. At 24 hours post-transfection, we vertically scratched monolayer cells by using a 200 μL pipette tip. Washed the cells three times with PBS, then replaced with serum-free culture medium for further culture. Stereo fluorescence microscopy (Leica, 160F) was used to document the scratch size at 0 hours, 18 and 24 hours. Images were processed by ImageJ.

Immunostaining

For assessing tubulin expression, U251 cells were planted into chamber slides (Saining, 1093000) and transfected at the desired density. At 24 hours post-transfection, washed the cells once on ice with pre-cooled PBS, then fixed with pre-cooled methanol at -20℃ for 20 minutes. The fixed cells were permeabilized with pre-cooled acetone at -20℃ for 1 minute. Removed acetone, incubated 0.5% BSA/PBS blocking solution at room temperature for 15 minutes. Cells were then incubated with 1:200 anti-α-tubulin mouse monoclonal antibody (EASYBIO, BE0031) at 37℃ for 45 minutes. After rinsing once with PBS, cells were incubated with 1:400 Alexa Fluor™ 488 goat anti-mouse IgG (H+L) secondary antibody (Invitrogen, A11029) at 37℃ for another 45 minutes. Washed three times with PBS, and sealed with Mounting Medium with DAPI (ZSGB Bio, ZLI-9557). Images were acquired by upright fluorescence microscopy (Leica, DM5000B) at 40× magnification, and processed by ImageJ.

Statistical analysis

Statistical analysis was performed using Graphpad Prism 6. The statistical significance

661   of differences between the two groups was determined by the independent unpaired

662   Student's t-test. Among three or more groups, one-way ANOVA analysis coupled with

663   Dunnett's test were used. All data are presented as the mean ± SEM. *P* value <0.05

664   indicates significant, with individual *P* values mentioned in the figure/figure legends.

 Figure Legends

Figure 1. Generation of *betaPix* conditional trap (*betaPix^{ct}*) allele by an

homologous recombination (HDR)-mediated knock-in method.

(A) Schematic diagram illustrates the HDR-mediated *Zwitch* strategy for

generating zebrafish knock-in allele at the *betaPix* locus.

(B) Genomic PCR analysis of the F_1 embryos confirming the right *Zwitch*

insertions.

(C) Sanger sequencing confirming the junction of *betaPix^{ct}* (after HDR-mediated

insertion) or *betaPix^{m}* (after Cre-mediated inversion) that are highlighted by

RED lines in (A).

(D) Representative stereomicroscopy images of erythrocytes stained with o-

dianisidine in *betaPix^{ct/ct}* and *betaPix^{m/m}* embryos at 48 hpf. Brain

hemorrhages, indicated with arrows, in Cre mRNA-injected embryos

(*betaPix^{m/m}*). Lateral views, anterior to the left.

(E) Quantification of hemorrhagic parameters in (D). Left panel showing

hemorrhage percentages with each dot representing an individual experiment

(n>100 each), and right panel showing hemorrhage areas with each dot

representing one embryo.

(F) qRT-PCR analysis showing the expression of *betaPix*, *RFP* and *alphaPix* in

betaPix^{ct/ct}, *betaPix^{ct/m}*, and *betaPix^{m/m}* embryos at 48 hpf. Each dot represents

one embryo. cryaa, αA-crystallin; PA, polyadenylation signal; SA, splice

acceptor; T2A, T2A self-cleaving peptide.

Figure 2. *betaPix^{m/m}* mutant have brain hemorrhages, central artery defects and abnormal glial structure that was partially rescued by Pak1 inhibitor IPA3 treatment.

(A) Left panel showing the maximum intensity projection of the glial structures in the hindbrain of *betaPix^{ct/ct}* and *betaPix^{m/m}* embryos at 48 hpf. Lateral view, anterior to left. Middle panel showing representative optical sections and right panel showing the higher magnifications of boxed area, presenting atypical glial structures with disoriented arrangements (yellow arrows) in *betaPix^{m/m}* embryos.

(B) Quantification of glial parameters in (A). Left panel showing the average glia length, and right panel showing glia length index normalized to individual head length, which each dot represents one embryo.

(C) Whole-mount RNA *in situ* hybridization revealed *nestin* and *pax2a* expression pattern in *betaPix^{ct/ct}* and *betaPix^{m/m}* embryos at 48 hpf. Dorsal view, anterior to the left.

(D) Optical sections of glial structure (green) and blood vessels (magenta) in the heads of siblings and CRISPR-mediated *betaPix* F_0 knockout embryos. Arteries in the hindbrain of *betaPix* KO mutants had developmental defects (white arrowheads), showing shorter distance between basilar artery (BA) and glial cell bodies.

(E) 3D reconstruction of the sox2-positive precursors (green) and vasculatures (magenta) in the heads of siblings and CRISPR-mediated *betaPix* F_0 knockout

embryos at 48 hpf. Box areas are shown in higher magnifications at the middle panels, with optical sections shown in the right panels. Arrows indicate CtA with enlarged perivascular space.

(F) Representative stereomicroscopy images of *o*-dianisidine staining of *betaPix^{ct/ct}* and *betaPix^{m/m}* embryos at 48 hpf that were treated with DMSO or PAK inhibitor IPA3. Brain hemorrhages indicated with the arrows.

(G) Quantification of brain hemorrhagic parameters in (F).

(H) 3D reconstruction of the glial structure (green) and vasculature (magenta) in the heads of *betaPix^{ct/ct}* and *betaPix^{m/m}* embryos at 48 hpf treated with DMSO or IPA3. Lateral view, anterior to left. Defects in CtAs (white arrows) and glia (yellow arrows) shown in *betaPix^{m/m}* embryos treated with DMSO or IPA3.

(I) Quantification of CtA parameters in (H).

(J) Quantification of glia parameters in (H).

Figure 3. Glial-specific *betaPix* knockouts recapture global *betaPix* mutant phenotypes.

(A) Representative stereomicroscopy images of erythrocytes stained with *o*-dianisidine in *betaPix^{ct/ct}* siblings and *gfap:GFP-Cre*; *betaPix^{ct/ct}* mutant embryos treated with DMSO or IPA3 at 48 hpf. Brain hemorrhages indicated with arrows in glial-specific *betaPix* knockouts. Lateral view with anterior to the left.

(B) 3D reconstruction of the vasculatures (magenta) in the heads at 48 hpf,

showing lateral view with anterior to the left. The box areas are shown in

higher magnifications of brain vasculatures at the right panels. CtA defects

(yellow arrows) were evident in *gfap:GFP-Cre*; *betaPix^{ct/ct}* mutant embryos.

(C) Quantification of brain hemorrhages (A) and CtA parameters (B). Left panel

showing hemorrhage percentages with each dot representing an individual

experiment (n>100 each). Middle panel showing hemorrhage areas with each

dot representing one embryo. Right panel showing CtA length index

normalized to individual head length, with each dot representing one embryo.

Figure 4. Single cell transcriptome reveals that a subcluster of glial progenitor and

stathmin family members are associated with *betaPix* mutation.

(A) Experimental strategy for single cell RNA sequencing of embryonic heads

from wild-type siblings and *betaPix* CRISPR mutants at 1 dpf and 2 dpf.

(B) UMAP visualization and clustering of cells labeled by cell type. Four samples

were aggregated and analyzed together.

(C) Proportions of 24 cell clusters were differentially distributed among four

sample groups. ctrl_1d, PBS-injected siblings at 1 dpf; ko_1d, *betaPix*

CRISPR mutants at 1 dpf; ctrl_2d, PBS-injected siblings at 2 dpf; ko_2d,

betaPix CRISPR mutants at 2 dpf.

(D) Enriched GO terms for differentially expressed genes for progenitor sub-

cluster comparing ko_2d to ctrl_2d groups.

(E) Violin plots showing the expression of the stathmin family genes by all cells

among four sample groups (left panel) or by progenitor sub-cluster among

four sample groups (right panel).

(F) qRT-PCR analysis showing expression of *betaPix*, *stmn1a*, *stmn1b*, *stmn4l* and

ppfia3 in glia of FACS-sorted *betaPix*$^{ct/ct}$ siblings and *betaPix*$^{m/m}$ mutants at 48

hpf. Each dot represents cells sorted from one embryo.

(G) qRT-PCR analysis showing expression of *betaPix*, *stmn1a*, *stmn1b*, *stmn4l* and

ppfia3 in glia of FACS-sorted *betaPix*$^{ct/ct}$ siblings and *gfap:GFP-Cre*;

betaPix$^{ct/ct}$ mutants at 48 hpf. Each dot represents cells sorted from one

embryo.

(H) Whole-mount RNA *in situ* hybridization revealing down-regulation of *stmn1a*,

stmn1b and *stmn4l* in *betaPix*$^{ct/ct}$ siblings and *gfap:GFP-Cre*; *betaPix*$^{ct/ct}$

mutant embryos at 48 hpf.

Figure 5. Stathmin acts downstream of *betaPix* in glial migration *via* regulating

tubulin polymerization.

(A) Whole-mount RNA *in situ* hybridization showing that *stmn1a, stmn1b and*

stmn4l expression in *gfap:GFP-Cre*; *betaPix*$^{ct/ct}$ embryos were partially

rescued by Pak1 inhibitor IPA3 treatment at 48 hpf. Dorsal views, anterior to

the left.

773  (B) qRT-PCR analysis showing that *stmn1a* and *stmn4l* expression were rescued in

774    *gfap:GFP-Cre*; *betaPix$^{ct/ct}$* mutants by IPA3 treatment at 48 hpf. Each dot

775    represents one embryo.

776  (C) Representative stereomicroscopy images of erythrocytes stained with *o*-

777    dianisidine in siblings and *stmn1a/1b/4l* CRISPR mutants at 48 hpf. Brain

778    hemorrhages, indicated with arrows, appeared in *stmn1a/1b/4l* mutants.

779    Lateral views with anterior to the left.

780  (D) Quantification of hemorrhagic parameters in (C). Left panel showing

781    hemorrhage percentages with each dot representing an individual experiment

782    (n>100 each), and right panel showing hemorrhage areas with each dot

783    representing one embryo.

784  (E) 3D reconstruction of glial structure (green) and vasculature (magenta) in the

785    heads at 48 hpf. Lateral view with anterior to the left. CtA defects (white

786    arrows) appeared in *stmn1a/1b/4l* mutants.

787  (F) Quantification of CtA and glia parameters in (E). Length index normalized to

788    individual head length, with each dot representing one embryo.

789  (G) Representative stereomicroscopy images of erythrocytes stained with *o*-

790    dianisidine in *bbh^{fn40a}* and *bbh^{fn40a}*; Tg(*gfap:GFP-stmn1b*) embryos at 48 hpf.

791    Brain hemorrhages, indicated with arrows, decreased in *bbh^{fn40a}* mutants with

792    glia-specific overexpression of *stmn1b*, compared with *bbh^{fn40a}* mutant

793    siblings. Lateral views with anterior to the left.

(H) Quantification of hemorrhagic parameters of (G). Left panel showing hemorrhage percentages with each dot representing an individual experiment (n>100 each), and right panel showing hemorrhage area with each dot representing one embryo.

(I) 3D reconstruction of the *gfap:GFP-stmn1b* overexpression (green) and vasculature (magenta) in the heads of *bbh^{fn40a}* siblings and *bbh^{fn40a}*; Tg(*gfap:GFP-stmn1b*) mutants at 48 hpf. Lateral view with anterior to left. White arrows indicate CtA defects.

(J) Quantification of CtA and glia parameters in (I). Length index normalized to individual head length, with each dot representing one embryo.

(K) Representative stereomicroscopy images of U251 cells at 0 and 24 hours after wounding. U251 cells were transfected with negative control siRNA or *betaPIX* siRNA separately, in combination with pcDNA3.1 vector, *betaPIX* overexpression plasmid, and *STMN1* overexpression plasmid. The wound edges are highlighted by dashed lines, with arrow lines indicating the wound width.

(L) Quantification of wound closure in (K), showing $^*P<0.05$, $^{***}P<0.005$ compared to negative control siRNA with empty vector transfection. $^{\#\#}P<0.01$, $^{\#\#\#}P<0.005$ compared to *betaPix* knockdown with empty vector transfection.

(M) Representative immunofluorescence image of alpha-tubulin and DAPI signals in U251 cells. U251 cells were transfected with negative control siRNA or *betaPIX* siRNA separately, in combination with pcDNA3.1 vector, *betaPIX* overexpression plasmid, and *STMN1* overexpression plasmid. Box

areas are shown in higher magnifications. Arrows indicate protrusions at the cell periphery.

(N) Quantification of cell percentages with protrusions in (M). [***]$P<0.005$ compared to negative control siRNA with empty vector transfection. [#]$P<0.05$, [##]$P<0.01$ compared to *betaPix* knockdown with empty vector transfection.

Figure 6. *Zfhx3/4* acts downstream of *betaPix* to regulate vascular integrity development.

(A) Whole-mount RNA *in situ* hybridization revealing that *vegfaa* decreased in *bbh*[fn40a] mutants compared with siblings at 36 hpf. Lateral views with anterior to the left. Arrows indicate *vegfaa* expression in the CtAs.

(B) qRT-PCR analysis revealing that *ZFHX3*, *ZFHX4 and* VEGFA decreased in U251 cells transfected with *betaPIX* siRNA, which were rescued by *betaPIX* overexpression.

(C) Whole-mount RNA *in situ* hybridization showing that *vegfaa* decreased in CRISPR-mediated *zfhx3/4* F$_0$ knockout embryos at 36 hpf. Lateral views with anterior to the left. Arrows indicate *vegfaa* expression in the CtAs.

(D) Representative stereomicroscopy images of erythrocytes stained with *o-dianisidine* in siblings and CRISPR-mediated *zfhx3/4* F$_0$ knockout embryos at 48 hpf. Arrows indicated brain hemorrhages in the knockout brains. Lateral views with anterior to the left.

(E) Quantification of hemorrhagic parameters in (D). Left panel showing
hemorrhage percentages with each dot representing an individual experiment
(n>100 each), and right panel showing hemorrhage areas with each dot
representing one embryo.

(F) 3D reconstruction of the glial structure (green) and vasculature (magenta) in
the hindbrains of siblings and CRISPR-mediated *zfhx3/4* F_0 knockouts at 48
hpf. Lateral view with anterior to the left. White arrows indicate CtA defects
in *zfhx3/4* knockout embryos.

(G) Quantification of CtA and glia length parameters of (F). Length index
normalized to individual head length, with each dot representing one embryo.

(H) Representative stereomicroscopy images of erythrocytes stained with *o-*
dianisidine in CRISPR-mediated *betaPix* F0 knockout embryos with or
without *zfhx4* mRNA injection at 48 hpf. Arrows indicate brain hemorrhages.
Lateral views with anterior to the left.

(I) Quantification of hemorrhagic parameters in (H). Left panel showing
hemorrhage percentages with each dot representing an individual experiment
(n>100 each), and right panel showing hemorrhage area with each dot
representing one embryo.

(J) 3D reconstruction of the glial structure (green) and vasculature (magenta) in
CRISPR-mediated *betaPix* F_0 knockout embryos with or without *zfhx4* mRNA
injection at 48 hpf. White arrows indicate CtAs and yellow arrows indicate
glia.

(K) Quantification of CtA and glia length parameters in (J). Length index

normalized to individual head length, with each dot representing one embryo.

(L) Whole-mount RNA *in situ* hybridization revealed that *zfhx4* and *vegfaa*

decreased in CRISPR-mediated *betaPix* F$_0$ knockout embryos at 36 hpf, which

were rescued by *zfhx4* mRNA injection. Lateral views with anterior to the left.

Arrows indicate hindbrain regions.

(M) Dot plots of several angiogenesis-associated genes expression in endothelial

cell cluster. Dot size indicates the percentage of cells with gene expression, and

dot color represents the average gene expression level.

Figure 7. Working Model on the function of glial *betaPix* in zebrafish vascular

integrity development of the hindbrain. *betaPix* is enriched in glia and regulates the

PAK1-Stathmin axis on microtublin stabilization, thus fine-tuning glial cell migration

and their interactions with vanscular endothelial cells; and in paralelle, betaPix may

also regulate Zfhx3/4-Vegfaa signaling in glia, which then modulates angiogenesis

during cerebral vessel development and maturation. Deletion of *betaPix* affects glial

cell migration and interaction with cerebral endothelial cells. MT, Microtubule.

Figure S1. Generation of betaPix conditional trap (*betaPix^{ct}*) allele by an HDR-

mediated knock-in method.

(A) Sanger sequencing results of the original pZwitch (+3) and the modified

vector with glycine-serine-glycine spacer (GSG) in front of the 2A peptide.

(B) Sanger sequencing results of the genomic PCR products at both junction sites

from the *betaPix^ct* F$_1$ offspring showing the correct knock-in in the betaPix

locus.

(C) The screening process for *betaPix^ct* knock-in zebrafish showing about 1%

efficiency with correct homologous recombinations.

(D) Representative stereomicroscopy images of *o*-dianisidine staining in *bbh^fn40a*

embryos at 48 hpf. Lateral view, anterior to the left in left panel; Dorsal view,

anterior to the top in right panel.

(E) Quantification of hemorrhagic parameters in (D). Left panel showing

hemorrhage percentages with each dot representing an individual experiment

(n>100 each), and right panel showing hemorrhage areas with each dot

representing one embryo.

(F) 3D reconstruction of the *betaPix* expression (magenta) in *betaPix^ct/+* brains of

embryos microinjected with PBS or *Cre* mRNA at 24 hpf, 48 hpf, 72 hpf and

144 hpf. Cre-induced *RFP* expression indicates successful inversion of gene

trap system (arrows). Dorsal view, anterior to the top.

(G) 3D reconstruction of the *neurod1* transgenic expression (green) and *betaPix*

expression (magenta) in *betaPix^ct/+* brains of embryos microinjected with PBS

or *Cre* mRNA at 24 hpf, 48 hpf, 72 hpf and 144 hpf. Cre-induced *RFP*

expression indicates successful inversion of gene trap system (arrows),

marking strong *betaPix* expression at the midbrain hindbrain boundary

(MHB), and milder expression at hindbrain at early stages. At 144 hpf, *betaPix*

expresses ubiquitously in the brain, outlining the cerebellum (arrowheads).

Dorsal view, anterior to the top.

Figure S2. *betaPix^{m/m}* mutant had brain hemorrhages, central arteries defects, and abnormal glial structure.

(A) 3D reconstruction of the vasculature (green) in the heads with lateral view, anterior to left. Endogenous *betaPix* expression (magenta) only shown in Cre mRNA-injected mutant embryos. The box areas are shown in higher magnifications of vasculatures at the right panels. Central arteries (CtA) defects (white arrows) were evident in *betaPix^{m/m}* embryos at 48 hpf. Arrow lines indicating the CtA length.

(B) Quantification of CtA parameters in (A). Left panel showing the average CtA length, and right panel showing the CtA length index normalized to individual head length, with each dot representing one embryo.

(C) 3D reconstruction of the glia structure. Box areas showing higher magnifications of glia at the right panels. D, dorsal; V, ventral. Arrow lines indicating the glia length.

Figure S3. CRISPR-mediated *betaPix* F_0 knockouts had similar phenotypes as *betaPix^{m/m}* mutants.

(A) Schematic diagram illustrates guide RNA sites for CRISPR-mediated *betaPix* F_0 knockout.

(B) qRT-PCR revealing that *betaPix* decreased in CRISPR-mediated *betaPix* F_0 knockout embryos compared with wild-type siblings at 48 hpf. Each dot represents one embryo.

(C) Maximal intensity projection of the glial structure of the hindbrains in siblings and CRISPR-mediated *betaPix* F_0 knockouts at 48 hpf, lateral view with anterior to left.

(D) Whole-mount RNA *in situ* hybridization revealing up-regulated *nestin* and down-regulated *pax2a* expression in CRISPR-mediated *betaPix* F_0 knockout embryos at 48 hpf. Dorsal view with the anterior to the left.

(E) Quantification of hemorrhagic parameters of siblings and CRISPR-mediated *betaPix* F_0 knockout embryos at 48 hpf. Left panel showing percentages of brain hemorrhage with each dot representing an individual experiment (n>100 each), and right panel showing hemorrhage areas with each dot representing one embryo.

(F) Quantification of CtA parameters of siblings and CRISPR-mediated *betaPix* F_0 knockout embryos at 48 hpf. Left panel showing average CtA length, and right panel showing CtA length index normalized to individual head length, with each dot representing one embryo.

(G) Quantification of glial parameters of siblings and CRISPR-mediated *betaPix* F_0 knockout embryos at 48 hpf. Left panel showing average glia length, and right panel showing glia length index normalized to individual head length, with each dot representing one embryo.

Figure S4. Generating glial-specific *betaPix* knockout zebrafish.

(A) Schematic diagram illustrates establishment of glial-specific *betaPix* knockout

zebrafish.

(B) 3D reconstruction of the heads at 48 hpf with lateral view, anterior to

left. Glial-specific *Cre* expression (green) was only shown in *gfap:GFP-Cre*;

betaPix^{ct/ct} mutant embryos, which overlaps with betaPix:RFP expression

(magenta).

(C) Genomic PCR analysis of the *betaPix^{ct}*- or *betaPix^{m}*-unique sequences in

betaPix^{ct/ct} siblings, *gfap:GFP-Cre*; *betaPix^{ct/+}* heterozygous mutants, and

gfap:GFP-Cre; *betaPix^{ct/ct}* homozygous mutants.

(D) qRT-PCR analysis revealing *Cre* expression in *gfap:GFP*-positive glial

population and *gfap:GFP*-negative non-glial population that were sorted from

gfap:GFP-Cre; *betaPix^{ct/ct}* embryos. Each dot represents one embryo.

(E) qRT-PCR analysis showing *betaPix, betaPix-RFP* and *alphaPix* expression in

betaPix^{ct/ct} control siblings and *gfap:GFP-Cre*; *betaPix^{ct/ct}* mutant embryos at

48 hpf. Each dot represents one embryo.

(F) Whole-mount RNA *in situ* hybridization revealing that *alphaPix* was

comparable but *betaPix* decreased in *gfap:GFP-Cre*; *betaPix^{ct/ct} mutants*

compared with *betaPix^{ct/ct}* siblings at 48 hpf. Lateral view with anterior to the

left.

Figure S5. Neither endothelial-, neuronal- nor mural-specific deletion of *betaPix*

caused brain hemorrhages and abnormal CtA phenotypes.

(A) Representative stereomicroscopy images of *o*-dianisidine staining in

betaPix$^{ct/ct}$, *kdrl:Cre*; *betaPix*$^{ct/ct}$, *huC:GFP-Cre*; *betaPix*$^{ct/ct}$, and *acta2:GFP-*

Cre; *betaPix*$^{ct/ct}$ embryos at 48 hpf.

(B) Quantification of hemorrhage parameters in (A).

(C) 3D reconstruction of the vasculature of *betaPix*$^{ct/ct}$, *kdrl:Cre*; *betaPix*$^{ct/ct}$,

huC:GFP-Cre; *betaPix*$^{ct/ct}$, and *acta2:GFP-Cre*; *betaPix*$^{ct/ct}$ embryos at 48 hpf.

Lateral view with anterior to left. The boxed areas of the hindbrains are shown

in higher magnifications at the right panels.

(D) Quantification of CtA parameters in (C).

Figure S6. Glial progenitor sub-cluster of *betaPix* knockouts has down-regulation

of s*tathmin* family members and up-regulation of *pak1* gene.

(A) UMAP visualization and clustering of cells labeled by cell type across four

sample groups.

(B) Dot plot of marker genes expression in each cell type. Dot size indicates the

percentage of cells expressed, and dot color represents the average expression

level.

(C) UMAP feature plots displaying relative expression levels of selected

transcripts among four sample groups. Cells are colored by expression level.

Figure S7. CRISPR-mediated *stmn1a/1b/4l* F$_0$ knockouts have similar but milder phenotypes as *betaPix* knockouts.

(A) qRT-PCR analysis showing that IPA3 treatment rescued *stmn1a* and *stmn4l* expression in CRISPR-mediated *betaPix* F$_0$ knockouts at 48 hpf. Each dot represents one embryo.

(B) Schematic diagram illustrates 4-guide RNAs mediated F$_0$ knockout strategy in the loci of *stmn1a*, *stmn1b* and *stmn4l*.

(C) Whole-mount RNA *in situ* hybridization confirming that *stmn1a, stmn1b* and *stmn4l* decreased in CRISPR-mediated *stmn1a/1b/4l* F$_0$ knockouts at 48 hpf. Dorsal views with anterior to the left.

(D) qRT-PCR analysis confirming that *stmn1a, stmn1b* and *stmn4l* decreased in mutant embryos of CRISPR-mediated *stmn1a/1b/4l* F$_0$ knockouts at 48 hpf. Each dot represents one embryo.

(E) qRT-PCR analysis confirming that *stmn1a, stmn1b* and *stmn4l* decreased in the FACS-sorted glia of CRISPR-mediated *stmn1a/1b/4l* F$_0$ knockouts at 48 hpf. Each dot represents sorted cells from one embryo.

(F) Optical sections of glial structure in the hindbrains of siblings and CRISPR-mediated *stmn1a/1b/4l* F$_0$ knockouts at 48 hpf. Lateral view with anterior to the left. Abnormal glial structures with disoriented arrangements (yellow arrows) in *stmn1a/1b/4l* mutant embryos.

(G) Quantification of average CtA and glial length in Figure 5 (E). Each dot represents one embryo.

(H) Whole-mount RNA *in situ* hybridization revealed that *nestin* increased while

pax2a decreased in CRISPR-mediated *stmn1a/1b/4l* F_0 knockout embryos at

48 hpf. Dorsal views with anterior to the left.

(I) 3D reconstruction and whole-mount RNA *in situ* hybridization showing that

gfap:GFP-stmn1b transgenic overexpression rescued glial defects in *bbh^{fn40a}*;

Tg(*gfap:GFP-stmn1b*) embryos at 48 hpf. Lateral views with anterior to the

left.

(J) Whole-mount RNA *in situ* hybridization showing that *nestin* and *pax2a* were

normally expressed in *bbh^{fn40a}*; Tg(*gfap:GFP-stmn1b*) embryos at 48 hpf.

Dorsal views with anterior to the left.

(K) Quantification of average CtA and glia length in Figure 5 (I). Each dot

represents one embryo.

(L) qRT-PCR analysis confirming the efficacy of *betaPix* transgenic over-

expression and *betaPix* siRNA knockdown in U251 cells.

(M) Representative stereomicroscopy images of A172 cells at 0 and 18 hours

after wounding. *betaPIX* siRNA treatment decreased A172 cell migration,

which were rescued by either *betaPIX* or STMN1 overexpression. The wound

edges are highlighted by dashed lines, with arrow lines indicating the wound

width.

(N) Quantification of wound closures in (M). $^{**}P<0.01$ compared to negative

control siRNA and empty vector transfection. $^{\#\#}P<0.01$, $^{\#\#\#}P<0.005$ compared

to *betaPix* knockdown and empty vector transfection.

Figure S8. *Zfhx3/4* **acts downstream of** *betaPix* **to regulate vascular integrity development.**

(A) Violin plots showing that *zfhx3* and *zfhx4* decreased in progenitor sub-cluster of *betaPix* knockouts.

(B) qRT-PCR analysis revealing that *zfhx3* and *zfhx4* decreased in FACS-sorted glia of *gfap:GFP-Cre*; *betaPix^{ct/ct}* mutants compared with *betaPix^{ct/ct}* siblings at 48 hpf. Each dot represents the cells sorted from one embryo.

(C) Schematic diagram illustrates 4-guide RNAs mediated F_0 knockout strategy in the *zfhx3* and *zfhx4* loci.

(D) Whole-mount RNA *in situ* hybridization confirming *zfhx3/4* efficiency in CRISPR-mediated *zfhx3/4* F_0 knockout embryos at 48 hpf.

(E) qRT-PCR analysis revealing that *zfhx3*, *zfhx4*, and *vegfaa* decreased in CRISPR-mediated *zfhx3/4* F_0 knockout embryos at 36 hpf.

(F) Whole-mount RNA *in situ* hybridization revealing that *nestin* increased in CRISPR-mediated *zfhx3/4* F_0 knockout embryos at 36 hpf. Arrows indicate the hindbrain regions.

(G) qRT-PCR analysis confirming overexpression of *zfhx4* in CRISPR-mediated *betaPix* F_0 knockout embryos with *zfhx4* mRNA injection at 70% epiboly.

(H) Representative stereomicroscopy images of erythrocytes stained with *o-dianisidine* showing that brain hemorrhages decreased in *bbh^{fn40a}* mutants with zfhx4 mRNA injection at 48 hpf. Arrows indicate brain hemorrhages. Lateral views with anterior to the left.

1056   (I) Quantification of hemorrhagic parameters in (H). Left panel showing

1057   hemorrhage percentages with each dot representing an individual experiment

1058   ($n>100$ each), and right panel showing hemorrhage area with each dot

1059   representing one embryo.

References

1. Sheth, K.N. (2022). Spontaneous Intracerebral Hemorrhage. New England Journal of Medicine *387*, 1589-1596. doi:10.1056/NEJMra2201449.

2. Chiang, P.T., Tsai, L.K., and Tsai, H.H. (2025). New targets in spontaneous intracerebral hemorrhage. Curr Opin Neurol 38, 10-17. 10.1097/wco.0000000000001325.

3. Thangameeran, S.I.M., Pang, C.Y., Lee, C.H., Tsai, S.T., Hu, W.F., and Liew, H.K. (2023). Experimental animal models and evaluation techniques in intracerebral hemorrhage. Tzu Chi Med J *35*, 1-10. 10.4103/tcmj.tcmj_119_22.

4. Isogai, S., Horiguchi, M., and Weinstein, B.M. (2001). The vascular anatomy of the developing zebrafish: an atlas of embryonic and early larval development. Dev Biol *230*, 278-301. 10.1006/dbio.2000.9995.

5. Buchner, D.A., Su, F., Yamaoka, J.S., Kamei, M., Shavit, J.A., Barthel, L.K., McGee, B., Amigo, J.D., Kim, S., Hanosh, A.W., et al. (2007). pak2a mutations cause cerebral hemorrhage in redhead zebrafish. Proceedings of the National Academy of Sciences *104*, 13996-14001. 10.1073/pnas.0700947104.

6. Jin, S.W., Herzog, W., Santoro, M.M., Mitchell, T.S., Frantsve, J., Jungblut, B., Beis, D., Scott, I.C., D'Amico, L.A., Ober, E.A., et al. (2007). A transgene-assisted genetic screen identifies essential regulators of vascular development in vertebrate embryos. Dev Biol 307, 29-42. 10.1016/j.ydbio.2007.03.526.

7.   Hegarty, J.M., Yang, H., and Chi, N.C. (2013). UBIAD1-mediated vitamin K2

synthesis is required for vascular endothelial cell survival and development.

Development *140*, 1713-1719. 10.1242/dev.093112.

8.   Stainier, D.Y., Fouquet, B., Chen, J.N., Warren, K.S., Weinstein, B.M., Meiler,

S.E., Mohideen, M.A., Neuhauss, S.C., Solnica-Krezel, L., Schier, A.F., et al.

(1996). Mutations affecting the formation and function of the cardiovascular

system in the zebrafish embryo. Development *123*, 285-292.

9.   Chen, J.N., van Bebber, F., Goldstein, A.M., Serluca, F.C., Jackson, D., Childs,

S., Serbedzija, G., Warren, K.S., Mably, J.D., Lindahl, P., et al. (2001). Genetic

steps to organ laterality in zebrafish. Comp Funct Genomics *2*, 60-68.

10.1002/cfg.74.

10.   Liu, J., Fraser, S.D., Faloon, P.W., Rollins, E.L., Vom Berg, J., Starovic-

Subota, O., Laliberte, A.L., Chen, J.N., Serluca, F.C., and Childs, S.J. (2007).

A betaPix Pak2a signaling pathway regulates cerebral vascular stability in

zebrafish. Proc Natl Acad Sci U S A 104, 13990-13995.

10.1073/pnas.0700825104.

11.   Manser, E., Loo, T.H., Koh, C.G., Zhao, Z.S., Chen, X.Q., Tan, L., Tan, I.,

Leung, T., and Lim, L. (1998). PAK kinases are directly coupled to the PIX

family of nucleotide exchange factors. Mol Cell 1, 183-192. 10.1016/s1097-

2765(00)80019-2

12.   Zhou, W., Li, X., and Premont, R.T. (2016). Expanding functions of GIT Arf

GTPase-activating proteins, PIX Rho guanine nucleotide exchange factors and

GIT-PIX complexes. J Cell Sci *129*, 1963-1974. 10.1242/jcs.179465.

13.   Morimoto, K., Tabata, H., Takahashi, R., and Nakajima, K. (2024).

Interactions between neural cells and blood vessels in central nervous system

development. Bioessays *46*, e2300091. 10.1002/bies.202300091.

14.   Grinchevskaya, L.R., Salikhova, D.I., Silachev, D.N., and Goldshtein, D.V.

(2024). Neural and Glial Regulation of Angiogenesis in CNS in Ischemic

Stroke. Bull Exp Biol Med *177*, 528-533. 10.1007/s10517-024-06219-4.

15.   McCarty, J.H. (2009). Cell adhesion and signaling networks in brain

neurovascular units. Curr Opin Hematol *16*, 209-214.

10.1097/MOH.0b013e32832a07eb.

16.   McCarty, J.H. (2009). Integrin-mediated regulation of neurovascular

development, physiology and disease. Cell Adh Migr *3*, 211-215.

10.4161/cam.3.2.7767.

17.   Liu, J., Zeng, L., Kennedy, R.M., Gruenig, N.M., and Childs, S.J. (2012). β

Pix plays a dual role in cerebral vascular stability and angiogenesis, and

interacts with integrin αvβ8. Dev Biol *363*, 95-105.

10.1016/j.ydbio.2011.12.022.

18.   Yang, R., Zhang, Y., Huang, D., Luo, X., Zhang, L., Zhu, X., Zhang, X., Liu,

Z., Han, J.Y., and Xiong, J.W. (2017). Miconazole protects blood vessels from

MMP9-dependent rupture and hemorrhage. Dis Model Mech *10*, 337-348.

10.1242/dmm.027268.

19.   Sugimoto, K., Hui, S.P., Sheng, D.Z., and Kikuchi, K. (2017). Dissection of

zebrafish shha function using site-specific targeting with a Cre-dependent

genetic switch. Elife *6*. 10.7554/eLife.24635.

20.   Ogawa, M., Geng, F.S., Humphreys, D.T., Kristianto, E., Sheng, D.Z., Hui,

S.P., Zhang, Y., Sugimoto, K., Nakayama, M., Zheng, D., et al. (2021). Krü

ppel-like factor 1 is a core cardiomyogenic trigger in zebrafish. Science *372*,

201-205. 10.1126/science.abe2762.

21.   Ogawa, M., and Kikuchi, K. (2024). Generation of Conditional Knockout

Zebrafish Using an Invertible Gene-Trap Cassette. Methods Mol Biol *2707*,

205-214. 10.1007/978-1-0716-3401-1_13.

22.   Wierson, W.A., Simone, B.W., WareJoncas, Z., Mann, C., Welker, J.M., Kar,

B., Emch, M.J., Friedberg, I., Gendron, W.A.C., Barry, M.A., et al. (2019).

Expanding the CRISPR Toolbox with ErCas12a in Zebrafish and Human

Cells. Crispr j *2*, 417-433. 10.1089/crispr.2019.0026.

23.   Wu, R.S., Lam, II, Clay, H., Duong, D.N., Deo, R.C., and Coughlin, S.R.

(2018). A Rapid Method for Directed Gene Knockout for Screening in G0

Zebrafish. Dev Cell *46*, 112-125 e114. 10.1016/j.devcel.2018.06.003.

24.   Quiñonez-Silvero, C., Hübner, K., and Herzog, W. (2020). Development of the

brain vasculature and the blood-brain barrier in zebrafish. Dev Biol *457*, 181-

190. 10.1016/j.ydbio.2019.03.005.

25.   Deacon, S.W., Beeser, A., Fukui, J.A., Rennefahrt, U.E., Myers, C., Chernoff,

J., and Peterson, J.R. (2008). An isoform-selective, small-molecule inhibitor

targets the autoregulatory mechanism of p21-activated kinase. Chem Biol *15*,

322-331. 10.1016/j.chembiol.2008.03.005.

26.   Bernardos, R.L., and Raymond, P.A. (2006). GFAP transgenic zebrafish. Gene

Expr Patterns *6*, 1007-1013. 10.1016/j.modgep.2006.04.006.

27.   Kim, C.H., Ueshima, E., Muraoka, O., Tanaka, H., Yeo, S.Y., Huh, T.L., and

Miki, N. (1996). Zebrafish elav/HuC homologue as a very early neuronal

marker. Neurosci Lett *216*, 109-112. 10.1016/0304-3940(96)13021-4.

28.   Raj, B., Wagner, D.E., McKenna, A., Pandey, S., Klein, A.M., Shendure, J.,

Gagnon, J.A., and Schier, A.F. (2018). Simultaneous single-cell profiling of

lineages and cell types in the vertebrate brain. Nature Biotechnology *36*, 442-

450. 10.1038/nbt.4103.

29.   Raj, B., Farrell, J.A., Liu, J., El Kholtei, J., Carte, A.N., Navajas Acedo, J., Du,

L.Y., McKenna, A., Relić, Đ., Leslie, J.M., and Schier, A.F. (2020).

Emergence of Neuronal Diversity during Vertebrate Brain Development.

Neuron *108*, 1058-1074.e1056. 10.1016/j.neuron.2020.09.023.

30.   Farnsworth, D.R., Saunders, L.M., and Miller, A.C. (2020). A single-cell

transcriptome atlas for zebrafish development. Dev Biol 459, 100-108.

10.1016/j.ydbio.2019.11.008.

31.   Etienne-Manneville, S. (2013). Microtubules in cell migration. Annu Rev Cell

Dev Biol *29*, 471-499. 10.1146/annurev-cellbio-101011-155711.

32.   Kwon, Y., Jeon, Y.W., Kwon, M., Cho, Y., Park, D., and Shin, J.E. (2020). β

Pix-d promotes tubulin acetylation and neurite outgrowth through a

PAK/Stathmin1 signaling pathway. PLoS One *15*, e0230814.

10.1371/journal.pone.0230814.

33.   Omelchenko, T., Rabadan, M.A., Hernandez-Martinez, R., Grego-Bessa, J.,

1172   Anderson, K.V., and Hall, A. (2014). beta-Pix directs collective migration of

1173   anterior visceral endoderm cells in the early mouse embryo. Genes Dev *28*,

1174   2764-2777. 10.1101/gad.251371.114.

1175   34.   Hsu, R.M., Hsieh, Y.J., Yang, T.H., Chiang, Y.C., Kan, C.Y., Lin, Y.T., Chen,

1176   J.T., and Yu, J.S. (2014). Binding of the extreme carboxyl-terminus of PAK-

1177   interacting exchange factor β (βPIX) to myosin 18A (MYO18A) is required

1178   for epithelial cell migration. Biochim Biophys Acta *1843*, 2513-2527.

1179   10.1016/j.bbamcr.2014.06.023.

1180   35.   López Tobón, A., Suresh, M., Jin, J., Vitriolo, A., Pietralla, T., Tedford, K.,

1181   Bossenz, M., Mahnken, K., Kiefer, F., Testa, G., et al. (2018). The guanine

1182   nucleotide exchange factor Arhgef7/βPix promotes axon formation upstream

1183   of TC10. Scientific Reports *8*, 8811. 10.1038/s41598-018-27081-1.

1184   36.   Cheng, K., Chahdi, A., Larabee, S.M., Tolaymat, M., Sundel, M.H.,

1185   Drachenberg, C.B., Zhan, M., Hu, S., Said, A.H., Shang, A.C., et al. (2023).

1186   Muscarinic receptor agonist-induced βPix binding to β-catenin promotes

1187   colon neoplasia. Sci Rep *13*, 16920. 10.1038/s41598-023-44158-8.

1188   37.   Connor, K., Murray, D.W., Jarzabek, M.A., Tran, N.L., White, K., Dicker, P.,

1189   Sweeney, K.J., O'Halloran, P.J., MacCarthy, B., Shiels, L.P., et al. (2020).

1190   Targeting the RhoGEF βPIX/COOL-1 in Glioblastoma: Proof of Concept

1191   Studies. Cancers (Basel) *12*. 10.3390/cancers12123531.

1192   38.   Ward, J.D., Ha, J.H., Jayaraman, M., and Dhanasekaran, D.N. (2015). LPA-

1193   mediated migration of ovarian cancer cells involves translocalization of Gαi2

1194   to invadopodia and association with Src and β-pix. Cancer Lett *356*, 382-391.

1195   10.1016/j.canlet.2014.09.030.

1196   39.   Bussmann, J., Wolfe, S.A., and Siekmann, A.F. (2011). Arterial-venous

1197   network formation during brain vascularization involves hemodynamic

1198   regulation of chemokine signaling. Development *138*, 1717-1726.

1199   10.1242/dev.059881.

1200   40.   Eubelen, M., Bostaille, N., Cabochette, P., Gauquier, A., Tebabi, P., Dumitru,

1201   A.C., Koehler, M., Gut, P., Alsteens, D., Stainier, D.Y.R., et al. (2018). A

1202   molecular mechanism for Wnt ligand-specific signaling. Science *361*.

1203   10.1126/science.aat1178.

1204   41.   Pitulescu, M.E., and Adams, R.H. (2014). Regulation of signaling interactions

1205   and receptor endocytosis in growing blood vessels. Cell Adh Migr *8*, 366-377.

1206   10.4161/19336918.2014.970010.

1207   42.   Wang, J., Chen, Y., Zeng, Z., Feng, R., Wang, Q., Zhang, Q., Sun, K., Chen,

1208   A.F., Lu, Y., and Yu, Y. (2021). HMGA2 contributes to vascular development

1209   and sprouting angiogenesis by promoting IGFBP2 production. Exp Cell Res

1210   *408*, 112831. 10.1016/j.yexcr.2021.112831.

1211   43.   Fu, C., An, N., Liu, J., A, J., Zhang, B., Liu, M., Zhang, Z., Fu, L., Tian, X.,

1212   Wang, D., and Dong, J.T. (2020). The transcription factor ZFHX3 is crucial

1213   for the angiogenic function of hypoxia-inducible factor 1α in liver cancer

1214 cells. J Biol Chem *295*, 7060-7074. 10.1074/jbc.RA119.012131.

1215 44. Gudbjartsson, D.F., Holm, H., Gretarsdottir, S., Thorleifsson, G., Walters,

1216 G.B., Thorgeirsson, G., Gulcher, J., Mathiesen, E.B., Njølstad, I., Nyrnes, A.,

1217 et al. (2009). A sequence variant in ZFHX3 on 16q22 associates with atrial

1218 fibrillation and ischemic stroke. Nat Genet *41*, 876-878. 10.1038/ng.417.

1219 45. Soriano-Tárraga, C., Lazcano, U., Giralt-Steinhauer, E., Avellaneda-Gómez,

1220 C., Ois, Á., Rodríguez-Campello, A., Cuadrado-Godia, E., Gómez-González,

1221 A., Fernández-Sanlés, A., Elosua, R., et al. (2020). Identification of 20 novel

1222 loci associated with ischaemic stroke. Epigenome-wide association study.

1223 Epigenetics *15*, 988-997. 10.1080/15592294.2020.1746507.

1224 46. Malik, R., Traylor, M., Pulit, S.L., Bevan, S., Hopewell, J.C., Holliday, E.G.,

1225 Zhao, W., Abrantes, P., Amouyel, P., Attia, J.R., et al. (2016). Low-frequency

1226 and common genetic variation in ischemic stroke: The METASTROKE

1227 collaboration. Neurology *86*, 1217-1226. 10.1212/wnl.0000000000002528.

1228 47. Kalvaitytė, M., and Balciunas, D. (2022). Conditional mutagenesis strategies

1229 in zebrafish. Trends Genet *38*, 856-868. 10.1016/j.tig.2022.04.007.

1230 48. Obermeier, B., Daneman, R., and Ransohoff, R.M. (2013). Development,

1231 maintenance and disruption of the blood-brain barrier. Nat Med *19*, 1584-

1232 1596. 10.1038/nm.3407.

1233 49. Langen, U.H., Ayloo, S., and Gu, C. (2019). Development and Cell Biology of

1234 the Blood-Brain Barrier. Annu Rev Cell Dev Biol *35*, 591-613.

1235 10.1146/annurev-cellbio-100617-062608.

1236   50.   Fleming, A., Diekmann, H., and Goldsmith, P. (2013). Functional

1237   characterisation of the maturation of the blood-brain barrier in larval zebrafish.

1238   PLoS One *8*, e77548. 10.1371/journal.pone.0077548.

1239   51.   Xie, J., Farage, E., Sugimoto, M., and Anand-Apte, B. (2010). A novel

1240   transgenic zebrafish model for blood-brain and blood-retinal barrier

1241   development. BMC Dev Biol *10*, 76. 10.1186/1471-213x-10-76.

1242   52.   Armulik, A., Genove, G., and Betsholtz, C. (2011). Pericytes: developmental,

1243   physiological, and pathological perspectives, problems, and promises. Dev

1244   Cell *21*, 193-215. 10.1016/j.devcel.2011.07.001.

1245   53.   Armulik, A., Genové, G., Mäe, M., Nisancioglu, M.H., Wallgard, E., Niaudet,

1246   C., He, L., Norlin, J., Lindblom, P., Strittmatter, K., et al. (2010). Pericytes

1247   regulate the blood-brain barrier. Nature *468*, 557-561. 10.1038/nature09522.

1248   54.   Bell, R.D., Winkler, E.A., Sagare, A.P., Singh, I., LaRue, B., Deane, R., and

1249   Zlokovic, B.V. (2010). Pericytes control key neurovascular functions and

1250   neuronal phenotype in the adult brain and during brain aging. Neuron *68*, 409-

1251   427. 10.1016/j.neuron.2010.09.043.

1252   55.   Daneman, R., Zhou, L., Kebede, A.A., and Barres, B.A. (2010). Pericytes are

1253   required for blood-brain barrier integrity during embryogenesis. Nature *468*,

1254   562-566. 10.1038/nature09513.

1255   56.   Antonelli-Orlidge, A., Saunders, K.B., Smith, S.R., and D'Amore, P.A. (1989).

1256   An activated form of transforming growth factor beta is produced by

1257   cocultures of endothelial cells and pericytes. Proc Natl Acad Sci U S A *86*,

1258   4544-4548. 10.1073/pnas.86.12.4544.

57.   Dohgu, S., Takata, F., Yamauchi, A., Nakagawa, S., Egawa, T., Naito, M., Tsuruo, T., Sawada, Y., Niwa, M., and Kataoka, Y. (2005). Brain pericytes contribute to the induction and up-regulation of blood-brain barrier functions through transforming growth factor-beta production. Brain Res *1038*, 208-215. 10.1016/j.brainres.2005.01.027.

58.   Dave, J.M., Mirabella, T., Weatherbee, S.D., and Greif, D.M. (2018). Pericyte ALK5/TIMP3 Axis Contributes to Endothelial Morphogenesis in the Developing Brain. Dev Cell *44*, 665-678.e666. 10.1016/j.devcel.2018.01.018.

59.   Reyahi, A., Nik, A.M., Ghiami, M., Gritli-Linde, A., Pontén, F., Johansson, B.R., and Carlsson, P. (2015). Foxf2 Is Required for Brain Pericyte Differentiation and Development and Maintenance of the Blood-Brain Barrier. Dev Cell *34*, 19-32. 10.1016/j.devcel.2015.05.008.

60.   Kim, Y., Imdad, R.Y., Stephenson, A.H., Sprague, R.S., and Lonigro, A.J. (1998). Vascular endothelial growth factor mRNA in pericytes is upregulated by phorbol myristate acetate. Hypertension *31*, 511-515. 10.1161/01.hyp.31.1.511.

61.   Franco, M., Roswall, P., Cortez, E., Hanahan, D., and Pietras, K. (2011). Pericytes promote endothelial cell survival through induction of autocrine VEGF-A signaling and Bcl-w expression. Blood *118*, 2906-2917. 10.1182/blood-2011-01-331694.

62.   Sato, T.N., Tozawa, Y., Deutsch, U., Wolburg-Buchholz, K., Fujiwara, Y.,

Gendron-Maguire, M., Gridley, T., Wolburg, H., Risau, W., and Qin, Y. (1995).

Distinct roles of the receptor tyrosine kinases Tie-1 and Tie-2 in blood vessel

formation. Nature *376*, 70-74. 10.1038/376070a0.

63.   Thurston, G., Suri, C., Smith, K., McClain, J., Sato, T.N., Yancopoulos, G.D.,

and McDonald, D.M. (1999). Leakage-resistant blood vessels in mice

transgenically overexpressing angiopoietin-1. Science *286*, 2511-2514.

10.1126/science.286.5449.2511.

64.   Sundberg, C., Kowanetz, M., Brown, L.F., Detmar, M., and Dvorak, H.F.

(2002). Stable expression of angiopoietin-1 and other markers by cultured

pericytes: phenotypic similarities to a subpopulation of cells in maturing

vessels during later stages of angiogenesis in vivo. Lab Invest *82*, 387-401.

10.1038/labinvest.3780433.

65.   Wan, Y., Jin, H.J., Zhu, Y.Y., Fang, Z., Mao, L., He, Q., Xia, Y.P., Li, M., Li,

Y., Chen, X., and Hu, B. (2018). MicroRNA-149-5p regulates blood-brain

barrier permeability after transient middle cerebral artery occlusion in rats by

targeting S1PR2 of pericytes. Faseb j *32*, 3133-3148. 10.1096/fj.201701121R.

66.   Li, F., Lan, Y., Wang, Y., Wang, J., Yang, G., Meng, F., Han, H., Meng, A.,

Wang, Y., and Yang, X. (2011). Endothelial Smad4 maintains cerebrovascular

integrity by activating N-cadherin through cooperation with Notch. Dev Cell

20, 291-302. 10.1016/j.devcel.2011.01.011.

67.   Liu, H., Zhang, W., Kennard, S., Caldwell, R.B., and Lilly, B. (2010). Notch3

is critical for proper angiogenesis and mural cell investment. Circ Res *107*,

860-870. 10.1161/circresaha.110.218271.

68.   Gundersen, G.A., Vindedal, G.F., Skare, O., and Nagelhus, E.A. (2014).

Evidence that pericytes regulate aquaporin-4 polarization in mouse cortical

astrocytes. Brain Struct Funct *219*, 2181-2186. 10.1007/s00429-013-0629-0.

69.   Jeanne, M., Jorgensen, J., and Gould, D.B. (2015). Molecular and Genetic

Analyses of Collagen Type IV Mutant Mouse Models of Spontaneous

Intracerebral Hemorrhage Identify Mechanisms for Stroke Prevention.

Circulation *131*, 1555-1565. 10.1161/circulationaha.114.013395.

70.   Gautam, J., Zhang, X., and Yao, Y. (2016). The role of pericytic laminin in

blood brain barrier integrity maintenance. Sci Rep *6*, 36450.

10.1038/srep36450.

71.   Gautam, J., Cao, Y., and Yao, Y. (2020). Pericytic Laminin Maintains Blood-

Brain Barrier Integrity in an Age-Dependent Manner. Transl Stroke Res *11*,

228-242. 10.1007/s12975-019-00709-8.

72.   Özkan, E., Çetin-Taş, Y., Şekerdağ, E., Kızılırmak, A.B., Taş, A., Yıldız, E.,

Yapıcı-Eser, H., Karahüseyinoğlu, S., Zeybel, M., and Gürsoy-Özdemir, Y.

(2021). Blood-brain barrier leakage and perivascular collagen accumulation

precede microvessel rarefaction and memory impairment in a chronic

hypertension animal model. Metab Brain Dis *36*, 2553-2566. 10.1007/s11011-

021-00767-8.

73.   Stewart, P.A., and Wiley, M.J. (1981). Developing nervous tissue induces

formation of blood-brain barrier characteristics in invading endothelial cells: a

study using quail--chick transplantation chimeras. Dev Biol *84*, 183-192.

10.1016/0012-1606(81)90382-1.

74. Hellström, M., Kalén, M., Lindahl, P., Abramsson, A., and Betsholtz, C.

(1999). Role of PDGF-B and PDGFR-beta in recruitment of vascular smooth

muscle cells and pericytes during embryonic blood vessel formation in the

mouse. Development *126*, 3047-3055.

75. Bjarnegård, M., Enge, M., Norlin, J., Gustafsdottir, S., Fredriksson, S.,

Abramsson, A., Takemoto, M., Gustafsson, E., Fässler, R., and Betsholtz, C.

(2004). Endothelium-specific ablation of PDGFB leads to pericyte loss and

glomerular, cardiac and placental abnormalities. Development *131*, 1847-

1857. 10.1242/dev.01080.

76. Mi, H., Haeberle, H., and Barres, B.A. (2001). Induction of astrocyte

differentiation by endothelial cells. J Neurosci *21*, 1538-1547.

10.1523/jneurosci.21-05-01538.2001.

77. Paredes, I., Vieira, J.R., Shah, B., Ramunno, C.F., Dyckow, J., Adler, H.,

Richter, M., Schermann, G., Giannakouri, E., Schirmer, L., et al. (2021).

Oligodendrocyte precursor cell specification is regulated by bidirectional

neural progenitor-endothelial cell crosstalk. Nat Neurosci *24*, 478-488.

10.1038/s41593-020-00788-z.

78. Daneman, R., Agalliu, D., Zhou, L., Kuhnert, F., Kuo, C.J., and Barres, B.A.

(2009). Wnt/beta-catenin signaling is required for CNS, but not non-CNS,

angiogenesis. Proc Natl Acad Sci U S A *106*, 641-646.

10.1073/pnas.0805165106.

79.   Kuhnert, F., Mancuso, M.R., Shamloo, A., Wang, H.T., Choksi, V., Florek, M.,

Su, H., Fruttiger, M., Young, W.L., Heilshorn, S.C., and Kuo, C.J. (2010).

Essential regulation of CNS angiogenesis by the orphan G protein-coupled

receptor GPR124. Science *330*, 985-989. 10.1126/science.1196554.

80.   Cullen, M., Elzarrad, M.K., Seaman, S., Zudaire, E., Stevens, J., Yang, M.Y.,

Li, X., Chaudhary, A., Xu, L., Hilton, M.B., et al. (2011). GPR124, an orphan

G protein-coupled receptor, is required for CNS-specific vascularization and

establishment of the blood-brain barrier. Proc Natl Acad Sci U S A *108*, 5759-

5764. 10.1073/pnas.1017192108.

81.   Zhou, Y., and Nathans, J. (2014). Gpr124 controls CNS angiogenesis and

blood-brain barrier integrity by promoting ligand-specific canonical wnt

signaling. Dev Cell *31*, 248-256. 10.1016/j.devcel.2014.08.018.

82.   Thomas, M., and Augustin, H.G. (2009). The role of the Angiopoietins in

vascular morphogenesis. Angiogenesis *12*, 125-137. 10.1007/s10456-009-

9147-3.

83.   Nico, B., Frigeri, A., Nicchia, G.P., Quondamatteo, F., Herken, R., Errede, M.,

Ribatti, D., Svelto, M., and Roncali, L. (2001). Role of aquaporin-4 water

channel in the development and integrity of the blood-brain barrier. J Cell Sci

114, 1297-1307. 10.1242/jcs.114.7.1297.

84.   Nicchia, G.P., Nico, B., Camassa, L.M., Mola, M.G., Loh, N., Dermietzel, R.,

Spray, D.C., Svelto, M., and Frigeri, A. (2004). The role of aquaporin-4 in the

blood-brain barrier development and integrity: studies in animal and cell

culture models. Neuroscience *129*, 935-945.

10.1016/j.neuroscience.2004.07.055.

85.   Hartfuss, E., Galli, R., Heins, N., and Götz, M. (2001). Characterization of

CNS precursor subtypes and radial glia. Dev Biol *229*, 15-30.

10.1006/dbio.2000.9962.

86.   Malatesta, P., Appolloni, I., and Calzolari, F. (2008). Radial glia and neural

stem cells. Cell Tissue Res *331*, 165-178. 10.1007/s00441-007-0481-8.

87.   Qian, X., Shen, Q., Goderie, S.K., He, W., Capela, A., Davis, A.A., and

Temple, S. (2000). Timing of CNS cell generation: a programmed sequence of

neuron and glial cell production from isolated murine cortical stem cells.

Neuron *28*, 69-80. 10.1016/s0896-6273(00)00086-6.

88.   Price, J., Williams, B.P., and Götz, M. (1995). The generation of cellular

diversity in the cerebral cortex. Ciba Found Symp *193*, 71-84; discussion 117-

126. 10.1002/9780470514795.ch4.

89.   Grupp, L., Wolburg, H., and Mack, A.F. (2010). Astroglial structures in the

zebrafish brain. J Comp Neurol *518*, 4277-4287. 10.1002/cne.22481.

90.   Chen, J., Poskanzer, K.E., Freeman, M.R., and Monk, K.R. (2020). Live-

imaging of astrocyte morphogenesis and function in zebrafish neural circuits.

Nat Neurosci *23*, 1297-1306. 10.1038/s41593-020-0703-x.

91.   Johnson, K., Barragan, J., Bashiruddin, S., Smith, C.J., Tyrrell, C., Parsons,

M.J., Doris, R., Kucenas, S., Downes, G.B., Velez, C.M., et al. (2016). Gfap-

positive radial glial cells are an essential progenitor population for later-born

neurons and glia in the zebrafish spinal cord. Glia *64*, 1170-1189.

10.1002/glia.22990.

92.   Umans, R.A., Pollock, C., Mills, W.A., 3rd, Clark, K.C., Pan, Y.A., and

Sontheimer, H. (2021). Using Zebrafish to Elucidate Glial-Vascular

Interactions During CNS Development. Front Cell Dev Biol *9*, 654338.

10.3389/fcell.2021.654338.

93.   Matsuoka, R.L., Rossi, A., Stone, O.A., and Stainier, D.Y.R. (2017). CNS-

resident progenitors direct the vascularization of neighboring tissues. Proc

Natl Acad Sci U S A *114*, 10137-10142. 10.1073/pnas.1619300114.

94.   Alvarez, J.I., Dodelet-Devillers, A., Kebir, H., Ifergan, I., Fabre, P.J., Terouz,

S., Sabbagh, M., Wosik, K., Bourbonnière, L., Bernard, M., et al. (2011). The

Hedgehog pathway promotes blood-brain barrier integrity and CNS immune

quiescence. Science *334*, 1727-1731. 10.1126/science.1206936.

95.   Brilha, S., Ong, C.W.M., Weksler, B., Romero, N., Couraud, P.O., and

Friedland, J.S. (2017). Matrix metalloproteinase-9 activity and a

downregulated Hedgehog pathway impair blood-brain barrier function in an in

vitro model of CNS tuberculosis. Sci Rep *7*, 16031. 10.1038/s41598-017-

16250-3.

96.   Shi, T., Yue, S., Xie, C., Li, X., Yang, D., Hu, L., Zhong, Y., Zhang, Y., and

Liu, W. (2024). MMP-2-mediated Scube2 degradation promotes blood-brain

barrier disruption by blocking the interaction between astrocytes and

endothelial cells via inhibiting Sonic hedgehog pathway during early cerebral

ischemia. J Neurochem *168*, 1877-1894. 10.1111/jnc.16021.

97.   Xing, G., Zhao, T., Zhang, X., Li, H., Li, X., Cui, P., Li, M., Li, D., Zhang, N.,

and Jiang, W. (2020). Astrocytic Sonic Hedgehog Alleviates Intracerebral

Hemorrhagic Brain Injury via Modulation of Blood-Brain Barrier Integrity.

Front Cell Neurosci *14*, 575690. 10.3389/fncel.2020.575690.

98.   Lawson, N.D., Vogel, A.M., and Weinstein, B.M. (2002). sonic hedgehog and

vascular endothelial growth factor act upstream of the Notch pathway during

arterial endothelial differentiation. Dev Cell *3*, 127-136. 10.1016/s1534-

5807(02)00198-3.

99.   Mannan, A., Dhiamn, S., Garg, N., and Singh, T.G. (2023). Pharmacological

modulation of Sonic Hedgehog signaling pathways in Angiogenesis: A

mechanistic perspective. Dev Biol *504*, 58-74. 10.1016/j.ydbio.2023.09.009.

100.   Argaw, A.T., Asp, L., Zhang, J., Navrazhina, K., Pham, T., Mariani, J.N.,

Mahase, S., Dutta, D.J., Seto, J., Kramer, E.G., et al. (2012). Astrocyte-derived

VEGF-A drives blood-brain barrier disruption in CNS inflammatory disease. J

Clin Invest *122*, 2454-2468. 10.1172/jci60842.

101.   Chapouly, C., Tadesse Argaw, A., Horng, S., Castro, K., Zhang, J., Asp, L.,

Loo, H., Laitman, B.M., Mariani, J.N., Straus Farber, R., et al. (2015).

Astrocytic TYMP and VEGFA drive blood-brain barrier opening in

inflammatory central nervous system lesions. Brain *138*, 1548-1567.

10.1093/brain/awv077.

102.   Matsumoto, T., and Mugishima, H. (2006). Signal transduction via vascular

1435   endothelial growth factor (VEGF) receptors and their roles in atherogenesis. J

1436   Atheroscler Thromb *13*, 130-135. 10.5551/jat.13.130.

1437   103.   Jakobsson, L., Franco, C.A., Bentley, K., Collins, R.T., Ponsioen, B., Aspalter,

1438   I.M., Rosewell, I., Busse, M., Thurston, G., Medvinsky, A., et al. (2010).

1439   Endothelial cells dynamically compete for the tip cell position during

1440   angiogenic sprouting. Nature Cell Biology *12*, 943-953. 10.1038/ncb2103.

1441   104.   Liu, Z.J., Shirakawa, T., Li, Y., Soma, A., Oka, M., Dotto, G.P., Fairman,

1442   R.M., Velazquez, O.C., and Herlyn, M. (2003). Regulation of Notch1 and Dll4

1443   by vascular endothelial growth factor in arterial endothelial cells: implications

1444   for modulating arteriogenesis and angiogenesis. Mol Cell Biol *23*, 14-25.

1445   10.1128/mcb.23.1.14-25.2003.

1446   105.   Olsson, A.K., Dimberg, A., Kreuger, J., and Claesson-Welsh, L. (2006). VEGF

1447   receptor signalling - in control of vascular function. Nat Rev Mol Cell Biol *7*,

1448   359-371. 10.1038/nrm1911.

1449   106.   Xu, L., Nirwane, A., and Yao, Y. (2019). Basement membrane and blood-brain

1450   barrier. Stroke Vasc Neurol *4*, 78-82. 10.1136/svn-2018-000198.

1451   107.   Halder, S.K., Sapkota, A., and Milner, R. (2022). The impact of genetic

1452   manipulation of laminin and integrins at the blood-brain barrier. Fluids

1453   Barriers CNS *19*, 50. 10.1186/s12987-022-00346-8.

1454   108.   Yao, Y., Chen, Z.L., Norris, E.H., and Strickland, S. (2014). Astrocytic laminin

1455   regulates pericyte differentiation and maintains blood brain barrier integrity.

Nat Commun *5*, 3413. 10.1038/ncomms4413.

109.   Milner, R., and Campbell, I.L. (2002). Developmental regulation of beta1

integrins during angiogenesis in the central nervous system. Mol Cell

Neurosci *20*, 616-626. 10.1006/mcne.2002.1151.

110.   Nishimura, S.L., Boylen, K.P., Einheber, S., Milner, T.A., Ramos, D.M., and

Pytela, R. (1998). Synaptic and glial localization of the integrin alphavbeta8 in

mouse and rat brain. Brain Res *791*, 271-282. 10.1016/s0006-8993(98)00118-

8.

111.   McCarty, J.H., Lacy-Hulbert, A., Charest, A., Bronson, R.T., Crowley, D.,

Housman, D., Savill, J., Roes, J., and Hynes, R.O. (2005). Selective ablation

of alphav integrins in the central nervous system leads to cerebral hemorrhage,

seizures, axonal degeneration and premature death. Development *132*, 165-

176. 10.1242/dev.01551.

112.   Beis, D., Bartman, T., Jin, S.W., Scott, I.C., D'Amico, L.A., Ober, E.A.,

Verkade, H., Frantsve, J., Field, H.A., Wehman, A., et al. (2005). Genetic and

cellular analyses of zebrafish atrioventricular cushion and valve development.

Development *132*, 4193-4204. 10.1242/dev.01970.

113.   Jin, S.W., Beis, D., Mitchell, T., Chen, J.N., and Stainier, D.Y. (2005). Cellular

and molecular analyses of vascular tube and lumen formation in zebrafish.

Development *132*, 5199-5209. 10.1242/dev.02087.

114.   Chang, N., Sun, C., Gao, L., Zhu, D., Xu, X., Zhu, X., Xiong, J.W., and Xi,

J.J. (2013). Genome editing with RNA-guided Cas9 nuclease in zebrafish

embryos. Cell Res *23*, 465-472. 10.1038/cr.2013.45.

115.   Jao, L.E., Wente, S.R., and Chen, W. (2013). Efficient multiplex biallelic

zebrafish genome editing using a CRISPR nuclease system. Proc Natl Acad

Sci U S A 110, 13904-13909. 10.1073/pnas.1308335110.

116.   Kawakami, K. (2004). Transgenesis and gene trap methods in zebrafish by

using the Tol2 transposable element. Methods Cell Biol *77*, 201-222.

10.1016/s0091-679x(04)77011-9.

117.   Lei, L., Yan, S.Y., Yang, R., Chen, J.Y., Li, Y., Bu, Y., Chang, N., Zhou, Q.,

Zhu, X., Li, C.Y., and Xiong, J.W. (2017). Spliceosomal protein eftud2

mutation leads to p53-dependent apoptosis in zebrafish neural progenitors.

Nucleic Acids Res *45*, 3422-3436. 10.1093/nar/gkw1043.

118.   Pang, M., Bai, L., Zong, W., Wang, X., Bu, Y., Xiong, C., Zheng, J., Li, J.,

Gao, W., Feng, Z., et al. (2020). Light-sheet fluorescence imaging charts the

gastrula origin of vascular endothelial cells in early zebrafish embryos. Cell

Discovery *6*, 74. 10.1038/s41421-020-00204-7.

1493

Figure 1

Figure 2

A *Tg(gfap*:GFP)
48 hpf
betaPix^ct/ct
betaPix^m/m
Scale bar:100 μm

B
P=0.0017
Glia length (μm)
betaPix^ct/ct   betaPix^m/m

P<0.0001
Normalized glia length index
betaPix^ct/ct   betaPix^m/m

C
nestin   *pax2a*
48 hpf
betaPix^ct/ct
17/18   17/17
betaPix^m/m
15/20   17/21
Scale bar: 500 μm

D *Tg(gfap*:GFP); *Tg(kdrl:mCherry)*
48 hpf
sibling
Z=0 BA   Z=1 BA   Z=2 BA   Z=3 BA
betaPix KO
Z=0 BA   Z=1 BA   Z=2 BA   Z=3 BA
Scale bar: 200 μm

E *Tg(sox2*:GFP); *Tg(kdrl:mCherry)*
48 hpf
sibling
betaPix KO
Scale bar: 150 μm

F *o-*dianisidine
48 hpf
betaPix^ct/ct
DMSO
IPA3
betaPix^m/m
DMSO
IPA3
Scale bar:150μm

G
P<0.0001
P=0.0016
Hemorrhage %
DMSO   IPA3

P<0.0001
P=0.0007
Hemorrhage area (mm²)
DMSO   IPA3
● betaPix^ct/ct   ● betaPix^m/m

H *Tg(gfap:GFP)*; *Tg(kdrl:mCherry)*
48 hpf
betaPix^ct/ct
DMSO
IPA3
betaPix^m/m
DMSO
IPA3
Scale bar:150μm

I
P=0.0001
P=0.0176
Normalized CtA length index
DMSO   IPA3

J
P<0.0001
P<0.0001
Normalized glia length index
DMSO   IPA3
● betaPix^ct/ct   ● betaPix^m/m

Figure 3

A — *o*-dianisidine

48 hpf

betaPix^ct/ct^ — DMSO, IPA3

gfap:GFP-Cre; betaPix^ct/ct^ — DMSO, IPA3

Scale bar:150μm

B — *Tg(kdrl:mCherry)*

betaPix^ct/ct^ — DMSO, IPA3

gfap:GFP-Cre; betaPix^ct/ct^ — DMSO, IPA3

Scale bar:150 μm

C

Hemorrhage % — $P<0.0001$ $P<0.0001$ DMSO IPA3

Hemorrhage area (mm^2) — $P<0.0001$ $P<0.0001$ DMSO IPA3

Normalized CtA length index — $P<0.0001$ $P=0.0079$ DMSO IPA3

● *betaPix*^ct/ct^
● *gfap:GFP-Cre; betaPix*^ct/ct^

Figure 4

A Inject CRISPR system containing PBS or betaPix-gRNAs

Hemorrhage initiation

hours post fertilization

0   24   36   48

Harvest head   Harvest head

B

Erythrocytes   Endothelial
Cartilage   Olfactory bulb
Pharyneal arch
Muscle   Retina neurogenesis
Cranial ganglion   Dorsal diencephalon
Pigment cell (iridophore)   Neuron   Hindbrain
Retina (photoreceptor precursor cells)   Ventral diencephalon   Progenitors
Pigment cell (neural crest derived)   Neural crest retina (amacrine)   Ventral midbrain
    Floor plate
Otic vesicle
Periderm   Neutrophils
Epidermis

UMAP2

UMAP1

C

Ratio of Cells

Conditions: ctrl_1d ko_1d ctrl_2d ko_2d

Cell Type
- Neuron
- Pharyneal arch
- Progenitors
- Retina neurogenesis
- Erythrocytes
- Ventral diencephalon
- Cartilage
- Ventral midbrain
- Dorsal diencephalon
- Retina (photoreceptor precursor cells)
- Muscle
- Epidermis
- Olfactory bulb
- Cranial ganglion
- Otic vesicle
- Hindbrain
- Endothelial
- retina (amacrine)
- Periderm
- Neural crest
- Neutrophils
- Pigment cell (iridophore)
- Pigment cell (neural crest derived)
- Floor plate

D Progenitors ctrl_2d vs ko_2d enrich GO terms

- chromatin silencing
- negative regulation of transcription, DNA-templated
- nucleosome positioning
- nucleosome assembly
- negative regulation of DNA metabolic process
- regulation of DNA recombination
- chromatin organization
- DNA packaging
- microtubule depolymerization
- negative regulation of cellular component organization
- regulation of microtubule polymerization or depolymerization
- defense response to Gram-positive bacterium
- microtubule polymerization or depolymerization
- chaperone cofactor-dependent protein refolding
- neurotransmitter secretion
- regulation of microtubule cytoskeleton organization
- cellular protein complex disassembly
- neurotransmitter transport
- response to heat
- microtubule cytoskeleton organization

-log10(p.adjust)   0   5   10   15   20   25

E

stmn1a, stmn1b, stmn2a, stmn2b, stmn3, stmn4, stmn4l

Conditions: ctrl_1d ko_1d ctrl_2d ko_2d

Progenitors.ctrl_1d Progenitors.ko_1d Progenitors.ctrl_2d Progenitors.ko_2d

F
- $betaPix^{ct/ct}$ glia
- $betaPix^{m/m}$ glia

Relative expression

betaPix   stmn1a   stmn1b   stmn4l   ppfia3

****   ****   ***   ****   ****

G
- $betaPix^{ct/ct}$ glia
- $gfap:GFP-Cre; betaPix^{ct/ct}$ glia

Relative expression

betaPix   stmn1a   stmn1b   stmn4l   ppfia3

****   ***   ****   ****   ****

H

	stmn1a	*stmn1b*	*stmn4l*
$betaPix^{ct/ct}$	48 hpf 24/24	24/24	24/24
$gfap:GFP-Cre; betaPix^{ct/ct}$	20/20	19/20	25/29

Scale bar: 500 μm

Figure 5

Figure 6

Figure 7