Glial *betaPix* is essential for blood vessel development in the zebrafish brain

ShihChing Chiu[1], Qinchao Zhou[1], Chenglu Xiao[2], Linlu Bai[1], Xiaojun Zhu[1,*], Wanqiu

Ding[1,*], and Jing-Wei Xiong[1,2,*,#]

[1]Beijing Key Laboratory of Cardiometabolic Molecular Medicine, Institute of

Molecular Medicine, and College of Future Technology, Peking University, Beijing

100871, China; [2]School of Basic Medical Sciences, The Second Affiliated Hospital,

Institute of Biomedical Innovation, The MOE Basic Research and Innovation Center

for the Targeted Therapeutics of Solid Tumors, Jiangxi Medical College, Nanchang

University, Nanchang 330031, China

*Corresponding authors: WD (dingwq@pku.edu.cn), XZ (zhuxiaojun@pku.edu.cn),

and JWX (jingwei_xiong@pku.edu.cn)

#Lead Contact: Jing-Wei Xiong, Ph.D. (jingwei_xiong@pku.edu.cn)

Abstract

The formation of blood-brain barrier and vascular integrity depends on the

coordinative development of different cell types in the brain. Previous studies have

shown that zebrafish *bubblehead* (*bbh*) mutant, which has mutation in the *betaPix*

locus, develops spontaneous intracerebral hemorrhage during early development.

However, it remains unclear in which brain cells *betaPix* may function. Here, we

established a highly efficient conditional knockout method in zebrafish by using

homology-directed repair (HDR)-mediated knockin and knockout technology, and

generated *betaPix* conditional trap (*betaPix^{ct}*) allele in zebrafish. We found that

betaPix in glia, but neither neurons, endothelial cells, nor pericytes, was critical for

glial and vascular development and integrity, thus contributing to the formation of

blood-brain barrier. Single-cell transcriptome profiling revealed that microtubule

aggregation signaling stathmins and pro-angiogenic transcription factors *Zfhx3/4* were

down-regulated in glial and neuronal progenitors, and further genetic analysis

suggested that betaPix may act upstream on the PAK1-Stathmin and Zfhx3/4-Vegfaa

signaling to regulate glia migration and angiogenesis. Therefore, this work reveals

that glial *betaPix* plays an important role in brain vascular development in zebrafish

embryos and possibly human cells.

Keywords: *betaPix*; hemorrhage; glial migration; knock-in; stathmin

Introduction

Intracerebral hemorrhage (ICH) is a life-threatening stroke type with the worst prognosis and few proven clinical treatments. Most patients who survive intracerebral hemorrhage end up with disabilities and are at risk for recurrency, cognitive decline, and systemic vascular issues, making this condition particularly significant among neurological disorders [1,2]. Several rodent models have been used for modeling ICH, including autologous blood injection, collagenase injection, thrombin injection, and micro-balloon inflation techniques [3]. Zebrafish (Danio rerio) exhibit a closed circulatory system and regulatory pathways that are highly conserved among vertebrates [4] with relatively cheap and easy to maintain disease models. Over the past decades, varieties of zebrafish mutants that spontaneously developed brain hemorrhage have been generated in mutagenesis screens, such as *redhead* [5], *reddish* [6,7] and *bubblehead* (*bbh*) [8,9]. Elucidating mutated gene function in these zebrafish models enables us to gain insights into the genetic basis and pathophysiological mechanisms of ICH development. The *bbh* mutant was identified independently in two large-scale ethylnitrosourea-induced mutagenesis screens [8,9] and positional cloning revealed that p21-activated kinase (Pak)-interacting exchange factor beta (*betaPix*) gene was mutated in *bbh* mutants [10]. *betaPix* contains SH3 domain that binds group I PAKs, and Dbl homology (DH) and pleckstrin homology (PH) domains that function as a RhoGEF to interact with Rac/Cdc42 small GTPases [11], thus participating in multiple cellular pathways to regulate cell polarity, adhesion and migration [12].

Various studies have reported that brain vessels are highly supported by neurons and

glial cells during development and regeneration [13,14]. Glial-specific *betaPix* have been

proposed to interact with *αvβ8* integrin and the Band 4.1s in glia, which link to

multiple intracellular signaling effectors such as *Wnt7a* and *Wnt7b* [15], but this

hypothesis was not experimentally addressed [16]. To this end, *betaPix* has been shown

to interact with *αvβ8* integrins and mediate focal adhesion formation and thus

modulate cerebral vascular stability in *bbh* zebrafish mutant [17]. Integrins are the main

molecular link between cells and the extracellular matrix (ECM) which serves as

scaffolds in the perivascular space. Based on a chemical suppressor screening of *bbh*

hemorrhages, we have previously reported that a small molecule called miconazole

downregulated the pErk-matrix metalloproteinase 9 (Mmp9) signaling to reduce ECM

degradation, thus improving vascular integrity [18]. However, how *betaPix* cell-type

specifically contributes to vascular integrity and maturation remains unclear. Here, we

generated *betaPix* conditional trap (*betaPix^ct*) alleles by using a CRISPR/HDR-based

Zwitch method, and found that *betaPix* acted mainly in glia to regulate glial and

vascular development and integrity *via* Stathmin and Zfhx3/4 signaling. Therefore,

this work provides the first experimental evidence that *betaPix* acts in glia for

vascular integrity development, and its functional conservation between zebrafish and

human cells may further guide us to decipher the genetic basis of ICH in the future.

Results

Generating *betaPix* conditional trap (*betaPix^ct^*) allele by an HDR-mediated

knockin and knockout method

To study *betaPix* function in cell-specific manner, we utilized a donor vector with an

invertible gene trap cassette *via* HDR (Zwitch) [19-21] for generating *betaPix* conditional

trap (*betaPix^ct^*) alleles. Zwitch consists of a splice acceptor conjugated to a red

fluorescent protein *via* a 2A self-cleaving peptide, flanked by two LoxP sites and two

Lox 5171 sites in reciprocal orientations. Outside this conditional trap cassette, we

included a lens-specific enhanced green fluorescence protein *α*-crystallin:EGFP for

screening knockout founders, as well as inserted both left and right arms for targeting

homologous sequences on the genome. We modified the Zwitch vector by adding a

glycine-serine-glycine (GSG) spacer in front of the 2A peptide to enhance cleavage

efficiency, and adding universal guide RNA (UgRNA)[22] target sequences outside the

left and right homologous arms for linearizing donor vector (Figures 1A and S1A).

We chose CRISPR/Cas9 system to generate double-strand breaks (DSBs) on intron 5

of the *betaPix* locus, and chose the most efficient guide RNAs based on their

efficiency to induce CRISPR indels. We used either long arms such as ~1000 bp or

short arms such as 24 bp that located upstream and downstream the guide RNA sites.

We then co-injected the donor vectors with targeting guide RNA, universal guide

RNA, and Cas9 mRNA into one-cell-stage embryos. At 4 days post-fertilization (dpf),

we selected *α*-crystallin-EGFP-positive embryos (F_0) for raising to adulthood, and

pre-selected F_0 founders by examining inheritable EGFP reporter expression of F_1

embryos. To confirm the correct homologous recombination in the *betaPix* locus, we

identified potential founders by genomic PCR to examine Zwitch insertion in the

betaPix locus. Precise knock-in genotypes were further verified using Sanger

sequencing (Figure 1B, 1C and S1B). Among 184 adult F_0 founders with

α-crystallin-EGFP expression, we found 84 founders had EGFP-positive F_1 embryos,

and 7 founders had expected PCR fragments around both 5' and 3' arms, which two

founders had correct insertions in the *betaPix* locus by Sanger sequencing (Figure

S1C), thus achieving ~1% efficiency on generating conditional alleles. Briefly, we

established an efficient method for generating conditional knockin/knockout mutants

in general, and generated a conditional gene trap line Ki(*betaPix*:Zwitch) in the

betaPix locus in particular, which is referred to as *betaPix*^{ct} hereafter.

A previous work has shown that *bubblehead* (*bbh*^{fn40a}) mutant has a global reduction

in *betaPix* transcripts, and *bbh*^{m292} mutant has a hypomorphic mutation in *betaPix,*

thus establishing that *betaPix* is responsible for *bubblehead* mutant phenotypes[10].

bbh^{fn40a} mutants developed cerebral hemorrhages phenotype between 36 to 52 hpf

(Figure S1D and S1E). To characterize the *betaPix*^{ct} allele, we injected one-cell-stage

betaPix$^{ct/ct}$ embryos with *Cre* mRNA to induce global inversion of Zwitch and

deletion of *betaPix*. As expected, we found *Cre*-induced precise inversion in

betaPix$^{m/m}$ as confirmed by Sanger sequencing (Figure 1C). By using *o*-dianisidine

staining to label hemoglobins, we found severe brain hemorrhages developed in

betaPix$^{m/m}$ mutant with the time window similar to *bbh*^{fn40a} mutants (Figure 1D and

1E). Consistently, qPCR analysis showed that *betaPix* decreased and *RFP* increased

in heterozygous *betaPix^{m/+}* and homozygous *betaPix^{m/m}* mutants compared with

betaPix^{ct/ct} wild-type siblings, while *alphaPix* expression was not affected (Figure 1F).

Of note, *TagRFP* expression in *betaPix^{m/+}* (after Cre-mediated recombination in

betaPix^{ct/+} embryos) reflected endogenous *betaPix* expression in the boundary of

midbrains and hindbrains as well as the hindbrains by light-sheet fluorescence

microscopy imaging (Figure S1F and S1G). Together, these results suggest that the

conditional trap *betaPix^{ct/ct}* zebrafish is functional for visualizing endogenous *betaPix*

expression (knockin) and performing loss-of-function study (conditional knockout).

betaPix^{m/m} mutant has brain hemorrhages, central arteries defects and abnormal

glial structure that is partially rescued by Pak1 inhibitor IPA3 treatment

The main phenotypes in *bbh* mutants consist of brain hemorrhage and hydrocephalus

as early as 36 hpf, as well as poor endothelial–mesenchymal contacts and defective

central arteries (CtAs) sprouting in the hindbrain [10,17]. To demonstrate the

practicability of *betaPix* conditional trap lines, we assessed reported phenotypes

associated with *betaPix* deficiency using homozygous *betaPix^{ct/ct}* zebrafish. Global

betaPix inactivation with *Cre* mRNA injection resulted in severe cerebral

hemorrhages (Figure 1D and 1E). By crossing the *betaPix^{m/m}* mutant line with

Tg(*kdrl*:GFP) transgenic line, we found defective angiogenesis in hindbrain central

arteries *via* light-sheet fluorescence microscopy imaging (Figure S2A and S2B),

which is consistent with *betaPix* mutant phenotypes as reported [10,18].

It has been suggested that endothelial *betaPix* might not contribute to the occurrence

of brain hemorrhage [10]. During brain development, neurons and glia are important

perivascular cells and they orchestrate with endothelial cells in a temporal and spatial

pattern. Transgenic zebrafish that express fluorescent proteins driven by the glial

fibrillary acidic protein (*gfap*) promoter have been widely used for labeling glial

population [26]. By crossing the *betaPix^{m/m}* mutant line with Tg(*gfap*:GFP) transgenic

line, we found that radial glial structure of the hindbrain had disoriented arrangement

and shorten process length in *betaPix^{m/m}* embryos compared to siblings by light-sheet

fluorescence microscopy imaging (Figure 2A, 2B and S2C). RNA *in situ*

hybridization showed that neuronal and glial precursors marker *nestin* increased while

differentiated neuronal marker *pax2a* decreased at the hindbrain of *betaPix^{m/m}*

embryos, suggesting delayed neuronal development and differentiation after *betaPix*

inactivation (Figure 2C). By using global *betaPix* knockout with multiple guide-RNA

simultaneously [23], we found that CRISPR-induced *betaPix* mutant F_0 embryos had

almost identical phenotypes to those in *betaPix^{fn40a}* mutants (Figure S3A and S3B),

such as cerebral hemorrhages, central arteries defects, and abnormal hindbrain glial

and neuronal precursor development (Figure S3C to S3G), further confirming the

loss-of-function phenotypes in *betaPix^{m/m}* mutants.

Vascularisation of the central arteries in the zebrafish hindbrain starts at 29 hpf. Tip

cells start to form in the dorsal side of the primordial hindbrain channels (PHBC),

sprout and migrate into the hindbrain tissue, moving towards the center around 36 hpf

to establish connections with the basilar artery. A characteristic arch architecture

forms at 48 hpf as boundaries dividing the hindbrain into rhombomeres[24]. In

Tg(*gfap*:GFP);Tg(*kdrl*:mCherry) double transgenic embryos, CRISPR-induced

betaPix knockouts showed that CtA sproutings were restricted to the bottom of glial

processes that failed to migrate upwards to form an arched structure in compared with

non-injected siblings (Figure 2D). The transcription factor SRY box-2 (Sox2) is

another marker gene for neuron and glial precursors. In Tg(*sox2*:GFP);

Tg(*kdrl*:mCherry) transgenic background, *Sox2*-positive precursor cells tightly

wrapped around the central artery in control siblings, but *Sox2* precursor cells had

loose contacts with the central arteries, with larger perivascular distances in

CRISPR-induced *betaPix* mutants (Figure 2E). These results implicate the impaired

interactions between endothelial cells and neuronal/glial cells during development

after *betaPix* deletion.

p21-activated kinase (Pak) is a binding partner for *betaPix*. *Pak2a* has been shown

mediating downstream signaling in bbh^{m292} mutants. Group I PAK allosteric

inhibitor IPA-3 covalently modifies and stabilizes the autoinhibitory N-terminal

region of PAK1, PAK2 and PAK3 [25]. As expected, IPA-3 treatment significantly

decreased incidence and intensity of cerebral hemorrhages in $betaPix^{m/m}$ mutant

embryos (Figure 2F and 2G), while IPA-3 treatment had no effect on hemorrhage

induction in $betaPix^{ct/ct}$ control siblings. In addition, central arterial defects were

partially rescued in *betaPix^(m/m)* mutant embryos after IPA-3 treatment, showing

increased endothelial protruding into the hindbrain and more arch structure formation

(Figure 2H and 2I). IPA-3 treatment also decreased abnormal glial process

arrangements, which the lengths of glial processes statistically reached the levels of

the control siblings (Figure 2J). Thus, these data suggest that the *betaPix-Pak1/2*

signaling regulates brain vascular integrity during development.

Glial-specific *betaPix* knockouts recapture its global knockout phenotypes

To investigate *betaPix* function in different types of perivascular cells, we generated

glial- or neuronal-specific transgenic lines using either *gfap* [26] or *huC* [27] promoters to

drive *EGFP-Cre* fusion gene expression under *betaPix^(ct/ct)* background, respectively

(Figure S4A). EGFP signals reported *Cre* expression while RFP signals efficiently

reported *Cre*-induced gene trap cassette inversion and disruption of *betaPix*

expressions (Figure S4B to S4F). Interestingly, glial-specific *betaPix* knockouts

developed severe cerebral hemorrhages and abnormal central arteries vascularization,

which were partially rescued by IPA-3 treatment (Figure 3A, 3B and 3C),

highlighting *betaPix* function in glia *via* Pak1/2 signaling. On the other hand, neither

vascular-, neuronal-, nor mural-specific deletion of *betaPix* had evident *betaPix*

mutant phenotypes (Figure S5A to S5D). These results suggest that glial *betaPix*

plays crucial roles in zebrafish embryonic vascular integrity and glial development *via*

regulating *Pak* activities.

Single-cell transcriptome profiling reveals that *gfap*-positive progenitors were

affected in *betaPix* knockouts

To investigate the interplays between glial and hemorrhagic pathology caused by

betaPix loss-of-function, we profiled the cranial tissues of CRISPR-edited zebrafish at

1 and 2 dpf using the 10X Chromium single-cell RNA sequencing (scRNA-seq)

platform (Figure 4A). Multi-guide targeting was able to generate almost 100% F_0 null

mutants, allowing us to select mutant embryos before brain hemorrhages starting from

36 hpf. Low-quality cells were excluded based on the numbers of genes detected and

percentages of reads mapped to mitochondrial genes per sample. A total of 38,670

cells passed quality control and were used for subsequent analyses. Uniform manifold

approximation and projection (UMAP) identified 71 cell clusters, which represented

24 zebrafish cranial cell types based on known marker gene expression profiles

(Figure 4B and S6A). Enriched gene markers were compared with previously

annotated gene markers in the ZFIN database and literatures[28,29]. By comparing the

proportion of cells in each sample, we found that most neuronal clusters had increased

relative proportions, while the numbers of glial and neuronal progenitors, endothelial

cells, erythrocytes, neural crests, muscles, cartilages, retinas (photoreceptor precursor

cells), olfactory bulbs, or epidermis and pharyneal arches were reduced in *betaPix*

knockout heads at 2 dpf (Figure 4C and S6B). This is consistent with the open public

databases of single-cell transcriptome atlas where *betaPix* is weakly expressed in a

broad range of cells including glia, neurons and neuronal precursors, retinas,

endothelial cells, pharynx, exocrine pancreas, olfactory cells and heart cells [30]. Given

that *betaPix* is critical in glia during brain vascular integrity development as shown

above (Figures 1 and 2), we examined the *gfap* expression among each cluster, and

found relatively high *gfap* expression in clusters including glial and neuronal

progenitors, hindbrain, ventral diencephalon, ventral midbrain and floor plate (Figure

S6A, indicated by the arrow). We next focused on the progenitor cluster owing to the

enriched *gfap* expression and the significantly reduced numbers of cells in this cluster

by *betaPix* deficiency. Furthermore, this progenitor cluster exhibited high-level

expression of cell proliferation and cell cycle related genes (*mki67*, *pcna*, *ccnd1*, *rrm1*

and *rrm2*) as well as key glial-associated genes (*gfap*, *fabp7a*, *her4*.1, *cx43*, *id1*,

fgfbp3, *atp1a1b* and *mdka*) (Table S1). From differentially expressed genes (DEGs) in

this progenitor cluster between controls and *betaPix* knockouts at 2 dpf, gene

ontology (GO) terms revealed three major categories: epigenetic remodeling,

microtubule organizations, and neurotransmitter secretion/transportation (Figure 4D).

Of these signaling genes, we were particularly interested in microtubule organizing

genes for further studies.

Stathmin acts downstream of *betaPix* in glial migration *via* regulating tubulin

polymerization

Microtubules are essential cytoskeletal elements composed of α/β-tubulin

heterodimers. The regulation of microtubule polymerization affects important cellular

functions such as mitosis, motor transport, and migration [31]. Based on the above

scRNA-seq data (Figure 4E and S6C), we examined if microtubule-destabilizing

protein Stathmins were affected in *betaPix* mutants. We found that *stathmin1a*

(*stmn1a*), *stathmin1b* (*stmn1b*), and *stathmin4l* (*stmn4l*) were relatively abundant in

both the whole brain and the progenitor cluster, and decreased specifically at 2 dpf in

betaPix knockout progenitor cells (Figure 4E). Subsequent qRT-PCR revealed that

these *Stathmin* expression decreased in both global *betaPix* knockouts (*betaPix$^{m/m}$*)

and glial-specific *betaPix* knockout glia (*gfap*:EGFP-Cre;*betaPix$^{ct/ct}$*) (Figure 4F and

4G), which was further verified by RNA *in situ* hybridization analysis (Figure 4H).

We next investigated whether *Stathmins* act downstream to *betaPix* signaling on

regulating vascular integrity development. To this end, we found that *Pak1/2*

inhibition enabled to rescue down-regulated expressions of *stmn1a*, *1b* and *4l* in either

global *betaPix$^{m/m}$* or glial-specific *betaPix* knockout zebrafish (Figure 5A, 5B and

S7A), which is consistent with previous findings that *Stathmin-1* acts downstream to

betaPix-Pak signaling in neurite outgrowth in mice[32].

To test whether *Stathmin* family genes are important for vascular stability, we utilized

CRISPR-induced F_0 knockout system simultaneously targeting the three stathmin

genes. While singular stathmin gene knockout led to no evident phenotypes (data not

shown), we found that simultaneously inactivating *stmn1a*, *stmn1b* and *stmn4l* caused

brain hemorrhages and hydrocephalus in the brain, similar to that from *betaPix^{fn40a}*

and betaPix$^{m/m}$ mutants (Figure 5C, 5D, and S7B to S7E). By labeling both glia and

vasculature with Tg(*gfap*:GFP; *kdrl*:mCherrry) transgenic embryos, we found that

triple stathmin gene knockouts had impaired CtAs, abnormal glia and neuronal

development in the hindbrain region (Figure 5E, 5F, and S7F to S7H). Furthermore,

overexpressing *stmn1b* by the *gfap* promoter partially rescued brain hemorrhages and

delayed neuronal development in *bbh^{fn40a}* mutants (Figure 5G, 5H, S7I and S7J), of

which glial processes elongation also improved, but CtAs defects failed to restore

(Figure 5I, 5J and S7K). Thus, these data support the critical role of *stathmins* in glia.

To examine the direct role of *betaPix-stathmin* in glia, we turned to use human

glioblastoma cell lines and manipulated the *betaPIX* levels by siRNA knockdown or

transgenic overexpression. We found that *betaPIX* knockdown attenuated glial

migration (Figure 5K, 5L, and S7L to S7N), which is consistent with previous studies

on *betaPix* function in the endoderm [33], epithelial cells [34], neurons [35] and multiple

carcinoma cell lines including glioblastoma [36-38]. Either *betaPIX* or *STMN1*

overexpression led to a partial restoration of the impaired glial migration, suggesting

Stathmins as downstream effectors of *betaPix*. Furthermore, immunofluorescence

analysis revealed that *betaPIX* knockdown led to abnormal accumulation of

microtubules at the cell periphery, whereas ectopic expression of *betaPIX* and

Stathmin partially rescued these microtubule defects (Figure 5M and

5N). Together, these results suggest that *betaPix* depletion disrupts

microtubule dynamics and leads to impaired glial development acting upstream to the

Pak-Stathmin signaling, subsequently disrupting vascular integrity and resulting in

brain hemorrhages. While the brain hemorrhages were only partially rescued and

abnormal CtAs defects failed to restore, indicating that additional pathways may act

downstream of *betaPix*.

Zfhx3/4 acts downstream of *betaPix* in regulating vascular integrity development.

Previous studies have identified multiple signaling pathways important for zebrafish

hindbrain angiogenic sprouting, such as VEGF, Notch, Wnt, Integrin β1,

angiopoietin/TIE, and insulin-like growth factor signaling [39-42]. To investigate whether

angiogenic signal is disrupted by *betaPix* depletion, we examined several angiogenic

gene expression pattern by performing RNA *in situ* hybridization of embryos at 36

hpf. The expression level of *Vegfaa* in sprouting CtAs decreased in *bbh*[fn40a] mutants

(Figure 6A). Consistently, *betaPix* knockdown significantly reduced *VEGFA*

expression in cultured glial cells, which was rescued by ectopic expression of *betaPix*

(Figure 6B; right panel). To explore how *betaPix* regulate *Vegf* signaling in glia, we

re-examined the GO analysis of single-cell sequencing data and found a significant

enrichment of epigenetic and transcriptional regulation in glial progenitor cluster

(Figure 4D). Among these candidate genes, transcription factor Zinc Finger

Homeobox 3 (ZFHX3) is known to mediate hypoxia-induced angiogenesis in

hepatocellular carcinoma *via* transcriptional activation of VEGFA [43]. Furthermore,

ZFHX3 has been associated with stroke in multiple genome-wide association studies

[44-46]. Interestingly, we found that transcription factors *ZFHX3* and *ZFHX4*

significantly downregulated after *betaPIX* inactivation in cultured glial cells (Figure

6B). The transcription levels of *Zfhx3* and *Zfhx4* also downregulated in glial

progenitor cluster of our single cell RNA sequencing data upon *betaPix* knockout at 2

dpf (Figure S8A), as well as the glia-specific *betaPix* mutants (Figure S8B), thus

suggesting that Zfhx3/4 might acts as downstream effectors of *betaPix*.

To determine whether *Zfhx3/4* is critical for vascular integrity by interacting with

betaPix in zebrafish, we depleted *Zfhx3* and *Zfhx4* simultaneously *via*

CRISPR-mediated F_0 knockouts (Figure S8C to S8E). We found that *Zfhx3/4*

knockouts decreased *Vegfaa* expression at sprouting CtAs at 36 hpf (Figure 6C).

Zfhx3/4 knockouts at 48 hpf had under-developed CtAs and reduced penetration of

endothelial cells into the hindbrain (Figure 6F and 6G), developed cerebral

hemorrhages during 36 to 52 hpf, which are comparable with *betaPix* mutant

phenotypes (Figure 6D to 6G). Despite up-regulated expression level of *nestin*, glial

arrangement and processes elongation remained statistically unaffected after *Zfhx3/4*

inactivation (Figure 6F, 6G and S8F). We next investigated whether increasing

expression level of *Vegfaa* in *betaPix* mutants was able to rescue impaired vascular

integrity. By using *Zfhx4* mRNA microinjection into one-cell stage embryos in either

bbh^{fn40a} mutants or CRISPR-induced *betaPix* knockouts, we found that ectopic

expression of *Zfhx4* increased *Vegfaa* expression level (Figure 6L), thus leading to a

drastic reduction of both percentages and volumes of brain hemorrhage shown in

these global *betaPix* mutants (Figure 6H, 6I, and S8G to S8I), as well as improved

CtAs and glial development (Figure 6J and 6K). To further explore whether

angiogenic potential altered in the endothelial cells, we re-examined our scRNA-seq

data and noted significant reduction of endothelial proportions among *betaPix*

knockouts (Figure S6B), which showed similar trend as glial progenitors. In this

endothelial cluster, CRISPR-induced *betaPix* knockouts had down-regulated

angiogenic gene expression at 2 dpf (Figure 6M). Despite the lack of endothelial

phenotype in cellular resolution in our present results, the disruptive transcriptome

signature in endothelial cluster suggests a potential link to the defective central

arterial development. Together, these data support the critical role of *Zfhx3/4* in

vascular integrity and glial development probably acting upstream to VEGFA and

downstream to *betaPix*.

Discussion

Conditional knockout technology in mice has been widely used to modify and

determine targeted gene function in a cell-specific manner. Spatio-temporal specific

modification is beneficial for studying embryonic lethality genes or investigating the

function of targeted genes in specific cell populations. Despite being broadly utilized

in mice, conditional knockouts in other animal models are limited due to technical

difficulties. To this end, multiple research groups have designed different strategies

for establishing floxed-mutant alleles, gene traps, and inducible Cas9 in zebrafish [47].

In particular, the Zwitch gene trap-based approach has been established successfully

in zebrafish, enabling to simultaneously produce knockin reporter driven by the

endogenous gene promoter and disrupting endogenous gene function in specific cell

types [19-21]. We adapted this strategy with the aid of CRISPR/Cas9 technology, and

added a GSG spacer to enhance cleavage efficiency and universal guide RNA target

sites on both homologous arms for producing linearized targeting vector. Moreover,

we also found that utilizing either short or long homologous arms enables to achieve

precise targeted integrations with this donor vector. Our short arms-based method can

be broadly adapted because of less restrictions on arm sequences and easily

engineering the donor vector. From the *betaPix* locus and other five unpublished loci

that we already generated, we achieved Zwitch alleles with ~1% correct insertions

based on homologous recombination as well as *Cre*-induced, cell-specific deletion

and functional analysis of targeted genes in both embryos and adult hearts. Therefore,

our CRISPR-based Zwitch method can be widely applied for generating conditional

mutant zebrafish in particular and possibly extended for other animal mutants in

general.

The maturation hallmark of central nervous system (CNS) vasculature is acquisition

of blood brain barrier (BBB) properties, establishing a stable environment crucial for

brain homeostasis in response to extrinsic factors and physiological changes. The core

features of the BBB include specialized tight junctions, highly selective

transporters and limited immune cell trafficking [48,49]. In zebrafish, BBB can be

functionally characterized from 3 dpf [50,51]. *Bubblehead* mutants acquire vessel rupture

phenotype early before BBB maturation, presenting a good hemorrhagic model for

studying early BBB development in this work. Multiple types of cells contribute to

diverse aspects of BBB development and maintenance, leading to proper vasculature

function across developmental timeline. Endothelial cells form the continuous inner

layer of blood vessels with junctional proteins and selective transporters, which are

the main subject of permeability regulation in BBB. It is well accepted that

endothelial cells do not show a predetermined role and that the brain environment is

sufficient to induce endothelial barrier [52]. Nevertheless, accumulative evidences point

to critical role of endothelial angiocrine signals in regulating several aspects of BBB

development. For example, Wnt7a/b ligands secreted by neural progenitor cells bind

to Frizzled receptors on endothelial cells, activating canonical Wnt signaling and

downstream genes. WNT ligands knockouts lead to angiogenic defects and BBB

leakages specifically in the CNS [53]. Endothelial GPR124, as a coactivator of canonical

Wnt signaling, has important function in CNS-specific angiogenesis and BBB

establishment [54-56]. Other signaling ligands such as angiopoietins and Ephrin family

ligands orchestrate with Tie or EphB receptors on endothelial cells, respectively[57].

Complex interactions between BBB cell types orchestrate proper angiogenesis and

CNS development in a spatio-temporal manner. Astrocytes ensheathe capillaries

through polarized end feet that is enriched with aquaporin-4 (Aqp4) proteins,

colocalized with inwardly rectified K^+ channels[58,59]. Functional coupling of ion and

water fluxes plays a critical role in regulating local osmotic equilibrium. During

development, radial glia is a neuroepithelial origin with heterogeneous populations

that are able to generate neurons, astrocytes, and oligodendrocytes[60-63]. In zebrafish,

radial glia has long been considered serving an astrocytic role. Radial glia in zebrafish

enriches with several key molecular markers for astrocytes and tight junctions while

Aqp4[+] radial glial processes rarely contact the vasculature[64]. A more recent article has

reported that zebrafish *Glast*-expressing radial glia transform into astrocyte-like cells,

displaying dense cellular processes, tiling behavior and proximity to synapses [65].

Whether end feet of these astrocyte-like cells enwrap capillary blood vessels and

resemble mammalian astrocytes warrants future investigations. In agreement with

mammalian astrocytes, two independent groups have shown that ablation of pan-glia

results in progressive brain hemorrhage[66,67] or deficiency of spinal cord arteries

development in zebrafish [68], demonstrating important role of zebrafish glia in BBB

formation. Multiple glia/astrocyte-derived signals have been shown to contribute to

endothelial barrier properties development. Astrocyte-expressed sonic hedgehog (Shh)

bind to hedgehog (Hh) receptors on endothelial cells and contribute to the BBB

functions by promoting junctional protein expression and the quiescence of the

immune system [69-72]. Before BBB formation, notochord-derived Shh activity promote

arterial cell fate on developing endothelial cells [73]. Shh stimulates the production of

angiogenic cytokines, angiopoietins and interleukins, *via* downstream transcription

factor *Gli* or non-canonical pathways crosstalks with iNOS/Netrin-1/PKC,

RhoA/Rock, ERK/MAPK, PI3K/Akt, Wnt/β-catenin, and Notch signaling pathways [74].

By establishing conditional alleles of *betaPix*, we determined the critical roles of

betaPix in glia, but not in other brain cells, on its cerebral vessel development and

integrity. However, it remains unclear how glial betaPix is related to the above

endothelial signaling to regulate endothelial cell function and blood vessel integrity.

It has been known that Pix and Pak participate in multiple signal transduction

pathways [12]. We are interested in distinctive zebrafish phenotype of the *betaPix*

mutants that exhibit severe cranial hemorrhage and hydrocephalus. Previous studies

have reported *betaPix-Pak2a* signaling and downstream regulation in focal adhesion

essential for vascular stabilization [10,17]. The experiments reported here establish glial

betaPix mediate vascular integrity development *via* two downstream effectors

(Figures 6M and 7). IPA-3 has been shown to decrease the level of phosphorylated

PAK1 in a rat model of subarachnoid hemorrhage [75], and presented anti-angiogenic

activity in zebrafish [76]. Moreover, constitutively active human *Pak1* mRNA

administration induced cranial hemorrhage in zebrafish [77]. While previous researches

focused on *Pak2*, our data suggest that *Pak1* also works with *betaPix* in vascular

integrity development.

Stathmin was first discovered in a study of leukemia, of which *Stathmins are* rapidly

phosphorylated when induced cells undergo terminal differentiation and stopped

proliferation [78]. Subsequent studies have showed downregulated *Stathmin* expressions

after stopping proliferation of leukemia cells by chemical reagents [79]. Similar

expression changes of *Stathmins* are also found in other tumor cell lines such as breast

cancer and ovarian cancer [80,81]. *Stathmins* have also been associated with the

development of the central nervous system. Systemic *Stathmin* knockout mice

develop pan-neural axonopathy upon aging [82]. Knockdown or overexpression of

Stathmin significantly reduced the dendritic growth of Purkinje cells [83]. Moreover,

Stathmin is downregulated with the differentiation of oligodendrocyte progenitor cells

in vitro [84]. In zebrafish, *Stathmin-1* and *Stathmin-2* deficiencies resulted in defected

optic nerves development with ectopic branches and axonal adhesions gaps [85].

Knockdown or overexpression of *Stathmin-4* led to premature differentiation of dorsal

midbrain neural progenitors [86]. Systemic knockout of *Stmn4* showed cell cycle arrest

in zebrafish retinal progenitor cells [87]. These studies are consistent with our results,

suggesting that transcriptionally downregulated *Stathmins* are highly correlated with

the impaired glial development in the phenotypic analysis of this study. Previous

studies have revealed the role of *betaPix*-d isoform in mouse hippocampal neuron

development *via* regulating microtubule stability and PAK-induced *Stathmin*

phosphorylation [32]. It remains identified whether deletion of glial- or neuronal-*betaPix*

leads to brain hemorrhage. Our study found that *stathmin* knockouts impair

development in a variety of cell types in the zebrafish hindbrain. Importantly,

hemorrhage and hydrocephalus of *stathmin* knockouts partially phenocopy *betaPix*

mutants. *In vivo* glial-specific *Stmn1b* overexpression confers partial rescue of the

glial and neuronal but not vascular developmental defects in *bbh* mutants.

Furthermore, ectopic expression of *betaPix* and *Stathmin* alleviate the impaired cell

migration and cell shape maintenance in betaPix siRNA glioblastoma cells. Therefore,

this work suggests that the *betaPix-Pak1-Stathmin* signaling regulates vascular

integrity *via* glial development.

Zfhx4 administration confers partial protection against *betaPix* depletion-induced

hemorrhage, establishing the novel association between the two. However, additional

components of this system remain to be discovered. It has been reported that

cAMP-response element binding protein (CREB) binds to the CRE consensus site at

the ZFHX3 promoter [88]. A variety of protein kinases modulate CREB activation *via*

phosphorylation at Ser133 [89] including protein kinase A (PKA), phosphatidylinositol

3-kinase (PI3K)/Akt, mitogen- and stress-activated kinase 1/2 (MSK1/2), and

Ca2+/calmodulin-dependent protein kinases and *Pak1* [90-92]. Our results demonstrate

that *Pak1* inhibition attenuated the majority of phenotypes induced by *betaPix*

deficiency. Thus, we speculate *Zfhx3/4* are closely associated with pathogenic

mechanisms downstream of *betaPix* signals by modulating expressions of

angiogenetic ligands. Further investigations are required to address this hypothesis.

VEGF is the main pro-angiogenic growth factor produced by various cell types that

controls multiple vascular development steps, such as proliferation, migration,

permeability and survival *via* interacting with VEGF receptors on endothelial cells.

Inactivation of VEGFA in astrocytes leads to BBB disruption in multiple sclerosis

models [93,94]. In early development of avascular neural tube, VEGF derived from

neuroectoderm serve as a central role in inducing tip cells sprouting from the

perineural vascular plexus (PVNP) and formation of vasculature lumens from stem

cells [95,96]. Ablating neuroglia reduces *vegfab* signals and leads to ectopic

intersegmental vessels sprouting and vertebral arteries in zebrafish [68]. VEGF signaling

orchestrates elaborately with other signaling pathways such as delta-like 4

(Dll4)/Notch signaling [97] and activates downstream components including the

Ras/Raf/MEK, PI3K/AKT, and p38/MAPK/HSP27 pathways [98]. Our data suggest

glial *betaPix* as an important factor in vascular integrity development. Together with

previous studies, *betaPix-Pak1-stathmin* signaling regulates glial growth and

differentiation[32] and *betaPix* might regulate *Vegfaa* secretion *via* downstream

transcription factors *Zfhx3/4*. Thus, this work presents novel glial *betaPix* signaling in

stabilizing and maintaining vascular integrity, and loss of this function causes

hemorrhage and subsequent disruption of the BBB (Figure 7). However, the

molecular mechanism of *betaPix* to *Zfhx3/4*, the sufficiency of *Vegfaa* and the

corresponding ligand-receptor interaction between glia and endothelial cells certainly

warrant future investigations.

Basement membrane is a non-cellular component consisting of several extracellular

matrix (ECM) proteins; serve as hub for structural supports, anchoring and

intercellular communication. Components of basement membrane are synthesized in

surrounding cells with diverse expression patterns [99]. For example, mice with deletion

of Collagen IV in either brain microvascular endothelial cells (BMEC)- or pericytes

lead to fully penetrant intracerebral hemorrhage, while COL4A1 deletion in astrocytes

results in mild hemorrhage phenotypes [100]. Differential cell-specific expression

pattern of laminin isoforms contribute differently to BBB formation [101]. As astrocytic

laminins mediate pericyte differentiation and regulate capillary permeability [102],

intercellular interactions with the ECM occur by cell surface integrin receptors.

Integrin β1 on endothelial cells are critical during angiogenesis[103]. On the other hand,

integrin αvβ8 has glial-specific enrichment which mediate focal adhesion to modulate

vascular integrity[17,104,105]. Notably, degradation of ECM and focal adhesions at the

perivascular space is critical to the hemorrhagic phenotypes of *bbh* mutants [17,18]. It

would be of great interest to determine whether glial *betaPix* account for ECM

breakdown in *bbh* mutants. Apart from *Vegfaa*, multiple pro-angiogenic growth

factors and signaling ligands contribute to effective glial–vascular communication in

the perivascular space. Interestingly, ZFHX3 has been reported activating a set of

procollagens, integrin and platelet-derived growth factor receptor β (*Pdgfrb*) genes in

response to retinoic acid stimulus, protecting cerebellar neurons from oxidative stress

[88]. In the future, it will be interesting to examine whether *Zfhx* signaling modulate

adhesion molecules in the pathogenesis of *betaPix* mutants.

Acknowledgements

The authors thank Drs. Bo Zhang, Jiulin Du, Jia Li, Liangyi Chen, Kazu Kikuchi, and

Jian Chen for providing fish lines, plasmid clones, and human glial cell lines; the

members of Dr. Xiong's laboratory for helpful discussions and technical assistance;

and the National Center for Protein Sciences at Peking University, particularly Dr.

Liying Du at the Flow Cytometry Core for technical help on the Beckman Coulter

MoFlo XDP; and Dr. Hua Liang at National Biomedical Imaging Center, Peking

University for assistance with the Luxendo Multi-View Selective-Plane Illumination

Microscopy (Bruker). This work is supported by grants from the National Key R&D

Program of China (2019YFA0801602 and 2023YFA1800600); the National Natural

Science Foundation of China (32230032 and 31730061).

Declaration of Interest

The authors declare no competing financial interests.

Materials and Methods

Fish maintenance

Zebrafish were raised and handled in accordance with the guidelines of the Peking University Animal Care and Use Committee accredited by the AAALAC.

The bbh^{fn40a} mutant was isolated from a large-scale mutagenesis screen of the zebrafish genome at Massachusetts General Hospital, Boston[9]. Tg(*acta2:GFP-Cre*) was purchased from Xinjia Pharmaceutical Technology Co., Ltd (Nanjing, China); Tg(*neurod:EGFP*) were kindly provided by Dr. Bo Zhang (Peking University, China); Tg(*huC:GFP*) were kindly provided by Dr. Liangyi Chen (Peking University, China); Tg(*kdrl:GFP*) [106] and Tg(*kdrl:mCherry*) [107] were described previously.

Cell culture and transfection

U251 and A172 human glioblastoma cell lines, kindly provided by Dr. Jian Chen at Chinese Institute for Brain Research, Beijing, were cultured in high glucose DMEM medium (Hyclone) supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin, at 37°C in 5% CO_2 humidified incubator. The *betaPIX* siRNA[37] and *STMN1* siRNA[108] targeting glioblastoma cell lines as previously reported were purchased from GenePharma (Shanghai, China). The sequences of the siRNAs are listed in **Table S2**. The full-length coding cDNAs of human *betaPIX* or *STMN1* were isolated from U251 cDNA library and subcloned into the pcDNA3.1 vector backbone. Transfection was conducted with Lipofectamine 3000 (Invitrogen) according to manufacturer's instructions at a final siRNA concentration of 30 nM.

mRNA/gRNA synthesis and microinjections

mRNA and gRNA were synthesized as described previously [109]. In brief, linearized

pT3TS-nls-zCas9-nls [110], pXT7-*Cre*, and pXT7-*Zfhx4* plasmid DNA were purified as

templates using TIANquick Mini Purification Kit (TIANGEN, DP203, China). Next,

in vitro transcription reactions were performed using the mMESSAGE mMACHINE

T3 (for pT3TS-nls-zCas9-nls) or T7 (for pXT7-*Cre* and pXT7-*Zfhx4*) kits (Life

Technologies) according to manufacturer's instructions. Templates for gRNA were

generated by complementary annealing and elongation of two oligos. Forward oligo

contained a T7 promoter and gRNA target sequence, and reverse oligo contained the

universal gRNA scaffold [23]. The resulting double-stranded DNA served as the

templates for *in vitro* transcription using HiScribe® T7 High Yield RNA Synthesis

Kit (NEB). The sequences of the gRNAs are listed in **Table S3**. For global *betaPix*

inactivation, *Cre* mRNA with a working concentration of 200 ng/μL were injected

into the *betaPix$^{ct/ct}$* embryos at one-cell stage. For CRISPR-mediated knockouts, 300

ng/μL Cas9 mRNA and 20-50 ng/μL gRNAs were injected into zebrafish embryos at

one-cell stage. For *Zfhx4* rescue experiments, *Zfhx4* mRNA with a working

concentration of 400 ng/μL were injected into the *bbh^{fn40a}* mutants or in combination

with CRISPR-mediated *betaPix* knockout system at one-cell stage. Injected embryos

were scored for mutant phenotypes at 36 or 48 hpf.

Construction of *betaPix* conditional knockout zebrafish

The pZwitch plasmid clone was kindly provided by Dr. Kazu Kikuchi (National

Cerebral and Cardiovascular Center Research Institute, Suita, Japan). GSG spacer was

added to pZwitch+3 between P2A and TagRFP coding sequences by overlapping

primer pairs. Next, a highly efficient gRNA was found from the fifth intron of *betaPix*.

Primer pairs were designed to amplify the left and right homologous arms from the

gRNA site for 1000 bp and 24 bp with following modifications: 1) The 5' end of the

forward primer of the left homologous arm were added with an *Nhe*1 site and a

universal gRNA site which provided an *in vivo* linearization cleavage site; a *Mlu*1 site

to the 5' end of the reverse primer of the left arm. 2) Added *EcoR*1 site to the 5' end of

the forward primer of the right homologous arm; added *Xho1* site and universal

gRNA site to the 5' end of the reverse primer of the right homologous arm. Then we

cloned the inverted homologous arms into polyclonal sites of the modified pZwitch+3,

and purified plasmid DNA with EndoFree Mini Plasmid Kit (Tiangen, DP118). *In*

vivo knock-in system was composed of 300 ng/μL Cas9 mRNA, 50 ng/μL

betaPix-intron5 gRNA, 50 ng/μL Universal gRNA, and 20 ng/μL donor vector.

Microinjected *in vivo* knock-in system into wildtype embryos at one-cell stage. Fish

founders were screened for *α*-crystallin reporter expression, raised to adulthood, and

re-screened for germline transmission and precise knock-in by Sanger sequencing.

Construction of transgenic reporters

The glial reporter plasmid *gfap*-EGFP [26] was kindly provided by Dr Jiulin Du

(Shanghai Institute for Biological Sciences, Shanghai, China). The *gfap* promoter was

used to establish Tg(*gfap:GFP-Cre*) plasmid. The *stmn1b* coding sequences were

amplified from wild-type zebrafish cDNA library, which were used to establish

Tg(*gfap:GFP-stmn1b*) plasmid.

Transgenic reporters were generated by using *Tol2*-based transgenesis [111]. In brief,

100 ng/μL *Tol2* transposase mRNA in combination with 20 ng/μL donor plasmid

DNA were co-injected into the embryos at one-cell stage. Transgenic founders were

screened for specific transgenic GFP expression, raised to adulthood, and re-screened

for germline transmission.

Inhibitor treatment

IPA-3 (Proteintech, CM05727) were dissolved in DMSO to form a 10 mM stock, and

then diluted with E3 medium to 3 μM in 6-well plates. Zebrafish embryos were

treated from 24 to 48 hpf, and then washed with E3 medium for phenotypic analysis.

***o*-dianisidine staining**

o-dianisidine staining was performed as described previously [18]. In brief, embryos

were dechorionated, anesthetized and incubated in fresh staining solution (0.6 mg/mL

o-dianisidine, 0.01 M sodium acetate pH4.5, 0.65% H_2O_2 and 40% Ethanol) for 20

minutes in dark. Washed three times with methanol, and performed benzyl

alcohol/benzyl benzoate (BABB) tissue clearing before imaging with stereo

microscope (Leica, 160F).

Whole-mount *in situ* hybridization (WISH)

Whole-mount *in situ* RNA hybridization was performed as described previously [112]. Antisense probes were synthesized using a digoxigenin RNA labeling kit (Roche, 11277073910). Primer sequences for all WISH probes used in this paper are provided in **Table S4**.

Quantitative real-time RT-PCR

Total RNA was isolated from embryos using Trizol (invitrogen) and cDNA was generated using HiScript® III RT SuperMix (Vazyme) according to the manufacturer's instruction. Quantitative real-time PCR was performed using Lightcycler (Roche) and ChamQ SYBR qPCR Master Mix (Vazyme). Primer sequences are listed in Supplementary Materials **Table S5**. Gene expression level was normalized against GAPDH level.

Light-sheet fluorescence microscopy imaging

Light-sheet microscopy imaging was performed as described previously [113]. In brief, transgenic zebrafish embryos were collected and maintained at 28.5°C. At the required developmental stages, the embryos were carefully dechorionated, paralyzed with tricaine, and transferred into 1% ultrapure low melting point agarose (16520-050; Invitrogen). The embryo was then drawn into a glass tube in top-down position using a 1 mL syringe with an 18G blunt needle. After agarose coagulation, a wire was inserted from bottom to push the agarose with embryo upwards, removed excess agar

until the head area is exposed from the top of the glass tube. Finally, the glass tube

was fixed in a sample holder for subsequent imaging.

Imaging was carried out with Luxendo Multi-View Selective-Plane Illumination

Microscopy (Bruker). Optical calibrations were adjusted according to the instruction

manual.

The optical plates are emitted from two Nikon CFI Plan Fluor 10x W 0.3 NA

immersion objectives in opposite directions. The detection is completed by two

Olympus 20x 1.0 NA immersion lenses. The parameters are set as follows: the green

fluorescence channel uses a laser of 488 nm coupled with BP497-554 filter, the red

fluorescent channel uses laser 561 nm coupled with BP580-627 filter, with an

exposure time of 100 ms, a delay of 11 ms, a line mode of 50 px, and z-Stack interval

of 3 μm. The raw data were stored in H5 file format, processed into TIFF format and

merged Multi-View dataset using MATLAB software. After pre-processing,

multi-fluorescence channels were merged in ImageJ, and 3D visualization and

measurement were performed by Imaris.

Single-cell RNA sequencing

For knockout samples, 300 ng/μL Cas9 mRNA and a mixture of number 1 to 4

betaPix gRNAs with 50 ng/μL each were co-injected into wildtype embryos at

one-cell stage. Injection with equivalent concentration of Cas9 mRNA with PBS

served as siblings. After microinjection, embryos were collected and maintained at

28.5°C. At 24 or 48 hpf, zebrafish were dechorionated, paralyzed and transferred onto

agarose plate. Heads were harvested with dissecting scissors in cold sterile 1X PBS

with pooling 200 heads for each group. Heads were then dissociated in 900 μL

Accutase cell detachment solution (Sigma-Aldrich, A6964) at 28.5°C for 3 hours and

re-suspended by pipetting every 30 min. Once digestion was complete, 100 μL FBS

was added to cell suspension and centrifuged at 4℃, 500 g for 3 min. The cells were

gently re-suspended in cold 500 μL 2.5% FBS in 1X PBS and filtered through a 40

μm strainer. PI and Hoechst33342 were stained for distinguishing living cells.

The single, living cells were sorted by Aria SORP (BD biosciences) into 1.5 ml tubes.

The cell counts and vitality were verified by AOPI staining coupled with an

Automated Cell Counter (Countstar BioTech). Around 10,000-12,000 cells were

loaded for each group. Single cells were barcoded with Chromium Next GEM Single

Cell 3 'Reagent Kits V3.1 kit (10x Genomics, 1000269) in 10× Chromium Controller

(10× Genomics). After qualified by peak shapes, fragment size and tailing with

Fragment Analyzer System kit (Agilent Technologies, DNF-915), single-cell

transcriptome libraries were sequenced via Illumina High Throughput Sequencing

PE150 (Novogene, Beijing). The sequencing data were analyzed using the

CellRanger-6.1.1 (10x Genomics) and mapped to reference genome

GRCz11-GRCz11.103. The output numbers of reads in four sample groups are

494,735,089 for ctrl-1d, 615,837,087 for ko-1d, 554,470,209 for ctrl-2d and

583,809,243 for ko-2d.

Low-quality cells were excluded from subsequent analyses under the following

conditions: when the number of expressed genes was less than 500, when there were

abnormally high counts of UMIs or genes (outliers of a normal distribution), or when

the mitochondrial content exceeded 9%. A total of 38,670 cells were qualified for

subsequent analyses. Unsupervised clustering was performed using Seurat (version:

4.0.2) with a resolution of 2.5, resulted in 71 cell populations and further annotated

into 24 zebrafish major cranial cell types. Differential expression analysis in Seurat v4

was used to identify cluster/cell type markers by Wilcoxon rank sum test.

Scratch assay

U251 cells were planted into culture plate and performed transfection at the desired

density. At 24 hours post-transfection, we vertically scratched monolayer cells by

using a 200 μL pipette tip. Washed the cells three times with PBS, then replaced with

serum-free culture medium for further culture. Stereo fluorescence microscopy (Leica,

160F) was used to document the scratch size at 0 hours, 18 and 24 hours. Images were

processed by ImageJ.

Immunostaining

For assessing tubulin expression, U251 cells were planted into chamber slides

(Saining, 1093000) and transfected at the desired density. At 24 hours

post-transfection, washed the cells once on ice with pre-cooled PBS, then fixed with

pre-cooled methanol at -20℃ for 20 minutes. The fixed cells were permeabilized with

pre-cooled acetone at -20℃ for 1 minute. Removed acetone, incubated 0.5%

BSA/PBS blocking solution at room temperature for 15 minutes. Cells were then

incubated with 1:200 anti-α-tubulin mouse monoclonal antibody (EASYBIO, BE0031)

at 37℃ for 45 minutes. After rinsing once with PBS, cells were incubated with 1:400

Alexa Fluor™ 488 goat anti-mouse IgG (H+L) secondary antibody (Invitrogen,

A11029) at 37℃ for another 45 minutes. Washed three times with PBS, and sealed

with Mounting Medium with DAPI (ZSGB Bio, ZLI-9557). Images were acquired by

upright fluorescence microscopy (Leica, DM5000B) at 40× magnification, and

processed by ImageJ.

Statistical analysis

Statistical analysis was performed using Graphpad Prism 6. The statistical

significance of differences between the two groups was determined by the

independent unpaired Student's t-test. Among three or more groups, one-way

ANOVA analysis coupled with Dunnett's test were used. All data are presented as the

mean ± SEM. P value <0.05 indicates significant, with individual P values mentioned

in the figure/figure legends.

 Figure Legends

 Figure 1. Generation of *betaPix* conditional trap (*betaPix^{ct}*) allele by an

 homologous recombination (HDR)-mediated knock-in method.

(A) Schematic diagram illustrates the HDR-mediated *Zwitch* strategy for

generating zebrafish knock-in allele at the *betaPix* locus.

(B) Genomic PCR analysis of the F$_1$ embryos confirming the right *Zwitch*

insertions.

(C) Sanger sequencing confirming the junction of *betaPix^{ct}* (after HDR-mediated

insertion) or *betaPix^{m}* (after Cre-mediated inversion) that are highlighted by

RED lines in (A).

(D) Representative stereomicroscopy images of erythrocytes stained with

o-dianisidine in *betaPix$^{ct/ct}$* and *betaPix$^{m/m}$* embryos at 48 hpf. Brain

hemorrhages, indicated with arrow, in *Cre* mRNA-injected embryos

(*betaPix$^{m/m}$*). Lateral views, anterior to the left.

(E) Quantification of hemorrhagic parameters in (D). Left panel showing

hemorrhage percentages, with independent experiments as dots. Right panel

showing hemorrhage areas, with each dot representing one embryo, #

represents the numbers of embryos scored for each analysis, and three or more

individual experiments conducted. Data are presented in mean ± SEM;

unpaired Student's t test with individual *P* values mentioned in the figure.

(F) qRT-PCR analysis showing the expression of *betaPix, RFP,* and *alphaPix* in

betaPix$^{ct/ct}$, *betaPix$^{ct/m}$,* and *betaPix$^{m/m}$* embryos at 48 hpf. Each dot represents

one embryo. Data are presented in mean ± SEM; one-way ANOVA analysis

with Dunnett's test, individual P values mentioned in the figure. cryaa, α

A-crystallin; PA, polyadenylation signal; SA, splice acceptor; T2A,

T2A self-cleaving peptide. Individual scale bars indicated in the figure.

Figure 2. *betaPix^{m/m}* mutant have brain hemorrhages, central artery defects and

abnormal glial structure that was partially rescued by Pak1 inhibitor IPA-

3 treatment.

(A) Left panel showing the maximum intensity projection of the glial structures in

the hindbrain of *betaPix^{ct/ct}* and *betaPix^{m/m}* embryos at 48 hpf. Lateral view,

anterior to left. Middle panel showing representative optical sections and right

panel showing the higher magnifications of boxed area, presenting atypical

glial structures with disoriented arrangements (yellow arrows) in *betaPix^{m/m}*

embryos.

(B) Quantification of glial parameters in (A). Left panel showing the average glia

length, and right panel showing glia length index normalized to individual

head length, which each dot represents one embryo. Data are presented in

mean ± SEM; unpaired Student's t test with individual P values mentioned in

the figure.

(C) Whole-mount RNA *in situ* hybridization revealed *nestin* and *pax2a* expression

pattern in *betaPix^{ct/ct}* and *betaPix^{m/m}* embryos at 48 hpf. Dorsal view, anterior

to the left.

(D) Optical sections of glial structure (green) and blood vessels (magenta) in the

heads of siblings and CRISPR-mediated *betaPix* F_0 knockout embryos.

Arteries in the hindbrain of *betaPix* KO mutants had developmental defects

(white arrowheads), showing shorter distance between basilar artery and glial

cell bodies.

(E) 3D reconstruction of the *sox2*-positive precursors (green) and vasculatures

(magenta) in the heads of siblings and CRISPR-mediated *betaPix* F_0 knockout

embryos at 48 hpf. Box areas are shown in higher magnifications at the middle

panels, with optical sections shown in the right panels. Arrows indicate CtA

with enlarged perivascular space.

(F) Representative stereomicroscopy images of *o*-dianisidine staining of

betaPix$^{ct/ct}$ and *betaPix$^{m/m}$* embryos at 48 hpf that were treated with DMSO or

PAK inhibitor IPA3. Brain hemorrhages indicated with arrows.

(G) Quantification of brain hemorrhagic parameters in (F). Left panel showing

hemorrhage percentages, with independent experiments as dots. Right panel

showing hemorrhage areas, with each dot representing one embryo, #

represents the numbers of embryos scored for each analysis, and three or more

individual experiments conducted. Data are presented in mean ± SEM;

one-way ANOVA analysis with Dunnett's test, individual *P* values mentioned

in the figure.

(H) Left panels showing 3D reconstruction of the glial structure (green) and

vasculature (magenta) in the heads of *betaPix$^{ct/ct}$* and *betaPix$^{m/m}$* embryos at 48

hpf treated with DMSO or IPA3. Lateral view, anterior to left. Box areas are

shown in higher magnifications at the middle panels. Defects in hindbrain

central arteries indicated in white arrows, while defects in radial glia indicated

in yellow arrows. Right panels showing schematic diagrams. Glia (grey) and

CtAs (pink) develop normally in DMSO or IPA3-treated *betaPix$^{ct/ct}$* embryos,

with fine radial glial processes and characteristic arch vasculture. Yet in

betaPix$^{m/m}$ embryos, abnormal development of the glia and central artery

presented. In IPA-3-treated *betaPix$^{m/m}$* embryos, central arterial defects were

partially rescued with relatively complete arch architecture, and glial processes

defects significantly rescued.

(I) Quantification of CtA parameters in (H). Left panel showing the average CtA

length, and right panel showing the CtA length index normalized to individual

head length, with each dot representing one embryo. Data are presented in

mean ± SEM; one-way ANOVA analysis with Dunnett's test, individual *P*

values mentioned in the figure.

(J) Quantification of glia parameters in (H). Left panel showing the average glia

length, and right panel showing glia length index normalized to individual

head length, which each dot represents one embryo. Individual scale bars

indicated in the figure. Data are presented in mean ± SEM; one-way ANOVA

analysis with Dunnett's test, individual *P* values mentioned in the figure.

Individual scale bars indicated in the figure. BA, basilar artery.

Figure 3. Glial-specific *betaPix* knockouts recapture global *betaPix* mutant

phenotypes.

(A) Representative stereomicroscopy images of erythrocytes stained with

   o-dianisidine in *betaPix^{ct/ct}* siblings and *gfap:GFP-Cre*; *betaPix^{ct/ct}* mutant

   embryos treated with DMSO or IPA3 at 48 hpf. Brain hemorrhages indicated

   with arrows in glial-specific *betaPix* knockouts. Lateral view with anterior to

   the left.

(B) Left panels showing 3D reconstruction of the vasculature (magenta) in the

   heads at 48 hpf, lateral view with anterior to the left. Box areas are shown in

   higher magnifications of brain vasculatures at the right panels. CtA defects

   indicated in yellow arrows in *gfap:GFP-Cre*; *betaPix^{ct/ct}* mutant embryos.

(C) Quantification of brain hemorrhages in (A) and CtA parameters in (B). Left

   panel showing hemorrhage percentages, with independent experiments as dots.

   Middle panel showing hemorrhage areas with each dot representing one

   embryo. # represents the numbers of embryos scored for each analysis, three

   or more individual experiments conducted. Right panel showing CtA length

   index normalized to individual head length, with each dot representing one

   embryo. Data are presented in mean ± SEM; one-way ANOVA analysis with

   Dunnett's test, individual *P* values mentioned in the figure. Individual scale

   bars indicated in the figure.

Figure 4. Single cell transcriptome reveals that a subcluster of glial progenitor

and stathmin family members are associated with *betaPix* mutation.

(A) Experimental strategy for single cell RNA sequencing of embryonic heads

from wild-type siblings and *betaPix* CRISPR mutants at 1 dpf and 2 dpf.

(B) UMAP visualization and clustering of cells labeled by cell type. Four samples

were aggregated and analyzed together.

(C) Proportions of 24 cell clusters were differentially distributed among four

sample groups. ctrl_1d, PBS-injected siblings at 1 dpf; ko_1d, *betaPix*

CRISPR mutants at 1 dpf; ctrl_2d, PBS-injected siblings at 2 dpf; ko_2d,

betaPix CRISPR mutants at 2 dpf.

(D) Enriched GO terms for differentially expressed genes for progenitor

sub-cluster comparing ko_2d to ctrl_2d groups.

(E) Violin plots showing the expression of the stathmin family genes by all cells

among four sample groups (left panel) or by progenitor sub-cluster among

four sample groups (right panel).

(F) qRT-PCR analysis showing expression of *betaPix, stmn1a, stmn1b, stmn4l*

and *ppfia3* in glia of FACS-sorted *betaPix^{ct/ct}* siblings and *betaPix^{m/m}* mutants

at 48 hpf. Each dot represents cells sorted from one embryo. Data are

presented in mean ± SEM. ****P* <0.005; *****P*<0.001; unpaired

Student's t test.

(G) qRT-PCR analysis showing expression of *betaPix, stmn1a, stmn1b, stmn4l*

and *ppfia3* in glia of FACS-sorted *betaPix^{ct/ct}* siblings and *gfap:GFP-Cre;*

betaPix$^{ct/ct}$ mutants at 48 hpf. Each dot represents cells sorted from one embryo.

Data are presented in mean ± SEM. ***P <0.005; ****P<0.001; unpaired

Student's t test.

(H) Whole-mount RNA *in situ* hybridization revealing down-regulation of *stmn1a*,

stmn1b and *stmn4l* in *betaPix$^{ct/ct}$* siblings and *gfap:GFP-Cre*; *betaPix$^{ct/ct}$*

mutant embryos at 48 hpf. Individual scale bars indicated in the figure.

Figure 5. *Stathmin* acts downstream of *betaPix* in glial migration *via* regulating

tubulin polymerization.

(A) Whole-mount RNA *in situ* hybridization showing that *stmn1a, stmn1b and*

stmn4l expression in *gfap:GFP-Cre*; *betaPix$^{ct/ct}$* embryos were partially

rescued by Pak1 inhibitor IPA3 treatment at 48 hpf. Dorsal views, anterior to

the left.

(B) qRT-PCR analysis showing that *stmn1a* and *stmn4l* expression were rescued

in *gfap:GFP-Cre*; *betaPix$^{ct/ct}$* mutants by IPA3 treatment at 48 hpf. Each dot

represents one embryo. Data are presented in mean ± SEM; one-way ANOVA

analysis with Dunnett's test, individual P values mentioned in the figure.

(C) Representative stereomicroscopy images of erythrocytes stained with

o-dianisidine in siblings and *stmn1a/1b/4l* CRISPR mutants at 48 hpf. Brain

hemorrhages, indicated with arrows, appeared in *stmn1a/1b/4l* mutants.

Lateral views with anterior to the left.

(D) Quantification of hemorrhagic parameters in (C). Left panel showing

hemorrhage percentages, with independent experiment as dot. Right panel

showing hemorrhage areas with each dot representing one embryo. #

represents the numbers of embryos scored for each analysis, three or more

individual experiments conducted. Data are presented in mean ± SEM;

unpaired Student's t test with individual P values mentioned in the figure.

(E) 3D reconstruction of glial structure (green) and vasculature (magenta) in the

heads at 48 hpf. Lateral view with anterior to the left. CtA defects (white

arrows) indicated in *stmn1a/1b/4l* mutants.

(F) Quantification of CtA and glia parameters in (E). Length index normalized to

individual head length, with each dot representing one embryo. Data are

presented in mean ± SEM; unpaired Student's t test with individual P values

mentioned in the figure.

(G) Representative stereomicroscopy images of erythrocytes stained with

o-dianisidine in *bbh*[fn40a] and *bbh*[fn40a]; Tg(*gfap:GFP-stmn1b*) embryos at 48 hpf.

Brain hemorrhages, indicated with arrows, decreased in *bbh*[fn40a] mutants with

glia-specific overexpression of *stmn1b*, compared with *bbh*[fn40a] mutant siblings.

Lateral views with anterior to the left.

(H) Quantification of hemorrhagic parameters of (G). Left panel showing

hemorrhage percentages, with independent experiment as dot. Right panel

showing hemorrhage areas with each dot representing one embryo. #

represents the numbers of embryos scored for each analysis, three or more

924 individual experiments conducted. Data are presented in mean ± SEM;

925 unpaired Student's t test with individual P values mentioned in the figure.

926 (I) 3D reconstruction of the *gfap:GFP-stmn1b* overexpression (green) and

927 vasculature (magenta) in the heads of bbh^{fn40a} siblings and bbh^{fn40a};

928 Tg(*gfap:GFP-stmn1b*) mutants at 48 hpf. Lateral view with anterior to

929 left. White arrows indicate CtA defects.

930 (J) Quantification of CtA and glia parameters in (I). Length index normalized to

931 individual head length, with each dot representing one embryo. Data are

932 presented in mean ± SEM; unpaired Student's t test with individual P values

933 mentioned in the figure.

934 (K) Representative stereomicroscopy images of U251 cells at 0 and 24 hours after

935 wounding. U251 cells were transfected with negative control siRNA or

936 *betaPIX* siRNA separately, in combination with pcDNA3.1 vector, *betaPIX*

937 overexpression plasmid, and *STMN1* overexpression plasmid. The wound

938 edges are highlighted by dashed lines, with arrow lines indicating the wound

939 width.

940 (L) Quantification of wound closure in (K), showing $^{*}P{<}0.05$, $^{***}P{<}0.005$

941 compared to negative control siRNA with empty vector transfection. $^{\#\#}P{<}0.01$,

942 $^{\#\#\#}P{<}0.005$ compared to *betaPix* knockdown with empty vector transfection.

943 Data are presented in mean ± SEM; one-way ANOVA analysis with Dunnett's

944 test.

(M) Representative immunofluorescence image of alpha-tubulin and DAPI

signals in U251 cells. U251 cells were transfected with negative control

siRNA or *betaPIX* siRNA separately, in combination with pcDNA3.1 vector,

betaPIX overexpression plasmid, and *STMN1* overexpression plasmid. Box

areas are shown in higher magnifications. Arrows indicate protrusions at the

cell periphery.

(N) Quantification of cell percentages with protrusions in (M). ***$P<0.005$

compared to negative control siRNA with empty vector transfection. #$P<0.05$,

##$P<0.01$ compared to *betaPix* knockdown with empty vector transfection.

Data are presented in mean ± SEM; one-way ANOVA analysis with Dunnett's

test. Individual scale bars indicated in the figure.

Figure 6. *Zfhx3/4* acts downstream of *betaPix* to regulate vascular integrity

development.

(A) Whole-mount RNA *in situ* hybridization revealing that *Vegfaa* decreased in

bbh^{fn40a} mutants compared with siblings at 36 hpf. Lateral views with anterior

to the left. Arrows indicate *vegfaa* expression in the CtAs that shown depletion

in *bbh*^{fn40a} mutants.

(B) qRT-PCR analysis revealing that *ZFHX3*, *ZFHX4 and* VEGFA decreased in

U251 cells transfected with *betaPIX* siRNA, which were rescued by *betaPIX*

overexpression. Data are presented in mean ± SEM; one-way ANOVA

analysis with Dunnett's test, individual *P* values mentioned in the figure.

(C) Whole-mount RNA *in situ* hybridization showing that *vegfaa* decreased in

CRISPR-mediated *zfhx3/4* F_0 knockout embryos at 36 hpf. Lateral views with

anterior to the left. Arrows indicate *vegfaa* expression in the CtAs that shown

reduction in *zfhx3/4* knockouts.

(D) Representative stereomicroscopy images of erythrocytes stained with

o-dianisidine in siblings and CRISPR-mediated *zfhx3/4* F_0 knockout embryos

at 48 hpf. Arrows indicated brain hemorrhages in the knockout brains. Lateral

views with anterior to the left.

(E) Quantification of hemorrhagic parameters in (D). Left panel showing

hemorrhage percentages, with independent experiment as dot. Right panel

showing hemorrhage areas with each dot representing one embryo. #

represents the numbers of embryos scored for each analysis, three or more

individual experiments conducted. Data are presented in mean ± SEM;

unpaired Student's t test with individual *P* values mentioned in the figure.

(F) Left panel showing 3D reconstruction of the glial structure (green) and

vasculature (magenta) in the hindbrain of siblings and CRISPR-mediated

zfhx3/4 F_0 knockouts at 48 hpf. Lateral view with anterior to the left. White

arrows indicate CtA defects. Right panels showing schematic diagrams. Glia

(grey) developed normally in both treatment, while defects in central artery

(pink) presented in *zfhx3/4* knockout embryos.

(G) Quantification of CtA and glia length parameters of (F). Length index

normalized to individual head length, with each dot representing one embryo.

Data are presented in mean ± SEM; unpaired Student's t test with individual *P*

values mentioned in the figure.

(H) Representative stereomicroscopy images of erythrocytes stained with

o-dianisidine in CRISPR-mediated *betaPix* F0 knockout embryos with or

without *zfhx4* mRNA injection at 48 hpf. Arrows indicate brain hemorrhages.

Lateral views with anterior to the left.

(I) Quantification of hemorrhagic parameters in (H). Left panel showing

hemorrhage percentages, with independent experiment as dot. Right panel

showing hemorrhage areas with each dot representing one embryo. #

represents the numbers of embryos scored for each analysis, three or more

individual experiments conducted. Data are presented in mean ± SEM;

unpaired Student's t test with individual *P* values mentioned in the figure.

(J) Left panel showing 3D reconstruction of the glial structure (green) and

vasculature (magenta) in CRISPR-mediated *betaPix* F_0 knockout embryos

with or without *zfhx4* mRNA injection at 48 hpf. White arrows indicate CtAs

and yellow arrows indicate glia. Right panels showing schematic diagrams.

Glia (grey) and CtA (pink) developmental defects rescued in *zfhx4* treatment.

(K) Quantification of CtA and glia length parameters in (J). Length index

normalized to individual head length, with each dot representing one embryo.

Data are presented in mean ± SEM; unpaired Student's t test with individual *P*

values mentioned in the figure.

(L) Whole-mount RNA *in situ* hybridization revealed that *Zfhx4* and *Vegfaa*

decreased in CRISPR-mediated *betaPix* F_0 knockout embryos at 36 hpf, which

were rescued by *Zfhx4* mRNA injection. Lateral views with anterior to the left.

Arrows indicate hindbrain regions.

(M) Dot plots of several angiogenesis-associated genes expression in endothelial

cell cluster. Dot size indicates the percentage of cells with gene expression,

and dot color represents the average gene expression level. Individual scale

bars indicated in the figure.

1018

Figure 7. Schematic diagram on the function of glial *betaPix* in zebrafish

vascular integrity development of the hindbrain. *betaPix* is enriched in glia and

regulates the PAK1-Stathmin axis on microtublin stabilization, thus fine-tuning glial

cell migration and their interactions with vanscular endothelial cells; and in paralelle,

betaPix may also regulate Zfhx3/4-Vegfaa signaling in glia, which then modulates

angiogenesis during cerebral vessel development and maturation. Deletion of *betaPix*

affects glial cell migration and interaction with cerebral endothelial cells. MT,

Microtubule.

1027

Figure S1. Generation of *betaPix* conditional trap (*betaPix^ct*) allele by an

HDR-mediated knock-in method.

(A) Sanger sequencing results of the original pZwitch (+3) and the modified

vector with glycine-serine-glycine spacer (GSG) in front of the 2A peptide.

(B) Sanger sequencing results of the genomic PCR products at both junction sites

from the *betaPix^{ct}* F$_1$ offspring showing the correct knock-in in the betaPix

locus.

(C) The screening process for *betaPix^{ct}* knock-in zebrafish showing about 1%

efficiency with correct homologous recombinations.

(D) Representative stereomicroscopy images of *o*-dianisidine staining in *bbh^{fn40a}*

embryos at 48 hpf. Lateral view, anterior to the left in left panel; Dorsal view,

anterior to the top in right panel.

(E) Quantification of hemorrhagic parameters in (D). Left panel showing

hemorrhage percentages, with independent experiment as dot. Right panel

showing hemorrhage areas with each dot representing one embryo. #

represents the numbers of embryos scored for each analysis, three or more

individual experiments conducted. Data are presented in mean ± SEM;

unpaired Student's t test with individual *P* values mentioned in the figure.

(F) 3D reconstruction of the *betaPix* expression (magenta) in *betaPix^{ct/+}* brains of

embryos microinjected with PBS or *Cre* mRNA at 24 hpf, 48 hpf, 72 hpf and

144 hpf. Cre-induced *RFP* expression indicates successful inversion of gene

trap system (arrows). Dorsal view, anterior to the top.

(G) 3D reconstruction of the *neurod1* transgenic expression (green) and *betaPix*

expression (magenta) in *betaPix^{ct/+}* brains of embryos microinjected with PBS

or *Cre* mRNA at 24 hpf, 48 hpf, 72 hpf and 144 hpf. Cre-induced *RFP*

expression indicates successful inversion of gene trap system (arrows),

marking strong *betaPix* expression at the midbrain hindbrain boundary (MHB),

and milder expression at hindbrain at early stages. At 144 hpf, *betaPix*

expresses ubiquitously in the brain, outlining the cerebellum (arrowheads).

Dorsal view, anterior to the top. Individual scale bars indicated in the figure.

1058

Figure S2. *betaPix^{m/m}* mutant had brain hemorrhages, central arteries defects,

and abnormal glial structure.

(A) 3D reconstruction of the vasculature (green) in the heads with lateral view,

anterior to left. Endogenous *betaPix* expression (magenta) only shown in Cre

mRNA-injected mutant embryos. Box areas are shown in higher

magnifications of vasculatures at the right panels. Central arteries (CtA)

defects (white arrows) were evident in *betaPix^{m/m}* embryos at 48 hpf. Arrow

lines indicating the CtA length.

(B) Quantification of CtA parameters in (A). Left panel showing the average CtA

length, and right panel showing the CtA length index normalized to individual

head length, with each dot representing one embryo. Data are presented in

mean ± SEM; unpaired Student's t test with individual P values mentioned in

the figure.

(C) 3D reconstruction of the glia structure. Box areas showing higher

magnifications of glia at the right panels. D, dorsal; V, ventral. Arrow lines

indicating the glia length. Individual scale bars indicated in the figure.

1075

Figure S3. CRISPR-mediated *betaPix* F_0 knockouts had similar phenotypes as

***betaPix$^{m/m}$* mutants.**

(A) Schematic diagram illustrates guide RNA sites for CRISPR-mediated *betaPix*

F_0 knockout. E, exon.

(B) qRT-PCR revealing that *betaPix* decreased in CRISPR-mediated *betaPix* F_0

knockout embryos compared with wild-type siblings at 48 hpf. Each dot

represents one embryo. Data are presented in mean ± SEM; unpaired

Student's t test with individual *P* values mentioned in the figure.

(C) Maximal intensity projection of the glial structure of the hindbrains in siblings

and CRISPR-mediated *betaPix* F_0 knockouts at 48 hpf, lateral view with

anterior to left.

(D) Whole-mount RNA *in situ* hybridization revealing up-regulated *nestin* and

down-regulated *pax2a* expression in CRISPR-mediated *betaPix* F_0 knockout

embryos at 48 hpf. Dorsal view with the anterior to the left.

(E) Quantification of hemorrhagic parameters of siblings and CRISPR-mediated

betaPix F_0 knockout embryos at 48 hpf. Left panel showing hemorrhage

percentages, with independent experiment as dot. Right panel showing

hemorrhage areas with each dot representing one embryo. # represents the

numbers of embryos scored for each analysis, three or more individual

experiments conducted. Data are presented in mean ± SEM; unpaired

Student's t test with individual *P* values mentioned in the figure.

(F) Quantification of CtA parameters of siblings and CRISPR-mediated *betaPix*

F_0 knockout embryos at 48 hpf. Left panel showing average CtA length, and

right panel showing CtA length index normalized to individual head length,

with each dot representing one embryo. Data are presented in mean ± SEM;

unpaired Student's t test with individual *P* values mentioned in the figure.

(G) Quantification of glial parameters of siblings and CRISPR-mediated *betaPix*

F_0 knockout embryos at 48 hpf. Left panel showing average glia length, and

right panel showing glia length index normalized to individual head length,

with each dot representing one embryo. Data are presented in mean ± SEM;

unpaired Student's t test with individual *P* values mentioned in the

figure. Individual scale bars indicated in the figure.

1108

Figure S4. Generating glial-specific *betaPix* knockout zebrafish.

(A) Schematic diagram illustrates establishment of glial-specific *betaPix* knockout

zebrafish.

(B) 3D reconstruction of the heads at 48 hpf with lateral view, anterior to

left. Glial-specific *Cre* expression (green) was only shown in *gfap:GFP-Cre*;

betaPix^{ct/ct} mutant embryos, which overlaps with betaPix:RFP expression

(magenta).

(C) Genomic PCR analysis of the *betaPix^{ct}*- or *betaPix^{m}*-unique sequences in

betaPix^{ct/ct} siblings, *gfap:GFP-Cre*; *betaPix^{ct/+}* heterozygous mutants, and

gfap:GFP-Cre; *betaPix^{ct/ct}* homozygous mutants.

(D) qRT-PCR analysis revealing *Cre* expression in *gfap:GFP*-positive glial

population and *gfap:GFP*-negative non-glial population that were sorted from

gfap:GFP-Cre; *betaPix$^{ct/ct}$* embryos. Each dot represents one embryo. Data are

presented in mean ± SEM; unpaired Student's t test with individual *P* values

mentioned in the figure.

(E) qRT-PCR analysis showing *betaPix, betaPix-RFP* and *alphaPix* expression in

betaPix$^{ct/ct}$ control siblings and *gfap:GFP-Cre*; *betaPix$^{ct/ct}$* mutant embryos at

48 hpf. Each dot represents one embryo. Data are presented in mean ± SEM;

unpaired Student's t test with individual *P* values mentioned in the figure.

(F) Whole-mount RNA *in situ* hybridization revealing that *alphaPix* was

comparable but *betaPix* decreased in *gfap:GFP-Cre*; *betaPix$^{ct/ct}$* mutants

compared with *betaPix$^{ct/ct}$* siblings at 48 hpf. Lateral view with anterior to the

left. Individual scale bars indicated in the figure.

1132

Figure S5. Neither endothelial-, neuronal- nor mural-specific deletion of *betaPix*

caused brain hemorrhages and abnormal CtA phenotypes.

(A) Representative stereomicroscopy images of *o*-dianisidine staining in

betaPix$^{ct/ct}$, *kdrl:Cre*; *betaPix$^{ct/ct}$*, *huC:GFP-Cre*; *betaPix$^{ct/ct}$*, and

acta2:GFP-Cre; *betaPix$^{ct/ct}$* embryos at 48 hpf.

(B) Quantification of hemorrhage parameters in (A). Up panel showing

hemorrhage percentages, with independent experiment as dot. Down panel

showing hemorrhage areas with each dot representing one embryo. #

represents the numbers of embryos scored for each analysis, three or more

individual experiments conducted. Data are presented in mean ± SEM;

unpaired Student's t test with individual *P* values mentioned in the figure.

(C) 3D reconstruction of the vasculature of *betaPix*ct/ct, *kdrl:Cre*; *betaPix*ct/ct,

huC:GFP-Cre; *betaPix*ct/ct, and *acta2:GFP-Cre*; *betaPix*ct/ct embryos at 48 hpf.

Lateral view with anterior to left. Boxed areas of the hindbrains are shown in

higher magnifications at the right panels.

(D) Quantification of CtA parameters in (C). Up panel showing average CtA

length, and down panel showing CtA length index normalized to individual

head length, with each dot representing one embryo. Data are presented in

mean ± SEM; unpaired Student's t test with individual *P* values mentioned in

the figure. Individual scale bars indicated in the figure.

1153

Figure S6. Glial progenitor sub-cluster of *betaPix* knockouts has

down-regulation of *stathmin* family members and up-regulation of *pak1* gene.

(A) Dot plot of marker genes expression in each cell type. Dot size indicates the

percentage of cells expressed, and dot color represents the average expression

level. Box area and arrow highlighting *gfap* expressions. Clusters with high

gfap levels including neuronal and glial progenitors, hindbrain, ventral

diencephalon, ventral midbrain and floor plate.

(B) UMAP visualization and clustering of cells labeled by cell type across four

sample groups.

(C) UMAP feature plots displaying relative expression levels of selected

transcripts among four sample groups. Cells are colored by expression level.

1165

Figure S7. CRISPR-mediated *stmn1a/1b/4l* F$_0$ knockouts have similar but milder

phenotypes as *betaPix* knockouts.

(A) qRT-PCR analysis showing that IPA3 treatment rescued *stmn1a* and *stmn4l*

expression in CRISPR-mediated *betaPix* F$_0$ knockouts at 48 hpf. Each dot

represents one embryo. Data are presented in mean ± SEM; one-way ANOVA

analysis with Dunnett's test, individual P values mentioned in the figure.

(B) Schematic diagram illustrates 4-guide RNAs mediated F$_0$ knockout strategy in

the loci of *stmn1a*, *stmn1b* and *stmn4l*.

(C) Whole-mount RNA *in situ* hybridization confirming that *stmn1a, stmn1b* and

stmn4l decreased in CRISPR-mediated *stmn1a/1b/4l* F$_0$ knockouts at 48 hpf.

Dorsal views with anterior to the left.

(D) qRT-PCR analysis confirming that *stmn1a, stmn1b* and *stmn4l* decreased in

mutant embryos of CRISPR-mediated *stmn1a/1b/4l* F$_0$ knockouts at 48 hpf.

Each dot represents one embryo. Data are presented in mean ± SEM; unpaired

Student's t test with individual *P* values mentioned in the figure.

(E) qRT-PCR analysis confirming that *stmn1a, stmn1b* and *stmn4l* decreased in

the FACS-sorted glia of CRISPR-mediated *stmn1a/1b/4l* F$_0$ knockouts at 48

hpf. Each dot represents sorted cells from one embryo. Data are presented in

mean ± SEM; unpaired Student's t test with individual P values mentioned in

the figure.

(F) Optical sections of glial structure in the hindbrains of siblings and

CRISPR-mediated *stmn1a/1b/4l* F_0 knockouts at 48 hpf. Lateral view with

anterior to the left. Abnormal glial structures with disoriented arrangements

(yellow arrows) in *stmn1a/1b/4l* mutant embryos.

(G) Quantification of average CtA and glial length in Figure 5 (E). Each dot

represents one embryo. Data are presented in mean ± SEM; unpaired

Student's t test with individual P values mentioned in the figure.

(H) Whole-mount RNA *in situ* hybridization revealed that *nestin* increased while

pax2a decreased in CRISPR-mediated *stmn1a/1b/4l* F_0 knockout embryos at

48 hpf. Dorsal views with anterior to the left.

(I) 3D reconstruction and whole-mount RNA *in situ* hybridization showing that

gfap:GFP-stmn1b transgenic overexpression rescued glial defects in *bbh*^{fn40a};

Tg(*gfap:GFP-stmn1b*) embryos at 48 hpf. Lateral views with anterior to the

left.

(J) Whole-mount RNA *in situ* hybridization showing that *nestin* and *pax2a* were

normally expressed in *bbh*^{fn40a}; Tg(*gfap:GFP-stmn1b*) embryos at 48 hpf.

Dorsal views with anterior to the left.

(K) Quantification of average CtA and glia length in Figure 5 (I). Each dot

represents one embryo. Data are presented in mean ± SEM; unpaired

Student's t test with individual P values mentioned in the figure.

(L) qRT-PCR analysis confirming the efficacy of *betaPix* transgenic

over-expression and *betaPix* siRNA knockdown in U251 cells. Data are

presented in mean ± SEM; one-way ANOVA analysis with Dunnett's test,

individual P values mentioned in the figure.

(M) Representative stereomicroscopy images of A172 cells at 0 and 18 hours

after wounding. *betaPIX* siRNA treatment decreased A172 cell migration,

which were rescued by either *betaPIX* or STMN1 overexpression. The wound

edges are highlighted by dashed lines, with arrow lines indicating the wound

width.

(N) Quantification of wound closures in (M). Data are presented in mean ± SEM;

one-way ANOVA analysis with Dunnett's test. **$P<0.01$ compared to negative

control siRNA and empty vector transfection. ##$P<0.01$, ###$P<0.005$ compared

to *betaPix* knockdown and empty vector transfecti on. Individual scale bars

indicated in the figure.

1220

Figure S8. *Zfhx3/4* acts downstream of *betaPix* to regulate vascular integrity

development.

(A) Violin plots showing that *Zfhx3* and *Zfhx4* decreased in progenitor sub-cluster

of *betaPix* knockouts.

(B) qRT-PCR analysis revealing that *Zfhx3* and *Zfhx4* decreased in FACS-sorted

glia of *gfap:GFP-Cre*; *betaPix^{ct/ct}* mutants compared with *betaPix^{ct/ct}* siblings

at 48 hpf. Each dot represents the cells sorted from one embryo. Data are

presented in mean ± SEM; unpaired Student's t test with individual P values

mentioned in the figure.

(C) Schematic diagram illustrates 4-guide RNAs mediated F_0 knockout strategy in

the *Zfhx3* and *Zfhx4* loci.

(D) Whole-mount RNA *in situ* hybridization confirming *Zfhx3/4* efficiency in

CRISPR-mediated *Zfhx3/4* F_0 knockout embryos at 48 hpf. Arrows indicate

significant reduction of *Zfhx3* and *Zfhx4* levels in hindbrain by *Zfhx3/4* double

knockout.

(E) qRT-PCR analysis revealing that *Zfhx3*, *Zfhx4*, and *Vegfaa* decreased in

CRISPR-mediated *Zfhx3/4* F_0 knockout embryos at 36 hpf. Data are presented

in mean ± SEM; unpaired Student's t test with individual P values mentioned

in the figure.

(F) Whole-mount RNA *in situ* hybridization revealing that *nestin* increased in

CRISPR-mediated *Zfhx3/4* F_0 knockout embryos at 36 hpf. Arrows indicate

the hindbrain regions.

(G) qRT-PCR analysis confirming overexpression of *Zfhx4* in CRISPR-mediated

betaPix F_0 knockout embryos with *Zfhx4* mRNA injection at 70% epiboly.

Data are presented in mean ± SEM; unpaired Student's t test with individual P

values mentioned in the figure.

(H) Representative stereomicroscopy images of erythrocytes stained with

o-dianisidine showing that brain hemorrhages (arrows) decreased in bbh^{fn40a}

mutants with *Zfhx4* mRNA injection at 48 hpf. Lateral views with anterior to

the left.

(I) Quantification of hemorrhagic parameters in (H). Left panel showing

hemorrhage percentages, with independent experiment as dot. Right panel

showing hemorrhage areas with each dot representing one embryo. #

represents the numbers of embryos scored for each analysis, three or more

individual experiments conducted. Data are presented in mean $\pm$ SEM;

unpaired Student's t test with individual *P* values mentioned in the figure.

Individual scale bars indicated in the figure.

References

References

1. Sheth, K.N. (2022). Spontaneous Intracerebral Hemorrhage. New England Journal of
Medicine *387*, 1589–1596. doi:10.1056/NEJMra2201449.

2. Chiang, P.-T., Tsai, L.-K., and Tsai, H.-H. (9900). New targets in spontaneous intracerebral
hemorrhage. Current Opinion in Neurology.

3. Thangameeran, S.I.M., Pang, C.Y., Lee, C.H., Tsai, S.T., Hu, W.F., and Liew, H.K. (2023).
Experimental animal models and evaluation techniques in intracerebral hemorrhage. Tzu Chi
Med J *35*, 1–10. 10.4103/tcmj.tcmj_119_22.

4. Isogai, S., Horiguchi, M., and Weinstein, B.M. (2001). The vascular anatomy of the
developing zebrafish: an atlas of embryonic and early larval development. Dev Biol *230*,
278–301. 10.1006/dbio.2000.9995.

5. Buchner, D.A., Su, F., Yamaoka, J.S., Kamei, M., Shavit, J.A., Barthel, L.K., McGee, B.,
Amigo, J.D., Kim, S., Hanosh, A.W., et al. (2007). pak2a mutations cause cerebral
hemorrhage in redhead zebrafish. Proceedings of the National Academy of Sciences *104*,
13996–14001. 10.1073/pnas.0700947104.

6. Jin, S.-W., Herzog, W., Santoro, M.M., Mitchell, T.S., Frantsve, J., Jungblut, B., Beis, D.,
Scott, I.C., D'Amico, L.A., Ober, E.A., et al. (2007). A transgene-assisted genetic screen
identifies essential regulators of vascular development in vertebrate embryos. Developmental
Biology *307*, 29–42.

7. Hegarty, J.M., Yang, H., and Chi, N.C. (2013). UBIAD1-mediated vitamin K2 synthesis is
required for vascular endothelial cell survival and development. Development *140*,
1713–1719. 10.1242/dev.093112.

8. Stainier, D.Y., Fouquet, B., Chen, J.N., Warren, K.S., Weinstein, B.M., Meiler, S.E.,
Mohideen, M.A., Neuhauss, S.C., Solnica-Krezel, L., Schier, A.F., et al. (1996). Mutations
affecting the formation and function of the cardiovascular system in the zebrafish embryo.
Development *123*, 285–292.

9. Chen, J.N., van Bebber, F., Goldstein, A.M., Serluca, F.C., Jackson, D., Childs, S., Serbedzija,
G., Warren, K.S., Mably, J.D., Lindahl, P., et al. (2001). Genetic steps to organ laterality in
zebrafish. Comp Funct Genomics *2*, 60–68. 10.1002/cfg.74.

10. Liu, J., Fraser, S.D., Faloon, P.W., Rollins, E.L., Vom Berg, J., Starovic-Subota, O., Laliberte,
A.L., Chen, J.N., Serluca, F.C., and Childs, S.J. (2007). A betaPix Pak2a signaling pathway
regulates cerebral vascular stability in zebrafish. Proc Natl Acad Sci U S A *104*, 13990–13995.
10.1073/pnas.0700825104.

11. Manser, E., Loo, T.-H., Koh, C.-G., Zhao, Z.-S., Chen, X.-Q., Tan, L., Tan, I., Leung, T., and
Lim, L. (1998). PAK Kinases Are Directly Coupled to the PIX Family of Nucleotide
Exchange Factors. Molecular Cell *1*, 183–192.

12. Zhou, W., Li, X., and Premont, R.T. (2016). Expanding functions of GIT Arf
GTPase-activating proteins, PIX Rho guanine nucleotide exchange factors and GIT-PIX
complexes. J Cell Sci *129*, 1963–1974. 10.1242/jcs.179465.

13. Morimoto, K., Tabata, H., Takahashi, R., and Nakajima, K. (2024). Interactions between
neural cells and blood vessels in central nervous system development. Bioessays *46*,
e2300091. 10.1002/bies.202300091.

14. Grinchevskaya, L.R., Salikhova, D.I., Silachev, D.N., and Goldshtein, D.V. (2024). Neural

and Glial Regulation of Angiogenesis in CNS in Ischemic Stroke. Bull Exp Biol Med *177*,
528–533. 10.1007/s10517-024-06219-4.

15.   McCarty, J.H. (2009). Cell adhesion and signaling networks in brain neurovascular units. Curr
Opin Hematol *16*, 209–214. 10.1097/MOH.0b013e32832a07eb.

16.   McCarty, J.H. (2009). Integrin-mediated regulation of neurovascular development, physiology
and disease. Cell Adh Migr *3*, 211–215. 10.4161/cam.3.2.7767.

17.   Liu, J., Zeng, L., Kennedy, R.M., Gruenig, N.M., and Childs, S.J. (2012). βPix plays a dual
role in cerebral vascular stability and angiogenesis, and interacts with integrin αvβ8. Dev Biol
363, 95–105. 10.1016/j.ydbio.2011.12.022.

18.   Yang, R., Zhang, Y., Huang, D., Luo, X., Zhang, L., Zhu, X., Zhang, X., Liu, Z., Han, J.Y.,
and Xiong, J.W. (2017). Miconazole protects blood vessels from MMP9-dependent rupture
and hemorrhage. Dis Model Mech *10*, 337–348. 10.1242/dmm.027268.

19.   Sugimoto, K., Hui, S.P., Sheng, D.Z., and Kikuchi, K. (2017). Dissection of zebrafish shha
function using site-specific targeting with a Cre-dependent genetic switch. Elife *6*.
10.7554/eLife.24635.

20.   Ogawa, M., Geng, F.S., Humphreys, D.T., Kristianto, E., Sheng, D.Z., Hui, S.P., Zhang, Y.,
Sugimoto, K., Nakayama, M., Zheng, D., et al. (2021). Krüppel-like factor 1 is a core
cardiomyogenic trigger in zebrafish. Science *372*, 201–205. 10.1126/science.abe2762.

21.   Ogawa, M., and Kikuchi, K. (2024). Generation of Conditional Knockout Zebrafish Using an
Invertible Gene-Trap Cassette. Methods Mol Biol *2707*, 205–214.
10.1007/978-1-0716-3401-1_13.

22.   Wierson, W.A., Simone, B.W., WareJoncas, Z., Mann, C., Welker, J.M., Kar, B., Emch, M.J.,
Friedberg, I., Gendron, W.A.C., Barry, M.A., et al. (2019). Expanding the CRISPR Toolbox
with ErCas12a in Zebrafish and Human Cells. Crispr j *2*, 417–433. 10.1089/crispr.2019.0026.

23.   Wu, R.S., Lam, II, Clay, H., Duong, D.N., Deo, R.C., and Coughlin, S.R. (2018). A Rapid
Method for Directed Gene Knockout for Screening in G0 Zebrafish. Dev Cell *46*, 112–125
e114. 10.1016/j.devcel.2018.06.003.

24.   Quiñonez-Silvero, C., Hübner, K., and Herzog, W. (2020). Development of the brain
vasculature and the blood-brain barrier in zebrafish. Dev Biol *457*, 181–190.
10.1016/j.ydbio.2019.03.005.

25.   Deacon, S.W., Beeser, A., Fukui, J.A., Rennefahrt, U.E., Myers, C., Chernoff, J., and Peterson,
J.R. (2008). An isoform-selective, small-molecule inhibitor targets the autoregulatory
mechanism of p21-activated kinase. Chem Biol *15*, 322–331.
10.1016/j.chembiol.2008.03.005.

26.   Bernardos, R.L., and Raymond, P.A. (2006). GFAP transgenic zebrafish. Gene Expr Patterns
6, 1007–1013. 10.1016/j.modgep.2006.04.006.

27.   Kim, C.H., Ueshima, E., Muraoka, O., Tanaka, H., Yeo, S.Y., Huh, T.L., and Miki, N. (1996).
Zebrafish elav/HuC homologue as a very early neuronal marker. Neurosci Lett *216*, 109–112.
10.1016/0304-3940(96)13021-4.

28.   Raj, B., Wagner, D.E., McKenna, A., Pandey, S., Klein, A.M., Shendure, J., Gagnon, J.A.,
and Schier, A.F. (2018). Simultaneous single-cell profiling of lineages and cell types in the
vertebrate brain. Nature Biotechnology *36*, 442–450. 10.1038/nbt.4103.

29.   Raj, B., Farrell, J.A., Liu, J., El Kholtei, J., Carte, A.N., Navajas Acedo, J., Du, L.Y.,
McKenna, A., Relić, Đ., Leslie, J.M., and Schier, A.F. (2020). Emergence of Neuronal

Diversity during Vertebrate Brain Development. Neuron *108*, 1058–1074.e1056.

10.1016/j.neuron.2020.09.023.

30.   Farnsworth, D.R., Saunders, L.M., and Miller, A.C. (2020). A single-cell transcriptome atlas

for zebrafish development. Developmental Biology *459*, 100–108.

31.   Etienne-Manneville, S. (2013). Microtubules in cell migration. Annu Rev Cell Dev Biol *29*,

471–499. 10.1146/annurev-cellbio-101011-155711.

32.   Kwon, Y., Jeon, Y.W., Kwon, M., Cho, Y., Park, D., and Shin, J.E. (2020). βPix-d promotes

tubulin acetylation and neurite outgrowth through a PAK/Stathmin1 signaling pathway. PLoS

One *15*, e0230814. 10.1371/journal.pone.0230814.

33.   Omelchenko, T., Rabadan, M.A., Hernandez-Martinez, R., Grego-Bessa, J., Anderson, K.V.,

and Hall, A. (2014). beta-Pix directs collective migration of anterior visceral endoderm cells

in the early mouse embryo. Genes Dev *28*, 2764–2777. 10.1101/gad.251371.114.

34.   Hsu, R.M., Hsieh, Y.J., Yang, T.H., Chiang, Y.C., Kan, C.Y., Lin, Y.T., Chen, J.T., and Yu,

J.S. (2014). Binding of the extreme carboxyl-terminus of PAK-interacting exchange factor β

(βPIX) to myosin 18A (MYO18A) is required for epithelial cell migration. Biochim Biophys

Acta *1843*, 2513–2527. 10.1016/j.bbamcr.2014.06.023.

35.   López Tobón, A., Suresh, M., Jin, J., Vitriolo, A., Pietralla, T., Tedford, K., Bossenz, M.,

Mahnken, K., Kiefer, F., Testa, G., et al. (2018). The guanine nucleotide exchange factor

Arhgef7/βPix promotes axon formation upstream of TC10. Scientific Reports *8*, 8811.

10.1038/s41598-018-27081-1.

36.   Cheng, K., Chahdi, A., Larabee, S.M., Tolaymat, M., Sundel, M.H., Drachenberg, C.B., Zhan,

M., Hu, S., Said, A.H., Shang, A.C., et al. (2023). Muscarinic receptor agonist-induced βPix

binding to β-catenin promotes colon neoplasia. Sci Rep *13*, 16920.

10.1038/s41598-023-44158-8.

37.   Connor, K., Murray, D.W., Jarzabek, M.A., Tran, N.L., White, K., Dicker, P., Sweeney, K.J.,

O'Halloran, P.J., MacCarthy, B., Shiels, L.P., et al. (2020). Targeting the RhoGEF

βPIX/COOL-1 in Glioblastoma: Proof of Concept Studies. Cancers (Basel) *12*.

10.3390/cancers12123531.

38.   Ward, J.D., Ha, J.H., Jayaraman, M., and Dhanasekaran, D.N. (2015). LPA-mediated

migration of ovarian cancer cells involves translocalization of Gαi2 to invadopodia and

association with Src and β-pix. Cancer Lett *356*, 382–391. 10.1016/j.canlet.2014.09.030.

39.   Bussmann, J., Wolfe, S.A., and Siekmann, A.F. (2011). Arterial-venous network formation

during brain vascularization involves hemodynamic regulation of chemokine signaling.

Development *138*, 1717–1726. 10.1242/dev.059881.

40.   Eubelen, M., Bostaille, N., Cabochette, P., Gauquier, A., Tebabi, P., Dumitru, A.C., Koehler,

M., Gut, P., Alsteens, D., Stainier, D.Y.R., et al. (2018). A molecular mechanism for Wnt

ligand-specific signaling. Science *361*. 10.1126/science.aat1178.

41.   Pitulescu, M.E., and Adams, R.H. (2014). Regulation of signaling interactions and receptor

endocytosis in growing blood vessels. Cell Adh Migr *8*, 366–377.

10.4161/19336918.2014.970010.

42.   Wang, J., Chen, Y., Zeng, Z., Feng, R., Wang, Q., Zhang, Q., Sun, K., Chen, A.F., Lu, Y., and

Yu, Y. (2021). HMGA2 contributes to vascular development and sprouting angiogenesis by

promoting IGFBP2 production. Exp Cell Res *408*, 112831. 10.1016/j.yexcr.2021.112831.

43.   Fu, C., An, N., Liu, J., A, J., Zhang, B., Liu, M., Zhang, Z., Fu, L., Tian, X., Wang, D., and

1389   Dong, J.T. (2020). The transcription factor ZFHX3 is crucial for the angiogenic function of

1390   hypoxia-inducible factor 1α in liver cancer cells. J Biol Chem *295*, 7060–7074.

1391   10.1074/jbc.RA119.012131.

1392   44.   Gudbjartsson, D.F., Holm, H., Gretarsdottir, S., Thorleifsson, G., Walters, G.B., Thorgeirsson,

1393   G., Gulcher, J., Mathiesen, E.B., Njølstad, I., Nyrnes, A., et al. (2009). A sequence variant in

1394   ZFHX3 on 16q22 associates with atrial fibrillation and ischemic stroke. Nat Genet *41*,

1395   876–878. 10.1038/ng.417.

1396   45.   Soriano-Tárraga, C., Lazcano, U., Giralt-Steinhauer, E., Avellaneda-Gómez, C., Ois, Á.,

1397   Rodríguez-Campello, A., Cuadrado-Godia, E., Gómez-González, A., Fernández-Sanlés, A.,

1398   Elosua, R., et al. (2020). Identification of 20 novel loci associated with ischaemic stroke.

1399   Epigenome-wide association study. Epigenetics *15*, 988–997.

1400   10.1080/15592294.2020.1746507.

1401   46.   Malik, R., Traylor, M., Pulit, S.L., Bevan, S., Hopewell, J.C., Holliday, E.G., Zhao, W.,

1402   Abrantes, P., Amouyel, P., Attia, J.R., et al. (2016). Low-frequency and common genetic

1403   variation in ischemic stroke: The METASTROKE collaboration. Neurology *86*, 1217–1226.

1404   10.1212/wnl.0000000000002528.

1405   47.   Kalvaitytė, M., and Balciunas, D. (2022). Conditional mutagenesis strategies in zebrafish.

1406   Trends Genet *38*, 856–868. 10.1016/j.tig.2022.04.007.

1407   48.   Obermeier, B., Daneman, R., and Ransohoff, R.M. (2013). Development, maintenance and

1408   disruption of the blood-brain barrier. Nat Med *19*, 1584–1596. 10.1038/nm.3407.

1409   49.   Langen, U.H., Ayloo, S., and Gu, C. (2019). Development and Cell Biology of the

1410   Blood-Brain Barrier. Annu Rev Cell Dev Biol *35*, 591–613.

1411   10.1146/annurev-cellbio-100617-062608.

1412   50.   Fleming, A., Diekmann, H., and Goldsmith, P. (2013). Functional characterisation of the

1413   maturation of the blood-brain barrier in larval zebrafish. PLoS One *8*, e77548.

1414   10.1371/journal.pone.0077548.

1415   51.   Xie, J., Farage, E., Sugimoto, M., and Anand-Apte, B. (2010). A novel transgenic zebrafish

1416   model for blood-brain and blood-retinal barrier development. BMC Dev Biol *10*, 76.

1417   10.1186/1471-213x-10-76.

1418   52.   Stewart, P.A., and Wiley, M.J. (1981). Developing nervous tissue induces formation of

1419   blood-brain barrier characteristics in invading endothelial cells: a study using quail--chick

1420   transplantation chimeras. Dev Biol *84*, 183–192. 10.1016/0012-1606(81)90382-1.

1421   53.   Daneman, R., Agalliu, D., Zhou, L., Kuhnert, F., Kuo, C.J., and Barres, B.A. (2009).

1422   Wnt/beta-catenin signaling is required for CNS, but not non-CNS, angiogenesis. Proc Natl

1423   Acad Sci U S A *106*, 641–646. 10.1073/pnas.0805165106.

1424   54.   Kuhnert, F., Mancuso, M.R., Shamloo, A., Wang, H.T., Choksi, V., Florek, M., Su, H.,

1425   Fruttiger, M., Young, W.L., Heilshorn, S.C., and Kuo, C.J. (2010). Essential regulation of

1426   CNS angiogenesis by the orphan G protein-coupled receptor GPR124. Science *330*, 985–989.

1427   10.1126/science.1196554.

1428   55.   Cullen, M., Elzarrad, M.K., Seaman, S., Zudaire, E., Stevens, J., Yang, M.Y., Li, X.,

1429   Chaudhary, A., Xu, L., Hilton, M.B., et al. (2011). GPR124, an orphan G protein-coupled

1430   receptor, is required for CNS-specific vascularization and establishment of the blood-brain

1431   barrier. Proc Natl Acad Sci U S A *108*, 5759–5764. 10.1073/pnas.1017192108.

1432   56.   Zhou, Y., and Nathans, J. (2014). Gpr124 controls CNS angiogenesis and blood-brain barrier

1433		integrity by promoting ligand-specific canonical wnt signaling. Dev Cell *31*, 248–256.
1434		10.1016/j.devcel.2014.08.018.
1435	57.	Thomas, M., and Augustin, H.G. (2009). The role of the Angiopoietins in vascular
1436		morphogenesis. Angiogenesis *12*, 125–137. 10.1007/s10456-009-9147-3.
1437	58.	Nico, B., Frigeri, A., Nicchia, G.P., Quondamatteo, F., Herken, R., Errede, M., Ribatti, D.,
1438		Svelto, M., and Roncali, L. (2001). Role of aquaporin-4 water channel in the development and
1439		integrity of the blood-brain barrier. J Cell Sci *114*, 1297–1307. 10.1242/jcs.114.7.1297.
1440	59.	Nicchia, G.P., Nico, B., Camassa, L.M., Mola, M.G., Loh, N., Dermietzel, R., Spray, D.C.,
1441		Svelto, M., and Frigeri, A. (2004). The role of aquaporin-4 in the blood-brain barrier
1442		development and integrity: studies in animal and cell culture models. Neuroscience *129*,
1443		935–945. 10.1016/j.neuroscience.2004.07.055.
1444	60.	Hartfuss, E., Galli, R., Heins, N., and Götz, M. (2001). Characterization of CNS precursor
1445		subtypes and radial glia. Dev Biol *229*, 15–30. 10.1006/dbio.2000.9962.
1446	61.	Malatesta, P., Appolloni, I., and Calzolari, F. (2008). Radial glia and neural stem cells. Cell
1447		Tissue Res *331*, 165–178. 10.1007/s00441-007-0481-8.
1448	62.	Qian, X., Shen, Q., Goderie, S.K., He, W., Capela, A., Davis, A.A., and Temple, S. (2000).
1449		Timing of CNS cell generation: a programmed sequence of neuron and glial cell production
1450		from isolated murine cortical stem cells. Neuron *28*, 69–80. 10.1016/s0896-6273(00)00086-6.
1451	63.	Price, J., Williams, B.P., and Götz, M. (1995). The generation of cellular diversity in the
1452		cerebral cortex. Ciba Found Symp *193*, 71–84; discussion 117–126.
1453		10.1002/9780470514795.ch4.
1454	64.	Grupp, L., Wolburg, H., and Mack, A.F. (2010). Astroglial structures in the zebrafish brain. J
1455		Comp Neurol *518*, 4277–4287. 10.1002/cne.22481.
1456	65.	Chen, J., Poskanzer, K.E., Freeman, M.R., and Monk, K.R. (2020). Live-imaging of astrocyte
1457		morphogenesis and function in zebrafish neural circuits. Nat Neurosci *23*, 1297–1306.
1458		10.1038/s41593-020-0703-x.
1459	66.	Johnson, K., Barragan, J., Bashiruddin, S., Smith, C.J., Tyrrell, C., Parsons, M.J., Doris, R.,
1460		Kucenas, S., Downes, G.B., Velez, C.M., et al. (2016). Gfap-positive radial glial cells are an
1461		essential progenitor population for later-born neurons and glia in the zebrafish spinal cord.
1462		Glia *64*, 1170–1189. 10.1002/glia.22990.
1463	67.	Umans, R.A., Pollock, C., Mills, W.A., 3rd, Clark, K.C., Pan, Y.A., and Sontheimer, H.
1464		(2021). Using Zebrafish to Elucidate Glial-Vascular Interactions During CNS Development.
1465		Front Cell Dev Biol *9*, 654338. 10.3389/fcell.2021.654338.
1466	68.	Matsuoka, R.L., Rossi, A., Stone, O.A., and Stainier, D.Y.R. (2017). CNS-resident
1467		progenitors direct the vascularization of neighboring tissues. Proc Natl Acad Sci U S A *114*,
1468		10137–10142. 10.1073/pnas.1619300114.
1469	69.	Alvarez, J.I., Dodelet-Devillers, A., Kebir, H., Ifergan, I., Fabre, P.J., Terouz, S., Sabbagh, M.,
1470		Wosik, K., Bourbonnière, L., Bernard, M., et al. (2011). The Hedgehog pathway promotes
1471		blood-brain barrier integrity and CNS immune quiescence. Science *334*, 1727–1731.
1472		10.1126/science.1206936.
1473	70.	Brilha, S., Ong, C.W.M., Weksler, B., Romero, N., Couraud, P.O., and Friedland, J.S. (2017).
1474		Matrix metalloproteinase-9 activity and a downregulated Hedgehog pathway impair
1475		blood-brain barrier function in an in vitro model of CNS tuberculosis. Sci Rep *7*, 16031.
1476		10.1038/s41598-017-16250-3.

1477   71.   Shi, T., Yue, S., Xie, C., Li, X., Yang, D., Hu, L., Zhong, Y., Zhang, Y., and Liu, W. (2024).
1478   MMP-2-mediated Scube2 degradation promotes blood-brain barrier disruption by blocking the
1479   interaction between astrocytes and endothelial cells via inhibiting Sonic hedgehog pathway
1480   during early cerebral ischemia. J Neurochem *168*, 1877–1894. 10.1111/jnc.16021.

1481   72.   Xing, G., Zhao, T., Zhang, X., Li, H., Li, X., Cui, P., Li, M., Li, D., Zhang, N., and Jiang, W.
1482   (2020). Astrocytic Sonic Hedgehog Alleviates Intracerebral Hemorrhagic Brain Injury via
1483   Modulation of Blood-Brain Barrier Integrity. Front Cell Neurosci *14*, 575690.
1484   10.3389/fncel.2020.575690.

1485   73.   Lawson, N.D., Vogel, A.M., and Weinstein, B.M. (2002). sonic hedgehog and vascular
1486   endothelial growth factor act upstream of the Notch pathway during arterial endothelial
1487   differentiation. Dev Cell *3*, 127–136. 10.1016/s1534-5807(02)00198-3.

1488   74.   Mannan, A., Dhiamn, S., Garg, N., and Singh, T.G. (2023). Pharmacological modulation of
1489   Sonic Hedgehog signaling pathways in Angiogenesis: A mechanistic perspective. Dev Biol
1490   *504*, 58–74. 10.1016/j.ydbio.2023.09.009.

1491   75.   Yan, J., Manaenko, A., Chen, S., Klebe, D., Ma, Q., Caner, B., Fujii, M., Zhou, C., and Zhang,
1492   J.H. (2013). Role of SCH79797 in maintaining vascular integrity in rat model of subarachnoid
1493   hemorrhage. Stroke *44*, 1410–1417. 10.1161/strokeaha.113.678474.

1494   76.   Jagadeeshan, S., Sagayaraj, R.V., Paneerselvan, N., Ghouse, S.S., and Malathi, R. (2017).
1495   Toxicity and anti-angiogenicity evaluation of Pak1 inhibitor IPA-3 using zebrafish embryo
1496   model. Cell Biol Toxicol *33*, 41–56. 10.1007/s10565-016-9358-5.

1497   77.   Zou, J., Li, W.Q., Li, Q., Li, X.Q., Zhang, J.T., Liu, G.Q., Chen, J., Qiu, X.X., Tian, F.J.,
1498   Wang, Z.Z., et al. (2011). Two functional microRNA-126s repress a novel target gene
1499   p21-activated kinase 1 to regulate vascular integrity in zebrafish. Circ Res *108*, 201–209.
1500   10.1161/circresaha.110.225045.

1501   78.   Braverman, R., Bhattacharya, B., Feuerstein, N., and Cooper, H.L. (1986). Identification and
1502   characterization of the nonphosphorylated precursor of pp17, a phosphoprotein associated
1503   with phorbol ester induction of growth arrest and monocytic differentiation in HL-60
1504   promyelocytic leukemia cells. J Biol Chem *261*, 14342–14348.

1505   79.   Luo, X.N., Mookerjee, B., Ferrari, A., Mistry, S., and Atweh, G.F. (1994). Regulation of
1506   phosphoprotein p18 in leukemic cells. Cell cycle regulated phosphorylation by p34cdc2
1507   kinase. J Biol Chem *269*, 10312–10318.

1508   80.   Curmi, P.A., Noguès, C., Lachkar, S., Carelle, N., Gonthier, M.P., Sobel, A., Lidereau, R., and
1509   Bièche, I. (2000). Overexpression of stathmin in breast carcinomas points out to highly
1510   proliferative tumours. Br J Cancer *82*, 142–150. 10.1054/bjoc.1999.0891.

1511   81.   Price, D.K., Ball, J.R., Bahrani-Mostafavi, Z., Vachris, J.C., Kaufman, J.S., Naumann, R.W.,
1512   Higgins, R.V., and Hall, J.B. (2000). The phosphoprotein Op18/stathmin is differentially
1513   expressed in ovarian cancer. Cancer Invest *18*, 722–730. 10.3109/07357900009012204.

1514   82.   Liedtke, W., Leman, E.E., Fyffe, R.E., Raine, C.S., and Schubart, U.K. (2002).
1515   Stathmin-deficient mice develop an age-dependent axonopathy of the central and peripheral
1516   nervous systems. Am J Pathol *160*, 469–480. 10.1016/s0002-9440(10)64866-3.

1517   83.   Belmont, L.D., and Mitchison, T.J. (1996). Identification of a protein that interacts with
1518   tubulin dimers and increases the catastrophe rate of microtubules. Cell *84*, 623–631.
1519   10.1016/s0092-8674(00)81037-5.

84.   Liu, A., Muggironi, M., Marin-Husstege, M., and Casaccia-Bonnefil, P. (2003).

Oligodendrocyte process outgrowth in vitro is modulated by epigenetic regulation of
cytoskeletal severing proteins. Glia *44*, 264–274. 10.1002/glia.10290.

85.   Carretero-Rodriguez, L., Guðjónsdóttir, R., Poparic, I., Reilly, M.L., Chol, M., Bianco, I.H.,
Chiapello, M., Feret, R., Deery, M.J., and Guthrie, S. (2021). The Rac-GAP alpha2-Chimaerin
Signals via CRMP2 and Stathmins in the Development of the Ocular Motor System. J
Neurosci *41*, 6652–6672. 10.1523/jneurosci.0983-19.2021.

86.   Lin, M.J., and Lee, S.J. (2016). Stathmin-like 4 is critical for the maintenance of neural
progenitor cells in dorsal midbrain of zebrafish larvae. Sci Rep *6*, 36188. 10.1038/srep36188.

87.   Jing, Y., Luo, Y., Li, L., Liu, M., and Liu, J.X. (2024). Deficiency of copper responsive gene
stmn4 induces retinal developmental defects. Cell Biol Toxicol *40*, 2.
10.1007/s10565-024-09847-8.

88.   Kim, T.S., Kawaguchi, M., Suzuki, M., Jung, C.G., Asai, K., Shibamoto, Y., Lavin, M.F.,
Khanna, K.K., and Miura, Y. (2010). The ZFHX3 (ATBF1) transcription factor induces
PDGFRB, which activates ATM in the cytoplasm to protect cerebellar neurons from oxidative
stress. Dis Model Mech *3*, 752–762. 10.1242/dmm.004689.

89.   Gonzalez, G.A., and Montminy, M.R. (1989). Cyclic AMP stimulates somatostatin gene
transcription by phosphorylation of CREB at serine 133. Cell *59*, 675–680.
10.1016/0092-8674(89)90013-5.

90.   Kim, J., and Kaang, B.K. (2022). Cyclic AMP response element-binding protein (CREB)
transcription factor in astrocytic synaptic communication. Front Synaptic Neurosci *14*,
1059918. 10.3389/fnsyn.2022.1059918.

91.   Shaywitz, A.J., and Greenberg, M.E. (1999). CREB: a stimulus-induced transcription factor
activated by a diverse array of extracellular signals. Annu Rev Biochem *68*, 821–861.
10.1146/annurev.biochem.68.1.821.

92.   Chung, J.H., Kim, D.H., Kim, Y.S., Son, B.S., Kim, D., Hwang, C., Shin, D., Noh, S.G., Han,
J.H., Kim, D.K., et al. (2018). Upregulation of P21-Activated Kinase 1 (PAK1)/CREB Axis in
Squamous Non-Small Cell Lung Carcinoma. Cell Physiol Biochem *50*, 304–316.
10.1159/000494007.

93.   Argaw, A.T., Asp, L., Zhang, J., Navrazhina, K., Pham, T., Mariani, J.N., Mahase, S., Dutta,
D.J., Seto, J., Kramer, E.G., et al. (2012). Astrocyte-derived VEGF-A drives blood-brain
barrier disruption in CNS inflammatory disease. J Clin Invest *122*, 2454–2468.
10.1172/jci60842.

94.   Chapouly, C., Tadesse Argaw, A., Horng, S., Castro, K., Zhang, J., Asp, L., Loo, H., Laitman,
B.M., Mariani, J.N., Straus Farber, R., et al. (2015). Astrocytic TYMP and VEGFA drive
blood-brain barrier opening in inflammatory central nervous system lesions. Brain *138*,
1548–1567. 10.1093/brain/awv077.

95.   Matsumoto, T., and Mugishima, H. (2006). Signal transduction via vascular endothelial
growth factor (VEGF) receptors and their roles in atherogenesis. J Atheroscler Thromb *13*,
130–135. 10.5551/jat.13.130.

96.   Jakobsson, L., Franco, C.A., Bentley, K., Collins, R.T., Ponsioen, B., Aspalter, I.M., Rosewell,
I., Busse, M., Thurston, G., Medvinsky, A., et al. (2010). Endothelial cells dynamically
compete for the tip cell position during angiogenic sprouting. Nature Cell Biology *12*,
943–953. 10.1038/ncb2103.

97.   Liu, Z.J., Shirakawa, T., Li, Y., Soma, A., Oka, M., Dotto, G.P., Fairman, R.M., Velazquez,

O.C., and Herlyn, M. (2003). Regulation of Notch1 and Dll4 by vascular endothelial growth
factor in arterial endothelial cells: implications for modulating arteriogenesis and angiogenesis.
Mol Cell Biol *23*, 14–25. 10.1128/mcb.23.1.14-25.2003.

98.   Olsson, A.K., Dimberg, A., Kreuger, J., and Claesson-Welsh, L. (2006). VEGF receptor
signalling - in control of vascular function. Nat Rev Mol Cell Biol *7*, 359–371.
10.1038/nrm1911.

99.   Xu, L., Nirwane, A., and Yao, Y. (2019). Basement membrane and blood-brain barrier. Stroke
Vasc Neurol *4*, 78–82. 10.1136/svn-2018-000198.

100. Jeanne, M., Jorgensen, J., and Gould, D.B. (2015). Molecular and Genetic Analyses of
Collagen Type IV Mutant Mouse Models of Spontaneous Intracerebral Hemorrhage Identify
Mechanisms for Stroke Prevention. Circulation *131*, 1555–1565.
10.1161/circulationaha.114.013395.

101. Halder, S.K., Sapkota, A., and Milner, R. (2022). The impact of genetic manipulation of
laminin and integrins at the blood-brain barrier. Fluids Barriers CNS *19*, 50.
10.1186/s12987-022-00346-8.

102. Yao, Y., Chen, Z.L., Norris, E.H., and Strickland, S. (2014). Astrocytic laminin regulates
pericyte differentiation and maintains blood brain barrier integrity. Nat Commun *5*, 3413.
10.1038/ncomms4413.

103. Milner, R., and Campbell, I.L. (2002). Developmental regulation of beta1 integrins during
angiogenesis in the central nervous system. Mol Cell Neurosci *20*, 616–626.
10.1006/mcne.2002.1151.

104. Nishimura, S.L., Boylen, K.P., Einheber, S., Milner, T.A., Ramos, D.M., and Pytela, R.
(1998). Synaptic and glial localization of the integrin alphavbeta8 in mouse and rat brain.
Brain Res *791*, 271–282. 10.1016/s0006-8993(98)00118-8.

105. McCarty, J.H., Lacy-Hulbert, A., Charest, A., Bronson, R.T., Crowley, D., Housman, D.,
Savill, J., Roes, J., and Hynes, R.O. (2005). Selective ablation of alphav integrins in the
central nervous system leads to cerebral hemorrhage, seizures, axonal degeneration and
premature death. Development *132*, 165–176. 10.1242/dev.01551.

106. Beis, D., Bartman, T., Jin, S.W., Scott, I.C., D'Amico, L.A., Ober, E.A., Verkade, H., Frantsve,
J., Field, H.A., Wehman, A., et al. (2005). Genetic and cellular analyses of zebrafish
atrioventricular cushion and valve development. Development *132*, 4193–4204.
10.1242/dev.01970.

107. Jin, S.W., Beis, D., Mitchell, T., Chen, J.N., and Stainier, D.Y. (2005). Cellular and molecular
analyses of vascular tube and lumen formation in zebrafish. Development *132*, 5199–5209.
10.1242/dev.02087.

108. Jeon, T.Y., Han, M.E., Lee, Y.W., Lee, Y.S., Kim, G.H., Song, G.A., Hur, G.Y., Kim, J.Y.,
Kim, H.J., Yoon, S., et al. (2010). Overexpression of stathmin1 in the diffuse type of gastric
cancer and its roles in proliferation and migration of gastric cancer cells. Br J Cancer *102*,
710–718. 10.1038/sj.bjc.6605537.

109. Chang, N., Sun, C., Gao, L., Zhu, D., Xu, X., Zhu, X., Xiong, J.W., and Xi, J.J. (2013).
Genome editing with RNA-guided Cas9 nuclease in zebrafish embryos. Cell Res *23*, 465–472.
10.1038/cr.2013.45.

110. Jao, L.-E., Wente, S.R., and Chen, W. (2013). Efficient multiplex biallelic zebrafish genome
editing using a CRISPR nuclease system. Proceedings of the National Academy of Sciences

110, 13904–13909. doi:10.1073/pnas.1308335110.

111.   Kawakami, K. (2004). Transgenesis and gene trap methods in zebrafish by using the Tol2

transposable element. Methods Cell Biol *77*, 201–222. 10.1016/s0091-679x(04)77011-9.

112.   Lei, L., Yan, S.Y., Yang, R., Chen, J.Y., Li, Y., Bu, Y., Chang, N., Zhou, Q., Zhu, X., Li,

C.Y., and Xiong, J.W. (2017). Spliceosomal protein eftud2 mutation leads to p53-dependent

apoptosis in zebrafish neural progenitors. Nucleic Acids Res *45*, 3422–3436.

10.1093/nar/gkw1043.

113.   Pang, M., Bai, L., Zong, W., Wang, X., Bu, Y., Xiong, C., Zheng, J., Li, J., Gao, W., Feng, Z.,

et al. (2020). Light-sheet fluorescence imaging charts the gastrula origin of vascular

endothelial cells in early zebrafish embryos. Cell Discovery *6*, 74.

10.1038/s41421-020-00204-7.

1620

Figure 1

Figure 2

Scale bar:100 μm

Scale bar: 500 μm

Scale bar: 200 μm

Scale bar: 150 μm

Scale bar:150μm

Scale bar:150μm

Figure 3

A *o*-dianisidine

48 hpf

betaPix^ct/ct^ DMSO / IPA3

gfap:GFP-Cre; betaPix^ct/ct^ DMSO / IPA3

Scale bar:150μm

B *Tg(kdrl:mCherry)*

betaPix^ct/ct^ DMSO / IPA3

gfap:GFP-Cre; betaPix^ct/ct^ DMSO / IPA3

Scale bar:150 μm

C

Hemorrhage % — *P*<0.0001 / *P*<0.0001 — #=489, #=370, #=104, #=102

Hemorrhage area (mm^2) — *P*<0.0001 / *P*<0.0001 — #=4

Normalized CtA length index — *P*<0.0001 / *P*=0.0079

● *betaPix*^ct/ct^
● *gfap:GFP-Cre; betaPix*^ct/ct^

DMSO IPA3

Figure 4

Scale bar: 500 μm

Figure 5

Figure 6

Figure 7

Figure S1

A

Splice acceptor | GSG | P2A | TagRFP

TTTTTCAGTCGATCGACAAACTCTTCGCGGTCGGAAGAAGAAGAAGAG CTACTAACTTCAGCCTGCTGAAGCAGGCTGGAGACGTGGAGGAGAACCCTGGACCTGACGTCTGTGTCTAAGGGCGAAGAG

Zwitch (+3)

TTTTTCAGTCGATCGACAAACTCTTCGCGGTCGGAAGAAGAAGAAGAGAGCGGAGCTACTAACTTCAGCCTGCTGAAGCAGGCTGGAGACGTGGAGGAGAACCCTGGACCTGACGTCTGTGTCTAAGGGCGAAGAG

Zwitch (+3) add GSG spacer

B

5'junction (F1+R1)

genome | Upstream (left) homology arm | loxP | plasmid

AGGGAAACTCCATCATCAACAAAGACAGGAAGTGCAGCAGTCAGAAGCACCACTTCCCCTTTCCTCTGACGCGTAAACTTCGTATAGCATACATTATACGAAGTTATCCAAGCTTCACCATCGACCCGAAT

3'junction (F2+R2)

plasmid | FRT | Downstream (right) homology arm | genome

GGCTGGACGTAAACTCCTCTTCAGACCTGAAGTTCCTATACTTTCTAGAGAATAGGAACTTCGGAATTCAAATGGCTTCACACTCCAGTTTATTCCACTTTCCGAGAGAGACTTAGAGATCGACTCTTTTA

C

Screening process	Number of positive fish	ratio
EGFP screening at F0	184	/
EGFP screening at F1	84	45.65%
Genomic PCR diagnosis	7	8.33%
Sequence diagnosis	2	28.57%
Overall	**2**	**1.09%**

D

o-dianisidine

48 hpf

bbh^{fn40a}

Scale bar: 500 μm

E

sibling #=135, *bbh^{fn40a}* #=244, *P*<0.0001

sibling #=6, *bbh^{fn40a}* #=6, *P*<0.0001

● sibling
● *bbh^{fn40a}*

F

24 hpf | 48 hpf | 72 hpf | 144 hpf

cryaa:Venus
betaPix:RFP

betaPix^{cl/+}

PBS

Cre mRNA

Scale bar: 150 μm

G

24 hpf | 48 hpf | 72 hpf | 144 hpf

neurod1:EGFP
betaPix:RFP

betaPix^{cl/+}

PBS

Cre mRNA

Scale bar: 150 μm

Figure S2

A *Tg(kdrl*:GFP); KI(*betaPix*:RFP)

*betaPix^ct/ct^*

48 hpf

*betaPix^m/m^*

Scale bar: 100 μm

B

P=0.0009

CtA length (μm)

betaPix^ct/ct^   betaPix^m/m^

P=0.0003

Normalized CtA length index

0.6

0.5

0.4

0.3

0.2

betaPix^ct/ct^   betaPix^m/m^

C *Tg(gfap*:GFP)

48 hpf   D

*betaPix^ct/ct^*

V

D

*betaPix^m/m^*

V

Scale bar: 100 μm

Figure S3

A betaPix locus

E1 E5 E10

RhoGEF domain

■ exon
| gRNA target site

B
Relative betaPix expression

P<0.0001

sibling   betaPix KO

C Tg(gfap:GFP)

48 hpf

sibling

betaPix KO

Scale bar: 100μm

D
nestin   pax2a

48 hpf

sibling

13/15   16/16

betaPix KO

10/14   12/15

Scale bar: 500 μm

E
Hemorrhage %

P<0.0001

sibling
#=424

betaPix KO
#=268

Hemorrhage area (mm²)

P<0.0001

sibling
#=8

betaPix KO
#=8

F
CtA length (μm)

P=0.0064

sibling   betaPix KO

Normalized CtA length index

P=0.0005

sibling   betaPixKO

G
Glia length (μm)

P=0.0003

sibling   betaPixKO

Normalized glia length index

P=0.0179

sibling   betaPix KO

Figure S4

A

ptol2-gfap-gfp-cre donor vector

Tol2 | *gfap* promoter | eGFP | T2A | Cre | polyA | Tol2

Ki(betaPix:Zwitch)^flox/flox X *Ki(betaPix:Zwitch)^flox/flox*

Microinjection (*tol2* mRNA+ ptol2-gfap-gfp-cre donor vector)

Ki(betaPix:Zwitch)^flox/flox; Tg(gfap:GFP-Cre) founder   *Ki(betaPix:Zwitch)^flox/flox*

X

Tg(gfap:GFP-Cre)^tg/+; Ki(betaPix:Zwitch)^flox/flox
Ki(betaPix:Zwitch)^flox/flox

B

Tg(gfap:Cre-GFP); Ki(betaPix:RFP)

betaPix^ct/ct

48 hpf

gfap:GFP-Cre; betaPix^ct/ct

Scale bar: 150 μm

C

betaPix^ct/+ | *gfap:GFP-Cre; betaPix^ct/+* | *gfap:GFP-Cre; betaPix^ct/ct*

Size (bp)

2000
1500 — betaPix^ct

750
500 — betaPix^m

D

Relative *Cre* expression

P=0.0088

non-glia   glia

E

betaPix

Relative expression

P<0.0001

betaPix^ct/ct   *gfap:GFP-Cre; betaPix^ct/ct*

betaPix-RFP

Relative expression

P<0.0001

betaPix^ct/ct   *gfap:GFP-Cre; betaPix^ct/ct*

alphaPix

Relative expression

P=0.8092

betaPix^ct/ct   *gfap:GFP-Cre; betaPix^ct/ct*

F

alphaPix | *betaPix*

betaPix^ct/ct

48 hpf

28/28 | 27/28

gfap:GFP-Cre; betaPix^ct/ct

25/25 | 22/22

Figure S5

Scale bar: 200 μm

Figure S6

Figure S8

Supplementary Table 1.xlsx

Supplementary Table 2 | Primers used for siRNA

Gene	Forward
betaPIX	AGCCUUCAGAUGAGGAGUUCGCGUC
STMN1	UAAAGAGAACCGAGAGGCA

Supplementary Table 3 | Primers used for guide RNA synthesis

oligo	
betaPix-1	TAATACGACTCACTATAGGATGCCTGCAAAAATCCTTGTTTTAGAGCTAGAAATAGC
betaPix-2	TAATACGACTCACTATAGGACGAGTTATCCTTCAGCAGTTTTAGAGCTAGAAATAGC
betaPix-3	TAATACGACTCACTATAGGCGCTCTCAATGGCAAAACGTTTTAGAGCTAGAAATAGC
betaPix-4	TAATACGACTCACTATAGGAGAGTTCATGGAAAGTAAGTTTTAGAGCTAGAAATAGC
stmn1a-1	TAATACGACTCACTATAGAGCTGGACAAGCGTGCTTCGTTTTAGAGCTAGAAATAGC
stmn1a-2	TAATACGACTCACTATAGAAGAAGGACCTTTCCTTGGGTTTTAGAGCTAGAAATAGC
stmn1a-3	TAATACGACTCACTATAAGAAAAGCGTGAGCATGAAAGTTTTAGAGCTAGAAATAGC
stmn1a-4	TAATACGACTCACTATAGAAACTGAACCAGAAAATGGGTTTTAGAGCTAGAAATAGC
stmn1b-1	TAATACGACTCACTATAGGACAAGCGTGCCTCAGGACGTTTTAGAGCTAGAAATAGC
stmn1b-2	TAATACGACTCACTATAGGGAATTCTCCCTTGGCATCGTTTTAGAGCTAGAAATAGC
stmn1b-3	TAATACGACTCACTATAGGAAAGTGGGAATTCTCCCTGTTTTAGAGCTAGAAATAGC
stmn1b-4	TAATACGACTCACTATAGGAGGTCCTTCAGAAAGCTAGTTTTAGAGCTAGAAATAGC
stmn4l-1	TAATACGACTCACTATAGAGCAGAAGAGAGACACTAGGTTTTAGAGCTAGAAATAGC
stmn4l-2	TAATACGACTCACTATAGGGAATTATCAAAGACATGGGTTTTAGAGCTAGAAATAGC
stmn4l-3	TAATACGACTCACTATAAGTGAAGGGTCCCGGCGAGGGTTTTAGAGCTAGAAATAGC
stmn4l-4	TAATACGACTCACTATAGGCTCACATCGCCGCCATGCGTTTTAGAGCTAGAAATAGC
zfhx3-1	TAATACGACTCACTATAGGGAGAGGGAGCTCACCAAAGTTTTAGAGCTAGAAATAGC
zfhx3-2	TAATACGACTCACTATAGGGCGATCCCTCACTTGGAGGTTTTAGAGCTAGAAATAGC
zfhx3-3	TAATACGACTCACTATAGGGCTCCACAGTTGGCAGCTGTTTTAGAGCTAGAAATAGC
zfhx3-4	TAATACGACTCACTATAGGCCCACTTCGTGAGCAATGGTTTTAGAGCTAGAAATAGC
zfhx4-1	TAATACGACTCACTATAGGTTGTCTTCATCTGGAGGAGTTTTAGAGCTAGAAATAGC
zfhx4-2	TAATACGACTCACTATAGGAACTGAAACCACTAAAGGGTTTTAGAGCTAGAAATAGC
zfhx4-3	TAATACGACTCACTATAGGAAGTGCGGAAAGAACGCTGTTTTAGAGCTAGAAATAGC
zfhx4-4	TAATACGACTCACTATAGGCTGGCAAGTTGAGTACATGTTTTAGAGCTAGAAATAGC
UgRNA Universal	TAATACGACTCACTATAGGGAGGCGTTCGGGCCACAGGTTTTAGAGCTAGAAATAGC
bottom-str and Ultramer	AAAAGCACCGACTCGGTGCCACTTTTTCAAGTTGATAACGGACTAGCCTTATTTTAACTTGCTATTTCTAGCTCTAAAAC

Supplementary Table 4 | Primers used for WISH probe synthesis

Gene	Forward	Reverse*
alphaPix	GGACACTCTCGATCCCAAACTGAC	GCCACACACTCCCGTCTCAC
betaPix	GCCATACACTGCCCTCTCACC	GCAGCTTTGCCAGACCACATC
RFP	GTGTCTAAGGGCGAAGAGCTGAT	CAGTTTGCTAGGGAGGTCGCAG
stmn1a	GGCTGCTACAAGTGACATTCAGG	GCAGACAGTAACAGGCATGTTAAGC
stmn1b	GCGTCCTCTGGAGCTGATATCC	GGCAGTGGCAAAACTGGCAC
stmn2a	CTGGAGGCGGCTGAGGAC	CAAGCTGTCTATCCGAAGGGTCC
stmn4l	ATGACTCTCGCCGCGTACAG	GATGCAAGTCAACTGTGGCCAC
vegfaa	ACTCACCGCAACACTCCACT	GAGCAAGGCTCACAGTGGTT
zfhx3	CTGAGCAGCACGAAGTGGATG	CAATCTGATGGAGTCCGATCCTACAC
zfhx4	GTTACAGGACAACGCGCATATCC	CACCCTTGACAGCGTTCACATTC

*The T7 sequence TAATACGACTCACTATAGGG was added to the 5' end of each reverse primer.

Supplementary Table 5 | Primers used for qRT-PCR

Zebrafish Gene	Forward	Reverse
alphaPix	ATGTTAAACCTTTAACCATGCCTGG	TTAGTGACCTCCTTCATCCCAGGAA
betaPix	GACCAAAGGCTCCGACAAAC	GGCCTTGGGTGGACTCTTC
cre	AATTTACTGACCGTACACCAAA	ATAACCAGTGAAACAGCATTGC
gapdh	GATACACGGAGCACCAGGTT	GCCATCAGGTCACATACACG
pak1	CTCTGCTGGGAGTATAGTCAGCG	CTGCTGCTGCAGGTTCATCTG
ppfia3	CCAGACGAGGATGGCACTAAGAC	CCCTTGGAGATGCATCGCTCTG
RFP	GTGTCTAAGGGCGAAGAGCTGAT	GGTGGTTGTTCACGGTGCC
stmn1a	GGCTGCTACAAGTGACATTCAGG	CATAGCTGCCATAATTGCTGTACGG
stmn1b	GCGTCCTCTGGAGCTGATATCC	CTGCCTCATGGTTCTTGCGTC
stmn4l	CACATTCCCACCTCGCCG	CTCAGCATGCTTGTCCTTCTCC
vegfaa	TCCAGGAGTATCCCGATGAG	GCTTTGACTTCTGCCTTTGG
zfhx3	GCCACTGTGGCAGCATCAG	CAGGCTCAGCATCTGCCAC
zfhx4	GAGCAGGTGCAGCAATGCC	CGCAGCTGAATGCAATGCAAG
ZO-1	CCGAACCCATCAACCGCATC	GAGTGAGATCCTTGGCACTCGG

Human Gene	Forward	Reverse
betaPIX	CCTCACGGAACACAGTGAGGAG	GTCAGGATCTGCAGCTCAAGC
gapdh	CGCTGAGTACGTCGTGGAGTC	CAGGAGGCATTGCTGATGATCTTG
pak1	GTGAAGGCTGTGTCTGAGACTC	GGAAGTGGTTCAATCACAGACCG
stmn1	GAACTGGAGAAGCGTGCCTCAG	CTTCAGTCTCGTCAGCAGGGTC
VEGFA	CCATGCCAAGTGGTCCCAG	GGTCTCGATTGGATGGCAGTAGC
ZFHX3	CCACGGACCCAGAAGAAGCC	CTTCTCTGCTTGGCTGGACG
ZFHX4	CAGAGGAGGACAACCTCAGTGAG	CCTCACTTTGTTCTTCTGAGGTCG
ZO-1	CCTGGGGGCTGTCTCAACTCC	CATGCTGGGCCGAAGAATCC