Sex differentiation, divergence, and introgression in the leishmaniasis vector *Lutzomyia longipalpis*

Authors:

Adam MM Stuckert[1]*, David Turissini[2], Emmanuel RR D'Agostino[3], David Peede[4], Maya Parvathaneni[2], Jonathan A. Rader[2], Carlos Muskus[5], Eduar Bejarano[6], Jeremy R. Wang[7], Rafael Vivero[5,8], Daniel R. Matute[2]*

[1]Department of Biology and Biochemistry, University of Houston, Houston, TX 77204, USA
[2]Biology Department, University of North Carolina, Chapel Hill, NC 27514, USA
[3]Department of Ecology and Evolutionary Biology, Princeton University, Princeton, NJ 08544, USA
[4]Department of Ecology, Evolution, and Organismal Biology, Brown University, Providence, RI 02912, USA
[5]PECET (Programa de Estudio y Control de Enfermedades Tropicales), Unidad de Entomología Médica, Universidad de Antioquia, SIU-Sede de Investigación Universitaria
[6]Grupo de investigaciones Biomédicas, Línea de Entomología Médica, Universidad de Sucre, Sincelejo, Colombia
[7]Department of Pathology, University of North Carolina, Chapel Hill, NC 27514, USA
[8]Grupo de Microdiversidad and Bioprospección, Facultad de Ciencias, Universidad Nacional de Colombia, Sede Medellín, Medellín, Colombia

Email: astuckert@uh.edu and dmatute@email.unc.edu

Keywords: Diptera, Psychodidae, sex-determination, genomics, disease vector

SIGNIFICANCE: Leishmaniasis has emerged as an important public health issue. Transmission occurs exclusively through Phlebotomine sandflies of the genera *Lutzomyia* and *Phlebotomus*, yet our ability to prevent its spread is constrained by limited knowledge of the biological and ecological factors that make these insects effective vectors. In this manuscript, we leverage whole-genome sequencing to investigate the evolutionary forces shaping genetic diversity in *Lutzomyia longipalpis*, a key vector of leishmaniasis.

<u>ABSTRACT</u>

An estimated one billion people are at risk of acquiring the disease leishmaniasis
annually. This disease, among others, is transmitted to humans via sandfly vectors of the family
Psychodidae. In the Neotropics, *Lutzomyia longipalpis* is one of the most common vectors of
leishmaniasis, but little is known about its genetics and population structure, hampering control
efforts. We generated sex-specific genome assemblies of *Lu. longipalpis* from Colombia, and
examined population genomics of this species. We found that sex accounts for ~11% of
genome-wide variation, primarily driven by differentiation between the sexes in an XY sex
chromosome. Analyses of transposable elements indicate that the size difference in the sex
chromosome is driven by more transposable elements in the male chromosome. Further, we
examined population genomics and structure across the South American continent. We
identified significant genetic divergence between Colombian and Brazilian sandflies and
population structure within *Lu. longipalpis sensu lato* samples collected from the same sites in
Brazil. These results strongly suggest the presence of multiple, sympatric species within a
cryptic species complex, concordant with other published phenotypic evidence.

INTRODUCTION

Phlebotomine sandflies of the genera *Lutzomyia* and *Phlebotomus* are the only known vectors of *Leishmania*, a protozoan parasite that is transmitted to humans[1,2]. *Leishmania* infections can lead to the clinical disease leishmaniasis, which can cause skin ulcers and scarring, destruction of mucous membranes, extreme organ damage (e.g., liver, spleen, and bone marrow), and can be fatal in some cases. *Leishmania* is endemic to 98 countries and an estimated one billion people are at risk of infection[3,4]. Despite its high prevalence and disease burden, leishmaniasis is poorly understood, and thus is classified as a neglected tropical disease[5,6]. Although leishmaniasis is largely regarded as a tropical disease, *Lutzomyia* ranges are expanding northward[7] and the U.S. is experiencing increasing caseloads of leishmaniasis[7–9]. Ecological niche modeling predicts that leishmaniasis is likely to become more common in the U.S. due to climate change[10], and this is already evident via increasing clinical cases[11]. Treating leishmaniasis is difficult and often prohibitively expensive for individuals in underserved areas with endemic *Leishmania,* thus, preventing the spread of leishmaniasis is critically important[12]. Perhaps the most effective way to prevent the transmission of vector-transmitted diseases like leishmaniasis is to control the vector itself[13]. Despite the clear importance of *Leishmania'*s vectors for human well-being, preventive efforts are hampered by our lack of knowledge of the genetic and ecological factors that make sandflies effective vectors of *Leishmania*. Compounding the problem, functional studies of leishmaniasis vector biology are hindered by a variety of gaping holes in scientific knowledge, including: taxonomy and phylogenetics of sandflies, genetic characteristics of sandfly susceptibility to leishmaniasis, and a lack of genomic resources.

In the Neotropics, there are ~40 *Lutzomyia* species that transmit *Leishmania*, but one of the most commonly implicated and widespread is *Lutzomyia longipalpis.* This species has a range from Argentina through Mexico[2,14]. Complicating our understanding of this species' biology, a variety of lines of evidence suggest that *Lu. longipalpis* is actually a species complex containing a number of cryptic species with substantial phenotypic divergence across its geographic range[15,16]. This evidence includes variation in microsatellites of key genes[17–19], courtship song[20], morphology[21,22], and pheromones[18]. Nevertheless, this "species" is thought to be the major vector of *Leishmania infantum,* the causative agent of leishmaniasis in the Neotropics[23]. This species inhabits a wide variety of microhabitats, including silvan habitats and inside of rural houses[24–29]. Like most phlebotomine sandflies, females of this species are hematophagous (feed on blood), and can spread leishmaniasis to birds, dogs, pigs, horses, humans, and a variety of other mammalian hosts during feeding, meaning that common subsistence animals or pets can be intermediate hosts and promote vector populations[30]. Dramatic environmental changes in human land use (e.g. deforestation, roads, etc.) have caused concomitant changes in *Lu. longipalpis* distribution and habitat use. Of particular note is that *Lu. longipalpis* has shown a shift in recent decades to live and breed in highly populated urban areas[27]. Given that this species is a common vector of leishmaniasis, this puts an ever-increasing number of people at risk of acquiring leishmaniasis.

In this piece, we address fundamental genetic questions in the widespread leishmaniasis vector *Lutzomyia longipalpis.* First, we present highly-contiguous chromosomal and sex-specific genome assemblies of *Lu. longipalpis.* We then use these new assemblies as tools to examine

population genomics of this sandfly from the Andean region of Colombia. We then further use
publicly available genomic data from Brazilian *Lu. longipalpis* to examine population genomics
of this vector species across the South American continent. As predicted by previous work
suggesting *Lu. longipalpis sensu lato* is a cryptic species complex, we find extensive population
differentiation between Colombian and Brazilian populations. Additionally, our population
genomic analyses indicate extremely low nucleotide diversity, elevated runs of homozygosity,
and lower effective population sizes in Colombia relative to any Brazilian population. Our results
provide a roadmap for the implementation of genomic analyses in sandflies and to ultimately
understand the molecular and genetic underpinnings of the interactions between vectors and
parasites.
109
110
RESULTS
Genome sequencing:
We produced sex-specific genomes for female and male Colombian sandflies (Table 1).
Table S1 lists all the accession numbers. The initial female contig assembly built from PacBio
HiFi reads was 148 Mbp in length spread across 2,254 contigs and had an N50 of 1.25 Mbp.
The male contig assembly built from PacBio HiFi reads was 159.7 Mbp in length spread across
1,835 contigs with an N50 of 1.14 Mbp. The inclusion of HiC data improved our genome
assembly. Scaffolding our assemblies with HiC data yielded putative chromosome-level
assemblies for both female and male sandflies. The female assembly was 151.2 Mbp in length
and contained four chromosomes with lengths of 47.7, 37.2, 29.0, and 22.8 Mbp and
represented over 95% of the total genome. The female assembly contained >95% of complete
dipteran orthologs: 94.16% in complete single copies, 1.13% in complete duplicated copies,
0.46% fragmented, and 4.26% missing. The male assembly was 159.7 Mbp in length and
contained four chromosomes of 49.0, 36.8, 29.3, and 22.5 Mb, representing over 95% of the
total genome. The male assembly contained just under 95% of complete dipteran orthologs:
93.49% in complete single copies, 1.19% in complete duplicated copies, 0.33% fragmented,
and 4.99% missing.
Next, we identified the differences between the male and female genomes (Figure 1A).
We identified 640,461 unique kmers associated with sex-limited chromosomes using SRY on
our Illumina data. These kmers mapped to 111,862 HiFi reads spanning 888 Mbp in the male
dataset. Our initial draft "Y" chromosome assembly was 26.7 Mbp across 740 contigs. After
scaffolding, we placed 691 contigs (93.4% of contigs) into 50 scaffolds in an assembly that was
26.8 Mbp in length (including placed gaps) with a largest scaffold of 26.2 Mbp (thus the other 49
scaffolds contain ~0.6 Mbp). Thus, we were able to scaffold nearly the entire section of the "Y"
chromosome that is limited to males into a single scaffold, with minimal unplaced content
outside of this scaffold.
137
138
139
140   **Table 1. Genome statistics.** The Brazilian assembly comes from[31], and is derived from a
141   colony collected in the 1980s from Jacobina, Bahia State, Brazil. Our female and male
142   assemblies are derived from wild-caught sandflies from Ricaurte, Colombia.

	Brazilian *Lu. longipalpis* assembly	Female contig	Female scaffolded	Male contig	Male scaffolded
Summary					
Assembled bases (Mbp)	159.7	1997	151.2	159.7	159.7
Number of contigs	383	151.2	2002	1835	1849
Contig N50 (Mbp)	1.04	1	1	1.1	1.1
Number of scaffolds	138	-	1742	-	1618
Scaffold N50 (Mbp)	40.6	-	37.2	-	36.8
Percentage of assembly in scaffolds	93.2	-	91	-	86.7
BUSCO (percentages)					
Single copies	90.5	94.12	94.16	93.9	93.49
Duplicated copies	6.73	1.13	1.13	1.19	1.19
Fragmented copies	0.33	0.46	0.46	0.4	0.33
Incomplete copies	0	0	0	0	0
Missing	2.89	4.29	4.26	5.02	4.99
RepeatMasker (Mbp)					
Percentage of repeats (%)	-	-	15.69	-	15.4
Repeat bases (Mbp)	-	-	23.7	-	24.6
Retroelements	-	-	4.37	-	4.92
DNA transposons	-	-	19.3	-	19.6
Small RNA	-	-	0.39	-	0.0319
Simple repeat	-	-	1.3	-	1.29
Low complexity	-	-	0.47	-	0.5
Total bases masked	-	-	25.69	-	26.67

Comparison between the sexes indicates increased transposable element content in males on
the sex chromosome:
Transposable elements (TEs) are considered important drivers of sex chromosome
evolution[32,33], so we quantified transposable element content in the female and male
assemblies. RepeatMasker identified and masked a total of 16.99% (female) and 16.70% (male)
of the genome, 15.69% and 15.4% of which were TEs in the female and male assemblies
respectively (Figure 1B). Most repeats were DNA transposons (12.8% and 12.3% of the
genome in females and males respectively) or Retroelements (2.9% and 3.1% of the genome in
females and males respectively, Figure 1C). A small proportion of the genome were simple
repeats (0.9% and 0.8% of the genome in females and males respectively). The difference in
relative proportion of transposable elements between the sexes is largely driven by
chromosome 1. This chromosome is not only longer in males (49 Mbp vs 47.7 Mbp), but also
has a larger overall proportion of bases masked in males (15.89% vs 13.60%, Figure 1C). The

male chromosome 1 has a total of 7.8 Mbp of transposable elements masked by our repeat
masking pipeline, whereas the female chromosome has 6.499 Mbp of transposable elements
masked. The difference, 1.3 Mbp, corresponds to the total difference in size between the
assembled male and female chromosome 1 (see Figure S1). In the three autosomal
chromosomes, the proportion of transposable elements to assembled chromosome length is
<0.5% different between females and males.

164
165   **Figure 1. The sex chromosomes of *Lu. longipalpis*. A.** Synteny plot of alignments between
166   assembled male (presumptive) X and Y chromosome sequences. **B.** Number of repeats in the
167   female and male genome assemblies based on genomic position. **C.** Violin plots of TEs in the
168   female and male genome assemblies and their length.
169
170
171

Genome annotation:
We annotated both sexes of our Colombian *Lu. longipalpis.* In total, we annotated
14,559 genes in the male assembly; 10,874 of these predicted genes were in the four
chromosomes (4008, 2607, 2371, and 1888 in chromosomes 1-4 respectively), as opposed to in
the contigs. In the female assembly, we annotated 12,029 genes in total; 10,934 of these
predicted genes were in the four chromosomes (3,922; 2,662; 2,371; and 1,979 in
chromosomes 1-4 respectively). The predicted genes are approximately equal to RefSeq
predictions in *Lu. longipalpis* from other localities (GCF_024334085.1), which has 12,045
predicted genes. In both sexes of our assemblies, the majority of predicted genes are found in
the chromosomes. However, our male assembly has approximately 2,000 more predicted genes
than existent assemblies. This is due to the presence of multiple copies of genes between the
heteromorphic chromosomes — most of which are represented as unplaced contigs. While
females had more predicted genes overall in the regions of the genome that are located in the
chromosomes, males had a greater number of predicted genes owing to the predicted genes
found in un-scaffolded contigs. Further, males had more predicted genes in chromosome 1, the
heteromorphic chromosome. We identified a total of 3,462 genes annotated in the highly
differentiated region of chromosome 1 (average $F_{ST} > 0.3$ between females and males) in the
male assembly. Of these, we tentatively assigned identifications to 3,611 genes based on
protein evidence from OrthDB version 11 and VectorBase version 67. Among the genes found
in this region of high differentiation between females and males within our Colombian population
are a number of genes related to courtship (e.g., *takeout*), sex-differentiation (e.g., *sex-lethal,*
fruitless, doublesex- and mab-3-related transcription factor A2, lethal(2)), testes-specific genes
(e.g., ejaculatory bulb-specific protein 3, ejaculatory bulb-specific protein 3-like, testis-expressed
protein 10 homolog, motile sperm domain-containing protein), putatively related to feeding and
finding mates (e.g., gustatory receptor genes, gustatory receptor for sugar taste 64f, odorant
receptor genes), or related to blood meals (e.g., venom allergen 5).
Nucleotide diversity across the genome is similar in females (π = 0.00177 ± 0.00233 SD
female assembly; π = 0.00170 ± 0.00228 SD male assembly) and males from Colombia (π =
0.00171 ± 0.00236 SD female assembly; π = 0.00159 ± 0.00225 SD male assembly; Figures
2A-2B). However, concordant with other results, nucleotide diversity was lower on chromosome
1 in female sandflies (π = 0.00138 ± 0.00192 SD female assembly; π = 0.00131 ± 0.00196 SD
male assembly) than males (π = 0.00165 ± 0.00239 SD female assembly; π = 0.00160 ±
0.00225 SD male assembly).
Next, we studied the extent of differentiation between the *X*-chromosome and the
potentially male-restricted chromosome using PCA. We saw evidence of genomic differentiation
between female and male genomes in our study. PC1, which corresponds to the sex of the
sample, explained ~11.2% of genome-wide differentiation. This signal was clearly driven by
chromosome 1, which showed a similar pattern in which sex explains ~25.1% of variance in the
PCA (Figure 2C). This signal was entirely absent when we did PCA for the putative
"autosomes", chromosomes 2-4, with comparatively very little of the variation explained (PC1:
4%, PC2: 2% Figure 2D) and no correspondence to the sex of the sample. Genealogies based
on chromosome 1 showed a similar result (Figure S2). Additionally, we saw a long, extended
region of differentiation (i.e., elevated F_{ST}) that occurred between female and male sandflies on
chromosome 1 (Figure 2E). This region also shows an extended region of decreased nucleotide

diversity in female sandflies (Figure 2A), but there was no concomitant decrease in nucleotide
diversity within that region in male sandflies (Figure 2B). Given this finding, we compared π
between sexes, with the specific hypothesis that π was lower on chromosome 1 in females than
on autosomes. A linear model indicated that sex ($t = 0.744$, $p = 0.4821$) and the interaction of
sex*chromosome ($t = -0.984$, $p = 0.4821$) were not a significant explanatory variable, but that
chromosome was ($t = 4.260$, $p = 0.381$), and it was driven by lower π in the sex chromosome.

Figure 2. Sex chromosome differentiation in *Lu. longipalpis*. A. π in female sandflies from
Colombia. **B.** π in male sandflies from Colombia. **C.** PCA of only chromosome 1. **D.** PCA of the
autosomes, chromosomes 2-4. **E.** F_{ST} between the sexes. **F.** Manhattan plot of genome-wide
association study between the sexes in Colombian sandflies. **G.** Coverage represented as a

ratio of average male depth divided by average female depth, calculated as a 2,000bp mean.
Two regions in chromosome 1 have been removed that have average log coverage of >30 (with
mean positions at 5847001 and 45239001). **H.** GO biological processes analysis from GWAS
results.

Genome wide association studies revealed autosomal genes associated with sex

Next, we did a GWAS to identify genetic regions associated with sex determination. We
identified 2,641 genic SNPs, and 3,165 intergenic SNPs associated with sex, the vast majority
of which were located in the sex chromosome. The highly differentiated region between sexes in
chromosome 1, the same that shows differentiation by our other analyses, showed a strong
association with sex (Figure 2F). Nonetheless, there were autosomal markers strongly
associated with sex. Sixty-seven SNPs were also associated with significant differences in
genomic coverage between the sexes. Eight genic, autosomal SNPs mapped to three predicted
genes in the three autosomal chromosomes, one of which has no detectable ortholog in any
other dipteran (g5391). A second gene (g7272) showed strong similarity to *mdg3*
retrotransposons in *D. melanogaster* (FBgn0043882). This gene also showed a difference in
genomic coverage between the sexes in our short read sequencing dataset, with a
log2(male:female) coverage ~0.78. The rest of markers showed a uniform distribution of
coverage around 1 (Figure 2G). The third and final gene with genic SNPs associated with sex
was Apoptotic protease-activating factor 1 (*apaf-1*), which plays a role in sex-specific cell
apoptosis and male infertility[34,35]. Twenty-four intergenic SNPs were in the autosomes, and they
are in close proximity to seven genes and three transposable elements. Table S2 lists their
name, location, association *p*-value, and known orthologs in other dipterans. No gene involved
in primary sex determination showed a significant association. Of note, we saw a significant
enrichment for genes involved in regulation of nervous system development (Figure 2H,
GO:0051962: 49.4X; GO:0051960: 32.5X, FDR ≤ 9.6 × 10^{-4}). Table S3 lists other GO categories
with their associated enrichment values.

Population genomics across the continent:
Lutzomyia longipalpis has been proposed to be differentiated into distinct populations
and even separate species across South and Central America[15,36,37]. We used a two-pronged
approach to study the population structure between our Colombian samples and samples of
females from several Brazilian populations[31]. Notably, Sobral was split *a priori* by Labbé et al.[31]
into Sobral 1S and Sobral 2S. First, we used phylogenetic trees. An unrooted genealogy based
on 100kb windows revealed a clustering determined by geographical location consistent with
previous studies[31]. Jacobina and Sobral 1S were each formed by two distinct and unrelated
clusters (Figure 3A-3B). We further examined gene concordance between populations as an
estimate of gene flow. We created a single "population" tree (akin to a species tree) from our
multiple individual sampling of both the whole genome, and of BUSCO genes. When forced to
recapitulate the populations, we found poor gene concordance support for several branches,

indicating poor alignment between individual gene trees and our consensus population tree
(Figure S3). This phylogenetic discordance may occur in instances of recent divergence,
incomplete lineage sorting, hybridization, and poor tree inference which may be occurring in this
population tree[38]. This system may be an amalgamation of all of these factors. In summary, the
two analyses were consistent and suggested significant differentiation between lineages from
different locales, particularly between Colombian and Brazilian samples. Further, we observed
evidence for differentiation between samples from the same locale in Jacobina, Sobral 1S, and
Sobral 2S (Figure 3A) which indicated the existence of differentiated and syntopic lineages.
Second, we used PCA to represent the genetic variation across the continent (Figure 3B). The
first three principal components (PCs) explain 61% of the genetic variance of South American
Lu. longipalpis (Figure S4). PC1 accounted for ~34% of genome-wide variation, and appears to
reflect differentiation between Colombia and Brazil, and between some Brazilian lineages (i.e.,
Lapinha cave; Figure 3C). PC2 and PC3 also capture considerable variance (16.4 and 10.6%
respectively; see Figure S5). PC2 separated Marajo and Sobral 2S from Ricaurte, Sobral 1S,
Laphina with Jacobina samples split between the two groups. Sobral 2S, Marajo, and Ricaurte
are similar in PC3, but Jacobina and Sobral 1S are divided into two separate clusters from each
locality.

284
Figure 3. Partition of genetic diversity in *Lu. longipalpis* across South America. A.
Phylogeny of sampled *Lutzomyia longipalpis* across South America. **B.** *Sampling localities with*
*genomic data included in population genetics analyses. **C.** Principal component analysis of*
*Colombian sandflies from Colombian (this study) and Brazilian sandflies. **D.** Structure plot of*
genotype probabilities across these same samples. Colombian flies divide into two genotypes
driven by sex. All Brazilian sandflies are females, while our Colombian samples also contain
males. Panels A and C have the same legend for sampling locality, D has a different color
schema due to the much larger number of genotypes predicted. Brazilian populations are from
Labbe et al.[31].
294
Next, we measured the extent of gene flow between these genetic clusters. Given the
strong evidence of reproductive isolation between lineages[21,37], we hypothesized reduced—or

even nonexistent—gene flow between the populations we identified. We used *PCAngsd*, which
identifies the most likely number of clusters, but also estimates admixture proportions from allele
frequencies. Genetic variance was best explained by the existence of 10 clusters. Two of these
clusters correspond to male and female *Lu. longipalpis* in Colombia, with little contribution to
other lineages (a probability of <1% all Brazilian samples, except for the Colombian genotype
that corresponds to males, which had a 73.9% probability in one Marajo sample, leading us to
believe it was a male sandfly). This analysis (Figure 3D) also separated Colombian and
Brazilian sandflies. The genetic variation in Brazilian samples was best explained by the
existence of eight groups which correspond largely to the five geographic sites from which they
were collected. The only exceptions were Sobral 1S, Sobral 2S, and Jacobina where we
detected at least two main clusters per locality. The majority of individuals show limited
evidence of admixed ancestry. In Colombia, Laphina, Jacobina, and all but one Marajo sandfly,
admixture proportions are <10% per individual. This is particularly notable in the sites with
syntopic lineages which could potentially have the chance to interbreed and yet do not appear
to (Jacobina, some individuals within Sobral), as evidenced by divergent genotype clusters
driving nearly the entirety of individual genetic makeup (e.g., blue vs teal individuals in Jacobina;
Figure 3D). However, Sobral showed significant admixture within certain individuals (those of
the "2S" lineage, Figure S6). We then proceeded to examine gene flow between populations
using fbranch statistics. These analyses indicated high gene flow from Sobral 2S into Jacobina,
Jacobina into Sobral 1S and Sobral 1S into Laphina (with Sobral "1S" and "2S" being *a priori*
identified as different *Lu. longipalpis* lineages by[31]; Figure S7). Thus, there is either significant
gene flow between populations or our consensus population tree does not adequately reflect
cryptic diversification and potential syntopic lineages. Indeed, the consensus phylogeny of all
individuals indicates the latter, as Sobral 2S individuals cluster with some Jacobina individuals.
Overall, these results confirm that the different lineages of *Lu. longipalpis* are divergent,
indicates advanced divergence towards speciation, and likely the presence of syntopic and
hybridizing lineages.
324
325
326

Figure 4. Genetic diversity in different lineages of *Lu. longipalpis*. A. Histograms of π in
10kb window sizes along the genome. **B.** π in Colombia **C**. π in Jacobina **D.** π in Marajo **E.** π in
Sobral 2S. **F**. π along the genome in Jacobina. **G.** π along the genome in Marajo. Figure 2A and
2B show the π in Colombia. Figure S8 shows π along the genome for the remaining four
populations. **H.** Pairwise comparisons show differences in π among pairs of populations.

The heterozygosity in the whole *L. longipalpis* clade was 0.06 (Figure 4A), but given the
population structure described above, we studied the partitioning of genetic diversity within each
Lu. longipalpis lineage. First, we quantified the extent of heterozygosity in each lineage (Figure
4B-E). Some of the populations are less genetically diverse than others. Colombia shows the
lowest heterozygosity (0.0036); Sobral-1S and Marajo showed the highest heterozygosity
(~0.01) and were three times more diverse than Colombia. In general, the trends of
heterozygosity along the genome were similar across populations (Figures 2A, 2B, 4F, 4G,
Figure S8). All pairwise comparisons were significant (Figure 4H), indicating differences in the
level of genetic diversity across *Lu. longipalpis* lineages. Second, we looked for runs of
homozygosity (ROH), which are identical haplotypes inherited from each parent and thus are a
measure of inbreeding[39]. The total length of ROH was correlated with the number of tracks in an
individual (Figure 5A). Colombian *Lu. longipalpis* had more runs of homozygosity (ROHs) than
any of the Brazilian populations (Figure 5B, C, Figure S9). The pattern is not driven by a single
individual and is a property of the whole population (Figure S10) and extends beyond the mean
level of linkage disequilibrium (Figure S11). Collectively, these results indicate that the genetic
diversity, both at the level of nucleotide and haplotype, of Colombian *Lu. longipalpis* is lower
than that of Brazilian populations. Based on this, we hypothesized that the demographic
trajectory of *Lu. longipalpis* in Colombia was different from that of the populations in Brazil and
thus we examined the trajectory of the effective population size over time of the different
lineages of *Lu. longipalpis*. We found that all Brazilian populations experienced an effective
population size increase on average around 122,410 generations ago (Figure 6). Captive
sandflies have a generation of 6-7 weeks[40], which would suggest that the population increase
occurred between 14,086 and 16,433 years ago (for 6 week or 7 week generations,
respectively). While we found the same trend in all Brazilian populations, the historic effective
population size of the Colombian population shows no evidence of dramatic change. As we
hypothesized, the Colombian and Brazilian populations of *Lu. longipalpis* have undergone
different demographic trajectories which may help explain their current levels of genetic
diversity.

361
362
Figure 5. Runs of homozygosity in *Lu. longipalpis*. **A**. Plot depicting the total number of ROH
regions over 50kb size and their total length by populations. **B**. Size distribution across genomic
regions for Colombian sandflies (binwidth = 500000). **C**. Size distribution across genomic
regions for sandflies from Marajo, Brazil (binwidth = 500000). Figure S9 shows the ROH
distribution per chromosome for other populations. Figure S10 shows ROH histograms by
individual for each population.
369

Figure 6. Demographic history of sandflies indicating temporal patterns of predicted effective population size. We assumed a generation time of six weeks.

DISCUSSION

Phlebotomine sandflies are disease vectors for several human and animal pathogens, and as a result, pose a serious human health threat[41–43]. One of the more notable diseases they transmit is the neglected tropical disease leishmaniasis, and in the Neotropics, one of the most common and important disease vectors of leishmaniasis is *Lutzomyia longipalpis*[44,45]. This sandfly species is widespread across the South American continent[2,14], poorly characterized[15,16], and is thought to be increasingly associated with anthropogenic and urbanized habitats[27]. Despite the clear human health risk posed by these vector species, little is known about their basic biology, hampering control and mitigation efforts[46]. Although there are some genomic resources[31,47], these are limited to Brazilian populations which represent geographically distant populations within a broader cryptic species complex of *Lu. longipalpis*. Here, we present highly continuous genome assemblies for both female and male sandflies from Colombia, which will be a critical resource for future work. We identified signatures of differentiation between the sexes and high divergence across the continent. Further, we discovered that the demographic histories of the different lineages of *Lu. longipalpis sensu lato* are complex, and that genetic

information points to the presence of multiple syntopic lineages. We discuss each of these
findings in the next paragraphs.

Our first finding is that females and males differ in their genomic composition. Although
recent genomic work has indicated no heteromorphy in the Psychodidae[48], prior work identified
the presence of heteromorphic chromosomes in sandflies[49–51]. Here, we sequenced individuals
from both sexes and produced highly continuous assemblies of both sexes. We found that the
longest chromosome in *Lu. longipalpis* is heteromorphic and ~1.3 Mbp longer in male sandflies
than in female sandflies. Our study builds on these previous observations and reveals that there
are differences in TE content and overall genomic composition between the *X* and *Y*
chromosomes. Further, our analysis of repeats within the genome indicates that this difference
in chromosomal length is likely driven by repeat content in male sandflies, which have an extra
~1.3 Mbp of repeats in this chromosome. Our results have implications for the study of sex
chromosome evolution in dipterans. Further, because only females are hematophagous, and
leishmaniasis is known to have negative impacts on longevity and fecundity of sandflies[52,53],
leishmaniasis is likely to be a strong selective force that may impart different selective regimes
on female and male genomes, which may be evident either between the sexes or potentially in
heteromorphic sex chromosomes.

We identified a large, highly differentiated region between the sexes in a heteromorphic
sex chromosome. This region shows a substantial dip in nucleotide diversity in females relative
to males and encompasses a suite of genes related to sex determination, including the
commonly characterized dipteran sex-determining genes *sex-lethal, fruitless* (*fru*)*, doublesex-*
and mab-3-related transcription factor A2 (*dsx*)*, lethal(2)*[54]. Some of these genes, *fru* and *dsx*,
have been implicated in sex determination in other phlebotomine species[55]. There are also a
variety of testes-specific genes (e.g., *ejaculatory bulb-specific protein 3*, *ejaculatory bulb-*
specific protein 3-like, *testis-expressed protein 10 homolog*, *motile sperm domain-containing*
protein). One gene in this region, *takeout,* shows a high degree of structural conservation and
has a conserved role in courtship across insects as diverse as fruit flies, mosquitos, and
moths[56]. Other genes in this region of differentiation may be related to basic ecology and life
history such as finding mates, finding prey, and effective hematophagy. Among these are
several odorant receptor and gustatory receptor genes. Genes in these families play a key role
in finding mates and lekking behavior[57], finding food or blood meals[58–60], and oviposition site
choice[61,62]. In other hematophagous lineages, odorant receptor genes are heavily regulated and
exhibit sex-specific and satiation-specific signatures of expression[63]. In *Lu. longipalpis,*
pheromones play a key role in attracting mates[18,64–66]; these show population-specific
signatures[18] and are perceived by odorant receptor genes which also show population specific
variation [57]. These may play a role in the cryptic *Lu. longipalpis* species complex via mate
choice, and may be a source of prezygotic isolation. Although these roles of odorant receptor
genes are well-known, these genes often have disparate functions among dipterans[67]. In sum,
there are a suite of genes in the genomic region that are highly differentiated between the sexes
that are related to ecology including finding food, finding mates, and oviposition that are
potentially playing important, sex-specific roles.

A complementary result, also related to sex differentiation in *Lutzomyia*, is the finding of
autosomal alleles associated with sex. In addition to these regions in chromosome 1, our
genome wide association study between female and male sandflies in Colombia yielded

significant peaks that were intragenic in three predicted genes, one per chromosome on
chromosomes 2-4. We were unable to identify the gene product on chromosome 2 via Uniprot,
OrthoDB, VectorBase, or NCBI blast. The predicted gene on chromosome 3 has also been
predicted in a previous assembly of *Lu. longipalpis* [31], but remains an uncharacterized protein
product. The intragenic SNPs our GWAS identified as associated with sex differences on
chromosome 4 is Apoptotic protease-activating factor 1 (*apaf-1*). The *C. elegans* homolog of
this gene (*CED-4*) interacts with the *C. elegans* gene *fem-1* which is crucial for male
development[34] and *apaf-1* deficient mice exhibit male infertility due to degeneration of
spermatogonia[35]. The *Drosophila* homolog of *apaf-1* is the gene Apaf-1–related killer (*ark,* or
sometimes *dark* for *Drosophila* Apaf-1–related killer), which is required for sperm formation[68].
The SNPs that we identified were present in intronic regions between predicted exons 3 and 4
in our gene annotation. However, the publicly available RNAseq data that we used for
annotation only contained female sandflies, potentially missing key, sex-specific isoforms of this
gene. We postulate that genomic differentiation between the sexes in this gene is related to sex-
specific differences in gametogenesis, or sex conflict (reviewed in [69]).
Perhaps most notably, our study provides strong evidence of divergence and population
structure across the continent. Our analyses combine novel population genomics data for
Colombian *Lu. longipalpis* populations with publicly-available whole genome sequence data,
which allowed us to carry out the first population genetics study of this critical vector species
across South America. Overall divergence between Brazilian and Colombian sandflies was
high, supportive of previous work suggesting that *Lu. longipalpis* is a cryptic species
complex[15,16,36,37]. In addition, we found evidence that Colombian *Lu. longipalpis* shows lower
genetic variability than all of the Brazilian populations. This result manifested in two different
facets, lower nucleotide diversity and a substantial increase in ROH tracts. ROH tracts were
more common in the *X*-chromosome in all populations but were particularly noticeable in
Colombia. It is unclear why Colombian flies exhibit such dramatic patterns of low diversity,
particularly because it does not appear to be an isolated population. A potential explanation is
reduced vagility in the Colombian lineage. For example if Colombian sandflies roam less during
their lifetime than Brazilian flies, they are more likely to experience matings with close relatives
and thus show more signs of inbreeding. An additional possibility is that Colombian *Lu.*
longipalpis exhibit facultative parthenogenesis, a trait seen in several dipteran species[70],
including in *Lu. mamedei*[71]. Both of these possibilities remain highly speculative and a
systematic survey of the reproductive biology of the different lineages of *Lu. longipalpis* is
needed to understand the patterns we observed. Our demographic analyses concur with the
inference of low genetic diversity in Colombia and indicate that all populations had a similar *Ne*
between 16,000 and 14,000 years ago depending on the generation time in the wild; Brazilian
populations underwent a large population expansion, Colombian *Lu. longipalpis* did not. This
timing is similar to the arrival and expansion of humans in South America[72]. Given that *Lu.*
longipalpis is one of the most anthropophilic species, population expansion in Brazilian sandflies
may have occurred as a result of human colonization and expansion.
473
Colombian sandflies (from our single sampling locality) showed structure only between
female and male sandflies, whereas multiple Brazilian populations (Jacobina, Sobral) were
highly structured in all our analyses. Together, these results indicate the presence of multiple

syntopic lineages—a result consistent with the previously hypothesized presence of cryptic
species within *Lu. longipalpis*, *sensu lato. Lutzomyia longipalpis* has been proposed to
encompass different chemotypes[64], and in some instances the proposals have been more
ambitious and have proposed the existence of a species complex which encompasses different
species with stark differences in mating behavior (e.g., mating song, pheromones and lekking
behavior). Notably, and despite the differentiation we describe for these lineages, we find some
evidence of admixture between lineages. Hybridization seems to be common in *Lutzomyia*
(reviewed in[73]), which poses the question of how frequently differentiating lineages exchange
alleles. Here we provide evidence for the existence of genome-wide admixture within *Lu.*
longipalpis. While this introgression does not appear sufficient to erode the genetic
differentiation between lineages, even when they are syntopic, this allele sharing might serve as
fodder for adaptation[74,75]. The importance of introgression has been previously hypothesized
across the *Lutzomyia* genus[76–78], and among *Lu. longipalpis* lineages[79,80]; our results represent
a formal genome-wide test of the occurrence of this phenomenon in this genus of disease
vectors.
Our manuscript leverages the power of whole genome sequencing to understand the
evolutionary processes that govern genetic variation in the critical leishmaniasis vector *Lu.*
longipalpis. Our results incorporate a genomic perspective into the study of differentiation within
Lu. longipalpis and represent a resource to not only understand sex chromosome evolution in
disease vectors, but also open the door to understanding the molecular and evolutionary
mechanisms of the interactions between the vectors, the *Leishmania* parasites, and the hosts
affected by the disease. Now that we have a better snapshot of the extent of genetic variation in
the vector, the next frontier is understanding whether the evolutionary interactions between
sandfly vectors and the *Leishmania* pathogen can provide the key to mitigating the spread of
leishmaniasis and the human health threat it poses.
502
503 **MATERIALS AND METHODS**
504 *Permits and ethics*. All sandflies collections were performed following the guidelines of the
505 Colombian decree No 001566 and 002662. Insects were collected under the permits 001566
506 and 002662 of the Autoridad de Licencias Ambientales. Sandflies were collected on private
507 property and permission was received from landowners before sampling.
508
Sample collection. Samples were collected from a semi-forested location outside of Ricaurte,
Cundinamarca, Colombia in November 2021 and October 2022**.** Sandflies were collected via
mechanical aspiration with a mouth aspirator and placed into vials to temporarily hold the
insects until processing. Prior to processing, vials containing the sandflies were put in a cooler
with ice to immobilize the specimens. Vials were emptied into a Petri dish where the sandflies
were selected and classified by species under a Leica dissecting scope. To identify sandfly
species, we used taxonomic keys[81,82]. Individual sandflies were removed with tweezers and
placed into individual 1.5mL tubes either in liquid nitrogen for genome assembly in 2022 or in
RNA-stabilizing buffer (RNALater[TM]; Ambion, USA) for whole genome resequencing in 2021.
Samples were exported to North Carolina, USA for sequencing.
519

Genome sequencing. Due to their small size (thorax length in females: mean = 0.052 ± 0.0025
mm; males: mean = 0.0483 ± 0.0027 mm;[83] individuals were pooled prior to genomic DNA
extraction for whole genome sequencing (females = 9, males = 10). We used two approaches to
obtain reads for the genome assemblies. First, we generated PacBio HiFi reads. We
homogenized sandflies with a pestle and incubated them at 55℃ for 30 minutes prior to
performing a standard phenol chloroform extraction. Extracted high molecular weight gDNA was
sent to the Genomics Core Facility at the Icahn School of Medicine at Mount Sinai, where
libraries were prepared using PacBio's ultra-low DNA input HiFi library prep which utilizes PCR
amplification of genomic material. Libraries were pooled and sequenced across two SMRT cells
on a Pacific Biosciences Sequel IIe to a coverage of 71.8x and 75.7x coverage for females and
males.
531   Second, we used chromatin configuration. We sent 25 female and 25 male *Lu.*
longipalpis in ethanol to Arima Genomics for DNA extraction and sequencing. Hi-C data was
generated using the Arima HiC kit and the Arima Library Prep Module according to the Arima
Genomics manufacturer's protocols. Briefly, Arima High Coverage Hi-C is designed to capture
the sequence and 3D conformation of genomes. Chromatin from tissues, cell lines, or blood is
first crosslinked to preserve genomic structure, then digested with a restriction enzyme cocktail
optimized for uniform coverage. The digested ends are labeled with biotinylated nucleotides
during a fill-in step and subsequently ligated to capture spatially proximal DNA fragments.
These proximally-ligated DNA fragments are purified, fragmented, and enriched for biotin-
labeled sequences.The female sample was sequenced to 189.5 million read pairs and the male
sample to 180 million read pairs on an Illumina NovaSeq on 'paired-end' mode.
542
Genome assemblies. We produced separate genome assemblies for the female and male
sandflies. While we initially attempted to use HiFiasm[84], we discovered that haplotype diversity
in our pooled data exceeded the allowed diversity in the assembly algorithm. Therefore, we
chose to use wtdbg2[85], an assembler with an algorithm that allows for greater diversity and
collapses haplotypes into a single, haploid assembly. We used the "wtpoa-cns" command in
wtdbg2 to polish the assembly.
549   We then scaffolded our initial draft genome assemblies with our Hi-C data. Splits were
calculated using Juicer version 1.6[86] and reads were mapped to the draft assemblies with BWA
version 0.7.17[87]. We then ran Samtools version 1.17 to sort the aligned reads and Picard's
"MarkDuplicates" version 2.27.5 to mark duplicates[88]. We indexed our initial contig assembly
using Samtools "faidx" prior to running the YaHS (Yet Another Hi-C Scaffolding tool) pipeline[89]
with a MAPQ score of 30. We used Juicer "pre" to create a HiC contact map file which we
imported into Juicebox Assembly Tools[90] to manually curate each assembly. Curation included
fixing one major translocation (female assembly) and splitting a chromosome that had an
incorrect telomere-to-telomere join (male assembly). We then used Juicer "post" to update the
manually curated final assemblies for females and males.
559   After examination of whole genome sequence results, we identified systemic and
dramatic differentiation between female and male sequences along the majority of chromosome
1. We were unable to initially phase a ZW or XY assembly using our assembly approaches,
likely due to the pooled nature of our data (i.e., because male sandflies were too small for
effective single-fly sequencing, and we were forced to pool multiple individuals). As a result, we

took a secondary approach to assemble what we predicted to be a Y chromosome in males
using the software SRY[91] to identify reads that are likely to be from sex-limited chromosomes.
We used female and male Illumina whole genome sequence reads to produce sex-specific
kmers. We then mapped these kmers to the raw male-only PacBio HiFi reads. We then
extracted those reads that were highlighted by presumptive sex-limited kmers from our Illumina
data and used wtdbg2[85] to assemble a Y chromosome from those HiFi reads. We then used
RagTag[92] to scaffold the male Y chromosome using the male X chromosome (chromosome 1)
as our backbone. We aligned this to the male "X" chromosome using minimap2[93] assuming
~0.1% sequence divergence ("-cx asm5") and visualized the alignment using "*pafr*" in R [94].
After producing our scaffolded assemblies, we identified repeat elements in the
assemblies. We used Repeat Modeler version 2.0.4[95] to model repeat elements within each
assembly. Most of these repeat elements were unidentified by Repeat Modeler, so we ran
Transposon Classifier RFS[96] to classify these repeat regions. We then created a dipteran and
Lu. longipalpis specific database of repeat elements. We combined the curated DFAM database
version 3.4[97] with the RepBase version 25.05 within RepeatMasker version 4.1.2-p1[98]. We used
the function "famdb.py" to extract all dipteran sequences from these databases, then added our
sex-specific sandfly repeat databases. We then identified and soft-masked repeats throughout
the genome using RepeatMasker version 4.1.2-p1[98] and specified our curated vertebrate and
Lu. longipalpis repeat library as the library of repeats.
We then annotated assemblies using Braker version 3.0.8[99–102]. We used a combination
of protein evidence and RNA seq evidence to annotate putative protein coding regions of the
genome. We used the OrthoDB v11 database[103] and the VectorBase 67 database[104,105] as input
protein data for annotation. Protein sequences were aligned to the genomes using miniprot[106].
We used the SRA toolkit[107] to download publicly-available RNA seq data from *Lu. longipalpis* to
annotate the genome. We downloaded samples with accession numbers SRR535756-
SRR535778. RNA seq reads were mapped to our genomes using HISAT2[108] and
sorted/manipulated using samtools[109]. We followed this with Braker3, utilizing the GeneMark-
ETP pipeline[110] to predict protein coding genes from RNAseq evidence. This was then
compared to the protein evidence, OrthoDB and VectorBase in this case, to find homologous
proteins that were then mapped back to the genome with ProtHint[101]. Augustus[111,112] then
predicted a second set of genes. We additionally used the diptera_odb10 database[113] in
Augustus to predict potentially missing protein products. We then used TSEBRA[114] to merge
gene predictions from Augustus and the GeneMark-ETP pipelines.
597
WGS resequencing. We resequenced a total of 63 *Lu. longipalpis* from Ricaurte,
Cundinamarca*,* Colombia. Of these, 37 sandflies were female and 26 were males sample
information in Table S1). Briefly, we conducted a homogenization and incubation similar to that
of the WGS gDNA extraction. Following the incubation, we followed a modified version of the
instructions on the Zymo Genomic DNA Clean and Concentrator kit, and quantified eluted DNA
using a Qubit. Our full extraction protocol can be found in the supplemental materials. Library
prep and sequencing were conducted at the University of North Carolina-Chapel Hill's High-
Throughput Sequencing Facility. Libraries were prepped using a ThruPLEX method for low DNA
concentrations (CHiP-seq). Following library preparation, pooled paired-end 150bp reads were
sequenced on an Illumina NovaSeq 6000 SP.

WGS mapping, variant calling, and diversity summary statistics. Next we measure the extent of
genetic diversity and differentiation within *Lu. longipalpis*. We followed the GATK best practices
for variant calling[115]. We used BBMap version 38.87 (https://github.com/BioInfoTools/BBMap) to
trim adapters from whole genome resequencing data and then mapped reads to the female
reference genome using BWA version 0.7.17[87]. Read groups were added using GATK's
AddOrReplaceReadGroups. For samples sequenced across two lanes, aligned data were then
merged using the "merge" tool in samtools version 1.14[109]. Aligned data were cleaned with
Picard's CleanSam, sorted with SortSam, and duplicates marked with MarkDuplicates (Picard
version 2.23.4[88]. Genotypes were then called with GATK HaplotypeCaller, before joint
genotyping data with GenotypeGVCFs. All GATK functions were from GATK version 4.1.9.0.
We merged bam files for each sample that were sequenced on multiple lanes using
samtools version 1.14 merge function[109]. We generated a Beagle file of genotype probabilities
using *angsd* version: 0.921-11-g20b0655, and then calculated principal components using the
angsd script "pcangsd.py"[116]. We then extracted eigenvalues and calculated the percent of the
variance in our data explained by each PC axis, which we then plotted in R version 4.1.1 ("Kick
Things") using ggplot2[117]. In addition to running this PCA pipeline with whole genome data, we
subsetted the alignment bam file in order to separately examine differentiation between the
presumptive sex chromosome (chromosome 1) and the autosomes (chromosomes 2-4) using
the same approach.
We examined basic population statistics (nucleotide diversity, π, and population
divergence [F_{ST}, Dxy]) using pixy[118], which accounts for invariant sites. We calculated pairwise
differences in nucleotide diversity and Weir-Cockerham F_{ST} values[119]. We were particularly
interested in potential differentiation between female and male sandflies, given that only females
are hematophagous and thus only females are vectors of leishmaniasis. Per pixy's instructions,
we first created a vcf file of invariant sites using the "-all-sites" flag in GATK's GenotypeGVCFs.
We then filtered the corresponding gvcf file to remove sites with a high proportion of missing
data (--max-missing 0.9), low coverage (--min-meanDP 3), and high overall depth (--max-
meanDP 40) using vcftools 0.1.15[120]. We then used pixy to calculate F_{ST} and Dxy across
1,000bp windows between female and male sandflies. The window size was chosen five the
extent of LD in *Lu. longipalpis*.) . Additionally, we subset our gvcf file into a female-only and a
male-only gvcf which we used to calculate nucleotide diversity (π) across 1,000bp windows in
female and male sandflies. We used PLINK 1.90b6.21[121] to calculate linkage disequilibrium. We
additionally calculated Tajima's D in 1,000 bp windows using VCFtools.
We detected runs of homozygosity (ROH) using PLINK 1.90b6.21[121] with the
parameters: --allow-extra-chr and --homozyg-kb 5. This command estimates the run length and
identifies contiguous segments of homozygosity greater than 5 kbp by assessing the observed
genotype data against expected patterns of homozygosity. ROH was detected separately for
each of Colombia and the five Brazil populations. With a 5 kbp minimum size, we identified a
minimum of 100 SNPs per region and on average there were 189-276 SNPs per region. We
repeated the same analyses using different parameter values for the minimum length of ROH in
kbp (5, 10, 25, 50, 100), and results were similar despite fewer ROH regions with higher kb
thresholds.

Phylogenetic trees. We used trees to address two questions. First, to examine the genealogy of
the largest chromosome relative to the presumptive autosomes in placement of females and
males in relation to each other, and second, to examine continent-wide evolutionary
relationships. We used the python script "vcf2phylp.py"[122] to split our vcf files into 100,000 bp
regions and convert them into the phylip format.
To examine the genealogy of the largest chromosome relative to the presumptive
autosomes, we created two separate trees using 1) only the largest chromosome or 2) the
presumptive autosomes in just Colombian sandflies. For both trees, we made a phylogenetic
tree for each window using IQ-TREE[123] version 2.2.2.7 with DNA specified ("-st DNA") and 1000
bootstraps ("-bb 1000"). We merged the resulting consensus tree files into a single tree and
used ASTRAL[124] version 5.7.8 to create a single genome-wide consensus tree. We additionally
created trees solely from chromosome 1 and the autosomes.
To examine evolutionary relationships across the continent, we made a single genome-
wide tree using the above methods. The approach for this question is identical to the one
described immediately above, with the exception that we created a single tree including all
chromosomes. Further, we examined gene concordance between the genome wide tree and
conserved, core dipteran genes from the BUSCO diptera_odb10 database[113]. To do this, we
used ASTRAL[124] version 5.7.8 to create a single genome-wide consensus tree that is
constrained to the populations sampled. We additionally extracted all the complete, single copy
dipteran BUSCO genes assembled in our male assembly, using the python script
"vcf2phylp.py"[122] to convert the vcf of these genomic regions into the phylip format. We created
single gene trees as per the windows above using IQ-TREE, and created a single consensus
population-level tree using ASTRAL as we did for our whole genome tree. We then used IQ-
TREE[123] version 2.2.2.7 to calculate gene concordance between each gene tree and the overall
consensus tree. Finally, we examined gene flow and the potential for introgression using
fbranch statistics in Dsuite [125]. Briefly, we used our consensus population tree and our variant
call file as input to examine fbranch statistics using "Dtrios" to examine all trios and "Fbranch" to
examine the output fbranch statistics. Given the clear divergence between Brazilian and
Colombian sandflies, we used Colombia as the outgroup in our analysis and focused on
Brazilian sandflies.
682
Population genomics of geographically disparate Lu. longipalpis *populations.* *Lutzomyia*
longipalpis has a broad geographic range from Argentina through Mexico[2,14]. Further, multilocus
sequence typing and phenotypic surveys have revealed strong population structure which might
correspond to cryptic speciation[17,18] and divergences in courtship song[20], morphology,[21] and
pheromones[18]. Recent work sequenced Brazilian *Lu. longipalpis* from four localities in Brazil[31].
Here we examined basic population genomic metrics between Colombian and Brazilian *Lu.*
longipalpis. We were particularly interested in differentiation between populations of putative *Lu.*
longipalpis, and as a result focus on metrics such as F_{ST} and Dxy between populations.
Out of the 68 Colombian samples, 63 sandflies (37 females and 26 males) passed our
quality metrics for population level analyses (>50% of reads mapping to our *Lu. longipalpis*
female assembly and > 5X average coverage). On average, 91.4% of female reads and 84.4%
of male reads mapped to the female assembly. Average coverage was 16.8x and 13.2x for
females and males respectively. When mapped to the male assembly, 84.3% of female reads

and 79.6% of male reads mapped and we had 14.6x and 11.7x coverage for females and males
respectively.
Additionally, we downloaded whole genome resequence data from[31] using the SRA
toolkit[107]. Table S1 lists the accession numbers of the previously published reads used in this
study. We then followed the same protocol that we used for our Colombian samples, the GATK
best-practices for variant calling [115]. We used BBMap version 38.87
(https://github.com/BioInfoTools/BBMap) to trim adaptors from whole genome resequencing
data and then mapped reads to the female reference genome using BWA version 0.7.17[87].
Read groups were added using GATK's AddOrReplaceReadGroups. For samples sequenced
across two lanes, aligned data were then merged using the "merge" tool in samtools version
1.14[109]. Aligned data were cleaned with Picard's CleanSam, sorted with SortSam, and
duplicates marked with MarkDuplicates (Picard version 2.23.4[88]. Genotypes were then called
with GATK HaplotypeCaller, before joint genotyping data with GenotypeGVCFs. All GATK
functions were from GATK version 4.1.9.0.
For samples across the continent of South America, we examined summary population
statistics (π, F_{st}, and Dxy) as we did for samples within Colombia (see above). We filtered our
gvcf file to remove sites with a high proportion of missing data (--max-missing 0.9), low
coverage (--min-meanDP 3), and high overall depth (--max-meanDP 40) using vcftools
0.1.15[120]. We then used pixy to calculate F_{ST} and Dxy across 1,000bp windows between
populations. Additionally, we subset our gvcf file into a female-only and a male-only gvcf which
we used to calculate nucleotide diversity (π) across 1,000bp windows in female and male
sandflies. We additionally calculated Tajima's D in 1,000 bp windows using VCFtools.
In order to examine population structure across the continent, we used angsd[116] to
examine the data with a PCA, structure, and *ADMIXTURE* based approaches. We used a
similar approach to the within-Colombia examinations. We generated a Beagle file of genotype
probabilities using *angsd* version: 0.921-11-g20b0655, and then calculated principal
components and genotype probabilities using the *angsd* script "pcangsd.py"[116]. This program
calculates ancestry based on individual allele frequency and uses a similar approach to delta K
in that it permutes iteratively across a range of ancestry possibilities. We then extracted
eigenvalues and calculated the percent variance in our data explained and plotted both this
PCA and a STRUCTURE plot in R version 4.1.1 ("Kick Things") using ggplot2[117].
To quantify if sex was responsible for differences in genome size, we used a linear
model (function "lm" in R) with average π per window in the sexes as the outcome and sex,
chromosome, and the interaction of sex and chromosome as fixed effects.
We estimated demographic histories using MSMC2[126] with one thread and ten time
segments of different lengths using the parameters '-t 1 -p 1-6+8-1+1-4'. MSMC2 was run
separately for individual samples since we had unphased genotypes. The MSCM2 input files
were generated using msmc-tools[127]. Separate input files were generated for each of the four
major chromosomes using 'generate_multihetsep.py' and then concatenated together into
separate genomic input files for each individual sample. 'generate_multihetsep.py' requires two
input files: a mask of sites with coverage for that individual and a genomic mappability mask.
The covered site masks were created in three steps run separately for each chromosome and
each individual sample using Samtools 1.21[128], BCFtools[128] and msmc-tools. The genomic
mappability mask was created using genmap[129]. MSMC2 results were plotted using R [130] using

a mutation rate of μ=1.25e-8 and an estimated generation time of 49 days[40] with time in years
calculated as 'time_boundary / μ / gen'. The effective population size was calculated as (1 / λ) /
(2 * μ) where lambda is the coalescence rate returned by MSMC2.
743
Genome wide association study for sex determination. Finally, we ran a genome-wide
association study (GWAS) for 133 samples from both Colombia and Brazil using sex as the
phenotype (113 females, 20 males) to identify genetic regions associated with sex
determination. First, we created the required plink input files from our vcf by filtering for just the
four chromosomes and using a minor allele frequency of 0.05. To run the GWAS, we used plink
2.0[121] using a generalized linear model framework with sex as a binary variable[131], and we
filtered out SNPs with 20% or more missing data (--geno 0.2) and a minor allele frequency of
0.05 (--maf 0.05). Results are presented from the continent wide analysis, but results are the
same using just the Colombian samples yielded qualitatively identical results.
753
754   **Data availability:**
755   All raw sequence data as well as genome assemblies are available on NCBI under BioProject
756   PRJNA1110158. Code can be found at https://github.com/AdamStuckert/Lu_longipalpis; and
757   will be deposited in Zenodo upon acceptance.
758
Author contributions:
AMMS: Conceived project, collected samples, conducted wet lab work, analyzed data, wrote
manuscript
DT: Analyzed data, wrote manuscript
ERRD: Conducted wet lab work, wrote manuscript
DP: Conducted wet lab work, wrote manuscript
MP: Conducted wet lab work, wrote manuscript
CM: Conceived project
JAR: Wrote manuscript
EB: Conceived project, collected samples
JRW: Conducted wet lab work, wrote manuscript
RV: Conceived project, collected samples, identified sandflies, wrote manuscript
DRM: Conceived project, collected samples, analyzed data, wrote manuscript
772
773   **Competing interests:**
774   The authors declare they have no competing interests.
775
776   **Acknowledgments:**
777   The research presented here was supported by the National Institute of General Medical
778   Sciences under award R35GM148244.

779
780 **References:**
781

1. Maroli, M., Feliciangeli, M. D., Bichaud, L., Charrel, R. N. & Gradoni, L. Phlebotomine

sandflies and the spreading of leishmaniases and other diseases of public health concern.

Med. Vet. Entomol. **27**, 123–147 (2013).

2. Young, D. G. & Duncan, M. M. Guide to the identification and geographic distribution of

Lutzomyia sand flies in Mexico, the West Indies, Central and South America (Diptera:

Psychodidae). *Memoirs of the American Entomological Institute* **54**, 1–887 (1994).

3. World Health Organization. Control of the leishmaniases. *World Health Organ. Tech. Rep.*

Ser. xii–xiii, 1–186, back cover (2010).

4. Alvar, J. *et al.* Leishmaniasis Worldwide and Global Estimates of Its Incidence. *PLoS One*

7, e35671 (2012).

5. Karimkhani, C. *et al.* Global burden of cutaneous leishmaniasis: a cross-sectional analysis

from the Global Burden of Disease Study 2013. *Lancet Infect. Dis.* **16**, 584–591 (2016).

6. Bern, C., Maguire, J. H. & Alvar, J. Complexities of assessing the disease burden

attributable to leishmaniasis. *PLoS Negl. Trop. Dis.* **2**, e313 (2008).

7. McIlwee, B. E., Weis, S. E. & Hosler, G. A. Incidence of Endemic Human Cutaneous

Leishmaniasis in the United States. *JAMA Dermatol.* **154**, 1032–1039 (2018).

8. Curtin, J. M. & Aronson, N. E. Leishmaniasis in the United States: Emerging issues in a

region of low endemicity. *Microorganisms* **9**, 578 (2021).

9. Gin, T. E., Lashnits, E., Wilson, J. M., Breitschwerdt, E. B. & Qurollo, B. Demographics and

travel history of imported and autochthonous cases of leishmaniosis in dogs in the United

States and Canada, 2006 to 2019. *J. Vet. Intern. Med.* **35**, 954–964 (2021).

10. González, C. *et al.* Climate change and risk of leishmaniasis in North America: predictions

from ecological niche models of vector and reservoir species. *PLoS Negl. Trop. Dis.* **4**,

e585 (2010).

11. Clarke, C. F., Bradley, K. K., Wright, J. H. & Glowicz, J. Case report: Emergence of autochthonous cutaneous leishmaniasis in northeastern Texas and southeastern Oklahoma. *Am. J. Trop. Med. Hyg.* **88**, 157–161 (2013).

12. Montenegro Quiñonez, C. A., Runge-Ranzinger, S., Rahman, K. M. & Horstick, O. Effectiveness of vector control methods for the control of cutaneous and visceral leishmaniasis: A meta-review. *PLoS Negl. Trop. Dis.* **15**, e0009309 (2021).

13. Wilson, A. L. *et al.* The importance of vector control for the control and elimination of vector-borne diseases. *PLoS Negl. Trop. Dis.* **14**, e0007831 (2020).

14. Soares, R. P. P. & Turco, S. J. *Lutzomyia longipalpis* (Diptera: Psychodidae: Phlebotominae): a review. *An. Acad. Bras. Cienc.* **75**, 301–330 (2003).

15. Bauzer, L. G. S. R., Souza, N. A., Maingon, R. D. C. & Peixoto, A. A. *Lutzomyia longipalpis* in Brazil: a complex or a single species? A mini-review. *Mem. Inst. Oswaldo Cruz* **102**, 1–12 (2007).

16. Maingon, R. D. C., Ward, R. D., Hamilton, J. G. C., Bauzer, L. G. S. R. & Peixoto, A. A. The *Lutzomyia longipalpis* species complex: does population sub-structure matter to Leishmania transmission? *Trends Parasitol.* **24**, 12–17 (2008).

17. Soto, S. I., Lehmann, T., Rowton, E. D., Vélez B, I. D. & Porter, C. H. Speciation and population structure in the morphospecies *Lutzomyia longipalpis* (Lutz & Neiva) as derived from the mitochondrial *ND4* gene. *Mol. Phylogenet. Evol.* **18**, 84–93 (2001).

18. Maingon, R. D. C. *et al.* Genetic identification of two sibling species of *Lutzomyia longipalpis* (Diptera: Psychodidae) that produce distinct male sex pheromones in Sobral, Ceará State, Brazil. *Mol. Ecol.* **12**, 1879–1894 (2003).

19. Gutierrez, M. A. C. *et al.* DNA barcoding of *Lutzomyia longipalpis* species complex (Diptera: Psychodidae), suggests the existence of 8 candidate species. *Acta Trop.* **221**, 105983 (2021).

20. Souza, N. A. *et al.* Analysis of the copulatory courtship songs of *Lutzomyia longipalpis* in six

populations from Brazil. *J. Med. Entomol.* **41**, 906–913 (2004).

21. Ward RD, Ribeiro AL, Ready PD, Murtagh A. Reproductive isolation between different

forms of *Lutzomyia longipalpis* (Lutz & Neiva),(Diptera: Psychodidae), the vector of

Leishmania donovani chagasi Cunha & Chagas and its significance to kala-azar distribution

in South America. *Memórias do Instituto Oswaldo Cruz* **78**, 269–280 (1983).

22. Mangabeira, O. Sobre a sistemática e biologia dos *Phlebotomus* do Ceará. *Rev. Bras.*

Malariol. Doencas Trop. **21**, 3–26 (1969).

23. Lainson, R. & Rangel, E. F. *Lutzomyia longipalpis* and the eco-epidemiology of American

visceral leishmaniasis, with particular reference to Brazil: a review. *Mem. Inst. Oswaldo*

Cruz **100**, 811–827 (2005).

24. Ferro, C., Pardo, R., Torres, M. & Morrison, A. C. Larval microhabitats of *Lutzomyia*

longipalpis (Diptera: Psychodidae) in an endemic focus of visceral leishmaniasis in

Colombia. *J. Med. Entomol.* **34**, 719–728 (1997).

25. Ferro, C. *et al.* Spatial Distribution of Sand Fly Vectors and Eco-Epidemiology of Cutaneous

Leishmaniasis Transmission in Colombia. *PLoS One* **10**, e0139391 (2015).

26. Costa, P. L. *et al.* Ecology of Lutzomyia longipalpis in an area of visceral leishmaniasis

transmission in north-eastern Brazil. *Acta Trop.* **126**, 99–102 (2013).

27. Salomón, O. D., Feliciangeli, M. D., Quintana, M. G., Afonso, M. M. dos S. & Rangel, E. F.

Lutzomyia longipalpis urbanisation and control. *Mem. Inst. Oswaldo Cruz* **110**, 831–846

(2015).

28. Berrozpe, P. E. *et al.* Spatiotemporal dynamics of *Lutzomyia longipalpis* and macro-habitat

characterization using satellite images in a leishmaniasis-endemic city in Argentina. *Med.*

Vet. Entomol. **33**, 89–98 (2019).

29. Morrison, A. C., Ferro, C., Morales, A., Tesh, R. B. & Wilson, M. L. Dispersal of the sand fly

Lutzomyia longipalpis (Diptera: Psychodidae) at an endemic focus of visceral leishmaniasis

in Colombia. *J. Med. Entomol.* **30**, 427–435 (1993).

30.   Afonso, M. M. D. S., Duarte, R., Miranda, J. C., Caranha, L. & Rangel, E. F. Studies on the

Feeding Habits of *Lutzomyia (Lutzomyia) longipalpis* (Lutz & Neiva, 1912) (Diptera:

Psychodidae: Phlebotominae) Populations from Endemic Areas of American Visceral

Leishmaniasis in Northeastern Brazil. *J. Trop. Med.* **2012**, 858657 (2012).

31.   Labbé, F. *et al.* Genomic analysis of two phlebotomine sand fly vectors of *Leishmania* from

the New and Old World. *PLoS Negl. Trop. Dis.* **17**, e0010862 (2023).

32.   Srikulnath, K., Ahmad, S. F., Singchat, W. & Panthum, T. Do Ty3/Gypsy transposable

elements play preferential roles in sex chromosome differentiation? *Life (Basel)* **12**, 522

(2022).

33.   Zhou, Q. *et al.* The epigenome of evolving *Drosophila* neo-sex chromosomes: dosage

compensation and heterochromatin formation. *PLoS Biol.* **11**, e1001711 (2013).

34.   Chan, S. L., Yee, K. S., Tan, K. M. & Yu, V. C. The *Caenorhabditis elegans* sex

determination protein FEM-1 is a CED-3 substrate that associates with CED-4 and

mediates apoptosis in mammalian cells. *J. Biol. Chem.* **275**, 17925–17928 (2000).

35.   Honarpour, N. *et al.* Adult Apaf-1-deficient mice exhibit male infertility. *Dev. Biol.* **218**, 248–

258 (2000).

36.   Uribe, S. The status of the *Lutzomyia longipalpis* species complex and possible implications

for Leishmania transmission. *Mem. Inst. Oswaldo Cruz* **94**, 729–734 (1999).

37.   Souza, N. A., Brazil, R. P. & Araki, A. S. The current status of the *Lutzomyia longipalpis*

(Diptera: Psychodidae: Phlebotominae) species complex. *Mem. Inst. Oswaldo Cruz* **112**,

161–174 (2017).

38.   Morales-Briones, D. F. *et al.* Disentangling sources of gene tree discordance in

phylogenomic data sets: Testing ancient hybridizations in Amaranthaceae s.L. *Syst. Biol.*

70, 219–235 (2021).

39.   Ceballos, F. C., Hazelhurst, S. & Ramsay, M. Assessing runs of Homozygosity: a

comparison of SNP Array and whole genome sequence low coverage data. *BMC*

Genomics **19**, 106 (2018).

40. Killick-Kendrick, R., Leaney, A. J. & Ready, P. D. The establishment, maintenance and

productivity of a laboratory colony of *Lutzomyia longipalpis* (Diptera: Psychodidae). *J. Med.*

Entomol. **13**, 429–440 (1977).

41. Killick-Kendrick, R. Phlebotomine vectors of the leishmaniases: a review. *Med. Vet.*

Entomol. **4**, 1–24 (1990).

42. Killick-Kendrick, R. The biology and control of phlebotomine sand flies. *Clin. Dermatol.* **17**,

279–289 (1999).

43. Ready, P. D. Biology of phlebotomine sand flies as vectors of disease agents. *Annu. Rev.*

Entomol. **58**, 227–250 (2013).

44. Salomon, O. D. *Lutzomyia longipalpis*, Gone with the Wind and Other Variables. *Neotrop.*

Entomol. **50**, 161–171 (2021).

45. Sousa-Paula, L. C. de, Otranto, D. & Dantas-Torres, F. *Lutzomyia longipalpis* (Sand Fly).

Trends Parasitol. **36**, 796–797 (2020).

46. Rêgo, F. & Soares, R. Lutzomyia longipalpis: an update on this sand fly vector. *An. Acad.*

Bras. Cienc. **93**, e20200254 (2021).

47. Huang, M. *et al.* Improved high quality sand fly assemblies enabled by ultra low input long

read sequencing. *Sci. Data* **11**, 918 (2024).

48. Vicoso, B. & Bachtrog, D. Numerous transitions of sex chromosomes in Diptera. *PLoS Biol.*

13, e1002078 (2015).

49. Kreutzer, R. D., Modi, G. B., Tesh, R. B. & Young, D. G. Brain cell karyotypes of six species

of New and Old World sand flies (Diptera: Psychodidae). *J. Med. Entomol.* **24**, 609–612

(1987).

50. Jiménez, M. E., Bello, F. J., Ferro, C. & Cárdenas, E. Brain cell karyotype of the

phlebotomine sand fly *Lutzomyia shannoni* (Dyar) (Diptera: Psychodidae). *Mem. Inst.*

909  *Oswaldo Cruz* **96**, 379–380 (2001).

910 51. Vigoder, F. M., Araripe, L. O. & Carvalho, A. B. Identification of the sex chromosome

911   system in a sand fly species, *Lutzomyia longipalpis* s.l. *G3* **11**, (2021).

912 52. El Sawaf, B. M., El Sattar, S. A., Shehata, M. G., Lane, R. P. & Morsy, T. A. Reduced

913   longevity and fecundity in Leishmania-infected sand flies. *Am. J. Trop. Med. Hyg.* **51**, 767–

914   770 (1994).

915 53. Agrela, I. F. & Feliciangeli, M. D. Effect of *Leishmania* spp infection on the survival, life

916   expectancy, fecundity and fertility of Lutzomyia longipalpiss.l. and Lutzomyia

917   pseudolongipalpis. *Mem. Inst. Oswaldo Cruz* **110**, 611–617 (2015).

918 54. Ashmore, J. S., Slippers, B., Duong, T. A. & Dittrich-Schröder, G. Understanding the

919   genetics of sex determination in insects and its relevance to genetic pest management.

920   *Insect Mol. Biol.* (2024) doi:10.1111/imb.12982.

921 55. Petrella, V. *et al.* Identification of sex determination genes and their evolution in

922   Phlebotominae sand flies (Diptera, Nematocera). *BMC Genomics* **20**, 522 (2019).

923 56. Saurabh, S., Vanaphan, N., Wen, W. & Dauwalder, B. High functional conservation of

924   takeout family members in a courtship model system. *PLoS One* **13**, e0204615 (2018).

925 57. Hickner, P. V. *et al.* Molecular signatures of sexual communication in the phlebotomine

926   sand flies. *PLoS Negl. Trop. Dis.* **14**, e0008967 (2020).

927 58. Takken, W. & Knols, B. G. Odor-mediated behavior of Afrotropical malaria mosquitoes.

928   *Annu. Rev. Entomol.* **44**, 131–157 (1999).

929 59. Foster, W. A. Mosquito sugar feeding and reproductive energetics. *Annu. Rev. Entomol.* **40**,

930   443–474 (1995).

931 60. Valenzuela, J. G., Francischetti, I. M. B., Pham, V. M., Garfield, M. K. & Ribeiro, J. M. C.

932   Exploring the salivary gland transcriptome and proteome of the *Anopheles stephensi*

933   mosquito. *Insect Biochem. Mol. Biol.* **33**, 717–732 (2003).

934 61. Dougherty, M. & Hamilton, G. Dodecanoic Acid Is the Oviposition Pheromone of *Lutzomyia*

longipalpis. *J. Chem. Ecol.* **23**, 2657–2671 (1997).

62. Dougherty, M. J., Guerin, P. M. & Ward, R. D. Identification of oviposition attractants for the

sandfly *Lutzomyia longipalpis* (Diptera: Psychodidae) in volatiles of faeces from

vertebrates. *Physiol. Entomol.* **20**, 23–32 (1995).

63. Ni, M. *et al.* Screening for odorant receptor genes expressed in *Aedes aegypti* involved in

host-seeking, blood-feeding and oviposition behaviors. *Parasit. Vectors* **15**, 71 (2022).

64. Spiegel, C. N. *et al.* The *Lutzomyia longipalpis* complex: a brief natural history of

aggregation-sex pheromone communication. *Parasit. Vectors* **9**, 580 (2016).

65. Hamilton, J. G. C., Maingon, R. D. C., Alexander, B., Ward, R. D. & Brazil, R. P. Analysis of

the sex pheromone extract of individual male *Lutzomyia longipalpis* sandflies from six

regions in Brazil. *Med. Vet. Entomol.* **19**, 480–488 (2005).

66. Watts, P. C. *et al.* Male sex pheromones and the phylogeographic structure of the

Lutzomyia longipalpis species complex (Diptera: Psychodidae) from Brazil and Venezuela.

Am. J. Trop. Med. Hyg. **73**, 734–743 (2005).

67. Maguire, S. E., Afify, A., Goff, L. A. & Potter, C. J. Odorant-receptor-mediated regulation of

chemosensory gene expression in the malaria mosquito *Anopheles gambiae*. *Cell Rep.* **38**,

110494 (2022).

68. Huh, J. R. *et al.* Multiple apoptotic caspase cascades are required in nonapoptotic roles for

Drosophila spermatid individualization. *PLoS Biol.* **2**, E15 (2004).

69. Kasimatis, K. R., Nelson, T. C. & Phillips, P. C. Genomic signatures of sexual conflict. *J.*

Hered. **108**, 780–790 (2017).

70. Sperling, A. L. & Glover, D. M. Parthenogenesis in dipterans: a genetic perspective. *Proc.*

Biol. Sci. **290**, 20230261 (2023).

71. Brazil, R. P. & Oliveira, S. M. Parthenogenesis in the sandfly *Lutzomyia mamedei* (Diptera:

Psychodidae). *Med. Vet. Entomol.* **13**, 463–464 (1999).

72. Goldberg, A., Mychajliw, A. M. & Hadly, E. A. Post-invasion demography of prehistoric

humans in South America. *Nature* **532**, 232–235 (2016).

73. Dos Reis, Y. V. & Alevi, K. C. C. Hybridization in Phlebotominae (Diptera: Psychodidae): A

mini-review. *Infect. Genet. Evol.* **86**, 104593 (2020).

74. Hedrick, P. W. Adaptive introgression in animals: examples and comparison to new

mutation and standing variation as sources of adaptive variation. *Mol. Ecol.* **22**, 4606–4618

(2013).

75. Arnold, M. L. & Martin, N. H. Adaptation by introgression. *J. Biol.* **8**, 82 (2009).

76. Mazzoni, C. J. *et al.* Multilocus analysis of introgression between two sand fly vectors of

leishmaniasis. *BMC Evol. Biol.* **8**, 141 (2008).

77. Mazzoni, C. J., Souza, N. A., Andrade-Coelho, C., Kyriacou, C. P. & Peixoto, A. A.

Molecular polymorphism, differentiation and introgression in the *period* gene between

Lutzomyia intermedia and *Lutzomyia whitmani*. *BMC Evol. Biol.* **6**, 85 (2006).

78. Testa, J. M., Montoya-Lerma, J., Cadena, H., Oviedo, M. & Ready, P. D. Molecular

identification of vectors of Leishmania in Colombia: mitochondrial introgression in the

Lutzomyia townsendi series. *Acta Trop.* **84**, 205–218 (2002).

79. Casaril, A. E. *et al.* Macrogeographic genetic structure of *Lutzomyia longipalpis* complex

populations using Next Generation Sequencing. *PLoS One* **14**, e0223277 (2019).

80. Araki, A. S. *et al.* Multilocus analysis of divergence and introgression in sympatric and

allopatric sibling species of the *Lutzomyia longipalpis* complex in Brazil. *PLoS Negl. Trop.*

Dis. **7**, e2495 (2013).

81. Galati, E. A. B. Phlebotominae (Diptera, Psychodidae): Classification, Morphology and

Terminology of Adults and Identification of American Taxa. in *Brazilian Sand Flies : Biology,*

Taxonomy, Medical Importance and Control (eds. Rangel, E. F. & Shaw, J. J.) 9–212

(Springer International Publishing, Cham, 2018).

82. Galati, E. A. B., Galvis-Ovallos, F., Lawyer, P., Léger, N. & Depaquit, J. An illustrated guide

for characters and terminology used in descriptions of Phlebotominae (Diptera,

Psychodidae). *Parasite* **24**, 26 (2017).

83. Vivero-Gomez, R. *et al.* Humidity and temperature preference in two Neotropical species of

sand flies. *Parasit. Vectors* **17**, 246 (2024).

84. Cheng, H., Concepcion, G. T., Feng, X., Zhang, H. & Li, H. Haplotype-resolved de novo

assembly using phased assembly graphs with hifiasm. *Nat. Methods* **18**, 170–175 (2021).

85. Ruan, J. & Li, H. Fast and accurate long-read assembly with wtdbg2. *Nat. Methods* (2019)

doi:10.1038/s41592-019-0669-3.

86. Durand, N. C. *et al.* Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-

C Experiments. *Cell Syst* **3**, 95–98 (2016).

87. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler

transform. *Bioinformatics* (2009).

88. Picard Toolkit. *Broad Institute, GitHub Repository* (2019).

89. Zhou, C., McCarthy, S. A. & Durbin, R. YaHS: yet another Hi-C scaffolding tool. *bioRxiv*

2022.06.09.495093 (2022) doi:10.1101/2022.06.09.495093.

90. Dudchenko, O. *et al.* The Juicebox Assembly Tools module facilitates de novo assembly of

mammalian genomes with chromosome-length scaffolds for under $1000.

doi:10.1101/254797.

91. Wang, X.-B., Liu, Q.-Y., Li, A.-L. & Ruan, J. SRY: An effective method for sorting long reads

of sex-limited chromosome. *bioRxiv* 2020.05.25.115592 (2020)

doi:10.1101/2020.05.25.115592.

92. Alonge, M. *et al.* Automated assembly scaffolding using RagTag elevates a new tomato

system for high-throughput genome editing. *Genome Biol.* **23**, 258 (2022).

93. Li, H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–

3100 (2018).

94. Winter, D. *Pafr: Read, Manipulate and Visualize 'Pairwise mApping Format' Data*. (2020).

95. Flynn, J. M. *et al.* RepeatModeler2 for automated genomic discovery of transposable

element families. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 9451–9457 (2020).

96. Riehl, K., Riccio, C., Miska, E. A. & Hemberg, M. TransposonUltimate: software for

transposon classification, annotation and detection. *bioRxiv* 2021.04.30.442214 (2021)

doi:10.1101/2021.04.30.442214.

97. Storer, J., Hubley, R., Rosen, J., Wheeler, T. J. & Smit, A. F. The Dfam community

resource of transposable element families, sequence models, and genome annotations.

Mob. DNA **12**, 2 (2021).

98. Smit, A. F. A., Hubley, R. & Green, P. *RepeatMasker Open-4.0*. (2013-2015).

99. Gabriel, L. *et al.* BRAKER3: Fully automated genome annotation using RNA-Seq and

protein evidence with GeneMark-ETP, AUGUSTUS and TSEBRA. *bioRxiv* (2023)

doi:10.1101/2023.06.10.544449.

100.Hoff, K. J., Lange, S., Lomsadze, A., Borodovsky, M. & Stanke, M. BRAKER1:

Unsupervised RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS.

Bioinformatics **32**, 767–769 (2016).

101.Brůna, T., Hoff, K. J., Lomsadze, A., Stanke, M. & Borodovsky, M. BRAKER2: automatic

eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by a

protein database. *NAR Genom Bioinform* **3**, lqaa108 (2021).

102.Hoff, K. J., Lomsadze, A., Borodovsky, M. & Stanke, M. Whole-Genome Annotation with

BRAKER. *Methods Mol. Biol.* **1962**, 65–95 (2019).

103.Kuznetsov, D. *et al.* OrthoDB v11: annotation of orthologs in the widest sampling of

organismal diversity. *Nucleic Acids Res.* **51**, D445–D451 (2023).

104.Butler, R. The design and development of Vectorbase: A bioinformatic resource center for

invertebrate vectors of human pathogens. (2010).

105.Giraldo-Calderón, G. I. *et al.* VectorBase: an updated bioinformatics resource for

invertebrate vectors and other organisms related with human diseases. *Nucleic Acids Res.*

43, D707–D713 (2014).

106. Li, H. Protein-to-genome alignment with miniprot. *Bioinformatics* **39**, (2023).

107. SRA Toolkit Development Team. *SRA Toolkit*. (2020).

108. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–915

(2019).

109. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–

2079 (2009).

110. Bruna, T., Lomsadze, A. & Borodovsky, M. GeneMark-ETP: Automatic Gene Finding in

Eukaryotic Genomes in Consistency with Extrinsic Data. *bioRxiv* (2024)

doi:10.1101/2023.01.13.524024.

111. Stanke, M. *et al.* AUGUSTUS: ab initio prediction of alternative transcripts. *Nucleic Acids*

Res. **34**, W435–9 (2006).

112. Hoff, K. J. & Stanke, M. Predicting Genes in Single Genomes with AUGUSTUS. *Curr.*

Protoc. Bioinformatics **65**, e57 (2019).

113. Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V. & Zdobnov, E. M.

BUSCO: Assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics* **31**, 3210–3212 (2015).

114. Gabriel, L., Hoff, K. J., Brůna, T., Borodovsky, M. & Stanke, M. TSEBRA: transcript selector

for BRAKER. *BMC Bioinformatics* **22**, 566 (2021).

115. Van der Auwera, G. A. *et al.* From FastQ data to high confidence variant calls: the Genome

Analysis Toolkit best practices pipeline. *Curr. Protoc. Bioinformatics* **43**, 11.10.1–11.10.33

(2013).

116. Korneliussen, T. S., Albrechtsen, A. & Nielsen, R. ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics* **15**, 1–13 (2014).

117. Wickham, H. ggplot2. *WIREs Comp Stat* **3**, 180–185 (2011).

118. Korunes, K. L. & Samuk, K. pixy: Unbiased estimation of nucleotide diversity and

divergence in the presence of missing data. *Mol. Ecol. Resour.* **21**, 1359–1368 (2021).

119. Weir, B. S. & Cockerham, C. C. Estimating f-statistics for the analysis of population

structure. *Evolution* **38**, 1358–1370 (1984).

120. Danecek, P. *et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158

(2011).

121. Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer

datasets. *Gigascience* **4**, 7 (2015).

122. Comeault, A. A., Kautt, A. F. & Matute, D. R. Genomic signatures of admixture and

selection are shared among populations of *Zaprionus indianus* across the western

hemisphere. *Mol. Ecol.* **30**, 6193–6210 (2021).

123. Nguyen, L.-T., Schmidt, H. A., von Haeseler, A. & Minh, B. Q. IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol. Biol. Evol.* **32**,

268–274 (2015).

124. Rabiee, M., Sayyari, E. & Mirarab, S. Multi-allele species reconstruction using ASTRAL.

Mol. Phylogenet. Evol. **130**, 286–296 (2019).

125. Malinsky, M., Matschiner, M. & Svardal, H. Dsuite - Fast D-statistics and related admixture

evidence from VCF files. *Mol. Ecol. Resour.* **21**, 584–595 (2021).

126. Wang, K., Mathieson, I., O'Connell, J. & Schiffels, S. Tracking human population structure

through time from whole genome sequences. *PLoS Genet.* **16**, e1008552 (2020).

127. Schiffels, S. *Msmc-Tools: Tools and Utilities for Msmc and msmc2.* (Github).

128. Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *Gigascience* **10**, giab008

(2021).

129. Pockrandt, C., Alzamel, M., Iliopoulos, C. S. & Reinert, K. GenMap: ultra-fast computation

of genome mappability. *Bioinformatics* **36**, 3687–3692 (2020).

130. R Core Team. *R: A Language and Environment for Statistical Computing.* (2024).

131. Hill, A. *et al.* Stepwise distributed open innovation contests for software development:

Acceleration of genome-wide association analysis. *Gigascience* **6**, 1–10 (2017).