Age-dependent expression and antiviral activity of interferon epsilon in respiratory epithelium

Mary McCabe[1], Helen E. Groves[1], Erin Getty[1], Emma Campbell[1], Connor G.G. Bamford[2], Guillermo Lopez Campos[1], Michael D. Shields[1], and Ultan F. Power[1*].

[1] Wellcome-Wolfson Institute for Experimental Medicine, Belfast, United Kingdom

[2] School of Biological Sciences & Institute for Global Food Security, Queen's University Belfast, Belfast, United Kingdom.

***Corresponding author**: Ultan F. Power PhD

Wellcome-Wolfson Institute for Experimental Medicine, School of Medicine, Dentistry & Biomedical Sciences, Queen's University Belfast, Medical Biology Centre, 97 Lisburn Road, Belfast BT9 7BL, Northern Ireland.

Tel: 44.28.9097.2285.

Email: u.power@qub.ac.uk

MM ORCID ID: 0000-0002-9307-5399

H.E.G ORCID ID: 0000-0001-9244-9961

E.G ORCID ID: 0009-0009-2027-7991 `

C.G.G.B ORCID ID: 0000-0003-4895-1983

G.L.C ORCID ID: 0000-0003-3011-0940

M.D.S ORCID ID: 0000-0002-3793-3571

U.F.P ORCID ID: 0000-0003-3246-3774

CRediT authorship contribution statement

Mary McCabe: Writing -Original Draft, Validation, Methodology, Investigation, Formal Analysis, Data Curation. **Helen E. Groves**: Conceptualization, Methodology, Supervision. **Erin Getty**: Investigation, Writing – Review & Editing. **Emma Campbell**: Investigation. **Connor G.G Bamford**: Methodology, Resources, Writing – Review & Editing, Supervision. **Guillermo Lopez Campos**: Methodology, Conceptualization, Writing – Review & Editing, Supervision. **Michael D. Shields**: Conceptualization, Writing – Review & Editing, Funding acquisition. **Ultan F. Power**: Conceptualisation, Resources, Writing – Review & Editing, Supervision, Funding acquisition.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

This work was supported by Wellcome Trust grant number 104516/Z/14/Z to HG; PHA HSCNI R&D Division funding (grant numbers RRG/3244105 and Com/4044/09), and Department for Economy, Northern Ireland to UFP.

Abstract

Respiratory syncytial virus (RSV) disease burden is greatest between six weeks and 6 months of life, with young age the most common risk factor among hospitalised children. A robust innate immune response in the airway epithelium is crucial for mitigating RSV-associated disease, but early-life immune responses to infection remain largely unexplored. RNA-seq analysis of RSV-infected primary nasal epithelial cell cultures from healthy infants at birth and one year revealed diminished expression of interferon epsilon (*IFNE*), a poorly characterised type I IFN, in newborns versus one-year. We hypothesised, therefore, that *IFNE* plays an important role during infant RSV infection. We found that *IFNE* is endogenously expressed in airway epithelial cell lines. Recombinant human IFNε (rhIFNε) induced an antiviral state against an RSV clinical isolate and related Sendai virus, but not SARS-CoV-2 under the conditions tested. The antiviral potency of rhIFNε was diminished relative to rhIFN beta (rhIFNβ1) or rhIFN lambda-1 (rhIFNλ1), as evidenced by IC_{50} data. Importantly, rhIFNε induced similar ISGs as rhIFNβ1 but demonstrated a transient temporal expression profile that differed from both rhIFNβ1 and rhIFNλ1. These results suggest that lower IFNε expression at birth may contribute to increased susceptibility to severe RSV-associated disease, offering insights into potential therapeutic interventions.

Keywords

Respiratory Syncytial Virus, Innate Immunity, Interferons, Interferon Epsilon, Sendai Virus, SARS-CoV-2

Introduction

Respiratory syncytial virus (RSV), a single-stranded negative-sense RNA virus of the *Pneumoviridae* family and *Orthopneumovirus* genus, is the commonest cause of acute lower respiratory tract infection (ALRTI) in infants globally [1]. Annually, ~33 million RSV LRTIs occur in children <5 years old, resulting in ~101,400 infant deaths [1]. Despite recent pharmacological advancements [2–6], treatment for severe RSV-associated disease remains palliative, with no RSV-specific antivirals or direct vaccination available for infants <6 months, the age group in which the disease burden is greatest [1, 7]. While epidemiological data has identified risk factors [8–14], most infants hospitalised share only the risk factor of age, suggesting host intrinsic features predispose them to severe disease. Because of the current incomplete understanding of RSV-host interactions in early life, we cannot predict which infants will experience severe disease and, thereby, prioritise prophylactic treatment. As RSV initially targets the ciliated cells of the nasopharynx [15–17], efficient innate immune responses play a vital role in determining RSV infection outcomes. Yet, whether and how these responses develop over time, and why infants <6 months are more frequently susceptible to severe disease, is unknown.

Interferons (IFN) are a crucial innate barrier against viral infection and dissemination [18, 19]. In the respiratory epithelium, type I (IFNα and -β) and type III (IFNλ1-λ4) IFNs play significant roles. Recently, Taveras *et al,* determined that all IFN types (I, II and III) from nasal samples were significantly increased in children (>6 years) versus infants (<6 months) and higher in milder outpatient RSV infections compared to severe inpatient infections [20]. Furthermore, we observed a threefold increase in mean IFNλ1 protein secretion at one year versus birth within paired birth and one-year primary nasopharyngeal cell samples (unpublished data, Groves HE, McCabe M, Coey J, Broadbent L, Guo-Parke H, Lopez Campos G, et al). These results suggest that strong IFN responses are a critical component of determining disease severity. As such, more

robust IFN responses to RSV infection with increasing chronological age may contribute to protection against severe disease in older infants and adults.

IFN epsilon (*IFNE*/IFNε) is a highly conserved type I IFN recognised for its multifaceted antimicrobial properties in the female reproductive tract (FRT) and its expression at other mucosal interfaces, including the lungs [21–24]. In contrast to other type I IFN subtypes, IFNε is 100-1000-fold less potent, and is expressed at high levels basally [25–27]. Interestingly, Thwaites *et al* described a trend for increased expression of type I IFNs (*IFNA1*, *IFNB1* and *IFNE*) in infants with moderate RSV ALRTI relative to severe cases [28] and *IFNE* was upregulated in children and young adults with SARS-CoV-2 infection [29]. Furthermore, *IFNE* was observed to be differentially expressed in children with long-coronavirus disease (COVID) symptoms [30]. Therefore, despite its conserved mucosal antiviral activity, understanding of the local expression and function of IFNε in airway epithelium against respiratory viruses, and its possible contributions to viral disease severity, remain limited.

In this study, we identified for the first time *IFNE* as differentially expressed with chronological age and confirmed its endogenous expression in immortalised airway epithelial cells. We assessed its antiviral activity against a clinical isolate of RSV and a related RNA virus, Sendai virus, and SARS-CoV-2. We determined the IC_{50} concentration of rhIFNε against RSV compared to other relevant mucosal IFNs and characterised the induction of downstream interferon-stimulated genes (ISGs). These results suggest an underappreciated role of IFNε in the induction of protective antiviral responses in the respiratory epithelium.

Materials and Methods

Cell Culture

BEAS-2B cells (kindly provided by Cliff Taggart, Queen's University Belfast) were maintained in 1

g/L (low) glucose DMEM with 10 units/mL Pen/Strep and 5% FBS. HEp-2 cells (kindly provided by

Prof. Ralph Tripp, University of Georgia) were maintained in 4.5 g/L (high) glucose DMEM with 10

units/mL Pen/Strep and 5% FBS. A549, Vero E6 and Vero's expressing ACE2 and TMPRSS2 (VAT)

cells were maintained in 4.5 g/L (high) glucose DMEM with 10 units/mL Pen/Strep and 10% FBS.

A549 expressing human ACE2 and TMPRSS2 (AAT) cells were maintained in 4.5 g/L (high) glucose

DMEM with 10 units/mL Pen/Strep, 10% FBS and maintained under antibiotic selection with

Hygromycin B (25 ug/mL) and Geneticin (25 ug/mL).

Nasal brushings were performed on healthy newborn term infants, (37–42 weeks gestation) and

preterm infants (28–34 weeks gestation) at birth and repeated in the same infants at one-year-

old, as previously described [31]. Harvested paediatric nasal cells were cultured to

differentiation using established methods [16][32] and infected with Multiplicity of Infection

(MOI) 3 in duplicate with a clinical isolate of RSV, designated BT2a, or mock-infected as previously

described [33].

RNA-Seq

127 Total RNA was extracted from WD-PNEC cultures at 96 hpi (High pure RNA isolation kit, Roche

128 diagnostics, Indianapolis, USA) and RNA quality was determined (Agilent RNA 6000 Nano Kit)

129 using the 2100 Bioanalyzer Instrument (Agilent Technologies). Total RNA sequencing was

130 conducted (unpublished data, Groves HE, McCabe M, Coey J, Broadbent L, Guo-Parke H, Lopez

131 Campos G, et al).

Virus Stocks and Titrations

RSV BT2a (clinical isolate) was isolated and characterised as previously described [33]. RSV

A2/mkate2 (a recombinant RSV reporter virus expressing Katushka, a far-red fluorescent protein)

was rescued from an infectious clone and helper plasmids kindly provided by Dr Martin Moore

(Emory University, Atlanta, USA). Sendai virus/eGFP (rSeV/eGFP) was generated by reverse

genetics [34]. SARS-CoV-2 Delta Variant was isolated from nasal swabs [35]. EMCV-Rueckert was

obtained as described previously [36]. Viral infections were performed at the MOI stated in each

figure legend by adding diluted virus suspensions to the apical compartment of WD-PNECs or

directly onto cell lines in serum-free medium. HEp-2 cells were used to titrate RSV BT2a, and Vero

E6 cells were used to titrate EMCV by tissue culture infectious dose 50 ($TCID_{50}$) assays. SARS-

CoV-2 plaque assays were performed on VAT cells [35].

<u>RNA extraction and RT-qPCR</u>

RNA was extracted according to the manufacturer's instructions (High pure RNA isolation kit,

Roche) and RNA quality was determined (Nanodrop One, Thermo Scientific®). cDNA synthesis

was performed in a 96-well thermocycler (Bio-Rad) using a High-Capacity cDNA Reverse

Transcription Kit (Applied Biosystems), according to the manufacturer's instructions. For the

target genes, *IFNE* (assay ID Hs00703565_s1*), IFNL1* (assay ID Hs00601677_g1), *IFNB1* (assay ID

Hs01077958_s1)*, MX1* (assay ID Hs00895608_m1)*, ISG15* (assay ID Hs01921425_s1)*, IFIT1*

(assay ID Hs03027069_s1)*, RSDA2* (assay ID Hs00369813_m1)*, IRF1* (assay ID Hs00971965_m1),

CXCL10 (assay ID Hs00171042_m1), YWHAZ (assay ID Hs01122445_g1), TBP (assay ID

Hs00427620_m1) and IPO8 (assay ID Hs00914057_m1) Thermo Fischer Scientific TaqMan ready-

to-use primer/probes were purchased and employed with the LightCycler 480 Probes Master Mix

(Roche). Relative quantification analysis of gene expression was performed using LightCycler 96

software (Roche). The Relative Quantification Cycle (Cq) of the sample was determined by

comparing the target gene to 3 reference housekeeping genes: *YWHAZ*, *TBP*, and *IPO8*.

<u>Recombinant human protein</u>

Cell lines were pretreated with recombinant human interferon epsilon (rhIFNε) (R&D systems,

Minneapolis, MN, USA), rhIFN beta (rhIFNβ1) (Peprotech®) and rhIFN lambda-1 (rhIFNλ1)

(Peprotech®) in low-serum DMEM for intervals stated within each figure legend. Cells were then

infected in serum-free DMEM. Following infection, cells were incubated for 2 h at 37 °C. After 2

h, the viral inoculum was removed and washed three-times with serum-free DMEM to remove

unattached virus, low-serum DMEM replaced and incubated at 37°C.

<u>Cell Imaging and Analysis</u>

The mean % fluorescence ratio of RSV A2/mkate2 and rSeV/eGFP infected cells was measured

using the brightfield and red/green fluorescence channels of the Celigo Imaging Cytometer

(Nexcelom). All data were presented as % fluorescence ratio ((brightfield confluence/red or green

confluence) *100).

<u>Generation of BEAS-2B IFNLR1$^{-/-}$ cells</u>

BEAS-2B cells overexpressing Cas9 were generated via lentiviral transduction of a third

generation lentiviral Cas9 expression plasmid (Addgene #52962), blasticidin selection (10 ug/mL)

and single-cell clonal isolation. Western blotting was used to confirm the stable expression of

Cas9 (Sigma-Aldrich, F3165). To generate *IFNLR1* BEAS-2B clonal cell line knockouts, a single-

guide RNA (sgRNA) (priCRISPR.IFNLR1.1(+) CACCGGTATTCGGACTCCACCCAG) was designed.

For each sgRNA, a forward and reverse oligonucleotide sequence was synthesised (Eurofins),

annealed and inserted into the lentiGuide-Puro vector via BsmB1-v2 (NEB) restriction digestion

and T4 DNA ligase (NEB) mediated ligation. Vector-sgRNA DNA was then transformed into DH10B

competent cells (Thermo Fischer Scientific) and spread onto Ampicillin selection plates (50

mg/mL). Single colonies were picked and miniprepped using PureLink™ HiPure Plasmid Miniprep

kit according to manufacturer's instructions (Invitrogen™). BEAS-2B *IFNLR1* knockout cell lines

were generated via lentiviral transduction of a third generation lentiGuide-Puro plasmid (Addgene

#84752) with inserted sgRNA sequence, puromycin (Gibco) selection and single-cell clonal

isolation.

<u>Statistical Analysis</u>

Statistical analyses included an unpaired two-tailed Student's t-test and one-way analysis of

variance (ANOVA) to assess the significance of test conditions, as stated in the figure legends.

For the time course profiles we calculated the Area under Curve (AUC) and then used this

summary measure for comparisons using one-way ANOVA. Statistical analysis was set at *

$p<0.05$, ** $p<0.01$, *** $p<0.001$, **** $p<0.0001$. Data presented as means ± standard error of the

means (SEM). Data were analysed using GraphPad® Prism 10 (GraphPad Software, Inc, La Jolla,

CA).

<u>Data Availability</u>

RNA-seq data has been deposited under moratorium pending publication and will be made

available to reviewers upon request. The GEO accession number for the RNA-seq data is:

GSE231666.

Results

Endogenous expression of *IFNE* increases during the first year of life.

In a recent study, we evaluated the impact of gestational and chronological age on RSV-induced airway epithelium innate immune responses using RNA-Seq (unpublished data, Groves HE, McCabe M, Coey J, Broadbent L, Guo-Parke H, Lopez Campos G, et al) (**Figure 1A**). RNA-seq analysis of RSV-infected WD-PNECs derived from infants sampled at birth and one-year-old revealed 63 differentially expressed genes (DEGs) (42 upregulated, 21 downregulated) with Log2Fold >|2| when comparing RSV-induced responses from 1-year- versus newborn-derived cultures (**Figure 1B**). Included in the 42 upregulated DEGs, *IFNE,* a poorly characterised type I IFN, was significantly increased following RSV infection versus uninfected controls (**Figure 1C**). Furthermore, *IFNE* expression was higher in 1 year- versus newborn-derived WD-PNECs, both at baseline and following RSV infection (**Figure 1C**). These results suggested that *IFNE* may play an unappreciated antiviral role in respiratory airway epithelium and contribute to more robust antiviral activities at 1 year compared to newborns.

230 **A.**

B.

C.

247 **Figure 1. Increase in *IFNE* expression during the first year of life.** (**A**) Summary of RNA-Seq experimental set-up. WD-PNEC cultures (n=12 paediatric donors) were infected with clinical isolate RSV BT2a (MOI=3) or mock-infected. RNA was extracted at 96 hpi and total RNA sequencing was conducted. (**B**) Volcano plot displaying differentially
248 expressed genes associated with RSV-infected cultures from 1-year-old and newborn donors, with *IFNE* expression highlighted. X-axis depicts the log2 differences in gene expression for each comparison of interest, with positive
249 values (in red) displaying significantly increased gene expression (log2 FC ≥ 2) and negative values (in blue) displaying significantly decreased gene expression (log2 FC ≤ -2). Y axis depicts adjusted p values (-log10 scale) for
250 each gene (higher p values representing greater statistical significance) after adjusting for batch by the year RNA-sequencing was completed. Green horizontal line indicates 0.05 p-value significance cut-off (**C**) *IFNE* Log2 Counts
251 per Million (CPM). Statistical analyses were corrected for multiple testing by applying the Benjamini-Hochberg (BH) p value correction (* p<0.05, ** p<0.01).

Endogenous expression of *IFNE* in immortalised airway epithelial cells.

Next, we assessed the expression of *IFNE* and two other mucosal IFNs within airway epithelial cells, including a type I (*IFNB1)* and a type III IFN (*IFNL1)*. *IFNB1* was chosen as a comparator as it was the sole type I IFN differentially expressed with infection within the RNA-seq data (unpublished data, Groves HE, McCabe M, Coey J, Broadbent L, Guo-Parke H, Lopez Campos G, et al). *IFNλ1* was chosen as a representative type III IFN as our group showed increased IFNλ1 release from primary cultures derived from infant respiratory tract samples following RSV infection [37]. Within common airway epithelial cell lines used to model RSV infection, *IFNE* was constitutively expressed at low levels, with the highest expression in adenocarcinoma human alveolar basal cells (A549) and A549 cells over-expressing human ACE2 and TMPRSS2 (AAT) (**Figure 2A**). Low levels of expression were evident in immortalised human bronchial epithelial cells (BEAS-2B) and human laryngeal carcinoma cells (HEp-2) (**Figure 2A**). This contrasted with *IFNB1* and *IFNL1*, which exhibited negligible expression under unstimulated sterile cell-culture conditions (**Figure 2B-C**). To further characterise *IFNE* expression, we infected A549 (**Figure 2D-F**) and BEAS-2B (**Figure 2G-I**) cells with a recombinant RSV reporter virus expressing a far-red fluorescent protein (RSV-A2/mKate2) or mock-infected them. At 48 hpi *IFNE, IFNB1* and *IFNL1* mRNA expression was higher within A549 cells (**Figure 2E-F**) compared to BEAS-2B cells (**Figure 2H-I**). *IFNE* expression was detected irrespective of infection in both cell lines (**Figure D, G**), with this unique endogenous pattern emphasized when comparing its expression to that of *IFNB1* or *IFNL1*.

Figure 2. *IFNε* is endogenously expressed in immortalised airway epithelial cells. BEAS-2B, A549, AAT and HEp-2 cells were seeded in 24 well plate. 24 h post seeding total RNA was extracted, reverse transcribed and subjected to qPCR analysis from BEAS-2B, A549, AAT and HEp-2 cells for *IFNE*, *IFNL1* and *IFNB1* expression (ROCHE, Lightcycler 480). Graphs show relative ratio values for (**A**) *IFNE*, (**B**) *IFNL1* and (**C**) *IFNB1* normalised using *YWHAZ*, *IPO8* and *TBP* as house-keeping genes. Data are presented as mean of triplicate technical replicates (±SEM) for n=3 independent experiments. (**D-F**) A549 or (**G-I**) BEAS-2B cells were mock-infected or infected with RSV-A2/mKate2 (MOI 1). At 48 hpi total RNA was extracted, reverse transcribed and subjected to qPCR analysis of *IFNE*, *IFNL1* and *IFNB1* expression (ROCHE, Lightcycler 480). Graphs show relative ratio values for *IFNE*, *IFNL1* and *IFNB1* normalised using *YWHAZ*, *IPO8* and *TBP* as housekeeping genes. Data are presented as the mean of duplicate technical replicates (±SEM) for n=3 independent experiments at 48 hpi. Statistical analyses were conducted using an unpaired two-tailed Student's t-test (* p<0.05, ** p<0.01).

IFNε has antiviral activity against RSV that is concentration- and cell type-dependent.

To assess if recombinant human IFNε (rhIFNε) protein can induce an antiviral effect within airway epithelial cell lines, A549 (**Figure 3A**) and BEAS-2B (**Figure 3B**) cells were pre-treated with rhIFNε, rhIFNβ1 and rhIFNλ1 protein (100 ng/mL) and infected with the highly IFN-sensitive virus encephalomyocarditis virus (EMCV) (MOI 1). IFNε pretreatment resulted in a 1.15-$\log_{10}$ reduction in EMCV titre versus the untreated in BEAS-2B cells (**Figure 3A**). This was similar to rhIFNλ1 pretreatment (1.7-log reduction) but much less effective than rhIFNβ1 pretreatment (3.5-$\log_{10}$ reduction). There was a non-statistically significant trend towards a reduction in virus titres within A549 cells (**Figure 3B**).

To explore the antiviral activity of rhIFNε against RSV, BEAS-2B (**Figure 3C**), HEp-2 (**Figure 3D**), A549 (**Figure 3E**) and AAT (**Figure 3F**) cells were pretreated with rhIFNε at a range of doses (100, 50, 5, 0.5 and 0.05 ng/mL) and infected with RSV-A2/mKate2 (MOI=0.3). A range was selected, as prior to this study, there was no evidence of IFNε's antiviral effect against respiratory viruses, including, but not limited to, RSV. At 24 hpi, the mean % fluorescence ratio was measured using the brightfield and red fluorescence channels of the Celigo Imaging Cytometer (Nexcelom). BSA pretreatment (5 μg/mL) was a negative control for the recombinant protein stabilisation carrier, and rhIFNβ1 and rhIFNλ1 pretreatment (100 ng/mL) were used as positive controls to ensure cell responsiveness to IFNs. Using fluorescence as a surrogate marker of viral replication, rhIFNε pretreatment resulted in a significant reduction in virus fluorescence in BEAS-2B (**Figure 3C**) and HEp-2 cells (**Figure 3D**), but not in A549 or AAT cells (**Figure 3E-F**), although there was a trend towards reduction in the latter cell lines. Reduction in virus replication was dose-dependent, with 100 ng/mL of rhIFNε resulting in the greatest reduction compared to untreated controls (1.8-fold reduction) in the most responsive cell line (BEAS-2B) (**Figure 3C**). This was less potent than rhIFNβ1 (27-fold reduction) but similar to rhIFNλ1 (1.9-fold reduction) (**Figure 3C**). RSV-A2/mKate2 fluorescence was much lower in AAT cells (**Figure 3F**) compared to their parental

counterpart A549 cells. To assess the antiviral effect against RSV BT2a [33], BEAS-2B cells were

pre-treated with 100 ng/mL of rhIFNε, rhIFNβ1and rhIFNλ1 and infected with RSV BT2a (MOI 0.3).

BEAS-2B cells were selected as they are a non-tumour transformed cell line and more responsive

to IFN stimulation. We found a 0.37-Log_{10} reduction at 24 hpi (**Figure 3G**) in RSV titres in rhIFNε

pre-treated cells compared to untreated controls. A similar reduction was seen with rhIFNβ1

(0.39-Log_{10} reduction) and rhIFNλ1 (0.44-Log_{10} reduction) pre-treatment at 24 hpi. Thereafter,

while mean virus growth kinetics were delayed slightly at 48 hpi, titres at 72 and 96 hpi were

indistinguishable from untreated- or BSA-controls (**Figure 3H**).

Figure 3. IFNε pretreatment reduces EMCV and RSV replication in BEAS-2B cells. (A) A549 and (**B**) BEAS-2B cells were pre-treated for 16 h with an IFN (100 ng/ml). Cells were then infected with EMCV (MOI=1) for 2 h at 37^{o}C. Data are presented as mean of duplicate technical replicates (±SEM) for n=4 independent experiments. Statistical analysis was conducted using one-way ANOVA. (**C**) BEAS-2B, (**D**) HEp-2 (**E**) A549, and (**F**) AAT cells were pre-treated for 16 h with IFNε (100, 50, 5, 0.5, and 0.05 ng/mL), IFNβ1 or IFNλ1 (100 ng/mL). Cells were then infected with RSV-A2/mKate2 (MOI=0.3). (C-F) At 24 hpi mean % fluorescence ratio was measured using the brightfield and red fluorescence channels of the Celigo Imaging Cytometer (Nexcelom). Data presented as % fluorescence ratio for mean of triplicate technical replicates (±SEM) n=3 independent experiments. Statistical analysis (C-F) was conducted using one-way ANOVA. (**G-H**) BEAS-2B cells were pre-treated for 16 h with 100 ng/mL of IFN. Cells were then infected with RSV BT2a (MOI=0.3). Supernatants were collected at 24, 48, 72, and 96 hpi and $TCID_{50}$ assays were conducted to determine RSV titres. Data are presented as the mean of duplicate technical replicates (±SEM) $\log_{10} TCID_{50}$/mL at (**G**) 24 hpi and a (**H**) 24-96 hpi time course for n=3 independent experiments. Statistical analysis for (**G**) was conducted using one-way ANOVA and for (**H**) using one-way ANOVA of AUC (* p<0.05, ** p<0.01, ***p<0.001, ****p<0.0001).

Relative antiviral activity of IFNε, IFNλ1 and IFNβ1 against RSV.

To further facilitate meaningful comparisons of the relative antiviral activity of rhIFNε with rhIFNβ1 and rhIFNλ1, we determined the half-maximal inhibitory concentration (IC$_{50}$) of these IFNs against RSV in BEAS-2B cells. Cells were pre-treated with decreasing concentrations of rhIFNε (**Figure 4A**), rhIFNλ1 (**Figure 4B**) and rhIFNβ1 (**Figure 4C**) and infected with RSV-A2/mKate2 (MOI=0.1). As anticipated, IFNε possessed the highest best-fit IC$_{50}$ value (9 ng/mL), followed by IFNλ1 (2.8 ng/ml) and IFNβ1 (0.24 ng/mL).

Given the limited understanding of how IFNε exerts its antiviral activity, we aimed to confirm that its presence would not interfere with the activity of other type I IFNs, specifically IFNβ1, in our study. Therefore, to further explore this without the interference from type III IFNs, *IFNLR1*$^{-/-}$ BEAS-2B cells (**Figure 4D**) were either pre-treated (**Figure 4E-F**), treated at infection (**Figure 4G-H)** or at 2 h post-infection (**Figure 4I-J**) with the IC$_{50}$ concentrations (**Figure 4E, G, I**) or 100 ng/mL (**Figure 4F, H, J**) of rhIFNε and rhIFNβ1. Cells were infected with RSV-A2/mKate2 (MOI=0.1). At 24 hpi, IC$_{50}$ concentrations of rhIFNε and rhIFNβ1 resulted in a reduction in viral spread with pre-treatment and with treatment at infection (**Figure 4E-H**). However, IC$_{50}$ concentrations of rhIFNε and rhIFNβ1 were not sufficient to significantly diminish RSV spread when treatment commenced 2 hpi (**Figure 4I)**. When pre-treated with both IFNs sequentially (rhIFNε and then rhIFNβ1 or vice versa) or treated with a combination of the two IFNs at the same time, the antiviral activity of rhIFNβ1 was not altered in the presence of rhIFNε. The greatest reduction in virus spread was seen when rhIFNβ1 was present regardless of the sequence of addition. Cells pre-treated or treated at infection with 100 ng/mL resulted in robust inhibition of viral spread, with those treated with 100 ng/mL rhIFNβ1 showing the greatest reduction (**Figure F-H**). Only 100 ng/mL of rhIFNβ1 inhibited viral spread significantly with treatment post-infection (**Figure 4J**). rhIFNλ1 pre-treatment (100 ng/mL) in *IFNLR1*$^{-/-}$ BEAS-2B cells did not result in a significant reduction in virus spread

348   (**Figure 4J**). These results demonstrated that rhIFNε does not interfere with the antiviral

349   activity of rhIFNβ1, despite both binding to the same receptor, at IC_{50} concentrations or 100

350   ng/mL.

Figure 4. IFNε possesses an increased IC$_{50}$ concentration against RSV compared to IFNβ1. BEAS-2B cells were pre-treated for 16 h with decreasing concentrations of (**A**) IFNε, (**B**) IFNλ1 and (**C**) IFNβ1. Cells were then infected with RSV-A2/mKate2 (MOI=0.1). At 24 hpi mean % fluorescence ratio was measured using the brightfield and red fluorescence channels of the Celigo Imaging Cytometer (Nexcelom). Statistical analysis (**A-C**) was conducted fitting a non-linear dose-response curve for mean of triplicate technical replicates for n=2 (IFNε and IFNλ1) or n=3 (IFNβ1) independent experiments in triplicate. (**D**) Summary of Experimental set-up for pretreatment, treatment at infection and treatment post-infection experiment. *IFNLR1* Cas9 knockout BEAS-2B cells were either pretreated (**E-F**), treated at infection (**G-H**) or treated 2 hours post infection (**I-J**) with the IC$_{50}$ concentration (**E, G, I**) or 100 ng/mL (**F, H, J**) of recombinant human IFNε and IFNβ1 protein. Cells were infected with RSV-A2/mKate2 (MOI=0.1). (**K**) *IFNLR1$^{-/-}$* BEAS-2B cells pretreated with 100 ng/mL of recombinant human IFNLR1. Data presented as % fluorescence ratio for as the mean of triplicate technical replicates (±SEM) n=3 independent experiments. Statistical analysis was conducted using one-way ANOVA (* p<0.05, **p<0.01, ***p<0.001, ****p<0.0001).

IFNε induces antiviral activity against Sendai Virus but not SARS-CoV-2.

To assess if an antiviral effect was evident beyond EMCV and RSV, A549 (**Figure 5A**) and BEAS-2B (**Figure 5B**) cells were pre-treated with a dose range of rhIFNε (100, 50, 5, 0.5 and 0.05 ng/mL) and infected with rSeV/eGFP (MOI=0.3). No significant reduction was observed in A549 cells, despite a reduction trend being observed (**Figure 5A**). At 24 hpi, a similar dose-dependent antiviral activity of rhIFNε was evident against rSeV/eGFP, as previously described for RSV A2/mkate2 in BEAS-2B cells (**Figure 5B**). The antiviral effect of rhIFNε pre-treatment (2.2-fold reduction) was like that of rhIFNλ1 (4-fold reduction) in BEAS-2B cells but less impressive compared to rhIFNβ1 (16.3-fold reduction) (**Figure 5B**). In contrast, AAT cells pretreated with rhIFNε, rhIFNβ1 or rhIFNλ1 (100 ng/mL) and infected with the SARS-CoV-2 Delta variant (MOI=0.3), had no significant reductions in viral replication with IFNε pre-treatment, a slight but significant reduction in virus replication with rhIFNλ1 pre-treatment, and a complete abrogation of virus replication following rhIFNβ1 pre-treatment at 48 hpi (**Figure 5C**) and over a time course of infection (**Figure 5D**). These results suggest that rhIFNε possesses antiviral activity against some RNA viruses, but not SARS-CoV-2 Delta Variant, at least at the dose tested, and its potency is clearly reduced compared to rhIFNβ1.

Figure 5. Relative potency of IFNε, IFNλ1 and IFNβ1 pre-treatment against Sendai virus and SARS-CoV-2.
(**A**) A549 and (**B**) BEAS-2B cells were pre-treated for 16 h with IFNε (100, 50, 5, 0.5, and 0.05 ng/ml), rhIFNβ1 or

rhIFNλ1 (100 ng/mL). Cells were infected with rSeV/eGFP (MOI=0.3). At 24 hpi mean % fluorescence ratio was measured using the brightfield and green fluorescence channels of the Celigo Imaging Cytometer (Nexcelom). Data presented as % fluorescence ratio for mean of triplicate technical replicates (±SEM) for n=3 independent

experiments. Statistical analysis was conducted using one-way ANOVA. (**C-D**) AAT cells were pre-treated for 16 h with rhIFNε, rhIFNλ1 and rhIFNβ1 (100 ng/mL). Cells were then infected with SARS-CoV-2 Delta variant

(MOI=0.3). Supernatants were collected at 24, 48, and 72 hpi and plaque assays were performed on VATs. Data presented as the mean of triplicate technical replicates (±SEM) $\log_{10}$ pfu/mL at (**C**) 48 hpi and a (**D**) 24-72 hpi

time course for n=3 independent experiments. Statistical analysis (**C**) was conducted using one-way ANOVA (**D**) Statistical analysis was conducted using one-way ANOVA of AUC (* p<0.05, **p<0.01, ***p<0.001, ****p<0.0001).

JAK-STAT inhibitor alters IFNε-mediated reduction of viral replication.

To confirm that IFNε was signalling via the classical type I IFN signal induction pathway (IFNAR-JAK-STAT) in airway epithelial cells, BEAS-2B (**Figure 6A-B**) and A549 (**Figure 6C-D**) cells were pre-treated with the JAK inhibitor Ruxolitinib for 2 h prior to rhIFNε (**Figure 6A, C**) and rhIFNβ1 (**Figure 6B, D**) pre-treatment (100 ng/mL). Pre-treatment with Ruxolitinib eliminated the reduction in viral replication with rhIFNε pretreatment in BEAS-2B cells (**Figure 6A**). A similar trend was seen for rhIFNβ1-pre-treated BEAS-2Bs (**Figure 6B**), although the concentration of Ruxolitinib used was not sufficient to completely block rhIFNβ1 signalling at 100 ng/mL. A549 cells did not produce a significant reduction in viral replication with rhIFNε treatment as described above (**Figure 6C**). However, pre-treatment of A549 cells with 100 ng/mL of rhIFNβ1 or rhIFNβ1 did result in a significant reduction in viral replication (**Figure 6D**). These results confirm the increased potency of rhIFNβ1 compared to rhIFNε at 100 ng/mL but that, like rhIFNβ1, rhIFNε utilises the JAK-STAT pathway to exert its antiviral effect against RSV.

423   **Figure 6. JAK-STAT inhibitor alters IFNε-mediated reduction of viral replication.** BEAS-2B and A549
cells were pre-treated with Ruxolitinib (INCB018424, Selleck Chemicals) or DMSO (2.5 µM) for 2 h before
424   16 h IFNε (**A-B**), and IFNβ1 (**C-D**) pre-treatment (100 ng/mL). Cells were then infected with RSV (RSV-
A2/mKate2) (MOI=0.1). At 24 hpi BEAS-2B (**A, C**) and A549 (**B, D**) cells mean % fluorescence ratio was
425   measured using the brightfield and red fluorescence channels of the Celigo Imaging Cytometer
(Nexcelom). Data presented as % fluorescence ratio for mean of triplicate technical replicates (±SEM)
for n=3 independent experiments for RSV, RSV+RUX, RSV+IFNε and RSV+RUX+IFNε. Data presented as
426   % fluorescence ratio of triplicate technical replicates for (±SEM) n=2 independent experiments for
RSV+DMSO and RSV+DMSO+IFNε. Statistical analysis was conducted using one-way ANOVA (* $p<0.05$
427   **p<0.01).

Differential ISG expression kinetics following rhIFNε, rhIFNλ1, and rhIFNβ1 treatment.

In our hands, IFNε was capable of inhibiting RSV replication only at early time points post-infection suggesting that the expression of downstream antiviral restriction factors may be transient. To assess if any differences were evident in the magnitude and expression kinetics of downstream antiviral restriction factors with rhIFNε treatment compared to rhIFNβ1 and rhIFNλ1, RT-qPCR was conducted on RNA extracted from BEAS-2B cells treated with rhIFNε, rhIFNλ1 and rhIFNβ1 (100 ng/mL) (**Figure 7A-F**). rhIFNε induced well known antiviral proteins, such as *MX1* (**Figure 7A**), *ISG15* (**Figure 7B**), *IFIT1* (**Figure 7C**) and *RSAD2* (**Figure 7D**). Similarly, IFNε also induced type I IFN-associated pro-inflammatory genes, such as *IRF1* (**Figure 7E**), and *CXCL10* (**Figure 7F**), which were previously reported to be induced only at low levels and transiently by rhIFNλ1 or other type III IFNs [38]. rhIFNε possessed unique ISG expression kinetics, with expression peaking early (4 h post-treatment), in comparison to rhIFNβ1, which induced a more potent and enduring response, and rhIFNλ1, which induced a slower yet incrementally increasing response through 24 h post-treatment. These results demonstrate that rhIFNε can induce a typical pro-inflammatory type I IFN antiviral gene signature in airway epithelial cells but that temporal expression of downstream ISGs is transient compared to rhIFNβ1 and rhIFNλ1 treatment.

Figure 7. Differential induction of antiviral and pro-inflammatory ISGs by IFNε. BEAS-2B cells were treated with IFNε, IFNλ1 or IFNβ1 (100 ng/ml) and total RNA was extracted immediately from untreated (t=0) and at indicated time points post-stimulation. RNA was subjected to RT-qPCR analysis of *IFIT1, ISG15, MX1, RSAD2, IRF1 and CXCL10* expression (ROCHE, Lightcycler 480). Graphs show relative ratio values for antiviral ISGs (**A**) *MX1*, (**B**) *ISG15*, (**C**) *IFIT1,* (**D**) *RSAD2 and pro-inflammatory ISGs* (**E**) *IRF1* and (**F**) CXCL10, normalised using *YWHAZ, IPO8* and *TBP* as housekeeping genes. Data presented (**A**-**D**) are mean of duplicate technical replicates (±SEM) fold change from mock for n=3 independent experiments and data presented (**E**-**F**) are mean of duplicate technical replicates (±SEM) fold change from mock for n=2 independent experiments.

Discussion

Conflicting data exists surrounding the role of IFN responses during RSV infection, particularly among those most vulnerable to severe RSV-related disease, infants aged 6 weeks to 6 months of life. Most of what is known describe the "classical" type I (IFNβ1 and IFNα's) and type III (IFNλ1-4) IFNs [39, 40]. However, little is understood about the "non-classical" IFN subtypes, such as IFNε. Previously we exploited a unique cohort of primary paediatric airway epithelial cell samples derived from the same healthy infants at birth and 1 year, to explore the development of innate immune responses to RSV infection in the first year of life (unpublished data, Groves HE, McCabe M, Coey J, Broadbent L, Guo-Parke H, Lopez Campos G, et al). We present here the differential expression of *IFNE* with chronological age, between healthy newborn and 1-year infants, both endogenously and following RSV infection in WD-PNECs cultures. This timespan coincides with those infants with high susceptibility rates to severe RSV-related disease (newborn) and those with lower frequencies of severe infection (1 year). Previous studies have reported endogenous and infection-induced *IFNE* expression in both the upper and lower respiratory tracts in humans and other mammals, with variations in expression linked to age and disease severity [24, 28, 29, 41, 42]. IFNε is conserved across various mammalian species [24, 43–46] but is a pseudogene in species susceptible to infections that are relevant to this study [47–49], alluding to a evolutionarily significant mucosal function. Collectively, these data suggest that IFNε may play a protective role in the respiratory epithelium during early-life viral infection and elevated levels in infants may be an indication of an improved ability to curtail and/or resolve infection.

As epithelial cells are the primary target of RSV infection *in vivo*, we assessed the expression of *IFNE* in immortalised epithelial cell lines commonly used to model RSV infection. *IFNE* was endogenously expressed at low levels in all cell lines tested. Recently, Martínez-Espinoza *et al* described increased *IFNE* expression during infection with RSV and hMPV, with peak *IFNE* expression at 48 hpi in A549 cells for both viruses [41]. We did not see this increase in *IFNE*

expression with RSV infection. This discrepancy may be due to various factors, including

differences in cell culturing conditions, viral strains used, assays performed, and other

experimental variables between authors. Our observations are consistent with its reported

unstimulated endogenous expression within the epithelium of the FRT and male reproductive

tract in humans and mice [21, 22, 50–52]. However, we did observe an increase in *IFNE*

expression within our RSV-infected WD-PNEC cultures, mirroring the increase Martinez-Espinoza

et al reported within human bronchial airway epithelial cells 3-5 days post RSV-infection [41].

Notably, Martínez-Espinoza *et al* suggested that RSV and hMPV induced *IFNE* expression in A549

cells via RIG-I, with contributions from MDA5 [41]. This suggests that viral RNA plays a role in the

induction of *IFNE* during active infection. The endogenous expression of *IFNE* is conserved across

multiple mucosal sites, concentrated primarily in the epithelial cells lining these surfaces.

Consequently, it may be regulated by distinct stimuli in a cell/tissue-specific manner, possibly by

distinct cell populations in these sites which fluctuate with tissue-specific stimuli. However, no

studies have been published regarding IFNε's role in human respiratory epithelium on a single-

cell level, and further work needs to be conducted regarding *IFNE* induction and regulation in

more physiologically relevant primary human airway epithelial models.

As a result of *IFNE's* endogenous epithelial expression, further investigation was prompted

regarding its possible contribution to localised antiviral activity. We showed that rhIFNε provided

antiviral protection against EMCV infection in BEAS-2B and A549 cells, consistent with similar

antiviral effects observed in cervical WISH cells [25]. Martínez-Espinoza *et al* demonstrated the

antiviral activity of rhIFNε against RSV and hMPV in A549 cells [41], which were shown to be highly

permissive to RSV infection [53]. Therefore, we chose to assess the antiviral properties of rhIFNε

in multiple respiratory cell lines. Pre-treatment with rhIFNε resulted in a significant dose-

dependent reduction in BEAS-2B and HEp-2 cell lines. BEAS-2B cells were observed to express

high levels of transcription factors and antiviral ISGs upon RSV infection and multiple IFN subtype

treatment, indicating a strong response to rhIFNε [53]. We did not observe a significant reduction

in virus fluorescence in A549 and AAT cells, when treated with 100 ng/mL of rhIFNε, consistent with Martínez-Espinoza *et al.*, who found that at least 250 ng/mL was needed for a significant effect on RSV fluorescence or titres in A549 cells [41]. In addition, prototypic lab-adapted strains, such as RSV A2/mkate2, have been reported to result in greater cytopathogenesis and induce greater concentrations of proinflammatory cytokines in the respiratory epithelium [33]. Therefore, we also chose to assess the antiviral activity of rhIFNε against a low passage clinical isolate of RSV, which more accurately reflects RSV infection *in vivo*. We found that IFNε pretreatment of BEAS-2B cells was antiviral against the clinical isolate RSV BT2a, with a significant reduction in infectious viral release at 24 hpi. However, without being replenished, this antiviral effect was quickly lost as the infection progressed. BEAS-2B cells were selected as the representative cell line due to their significant reduction in RSV A2/mKate2 infection following 100 ng/mL rhIFNε pretreatment and their non-cancerous immortalized state. Our data emphasizes the importance of cell line selection in studying viral IFN sensitivity, especially for lesser-understood IFN subtypes and suggests that while IFNε may not be the primary defender against viral infection in epithelial cells, they support its potential role as an endogenous antiviral mucosal defence cytokine.

The relative antiviral activities of each IFN were reflected in their IC_{50} values. Our data indicated that rhIFNε was ~3-fold less potent than rhIFNλ1 and ~38-fold less potent than rhIFNβ1, similar to those described for recombinant mouse IFNε (rmIFNε) against EMCV in WISH cells [25]. Interestingly, the IC_{50} concentration of rhIFNε and 100 ng/mL was sufficient to reduce viral replication under pretreatment and treatment-at-infection conditions, but not with treatment 2 h post RSV infection in BEAS-2B cells. However, its inability to provide protection against infection when added post-RSV infection most likely reflects its reduced antiviral potency and RSV's recognised ability to efficiently modulate IFN responses [54]. This is consistent with data showing a reduction in Zika virus replication in HRT8 cells with rmIFNε pretreatment, but not post-infection [50]. As expected, RSV was also able to antagonise IFNβ1's antiviral activity when added

post-infection at its IC_{50} concentration, as IFNε and IFNβ1 share the same signalling pathway. However, when added 2 h post-RSV infection at 100 ng/mL rhIFNβ1 was able to reduce viral replication significantly, most likely reflecting rhIFNβ1's greater potency. As anticipated the antiviral activity of rhIFNβ1 consistently superseded rhIFNε when added before, after or simultaneously with rhIFNε. Like IFNβ1, IFNε possesses a greater affinity for the IFNAR1 chain [25]. However, IFNε binds to IFNAR1 at a much lower affinity than IFNβ1 (100-1000-fold), which may partly explain the increased antiviral capacity of IFNβ1 in this context [25].

Infection with the related virus rSeV/eGFP confirmed that IFNε's antiviral effect was not restricted to RSV, with the greatest reduction in rSeV/eGFP fluorescence also observed within BEAS-2B cells. However, we did not observe an antiviral effect against the SARS-CoV-2 Delta variant in AAT cells at the very high dose tested (100 ng/mL), a variant that was previously described to have increased sensitivity to IFNs [55–58]. To our knowledge, this is the first report demonstrating SARS-CoV-2's insensitivity to rhIFNε [59]. Although our results suggest that rhIFNε was not effective in altering SARS-CoV-2 infection, it is important to note that we used a modified immortalized cell line highly susceptible to SARS-CoV-2 infection. Further studies in a more physiologically relevant nasal epithelial model are required, given evidence that IFNε may influence age-related responses to SARS-CoV-2 [29, 30]. It is possible that during SARS-CoV-2 infection, IFNε possess functions beyond direct antiviral activity, as suggested by its wider immunomodulatory functions [60]. Overall, our data demonstrate that respiratory viruses possess differential sensitivities to rhIFNε restriction and emphasise the importance of assessing the antiviral capacity against multiple virus families.

Our data using a JAK-STAT signalling inhibitor confirmed that IFNε signals through this pathway to exert its antiviral activity against RSV. These data provided the rationale to explore the magnitude and kinetics of the induction of downstream ISGs with rhIFNε treatment. Recently, IFNε was shown to induce common ISGs upregulated during RSV infection [41]. Our data showed that

rhIFNε treatment induced a unique ISG expression pattern with expression of four antiviral ISGs

associated with RSV infection (*MX1*, *ISG15*, *IFIT1* and *RSAD2*) peaking at 4 h post-stimulation.

IFNε's transient induction pattern was consistent with that observed against ZIKV infection in

human FRT cell lines [50] and this rapid transient ISG induction may be reflective of its short-term

reduction in viral titre at 24 hpi in BEAS-2B cells, but not at later time points. Furthermore, rhIFNε,

like rhIFNβ1, induced the expression of the transcription factor *IRF1* and the chemokine *CXCL10*.

rhIFNλ1 treatment did not induce *IRF1* or *CXCL10* expression to the same magnitude compared

to rhIFNε and rhIFNβ1. Rapid increased *IRF1* and *CXCL10 expression* are considered type I IFN-

associated [38]. However, type III IFNs were shown to induce *IRF1* and *CXCL10* expression at

lower levels [38]. Together our results demonstrated that rhIFNε can induce a strong classical

type I antiviral gene signature in airway epithelial cells but its potency and expression kinetics are

distinct compared to rhIFNβ1 and rhIFNλ1.

IFNε's unique low-level endogenous expression, its reduced potency, and transient ISG

expression kinetics, may strike a balance between the beneficial effects of type I IFN and

minimizing the tissue toxicity associated with its presence [61]. The question remains whether

the endogenous expression of IFNε results in a continuously primed airway epithelium, creating

an endogenous barrier to infection. If so, IFNε or the ISGs it induces could be leveraged as

potential biomarkers to identify those children at increased risk of severe disease in which no

high-risk factors have been identified (6 weeks and 6 months). Could the unique properties of

IFNε be further harnessed as an antiviral to non-destructively boost protective responses to

infection? Further work is required to explore IFNε's ability to boost or improve innate immune

responses within primary airway epithelial cell cultures.

<u>Conclusion</u>

IFNε is a highly conserved type I IFN expressed at multiple mucosal surfaces and is recognised

for its antimicrobial abilities in the FRT. We demonstrated differential expression of *IFNE* with age

and RSV infection in human nasal epithelial cells and confirmed its antiviral activity against a

clinical isolate of RSV. Although less potent than rhIFNβ1 or rhIFNλ1, rhIFNε's unique ISG

expression kinetics at 100 ng/mL and mild inflammatory profile suggest it may play a role in

mitigating RSV infection in airway epithelium. Reduced IFNε expression in newborns may

partially explain their heightened vulnerability to RSV, presenting an opportunity for therapeutic

interventions. Further studies are needed to explore IFNε's potential as an antiviral therapy or

prognostic tool for early-life respiratory infections.

618   **References**

619   [1]   Li Y, Wang X, Blau DM, et al. Global, regional, and national disease burden estimates of

620   acute lower respiratory infections due to respiratory syncytial virus in children younger

621   than 5 years in 2019: a systematic analysis. *Lancet* 2022; 399: 2047–2064.

622   [2]   Hammitt LL, Dagan R, Yuan Y, et al. Nirsevimab for Prevention of RSV in Healthy Late-

623   Preterm and Term Infants. *N Engl J Med* 2022; 386: 837–846.

624   [3]   Drysdale SB, Cathie K, Flamein F, et al. Nirsevimab for Prevention of Hospitalizations

625   Due to RSV in Infants. *N Engl J Med* 2023; 389: 2425–2435.

626   [4]   Ares-Gómez S, Mallah N, Santiago-Pérez M-I, et al. Effectiveness and impact of universal

627   prophylaxis with nirsevimab in infants against hospitalisation for respiratory syncytial

628   virus in Galicia, Spain: initial results of a population-based longitudinal study. *Lancet*

629   *Infect Dis* 2024; 24: 817–828.

630   [5]   Walsh EE, Pérez Marc G, Zareba AM, et al. Efficacy and Safety of a Bivalent RSV Prefusion

631   F Vaccine in Older Adults. *N Engl J Med* 2023; 388: 1465–1477.

632   [6]   Kampmann B, Madhi SA, Munjal I, et al. Bivalent Prefusion F Vaccine in Pregnancy to

633   Prevent RSV Illness in Infants. *N Engl J Med* 2023; 388: 1451–1464.

634   [7]   Hall CB, Weinberg GA, Iwane MK, et al. The burden of respiratory syncytial virus infection

635   in young children. *N Engl J Med* 2009; 360: 588–598.

636   [8]   MacDonald NE, Hall CB, Suffin SC, et al. Respiratory Syncytial Viral Infection in Infants

637   with Congenital Heart Disease. *N Engl J Med* 1982; 307: 397–400.

638   [9]   Chaw PS, Wong SWL, Cunningham S, et al. Acute Lower Respiratory Infections

639   Associated With Respiratory Syncytial Virus in Children With Underlying Congenital

640   Heart Disease: Systematic Review and Meta-analysis. *J Infect Dis* 2020; 222: S613–S619.

[10] Rose EB, Dahl RM, Havers FP, et al. Respiratory Syncytial Virus-Associated

Hospitalizations in Children With Neurological Disorders, 2006-2015. *J Pediatric Infect*

Dis Soc 2021; 10: 951–957.

[11] Moyes J, Cohen C, Pretorius M, et al. Epidemiology of Respiratory Syncytial Virus–

Associated Acute Lower Respiratory Tract Infection Hospitalizations Among HIV-Infected

and HIV-Uninfected South African Children, 2010–2011. *J Infect Dis* 2013; 208: S217–

S226.

[12] Kristensen K, Hjuler T, Ravn H, et al. Chronic Diseases, Chromosomal Abnormalities,

and Congenital Malformations as Risk Factors for Respiratory Syncytial Virus

Hospitalization: A Population-Based Cohort Study. *Clin Infect Dis* 2012; 54: 810–817.

[13] Shi T, Vennard S, Mahdy S, et al. Risk Factors for Poor Outcome or Death in Young

Children With Respiratory Syncytial Virus–Associated Acute Lower Respiratory Tract

Infection: A Systematic Review and Meta-Analysis. *J Infect Dis* 2022; 226: S10–S16.

[14] Chaw PS, Hua L, Cunningham S, et al. Respiratory Syncytial Virus-Associated Acute

Lower Respiratory Infections in Children With Bronchopulmonary Dysplasia: Systematic

Review and Meta-Analysis. *J Infect Dis* 2020; 222: S620–S627.

[15] Becker S, Soukup J, Yankaskas JR. Respiratory syncytial virus infection of human primary

nasal and bronchial epithelial cell cultures and bronchoalveolar macrophages. *Am J*

Respir Cell Mol Biol 1992; 6: 369–374.

[16] Villenave R, Thavagnanam S, Sarlang S, et al. In vitro modeling of respiratory syncytial

virus infection of pediatric bronchial epithelium, the primary target of infection in vivo.

Proc Natl Acad Sci U S A 2012; 109: 5040–5045.

[17] Zhang L, Peeples ME, Boucher RC, et al. Respiratory syncytial virus infection of human

airway epithelial cells is polarized, specific to ciliated cells, and without obvious

cytopathology. *J Virol* 2002; 76: 5654–5666.

[18]   Stanifer ML, Guo C, Doldan P, et al. Importance of Type I and III Interferons at Respiratory

and Intestinal Barrier Surfaces. *Front Immunol* 2020; 11: 608645.

[19]   Mesev E V, LeDesma RA, Ploss A. Decoding type I and III interferon signalling during viral

infection. *Nat Microbiol* 2019; 4: 914–924.

[20]   Taveras J, Garcia-Maurino C, Moore-Clingenpeel M, et al. Type III Interferons, Viral Loads,

Age, and Disease Severity in Young Children with Respiratory Syncytial Virus Infection. *J*

Infect Dis 2023; 227: 61–70.

[21]   Hardy MP, Owczarek CM, Jermiin LS, et al. Characterization of the type I interferon locus

and identification of novel genes. *Genomics* 2004; 84: 331–345.

[22]   Fung KY, Mangan NE, Cumming H, et al. Interferon-ε protects the female reproductive

tract from viral and bacterial infection. *Science* 2013; 339: 1088–1092.

[23]   Bourke NM, Achilles SL, Huang SU-S, et al. Spatiotemporal regulation of human IFN-ε

and innate immunity in the female reproductive tract. *JCI insight*; 7. Epub ahead of print

September 2022. DOI: 10.1172/jci.insight.135407.

[24]   Demers A, Kang G, Ma F, et al. The mucosal expression pattern of interferon-ε in rhesus

macaques. *J Leukoc Biol* 2014; 96: 1101–1107.

[25]   Stifter SA, Matthews AY, Mangan NE, et al. Defining the distinct, intrinsic properties of

the novel type I interferon, IFNε. *J Biol Chem* 2018; 293: 3168–3179.

[26]   de Geus ED, Volaric JS, Matthews AY, et al. Epithelially Restricted Interferon Epsilon

Protects Against Colitis. *Cell Mol Gastroenterol Hepatol* 2024; 17: 267–278.

[27]   Harris BD, Schreiter J, Chevrier M, et al. Human interferon-ε and interferon-κ exhibit low

potency and low affinity for cell-surface IFNAR and the poxvirus antagonist B18R. *J Biol*

Chem 2018; 293: 16057–16068.

[28]   Thwaites RS, Coates M, Ito K, et al. Reduced Nasal Viral Load and IFN Responses in

690   Infants with Respiratory Syncytial Virus Bronchiolitis and Respiratory Failure. *Am J*

691   *Respir Crit Care Med* 2018; 198: 1074–1084.

692   [29]   Pierangeli A, Gentile M, Oliveto G, et al. Comparison by Age of the Local Interferon

693   Response to SARS-CoV-2 Suggests a Role for IFN-ε and -ω. *Front Immunol* 2022; 13: 1–

694   10.

695   [30]   Fracella M, Mancino E, Nenna R, et al. Age-related transcript changes in type I interferon

696   signaling in children and adolescents with long COVID. *Eur J Immunol* 2024; 54: 1–10.

697   [31]   Groves HE, Guo Parke H, Broadbent L, et al. Characterisation of morphological

698   differences in well-differentiated nasal epithelial cell cultures from preterm and term

699   infants at birth and one-year. *PLoS One* 2018; 1–14.

700   [32]   Broadbent L, Villenave R, Hong G-P, et al. In Vitro Modeling of RSV Infection and

701   Cytopathogenesis in Well-Differentiated Human Primary Airway Epithelial Cells (WD-

702   PAECs). In: Tripp RA, Jorquera P (eds) *Human Respiratory Syncytial Virus*. Springer, 2016,

703   pp. 199–139.

704   [33]   Villenave R, O'Donoghue D, Thavagnanam S, et al. Differential cytopathogenesis of

705   respiratory syncytial virus prototypic and clinical isolates in primary pediatric bronchial

706   epithelial cells. *Virol J* 2011; 8: 43.

707   [34]   Rémi V, Olivier T, Surendran T, et al. Cytopathogenesis of Sendai Virus in Well-

708   Differentiated Primary Pediatric Bronchial Epithelial Cells . *J Virol* 2010; 84: 11718–

709   11728.

710   [35]   Bamford CGG, Broadbent L, Aranday-Cortes E, et al. Comparison of SARS-CoV-2

711   Evolution in Paediatric Primary Airway Epithelial Cell Cultures Compared with Vero-

Derived Cell Lines. *Viruses*; 14. Epub ahead of print February 2022. DOI:

10.3390/v14020325.

[36] Bamford CGG, Aranday-Cortes E, Filipe IC, et al. A polymorphic residue that attenuates

the antiviral potential of interferon lambda 4 in hominid lineages. *PLOS Pathog* 2018; 14:

e1007307.

[37] Guo-Parke H, Canning P, Douglas I, et al. Relative respiratory syncytial virus

cytopathogenesis in upper and lower respiratory tract epithelium. *Am J Respir Crit Care*

Med 2013; 188: 842–851.

[38] Forero A, Ozarkar S, Li H, et al. Differential Activation of the Transcription Factor

IRF1 Underlies the Distinct Immune Responses Elicited by Type I and Type III Interferons.

Immunity 2019; 51: 451-464.e6.

[39] Stephens LM, Varga SM. Function and modulation of type i interferons during respiratory

syncytial virus infection. *Vaccines* 2020; 8: 1–16.

[40] Broggi A, Granucci F, Zanoni I. Type III interferons: Balancing tissue tolerance and

resistance to pathogen invasion. *J Exp Med* 2019; 217: e20190295.

[41] Martínez-Espinoza I, Babawale PI, Miletello H, et al. Interferon Epsilon-Mediated Antiviral

Activity Against Human Metapneumovirus and Respiratory Syncytial Virus. *Vaccines*; 12.

Epub ahead of print October 2024. DOI: 10.3390/vaccines12101198.

[42] Fung KY, de Geus ED, Ying L, et al. Expression of Interferon Epsilon in Mucosal

Epithelium is Regulated by Elf3. *Mol Cell Biol* 2024; 44: 334–343.

[43] Chung O, Jung Y-E, Lee KW, et al. The Analyses of Cetacean Virus-Responsive Genes

Reveal Evolutionary Marks in Mucosal Immunity-Associated Genes. *Biochem Genet*

2022; 60: 2299–2312.

[44]   Abdel-Fattah M, Saeed H, El-Shennawy L, et al. The Arabian camel, Camelus

dromedarius interferon epsilon: Functional expression, in vitro refolding, purification

and cytotoxicity on breast cancer cell lines. *PLoS One* 2019; 14: e0213880.

[45]   Yang L, Xu L, Li Y, et al. Molecular and functional characterization of canine interferon-

epsilon. *J Interf cytokine Res Off J Int Soc Interf Cytokine Res* 2013; 33: 760–768.

[46]   Fischer CD, Wachoski-Dark GL, Grant DM, et al. Interferon epsilon is constitutively

expressed in equine endometrium and up-regulated during the luteal phase. *Anim*

Reprod Sci 2018; 195: 38–43.

[47]   Choo SW, Rayko M, Tan TK, et al. Pangolin genomes and the evolution of mammalian

scales and immunity. *Genome Res* 2016; 26: 1312–1322.

[48]   Ahmad HI, Jameel L, Zahra Q, et al. Molecular Evolution of Interferon-Epsilon (IFNε)

Pseudogene Modulates Innate and Specific Antiviral Immunity in Manis javanica. *J Zool*

Syst Evol Res; 2023. Epub ahead of print 2023. DOI: 10.1155/2023/2949008.

[49]   Shi W, Shi M, Que T-C, et al. Trafficked Malayan pangolins contain viral pathogens of

humans. *Nat Microbiol* 2022; 7: 1259–1269.

[50]   Coldbeck-Shackley RC, Romeo O, Rosli S, et al. Constitutive expression and distinct

properties of IFN-epsilon protect the female reproductive tract from Zika virus infection.

PLOS Pathog 2023; 19: e1010843.

[51]   Bourke NM, Mesiano S, Hertzog PJ. Spatiotemporal regulation of human IFNε and innate

immunity in the female reproductive tract.

[52]   Wijayarathna R, de Geus ED, Genovese R, et al. Interferon epsilon is produced in the

testis and protects the male reproductive tract against virus infection, inflammation and

damage. *PLoS Pathog* 2024; 20: e1012702.

[53]   Hillyer P, Shepard R, Uehling M, et al. Differential Responses by Human Respiratory

759   Epithelial Cell Lines to Respiratory Syncytial Virus Reflect Distinct Patterns of Infection

760   Control. *J Virol*; 92. Epub ahead of print August 2018. DOI: 10.1128/JVI.02202-17.

761   [54]   Jin H, Zhou H, Cheng X, et al. Recombinant Respiratory Syncytial Viruses with Deletions

762   in the NS1, NS2, SH, and M2-2 Genes Are Attenuated in Vitro and in Vivo. *Virology* 2000;

763   273: 210–218.

764   [55]   Guo K, Barrett BS, Morrison JH, et al. Interferon resistance of emerging SARS-CoV-2

765   variants. *Proc Natl Acad Sci U S A* 2022; 119: 1–8.

766   [56]   Nchioua R, Schundner A, Klute S, et al. Reduced replication but increased interferon

767   resistance of SARS-CoV-2 Omicron BA.1. *Life Sci alliance*; 6. Epub ahead of print June

768   2023. DOI: 10.26508/lsa.202201745.

769   [57]   Thorne LG, Bouhaddou M, Reuschl A-K, et al. Evolution of enhanced innate immune

770   evasion by SARS-CoV-2. *Nature* 2022; 602: 487–495.

771   [58]   Reuschl A-K, Thorne LG, Whelan MVX, et al. Evolution of enhanced innate immune

772   suppression by SARS-CoV-2 Omicron subvariants. *Nat Microbiol* 2024; 9: 451–463.

773   [59]   Bastard P, Gervais A, Le Voyer T, et al. Human autoantibodies neutralizing type I IFNs:

774   From 1981 to 2023. *Immunol Rev* 2024; 322: 98–112.

775   [60]   Mayall JR, Horvat JC, Mangan NE, et al. Interferon-epsilon is a novel regulator of NK cell

776   responses in the uterus. *EMBO Mol Med* 2024; 16: 267–293.

777   [61]   Rodero MP, Crow YJ. Type I interferon–mediated monogenic autoinflammation: The type

778   I interferonopathies, a conceptual overview. *J Exp Med* 2016; 213: 2527–2538.