Development and social dynamics of stone tool use in white-faced capuchin monkeys

Zoë Goldsborough[1,2,3,4,*], Meredith Carlson[4,5], Leonie Reetz[1,2], Margaret C. Crofoot[1,2,3,4], Brendan J. Barrett[1,2,3,4,**],

1 Department for the Ecology of Animal Societies, Max Planck Institute of Animal Behavior, 78467, Konstanz, Germany
2 Department of Biology, University of Konstanz, 78464, Konstanz, Germany
3 Center for the Advanced Study of Collective Behavior, University of Konstanz, 78464, Konstanz, Germany
4 Smithsonian Tropical Research Institute, Ancon, 0843-03092, Panama
5 Department of Anthropology, University of California, Davis, CA 95616, USA

Corresponding authors:
∗ zgoldsborough@outlook.com
∗∗ bbarrett@ab.mpg.de

Abstract

Percussive tool use for extractive foraging allows animals to access otherwise inaccessible resources and forage more efficiently, with potentially important implications for their fitness. The development of tool use proficiency has been well-documented in nut-cracking chimpanzees and robust capuchins, and in shellfish-cracking long-tailed macaques, where mothers and proficient tool users are the most important models. However, little is known about how tool use develops in populations where opportunities for social learning are scarce. White-faced capuchins (*Cebus capucinus imitator*) on Jicarón Island, Panama, provide a unique case to consider this question: stone tool use is entirely male-biased, meaning juveniles cannot learn from their mothers, and reduced group cohesion further limits social learning opportunities. Here, we investigate the acquisition and development of stone tool proficiency in this population using a year-long dataset from camera traps placed at two experimental anvils. We assess differences in proficiency between age classes, examine the development of tool use proficiency over time, and explore patterns of social attention during tool use. We show that juvenile capuchins are less proficient than subadults and adults, but their proficiency remains stable over the course of one year, suggesting that skill development may require prolonged practice or physical maturation. In contrast to other primates, social learning opportunities on Jicarón appear limited and scrounging is rare, yet we do find robust patterns in social attention when it occurs. Social attention to tool use mostly comes from juveniles too young to use tools themselves, who observe proficient subadults that tolerate scrounging. Our results contribute to the understanding of how complex tool use behaviors are acquired and maintained in primates, highlighting the role of social tolerance in the development of proficiency.

Introduction

Tool use is a striking example of behavioral flexibility in animals, and the use of tools for extractive foraging can increase fitness by allowing faster access to a greater variety of resources [1]. However, depending on the complexity of the tool use behavior, reaching proficiency is a process which can take years [2]. A behavior is considered tool use when an animal i) uses an object which is not part of their own body, ii) which is unattached to the surrounding environment (or attached and manipulatable [3]), and iii) the animal manipulates the object to reach a useful outcome [4]. To describe complexity of tool use behavior, a distinction can be made between first-order tool use (involving a single object, e.g., pounding a nut on a fixed surface) and second-order tool use (involving two objects, e.g., pounding a nut on a fixed surface using a hammerstone) [5]. Second-order tool use is thought to reflect greater cognitive complexity, as it requires more planning and physical coordination to correctly bring together all separate objects to obtain the desired outcome. Percussive stone technology—particularly extractive foraging using hammerstone and anvil tool use—is the focus of much research, given its high level of complexity, but also its relative rarity in the animal kingdom despite its prominent role in human evolution [6]. The only non-human wild primates that regularly use stone tools for extractive foraging are chimpanzees (*Pan troglodytes* [7]), long-tailed macaques (*Macaca fascicularis* spp. [8, 9]), robust capuchins (*Sapajus* spp. [10–13]), and gracile capuchins (*Cebus* spp. [14, 15]). Examining how stone tool use is acquired and how proficiency develops is key to understanding how a tool use tradition is maintained within a group, as well as how it spreads between groups.

For particularly complex techniques like percussive stone technology, acquisition of the skill may require sufficient exposure at a young age. In chimpanzees, insufficient exposure to nut-cracking during a critical learning period (between 3-10 years) may hinder or completely impede learning of the behavior in adulthood [16–18]. In robust capuchins [19] and macaques [20], acquisition of tool use also mostly occurs early in life. Exposure to tool use has an important social component: in both chimpanzees and robust capuchins, immature individuals are thought to acquire nut-cracking via social learning processes such as local enhancement, emulation, and social facilitation [16, 21–23]. Though social learning has been argued not to be critical to the acquisition of nut-cracking in captive chimpanzees [24], evidence from wild chimpanzees continues to support its important role [25].

In chimpanzees, most transmission of tool use behaviors occurs from mothers to their offspring [26, 27]. Juveniles spend a great deal of time with their mothers during the sensitive learning period, and mothers are highly tolerant of their offspring observing their tool use events and scrounging on opened nuts. There is also some evidence of female chimpanzees providing learning opportunities for their offspring through "scaffolding" (sharing nut-cracking materials such as hammers with their offspring) [28]. In contrast, robust capuchins mothers were not common models to their offspring. Rather, juveniles preferably observed tool use events by proficient tool users [22], who were older and also more dominant than themselves [21]. Social closeness or physical proximity did not affect who juveniles chose to observe. However, they did prefer individuals who were more tolerant of observers, and scrounging occurred in 75% of tool use events [21].

High inter-individual social tolerance is thought to be an important pre-condition for tool use acquisition [29], as observational learning of a complex skill involves close visual attention to the tool use activities. Being in such close proximity to an intolerant

tool-user is risky for an observer, who risks being chased off or attacked. With high social tolerance the opportunity to scrounge often arises, which not only gives naïve individuals the chance to observe the end-result of tool use, but also allows them to share in the payoff—thus increasing their motivation to acquire tool use themselves [30]. After acquisition, honing of tool use skill can be a lengthy process. Younger, active learners spend more time on acquiring tool use and are less proficient than experienced adults [20]. However, in bearded capuchins (*Sapajus libidinosus*), juveniles were not found to make more mistakes than adults during tool use. Rather, juveniles lower proficiency mainly resulted from less expertise in how to grip the hammerstone [31]. Due to differences in skill and/or strength, younger individuals may also be limited in what resources they can open [11]. Furthermore, percussive stone tool use might have an upper age limit. In chimpanzees, progressive old age led to a reduction in both nut-cracking frequency and efficiency [32].

While the ontogeny of stone tool use for nut-cracking is well studied in chimpanzees and robust capuchins, little is known about gracile capuchins, who have only recently been added to the list of non-human primate species habitually using stone tools. In particular, the white-faced capuchins (*Cebus capucinus imitator*) living in Coiba National Park, Panamá, are an important study system for studying tool use acquisition and proficiency development, as social learning opportunities seem to differ from other tool-using primates. White-faced capuchins on the islands of Coiba and Jicarón habitually use stone tools to crack open a variety of food items, including fruits like sea almonds (*Terminalia catappa*), coconuts (*Cocos nucifera*) and palm fruits (*Bactris major*, *Astrocaryum standleyanum*) as well as invertebrates like Halloween crabs (*Gecarcinus quadratus*), hermit crabs (*Coenobita compressus*), nerite snails (*Nerita sp.*) and other freshwater mollusks [14, 33]. However, on Jicarón, despite this tradition persisting for at least 17 years, it seems entirely localized to a 1.5 km stretch of coast, occupied by 2-3 neighboring capuchin groups [34]. Only one group of capuchins appears to use tools habitually on the full variety of resources, the other two neighboring groups have only been observed to sporadically use tools in the intertidal zone. In the most frequent tool-using group, stone tool use is fully male-biased: only males have ever been observed to use stone tools, despite females regularly using tools on the neighboring island of Coiba [35]. This male-bias means that juvenile capuchins on Jicarón cannot learn tool use from their mothers, as is common in chimpanzees. Furthermore, this group of tool-using capuchins appear to have reduced group cohesion (Goldsborough et al., in prep), further limiting social learning opportunities. Taken together, the tool-using capuchins on Jicarón present a unique situation where tool use is incredibly localized, yet persistent, and naïve individuals only have limited opportunity to observe tool use due to both less social cohesion as well as the male-biased nature of the behavior —which makes mothers unavailable as a model. Given these constraints, understanding how tool use is acquired and develops in this group provides insights into the flexibility of social and individual learning mechanisms in primates.

Herein, we provide a first exploration of the acquisition and development of stone tool use proficiency by one group of white-faced capuchins on Jicarón island. Using a year of data from camera traps on two experimental anvils, we compare features of tool use proficiency– efficiency and mistakes– between juvenile, subadult, and adult capuchins. Furthermore, we present a longitudinal investigation of the improvement in tool use skills by juveniles over a 1-year period, as well as an initial exploration of tool use skill in old age. Lastly, we consider potential mechanisms driving acquisition of stone tool use on Jicarón through examining social attention to tool use sequences. We explore who pays attention to tool use events, which tool users are most likely to receive social

attention, and how social attention relates to scrounging and tool user proficiency. In doing so, we provide a first reference of stone tool use proficiency development in white-faced capuchins, which will allow for comparison to other stone-tool-using primate species and lays the groundwork for future studies into the maintenance and spread of tool use behavior in this population.

Methods

Site and subjects

Jicarón island (2002 ha) is part of Coiba National Park, a UNESCO World Heritage site consisting of nine islands and 100 islets off the Pacific coast of the Veraguas Province, Panama. Jicarón island lies approximately 50 kilometers from the mainland, and is estimated to have been separated from the mainland for 14 000-18 000 years [36]. Jicarón island was inhabited by indigenous communities from 250 CE until the 16th century, served as a penal colony from 1919 until 2004 [37], but currently is entirely uninhabited with no human presence aside from occasional scientific and touristic expeditions. White-faced capuchins occur on Jicarón island, as well as on the largest island of Coiba and the small island of Rancheria. Jicarón has depauperate mammalian communities and a total absence of mammalian predators [38] and white-faced capuchins show increased terrestriality [39] and live at high densities [14, 40] compared to mainland populations. Two to three groups of capuchins on Jicarón – occupying a 1.5 km stretch of coast – use stone tools for extractive foraging [14], as well as at least one group on Coiba [33]. All known tool-using groups on Jicarón use tools for foraging in the intertidal zone, but only a single social group uses stone tools inland on a wider variety of resources. This group is the focus of this study, and their stone tool use appears to be entirely male-biased [35], with no observations of females using tools.

The most active tool-using group on Jicarón has been monitored using unbaited camera traps since 2017, but based on previous reports likely has been using tools since 2008 [14]. Stone tool use for extractive foraging has been documented to occur at three types of sites in this group, distinguishable through tool use activity and intensity, site erasure and debris accumulation (see [34] for more details). For the current study, we focus on tool use at high accumulation sites, from here on out referred to as "anvils", where capuchins habitually use tools and large amounts of debris accumulate over time. In 2022, the group was estimated to be comprised of around 20-25 individuals (5-6 adult males, 5-6 adult females, 4-5 subadults, 7-10 juveniles). These estimates are based on identifiable individuals (Table 1) as well as the maximum number of capuchins of a specific age-sex class observed together in one sequence. We are unable to obtain precise estimates of group size due to the nature of data collection via camera trapping, where the visual field is always limited, and juveniles are difficult to identify reliably from images alone.

Experimental anvil set-up

We set up two experimental anvil sites in the range of the tool-using group in January 2022: a wooden anvil at a site which has been heavily used since 2017 (location name CEBUS-02), and a stone anvil at an entirely novel location under sea almond trees (*Terminalia catappa*) where there was no evidence of prior tool use (EXP-ANV-01). To set up each experimental anvil, we first cleared the area of debris and placed a 1.8 m by 1.8 m piece of mesh on the ground, on which we placed the anvil. Anvils were selected from nearby based on the presence of a flat processing surface, as well as similarity in

Name	Code	Sex	Age estimate in years (in 2022)	Age class (2022)	Number of tool-using sequences	Notes
Known tool users						
Abraham	ABE	Male	30	Adult	0	Presumed alpha male, known tool user from previous data
Cystopher	CYS	Male	30	Adult	0	Presumed deceased
Snaggletooth McGee	SMG	Male	10-20	Adult	477	
Tom	TOM	Male	10-15	Adult	287	
Ink	INK	Male	10	Adult	6	
Larry	LAR	Male	7-8	Subadult	785	Border of old subadult/ young adult
Mick	MIC	Male	6-9	Subadult	401	Border of old subadult/ young adult
Spot	SPT	Male	7	Subadult	665	
Terry	TER	Male	5-6	Juvenile	227	Border of old juvenile/ young subadult
Balthasar	BAL	Male	4-5	Juvenile	384	
Zim	ZIM	Male	3-4	Juvenile	78	
Peaky	PEA	Male	3	Juvenile	254	
Joe	JOE	Male	<3	Juvenile	12	
Known non-tool users						
Mr. Email	MRE	Male		Adult		No tool use observed since 2017
Joker	JOK	Male		Subadult		
Olga	OLG	Female		Adult		Presumed deceased
Beatrice	BEA	Female		Adult		
Leona	LEO	Female		Adult		
Dottie	DOT	Female		Adult		
Sadie	SAD	Female		Adult		
Rosa	ROS	Female		(Sub)adult		
Blocky	BLO	Unknown		Juvenile		Very young (<2 years)
Mini	MIN	Unknown		Juvenile		Very young (<2 years)

Table 1: Overview of known individuals in the tool-using group, divided into known tool-users and known non-tool-users. Age estimates in years for tool users are rough approximations based on size and growth from 2017-2025. The borders for age classes are juveniles <6, subadults 6-8 years of age and adults >8 years.

size to the preexisting anvil at CEBUS-02, which had washed away prior to these experiments. Additionally, at each anvil, we supplied four uniquely marked hammerstones of varying sizes and weights sourced from nearby the anvils (Figure 1A). While both anvils were placed under sea almond trees, no nuts were included in the experimental set-up.

Two unbaited camera traps were attached to trees and aimed at each experimental anvil: a close-up video camera (Reconyx UltraFire XP9) recording a 60 second videos per trigger, and a wider-angle still (Reconyx Hyperfire HF2X) camera trap recording 30 images per trigger event without any between-trigger delays. Camera traps were placed in two deployments lasting approximately 6 months, and ran from January 2022 until January 2023. When camera traps were replaced in July 2022, both experimental anvils were excavated (Figure 1B), meaning all debris was collected down to the underlying mesh. Hammerstones and fragments found within the mesh perimeter were recovered and identified on the basis of markings and refits, when necessary. The remaining intact hammerstones were provided on the anvil during the second deployment.

Figure 1: Photographs of the experimental anvils, showing A) placement of CEBUS-02 in January 2022, and B) EXP-ANV-01 during excavation in July 2022. Photo credits belong to Brendan J. Barrett and Meredith Carlson.

Coding

All videos containing tool use events were coded frame-by-frame in BORIS v. 8.21.8 [41] by three coders (ZG, LR, and MC). We considered individual tool use sequences, where the start is defined as the moment an individual places an item on the anvil for processing and grabs the hammerstone. A sequence ends when either i) the item is opened and consumed ii) the individual relocates with the item to a place out of view of the camera or iii) the item is abandoned, which includes when the item was opened but not eaten. Per tool use sequence, we identified the age class (juvenile, subadult, or adult) of the tool user, and their identity when possible. We classified age based on visual appearance, with juveniles being individuals estimated to be <6 years old, subadults 6-8 years old, and adults >8 years old. We subsequently coded all behaviors by the tool user following an ethogram we established (Table S1, see also video ethogram https://keeper.mpdl.mpg.de/d/0c1b9853f3f342d8b3da/), coding features of tool use efficiency (e.g., number of successful pounds, number of misstrikes) and technique (e.g., type of pound [crouching, standing, jumping]). We also coded asocial (e.g., type of item processed, hammerstone properties) and social factors (e.g., displacements, social attention) on the level of the sequence (Table S2). A single video could contain one or more tool use sequences, and sequences could also span several videos if the individual was not finished processing when the video ended, and continued in the next triggered video.

To obtain additional information on social attention, sequences with other capuchins present besides the tool user were coded again by one coder (ZG) following a different protocol (Table S3). Per sequence, we coded the age-sex class of each other visible capuchin, as well as the age-sex class of capuchins paying social attention to, scrounging from, or displacing the tool user. Capuchins were identified whenever possible. Juveniles were usually not able to be sexed, as to do so requires close up images of their genitals for which the quality of the camera traps and image angle is insufficient. We also did not observe any subadult females present during tool use sequences, so only coded subadult males. Coders were all trained by ZG on a sample set of data before coding independently once inter-rater agreement on relevant measures (e.g., tool user age, outcome of sequence, number of pounds) was estimated to be good (Cohen's kappa

Statistical analyses

All statistical analyses were done in R v. 4.3.1 [42]. All Bayesian regression models were fit via the "brm" function in the brms package v. 2.16.1 [43]. We calculated contrasts to evaluate the credibility of the difference between categories (e.g., comparing juveniles' time to open an item to subadults' time) using the "hypothesis" function in brms. To identify credible effects, we considered what proportion of the posterior probability (PP) of the contrast was greater than 0, using a cut-off of 89% to reflect a reliable effect. In the aforementioned example, a posterior probability of 95% would reflect that the model estimates a 95% probability of juveniles taking more time than subadults to open an item.

Proficiency

To compare tool use proficiency between age classes, we first subsetted our data to only sequences where the item was successfully opened, as failure was too rare to study (occurring in fewer than 4% of observed tool use events). We further limited our analyses on proficiency to sequences where the tool user could clearly be identified. We also excluded sequences that were split across multiple videos, as several seconds of behavior can be missed in between triggers. Additionally, we only included sequences where sea almonds were the item being processed, since they constituted the vast majority (98%) of all tool use sequences and other items were underrepresented. In our dataset, each row represented a tool use sequence and variables were aggregated to the sequence level.

We quantified tool-using proficiency per sequence in four different ways: i) seconds needed to open an item, ii) number of pounds to open an item, iii) number of repositions of the item on the anvil in between strikes and peels of the sea almond's exocarp, iv) number of misstrikes and strikes that sent the item flying off the anvil. To compare sequence duration between age classes, we fit a generalized linear Gamma regression (*Model e1*), with sequence duration in seconds as the outcome. As predictors, we included the age class of the tool user (juvenile, subadult, or adult, with juvenile as reference level), the color of the sea almond (brown, green, red, or unknown, corresponding to its ripeness and with brown as reference level), the material of the anvil (wood or stone, with stone as reference), and a random effect of the identity of the tool user. We included the color of sea almond and the material of the anvil, as we expected both to potentially have an effect on the number of pounds required to open a sea almond. Fresh sea almond fruits are green, and turn yellow, then red, as they ripen [44]. After a while, cached fruits turn brown, and we expect these dried out fruits to be easier to crack than fresh, green ones. Stone and wood have different densities and hardness, likely also affecting how easily the sea almond cracks when smashed with a stone hammer.

To compare number of pounds to open an item between age classes, we fit another generalized linear model (*Model e2*), with the same structure as the model described above. As the predictor was the discreet number of pounds, we used a Poisson distribution. We also compared the rate of pounding (so number of pounds per second) between age classes by running a model with the same structure as *Model e2*, but with an offset of the log of sequence duration in seconds (*Model e2b*). We used the same model structure—without an offset—to consider the number of item repositions (*Model e3a* and item peels (*Model e3b*) during a sequence. Misstrikes, where a capuchin attempts to strike the item but misses it entirely, were not very common. Thus, to

compare its occurrence between age classes, we fit a zero-inflated Poisson model, with the same predictors as above (*Model e4a*). Lastly, to compare the occurrence of items flying off the anvil between age classes, which was similarly rare, we fit another zero-inflated Poisson model (*Model e4b*).

Development

To examine the development of tool use proficiency over time, we looked at known individuals and how their processing of one color of sea almond (brown, which is most frequently processed) changed over the course of the full year of coding. We visually considered how the number of pounds, number of repositions, and number of misstrikes changed, with the expectation that (sub)adults would remain rather stable, while juveniles would show fewer pounds, repositions, and mistakes as their proficiency increased. For the number of pounds, we also ran a Poisson GLM (*model dev1*) with as predictors an interaction of time (parameterized as the log of the seconds passed since January 1, 1970) with the identity of the tool user. Furthermore, we also considered a case study of one adult male, estimated to be the oldest in the capuchin group, to see if and how his tool-using behavior changed since the beginning of sampling in this project (in 2017).

Social attention

To examine social learning's role in tool use acquisition, we ran further analyses on a subset of sequences where other capuchins were present. Here, we also excluded sequences split across several videos, as well as sequences where one of the capuchins present was unable to be assigned an age-sex class, or where the ending of the tool use sequence was not captured on video (so the sequence likely continued after the video stopped recording). We also restructured our dataset to have each row reflect an individual present during a tool-using sequence, with a variable indicating whether they paid social attention yes/no.

First, to consider which age class is more likely to receive social attention, and which age-sex class is more likely to pay social attention to tool use, we fit a generalized Bernoulli logistic regression (*Model socatt1*). The outcome of this model was social attention yes/no, while the predictors included the age of the tool user (juvenile, subadult, adult, with juvenile as reference) and the age-sex class of the observer (juvenile, subadult male, adult female, and adult male, with juvenile as reference). We also included location (CEBUS-02 or EXP-ANV-01) and two social factors which may affect the probability of social attention occurring, namely the total number of capuchins present during the sequence, and the number of capuchins scrounging during the sequence. Lastly, we included an offset of the log of sequence duration, to account for longer sequences having more opportunities for social attention, and a random effect of sequence ID, to account for non-independence of observations from the same sequence.

Second, to consider whether more efficient tool users, or specific individuals, receive more social attention, we subsetted the data further to only sequences where the item was opened, and where the tool user could be identified. We ran another generalized Bernoulli logistic regression (*Model socatt1b*) with the outcome of social attention yes/no, and predictors of tool user age, age-sex of observer, the number of pounds used to open the item, and the number of mistakes during the sequence. We also included an offset of sequence duration, and also the identity of the tool user as a random effect.

All models were fit with regularizing Normal (0,1) priors for intercepts and predictors, and exponential (1) priors for standard deviations of varying effects. For each model, we did a prior predictive simulation to compare our chosen priors to default priors, and to evaluate the identifiability of parameters. The final models were ran with three chains of 3000 iterations each, with a 1500 iterations warm-up per chain. Our models were stable with large effective sample sizes (Bulk_ESS and Tail_ESS over 1000 for all estimates; [43] and Rhat values <1.01 [45]. Additionally, Pareto k estimates for all models were below 0.7. We visually assessed model fit and confirmed our choice of priors using the posterior predictive check function.

Data availability

Details of model output are available as supplemental information. All code and data necessary to replicate analyses are publicly available [46].

Results

Between January 2022 and January 2023, we recorded a total of 3657 tool use sequences at the two anvil sites spread across 215 days. CEBUS-02 was the more popular site, with 2574 observed sequences compared to 1083 at EXP-ANV-01. Both locations have gaps in tool use activity over the time period (Figure S1). At CEBUS-02, the video cameras ran out of battery prematurely due to the higher activity (cameras ran 2022-01-10 until 2022-06-29 and 2022-07-16 until 2022-09-30). At EXP-ANV-01, the capuchins only regularly started using the novel anvil in March 2022, and reduced their activity after a tree branch fell on anvil on 2022-10-10, though cameras ran the full time period (2022-01-10 until 2022-07-16 and 2022-07-17 until 2023-01-01). The average number of sequences observed on a day when tool use occurred was 17.01 (range 1-114). Sea almonds were the item being processed in the vast majority of tool use sequences (3601 sequences equal to 98%, Table 2).

Item		Count	Percentage
Sea almond	Brown	2381	**65.11**
	Green	553	**15.12**
	Red	78	**2.13**
	Unknown	589	**16.11**
Coconut		12	**0.33**
Fruit		1	**0.03**
Crab (halloween or hermit)		12	**0.33**
Other		16	**0.44**
Unknown		15	**0.41**

Table 2: Frequency of items being processed during all observed tool use sequences

Capuchins successfully opened the item in 89% of the sequences. Success rates were lower for juveniles (79%) compared to subadults and adults (93% and 90% respectively). The number of pounds needed to open a single item ranged from 1 to 34. In the remaining sequences, the tool user either abandoned the item (4%), relocated to another anvil with the item (3%), or the ending was not captured on camera (4%). In both sites, the same hammerstone was used in over 90% of the tool use sequences, and all age classes used the same hammerstones.

We were able to classify the tool user's age in all of the sequences, and reliably identify the tool user in 98% of the sequences (see Table 1 for the number of tool use sequences observed per individual). The tool user was the only individual visible in the video in 72% of all sequences. When other capuchins were present (1010 sequences), we observed social attention occurring in 217 sequences (21%). Both displacements and scrounging were rare, occurring in 12% and 25% of opportunities (meaning, other capuchins being present).

Tool-using proficiency

Efficiency (number of pounds and processing time)

The sample used for analyses consists of 2900 sequences where i) the tool user successfully opened a sea almond, ii) the tool user was identified, and iii) the sequence was not split across several videos ($n_{adult} = 662$, $n_{subadult} = 1612$, $n_{juvenile} = 626$). Our models estimated that juveniles need considerably longer to open a sea almond with stone tools than subadults and adults (Figure 2, for model estimates see Tables S4 & S5). Juveniles took an average of 19.30 seconds (95% CI [14.30, 25.53]) and 5.21 pounds (95% CI [4.39, 6.30]) to open an item, considerably longer than subadults, who took on average 9.78 seconds (95% CI [5.99, 15.33]) and 3.63 pounds (95% CI [2.75, 4.71]), and adults, who took 11.25 seconds (95% CI [7.17, 18.54]) and 3.82 pounds (95% CI [2.80, 5.05]). Juveniles taking more seconds and more pounds to open an item than subadults and adults were credible effects (all PP $>0 = >98\%$). Subadults and adults did not differ significantly from one another in both models.

The level of ripeness of the sea almond and the material of the anvil affected both seconds and pounds required to process a sea almond. Fresher sea almonds (green and red in color) took more time and more pounds to open than older (brown) sea almonds. We estimated green almonds to require 1.32 times more seconds (95% CI [1.27, 1.38], PP $>0 = 100\%$) and 1.26 times more pounds (95% CI [1.21, 1.32], PP $>0 = 100\%$) than brown almonds. Red almonds 1.49 times more seconds (95% CI [1.34, 1.65], PP $>0 = 100\%$) and 1.31 times more pounds (95% CI [1.17, 1.45], PP $>0 = 100\%$). Red and green sea almonds did not differ considerably from one another in processing times or number of pounds. The material of the anvil did not affect processing times, but the number of pounds was slightly higher (1.09 times, 95% CI [1.06, 1.14], PP $>0 = 100\%$) at the stone anvil than at the wooden anvil. The relationship between the number of pounds and the sequence duration (so the speed at which individuals pound) was positive between age classes (Figure 3). However juveniles showed a 0.75 times lower rate of pounding (number of pounds per second) than adults (95% CI [0.62, 0.93], PP $>0 = 98\%$) and 0.74 times lower than subadults (95% CI [0.61, 0.89], PP $>0 = 99\%$, for all estimates see Table S6).

Figure 2: Model estimates and mean values of pounds, repositions, and peels in one tool use sequence opening a single sea almond per age-class (juvenile, subadult, adult). Model estimates are reflected by violin plots, and observed means from data are represented as points.

Figure 3: Relationship between number of seconds and number of pounds needed to open a single item using stone tools, separately for each age class.

Repositioning and peeling

All age classes showed repositioning of the item on the anvil during a tool use sequence, or picking it up and peeling it before continuing to strike, but juveniles did it more than adults and subadults (Figure 2, for model estimates see Tables S7 & S8). Juveniles showed considerably higher incidences of repositioning and peeling ($\text{mean}_{\text{n\_reposition}} = 1.11$, 95% CI [0.64, 1.80]; $\text{mean}_{\text{n\_peel}} = 0.43$, 95% CI [0.27, 0.68]) than subadults ($\text{mean}_{\text{n\_reposition}} = 0.33$, 95% CI [0.16, 0.79]; $\text{mean}_{\text{n\_peel}} = 0.18$, 95% CI [0.09, 0.37]) and adults ($\text{mean}_{\text{n\_reposition}} = 0.32$, 95% CI [0.15, 0.79]; $\text{mean}_{\text{n\_peel}} = 0.18$, 95% CI [0.09, 0.40]). Juvenile tool users repositioning and peeling more than subadults and adults were credible effects (all PP $>0 = >98\%$). Tool users peeled fresher sea almonds, in particular green ones, more often than brown, older sea almonds (1.48 times more, 95% CI [1.23, 1.75], PP $>0 = 100\%$). The material of the anvil made no difference in the number of repositions or peels.

Mistakes

Tool users rarely made mistakes during tool use sequences (occurring 323 times in the sample of 2900 tool use sequences), and in 47% of cases the tool user making a mistake was juvenile. The most common type of mistake was the item flying off the anvil after being struck (occurring 161 times), followed by misstrikes (106 times), and dropping the hammerstone (occurring 56 times). Juvenile tool users were more likely to make misstrikes or sent the item flying off ($\text{mean}_{\text{n\_misstrikes}} = 0.14$, 95% CI [0.03, 0.51]; $\text{mean}_{\text{n\_itemflies}} = 0.32$, 95% CI [0.18, 0.52]) than subadults ($\text{mean}_{\text{n\_misstrikes}} = 0.04$, 95% CI [0.01, 0.26]; $\text{mean}_{\text{n\_itemflies}} = 0.07$, 95% CI [0.05, 0.14]) and adults ($\text{mean}_{\text{n\_misstrikes}} = 0.06$, 95% CI [0.01, 0.35]; $\text{mean}_{\text{n\_itemflies}} = 0.06$, 95% CI [0.03, 0.13]). Juveniles making more mistakes than subadults was a credible effect (misstrikes PP $>0 = 91\%$, items flying PP $>0 = 100\%$) as was juveniles sending more items flying than adults (misstrikes PP $>0 = 83\%$, items flying PP $>0 = 100\%$) The posterior probability of juveniles having more misstrikes than subadults and adults was over 84%, and 100% for juveniles having more items flying (for model estimates see Tables S9 & S10).

The material of the anvil also affected the likelihood of the item flying off: a sea almond was 3.42 times more likely to fly off of a stone than wooden anvil (95% CI [2.56, 4.62], PP $>0 = 100\%$). Green and red sea almonds were also 3.49 (95% CI [2.53, 4.81], PP $>0 = 100\%$) and 2.46 times (95% CI [1.27, 4.62], PP $>0 = 99\%$) more likely to fly off when struck than brown, dried out, sea almonds.

Tool use development

We observed little change in tool use proficiency over the year for all age classes. Adults' performance was the most stable, but juveniles and subadults also did not show considerably changes in the number of pounds (for model estimates see Table S11), repositions, and misstrikes over time (Figure 4). The reduced sampling near the end of the observation period, combined with some individuals (mostly adults) showing clustered activity rather than activity spread over time, means that there is more uncertainty at certain times of the year.

In light of tool use skill development as a function of age, it is important to mention one individual who was absent as a tool user in our sample: ABE. This adult male is estimated to be at least 30 years old, and the likely alpha male of the group, inferred by his size, him being recruited by other group members in conflicts, and the absence of him receiving aggression. While he was observed using tools 73 times in data collected

Figure 4: Development of number of pounds (purple), misstrikes (green), and repositions (blue) per individual over the full observation period for brown sea almonds only. The color of the box with the individuals code represent their age class (blue for juveniles, lilac for subadults, orange for adults). Points are raw data, with point shape corresponding to anvil location. The smooth curve is estimated using method "loess" and formula "y∼x".

2017-2019, he never used tools in the current observations. However, 13 times ABE displaced a tool user from the anvil (out of 175 total displacements), and then consumed the item they opened. No other individuals "stole" from tool users in this manner.

Social attention

After filtering, our sample contained 844 sequences that were completely captured within a single video, during which other capuchins aside from the tool user were present, who could all be assigned an age-sex class. The majority of other capuchins present were juveniles (833 out of 1346 capuchins present, 62%), followed by subadults (269, 20%) and adult males (135, 10%). Adult females were most rarely present during tool use sequences (109 times, 8%). In this filtered sample, social attention occurred in 183 (22%) of the sequences with capuchins present.

Individuals paying social attention were overwhelmingly juveniles, an adult male was observed paying social attention once, adult females five times and subadult males 18 times. Of the 208 occurrences of juveniles paying social attention, 23 were identified as known tool-using juveniles. The remaining unidentified juveniles were likely not tool users, as they were too young (<3 years) to easily be identified and we have not observed regular tool use by any individuals of this age.

Our model exploring factors predicting the probability of social attention also found juveniles to be the age class most likely to pay social attention. Given an adult male tool user at the most frequently used location (CEBUS-02), juveniles had a probability of showing social attention of 0.16 (95% CI [0.09, 0.26]), considerably higher than subadult males (0.05, 95% CI [0.02, 0.10]), adult males (0.02, 95% CI [0.01, 0.05]), and adult females (0.03, 95% CI [0.01, 0.07]). In terms of who received social attention, given a juvenile observer at CEBUS-02, juveniles were less likely to receive social attention [0.07, 95% CI: 0.04-0.10], than subadults [0.18, 95% CI: 0.14-0.23] and adults [0.16, 95% CI: 0.09-0.26], who did not differ significantly from one another (Figure 5). Juveniles being more likely to pay social attention and less likely to receive social attention than other age-sex classes was a credible effect (PP >0 for all contrasts = 100%).

Figure 5: Likelihood of paying social attention to a tool user, presented per age-sex class of the observer and age of the tool user. Model estimates are reflected by box plots, and observed means from data are represented as points.

When more capuchins were present during a tool-using sequence, the chance of social attention occurring was lower. However, in sequences where more individuals were scrounging, social attention was more likely to occur (Figure 6, for model estimates Table S12). When we consider the type of scrounging (whether it was tolerated by the tool user during the tool use event, or only occurred after the tool user had left), we see that social attention mostly occurred in combination with tolerated scrounging (tolerated scrounging occurred in 106 out of 183 sequences with social attention, compared to scrounging afterwards occurring in 13 sequences). Although sequences with other capuchins present were rarer at the EXP-ANV-01 site than CEBUS-02 (180 compared to 664), social attention was more likely at EXP-ANV-01 (Table S12). For example, given the most frequently occurring combination of a juvenile paying attention to a subadult tool user, the predicted likelihood of social attention at EXP-ANV-01 is

0.36 (95% CI [0.27-0.46] and at CEBUS-02 0.18 (95% CI [0.14, 0.23]). Paying social attention was not entirely without risks: in one tool use sequence a juvenile observer got too close to the tool user, and was hit on the foot with the hammerstone (Video S1).

Figure 6: Likelihood of paying social attention to a tool user, depending on number of capuchins present (left) and scrounging (right) in the sequence. Model estimates are reflected by the lines, and observed means from data are represented as points. Color of points reflects the size of the sample contributing to the mean.

To consider whether social attention is preferentially paid to more proficient tool users, we limited our sample to sequences where a) the item was successfully opened, b) we could confidently identify the tool user, and c) the tool user had more than 10 tool-using sequences in the sample (n = 1175 sequences). Our model estimated a slight trend of more social attention occurring in more efficient tool-using sequences (where fewer pounds needed to open the item), and this relationship was strongest if observers were juveniles (see Figure S2 and model estimates Table S13). Given a juvenile observer of a subadult tool user, the likelihood of the juvenile paying social attention was predicted to decrease with each additional pound the subadult tool user made. The individuals who received most social attention during tool use were the three subadult males, who also showed highest incidence of tool use with other individuals present (Figure 7). Individual subadults differed little from each other in how much social attention they received.

Figure 7: Likelihood of receiving social attention for different tool users. Model estimates are reflected by violin plots, and observed means from data are represented as points. Color reflects the age class of the tool user, and the numbers above the number of sequences with social attention (top) compared to the number of sequences with other capuchins present where no social attention occurred (bottom in italics).

Discussion

Stone tool use by white-faced capuchins on Jicarón is a complex behavior; proficiency takes years to develop. Compared to subadult and adult tool users, juvenile tool users need more time and more pounds to open an item using tools, more frequently reposition the item in between strikes, and make more mistakes. Subadults and adults show comparable levels of proficiency on all measures. Four identifiable tool-using juveniles did not markedly improve in tool use proficiency over the course of approximately one year. No other capuchins were present in the vast majority of tool use sequences and social attention to the tool user was rare (occurring in just 6% of all sequences). Nonetheless, we uncovered some robust patterns in who paid social attention to whom, and in what context. Juveniles were most likely to pay social attention to tool use, and most of these juveniles were too young to be tool users themselves. Social attention was directed at subadults and adults, with a slight preference towards more efficient tool users. Subadults received the most social attention. The more capuchins that were present during a tool-using sequence, the lower the probability that social attention occurred. Scrounging showed the opposite relationship: the more individuals scrounging, the higher the chance of social attention.

Differences in tool use proficiency

It is unsurprising that juveniles are less proficient tool users than adults and subadults, regardless of the measure of proficiency considered. This pattern is similar to what has previously been described in macaque stone tool use [20] and for this species of capuchin in other extractive foraging tasks [47]. However, juveniles were still very successful in opening items, and the difference in their efficiency compared with subadults and adults is small. Overall, white-faced capuchins had a slightly higher success rate than what is reported in studies of other nut-cracking primates. The item being abandoned entirely only occurred in 4% of all sequences, much lower than in chimpanzees (failure occurring in 8% of tool use events [48]) and robust capuchins (failure in 2-38% of events depending on population [11, 49, 50]). This does not necessarily reflect higher skill of the white-faced capuchins, but could also be an artifact of their high persistence and tenacity. It is important to note that we excluded tool use sequences that spanned more than one camera trap video. This was done to ensure no pounds or other behavior were missed between triggers, but also means we likely excluded the very worst tool-using sequences, where individuals took the longest to open the item, and some sequences were successful individuals rapidly opened many items in a row. As both of these variables are linked to age (juveniles are more likely to be less proficient and have sequences spanning multiple videos, while adults might process more rapidly and have multiple sequences in one video), their exclusion is not likely to significantly alter our results.

The clearest distinction between juvenile and (sub)adult tool use proficiency was the much higher incidence of mistakes by juveniles. Mistakes were rare in our sample overall, similar to bearded capuchins [31], but, contrary to their findings, we found that juveniles were considerably more likely to miss the item, or have the item fly off the anvil. Juveniles also more frequently repositioned the food item between strikes. More frequent repositioning by juveniles could reflect poorer control over the hammerstones than adults. This ties in with our surprising finding that at both sites, the same hammerstone was used in over 90% of the sequences, despite multiple other hammerstones being provided. These stones were large and heavy, and thus easier to handle for (sub)adults than juveniles. The (lack of) hammer selectivity relative to body size at the experimental anvil is an interesting avenue for future research, as it stands in stark contrast to findings in chimpanzees [51], macaques [52], and robust capuchins [53].

This may be related to the fact that large hammers might yield large dividends for ⁵⁰³ gracile capuchins, which are the smallest, lightest tool-using primates. ⁵⁰⁴

The same hammerstone may have been used so consistently at each site because there ⁵⁰⁵ was also little variation in the food items being opened. In nearly all sequences, ⁵⁰⁶ capuchins processed sea almonds. However, the level of ripeness of the sea almond does ⁵⁰⁷ appear to affect ease of processing: green, fresher sea almonds required more seconds ⁵⁰⁸ and more pounds to open than brown sea almonds, were peeled more frequently, and ⁵⁰⁹ were more likely to fly off the anvil when struck. This likely reflects the harder exocarp ⁵¹⁰ of the fruit at this stage, before it is dries out and becomes more brittle. The lack of ⁵¹¹ hammerstone selectivity for sea almond ripeness and higher incidence of peeling with ⁵¹² green sea almonds indicates that perhaps the additional step of peeling off the exocarp ⁵¹³ in between strikes helps overcome the difference in hardness, a technique which juveniles ⁵¹⁴ might also need to learn to become a more proficient tool user. However, juveniles also ⁵¹⁵ peeled more frequently than (sub)adults, so it is also possible that juveniles lack ⁵¹⁶ knowledge/ability to precisely strike the green sea almonds without them flying off the ⁵¹⁷ anvil – which green sea almonds are more likely to do — and that peeling the fruit ⁵¹⁸ makes it easier to strike. In either case, becoming a proficient stone tool user might not ⁵¹⁹ only involve learning *how* to process, but also *what* and *when*. Sea almonds have ⁵²⁰ variable fruiting seasons, and although we know little of seasonal fluctuations in food ⁵²¹ availability on Jicarón, our data does show sea almond processing peaking bimodally ⁵²² (Figure S1). Additionally, (sub)adults seem to focus more of their tool-using efforts at ⁵²³ specific times of the year (see Figure 4) whereas juveniles use tools at a more consistent ⁵²⁴ rate throughout the year. Future studies can elucidate the cause of these seasonal ⁵²⁵ fluctuations in tool use, and also explore whether adults indeed concentrate their ⁵²⁶ tool-using efforts when sea almonds are most abundant or, alternately, when other, ⁵²⁷ easier to consume food resources are unavailable. ⁵²⁸

Development of tool proficiency ⁵²⁹

The lack of change in tool use proficiency by juveniles over the course of our study ⁵³⁰ period suggests that honing these skills to adult levels of proficiency is likely a slow ⁵³¹ process, or one limited by physical features (like strength) which only mature in ⁵³² adulthood. White-faced capuchins are a long-lived species, with a lifespan up to 50 ⁵³³ years [5], similar to chimpanzees, where it has been shown that becoming a proficient ⁵³⁴ nut-cracker can take 3-7 years of practice [18, 54]. Our results, though not yet ⁵³⁵ sufficiently longitudinal for a full understanding of tool use ontogeny, suggests that in ⁵³⁶ white-faced capuchins, becoming a proficient stone tool user may be a similarly ⁵³⁷ extended process. Furthermore, our finding that the oldest tool-using male in the group, ⁵³⁸ Abraham, did not use tools at all during the 1 year period of this study, hints at a ⁵³⁹ potentially upper age limit for stone tool use in white-faced capuchins, as has been ⁵⁴⁰ documented in chimpanzees [32]. Abraham is the only individual to displace tool users ⁵⁴¹ from the anvil to steal the item they opened, which suggests he is still interested in the ⁵⁴² resources, but perhaps no longer able or willing to invest the energy and time to use ⁵⁴³ tools himself. Here, too, a more longitudinal approach would be very valuable to ⁵⁴⁴ examine the fluctuations in tool use proficiency across the lifespan. ⁵⁴⁵

The role of social attention in tool use acquisition ⁵⁴⁶

Stone tool use at the experimental anvils appeared to be a largely solitary activity. As ⁵⁴⁷ such, opportunities for social attention seem limited, and even when other capuchins ⁵⁴⁸ were present, social attention only occurred in 20% of the sequences. Data from camera ⁵⁴⁹ traps may underestimate social attention, as individuals out of frame could be looking ⁵⁵⁰

at the tool user without this being captured. However, attention from a distance likely only provides the observer with coarse information on the tool use behavior. The low frequency of close social attention in our sample is surprising compared to other nut-cracking primates. In one study on tufted capuchins, conspecifics observed 36% of all nut-cracking events [21], and in chimpanzees, juveniles observe hundreds of nut-cracking events from their mothers [28]. Yet on Jicarón, juvenile capuchins have no opportunity to learn tool use from their mothers, as females do not use tools [35]. Individuals paying social attention to tool use were mostly juveniles too young to be tool users themselves, suggesting that social observation may be most important for the acquisition of the tool use behavior [47]. Although scrounging was not as common as in tufted capuchins [21], the more individuals were scrounging, the higher the likelihood of social attention occurring. Furthermore, when scrounging did occur together with social attention, it was nearly always tolerated by the tool user. The tool users who received the most social attention, subadult males, are also known to be tolerant of juveniles [55]. Our results echo the idea that inter-individual tolerance plays a crucial role in social learning of complex behaviors [29]. Furthermore, capuchins might also rely on local enhancement and interaction with temporally enduring artifacts of tool use to learn about the tool use behavior [30]. In the absence of many opportunities for direct social learning, debris left on the anvil after a tool use sequence, such as partially processed food items, and wear on the hammerstone and anvil may provide important information to individuals visiting the site. Capuchins on Jicarón using this physical evidence of tool use as a cue could also explain the "burn-in period" at the novel anvil, EXP-ANV-01, since perhaps regular tool use by many individuals only started once enough debris had accumulated from sporadic tool use events, marking this location as a suitable anvil for tool use.

The importance of location

Here, we report on data collected from only two anvils in the tool-using groups' range. However, we have identified at least 10 different anvils that are habitually used for stone tool use, and over 300 ephemeral locations. While CEBUS-02 is the most frequently used anvil, which has, to the best of our knowledge, been in use for the longest time (at least since 2008), it is possible that most social learning of tool use occurs elsewhere, for instance during tool use in the intertidal zone [34]. We do see a difference in the likelihood of social attention between the two experimental anvils: CEBUS-02 has more opportunities for social attention, but social attention is more likely at EXP-ANV-01. This could be due to CEBUS-02 being an established, well-used anvil prior to the current study. We infer that this anvil is used so frequently because it is located in a productive foraging area (next to a stream), leading to more capuchins being present at the same time. In contrast, we purposefully placed the experimental anvil EXP-ANV-01 in a location without evidence of prior tool use, meaning this location likely held less interest for the capuchins. This inference is supported by the fact that it took capuchins taking several months to start regularly using this anvil. The availability of ripe sea almonds at this location may have also influenced the latency to capuchins visiting this anvil. Another, non-exclusive, possibility is that, either due to anvil material or other properties of the site, CEBUS-02 is simply a "better" anvil. In this case, there may be more competition over this site, and lower tolerance for observers. We found that on average, capuchins used more pounds at the stone anvil than the wooden anvil, and that items were more likely to fly off the stone than the wooden anvil, further supporting that the wooden anvil at CEBUS-02 has better properties for tool use. Anvil selectivity by tool users and the notion of "wear in" time of new anvils is another interesting avenue for future research, in order to examine what exact physical properties drive these differences in tool use efficiency.

Spread of tool use to other groups

The seemingly slow mastery of stone tool use by white-faced capuchins also has implications for the spread of this behavior. Despite being persistent over time, the stone tool use behavior on Jicarón is incredibly localized [14]. In mainland populations of white-faced capuchins, males are the dispersing sex and first dispersal occurs on average at 4.5 years of age in Santa Rosa [56] and 7 years of age in Lomas Barbudal [57]. On Jicarón, males of this age would not yet be proficient tool users based on our data, and as such, might cease using tools post-dispersal, limiting the spread of the behavior. However, it is unclear whether the typical pattern of male dispersal holds on Jicarón, or if, instead, both sexes, or only females disperse. Switches in dispersal tendency are known to occur under high density conditions [58]. It appears likely that at least some males remain in the natal group, as we have tracked some individuals from juvenile to (sub)adulthood. Further elucidating the dispersal tendency of these capuchins is crucial to better understand the maintenance and spread of the tool use behavior.

Conclusion

We present a first description of variation in and development of stone tool use proficiency in white-faced capuchin monkeys. Using camera traps placed at two experimental tool use anvils on Jicarón island, Panama, we collected nearly a year of tool-using sequences, mostly of capuchins processing sea almonds. Similar to other primates that show percussive stone tool use, white-faced capuchin juveniles are less proficient tool users than adults (in terms of efficiency and technique), and development of proficiency appears to be a slow process. Ripeness of the sea almond affected processing times and efficiency, with less ripe green sea almonds requiring more time and peeling of the skin in between pounds, and generating more mistakes. In contrast to findings from other tool-using primates, tool-using appeared to be a largely solitary activity, with limited opportunities for social attention and even fewer actual occurrences of social attention. However, when social attention to tool use did occur, there were clear patterns to it that echoed studies in other primates. Social attention was mostly paid by juveniles to proficient subadults, who were tolerant of scrounging. We also found an effect of location on the likelihood of social attention occurring, suggesting social learning opportunities differ between sites. In short, we lay the groundwork for both comparisons between stone tool-using primate species, as well as future research on other aspects of white-faced capuchin stone tool use like seasonality and hammerstone and anvil selectivity.

Acknowledgments

We are grateful to Evelyn del Rosario-Vargas, Pedro Luis Castillo-Caballero, Eliecer Vega-Patiño, Juan Rojas Garrido, James Chaves, and Tamara Dogandžić for their assistance with fieldwork, and STRI Panama and the Coiba research station for making this research possible. We further thank Lucia Torrez, Katrin Dieter, and Angie Ruiz for their logistical support.

Ethics statement

Data for this study were collected as minimally invasively as possible. We obtained permission for this study from the relevant authority Ministerio de Ambiente, Panama

Supporting information

Video ethogram: https://keeper.mpdl.mpg.de/d/0c1b9853f3f342d8b3da/

Video S1: https://youtu.be/tvuClVgwLRQ

Table S1: Behavioral ethogram of all behaviors coded for every tool use sequence.

Table S2: Overview of sequence-level variables coded for every tool use sequence.

Table S3: Behavioral ethogram of behaviors coded per sequence with capuchins present.

Figure S1: Number of tool use sequences observed per experimental anvil.

Table S4: Posterior mean model estimates of Model e1.

Table S5: Posterior mean model estimates of Model e2.

Table S6: Posterior mean model estimates of Model e2b.

Table S7: Posterior mean model estimates of Model e3a.

Table S8: Posterior mean model estimates of Model e3b.

Table S9: Posterior mean model estimates of Model e4a.

Table S10: Posterior mean model estimates of Model e4b.

Table S11: Posterior mean model estimates of Model dev1.

Table S12: Posterior mean model estimates of Model socatt1.

Table S13: Posterior mean model estimates of Model socatt1b.

Figure S2: Predicted probabilities from model socatt1b of probabilities of social attention occurring, depending on the number of pounds in the sequence.

References

1. Biro, D., Haslam, M. & Rutz, C. Tool use as adaptation. *Philosophical Transactions of the Royal Society B: Biological Sciences* **368.** Publisher: Royal Society, 20120408. doi:10.1098/rstb.2012.0408 (Nov. 2013).

2. Boesch, C., Bombjaková, D., Meier, A. & Mundry, R. Learning curves and teaching when acquiring nut-cracking in humans and chimpanzees. en. *Scientific Reports* **9.** Publisher: Nature Publishing Group, 1515. issn: 2045-2322. doi:10.1038/s41598-018-38392-8 (Feb. 2019).

3. Shumaker, R. W., Walkup, K. R., Beck, B. B. & Burghardt, G. M. *Animal Tool Behavior: The Use and Manufacture of Tools by Animals* revised and updated edition. English. isbn: 978-0-8018-9853-2 (Johns Hopkins University Press, Baltimore, May 2011).

4. Beck, B. B. *Animal Tool Behaviour: The Use and Manufacture of Tools by Animals* (Garland STM Pub., New York, NY, 1980).

5. Fragaszy, D. M., Visalberghi, E. & Fedigan, L. M. *The Complete Capuchin: The Biology of the Genus Cebus* en. Google-Books-ID: A6TmtS_qOwgC. isbn: 978-0-521-66768-5 (Cambridge University Press, June 2004).

6. De la Torre, I. & Hirata, S. Percussive technology in human evolution: an introduction to a comparative approach in fossil and living primates. *Philosophical Transactions of the Royal Society B: Biological Sciences* **370,** 20140346. issn: 0962-8436. doi:10.1098/rstb.2014.0346 (Nov. 2015).

7. Sugiyama, Y. & Koman, J. Tool-Using and -Making Behavior in Wild Chimpanzees at Bossou, Guinea. *Primates* **20,** 513–524 (1979).

8. Gumert, M. D. & Malaivijitnond, S. Marine prey processed with stone tools by burmese long-tailed macaques (Macaca fascicularis aurea) in intertidal habitats. en. *American Journal of Physical Anthropology* **149,** 447–457. issn: 1096-8644. doi:https://doi.org/10.1002/ajpa.22143 (2012).

9. Muhammad, R. *et al.* Influence of COVID-19 on the emergence of stone-tool use behavior in a population of common long-tailed macaques (Macaca fascicularis fascicularis) in Thailand. en. *American Journal of Primatology* **86,** e23580. issn: 1098-2345. doi:10.1002/ajp.23580 (2024).

10. Ottoni, E. B. & Mannu, M. Semifree-ranging Tufted Capuchins (Cebus apella) Spontaneously Use Tools to Crack Open Nuts. en. *International Journal of Primatology,* 12 (2001).

11. Spagnoletti, N., Visalberghi, E., Ottoni, E., Izar, P. & Fragaszy, D. Stone tool use by adult wild bearded capuchin monkeys (Cebus libidinosus). Frequency, efficiency and tool selectivity. *Journal of Human Evolution* **61,** 97–107. doi:10.1016/j.jhevol.2011.02.010 (2011).

12. Lima, G. C. B. *et al.* A new addition to the toolbox: stone tool use in blonde capuchin monkeys (Sapajus flavius). en. *Primates.* issn: 1610-7365. doi:10.1007/s10329-024-01143-7 (June 2024).

13. Canale, G. R., Guidorizzi, C. E., Kierulff, M. C. M. & Gatto, C. A. F. R. First record of tool use by wild populations of the yellow-breasted capuchin monkey (Cebus xanthosternos) and new records for the bearded capuchin (Cebus libidinosus). eng. *American Journal of Primatology* **71,** 366–372. issn: 1098-2345. doi:10.1002/ajp.20648 (May 2009).

14. Barrett, B. J., Monteza-Moreno, C. M., Dogandžić, T., Ibáñez, A. & Crofoot, M. C. Habitual stone-tool-aided extractive foraging in white-faced capuchins, Cebus capucinus. en. *Royal Society Open Science* **5,** 181002. doi:https://doi.org/10.1098/rsos.181002 (2018).

15. Araujo, P. E. *et al.* First record of tool use by a wild population of Cebus albifrons (Humboldt, 1812) (Primates, Cebidae) in Puerto Misahuallí, Napo, Ecuador. en. *Boletín Técnico. Serie Zoológica* **15.** Number: 16, 59–72. ISSN: 1390-3004 (2021).

16. Biro, D. *et al.* Cultural innovation and transmission of tool use in wild chimpanzees: evidence from field experiments. eng. *Animal Cognition* **6,** 213–223. ISSN: 1435-9448. doi:10.1007/s10071-003-0183-x (Dec. 2003).

17. Inoue-Nakamura, N. & Matsuzawa, T. Development of stone tool use by wild chimpanzees (Pan troglodytes). *Journal of Comparative Psychology* **111.** Place: US Publisher: American Psychological Association, 159–173. ISSN: 1939-2087. doi:10.1037/0735-7036.111.2.159 (1997).

18. Matsuzawa, T. in *Chimpanzee cultures* 351–370 (Harvard University Press, Cambridge, MA, US, 1994). ISBN: 978-0-674-11662-7.

19. De Resende, B. D., Ottoni, E. B. & Fragaszy, D. M. Ontogeny of manipulative behavior and nut-cracking in young tufted capuchin monkeys (*Cebus apella*): a Perception–action perspective. en. *Developmental Science* **11,** 828–840. ISSN: 1363-755X, 1467-7687. doi:10.1111/j.1467-7687.2008.00731.x (Nov. 2008).

20. Tan, A. W. Y. From play to proficiency: The ontogeny of stone-tool use in coastal-foraging long-tailed macaques (Macaca fascicularis) from a comparative perception-action perspective. *Journal of Comparative Psychology* **131.** Place: US Publisher: American Psychological Association, 89–114. ISSN: 1939-2087. doi:10.1037/com0000068 (2017).

21. Coelho, C. G. *et al.* Social learning strategies for nut-cracking by tufted capuchin monkeys (Sapajus spp.) en. *Animal Cognition* **18,** 911–919. ISSN: 1435-9448, 1435-9456. doi:10.1007/s10071-015-0861-5 (July 2015).

22. Ottoni, E. B., de Resende, B. D. & Izar, P. Watching the best nutcrackers: what capuchin monkeys (Cebus apella) know about others' tool-using skills. eng. *Animal Cognition* **8,** 215–219. ISSN: 1435-9448. doi:10.1007/s10071-004-0245-8 (Oct. 2005).

23. Eshchar, Y., Izar, P., Visalberghi, E., Resende, B. & Fragaszy, D. When and where to practice: social influences on the development of nut-cracking in bearded capuchins (Sapajus libidinosus). en. *Animal Cognition* **19,** 605–618. ISSN: 1435-9456. doi:10.1007/s10071-016-0965-6 (May 2016).

24. Neadle, D., Bandini, E. & Tennie, C. Testing the individual and social learning abilities of task-naïve captive chimpanzees (Pan troglodytes sp.) in a nut-cracking task. eng. *PeerJ* **8,** e8734. ISSN: 2167-8359. doi:10.7717/peerj.8734 (2020).

25. Koops, K., Soumah, A. G., van Leeuwen, K. L., Camara, H. D. & Matsuzawa, T. Field experiments find no evidence that chimpanzee nut cracking can be independently innovated. eng. *Nature Human Behaviour* **6,** 487–494. ISSN: 2397-3374. doi:10.1038/s41562-021-01272-9 (Apr. 2022).

26. Boesch, C. Teaching among wild chimpanzees. *Animal Behaviour* **41,** 530–532. ISSN: 0003-3472. doi:10.1016/S0003-3472(05)80857-7 (Mar. 1991).

27. Lonsdorf, E. V. Sex differences in the development of termite-fishing skills in the wild chimpanzees, *Pan troglodytes schweinfurthii*, of Gombe National Park, Tanzania. *Animal Behaviour* **70,** 673–683. ISSN: 0003-3472. doi:10.1016/j.anbehav.2004.12.014 (Sept. 2005).

28. Estienne, V., Cohen, H., Wittig, R. M. & Boesch, C. Maternal influence on the development of nut-cracking skills in the chimpanzees of the Taï forest, Côte d'Ivoire (Pan troglodytes verus). en. *American Journal of Primatology* **81,** e23022. ISSN: 1098-2345. doi:`10.1002/ajp.23022` (2019).

29. Van Schaik, C. P., Deaner, R. O. & Merrill, M. Y. The conditions for tool use in primates: implications for the evolution of material culture. en. *Journal of Human Evolution* **36,** 719–741. ISSN: 0047-2484. doi:`10.1006/jhev.1999.0304` (June 1999).

30. Fragaszy, D. M. *et al.* The fourth dimension of tool use: temporally enduring artefacts aid primates learning to use tools. *Philosophical Transactions of the Royal Society B: Biological Sciences* **368.** Publisher: Royal Society, 20120410. doi:`10.1098/rstb.2012.0410` (Nov. 2013).

31. Fragaszy, D. M. *et al.* Adult and juvenile bearded capuchin monkeys handle stone hammers differently during nut-cracking. en. *American Journal of Primatology* **82,** e23156. ISSN: 1098-2345. doi:`10.1002/ajp.23156` (2020).

32. Howard-Spink, E. *et al.* *Old age variably impacts chimpanzee engagement and efficiency in stone tool use* en. Pages: 2024.11.25.625128 Section: New Results. Dec. 2024. doi:`10.1101/2024.11.25.625128`.

33. Monteza-Moreno, C. M. *et al.* White-Faced Capuchin, Cebus capucinus imitator, Hammerstone and Anvil Tool Use in Riparian Habitats on Coiba Island, Panama. en. *International Journal of Primatology* **41,** 429–433. ISSN: 0164-0291, 1573-8604. doi:`10.1007/s10764-020-00156-5` (June 2020).

34. Goldsborough, Z. *et al.* Coupling of Coastal Activity with Tidal Cycles is Stronger in Tool-using Capuchins (Cebus capucinus imitator). *Royal Society Open Science* **10,** 230355. doi:`https://doi.org/10.1098/rsos.230355` (Sept. 2023).

35. Goldsborough, Z., Crofoot, M. C. & Barrett, B. J. Male-biased stone tool use by wild white-faced capuchins (Cebus capucinus imitator). *American Journal of Primatology* **86,** e23594. doi:`10.1002/ajp.23594` (2024).

36. Titcomb, M. & O'Dea, A. *Post-glacial Sea Level rise on the Isthmus of Panama* 2020.

37. Isaza, I. & Vrba, E. *Ocupación pre-colombina de las Islas del Parque Nacional Coiba. Panama: Informe final del Proyecto PRB08-003 archivos SENACYT* tech. rep. (2010).

38. Ibáñez, A., Pérez-Jordá, J., Juste, J. & Guillén, A. in. ed. S Castroviejo, 469–486 (Agencia Española de Cooperacíon Internacional, 1997).

39. Monteza-Moreno, C. M., Crofoot, M. C., Grote, M. N. & Jansen, P. A. Increased terrestriality in a Neotropical primate living on islands with reduced predation risk. en. *Journal of Human Evolution* **143,** 102768. ISSN: 00472484. doi:`10.1016/j.jhevol.2020.102768` (June 2020).

40. Milton, K. & Mittermeier, R. A. A brief survey of the primates of Coiba Island, Panama. en. *Primates* **18,** 931–936. ISSN: 0032-8332, 1610-7365. doi:`10.1007/BF02382942` (Oct. 1977).

41. Friard, O. & Gamba, M. BORIS: a free, versatile open-source event-logging software for video/audio coding and live observations. en. *Methods in Ecology and Evolution* **7,** 1325–1330. ISSN: 2041-210X. doi:`10.1111/2041-210X.12584` (2016).

42. Team, R. C. *R: A language and environment for statistical computing* Vienna, Austria, 2022.

43. Bürkner, P.-C. brms: An R Package for Bayesian Multilevel Models Using Stan. en. *Journal of Statistical Software* **80,** 1–28. ISSN: 1548-7660. doi:10.18637/jss.v080.i01 (Aug. 2017).

44. Perez, R. & Condit, R. *Tree Atlas of Panama* (Center for Tropical Forest Science, Smithsonian Tropical Research Institute).

45. Vehtari, A., Gelman, A., Simpson, D., Carpenter, B. & Bürkner, P.-C. Rank-normalization, folding, and localization: An improved $\widehat{R}$ for assessing convergence of MCMC. *Bayesian Analysis* **16.** arXiv:1903.08008 [stat]. ISSN: 1936-0975. doi:10.1214/20-BA1221 (June 2021).

46. Goldsborough, Z. *ZoeGold/tooluseproficiency: Data and Code Tool Use Proficiency Pre-print* Apr. 2025. doi:10.5281/zenodo.15173297.

47. Barrett, B. J., McElreath, R. L. & Perry, S. E. Pay-off-biased social learning underlies the diffusion of novel extractive foraging traditions in a wild primate. *Proceedings of the Royal Society B: Biological Sciences* **284.** Publisher: Royal Society, 20170358. doi:10.1098/rspb.2017.0358 (June 2017).

48. Berdugo, S., Cohen, E., Davis, A. J., Matsuzawa, T. & Carvalho, S. Reliable long-term individual variation in wild chimpanzee technological efficiency. en. *Nature Human Behaviour.* Publisher: Nature Publishing Group, 1–9. ISSN: 2397-3374. doi:10.1038/s41562-024-02071-8 (Dec. 2024).

49. Falótico, T. & Ottoni, E. B. The manifold use of pounding stone tools by wild capuchin monkeys of Serra da Capivara National Park, Brazil. en. Publisher: Brill. doi:10.1163/1568539X-00003357 (Jan. 2016).

50. Falótico, T., Macedo, A. C., de Jesus, M. A., Espinola, T. & Valença, T. Nut-cracking success and efficiency in two wild capuchin monkey populations. *Royal Society Open Science* **11.** Publisher: Royal Society, 240161. doi:10.1098/rsos.240161 (June 2024).

51. Boesch, C. & Boesch, H. Optimisation of Nut-Cracking With Natural Hammers By Wild Chimpanzees. en. Publisher: Brill. doi:10.1163/156853983X00192 (Jan. 1983).

52. Gumert, M. D. & Malaivijitnond, S. Long-tailed macaques select mass of stone tools according to food type. *Philosophical Transactions of the Royal Society B: Biological Sciences* **368,** 20120413. ISSN: 0962-8436. doi:10.1098/rstb.2012.0413 (Nov. 2013).

53. Visalberghi, E. *et al.* Selection of Effective Stone Tools by Wild Bearded Capuchin Monkeys. *Current Biology* **19,** 213–217. ISSN: 0960-9822. doi:10.1016/j.cub.2008.11.064 (Feb. 2009).

54. Boesch, C. & Boesch, H. Tool Use and Tool Making in Wild Chimpanzees. en. Publisher: Brill. doi:10.1159/000156428 (Feb. 1990).

55. Jack, K. M. & Riley, E. P. Male social tolerance, cooperation, and affiliation in male dispersing primates. *Behaviour* **151.** Publisher: Brill, 861–870. ISSN: 0005-7959 (2014).

56. Jack, K. M., Sheller, C. & Fedigan, L. M. Social factors influencing natal dispersal in male white-faced capuchins (Cebus capucinus). en. *American Journal of Primatology* **74,** 359–365. ISSN: 1098-2345. doi:10.1002/ajp.20974 (2012).

57. Perry, S., Godoy, I. & Lammers, W. en. in *Long-Term Field Studies of Primates* (eds Kappeler, P. M. & Watts, D. P.) 141–163 (Springer, Berlin, Heidelberg, 2012). ISBN: 978-3-642-22514-7. doi:10.1007/978-3-642-22514-7_7.

58. Clutton-Brock, T. H. & Lukas, D. The evolution of social philopatry and dispersal in female mammals. en. *Molecular Ecology* **21,** 472–492. ISSN: 1365-294X. doi:https://doi.org/10.1111/j.1365-294X.2011.05232.x (2012).