HCV infection induces ubiquitin-dependent degradation of LATS1, inactivating the

Hippo pathway and upregulating transcription of the CYR61 and CTGF genes

Maria Alethea Septianastiti [1, 2], Chieko Matsui [1], Zihan Xu [1], Fransisca Puspitasari [1], Lin

Deng [1], Takayuki Abe [3], Ikuo Shoji [1, #]

[1] Division of Infectious Disease Control, Center for Infectious Diseases, Kobe University

Graduate School of Medicine, Kobe, Japan

[2] Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta,

Indonesia

[3] Division of Virology, Niigata University Graduate School of Medical and Dental Sciences,

Niigata, Japan

Running Title: HCV inactivates Hippo via LATS1 degradation

Correspondence: Prof. Ikuo Shoji

E-mail: ishoji@med.kobe-u.ac.jp

Address: Division of Infectious Disease Control, Center for Infectious Diseases, Kobe

University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe, Japan.

Tel.: +81-78-382-5500; Fax: +81-78-382-5519.

Word count

Abstract: 226 words

Importance: 117 words

Text: 3377

Figure: 7

Keywords: HCV, LATS1, Itch, ubiquitin, Hippo pathway, YAP1

ABSTRACT

Hepatitis C virus (HCV) is a major cause of hepatocellular carcinoma (HCC). However, the mechanisms underlying HCV-related hepatocarcinogenesis are not fully understood. The Hippo signaling pathway plays a crucial role in tumor suppression. Here, we investigated the role of the Hippo pathway in HCV-related pathogenesis. We demonstrated that HCV infection induces degradation of LATS1, a key regulator of the Hippo pathway. LATS1 degradation was reversed by a proteasomal inhibitor, but not by lysosomal inhibitors, indicating that HCV promotes proteasomal degradation of LATS1. Cell-based ubiquitylation assays revealed that HCV infection induces K6- and K11-linked polyubiquitylation of LATS1. HCV-induced degradation of LATS1 was suppressed in si-Itch-transfected Huh-7.5 cells. Moreover, Itch overexpression enhanced LATS1 polyubiquitylation in HCV-infected Huh-7.5 cells. These results suggest that the Itch ubiquitin ligase is involved in LATS1 degradation. The JNK inhibitor SP600125 abolished LATS1 polyubiquitylation, suggesting that the HCV-induced ROS/JNK/Itch pathway promotes ubiquitin-dependent proteasomal degradation of LATS1. Moreover, cell fractionation assays and immunofluorescence staining revealed that HCV infection promotes nuclear translocation of YAP1 protein, suggesting suppression of the Hippo pathway. Additionally, expression of the YAP1 target genes CYR61 and CTGF, which are involved in tissue remodeling and proliferation, was upregulated in HCV-infected Huh-7.5 cells. Taken together, these findings suggest that the HCV-induced ROS/JNK/Itch pathway promotes the ubiquitin-dependent proteasomal degradation of LATS1 protein, leading to Hippo pathway suppression and increased expression of CYR61 and CTGF.

IMPORTANCE

HCV infection induces ubiquitin-dependent degradation of LATS1, leading to inactivation of the Hippo pathway and upregulation of the YAP1 target genes CYR61 and CTGF. We provide evidence that HCV infection promotes the Itch-mediated, ubiquitin-dependent degradation of LATS1, a key regulator of the Hippo pathway. HCV infection induces K6- and K11-linked polyubiquitylation of LATS1, leading its degradation. HCV-induced degradation of LATS1 promotes Hippo pathway inactivation and nuclear translocation of YAP1, thereby upregulating expression of the CYR61 and CTGF genes. We propose a novel molecular mechanism by which HCV infection induces activation of Itch and promotes LATS1 degradation, resulting in Hippo pathway inactivation. Understanding HCV-induced Hippo pathway inactivation may lead to new strategies for preventing or treating HCV-related disease.

INTRODUCTION

Hepatitis C virus (HCV) infection remains a global health burden, affecting an estimated 50 million people worldwide as of 2022 (1). HCV infection causes acute and chronic liver diseases, ranging from mild illness to severe complications, including liver cirrhosis and hepatocellular carcinoma (HCC). Although highly effective direct-acting antivirals (DAAs) are available for HCV treatment, the risk of HCC persists even after successful viral clearance (2). Notably, HCC can occur following a sustained virological response, at an annual incidence of approximately 1%, surpassing that of other organ cancers (3).

HCV is a positive-sense, single-stranded RNA virus classified in the *Flaviviridae* family, *Hepacivirus* genus. The 9.6-kb HCV RNA genome encodes a single polyprotein of approximately 3,010 amino acids, which is cleaved by viral proteases and host signalases into structural proteins (Core, E1, and E2) and nonstructural proteins (p7, NS2, NS3, NS4A, NS5A, and NS5B) (4). As HCV infection is a major cause of HCC, understanding HCV-induced modulation of cellular signaling pathways is crucial for elucidating the mechanism underlying HCV-induced hepatocarcinogenesis.

The Hippo pathway, a key regulator of cell proliferation, apoptosis, and organ size control, plays an essential role in tumor suppression. Dysregulation of the Hippo pathway is implicated in the development of several cancers, including liver cancer (5–7). The Hippo pathway involves a series of proteins that form a kinase cascade, including MST1/2, LATS1/2, the adaptor protein SAV1, MOB1A/B, and the transcription coactivators YAP1 and TEAD transcription factors. Activation of the Hippo pathway involves the phosphorylation of the Serine/Threonine kinase LATS1, which in turn phosphorylates YAP1 at Ser127 and promotes the binding of phosphorylated YAP1 to 14-3-3 proteins (8, 9). The association of 14-3-3 proteins with phosphorylated YAP1 prevents it from entering the nucleus. Conversely,

when the Hippo pathway is inactivated, unphosphorylated YAP1 translocates to the nucleus and induces the transcription of genes responsible for cell growth (10,11).

Several viruses have been shown to modulate the Hippo pathway (12–15). HCV E2 activates the Hippo pathway by interacting with CD81, while HCV NS4B suppresses the Hippo pathway via Scribble, promoting PI3K/AKT signaling and epithelial-mesenchymal transition (EMT) (16,17). Similarly, Kaposi sarcoma-associated herpesvirus (KSHV) vGPCR inhibits LATS1/2, enabling YAP1 nuclear translocation. Human papillomavirus 16 E6 alters YAP1 localization through interaction with PDZ domains (13,15).

We previously reported that HCV infection activates the ROS/JNK signaling pathway which in turn activates Itch, a HECT-type E3 ubiquitin ligase. Activated Itch promotes polyubiquitylation of VPS4A, resulting in efficient HCV particle release (18). Notably, Itch has been shown to promote the proteasomal degradation of LATS1, a critical component of the Hippo pathway (19,20).

In this study, we investigate the Hippo pathway's role in HCV-related pathogenesis. We demonstrate that HCV infection promotes the ubiquitin-dependent degradation of LATS1. We also demonstrate that Itch plays an important role in HCV-induced polyubiquitylation of LATS1, leading to Hippo pathway suppression and thereby contributing to HCV pathogenesis.

RESULTS

HCV promotes proteasomal degradation of LATS1 protein

To investigate the Hippo signaling pathway's role in HCV-related pathogenesis, we examined LATS1 protein levels in HCV J6/JFH1-infected Huh-7.5 cells. Immunoblot analysis revealed a reduction in LATS1 protein levels in HCV-infected cells at both 2and 4 days post-infection (Fig. 1A, first panel, lanes 2 and 4). However, reverse-transcription quantitative polymerase

chain reaction (RT-qPCR) analysis showed increased LATS1 mRNA levels (Fig. 1B),

indicating that HCV-induced reduction of LATS1 protein levels is not due to suppression of

LATS1 mRNA expression.

To further elucidate the mechanism underlying the HCV-induced reduction of LATS1,

we treated HCV-infected Huh-7.5 cells with two lysosomal inhibitors: NH_4Cl and

Bafilomycin A1. However, neither inhibitor restored LATS1 protein levels in HCV-infected

cells (Fig. 1C, first panel, lane 6; Fig. 1D, first panel, lane 6). Moreover, small interfering

RNA (siRNA)-mediated knockdown of LAMP2A, a key protein involved in chaperone-

mediated autophagy, also failed to restore LATS1 protein levels in HCV-infected cells (Fig.

1E, first panel, lane 4). These results suggest that HCV induces LATS1 reduction through a

pathway independent of lysosomal degradation.

To determine whether HCV infection promotes proteasomal degradation of LATS1,

we treated the cells with the proteasome inhibitor clasto-lactacystin β-lactone. Immunoblot

analysis revealed that clasto-lactacystin β-lactone restored LATS1 protein levels in HCV-

infected cells (Fig. 1F, first panel, lane 6). These results suggest that HCV infection promotes

proteasomal degradation of LATS1.

HCV induces K6- and K11-linked polyubiquitylation of LATS1

To investigate whether HCV infection promotes polyubiquitylation of LATS1, we

performed a cell-based ubiquitylation assay. Huh-7.5 cells were infected with HCV J6/JFH1

and subsequently transfected with pCAG-FLAG−LATS1 along with either pRK5-HA−

ubiquitin (Ub) wild-type or pRK5-HA−Ub mutants containing only a single lysine residue at

K6, K11, K27, K29, K33, K48, or K63. The cell-based ubiquitylation assays showed that

LATS1 was polyubiquitylated wild-type Ub, K6−Ub, and K11−Ub (Fig. 2, first panel, lanes

11, 12, and 13). These findings suggest that HCV infection promotes K6- and K11-linked polyubiquitylation of LATS1.

E3 ligase Itch, but not WWP1, plays a crucial role in HCV-induced polyubiquitylation and degradation of LATS1

To determine whether HCV-induced LATS1 polyubiquitylation depends specifically on Itch ubiquitin ligase, we generated an inactive mutant Itch by mutating cysteine 868 to alanine (pCAG-FLAG-Itch C868A), which abolishes E3 ligase activity. Huh-7.5 cells were co-transfected with pCAG-Myc-LATS1, pRK5-HA−Ub, and either pCAG-FLAG-Itch or pCAG-FLAG-Itch C868A. Immunoblot analysis revealed that LATS1 polyubiquitylation was increased upon co-transfection with pCAG-FLAG-Itch (Fig. 3A, first panel, lane 2) but not with the inactive mutant (Fig. 3A, first panel, lane 3). These results suggest that Itch mediates the polyubiquitylation of LATS1.

To determine whether other E3 ubiquitin ligases in the HECT family, such as WWP1, also play a role in the polyubiquitylation of LATS1, we performed a cell-based ubiquitylation assay using wild-type pCAG-FLAG−WWP1 and the inactive mutant pCAG-FLAG−WWP1 C890A. Immunoblot analysis showed that LATS1 ubiquitylation was increased upon co-transfection with wild-type pCAG-FLAG-WWP1 (Fig. 3B, first panel, lane 2), while the inactive mutant abolishes this effect (Fig. 3B, first panel, lane 3). Next, we examined whether HCV-induced LATS1 ubiquitylation is mediated by Itch or WWP1 in HCV-infected cells. Mock- and HCV-infected Huh-7.5 cells were co-transfected with pRK5-HA−Ub, pCAG-Myc-LATS1, and either pCAG-FLAG-Itch or pCAG-FLAG−WWP1. The cell-based ubiquitylation assay revealed that HCV infection enhanced LATS1 polyubiquitylation when Itch was overexpressed compared to when WWP1 was overexpressed (Fig. 3C, first panel, lanes 5 and 6).

To verify Itch's role in HCV-induced LATS1 polyubiquitylation and degradation, we

performed siRNA-mediated knockdown of either Itch or WWP1 in HCV-infected cells.

Immunoblot analysis demonstrated that knockdown of Itch restored LATS1 protein levels

(Fig. 3D, first panel, lanes 2 and 4), whereas knockdown of WWP1 further decreased LATS1

protein levels (Fig. 3E, first panel, lanes 2 and 4). These findings suggest that Itch is

responsible for HCV-induced ubiquitin-dependent degradation of LATS1.

HCV-induced ROS/JNK/Itch signaling pathway plays a role in polyubiquitylation of

LATS1

To further determine whether the HCV-induced ROS/JNK/Itch signaling pathway is

involved in the polyubiquitylation of LATS1, we treated cells with the JNK inhibitor

SP600125 and assayed for LATS1 polyubiquitylation. Immunoblot analysis revealed that the

JNK inhibitor SP600125 disrupted HCV-induced Itch-mediated LATS1 polyubiquitylation

(Fig. 4, first panel, lane 6). These results suggest that the HCV-induced ROS/JNK/Itch

pathway promotes polyubiquitylation of LATS1.

HCV infection promotes translocation of YAP1 from the cytoplasm to the nucleus.

LATS1 is the protein kinase that phosphorylates YAP1 at Ser127 and activates the Hippo

signaling pathway. Therefore, we hypothesized that HCV-induced degradation of LATS1

may suppress the Hippo pathway. To assess this hypothesis, we performed immunoblot

analysis of components of the Hippo pathway, LATS1 and YAP1, in mock- and HCV-

infected Huh-7.5 cells. Immunoblot analysis revealed that HCV infection promoted LATS1

degradation (Fig. 5A, first panel, lane 2), accompanied by a reduction in phosphorylated

LATS1 at Ser 909 (Fig. 5A, second panel, lane 2), suggesting that active LATS1 is decreased

in HCV-infected cells. The decrease in pLATS1 at Ser 909 subsequently resulted in reduced

YAP1 phosphorylation at Ser 127 (Fig. 5A, third panel, lane 2), whereas total YAP1 levels

remained unchanged (Fig. 5A, fourth panel, lane 2). These results suggest that HCV infection

reduced LATS1 and phosphorylated LATS1 at S909, resulting in decreased YAP1

phosphorylation at Ser 127, thereby inactivating the Hippo pathway.

To determine whether unphosphorylated YAP1 translocates to the nucleus, we

performed a cell fractionation assay coupled with immunoblot analysis. The assay revealed

an increased nuclear fraction of YAP1 in HCV-infected cells (Fig. 5B, first panel, lane 4)

compared to mock-infected cells (Fig. 5B, first panel, lane 3). Furthermore,

immunofluorescence staining revealed that YAP1 was predominantly localized to the nucleus

in HCV-infected Huh-7.5 cells (Fig. 5C, second panel), whereas YAP1 was more

cytoplasmically distributed in mock-infected cells (Fig. 5C, first panel). These results suggest

that HCV infection promotes LATS1 degradation, leading to reduced YAP1 phosphorylation

and facilitating YAP1 nuclear translocation, thereby inactivating the Hippo pathway.

mRNA levels of YAP1 target genes are upregulated upon HCV infection

To determine whether HCV-induced nuclear translocation of YAP1 affects downstream gene

expression, we performed RT-qPCR to measure the mRNA levels of canonical YAP1 target

genes, CYR61 and CTGF, in HCV-infected cells. RT-qPCR showed that HCV infection

significantly increased the mRNA levels of CYR61 and CTGF (Fig. 6A and 6B).

To determine whether HCV-induced upregulation of CYR61 and CTGF depends

specifically on YAP1 nuclear translocation, we used siRNA to knock down YAP1 expression

in HCV-infected Huh-7.5 cells. YAP1 mRNA levels were successfully reduced by

approximately 80% (Fig. 6C). Under HCV-infected conditions, the relative mRNA levels of

CYR61 and CTGF were significantly lower in siYAP1-transfected cells (Fig. 6D and 6E).

These results suggest that HCV promotes the upregulation of CYR61 and CTGF via YAP1

nuclear translocation. CYR61 and CTGF are known to play roles in tissue remodeling and

cell proliferation, suggesting that the HCV-induced upregulation of CYR61 and CTGF genes

may contribute to HCV-related pathogenesis.

218   In summary, we propose the model shown in Fig. 7. Under normal conditions, the

Hippo pathway remains active when LATS1 phosphorylates YAP1, and phosphorylated

YAP1 is retained in the cytoplasm. In HCV infection, HCV activates the ROS/JNK/Itch

pathway. Active Itch induces the polyubiquitylation and proteasomal degradation of LATS1.

The degradation of LATS1 inhibits YAP1 phosphorylation and promotes YAP1 nuclear

translocation, thereby upregulating transcription of the YAP1 target genes, CYR61 and

CTGF. HCV-induced suppression of the Hippo pathway may play a role in HCV

pathogenesis.

DISCUSSION

In this study, we found that HCV infection induced the degradation of LATS1 protein, a

serine/threonine kinase essential for regulating the tumor-suppressive Hippo pathway (Fig.

1). We demonstrated that HCV infection promoted polyubiquitylation and proteasomal

degradation of LATS1 (Fig. 2). In particular, our results showed that HCV infection induced

K6- and K11-linked polyubiquitylation of LATS1. We also demonstrated that HCV infection

promoted degradation of LATS1, which inactivated the Hippo pathway and facilitated YAP1

nuclear translocation (Fig. 5), and that HCV infection induced the upregulation of canonical

YAP1 target genes, CYR61 and CTGF (Fig. 6).

236   Both CYR61 and CTGF are crucial for hepatocellular carcinoma (HCC) progression,

exhibiting higher expression in HCC tissues than in normal liver tissues (21). CYR61 and

CTGF act as adhesive substrates that stimulate fibroblast signaling, enhancing angiogenesis

and wound healing (22). Notably, elevated CTGF levels in human HCCs have also been

linked to poor clinical outcomes (23). We obtained evidence suggesting that HCV infection

inactivates the Hippo pathway, leading to the upregulation of CYR61 and CTGF expression,

thereby contributing to HCC progression.

Our findings further elucidate the upstream regulatory mechanisms underlying HCV-

induced inactivation of the Hippo pathway. We demonstrated that HCV infection promotes

K6- and K11-linked polyubiquitylation and proteasomal degradation of LATS1. Previous

studies suggest that all non-K63 linkages, including K6, K11, K27, K29, and K33, can target

protein for proteasomal degradation (24, 25).

The NEDD4 subfamily of HECT-type E3 ubiquitin ligases is characterized by

multiple WW domains that specifically interact with the PPxY domain of LATS1. Notably,

certain members of this subfamily have been identified as key regulators of LATS1 protein

stability (19, 20, 24, 25). We hypothesized that the NEDD4 subfamily, such as Itch or

WWP1, is involved in HCV-induced polyubiquitylation and proteasomal degradation of

LATS1. Our results suggest that Itch, but not WWP1, plays a crucial role in HCV-induced

degradation of LATS1 protein (Figs. 3 and 4). Both Itch and WWP1 contributed to LATS1

polyubiquitylation in mock-infected cells. Importantly, siRNA-mediated knockdown of Itch

restored LATS1 protein in HCV-infected cells. In contrast, siRNA-mediated knockdown of

WWP1 further reduced LATS1 protein levels (Fig. 3E). Notably, condition-specific

regulatory shifts have been observed in some proteins. For example, the tumor suppressor

p53 is primarily degraded by Murine Double Minute 2 (Mdm2) under normal conditions,

whereas in HPV16 E6-positive cancer cells, degradation occurs entirely via the E6-AP

pathway instead of Mdm2 (26). These observations raise the possibility that HCV infection

predominantly facilitates Itch-mediated ubiquitylation and degradation of LATS1 via as-yet-

undetermined mechanisms. Further investigation is required to elucidate the details of these

regulatory mechanisms.

We previously showed that HCV activates the ROS/JNK signaling pathway, which

activates Itch and facilitates polyubiquitylation of VPS4A, enhancing HCV release (2). In this

study, we demonstrated evidence suggesting that HCV-induced ROS/JNK/Itch activation

promotes the ubiquitin-dependent degradation of LATS1, contributing to the inactivation of

the Hippo pathway and nuclear translocation of YAP1, thereby upregulating CYR61 and

CTGF genes. This study represents, to our knowledge, the first demonstration that HCV-

induced ROS/JNK/Itch activation promotes Hippo pathway inactivation, leading to

upregulation of CYR61 and CTGF genes. These insights contribute to our understanding of

the molecular mechanisms underlying HCV pathogenesis and hepatocarcinogenesis,

highlighting potential targets for therapeutic intervention.

MATERIALS AND METHODS

Cell culture and viruses

Human hepatoma cell line Huh-7.5 cells were generously provided by Dr. Charles M. Rice

(The Rockefeller University, New York, NY). The cells were cultured in Dulbecco's Modified

Eagle's Medium (DMEM) (high glucose) containing L-glutamine and phenol red (044-

29765; Fuji Film Wako Pure Chemical Industries, Osaka, Japan) supplemented with 50 IU/ml

penicillin, 50 µg/ml streptomycin (15-140-122; Gibco, Grand Island, NY), 10% heat-

inactivated fetal bovine serum (FBS) (S1760-500; Biowest, Nuaillé, France), and 0.1 mM

non-essential amino acids (Invitrogen, Carlsbad, CA). Cells were cultured at 37°C in a 5%

CO_2 incubator. Cells were washed with PBS (-) solution (05913; Nissui Pharmaceutical Co.,

Ltd., Tokyo, Japan) and transfected with plasmid DNA using FuGENE 6 transfection reagents

(E269A; Promega, Madison, WI). The pFL-J6/JFH1 plasmid, encoding the entire viral

genome of the chimeric HCV-2a strain J6/JFH1, was also kindly provided by Dr. Charles M.

Rice. HCV genome RNA was synthesized *in vitro* using pFL-J6/JFH1 as a template and

subsequently transfected into Huh-7.5 cells via electroporation. The virus produced in the

culture supernatant was then used for infection experiments (27, 28).

Expression plasmids

The cDNA fragment of LATS1 was amplified by PCR using the following specific primers:

forward 5'-TCGAGCTCAGCGGCCATGAAGAGGAGTGAAAAGCCAG-3' and reverse

5'-TCGAGCTCAGCGGCCATGAAGAGGAGTGAAAAGCCAG-3'. The amplified PCR

product was purified and inserted into the NotI site of the pCAG-FLAG plasmid using the In-

Fusion HD Cloning Kit (Clontech, Mountain View, CA). The sequences of the inserts were

verified by sequencing (Eurofins Genomics, Tokyo, Japan).

To construct the pCAG-Myc-LATS1 expression plasmid, pCAG-FLAG-LATS1 was

used as a template and the cDNA fragment of LATS1 was amplified using the following

specific primers: forward 5'-

AAGGTACCATGGAGCAAAAGCTCATTTCTGAAGAGGACCTGATGAAGAGGAGTG

AAAAGCCA-3' and reverse 5'-TTAGATCTTCAAACATATACTAGATCGCGATT-3'. The

amplified PCR product was purified and inserted into the KpnI and BglII sites of the pCAG-

MCS2 vector using the Ligation-Convenience Kit (Nippon Gene, Tokyo, Japan).

The expression plasmids pCAG-FLAG-Itch and pCAG-FLAG−WWP1 were

described previously (29). The C890A point mutant of WWP1 was generated by overlap

extension PCR using pCAG-FLAG−WWP1 as a template and the following specific primers:

forward 5'-AGAAGCCATACAGCTTTTAATCGCTTG-3' and reverse 5'-

CAAGCGATTAAAAGCTGTATGGCTTCT-3'. The sequences of the inserts were verified

by sequencing (Eurofins Genomics).

N-terminal HA-tagged ubiquitin (Ub) expression plasmids, including pRK5-HA−Ub-

WT, pRK5-HA−Ub-K6, pRK5-HA−Ub-K11, pRK5-HA−Ub-K27, pRK5-HA−Ub-K29,

pRK5-HA−Ub-K33, pRK5-HA−Ub-K48, and pRK5-HA−Ub-K63, were purchased from

Addgene (Watertown, MA).

Antibodies and reagents

The mouse monoclonal antibodies (MAbs) used in this study were anti-FLAG (M2) MAb

(F3165, Sigma-Aldrich), anti-β-actin (A5441; Sigma-Aldrich) anti-c-Myc (9E10) Mab (sc-

40; Santa Cruz Biotechnology, Dallas, TX), anti-Itch Mab (611198, BD Transduction

Laboratories, San Jose, CA), anti-Histone H3 (1G1) MAb (sc-517576; Santa Cruz

Biotechnology), and anti-YAP (63.7) MAb (sc-101199; Santa Cruz Biotechnology). The

rabbit monoclonal antibody used in this study was anti-LATS1 (C66B5) MAb (3477; Cell

Signaling Technology). The rabbit polyclonal antibodies (PAbs) used in this study included

anti-HA PAb (H-6908; Sigma-Aldrich), anti-phospho-LATS1 (Ser909) PAb (9157; Cell

Signaling Technology), anti-LAMP2A PAb (ab18528; Abcam, Cambridge, UK), anti-YAP

PAb (4912; Cell Signaling Technology), anti-phospho-YAP (Ser127) (4911; Cell Signaling

Technology), anti-NS5A PAb (2914-1; a kind gift from T. Wakita, National Institute of

Infectious Diseases, Tokyo, Japan), and anti-IκBα PAb (9242; Cell Signaling Technology).

Horseradish peroxidase (HRP)-conjugated anti-mouse IgG (7076S; Cell Signaling

Technology) and HRP-conjugated anti-rabbit IgG (7074S; Cell Signaling Technology) were

used as secondary antibodies. Clasto-lactacystin β-lactone (031-18201; Wako) was used as a

proteasome inhibitor. Ammonium chloride (NH_4Cl) was purchased from Fuji Film Wako

Pure Chemical Industries.

Immunoblot analysis

Immunoblot analysis was conducted as previously described (30). Cell lysates were separated

using 8% or 15% sodium dodecyl sulfate−polyacrylamide gel electrophoresis (SDS-PAGE)

and transferred onto a 0.45 µm Immobilon-P polyvinylidene difluoride (PVDF) membrane

(Millipore, Billerica, MA). The membranes were incubated with a primary antibody, followed

by an HRP-conjugated secondary antibody. Positive bands were detected using Amersham

enhanced chemiluminescence (ECL) western blotting detection reagents (Cytiva, RPN2106;

Sigma-Aldrich). Band intensities were quantified using ImageJ software (version 1.54h).

Cell-based ubiquitylation assay and immunoprecipitation

Cell-based ubiquitylation assays and immunoprecipitation experiments were performed as

previously described (29). Cultured cells were lysed in a lysis buffer containing 50 mM

HEPES (pH 7.5), 150 mM NaCl, 1 mM EDTA, 1% NP-40, 1 mM DTT, 1 mM PMSF, 10 mM

pyrophosphate, 10 mM glycerophosphate, 50 mM NaF, 1.5 mM Na_3VO_4, and a complete™,

EDTA-free protease inhibitor cocktail (Roche Molecular Biochemicals, Mannheim,

Germany). After sonication, the lysates were immunoprecipitated using either anti-FLAG M2

affinity gel (A2220; Sigma-Aldrich) or Protein A-Sepharose 4 Fast Flow (GE17-5280-04; GE

Healthcare, Buckinghamshire, UK) pre-incubated with the appropriate antibodies.

Immunoprecipitation was carried out at 4°C for 4 h or overnight. The immunoprecipitates

were washed five times with the lysis buffer and analyzed by immunoblotting using specific

antibodies, including anti-HA for ubiquitylation assays.

Cell fractionation assay

Cytoplasmic and nuclear extracts from HCV-infected and mock-infected Huh-7.5 cells were

separated using NE-PER™ Nuclear and Cytoplasmic Extraction Reagents (78835; Thermo

Fisher Scientific) following the manufacturer's instructions. The resulting lysates were then

analyzed by immunoblotting.

RNA extraction and RT-qPCR

Total RNA was extracted using the ReliaPrep RNA Cell Miniprep System (Promega) following the manufacturer's protocol. The GoScript Reverse Transcription System (Promega) was used for cDNA synthesis. RT-qPCR was performed on a StepOnePlus Real-Time PCR System (Applied Biosystems, Foster City, CA) using TB Green Premix Ex Taq II (TaKaRa Bio, Shiga, Japan) with SYBR Green chemistry. The primer sequences used were as follows: LATS1 (forward: 5'-CTCTGCACTGGCTTCAGATG-3', reverse: 5'-TCCGCTCTAATGGCTTCAGT-3'); YAP1 (sense: 5'- CCCGACAGGCCAGTACTGAT-3', antisense: 5'-CAGAGAAGCTGGAGAGGAATGAG-3'CYR61 (sense: 5'-GGTCAAAGTTACCGGGCAGT-3', antisense: 5'-GGAGGCATCGAATCCCAGC-3'); and CTGF (forward: 5'-CCTGGTCCAGACCACAGAGT-3', reverse: 5'-ATGTCTTCATGCTGGTGCAG-3'). Expression of the human GAPDH gene was measured as an internal control using the primers 5'-GCCATCAATGACCCCTTCATT-3' (forward) and 5'-TCTCGCTCCTGGAAGATGG-3'(reverse).

RNA interference and stable knockdown cells

The generation of shControl and shLAMP2A cells was described previously (31). HCV-infected and uninfected Huh-7.5 cells were transfected with either Itch siRNA (SI00141085; Qiagen,). WWP1 siRNA (JP00049582-001; Sigma-Aldrich), YAP siRNA (WD11034049-001; Sigma-Aldrich), or negative-control scrambled siRNA (1027281; Qiagen) using Lipofectamine RNAiMAX transfection reagent (Life Technologies, Carlsbad, CA) according to the manufacturer's instructions. The siRNA sequences were as follows: siWWP1, 5'-CUUCUACGAUCAUCAACUCTT-3'; siYAP1, 5'-TTCTTTATCTAGCTTGGTGGC-3'. All siRNAs were transfected once.

Immunofluorescence staining

Huh-7.5 cells cultured on glass coverslips were fixed with 4% paraformaldehyde at room temperature (RT) for 15 min. After being washed with phosphate-buffered saline (PBS) (05913; Nissui Pharmaceutical Co., Ltd.), the cells were permeabilized with PBS containing 0.1% Triton X-100 for 15 min at RT. To block nonspecific binding, the cells were incubated in PBS containing 1% bovine serum albumin (BSA) (01859-47; Nacalai Tesque) for 60 min. The cells were then incubated with 1% BSA in PBS containing mouse anti-YAP monoclonal antibody and rabbit anti-NS5A polyclonal antibody at RT for 60 min. Afterward, the cells were washed three times with PBS and further incubated with 1% BSA in PBS containing Alexa Fluor™ 594-conjugated anti-mouse IgG (A11005; Invitrogen) and Alexa Fluor™ 488-conjugated anti-rabbit IgG (A11008; Invitrogen) at RT for 60 min. Finally, the cells were washed four times with PBS, mounted on glass slides, and examined using a confocal microscope (LSM 700, Carl Zeiss, Oberkochen, Germany).

Statistical analysis

Results were expressed as means ± standard error of the mean (SEM). Statistical significance was evaluated using Student's t-test and was defined as a P value of <0.05.

Data Availability Statement

All data are presented in the main figures. The data that support the findings of this study is available at bioRxiv (https://www.biorxiv.org/). Raw sequencing data, microscopy images, materials, and sequence information are available upon request. Correspondence and requests for materials should be addressed to Professor Ikuo Shoji.

ACKNOWLEDGEMENTS

415 We sincerely appreciate Dr. C. M Rice (The Rockefeller University, New York) for

416 generously providing the Huh-7.5 cells and pFL-J6/JFH1 plasmid. We also thank Y. Kozaki

417 for her valuable secretarial assistance.

FUNDING INFORMATION

This research was supported by grants for Basic and Clinical Research on Hepatitis from the

Japan Agency for Medical Research and Development (AMED) grant No.

23fk0210090s1203 and 20fk0210040s0703, a KAKENHI grant No. 20K07514 and

22K15470, and the Program for the Nurturing of Next Generation Leaders Guiding Medical

Innovation in Asia, funded by the Ministry of Education, Culture, Sports, Science, and

Technology (MEXT) of Japan.

AUTHOR CONTRIBUTIONS

428 C.M. and I.S. conceived and designed the experiments. S.MA and C.M. carried out most of

429 the experiments. ZX., F.P., C.M., L.D., and T.A. assisted the constructions and the data

430 analysis. S.MA., C.M, and I.S. wrote the manuscript.

DECLARATION OF INTERESTS

433 The authors declare no conflict of interest.

    FIGURE LEGENDS

Figure 1. HCV induces proteasomal degradation of LATS1.

(A) Huh-7.5 cells were infected with HCV J6/JFH1 at a multiplicity of infection (MOI) of 2.

Cells were cultured and harvested at 2 and 4 days post-infection (dpi). Immunoblot analysis

was performed using the indicated antibodies, with β-actin levels serving as a loading control.

(B) Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of 2, cultured, and harvested at

2 and 4 dpi. Total cellular RNA was extracted, and LATS1 mRNA levels were quantified

using RT-qPCR. GAPDH mRNA levels were used for normalization. The value for day 2 in

the mock-infected group was expressed as 1.0. Data are presented as means ± SEM from

three independent experiments; p-value < 0.05 (*) was significant, compared with the

controls. (C) Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of 2, followed by

treatment with NH_4CL for 12 h. Immunoblotting was performed with the specified

antibodies, and β-actin levels served as a loading control. (D) Huh-7.5 cells were infected

with HCV J6/JFH1 at an MOI of 2, followed by treatment with Bafilomycin A1 for 12 h.

Immunoblotting was performed with the indicated antibodies, and β-actin was used as a

loading control.

(E) shControl- and shLAMP2A-Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of

2. Immunoblot analysis was performed using indicated antibodies with β-actin as a loading

control. (F) Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of 2. Cells were

treated with 30 μM clasto-lactacystin β-lactone for 12 h, harvested, and analyzed by

immunoblotting using the indicated antibodies. β-actin was used as a loading control.

Figure 2. HCV induces K6- and K11-linked polyubiquitylation of LATS1.

Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of 0.5 and co-transfected with

pCAG-FLAG-LATS1 together with the indicated HA-tagged ubiquitin mutant plasmids. At 2

days post-transfection, cells were harvested and cell lysates were immunoprecipitated with

anti-FLAG beads, followed by immunoblotting with the indicated antibodies. Input samples

were also analyzed by immunoblotting with the indicated antibodies, and β-actin was used as

a loading control.

Figure 3. E3 ligase Itch, but not WWP1, plays a crucial role in HCV-induced

polyubiquitylation and degradation of LATS1.

(A) Huh-7.5 cells were co-transfected with pCAG-Myc-LATS1, pRK5-HA-Ubiquitin,

together with either pCAG-FLAG-Itch WT or pCAG-FLAG-Itch C868A. At 2 days post-

transfection, the cells were harvested. The cell lysates were immunoprecipitated with anti-c-

Myc antibody-conjugated Sepharose A beads, followed by immunoblotting with the indicated

antibodies. Input samples were also analyzed by immunoblotting. β-actin was used as a

loading control. (B) Huh-7.5 cells were co-transfected with pCAG-Myc-LATS1, pRK5-

HA−Ubiquitin, together with either pCAG-FLAG−WWP1 WT or pCAG-FLAG−WWP1

C890A. At 2 days post-transfection, cells were harvested. The cell lysates were

immunoprecipitated with anti-c-Myc antibody−conjugated Sepharose A beads, followed by

immunoblotting with the indicated antibodies. Input samples were also immunoblotted, with

β-actin serving as a loading control.

(C) Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of 1 and co-transfected with

pCAG-Myc-LATS1 and pRK5-HA−Ubiquitin, together with either pCAG-FLAG-Itch or

pCAG-FLAG−WWP1. Cell lysates were immunoprecipitated using anti-c-Myc

antibody−conjugated Sepharose A beads, followed by immunoblotting with the indicated

antibodies. Input samples were also analyzed. β-actin was used as a loading control. (D) Huh-

7.5 cells (3×10^5 cells per well in a 12-well plate) were transfected with 48 pmol of either

control siRNA or Itch-specific siRNA. After 24 h, the cells were infected with HCV J6/JFH1

at an MOI of 2 and harvested at 2 days post-infection. The samples were analyzed by

immunoblotting with the indicated antibodies. β-actin was used as a loading control. (E) Huh-

7.5 cells (3×10^5 cells per well in a 12-well plate) were transfected with 48 pmol of either

control siRNA or WWP1-specific siRNA. At 24 h after siRNA-transfection, the cells were

infected with HCV J6/JFH1 at an MOI of 2. Cells were harvested at 2 days post-infection,

and the samples were subjected to immunoblotting with the indicated antibodies. β-actin was

used as a loading control.

Figure 4. HCV-induced ROS/JNK/Itch signaling pathway plays a role in

polyubiquitylation of LATS1 protein

Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of 1, then co-transfected with

pCAG-Myc-LATS1 and pRK5-HA-ubiquitin, together with pCAG-FLAG-Itch. After 2 days,

the cells were harvested with or without pretreatment with JNK inhibitor SP6000125 (30 μM

for 30 h), and the lysates were immunoprecipitated with anti-c-Myc antibody−conjugated

Sepharose A beads, followed by immunoblotting with the indicated antibodies. Input samples

were also analyzed by immunoblotting. β-actin was used as a loading control.

Figure 5. HCV infection promotes the translocation of YAP1 from the cytoplasm to the

nucleus.

(A) Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of 2. Following infection, the

cells were harvested and analyzed by immunoblotting using antibodies against components of

the Hippo pathway. (B) Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of 2. At 4

days after infection, the cells were harvested and subjected to cell fractionation analysis. Ikβα

was used as a loading control for the cytoplasmic fraction, and Histone H3 was used as a

loading control for the nuclear fraction. (C) Huh-7.5 cells were plated and cultured for 12 h

and infected with HCV J6/JFH1 at an MOI of 2. Cells were stained with anti-YAP1

monoclonal antibody followed by Alexa Fluor 594−conjugated goat anti-mouse IgG (red) and

anti-NS5A polyclonal antibody followed by Alexa Fluor 488−conjugated goat anti-rabbit IgG

(green). Nuclei were counterstained with Hoechst 33342 (blue). Images were acquired using

scanning laser confocal microscopy and processed with ImageJ software.

Figure 6. The mRNA levels of YAP1 target genes are significantly upregulated upon

HCV infection.

(A, B) Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of 2. Cells were cultured

and harvested at 2 and 4 days post-infection (dpi). Total cellular RNA was subsequently

extracted, and the mRNA levels of CYR61 and CTGF were quantified by RT-qPCR. GAPDH

mRNA levels were used for normalization. The mRNA level for day 2 in the mock-infected

group was arbitrarily expressed as 1.0. Data are presented as means ± SEM from three

independent experiments. A p-value < 0.05 (*) was considered significant. (C−E) Huh-7.5

cells at $(1.8 \times 10^5$ cells per well in a 24-well plate) were transfected with 24 pmol of either

control siRNA or YAP siRNA. At 24 h after siRNA transfection, the cells were infected with

HCV J6/JFH1 at an MOI of 2, cultured, and harvested at the indicated time points. Total RNA

was extracted, and YAP1, CYR61, and CTGF mRNA levels were quantified by RT-qPCR,

normalized to GAPDH mRNA levels. The mRNA level at day 2 in the mock-infected group

was arbitrarily set to 1.0. Data are expressed as mean ± SEM from three independent

experiments. A p-value < 0.05 (*) was considered significant.

Figure 7. A proposed model of HCV-induced inactivation of the Hippo pathway and

HCV-related pathogenesis.

Under normal conditions, the Hippo pathway remains predominantly active, with LATS1

phosphorylating YAP1, thereby retaining YAP1 in the cytoplasm and preventing its nuclear

entry. HCV infection activates the ROS/JNK/Itch signalling pathway, which is crucial for

HCV particle release. Activation of the ROS/JNK/Itch signalling pathway leads to the

phosphorylation and enhanced activity of Itch ubiquitin ligase. Activated Itch promotes the

ubiquitin-dependent proteasomal degradation of LATS1 protein, leading to the Hippo

pathway inactivation. As a result, YAP1 translocates from the cytoplasm to the nucleus and

upregulates the transcription of CYR61 and CTGF genes, contributing to HCV-related

pathogenesis.

REFERENCES

1. World Health Organization. 2024. Global Hepatitis Report 2024: Action for Access in Low- and Middle-Income Countries. World Health Organization, Geneva, Switzerland.

2. Bartenschlager R, Baumert TF, Bukh J, Houghton M, Lemon SM, Lindenbach BD, Lohmann V, Moradpour D, Pietschmann T, Rice CM, Thimme R, Wakita T. 2018. Critical challenges and emerging opportunities in hepatitis C virus research in an era of potent antiviral therapy: Considerations for scientists and funding agencies. Virus Res 248:53–62.

3. Baumert TF, Jühling F, Ono A, Hoshida Y. 2017. Hepatitis C-related hepatocellular carcinoma in the era of new generation antivirals. BMC Med 15(1):52.

4. Scheel TKH, Rice CM. 2013. Understanding the hepatitis C virus life cycle paves the way for highly effective therapies. Nat Med 19(7):837–49.

5. Sohn BH, Shim JJ, Kim SB, Jang KY, Kim SM, Kim JH, Hwang JE, Jang HJ, Lee HS, Kim SC, Jeong W, Kim SS, Park ES, Heo J, Kim YJ, Kim DG, Leem SH, Kaseb A, Hassan MM, Cha M, Chu IS, Johnson RL, Park YY, Lee JS. 2016. Inactivation of Hippo Pathway Is Significantly Associated with Poor Prognosis in Hepatocellular Carcinoma. Clin Cancer Res 22(5):1256–64.

6. Zheng T, Wang J, Jiang H, Liu L. 2011. Hippo signaling in oval cells and hepatocarcinogenesis. Cancer Lett 302(2):91–9.

7. Zhao B, Li L, Lei Q, Guan KL. 2010. The Hippo–YAP pathway in organ size control and tumorigenesis: an updated version. Genes Dev 24(9):862–74.

8. Zhao B, Li L, Tumaneng K, Wang CY, Guan KL. 2010. A coordinated phosphorylation by Lats and CK1 regulates YAP stability through SCF $^{\beta\text{-TRCP}}$. Genes Dev 24(1):72–85.

9. Zhao B, Wei X, Li W, Udan RS, Yang Q, Kim J, Xie J, Ikenoue T, Yu J, Li L, Zheng P, Ye K, Chinnaiyan A, Halder G, Lai ZC, Guan KL. 2007. Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell contact inhibition and tissue growth control. Genes Dev 21(21):2747–61.

10. Meng Z, Moroishi T, Guan KL. 2016. Mechanisms of Hippo pathway regulation. Genes Dev 30(1):1–17.

11. Kwon H, Kim J, Jho E. 2022. Role of the Hippo pathway and mechanisms for controlling cellular localization of YAP/TAZ. FEBS J 289(19):5798–818.

12. Wang Z, Lu W, Zhang Y, Zou F, Jin Z, Zhao T. 2020. The Hippo Pathway and Viral Infections. Front Microbiol 10:3033.

13. Liu G, Yu FX, Kim YC, Meng Z, Naipauer J, Looney DJ, Liu X, Gutkind JS, Mesri EA, Guan KL. 2015. Kaposi sarcoma-associated herpesvirus promotes tumorigenesis by modulating the Hippo pathway. Oncogene 34(27):3536–46.

14. Garcia G, Paul S, Beshara S, Ramanujan VK, Ramaiah A, Nielsen-Saines K, Li MMH, French SW, Morizono K, Kumar A, Arumugaswami V. 2020. Hippo Signaling Pathway Has a Critical Role in Zika Virus Replication and in the Pathogenesis of Neuroinflammation. Am J Pathol 190(4):844–61.

15. Webb Strickland S, Brimer N, Lyons C, Vande Pol SB. 2018. Human Papillomavirus E6 interaction with cellular PDZ domain proteins modulates YAP nuclear localization. Virology 516:127–38.

16. Xue Y, Mars WM, Bowen W, Singhi AD, Stoops J, Michalopoulos GK. 2018. Hepatitis C

Virus Mimics Effects of Glypican-3 on CD81 and Promotes Development of

Hepatocellular Carcinomas via Activation of Hippo Pathway in Hepatocytes. Am J Pathol

188(6):1469–77.

17. Hu B, Xie S, Hu Y, Chen W, Chen X, Zheng Y, Wu X. 2017. Hepatitis C virus NS4B

protein induces epithelial-mesenchymal transition by upregulation of Snail. Virol J

14(1):83.

18. Deng L, Liang Y, Ariffianto A, Matsui C, Abe T, Muramatsu M, Wakita T, Maki M,

Shibata H, Shoji I. 2022. Hepatitis C Virus-Induced ROS/JNK Signaling Pathway

Activates the E3 Ubiquitin Ligase Itch to Promote the Release of HCV Particles via

Polyubiquitylation of VPS4A. J Virol 96(6):e01811-21. A

19. Salah Z, Melino G, Aqeilan RI. 2011. Negative Regulation of the Hippo Pathway by E3

Ubiquitin Ligase ITCH Is Sufficient to Promote Tumorigenicity. Cancer Res 71(5):2010–

20.

20. Ho KC, Zhou Z, She YM, Chun A, Cyr TD, Yang X. 2011. Itch E3 ubiquitin ligase

regulates large tumor suppressor 1 stability. Proc Natl Acad Sci U S A 108(12):4870–5.

21. Zeng ZJ. 2004. Expressions of cysteine-rich61, connective tissue growth factor and Nov

genes in hepatocellular carcinoma and their clinical significance. World J Gastroenterol

10(23):3414.

22. Chen CC, Chen N, Lau LF. 2001. The Angiogenic Factors Cyr61 and Connective Tissue

Growth Factor Induce Adhesive Signaling in Primary Human Skin Fibroblasts. J Biol

Chem 276(13):10443–52.

23. Simile MM, Latte G, Demartis MI, Brozzetti S, Calvisi DF, Porcu A, Feo CF, Seddaiu

MA, Daino L, Berasain C, Tomasi ML, Avila MA, Feo F, Pascale RM. 2016. Post-

translational deregulation of YAP1 is genetically controlled in rat liver cancer and

determines the fate and stem-like behavior of the human disease. Oncotarget

7(31):49194–216.

24. Yeung B, Ho KC, Yang X. 2013. WWP1 E3 Ligase Targets LATS1 for Ubiquitin-

Mediated Degradation in Breast Cancer Cells. PLoS ONE 8(4):e61027.

25. Salah Z, Cohen S, Itzhaki E, Aqeilan R. 2013. NEDD4 E3 ligase inhibits the activity of

the Hippo pathway by targeting LATS1 for degradation. Cell Cycle 12(24):3817–23.

26. Hengstermann A, Linares LK, Ciechanover A, Whitaker NJ, Scheffner M. 2001.

Complete switch from Mdm2 to human papillomavirus E6-mediated degradation of p53

in cervical cancer cells. Proc Natl Acad Sci U S A 98(3):1218-1223.

27. Wakita T, Pietschmann T, Kato T, Date T, Miyamoto M, Zhao Z, Murthy K, Habermann

A, Krausslich HG, Mizokami M, Bartenschlager R, Liang TJ. 2005. Production of

infectious hepatitis C virus in tissue culture from a cloned viral genome. Nat Med

11(7):791–6.

28. Bungyoku Y, Shoji I, Makine T, Adachi T, Hayashida K, Nagano-Fujii M, Ide Y, Deng L,

Hotta H. 2009. Efficient production of infectious hepatitis C virus with adaptive

mutations in cultured hepatoma cells. J Gen Virol 90(7):1681–91.

29. Shirakura M, Murakami K, Ichimura T, Suzuki R, Shimoji T, Fukuda K, Abe K, Sato S,

Fukasawa M, Yamakawa Y, Nishijima M, Moriishi K, Matsuura Y, Wakita T, Suzuki T,

Howley PM, Miyamura T, Shoji I. 2007. E6AP Ubiquitin Ligase Mediates Ubiquitylation

and Degradation of Hepatitis C Virus Core Protein. J Virol 81(3):1174–85.

30. Matsui C, Shoji I, Kaneda S, Sianipar IR, Deng L, Hotta H. 2012. Hepatitis C Virus

Infection Suppresses GLUT2 Gene Expression via Downregulation of Hepatocyte

Nuclear Factor 1α. J Virol 86(23):12903–11.

30. Yuliandari P, Matsui C, Deng L, Abe T, Mori H, Taguwa S, Ono C, Fukuhara T, Matsuura

Y, Shoji I. 2022. Hepatitis C virus NS5A protein promotes the lysosomal degradation of

diacylglycerol O-acyltransferase 1 (DGAT1) via endosomal microautophagy. Autophagy

Rep 1(1):264–85.

Fig. 1 Maria et al.

Fig. 2 Maria et al.

Fig. 3 Maria et al.

HA-Ubiquitin   **+   +   +   +   +   +**
FLAG-Itch   **-   +   +   -   +   +**
SP600125   **-   -   +   -   -   +**
Myc-LATS1   **+   +   +   +   +   +**
HCV   **-   -   -   +   +   +**

(kDa)

180 –

130 –   HA-Ubiquitin   IP: Myc

130 –   LATS1

130 –   LATS1

130 –   myc-LATS1

100 –   FLAG-Itch   Input

63 –   NS5A

48 –   β-actin

Lane   **1   2   3   4   5   6**

Fig. 4 Maria et al.

A

B

C

YAP1   NS5A   Nucleus   Merge

Mock

HCV

Fig. 5 Maria et al.

Normal condition

HCV infection