HCV infection induces ubiquitin-dependent degradation of LATS1, inactivating the Hippo pathway and
upregulating transcription of the CYR61 and CTGF genes

Maria Alethea Septianastiti [1, 2], Chieko Matsui [1], Zihan Xu [1], Fransisca Puspitasari [1], Dewa
Nyoman Murti Adyaksa[1,2], Lin Deng [1], Takayuki Abe [3], Ikuo Shoji [1, #]

[1] Division of Infectious Disease Control, Center for Infectious Diseases, Kobe University
Graduate School of Medicine, Kobe, Japan

[2] Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta,
Indonesia

[3] Division of Virology, Niigata University Graduate School of Medical and Dental Sciences,
Niigata, Japan

Correspondence: Prof. Ikuo Shoji
E-mail: ishoji@med.kobe-u.ac.jp
Address: Division of Infectious Disease Control, Center for Infectious Diseases, Kobe
University Graduate School of Medicine, 7-5-1 Kusunoki-cho, Chuo-ku, Kobe, Japan.
Tel.: +81-78-382-5500; Fax: +81-78-382-5519.
19
Keywords: HCV, LATS1, Itch, ubiquitin, Hippo pathway, YAP1

Word count

Abstract: 209 words

Impact statement: 89 words

Text: 3425 words

Figure: 6

ABSTRACT

Hepatitis C virus (HCV) is often associated with chronic liver diseases and significant alterations in host cellular signaling. However, the molecular mechanisms underlying HCV-related liver pathogenesis remain to be elucidated. The Hippo signaling pathway, a key regulator of cell proliferation and survival, plays a critical role in maintaining liver homeostasis. Here we investigated the role of the Hippo pathway in HCV-related pathogenesis. We demonstrated that HCV infection induces degradation of LATS1, a key regulator of the Hippo pathway. Degradation of LATS1 protein was restored by a proteasomal inhibitor, but not a lysosome inhibitor, indicating that HCV promotes proteasomal degradation of LATS1 protein. HCV-induced degradation of LATS1 protein was suppressed in si-Itch-transfected Huh-7.5 cells. These results suggest that Itch ubiquitin ligase is involved in ubiquitin-dependent degradation of LATS1 protein. Cell fractionation assays and immunofluorescence staining revealed that HCV infection promoted nuclear translocation of YAP1 protein, suggesting that HCV infection suppresses the Hippo pathway. Furthermore, the transcription of YAP1 target genes, CYR61 and CTGF, that are involved in tissue remodeling and proliferation, was upregulated in HCV-infected Huh-7.5 cells and in HCV-infected patients. Taken together, we propose that HCV promotes the ubiquitin-dependent proteasomal degradation of LATS1 protein, leading to suppression of the Hippo pathway, thereby upregulating transcription of CYR61 and CTGF genes.

DATA SUMMARY

All data are presented in the main figures. Raw sequencing data, microscopy images, materials, and sequence information are available upon request. Correspondence and requests

for materials should be addressed to Professor Ikuo Shoji. The data that support the findings

of this study is available at bioRxiv (https://www.biorxiv.org/).

IMPACT STATEMENT

We demonstrate evidence suggesting that HCV infection promotes the Itch-mediated

ubiquitin-dependent degradation of LATS1 protein, a key factor for the Hippo pathway.

HCV-induced ubiquitin-dependent degradation of LATS1 protein promotes inactivation of

the Hippo pathway and nuclear translocation of YAP1 protein, thereby upregulating

transcription of CYR61 and CTGF genes. We propose a novel molecular mechanism in

which HCV infection promotes degradation of LATS1 protein, leading to inactivation of the

Hippo pathway. Understanding HCV-induced inactivation of the Hippo pathway may lead to

developing new strategies for preventing or treating HCV-related pathogenesis.

INTRODUCTION

Hepatitis C virus (HCV) infection remains a global health burden, affecting an estimated 50 million people worldwide as of 2022 (1). HCV infection often causes chronic liver diseases, ranging from mild illness to severe complications, including liver cirrhosis and hepatocellular carcinoma (HCC). Although highly effective direct-acting antivirals (DAAs) are available for HCV treatment, the risk of HCC persists even after successful viral clearance (2). Notably, HCC can occur following a sustained virological response, at an annual incidence of approximately 1%, surpassing the cancerous conditions in other organs (3). HCV is a positive-sense, single-stranded RNA virus classified in the *Flaviviridae* family, *Hepacivirus* genus. The 9.6-kb HCV RNA genome encodes a single polyprotein of approximately 3,010 amino acids, which is cleaved by viral proteases and host signalases into structural proteins (Core, E1, and E2) and nonstructural proteins (p7, NS2, NS3, NS4A, NS5A, and NS5B) (4).

Understanding HCV-induced modulation of host cellular signaling pathway is crucial for elucidating mechanisms that drive liver disease progression. The Hippo pathway, a key regulator of cell proliferation, apoptosis, and organ size control, plays an essential role in tissue homeostasis. Dysregulation of the Hippo pathway is implicated in the development of several cancers, including liver cancer (5–7). The Hippo pathway involves a series of proteins that form a kinase cascade, including MST1/2, LATS1/2, the adaptor protein SAV1, MOB1A/B, and the transcription coactivators YAP1 and TEAD transcription factors. Activation of the Hippo pathway involves the phosphorylation of the Serine/Threonine kinase LATS1, which in turn phosphorylates YAP1 at Ser127 and promotes the binding of phosphorylated YAP1 to 14-3-3 proteins (8, 9). The association of 14-3-3 proteins with phosphorylated YAP1 prevents it from entering the nucleus. Conversely, when the Hippo pathway is inactivated, unphosphorylated YAP1 translocates to the nucleus and induces the transcription of genes responsible for cell growth (10,11).

Several viruses have been shown to modulate the Hippo pathway (12–15). HCV E2

activates the Hippo pathway by interacting with CD81, while HCV NS4B suppresses the

Hippo pathway via Scribble, promoting PI3K/AKT signaling and epithelial-mesenchymal

transition (EMT) (16,17). Similarly, Kaposi sarcoma-associated herpesvirus (KSHV) vGPCR

inhibits LATS1/2, enabling YAP1 nuclear translocation. Human papillomavirus 16 E6 alters

YAP1 localization through interaction with PDZ domains (13,15).

We previously reported that HCV infection activates the ROS/JNK signaling pathway

which in turn activates Itch, a HECT-type E3 ubiquitin ligase. Activated Itch promotes

polyubiquitylation of VPS4A, resulting in efficient HCV particle release (18). Notably, Itch

has been shown to promote the proteasomal degradation of LATS1, a critical component of

the Hippo pathway (19,20). In this study, we investigate a role of the Hippo pathway in

HCV-related pathogenesis. We demonstrate that Itch plays an important role in HCV-induced

ubiquitin-dependent degradation of LATS1, leading to Hippo pathway suppression and

thereby contributing to HCV pathogenesis.

METHODS

Cell culture and viruses.

Human hepatoma cell line Huh-7.5 cells were generously provided by Dr. Charles M. Rice

(The Rockefeller University, New York, NY). The cells were cultured in Dulbecco's

Modified Eagle's Medium (DMEM) (high glucose) containing L-glutamine and phenol red

(044-29765; Fuji Film Wako Pure Chemical Industries, Osaka, Japan) supplemented with 50

IU/ml penicillin, 50 µg/ml streptomycin (15-140-122; Gibco, Grand Island, NY), 10% heat-

inactivated fetal bovine serum (FBS) (S1760-500; Biowest, Nuaillé, France), and 0.1 mM

non-essential amino acids (Invitrogen, Carlsbad, CA). Cells were cultured at 37°C in a 5%

CO☐ incubator. Cells were washed with PBS (-) solution (05913; Nissui Pharmaceutical Co.,

Ltd., Tokyo, Japan) and transfected with plasmid DNA using FuGENE 6 transfection

reagents (E269A; Promega, Madison, WI) following the manufacturer's instructions. The

pFL-J6/JFH1 plasmid, encoding the entire viral genome of the chimeric HCV-2a strain

J6/JFH1, was also kindly provided by Dr. Charles M. Rice. HCV genome RNA was

synthesized *in vitro* using pFL-J6/JFH1 as a template and subsequently transfected into Huh-

7.5 cells via electroporation. The virus produced in the culture supernatant was then used for

infection experiments (21, 22).

The Huh-7 cells stably harboring HCV genotype 1b (RCYM1) were described previously

(23)

Expression plasmids.

The cDNA fragment of LATS1 was amplified by PCR using the following specific primers:

forward 5'-TCGAGCTCAGCGGCCATGAAGAGGAGTGAAAAGCCAG-3' and reverse

5'-TCGAGCTCAGCGGCCATGAAGAGGAGTGAAAAGCCAG-3'. The amplified PCR

product was purified and inserted into the NotI site of the pCAG-FLAG plasmid using the In-

Fusion HD Cloning Kit (Clontech, Mountain View, CA). The sequences of the inserts were

verified by sequencing (Eurofins Genomics, Tokyo, Japan).

To construct the pCAG-Myc-LATS1 expression plasmid, pCAG-FLAG-LATS1 was

used as a template and the cDNA fragment of LATS1 was amplified using the following

specific primers: forward 5'-

AAGGTACCATGGAGCAAAAGCTCATTTCTGAAGAGGACCTGATGAAGAGGAGTG

AAAAGCCA-3' and reverse 5'-TTAGATCTTCAAACATATACTAGATCGCGATT-3'.

The amplified PCR product was purified and inserted into the KpnI and BglII sites of the

pCAG-MCS2 vector using the Ligation-Convenience Kit (Nippon Gene, Tokyo, Japan).

The expression plasmids pCAG-FLAG-Itch and pCAG-FLAG-WWP1 were

described previously (24). The C890A point mutant of WWP1 was generated by overlap

extension PCR using pCAG-FLAG-WWP1 as a template and the following specific primers:

forward 5'-AGAAGCCATACAGCTTTTAATCGCTTG-3' and reverse 5'-

CAAGCGATTAAAAGCTGTATGGCTTCT-3'. The sequences of the inserts were verified

by sequencing (Eurofins Genomics).

N-terminal HA-tagged ubiquitin (Ub) expression plasmid was purchased from

Addgene (Watertown, MA).

Antibodies and reagents.

The mouse monoclonal antibodies (MAbs) used in this study were anti-FLAG (M2) MAb

(F3165, Sigma-Aldrich), anti-β-actin (A5441; Sigma-Aldrich) anti-c-Myc (9E10) Mab (sc-

40; Santa Cruz Biotechnology, Dallas, TX), anti-Itch Mab (611198, BD Transduction

Laboratories, San Jose, CA), anti-Histone H3 (1G1) MAb (sc-517576; Santa Cruz

Biotechnology), and anti-YAP (63.7) MAb (sc-101199; Santa Cruz Biotechnology). The

rabbit monoclonal antibody used in this study was anti-LATS1 (C66B5) MAb (3477; Cell

Signaling Technology). The rabbit polyclonal antibodies (PAbs) used in this study included

anti-HA PAb (H-6908; Sigma-Aldrich), anti-phospho-LATS1 (Ser909) PAb (9157; Cell

Signaling Technology), anti-LAMP2A PAb (ab18528; Abcam, Cambridge, UK), anti-YAP

PAb (4912; Cell Signaling Technology), anti-phospho-YAP (Ser127) (4911; Cell Signaling

Technology), anti-NS5A PAb (2914-1; a kind gift from T. Wakita, National Institute of

Infectious Diseases, Tokyo, Japan), and anti-IκBα PAb (9242; Cell Signaling Technology).

Horseradish peroxidase (HRP)-conjugated anti-mouse IgG (7076S; Cell Signaling

Technology) and HRP-conjugated anti-rabbit IgG (7074S; Cell Signaling Technology) were

used as secondary antibodies. Clasto-lactacystin β-lactone (031-18201; Wako) was used as a

proteasome inhibitor. Ammonium chloride (NH$_4$Cl) was purchased from Fuji Film Wako

Pure Chemical Industries.

Immunoblot analysis.

Immunoblot analysis was conducted as previously described (25). Cell lysates were separated

using 8% or 15% sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE)

and transferred onto a 0.45 µm Immobilon-P polyvinylidene difluoride (PVDF) membrane

(Millipore, Billerica, MA). The membranes were incubated with a primary antibody,

followed by an HRP-conjugated secondary antibody. Positive bands were detected using

Amersham enhanced chemiluminescence (ECL) western blotting detection reagents (Cytiva,

RPN2106; Sigma-Aldrich). Band intensities were quantified using ImageJ software (version

1.54h).

Cell-based ubiquitylation assay and immunoprecipitation.

Cell-based ubiquitylation assays and immunoprecipitation experiments were performed as

previously described (24). Cultured cells were lysed in a lysis buffer containing 50 mM

HEPES (pH 7.5), 150 mM NaCl, 1 mM EDTA, 1% NP-40, 1 mM DTT, 1 mM PMSF, 10

mM pyrophosphate, 10 mM glycerophosphate, 50 mM NaF, 1.5 mM Na□VO□, and a

complete™, EDTA-free protease inhibitor cocktail (Roche Molecular Biochemicals,

Mannheim, Germany). After sonication, the lysates were immunoprecipitated using either

anti-FLAG M2 affinity gel (A2220; Sigma-Aldrich) or Protein A-Sepharose 4 Fast Flow

(GE17-5280-04; GE Healthcare, Buckinghamshire, UK) pre-incubated with the appropriate

antibodies. Immunoprecipitation was carried out at 4°C for 4 h or overnight. The

immunoprecipitates were washed five times with the lysis buffer and analyzed by

immunoblotting using specific antibodies, including anti-HA for ubiquitylation assays.

Cell fractionation assay.

Cytoplasmic and nuclear extracts from HCV-infected and mock-infected Huh-7.5 cells were

separated using NE-PER™ Nuclear and Cytoplasmic Extraction Reagents (78835; Thermo

Fisher Scientific) following the manufacturer's instructions. The resulting lysates were then

analyzed by immunoblotting.

CHX chase experiment.

To determine the half-life of LATS1 protein, cells were treated with 50 µg/mL

cycloheximide (CHX). Cells designated as the zero-time point were harvested immediately

after CHX treatment. For subsequent time points, cells were incubated CHX-containing

medium at 37ºC for 6, 12, 18, and 24 hours, as indicated.

RNA extraction and RT-qPCR.

Total RNA was extracted using the ReliaPrep RNA Cell Miniprep System (Promega)

following the manufacturer's protocol. The GoScript Reverse Transcription System

(Promega) was used for cDNA synthesis. RT-qPCR was performed on a StepOnePlus Real-

Time PCR System (Applied Biosystems, Foster City, CA) using TB Green Premix Ex Taq II

(TaKaRa Bio, Shiga, Japan) with SYBR Green chemistry. The primer sequences used were

as follows: LATS1 (forward: 5'-CTCTGCACTGGCTTCAGATG-3', reverse: 5'-

TCCGCTCTAATGGCTTCAGT-3'); YAP1 (forward: 5'-

CCCGACAGGCCAGTACTGAT-3', reverse: 5'-CAGAGAAGCTGGAGAGGAATGAG-

3'); CYR61 (forward: 5'-GGTCAAAGTTACCGGGCAGT-3', reverse: 5'-

GGAGGCATCGAATCCCAGC-3'); and CTGF (forward: 5'-

CCTGGTCCAGACCACAGAGT-3', reverse: 5'-ATGTCTTCATGCTGGTGCAG-3').

Expression of the human GAPDH gene was measured as an internal control using the primers

(forward: 5'-GCCATCAATGACCCCTTCATT-3', reverse: 5'-

TCTCGCTCCTGGAAGATGG-3'.

RNA interference and stable knockdown cells.

The generation of shControl and shLAMP2A cells was described previously (26). HCV-

infected and uninfected Huh-7.5 cells were transfected with either Itch siRNA (SI00141085;

Qiagen,). WWP1 siRNA (JP00049582-001; Sigma-Aldrich), YAP siRNA (WD11034049-

001; Sigma-Aldrich), or negative-control scrambled siRNA (1027281; Qiagen) using

Lipofectamine RNAiMAX transfection reagent (Life Technologies, Carlsbad, CA) according

to the manufacturer's instructions. The siRNA sequences were as follows: siWWP1, 5'-

CUUCUACGAUCAUCAACUCTT-3'; siYAP1, 5'-TTCTTTATCTAGCTTGGTGGC-3'. All

siRNAs were transfected once.

Immunofluorescence staining.

Huh-7.5 cells cultured on glass coverslips were fixed with 4% paraformaldehyde at room

temperature (RT) for 15 min. After being washed with phosphate-buffered saline (PBS)

(05913; Nissui Pharmaceutical Co., Ltd.), the cells were permeabilized with PBS containing

0.1% Triton X-100 for 15 min at RT. To block nonspecific binding, the cells were incubated

in PBS containing 1% bovine serum albumin (BSA) (01859-47; Nacalai Tesque) for 60 min.

The cells were then incubated with 1% BSA in PBS containing mouse anti-YAP monoclonal

antibody and rabbit anti-NS5A polyclonal antibody at RT for 60 min. Afterward, the cells

were washed three times with PBS and further incubated with 1% BSA in PBS containing

Alexa Fluor™ 594-conjugated anti-mouse IgG (A11005; Invitrogen) and Alexa Fluor™ 488-

conjugated anti-rabbit IgG (A11008; Invitrogen) at RT for 60 min. Finally, the cells were

washed four times with PBS, mounted on glass slides, and examined using a confocal

microscope (LSM 700, Carl Zeiss, Oberkochen, Germany).

Transcriptomic data analysis.

To evaluate the expression of CYR61 and CTGF mRNAs in HCV-infected liver tissue,

transcriptomic data were obtained from the GEO database (accession number GSE84346).

The dataset included samples from HCV-infected patients (n = 38) and healthy controls (n =

6). Normality of gene expression data was assessed using the Kolmogorov-Smirnov and

Shapiro-Wilk tests. As the data were not normally distributed, group comparisons were

performed using the non-parametric Mann-Whitney U test. A p-value <0.05 was considered

statistically significant.

Statistical analysis.

For RT-qPCR data, results were expressed as means ± standard error of the mean (SEM).

Statistical significance was evaluated using Student's t-test and was defined as a P value of

<0.05.

RESULTS

HCV promotes proteasomal degradation of LATS1 protein.

To investigate a role of the Hippo signaling pathway in HCV-related pathogenesis, we

examined LATS1 protein levels in HCV J6/JFH1-infected Huh-7.5 cells. Immunoblot

analysis revealed a reduction in LATS1 protein levels in HCV-infected cells at both 2 and 4

days post-infection (Fig. 1A, first panel, lanes 2 and 4). However, reverse-transcription

quantitative polymerase chain reaction (RT-qPCR) analysis showed increased LATS1 mRNA

levels (Fig. 1B), indicating that HCV-induced reduction of LATS1 protein levels is not due to

suppression of LATS1 mRNA expression.

To determine whether HCV-induced reduction of LATS1 protein is protein

degrdataion, we performed CHX-chase analyses. The CHX-chase analyses revealed that the

half-life of LATS1 protein was significantly reduced upon HCV infection (Fig. 1C, first

panel; Fig. 1D).

To further elucidate a mechanism underlying HCV-induced reduction of LATS1

protein, we treated HCV-infected Huh-7.5 cells with the lysosomal inhibitor bafilomycin A1.

The LATS1 protein levels in HCV-infected cells were not restored by treatment with

bafilomycin A1 (Fig. 1E, first panel, lane 6), suggesting that HCV-induced LATS1 protein

reduction is not due to the lysosomal degradation.

To determine whether HCV-induced LATS1 protein reduction is due to proteasomal

degradation, we treated the cells with the proteasome inhibitor clasto-lactacystin β-lactone.

Treatment of the cells with clasto-lactacystin β-lactone restored LATS1 protein levels in

HCV-infected cells (Fig. 1F, first panel, lane 6), indicating that HCV-induced LATS1 protein

reduction is due to proteasomal degradation. Collectively, these results indicate that HCV

promotes proteasomal degradation of LATS1 protein.

To determine whether HCV-induced LATS1 protein reduction is conserved among

different HCV genotypes, we utilized Huh-7 cells harboring the HCV genotype 1b genome-

length replicon (RCYM1). Immunoblot analyses revealed that LATS1 protein levels were

reduced compared to parental Huh-7 cells (Fig. 1G, first panel, lanes 2 and 4). These results

indicate that LATS1 protein is reduced in HCV-1b replicating cells.

E3 ligase Itch, but not WWP1, plays a crucial role in HCV-induced polyubiquitylation

and degradation of LATS1.

To determine whether HCV-induced LATS1 degradation depends specifically on Itch

ubiquitin ligase, we generated an inactive mutant Itch by mutating cysteine 868 to alanine

(pCAG-FLAG-Itch C868A), which abolishes E3 ligase activity. Huh-7.5 cells were co-

transfected with pCAG-Myc-LATS1, pRK5-HA-Ub, and either pCAG-FLAG-Itch or pCAG-

FLAG-Itch C868A. Immunoblot analysis revealed that LATS1 polyubiquitylation and

degradation were increased upon co-transfection with pCAG-FLAG-Itch (Fig. 2A, first and

second panel, lane 2) but not with the inactive mutant (Fig. 2A, first and second panel, lane

3). These results suggest that Itch mediates polyubiquitylation and degradation of LATS1.

To determine whether other E3 ubiquitin ligases in the HECT family, such as WWP1,

also play a role in the polyubiquitylation of LATS1, we performed a cell-based ubiquitylation

assay using wild-type pCAG-FLAG–WWP1 and the inactive mutant pCAG-FLAG–WWP1

C890A. Immunoblot analysis showed that LATS1 polyubiquitylation and degradation were

increased upon co-transfection with wild-type pCAG-FLAG-WWP1 (Fig. 2B, first and

second panel, lane 2), while the inactive mutant abolishes this effect (Fig. 2B, first and

second panel, lane 3).

To verify Itch's role in HCV-induced LATS1 degradation, we performed siRNA-

mediated knockdown of either Itch or WWP1 in HCV-infected cells. Immunoblot analysis

demonstrated that knockdown of Itch restored LATS1 protein levels (Fig. 2C, first panel,

lanes 2 and 4), whereas knockdown of WWP1 further decreased LATS1 protein levels (Fig.

2D, first panel, lanes 2 and 4). These findings suggest that Itch is responsible for HCV-

induced ubiquitin-dependent degradation of LATS1.

HCV infection promotes translocation of YAP1 from the cytoplasm to the nucleus.

LATS1 is the protein kinase that phosphorylates YAP1 at Ser127 and activates the Hippo

signaling pathway. Therefore, we hypothesized that HCV-induced degradation of LATS1

may suppress the Hippo pathway. To assess this hypothesis, we performed immunoblot

analysis of components of the Hippo pathway, LATS1 and YAP1, in mock- and HCV

J6/JFH1-infected Huh-7.5 cells. Immunoblot analysis revealed that HCV infection promoted

LATS1 degradation (Fig. 3A, first panel, lane 2), accompanied by a reduction in

phosphorylated LATS1 at Ser 909 (Fig. 3A, second panel, lane 2), suggesting that active

LATS1 is decreased in HCV-infected cells. The decrease in pLATS1 at Ser 909 subsequently

resulted in reduced phosphorylated YAP1 at Ser 127 (Fig. 3A, third panel, lane 2), whereas

total YAP1 levels remained unchanged (Fig. 3A, fourth panel, lane 2). These results suggest

that HCV infection reduced LATS1 and phosphorylated LATS1 at Ser 909, resulting in

decreased phosphorylated YAP1 at Ser 127, thereby inactivating the Hippo pathway.

To determine whether unphosphorylated YAP1 translocates to the nucleus, we

performed a cell fractionation assay coupled with immunoblot analysis. The cell fractionation

assay revealed an increased nuclear YAP1 in HCV-infected cells (Fig. 3B, first panel, lane 4)

compared to mock-infected cells (Fig. 3B, first panel, lane 3). Furthermore,

immunofluorescence staining revealed that YAP1 proteins were predominantly localized to

the nucleus in HCV-infected Huh-7.5 cells (Fig. 3C, second panel). The line scan analysis

showed that YAP1 exhibited cytoplasmic and nuclear distribution in mock-infected cells as

shown by the fluorescence and the corresponding line scan profile (Fig. 3C, first panel; Fig.

3D). In contrast, the YAP1 intensity peak overlapped with the nuclear signal peak in HCV-

infected cells, indicating YAP1 accumulation within the nucleus (Fig. 3C, second panel; Fig.

3E). These results suggest that HCV infection promotes LATS1 degradation, leading to

reduced YAP1 phosphorylation and facilitating YAP1 nuclear translocation, thereby

inactivating the Hippo pathway.

mRNA levels of YAP1 target genes are upregulated upon HCV infection.

To determine whether HCV-induced nuclear translocation of YAP1 affects downstream gene expression, we performed RT-qPCR to measure the mRNA levels of canonical YAP1 target genes, CYR61 and CTGF, in HCV-infected cells. RT-qPCR showed that HCV infection significantly increased the mRNA levels of CYR61 and CTGF (Fig. 4A and 4B).

To determine whether HCV-induced upregulation of CYR61 and CTGF depends specifically on YAP1 nuclear translocation, we used siRNA to knock down YAP1 mRNA in HCV-infected Huh-7.5 cells. YAP1 mRNA levels were successfully reduced by approximately 80% (Fig. 4C). The mRNA levels of CYR61 and CTGF in HCV-infected cells were significantly lower than in siYAP1-transfected HCV-infected cells (Fig. 4D and 4E). These results suggest that HCV promotes the upregulation of CYR61 and CTGF via YAP1 nuclear translocation. CYR61 and CTGF are known to play roles in tissue remodeling and cell proliferation, suggesting that the HCV-induced upregulation of CYR61 and CTGF genes contributes to HCV-related pathogenesis.

CYR61 and CTGF expression levels are elevated in patients with chronic HCV infection.

To further validate the HCV-induced upregulation of CYR61 and CTGF in a clinical setting, we analyzed publicly available transcriptomic data. Both CYR61 and CTGF mRNAs were significantly upregulated in HCV-infected individuals compared to healthy controls (Fig. 5A and 5B), confirming upregulation of CYR61 and CTGF mRNAs in HCV-infected patients.

In summary, we propose a model shown in Fig. 6. Under normal conditions, the Hippo pathway remains active when LATS1 phosphorylates YAP1, and phosphorylated YAP1 localizes in the cytoplasm. In HCV infection, HCV activates the ROS/JNK/Itch

359 pathway (18). Active Itch induces the polyubiquitylation and proteasomal degradation of

360 LATS1. The degradation of LATS1 inhibits YAP1 phosphorylation and promotes YAP1

361 nuclear translocation, thereby upregulating transcription of the YAP1 target genes, CYR61

362 and CTGF. HCV-induced suppression of the Hippo pathway plays a role in HCV

363 pathogenesis.

DISCUSSION

In this study, we found that HCV infection induced the degradation of LATS1 protein, a

serine/threonine kinase essential for regulating the tumor-suppressive Hippo pathway (Fig.

1). We demonstrated that HCV infection inactivated the Hippo pathway and facilitated YAP1

nuclear translocation, and that HCV infection induced the upregulation of canonical YAP1

target genes, CYR61 and CTGF (Figs. 3-5).

Both CYR61 and CTGF are crucial for hepatocellular carcinoma (HCC) progression,

exhibiting higher expression in HCC tissues than in normal liver tissues (27). CYR61 and

CTGF act as adhesive substrates that stimulate fibroblast signaling, enhancing angiogenesis

and wound healing (28). Notably, elevated CTGF levels in human HCCs have also been

linked to poor clinical outcomes (29). We obtained evidence suggesting that HCV infection

inactivates the Hippo pathway, leading to the upregulation of CYR61 and CTGF expression,

thereby contributing to HCV pathogenesis.

The NEDD4 subfamily of HECT-type E3 ubiquitin ligases is characterized by

multiple WW domains that specifically interact with the PPxY domain of LATS1. Notably,

certain members of this subfamily have been identified as key regulators of LATS1 protein

stability (19, 20, 30, 31). We hypothesized that the NEDD4 subfamily, such as Itch or

WWP1, is involved in HCV-induced polyubiquitylation and proteasomal degradation of

LATS1. We previously showed that HCV activates the ROS/JNK signaling pathway, which

activates Itch and facilitates polyubiquitylation of VPS4A, enhancing HCV release (18). Our

results suggest that Itch, but not WWP1, plays a crucial role in HCV-induced degradation of

LATS1 protein (Fig. 2). Both Itch and WWP1 contributed to LATS1 polyubiquitylation in

mock-infected cells. Importantly, siRNA-mediated knockdown of Itch restored LATS1

protein in HCV-infected cells. In contrast, siRNA-mediated knockdown of WWP1 further

reduced LATS1 protein levels. These findings suggest that Itch predominantly plays a role in

regulation of LATS1 protein during HCV infection, possibly because HCV activates Itch

activity. WWP1 depletion may trigger compensatory activation of Itch, leading to enhanced

LATS1 degradation rather than recovery.

Notably, condition-specific regulatory shifts have been observed in some proteins. For

example, the tumor suppressor p53 is primarily degraded by Murine Double Minute 2

(Mdm2) under normal conditions, whereas in HPV16 E6-positive cancer cells, p53

degradation occurs entirely via the E6-AP pathway instead of Mdm2 (32). These

observations raise the possibility that HCV infection predominantly facilitates Itch-mediated

ubiquitylation and degradation of LATS1 via as-yet-undetermined mechanisms. Further

investigation is required to elucidate the details of these regulatory mechanisms.

In this study, we demonstrated evidence suggesting that HCV infection promotes the

ubiquitin-dependent degradation of LATS1 protein. This degradation contributes to the

inactivation of the Hippo pathway, resulting in nuclear translocation of YAP1 and subsequent

upregulation of the YAP1 target genes CYR61 and CTGF genes. To our knowledge, this is

the first study to demonstrate that HCV-induced degradation of LATS1 leads to Hippo

pathway inactivation, thereby upregulating CYR61 and CTGF genes. These insights may

contribute to our understanding of the molecular mechanisms underlying HCV pathogenesis,

highlighting potential targets for therapeutic intervention.

AUTHOR STATEMENTS

Author contributions

C.M. and I.S. conceived and designed the experiments. S.MA and C.M. carried out most of the experiments. ZX., F.P., C.M., L.D., and T.A. assisted the constructions and the data analysis. D.N.M.A. performed bioinformatics analysis of public gene expression data. S.MA., C.M, and I.S. wrote the manuscript.

Conflicts of interest

The authors declare no conflict of interest.

Funding information

This research was supported by grants for Basic and Clinical Research on Hepatitis from the Japan Agency for Medical Research and Development (AMED) grant No. 23fk0210090s1203 and 20fk0210040s0703, a KAKENHI grant No. 20K07514 and 22K15470, and the Program for the Nurturing of Next Generation Leaders Guiding Medical Innovation in Asia, funded by the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan.

Acknowledgements

We sincerely appreciate Dr. C. M Rice (The Rockefeller University, New York) for generously providing the Huh-7.5 cells and pFL-J6/JFH1 plasmid. We also thank Y. Kozaki for her valuable secretarial assistance.

 FIGURE LEGENDS

 Figure 1. HCV induces proteasomal degradation of LATS1 protein. (A) Huh-7.5 cells

 were infected with HCV J6/JFH1 at a multiplicity of infection (MOI) of 2. Cells were

 cultured and harvested at 2 and 4 days post infection (dpi). Immunoblot analysis was

 performed using indicated antibodies. β-actin levels were used as a loading control. (B) Huh-

 7.5 cells were infected with HCV J6/JFH1 at an MOI of 2. Cells were cultured and harvested

 at 2 and 4 days dpi. Subsequently, total cellular RNA was extracted, and LATS1 mRNA

 levels were quantified by RT-qPCR. To normalize the LATS1 mRNA levels, GAPDH mRNA

 levels were used as an internal control. The value for day 2 in the mock-infected group was

 arbitrarily expressed as 1.0. Data are presented as means ± SEM from three independent

 experiments that yielded similar results; p-value < 0.05 (*) was significant, compared with

 the controls. (C) HCV-infected and mock-infected control cells were treated with 50 µg/mL

 cycloheximide (CHX) for 0, 6, 12, 18, and 24 hours. Immunoblot analysis was performed

 using indicated antibodies, and β-actin levels served as a loading control. (D) Specific signals

 were quantified by densitometry. The percentages of remaining LATS1 and β-actin at each

 time point were normalized to their respective basal level. Closed circles represent HCV-

 infected cells and open circles represent mock-infected control cells. Data shown are

 representative of three independent experiments that yielded similar results. (E) Huh-7.5 cells

 were infected with HCV J6/JFH1 at an MOI of 2, followed by treatment with Bafilomycin

 A1 for 12 hours. Immunoblotting was performed with indicated antibodies and β-actin levels

 served as a loading control. (F) Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of

 2. Cells were treated with 30µM Clasto-Lactacystin β-Lactone for 12 hours, harvested and

 analyzed by immunoblotting using the indicated antibodies. β-actin served as a loading

 control. The immunoblots are representative of three independent experiments that yielded

 similar results. (G) Huh-7 cells stably harboring the HCV genotype genome-length replicon

(RCYM1) and the parental Huh-7 cells were seeded and cultured for 2 or 4 days. Cells were

harvested and analyzed by immunoblotting using the indicated antibodies.

Figure 2. E3 ligase Itch, but not WWP1, plays a crucial role in HCV-induced

polyubiquitylation and degradation of LATS1 protein. (A) Huh-7.5 cells were co-

transfected with pCAG-Myc-LATS1, pRK5-HA-Ub, together with either pCAG-FLAG-Itch

WT or pCAG-FLAG-Itch C868A. At 2 days after transfection, the cells were harvested. The

cell lysates were immunoprecipitated with anti-c-Myc antibody, followed by immunoblotting

using anti-HA rabbit pAb and anti-c-Myc mouse mAb. Input samples were also analyzed by

immunoblotting using anti-c-Myc mouse mAb, and anti-FLAG mouse mAb. β-actin served

as a loading control. (B) Huh-7.5 cells were co-transfected with pCAG-Myc-LATS1, pRK5-

HA-Ubiquitin, together with either pCAG-FLAG-WWP1 WT or pCAG-FLAG-WWP1

C890A. At 2 days after transfection, cells were harvested. The cell lysates were

immunoprecipitated with anti-c-Myc antibody, followed by immunoblotting with anti-HA

rabbit pAb and anti-c-Myc mouse mAb. Input samples were also analyzed by

immunoblotting using anti-c-Myc mouse mAb, and anti-FLAG mouse mAb. β-actin served

as a loading control. (C) Huh-7.5 cells at 3×10^5 cells in a 12-well plate were transfected with

48 pmol of either control siRNA or Itch-specific siRNA. After 24 hours, the cells were

infected with HCV J6/JFH1 at an MOI of 2 and harvested at 2 days post-infection. The

samples were analyzed by immunoblotting with specified antibodies. β-actin served as a

loading control. (D) Huh-7.5 cells at 3×10^5 cells in a 12-well plate were transfected with 48

pmol of either control siRNA or WWP1-specific siRNA. At 24 hours after siRNA-

transfection, the cells were infected with HCV J6/JFH1 at an MOI of 2. Cells were harvested

at 2 days post-infection and the samples were subjected to immunoblotting with the indicated

antibodies. β-actin served as a loading control. The immunoblots are representative of three

independent experiments that yielded similar results.

Figure 3. HCV infection promotes the translocation of YAP1 from the cytoplasm to the

nucleus. (A) Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of 2. At 3 days after

infection, the cells were harvested and analyzed by immunoblotting using the indicated

antibodies targeting components of the Hippo pathway. β-actin served as a loading control.

(B) Huh-7.5 cells were infected with HCVJ6/JFH1 at an MOI of 2. At 3 days after infection,

the cells were harvested and subjected for cell fractionation analysis. IκBα served as a

loading control for the cytoplasmic fraction and Histone H3 served as a loading control for

the nuclear fraction. The immunoblots are representative of three independent experiments

that yielded similar results. (C) Huh-7.5 cells were plated and cultured for 12 hours and

infected with HCV J6/JFH1 at an MOI of 2. Cells were stained with anti-YAP1 monoclonal

antibody followed by Alexa Fluor 594-conjugated goat anti-mouse IgG (red) and anti-NS5A

polyclonal antibody followed by Alexa Fluor 488-conjugated goat anti-rabbit IgG (green).

Nuclei were counterstained with Hoechst 33342 (blue). Images were acquired using scanning

laser confocal microscopy and processed with ImageJ software. Scale bar: 10µm. The

immunofluorescence staining results are representative of three independent experiments. (D,

E) A straight line was drawn from the cytoplasm through the nucleus of a representative cell

in confocal immunofluorescence images from the mock (D) and HCV-infected (E) groups

using ImageJ. The corresponding line scan profiles display the fluorescence intensity (gray

value, arbitrary units) of YAP1 (red line) and Hoechst 33342 (blue line) along the measured

distance. The degree of overlap between the YAP1 and Hoechst 33342 signals indicates the

extent of YAP1 nuclear localization. Data shown are representative of three independent

experiments.

Figure 4. The mRNA levels of YAP1 target genes are significantly upregulated upon HCV infection. (A, B) Huh-7.5 cells were infected with HCV J6/JFH1 at an MOI of 2. Cells were cultured and harvested at 2 and 4 days post infection (dpi). Total cellular RNA was subsequently extracted, and the mRNA levels of CYR61 and CTGF were quantified by RT-qPCR. GAPDH mRNA levels were used for normalization. The mRNA level for day 2 in the mock-infected group was arbitrarily expressed as 1.0. Data are presented as means ± SEM from three independent experiments. The p-value < 0.05 (*) was significant. (C, D and E) Huh-7.5 cells at 1.8×10^5 cells in a twenty four-well plate were transfected with 24 pmol of either control siRNA or YAP1 siRNA. At 24 h after sinRNA-transfection, the cells were infected with HCV J6/JFH1 at an MOI of 2, cultured and harvested at the indicated time points. Total RNA was extracted, and YAP1, CYR61 and CTGF mRNA levels were quantified by RT-qPCR, normalized to GAPDH mRNA levels. The mRNA levels at day 2 in the mock-infected group was arbitrarily expressed as 1.0. Data are expressed as means ± SEM from three independent experiments. The p-value < 0.05 (*) was significant.

Figure 5. CYR61 and CTGF mRNA levels are increased in patients with chronic HCV infection. (A, B) CYR61 and CTGF mRNA levels in the liver tissue of patients with chronic HCV infection (n=38) compared to healthy individuals (n=6) (GEO accession number GSE 84346). Data are presented as the mean ± SD. The p-value < 0.05 (*) was significant.

Figure 6. A proposed model of HCV-induced inactivation of the Hippo pathway and HCV-related pathogenesis. Under normal condition, the Hippo pathway remains active. LATS1 phosphorylates YAP1, thereby retaining YAP1 in the cytoplasm and preventing its nuclear entry. HCV infection activates the ROS/JNK/Itch signaling pathway, which leads to

531   the phosphorylation and enhanced activity of Itch ubiquitin ligase. Activated Itch promotes

532   the ubiquitin-dependent proteasomal degradation of LATS1 protein, leading to the Hippo

533   pathway inactivation. As a result, YAP1 translocates from the cytoplasm to the nucleus and

534   upregulates the transcription of CYR61 and CTGF genes, contributing to HCV-related

535   pathogenesis.

REFERENCES

1.	Global Hepatitis Report 2024: Action for Access in Low- and Middle-Income Countries. 1st ed. Geneva: World Health Organization; 2024. 1 p.

2.	Bartenschlager R, Baumert TF, Bukh J, Houghton M, Lemon SM, Lindenbach BD, et al. Critical challenges and emerging opportunities in hepatitis C virus research in an era of potent antiviral therapy: Considerations for scientists and funding agencies. *Virus Res*. 2018;248:53–62.

3.	Baumert TF, Jühling F, Ono A, Hoshida Y. Hepatitis C-related hepatocellular carcinoma in the era of new generation antivirals. *BMC Med*. 2017;15(1):52.

4.	Scheel TKH, Rice CM. Understanding the hepatitis C virus life cycle paves the way for highly effective therapies. *Nat Med*. 2013;19(7):837–49.

5.	Sohn BH, Shim JJ, Kim SB, Jang KY, Kim SM, Kim JH, et al. Inactivation of Hippo Pathway Is Significantly Associated with Poor Prognosis in Hepatocellular Carcinoma. *Clin Cancer Res*. 2016;22(5):1256–64.

6.	Zheng T, Wang J, Jiang H, Liu L. Hippo signaling in oval cells and hepatocarcinogenesis. *Cancer Lett*. 2011;302(2):91–9.

7.	Zhao B, Li L, Lei Q, Guan KL. The Hippo–YAP pathway in organ size control and tumorigenesis: an updated version. *Genes Dev*. 2010;24(9):862–74.

8.	Zhao B, Li L, Tumaneng K, Wang CY, Guan KL. A coordinated phosphorylation by Lats and CK1 regulates YAP stability through SCF $^{\beta\text{-TRCP}}$. *Genes Dev*. 2010;24(1):72–85.

9.   Zhao B, Wei X, Li W, Udan RS, Yang Q, Kim J, et al. Inactivation of YAP

oncoprotein by the Hippo pathway is involved in cell contact inhibition and tissue growth

control. *Genes Dev*. 2007;21(21):2747–61.

10.   Meng Z, Moroishi T, Guan KL. Mechanisms of Hippo pathway regulation. *Genes*

Dev. 2016;30(1):1–17.

11.   Kwon H, Kim J, Jho E. Role of the Hippo pathway and mechanisms for controlling

cellular localization of YAP/TAZ. *FEBS J*. 2022;289(19):5798–818.

12.   Wang Z, Lu W, Zhang Y, Zou F, Jin Z, Zhao T. The Hippo Pathway and Viral

Infections. *Front Microbiol*. 2020;10:3033.

13.   Liu G, Yu FX, Kim YC, Meng Z, Naipauer J, Looney DJ, et al. Kaposi sarcoma-

associated herpesvirus promotes tumorigenesis by modulating the Hippo pathway. *Oncogene*.

2015;34(27):3536–46.

14.   Garcia G, Paul S, Beshara S, Ramanujan VK, Ramaiah A, Nielsen-Saines K, et al.

Hippo Signaling Pathway Has a Critical Role in Zika Virus Replication and in the

Pathogenesis of Neuroinflammation. *Am J Pathol*. 2020;190(4):844–61.

15.   Webb Strickland S, Brimer N, Lyons C, Vande Pol SB. Human Papillomavirus E6

interaction with cellular PDZ domain proteins modulates YAP nuclear localization. *Virology*.

2018;516:127–38.

16.   Xue Y, Mars WM, Bowen W, Singhi AD, Stoops J, Michalopoulos GK. Hepatitis C

Virus Mimics Effects of Glypican-3 on CD81 and Promotes Development of Hepatocellular

Carcinomas via Activation of Hippo Pathway in Hepatocytes. *Am J Pathol*.

2018;188(6):1469–77.

17.   Hu B, Xie S, Hu Y, Chen W, Chen X, Zheng Y, et al. Hepatitis C virus NS4B protein

induces epithelial-mesenchymal transition by upregulation of Snail. *Virol J*. 2017;14(1):83.

18.   Deng L, Liang Y, Ariffianto A, Matsui C, Abe T, Muramatsu M, et al. Hepatitis C

Virus-Induced ROS/JNK Signaling Pathway Activates the E3 Ubiquitin Ligase Itch to

Promote the Release of HCV Particles via Polyubiquitylation of VPS4. *J Virol*.

2022;96(6):e01811-21.

19.   Salah Z, Melino G, Aqeilan RI. Negative Regulation of the Hippo Pathway by E3

Ubiquitin Ligase ITCH Is Sufficient to Promote Tumorigenicity. *Cancer Res*.

2011;71(5):2010–20.

20.   Ho KC, Zhou Z, She YM, Chun A, Cyr TD, Yang X. Itch E3 ubiquitin ligase

regulates large tumor suppressor 1 stability. *Proc Natl Acad Sci*. 2011;108(12):4870–5.

21.   Wakita T, Pietschmann T, Kato T, Date T, Miyamoto M, Zhao Z, et al. Production of

infectious hepatitis C virus in tissue culture from a cloned viral genome. *Nat Med*.

2005;11(7):791–6.

22.   Bungyoku Y, Shoji I, Makine T, Adachi T, Hayashida K, Nagano-Fujii M, et al.

Efficient production of infectious hepatitis C virus with adaptive mutations in cultured

hepatoma cells. *J Gen Virol*. 2009;90(7):1681–91.

23.   Murakami K, Ishii K, Ishihara Y, Yoshizaki S, Tanaka K, Gotoh Y, et al. Production

of infectious hepatitis C virus particles in three-dimensional cultures of the cell line carrying

the genome-length dicistronic viral RNA of genotype 1b. *Virology*, 2006;351:381-92.

24.   Shirakura M, Murakami K, Ichimura T, Suzuki R, Shimoji T, Fukuda K, et al. E6AP

Ubiquitin Ligase Mediates Ubiquitylation and Degradation of Hepatitis C Virus Core Protein.

J Virol. 2007;81(3):1174–85.

25.   Matsui C, Shoji I, Kaneda S, Sianipar IR, Deng L, Hotta H. Hepatitis C Virus

Infection Suppresses GLUT2 Gene Expression via Downregulation of Hepatocyte Nuclear

Factor 1α. *J Virol*. 2012;86(23):12903–11.

26.   Yuliandari P, Matsui C, Deng L, Abe T, Mori H, Taguwa S, et al. Hepatitis C virus

NS5A protein promotes the lysosomal degradation of diacylglycerol O-acyltransferase 1

(DGAT1) via endosomal microautophagy. *Autophagy Rep*. 2022;1(1):264–85.

27.   Zeng ZJ. Expressions of cysteine-rich61, connective tissue growth factor and Nov

genes in hepatocellular carcinoma and their clinical significance. *World J Gastroenterol*.

2004;10(23):3414.

28.   Chen CC, Chen N, Lau LF. The Angiogenic Factors Cyr61 and Connective Tissue

Growth Factor Induce Adhesive Signaling in Primary Human Skin Fibroblasts. *J Biol Chem*.

2001;276(13):10443–52.

29.   Simile MM, Latte G, Demartis MI, Brozzetti S, Calvisi DF, Porcu A, et al. Post-

translational deregulation of YAP1 is genetically controlled in rat liver cancer and determines

the fate and stem-like behavior of the human disease. *Oncotarget*. 2016;7(31):49194–216.

30.   Yeung B, Ho KC, Yang X. WWP1 E3 Ligase Targets LATS1 for Ubiquitin-Mediated

Degradation in Breast Cancer Cells. Viglietto G, editor. *PLoS ONE*. 2013;8(4):e61027.

31.   Salah Z, Cohen S, Itzhaki E, Aqeilan R. NEDD4 E3 ligase inhibits the activity of the

Hippo pathway by targeting LATS1 for degradation. *Cell Cycle*. 2013;12(24):3817–23.

32.   Hengstermann A, Linares LK, Ciechanover A, Whitaker NJ, Scheffner M. Complete

switch from Mdm2 to human papillomavirus E6-mediated degradation of p53 in cervical

cancer cells. *Proc Natl Acad Sci U S A*. 2001;98(3):1218-23.

Fig. 1 Maria et al.

Fig. 2 Maria et al.

Fig. 4 Maria et al.

Normal condition

HCV infection