Title: Autism-associated *Scn2a* haploinsufficiency disrupts *in vivo* dendritic signaling and

impairs flexible decision-making

Authors: Hao Wu[1,2], Luqun Shen[1], Jonathan Indajang[1], Neil K. Savalia[1,2,3], Timothy G.

Johnson[1], Jiayin Qu[1], Kevin J. Bender[4], Alex C. Kwan[1,5,*]

Affiliations:

[1]Meinig School of Biomedical Engineering, Cornell University, Ithaca, NY, 14853, USA

[2]Interdepartmental Neuroscience Program, Yale University School of Medicine, New Haven, CT,

06511, USA

[3]Medical Scientist Training Program, Yale University School of Medicine, New Haven, CT,

06511, USA

[4]Center for Integrative Neuroscience, Department of Neurology, University of California, San

Francisco, CA, 94158, USA

[5]Department of Psychiatry, Weill Cornell Medicine, New York, NY, 10065, USA

[*]Correspondence to Alex Kwan, Ph.D., Room 111 Weill Hall, 526 Campus Road, Ithaca, NY,

14853, United States; E-mail: alex.kwan@cornell.edu

Keywords: sodium channel, calcium, dendrite, pyramidal cell, frontal cortex, behavioral

flexibility

ABSTRACT

SCN2A is a high-confidence risk gene for autism spectrum disorder. Loss-of-function mutations in *Scn2a* reduce dendritic excitability in neocortical pyramidal cells. However, the impact of *Scn2a* haploinsufficiency on dendritic signaling *in vivo*, particularly during behavior, is unknown. In this study, we used two-photon microscopy to image dendritic calcium transients in deep layer pyramidal cells in the mouse medial frontal cortex. *Scn2a*[+/-] mice had diminished coupling between apical and proximal dendritic compartments. Pyramidal tract neurons had abnormal event rates, while intratelencephalic neurons had compartment-specific alterations indicative of diminished dendritic integration. In a matching pennies task, *Scn2a*[+/-] mice were inflexible in the face of changing competitive pressure. Apical dendritic tuft in IT neurons typically encoded reward and strategy, but these task-specific representations were absent in *Scn2a*[+/-] mice. Collectively, the findings demonstrate that *Scn2a* haploinsufficiency weakens dendritic integration *in vivo* and disrupts the dendritic encoding of decision variables, potentially contributing to the cognitive rigidity in autism spectrum disorder.

INTRODUCTION

Autism spectrum disorder (ASD) is a neurodevelopmental condition characterized by deficits in perception, social interaction, and cognition. At the genetic level, studies of rare coding variants have identified numerous high-confidence risk genes for ASD, many of which are involved in signaling and scaffolding in the postsynaptic dendritic compartments[1-3]. Although the etiology of ASD remains poorly understood, growing evidence implicates deep layer glutamatergic neurons in the neocortex as key contributors[4]. Alterations in excitatory transmission have been observed in many ASD models[5], and the synaptic dysfunctions may be closely tied to abnormalities in dendritic integration[6]. Supporting this connection, structural alterations to dendrites and dendritic spines have been reported in animal models for ASD[7,8] or autism related disorders such as Fragile X and Angelman syndrome[9,10]. However, uncovering the precise mechanisms by which the genetic perturbations lead to dendritic dysfunctions and ultimately behavioral deficits, remain an ongoing challenge.

Among the most strongly implicated high-confidence risk genes for ASD is *SCN2A*[2,3]. Loss-of-function mutations in *SCN2A* are associated with ASD and intellectual disability, whereas other alterations including gain-of-function mutations are more commonly associated with early onset epilepsy[11-13]. *SCN2A* encodes $Na_v1.2$, the type II alpha subunit of a voltage-gated sodium channel found in many cell types and brain regions. In the excitatory pyramidal cells in the

neocortex, Na$_v$1.2 is initially localized to the axon initiation segment during early development,

but later becomes the dominant isoform expressed in the somatodendritic region in

adulthood[14,15]. A function of Na$_v$1.2 is to facilitate membrane excitability. Indeed, in mice with

Scn2a haploinsufficiency, a disruption of Na$_v$1.2 expression reduces dendrite excitability and

impairs synaptic plasticity[14]. Specifically, when the cell body is stimulated, calcium influx

associated with the backpropagation of action potentials is diminished, reflecting as a decrease

in the amplitude in the apical dendritic shaft, and inability to invade into the distal dendritic tuft[14].

While these results highlight the importance of *Scn2a* for dendritic function, the studies were

performed in acute brain slices. The extent to which these deficits may manifest *in vivo* and their

behavioral relevance are open questions.

Recent advances in two-photon microscopy have provided powerful new approaches for

visualizing calcium signals in the dendrites of cortical pyramidal neurons *in vivo*. The

optimization of genetically encoded calcium indicators has yielded reporters with high signal-to-

noise ratios, enabling imaging in subcellular compartments of neurons in live mice[16]. This

improvement has led to exciting discoveries about how synaptic inputs may be propagated and

integrated in dendritic compartments. For example, studies have mapped the feature selectivity

of inputs impinging at different dendritic locations[17-20], and revealed choice- and reward-related

dendritic signals in the sensorimotor cortex during decision-making tasks[21-25]. A particularly

valuable innovation is the development of multi-plane imaging. This technique relies on rapid

axial focusing via one of several possible methods such as electrically tunable lens, acousto-

optic deflectors, and remote focusing. Multi-plane imaging enables near simultaneous recording

of fluorescence transients at several depths, which is especially useful for quantifying the

coupling between multiple compartments of a neuron including various dendritic branches and

the soma *in vivo*[21,26,27].

The goal of this study is to determine the effects of *Scn2a* haploinsufficiency on dendritic

calcium signaling *in vivo*. Performing multi-plane imaging using an electrically tunable lens, we

visualized calcium transients in the distal apical tuft, apical trunk, and proximal trunk of

pyramidal neurons in the mouse medial frontal cortex. We found reduced coupling of dendritic

calcium signals in *Scn2a*$^{+/-}$ animals relative to controls. To assess how *Scn2a* haploinsufficiency

affects behavior, we developed a competitive decision-making task for head-fixed mice that

encourages adaptive strategy switching. *Scn2a*$^{+/-}$ mice were cognitively rigid, failing to adjust to

a computer opponent that changes its strategy. During this task, apical dendritic tuft in the

frontal cortex normally encoded reward- and strategy-related signals, but these representations

were absent in *Scn2a*[+/-] mice. Overall, the results suggest the dendritic integration, normally

mediated by $Na_v1.2$, is critical for flexible decision-making, and that these processes are

impaired in *Scn2a* haploinsufficiency.

RESULTS

Multi-plane imaging of dendritic calcium transients in frontal cortical PT and IT neurons

To study *Scn2a* haploinsufficiency, we compared heterozygous *Scn2a*[+/-] mice to littermate

Scn2a[+/+] controls. Our goal was to characterize dendritic calcium transients in pyramidal cells in

the medial frontal cortex (**Fig. 1A**). There are two major subpopulations of pyramidal cells:

pyramidal tract (PT) and intratelencephalic (IT) neurons. PT and IT neurons have distinct

features; they have different morphological characteristics, physiological properties, and long-

range projection targets[28-30]. Moreover, PT and IT neurons may have differential contributions to

neuropsychiatric conditions and their treatments[31-33]. To target a sparse number of PT neurons,

we injected a low titer of AAVrg-hSyn-Cre into the ipsilateral pons, and AAV-CAG-Flex-GCaMP6f

in the dorsal medial frontal cortex including the anterior cingulate cortex (ACAd) and medial

premotor cortex (MOs) (**Fig. 1B**). Fluorescence images of histological coronal sections

confirmed a sparse labeling of neurons restricted to the deep layers of the medial frontal cortex,

consistent with the expected laminar distribution of PT neurons (**Fig. 1B**).

Fig. 1. Imaging calcium transients in the dendritic compartments of single frontal cortical PT or IT neuron.

(A) The imaging setup involving two-photon imaging of an awake, head-fixed mouse. ETL, electrically tunable lens.

(B) Left, viral strategy to express GCaMP6f in frontal cortical PT neurons. Right, fixed coronal section showing GCaMP6f-expressing cells in the medial frontal cortex.

(C) Similar to (B) for frontal cortical IT neurons.

(D) Example two-photon images at different depths corresponding to apical tuft, apical trunk and proximal trunk of a frontal cortical PT neuron in a $Scn2a^{+/+}$ control mouse. Dash lines indicate the region of interest.

(E) ΔF/F traces from one ROI each from apical tuft, apical trunk and proximal trunk of a PT neuron in a $Scn2a^{+/+}$ control mouse. Right, magnified view of the boxed area.

(F) Similar to (E) for a $Scn2a^{+/-}$ mouse.

(G – I) Similar to (D – F) for a frontal cortical IT neuron.

ACAd, anterior cingulate area dorsal part. MOs, secondary motor cortex. CP, caudate putamen.

We imaged spontaneous calcium transients from apical dendritic compartments in the awake,

head-fixed mouse. We used two-photon microscopy, with the addition of an electrically tunable

lens (ETL) placed above the back aperture of the objective. By applying different electrical

currents, the ETL can flex quickly, which allows for a rapid change of the focal plane[34]. Prior to

time-lapse imaging, we would acquire a z-stack from the pial surface to -500 µm with a step size

of 5 µm to identify dendritic segments that belong to the same cell. Then time-lapse imaging

commenced, which lasted for 30 minutes. For each neuron, we would capture fluorescence

transients at three focal planes, targeting one depth each for apical tuft (-50 to -150 µm), apical

trunk (-250 to -400 µm), and proximal dendrite (<-400 µm) (**Fig. 1D**). The overall frame rate was

7.93 Hz. For analysis, we selected regions of interest (ROIs) corresponding to dendritic

segments (see **Methods**). We calculated the fractional change in fluorescence $\Delta F/F$ for each

ROI.

In control animals, we saw that the dendritic calcium transients between different compartments

of a frontal cortical PT neuron were generally synchronized (**Fig. 1E**). That is, whenever there

was a calcium transient in one compartment (e.g., an ROI in the plane of apical tuft), we would

observe a corresponding calcium transient in other compartments (e.g., ROIs for the apical

trunk and the proximal trunk). However, there were also cases when transients were observed

in one compartment but not another, such as the leftmost calcium event in the proximal trunk in

Fig. 1E, which was not detectable in the apical trunk or apical tuft. We noted the signal-to-noise

was typically worse in the ROIs at the apical tuft compared to the apical trunk and proximal

trunk (inset, **Fig. 1E**), which was expected given the few fluorophores within the small volume of

a dendritic segment in the apical tuft. Relatedly, the decay kinetics for the calcium transients

was noticeably longer in the apical and proximal trunks, also consistent with these

compartments having larger volume and thus needed more time for the clearance of calcium

after an influx event[35]. **Fig. 1F** shows example dendritic calcium transients imaged from a frontal

cortical PT neuron in a *Scn2a*$^{+/-}$ mouse.

Using the same imaging setup, we also characterized the spontaneous calcium transients in the

dendrites of frontal cortical IT neurons in the awake, head-fixed mouse. To target a sparse

number of IT neurons, we injected a low titer of AAVrg-hSyn-Cre into the contralateral striatum

and AAV-CAG-Flex-GCaMP6f in the medial frontal cortex (**Fig. 1C**). Fluorescence images of

fixed coronal sections showed sparse labeling of neurons in both superficial and deep layers of

the medial frontal cortex, in agreement with the broader distribution of IT neurons (**Fig. 1C**).

Because IT neurons tended to have cell bodies that lie more superficially, we targeted the three

focal planes slightly differently for apical tuft (-50 to -150 µm), apical trunk (-150 to -350 µm),

and proximal dendrite (<-350 µm) (**Fig. 1G**). **Fig. 1H** shows dendritic calcium transients imaged

from a frontal cortical IT neuron in a control animal. **Fig. 1I** shows an example from a *Scn2a*$^{+/-}$
mouse.

Scn2a* haploinsufficiency impairs dendritic coupling in PT and IT neurons *in vivo
Although the calcium transients were mostly synchronized across the different dendritic
compartments, we also occasionally observed isolated transients in one compartment only. To
quantify the coupling between dendrites, we calculated conditional probabilities. Starting from
the Δ*F/F(t)* for each ROI, we used the semi-automated OASIS algorithm to infer the underlying
electrical events[36]. Then, for instance, to determine the amount of transmission from the apical
trunk up to the apical tuft, we would consider each event in the apical trunk and tabulate if there
was a corresponding event in the apical tuft occurring within ±3 imaging frames (see **Methods**).
In an example drawn from actual data (**Fig. 2A**), there were 20 events detected in the apical
trunk including 2 events that lacked a counterpart in the apical tuft, hence the conditional
probability was 0.9. We considered pairwise correlation as another metric that could be used to
estimate coupling, but correlation coefficient may be prone to errors because the calcium
transients in different dendritic compartments have different decay kinetics.

For PT neurons, we imaged 23 cells in 6 *Scn2a*$^{+/+}$ mice (5 males, 1 female) and 31 cells in 6
Scn2a$^{+/-}$ mice (3 males, 4 females). We calculated conditional probabilities to estimate the
coupling between all imaged ROIs (**Fig. 2B – G**). We note that some ROIs could come from the
same imaging plane (e.g., several ROIs from the same plane for apical tuft of a neuron),
therefore for statistical analysis, we used a mixed effects model, which included random effects
terms to account for the hierarchical structure of the data where ROIs come from dendrites that
may originate from the same neuron and mouse (**Table S1**). In the backward direction, the
coupling between the proximal trunk and apical tuft was significantly reduced in *Scn2a*$^{+/-}$ mice
compared to controls (*Scn2a*$^{+/+}$: 0.75±0.04; *Scn2a*$^{+/-}$: 0.61±0.02; main effect of genotype: *P* =
0.015, mixed effects model; **Fig. 2B**). The other significant effect associated with *Scn2a*
haploinsufficiency was reduced coupling between the apical trunk and the apical tuft (*Scn2a*$^{+/+}$:
0.85±0.02; *Scn2a*$^{+/-}$ mice: 0.77±0.02; main effect of genotype: *P* = 0.005, mixed effects model;
Fig. 2D). There was no detected difference in the forward direction for dendritic integration in
frontal cortical PT neurons between *Scn2a*$^{+/-}$ and control mice (**Fig. 2E – G**).

Fig. 2. *Scn2a* haploinsufficiency impairs dendritic coupling in PT and IT neurons *in vivo*.

(A) Example ΔF/F traces and the corresponding inferred events from the apical tuft and apical trunk of a PT

neuron. Asterisk indicates one inferred event, or else the number indicates the number of inferred events. Red patch, time window centered around an event in apical trunk with no corresponding event in apical tuft.

(B) Conditional probability of an event in apical tuft given an event in the proximal trunk in PT neurons for $Scn2a^{+/+}$ control animals and $Scn2a^{+/-}$ mice. Each point is a ROI. Mean±SEM.

(C) Similar to (B) for event in apical trunk given event in proximal trunk.

(D) Similar to (B) for event in apical tuft given event in apical trunk.

(E) Similar to (B) for event in proximal trunk given event in apical tuft.

(F) Similar to (B) for event in proximal trunk given event in apical trunk.

(G) Similar to (B) for event in apical trunk given event in apical tuft.

(H) – (M) Similar to (B) – (G) for IT neurons.

*, $P<0.05$. **, $P<0.01$.

For IT neurons, we imaged 12 cells in 7 $Scn2a^{+/+}$ mice (4 males, 3 females) and 13 cells in 5

$Scn2a^{+/-}$ mice (1 male, 4 females). We found reduced coupling between dendritic compartments

in both forward and backward directions (**Fig. 2H – M**). Similar to PT neurons, we found a

significant deficit in the coupling between the proximal trunk and apical tuft in the backward

direction ($Scn2a^{+/+}$: 0.87±0.02; $Scn2a^{+/-}$: 0.71±0.06; main effect of genotype: $P = 0.014$, mixed

effects model; **Fig. 2H**). For the forward direction, propagation of dendritic signal was impaired

from the apical tuft to the proximal trunk ($Scn2a^{+/+}$: 0.86±0.03; $Scn2a^{+/-}$: 0.63±0.06; main effect

of genotype: $P = 0.002$, mixed effects model; **Fig. 2K**), and from the apical trunk to the proximal

trunk ($Scn2a^{+/+}$: 0.90±0.02; $Scn2a^{+/-}$: 0.74±0.04; main effect of genotype: $P = 0.001$, mixed

effects model; **Fig. 2L**). Altogether, these results showed that, in many cases, the coupling

between dendritic compartments reduced with the heterozygous loss of $Na_V1.2$. The coupling

was most adversely impacted between the apical tuft and the proximal trunk, which span the

greatest distance across the PT and IT neurons in our analysis.

From the events inferred from $\Delta F/F$ for each ROI, we could determine the rate of the dendritic

calcium transients. We found that there were cell type and compartment specific differences in

the dendritic calcium transients (genotype × cell type interaction: $P = 0.021$, compartment × cell

type interaction: $P = 0.0003$, mixed effects model; **Table S1**). For PT neurons, their dendrites

were overall more active in $Scn2a^{+/-}$ mice than control animals, irrespective of the dendritic

compartment ($P = 5 \times 10^{-7}$, Bonferroni *post hoc* test; **Fig. S1A**). For frontal cortical IT neurons,

there was no main difference across genotypes ($P = 0.8$, Bonferroni *post hoc* test), but the

frequency of calcium events varied across the dendritic compartments (**Fig. S1B**). Namely, we

found that in $Scn2a^{+/-}$ animals, the calcium event rate was higher in the apical tuft and apical

trunk than the proximal trunk (apical tuft vs. proximal trunk: $P = 0.033$, apical trunk vs. proximal

trunk: $P = 0.035$, apical tuft vs. apical trunk: $P = 0.6$, Bonferroni *post hoc* test). The reduced event

rate in the proximal trunk is consistent with the deficient forward transmission in IT neurons in

Scn2a[+/-] animals (**Fig. 2K, L**). This analysis demonstrates an aberrant elevation in dendritic calcium

events specifically in the PT neuron subpopulation because of *Scn2a* haploinsufficiency, which may

compensate for the deficient coupling across dendritic compartments.

Scn2a-haploinsufficient mice were inflexible against changing behavioral strategies

So far, our results have revealed how *Scn2a* haploinsufficiency affects *in vivo* spontaneous

activity in the dendrites, but we also want to know the potential alterations during behavior. The

dorsal medial frontal cortex in rodents is involved in adaptive decisions involving behavioral

flexibility[37]. We therefore tested the animals on various decision-making paradigms. Initially, we

trained *Scn2a*[+/-] mice to perform a probabilistic reward reversal task, but observed no alteration

in performance compared to control animals[38]. Next, we trained *Scn2a*[+/-] mice to play a standard

version of the matching pennies game that we previously adapted for head-fixed mice[39,40].

Matching pennies is a classic two-playing competitive game[41]. In this standard version, the two

players are the mouse and a computer, with the computer programmed to always play with the

same strategy. We again found that *Scn2a*[+/-] mice performed to a level that was not

distinguishable from control animals (**Fig. S2**). We surmised that these tasks may not

sufficiently assess flexible behavior, because the animal can rely on the same behavioral

strategy for the entire task. For the probabilistic reward reversal, mice can excel by using a

finite-state approach, e.g. choosing the high-reward option repeatedly, until encountering a few

reward omissions, then switching to the other option[42,43]. For matching pennies, since the

computer always used the same strategy, animals could counter by employing a strategy of

randomly choosing between the two options[39,44]. Instead, ASD-related cognitive deficits may be

accentuated in tasks that encourage flexibility in behavioral strategies.

We therefore designed a new task, still based on matching pennies, but the animal face a

computer opponent that can switch strategies. In other words, the rules of the game – payoff

matrix and trial timing – always remain the same, but the innovation is that we programmed

three different strategies for the computer opponent. Briefly, a head-fixed mouse would play

against a computer opponent (**Fig. 3A**). Each trial, the mouse would indicate a choice between

the left and right options using a directional tongue lick. The computer would also choose one of

the options. The payoff matrix is shown in **Fig. 3B**. If the mouse and computer chose the same

option (i.e., both picked left or both picked right), then the mouse would receive a reward of ~4

µL of water. If the mouse and computer chose different options (i.e., mouse picked left and

computer picked right, or vice versa), then the mouse would have no reward. A trial began when

a motor actuator advanced the two lick spouts to just in front of the mouse's head (**Fig. 3C**).

Then a sound cue was played, which was followed by a 2-s response window. Depending on

the mouse's choice (first lick during the response window) and the payoff matrix, water would be

delivered immediately if the mouse won the trial. Subsequently, the motor actuator would retract

and there was a random inter-trial interval between each trial. The computer can play with one

of three possible strategies. The "predictive" strategy is a strong player that would play by

predicting the mouse's next choice based on the history of past choices and outcomes in that

session (same as the strategy used in ref. [39], algorithm 2 in ref. [45], and competitor 1 in ref. [44];

see **Methods**). The "WSLS" strategy stands for win-stay, lose-switch, where the computer

would pick the same option if it won the last trial or switch options if it lost the last trial. Finally,

the "alternate" strategy is when the computer would choose left, right, left, right, and so on,

regardless of the mouse's previous choices and outcomes.

Fig. 3. *Scn2a*-haploinsufficient mice were inflexible against changing behavioral strategies.

(A) Behavioral setup.

(B) The payoff matrix of the matching pennies game. 1 denotes a reward for the mouse, and 0 denotes no reward.

(C) The trial timing.

(D) An example session when a mouse played matching pennies with a computer opponent that changed strategies between predictive (pink shading) and WSLS (green shading). Top row, the strategy used by the computer. Middle row, the mouse's choices and outcomes. Bottom row: computer's choices. Black line, reward.

(E) The switch probability during predictive and WSLS trials for *Scn2a*[+/+] control mice. Each point represents the mean value of a mouse. Red crosshair, Mean±SEM.

(F) Similar to (E) for *Scn2a*[+/-] mice**.**

(G) Bar plot of the switch probability when the computer used predictive strategy. Each point represents the mean value of a mouse. Mean±SEM.

(H) Similar to (G) when the computer used WSLS strategy.

(I) Bar plot of the switch probability difference between predictive trials and WSLS trials. Each point represents the mean value of a mouse. Mean±SEM.

(J) – (O) Similar to (D) – (I) for a computer opponent that changed strategies between predictive (pink shading) and alternate (brown shading). Note that for panel O, there is an outlier for *Scn2a*[+/+] mice at 36.6 not included in the plot.

We trained *Scn2a*[+/-] and control mice to play the matching pennies game against a computer

opponent using the predictive strategy until the animals became proficient (>35% reward rate

for 3 consecutive days; see **Methods and Materials**). Then we tested them on the matching

pennies game with the exact same rules, but now against a computer opponent that switched its

strategy every 90 – 110 trials during a session. **Fig. 3D** shows a control mouse playing

matching pennies when the computer opponent started with the predictive strategy, switched to

WSLS, and then toggled for the remainder of the session. We characterized the animals'

performance in 130 sessions from 38 *Scn2a*[+/+] mice (16 males, 22 females) and 108 sessions

from 25 *Scn2a*[+/-] mice (15 males, 10 females). The control animals had a probability of switching

their option at 36.5±1.5% during predictive blocks and 39.5±2.0% during WSLS blocks (**Fig.**

3E). The *Scn2a*[+/-] animals had a probability of switching their option at 35.6±1.2% during

predictive blocks and 35.0±1.07% during WSLS blocks (**Fig. 3F**). Although the differences in

performance on aggregate was obscured by variability across individuals (**Fig. 3G, H**), when we

specifically analyzed how each animal adapted by calculating its switching probability during

predictive block relative to during WSLS block, we detected a significant impairment for the

Scn2a[+/-] mice to adjust to the changing strategy (*Scn2a*[+/+]: -2.9±0.9%; *Scn2a*[+/-]: 0.6±0.9%; $P =$

0.032, Wilcoxon rank sum test; **Fig. 3I**). The majority of control animals switched more during

WSLS blocks than predictive blocks, although this was a trend and not significant ($P = 0.07$,

sign test), unlike *Scn2a*[+/-] animals ($P = 1$, sign test). This is a beneficial adaptation for the control

animals, because a player should switch more during WSLS. In fact, to maximize reward rate,

an optimal player should be switching half the time against the predictive strategy but switch all

the time against the WSLS strategy.

We also trained animals playing matching pennies when the computer opponent was swapping

between the predictive and alternate strategies (**Fig. 3J**). We had a smaller data set, which

included 60 sessions from 15 $Scn2a^{+/+}$ mice (7 males, 8 females) and 38 sessions from 8

$Scn2a^{+/-}$ mice (4 males, 4 females). The control animals had a probability of switching their

option at 34.1±1.9% during predictive blocks and 24.8±2.8% during alternate blocks (**Fig. 3K**).

The $Scn2a^{+/-}$ animals had a probability of switching their option at 36.5±1.8% during predictive

blocks and 33.3±2.7% during WSLS blocks (**Fig. 3L**). In this case, when we quantified how

each animal adapts by calculating its switching probability during the predictive blocks relative to

during the alternate blocks, we did not detect a significant difference between the genotype

($Scn2a^{+/+}$: 9.3±2.6; $Scn2a^{+/-}$: 3.2±1.2; P = 0.11, Wilcoxon rank sum test; **Fig. 3M-O**), likely owing

to the small sample size. Nevertheless, we note that the control animals switched less during

alternate blocks than predictive blocks (P = 0.007, sign test), whereas the $Scn2a^{+/-}$ animals

again did not adapt (P = 0.3, sign test). This shift in behavior for control mice is consistent with a

previous study, which found that when the computer's choice is not contingent on the animal's

decision, the animal would just stick to the same option repeatedly[45].

Altogether, these results demonstrate that control mice can adjust their behavior while playing

matching pennies with computer opponents employing different strategies. Notably, they

switched more frequently against the WSLS strategy but less against the alternate strategy,

showing behavioral flexibility in both directions from their likely over-trained responses to the

predictive strategy. $Scn2a^{+/-}$ mice, by contrast, showed significantly reduced adaptability, and

did not deviate from their initial trained behavior.

Task-related signals in the apical dendritic tuft of IT neurons

Although we observed reduced dendritic coupling in both PT and IT neurons in $Scn2a^{+/-}$ mice,

the loss of dendritic signaling in the forward direction was specific to IT neurons, suggesting a

potential deficit in the integration of task-related signals during behavior. We therefore

hypothesize that dendritic dysfunction in IT neurons may relate to the inflexibility in decision-

making in $Scn2a^{+/-}$ mice. To investigate this possibility, we imaged dendritic calcium signals

while the mouse was playing the matching pennies game against a computer opponent

switching between the predictive and WSLS strategies (**Fig. 4A**). The dual viral injection

strategy used for **Fig. 1** enabled imaging of the same dendritic tree from single neurons,

however the sparse labeling approach had a low success rate, often leading to zero or dense

labeling. Here, to increase the yield because we also need to invest time to train animals, we

leveraged the Cre-driver line *PlexinD1-2A-CreER* to target cortical IT neurons[46]. We injected

AAV-CAG-Flex-GCaMP6f into the ACAd and medial MOs portion of the medial frontal cortex of

PlexinD1-CreER$^{+/-}$::Scn2a$^{+/-}$ mice and *PlexinD1-CreER$^{+/-}$::Scn2a$^{+/+}$* control animals (**Fig. 4B**).

We administered tamoxifen to induce Cre-mediated expression of GCaMP6f. We focused on

imaging dendritic segments from the apical tuft located 50 – 100 µm below the pial surface (**Fig.**

4C).

We observed task-related signals in the apical dendritic tuft. **Fig. 4D** shows Δ*F/F* from two

example ROIs of a control mouse, aligned to the time of sound cue at the start of a trial.

Fluorescence increased in ROI 1 when the animal chose left and received a reward, relative to

all other conditions. This indicates a preference for the choice *x* outcome interaction, specifically

for left and reward. ROI 2 exhibited higher fluorescence when there was a reward together with

either a left or right choice, which reflected a reward-related signal. **Fig. 4E** shows two example

ROIs from a *PlexinD1-CreER$^{+/-}$::Scn2a$^{+/-}$* mouse. The Δ*F/F* transients at around -1 s and again

at 3 s corresponded to the onset times when the motorized actuator was advancing and

retracting the lick spouts. These were task-related signals but not selective for specific choice or

outcome.

Fig. 4. Task-related signals in the apical dendritic tuft of IT neurons.

In total, we imaged 389 ROIs during 18 sessions from 4 *PlexinD1-CreER*[+/-]*::Scn2a*[+/+] mice (2

males, 2 females) and 286 ROIs during 17 sessions from 4 *PlexinD1-CreER*[+/-]*::Scn2a*[+/-] mice (4

males). To quantify the fraction of ROIs that exhibited task-related signals, we determined

whether there was a significant difference in $\Delta F/F$ between the cue period (t = -0.5 – 0.5 s) and

baseline (t = -2 – -1 s) or between the outcome period (t = 1 – 2 s) and baseline (t = -2 – -1 s).

We found that most ROIs in the apical tuft had task-related signals, with the proportions not

different for animals with the partial loss of *Scn2a* than controls for cue period (control:

74.3±4.3%; *PlexinD1-CreER*[+/-]*::Scn2a*[+/-]: 60.2±7.5%; *P* = 0.1, Wilcoxon rank sum test) or

outcome period (control: 55.0±6.2%; *PlexinD1-CreER*[+/-]*::Scn2a*[+/-]: 47.8±7.7%; *P* =0.6, Wilcoxon

rank sum test; **Fig. 4F**). This analysis indicates that most dendritic segments of frontal cortical IT

neurons had task-related calcium transients while the mice were engaged in the matching

pennies task.

Reward- and strategy-related signals were absent in the dendrites of *Scn2a*-

haploinsufficient mice

To characterize the factors that account for the calcium signals in the apical tuft of IT neurons

during matching pennies, we used multiple linear regression. For each ROI, we fitted a

regression model to determine how the fluorescence transient $\Delta F/F(t)$ relates to the choices,

outcomes, and their interactions for the current and past trials, as well strategies and their

interactions (**Fig. 5A**). The regression analysis revealed that ~7% of the ROIs were modulated

by the outcome of the current trial in control animals (**Fig. 5B**). Unlike the reward-related spiking

activity changes in medial frontal cortex that can last for a few trials[47-49], here we observed that

the reward-related dendritic calcium signals were transient and largely restricted to the current

trial (i.e., R(n) and not R(n-1)). The strategy of the computer opponent was robustly

represented, across all trial times and especially around the cue when decisions were made. By

contrast, in *PlexinD1-CreER+/-::Scn2a+/-* mice, the reward- and strategy-related signals were

absent in the apical tuft of IT neurons (**Fig. 5C**). Therefore, although there was task-modulated

$\Delta F/F$ in the apical tuft of IT neurons, there was no selectivity for reward and strategy variables in

animals with *Scn2a* haploinsufficiency.

Fig. 5. Reward- and strategy-related signals were absent in the dendrites of *Scn2a*-haploinsufficient mice.

(A) The multiple linear regression model used to fit the ΔF/F of a ROI from the apical tuft of IT neurons.

(B) Fraction of ROIs in *PlexinD1-CreER+/-::Scn2a+/+* control animals with statistically significant ($P < 0.01$)

regression coefficients for the various predictor variables. Red shading indicates *P*-value calculated based on the

binomial test.

(C) Similar to (B) in *PlexinD1-CreER+/-::Scn2a+/-* mice.

DISCUSSION

The results in this study support three main conclusions. One, *Scn2a* haploinsufficiency impairs

the coupling between apical and proximal dendritic compartments in PT and IT pyramidal

neurons in the medial frontal cortex. Two, *Scn2a+/-* mice exhibited cognitive rigidity, where they

were unable to adapt to the changing strategy of a computer opponent in a matching pennies

game. Three, reward- and strategy-related signals in the dendrites of frontal cortical IT neurons

were abolished in *Scn2a*-haploinsufficient mice, hinting at a dendritic substrate for the inflexible

behavior during decision-making.

Numerous studies have investigated the coupling between dendritic and somatic compartments

in cortical pyramidal neurons *in vivo*. In anesthetized animals, layer 5 pyramidal neurons in the

barrel cortex exhibit varied coupling between the apical dendrites and proximal soma[50].

Specifically, large calcium signals in the apical dendrites are associated with burst firing in the

soma. These correlated events may be elicited by whisker stimulation or by backpropagating

action potentials. The fluorescence transients detected in our study are likely associated with

these burst firing events rather than single action potentials, as indicated by the relatively low

rate of calcium events and the sensitivity of the genetically encoded calcium indicator under our

imaging conditions[35]. Our observation of high level of dendritic coupling in control animals also

aligns with other prior works. A previous study suggests that layer 5 pyramidal neurons in the

anterior cingulate cortex may exhibit more reliable signal propagations between soma and

dendrite compared to those in the sensory cortex[51]. In the awake animal, increased background

excitatory activity would elevate dendritic excitability, which should strengthen the coupling

between somatic, proximal, and apical compartments[52,53]. Indeed, other recent studies in awake

mice reported a close correspondence of calcium transients between distal apical dendrites and

locations proximal to the cell body[21,26,54]. Our results are in agreement with these earlier studies

to show a high degree of dendritic coupling in the medial frontal cortex of awake animals.

Scn2a encodes the voltage-gated sodium channel $Na_v1.2$, which localizes to dendrites of

neocortical pyramidal cells after the first postnatal week, and is expected to support dendritic

excitability. Indeed, in brain slices, one could evoke dendritic calcium signal via backpropagation

by stimulating the cell body, and this signal was reduced with the heterozygous loss of $Na_v1.2$[14].

Our study adds to this finding by showing the impact of *Scn2a* haploinsufficiency on the

transmission of signals in dendrites *in vivo* in awake animals. We further found cell-type specific

differences. PT neurons had impaired coupling affecting transmission from the proximal and

apical trunk up to the apical tuft in *Scn2a*$^{+/-}$ animals. Accompanying this decoupling in the

backward direction is an aberrant elevation in the number of calcium events in the apical tuft. By

contrast, IT neurons had diminished dendritic signaling in both the forward and backward

directions, and its calcium event rates were not changed by *Scn2a* haploinsufficiency. The

aberrant calcium event rates may relate to the paradoxical increase in action potential

excitability seen in cortical pyramidal neurons following the deletion of *Scn2a*[55,56].

During behavior, reward- and strategy-related signals were diminished in frontal cortical IT

neurons of *Scn2a*$^{+/-}$ animals. Prior studies have shown that causally silencing neural activity in

the frontal cortical region can impair behavioral adjustments[37,57], therefore our observed loss of

task-related signals may underpin the cognitive rigidity. Interestingly, our results match a prior

study identifying reward-related signals in the dendrites of IT, but not PT, pyramidal neurons in

the somatosensory cortex in behaving mice[24]. Given that frontal cortical PT and IT neurons

underpin distinct aspects of motor and decision-related behaviors[32,58,59], the cell-type specific

deficits provide insights into the circuit mechanisms in ASD.

Despite the strong causal link from *SCN2A* to ASD, prior studies of the heterozygous *Scn2a*$^{+/-}$

mice reported only mild behavioral changes. For adult animals, there was no notable change in

many behavioral assays designed to test locomotion, anxiety, social interactions, and

memory[14,60]. There were some phenotypes for *Scn2a*$^{+/-}$ mice, such as decreased ultrasonic

vocalization[60], novelty-induced hyperactivity [61], enhanced social interaction [62], and elevated

long-term fear memory[61,62]. Phenotypes such as increased responses to lights-on stimuli,

daytime hypoactivity, and night-time hyperactivity have also been described in zebrafish models

with mutations in the related *Scn1lab*[63]. There have been reports of spatial learning deficits in

Scn2a-haploinsufficient mice[64,65]. A particularly promising trait for translational research is

impaired vestibulo-ocular reflex, which is cerebellum-dependent and can be observed in both

humans and rodents[66]. However, these behavioral phenotypes are difficult to relate to the

intellectual deficits observed in humans[11-13]. In this study, building on the classic matching

pennies game, we developed a task that incentivizes cognitive flexibility. Testing a large cohort

of *Scn2a*$^{+/-}$ mice and *Scn2a*$^{+/+}$ littermate controls, we demonstrated that control animals could

adjust bidirectionally to the shifting behavioral strategies, but mice with *Scn2a* haploinsufficiency

failed to adapt. We believe our findings have translational significance, because games such as

matching pennies tap into the social and strategic aspects of decision-making[41,67]. The same

competitive or cooperative game can be tested on animal models[68] and individuals with ASD[69,70].

In fact, one recent study reported a similar variant of matching pennies, where they

programmed computer opponents to play with either a cooperative or competitive strategy and

evaluated how human participants would adapt[71]. We expect cross-species tasks will facilitate

the translation of treatments from animal models to humans.

Understanding how ASD risk genes affect dendritic function is crucial because dendrites play a

central role in how neurons receive, integrate, and process information. Here our data provide

evidence that the heterozygous loss of the high-confidence risk gene *Scn2a* reduces dendritic

coupling in the medial frontal cortex and impairs flexible decision-making. By linking the risk

gene to dendritic alterations *in vivo* and during behavior, our results provide mechanistic insights

into the cortical circuit deficits underlying the cognitive symptoms observed in ASD, which

ultimately may advance diagnostic and treatment strategies for the disorder.

Acknowledgments

This work was supported by a Simons Foundation Autism Research Initiative Pilot Award

(A.C.K.). Additional support was provided by a China Scholarship Council-Yale World Scholars

Fellowship (H.W.), Cornell Engineering Learning Initiatives (J.I.), NIH training grants

T32GM007205 (N.K.S.), NIH fellowship F30MH129085 (N.K.S.), and NIH instrumentation grant

S10OD032251 (Cornell Biotechnology Resource Center Imaging Facility).

Contributions

H.W., L.S., and A.C.K planned the study. H.W. and L.S. performed experiments. J.I. and T.S.J.

assisted with the behavioral studies. N.K.S. assisted with imaging experiments. H.W. and L.S.

analyzed the data. J.Q. assisted with analysis of the behavioral data. K.J.B. helped with

experimental design and discussions. H.W. and A.C.K. drafted the manuscript. All authors

reviewed the manuscript before submission.

Competing interests

A.C.K. has been a scientific advisor or consultant for Boehringer Ingelheim, Empyrean

Neuroscience, Freedom Biosciences, and Xylo Bio. A.C.K. has received research support from

Intra-Cellular Therapies. K.J.B. is an advisor or consultant for Regel Therapeutics, Neurocrine

Biosciences, and Emugen Therapeutics. The other authors report no competing interest.

Data availability

Data and code associated with the study will be available on https://github.com/Kwan-Lab.

METHODS

Animals. *Scn2a*$^{+/-}$ mice were originally generated via homologous recombination[72] and sourced

from the University of California San Francisco. C57BL/6J (Stock #000664) and *PlexinD1-2A-*

CreER1 (B6;129S4-*Plxnd1*$^{tm1.1(cre/ERT2)Zjh}$/J, Stock #036296)[46] mice were purchased from

Jackson Laboratory. A *Scn2a*$^{+/-}$ mouse and a C57BL/6J mouse were used as breeding pairs to

generate *Scn2a*$^{+/+}$ and *Scn2a*$^{+/-}$ mice. A *Scn2a*$^{+/-}$ mouse and a *PlexinD1-2A-CreER*$^{+/+}$ mouse

were used as breeding pairs to generate *PlexinD1-2A-CreER^+/-::Scn2a^+/+* and *PlexinD1-2A-*

CreER^+/-::Scn2a^+/- mice. Ear punch was used for genotyping, in which the inserted gene *Neo^r*

was used to identify the heterozygous loss of *Scn2a.* Mice were housed in groups of 2 – 4

animals with 12-hr-dark and 12-hr-light cycle control (lights off at 7pm at Yale University; lights

off at 8pm at Cornell University). Animal procedures were approved by the Institutional Animal

Care and Use Committees at Yale University and Cornell University.

Surgical Procedures. Prior to surgery, the mouse was injected with dexamethasone (3 mg/kg,

i.m.; #024751, Henry Schein Animal Health). Animals were anesthetized with 2 - 3% isoflurane

in oxygen for induction before the surgery. The isoflurane was lowered to 1 - 1.5% during the

surgical procedures. The mouse was placed on a water-circulating heat pad (#RT-0515, Kent

Scientific) in a stereotaxic surgical frame (Model 900, David Kopf Instruments). The body

temperature was maintained at 38°C. Petrolatum ophthalmic ointment (#IS4398, Dechra) was

applied to cover the eyes of the mouse. The hair of the scalp was removed by scissors. The skin

was disinfected using an alcohol prep pad (#191089, McKesson), and pushed aside to expose

the skull. The scalp was wiped with ethanol and povidone-iodine. A small craniotomy was made

above each of the target regions using a handheld dental drill (#HP4-917, Foredom). A

borosilicate glass pipette was inserted into the brain and adeno-associated virus (AAV) was

delivered intracranially into the target region using an injector (Nanoject II/III Auto-Nanoliter

Injector, Drummond Scientific). The injection was performed with a series of 4.6 nL pulses with

30 s interval between pulses. To reduce backflow of the virus, we waited 5 minutes before

retracting the glass pipette. For each target region (i.e., medial frontal cortex, striatum, or pons),

four injections were made at sites corresponding to four vertices of a 0.2 mm-wide square

centered at the center coordinates of the target region. During the procedure, artificial

cerebrospinal fluid (ACSF, 135 mM NaCl, 5 mM HEPES, 5 mM KCl, 1.8 mM CaCl2, 1 mM

MgCl2; pH 7.3) was applied to keep the cortical surface moist. After injecting all targeted

regions, the craniotomies were covered with silicone elastomer (#0318, Smmoth-On, Inc.), and

the skin was sutured (#1265B, Surgical Specialties Corporation). Animal was given carprofen (5

mg/kg, s.c.; #002459, Henry Schein Animal Health) after the surgery and on each of the

following 3 days.

For imaging spontaneous dendritic calcium transients in PT neurons, 184 nL of AAVrg-hSyn-

Cre-WPRE-hGH (1:200 diluted in phosphate-buffered saline (PBS; #P4417, Sigma-Aldrich))

was injected into the pons (anteroposterior (AP): -3.4 mm, mediolateral (ML): -0.7 mm,

dorsoventral (DV): -5.2 mm; relative to bregma) and 147.2 nL of AAV1-CAG-Flex-GCaMP6f-

WPRE-SV40 (1:10 diluted in PBS) was injected into the ACAd and medial MOs region of the

medial frontal cortex (AP: 1.5 mm, ML: -0.4 mm; DV: -1.0 mm) of *Scn2a*$^{+/-}$ mice and *Scn2a*$^{+/+}$

littermate control mice. For imaging spontaneous dendritic calcium transients in IT neurons, 184

nL of AAVrg-hSyn-Cre-WPRE-hGH (1:200 diluted in PBS) was injected into the contralateral

striatum (AP: 0.6 mm, ML: 2.2 mm, DV: -2.8 mm) and 147.2nL of AAV1-CAG-Flex-GCaMP6f-

WPRE-SV40 (1:10 diluted in PBS) was injected into the medial frontal cortex (AP: 1.5 mm, ML: -

0.4mm; DV: -1.0mm) of *Scn2a*$^{+/-}$ mice and *Scn2a*$^{+/+}$ littermate control mice.

At 3 weeks after the viral injection, there was a second surgery to implant a glass window. The

pre- and post-operative steps were the same. An incision was made to remove skin and expose

the skull. The skull was cleaned by wiping with the ethanol and povidone-iodine. A 3 mm-

diameter circular craniotomy centering at the medial frontal cortex was made using the

handheld dental drill. ACSF was applied to keep the dura moist. A window was constructed

using two pieces of glass coverslips (3 mm diameter, 0.15 mm thickness; #640720, Warner

Instruments) that were bonded with ultraviolet light-curing optical adhesive (#NOA 61, Norland

Products) using an ultraviolet illuminator (#2182210, Loctite). This two-layer glass window was

placed into the craniotomy, and super glue adhesive (Henkel Loctite 454) was used to secure

the window to the surrounding skull. After 20 minutes when the glue was dry, a stainless steel

headplate (eMachineShop; design available at https://github.com/Kwan-Lab/behaivor-rigs) was

placed centered on the glass window using the super glue (All-purpose glue, Krazy) and a quick

adhesive cement system (Metabond, Parkell). Mouse would recover for two weeks after the

window implant prior to imaging experiments.

For imaging dendritic calcium transients during behavior, the procedures were performed in a

single surgery. The pre- and post-operative steps were the same. The hair of the scalp was

removed by scissors. An incision was made to remove skin and expose the skull. The skull was

cleaned by wiping with the ethanol and povidone-iodine. A 3 mm-diameter circular craniotomy

centering at the medial frontal cortex was made using the handheld dental drill. ACSF was

applied to keep the dura moist. To target IT neurons, 147.2 nL of AAV1-CAG-Flex-GCaMP6f-

WPRE-SV40 (1:10 diluted in PBS) was injected into the medial frontal cortex (AP: 1.5 mm, ML: -

0.4 mm; DV: -1.0 mm) of *PlexinD1-2A-CreER*$^{+/-}$*::Scn2a*$^{+/-}$ and *PlexinD1-2A-CreER*$^{+/-}$*::Scn2a*$^{+/+}$

mice. The other injection parameters were the same as described in the previous section. A

four-layer glass window was made by bonding 1 layer of 4 mm-diameter glass together with 3

layers of 3 mm-diameter glass using the ultraviolet light-curing optical adhesive. The thicker

glass window was needed to minimize the motion of the brain, which was more vigorous during

behavior than when imaging spontaneous activity. This glass window was placed in the

craniotomy and secured by the super glue adhesive. After 20 minutes, a stainless steel

headplate was placed centered on the glass window with the super glue and a quick adhesive

cement system.

Tamoxifen injection. Tamoxifen was used to induce Cre-dependent gene expression when

using *PlexinD1-2A-CreER$^{+/-}$::Scn2a$^{+/-}$* and *PlexinD1-2A-CreER$^{+/-}$::Scn2a$^{+/+}$* mice. Tamoxifen

(#T4648, Sigma-Aldrich) was dissolved in corn oil (#C8267, Sigma-Aldrich) at a concentration of

20 mg/mL. The solution was placed for 1 hour at 37 °C in an ultrasonic water bath to help

dissolve the compound. The solution was aliquoted into 2 mL Eppendorf tubes and stored at -

20°C. Prior to injection, the aliquoted tamoxifen was thawed at 4°C. Tamoxifen (75 mg/kg, i.p.)

was injected for 5 consecutive days starting from at 3 days after surgery. Animals started water

restriction and behavior training at least 7 days after the last tamoxifen injection.

Behavioral setup. The operant training setup for the head-fixed mouse was based on the

design from our prior studies[37,49]. Detailed instruction to construct the apparatus is available at

https://github.com/Kwan-Lab/behavioral-rigs. The mouse's head stabilized via the headplate

using a stainless-steel piece (eMachineShop). The mouse sat in an acrylic tube (8486K433;

McMaster-Carr), which limited gross movements but allowed postural adjustments. A lick port

with two water spouts was positioned in front of the mouse. The spouts were blunted 20-gauge

stainless-steel needles. The lick port was placed on a motorized linear actuator (0.8" stroke, 4.5

lb; B07ZJ4B3WW, DC House), which was powered by an L293D motor driver (497-2936-5-ND,

DigiKey). The actuator enabled us to move the lick port to a location accessible to the mouse at

the start of a trial and then retract the lick port at the end of the response window. When the

mouse licked either spout, the contact was recorded by a battery-powered lick detection

electronic circuit, which sent signals to the computer via a data acquisition unit (USB-201,

Measurement Computing) and logged by the Presentation software (Neurobehavioral System

Inc.). Water delivery from each lick spout was controlled by a solenoid fluid valve (MB202-V-A-3-

0-L-204; Gems Sensors & Controls). When there was a reward, ~4 µL of water was delivered by

sending a signal via the data acquisition unit to the water delivery electronic circuit to activate

the solenoid. A pair of speakers (S120, Logitech) were placed in front of the mouse to play the

sound cue. For training, the entire setup was placed inside a behavior box with walls lined with

soundproof acoustic foams (5692T49, McMaster-Carr).

Behavioral training – matching pennies. All the tasks were programming using the scripting

language in the Presentation software. Mice were water restricted for 48 hours before the

behavioral training. For the first training stage, mice were habituated to head restraint and the

lick spouts were always positioned in front of the mouse (i.e., the actuator did not move). They

could lick either lick spout for water until they are satiated. The first stage lasted 2–3 days, with

the aim of helping the mouse get used to the head restraint and the licking circuit. For the

second training stage, at the start of a trial, the actuator started to advance. We measured that

the actuator took 0.85 s to bring the lick spout to the front of the mouse. Therefore, 0.85 s after

the start of a trial, a 5 kHz sound cue lasting 0.2 s was played and a 2-s response window

began. During the response window, the first lick to either side of lick spouts will lead to the

immediate delivery of ~4 µL of water to the contacted spout and triggered a fixed 3-s period for

consumption. If no lick was detected, the fixed 3-s period would be applied at the end of the 2-s

response window. At the end of the fixed 3-s period, the actuator started to retract and an

intertrial interval was applied until the start of the next trial. The duration of the intertrial interval

was drawn randomly for each trial, from a truncated exponential distribution with $\lambda = 1/3$ and

boundaries of 1 and 5. Once the ITI concluded, the next trial would start. The session continued

until either the mouse misses for 10 consecutive trials, or if it made 1000 responses. The

second stage typically lasted 2 days, with the aim of helping the mouse get used to the moving

actuator. When a mouse could lick at least 100 trials per session, it proceeded to the matching

pennies game.

For matching pennies task, the timing of events in a trial was exactly the same as the second

stage. The first lick to either water spout during the 2-s response window would be the mouse's

choice (left or right) for that trial. According to the payoff matrix of matching pennies, if the

mouse's choice matched the computer's choice, ~4 µL of water will be delivered and triggered

the fixed 3-s period. If the mouse's choice did not match the computer's choice, there was no

reward but still triggered the fixed 3-s period. In this standard version of the matching pennies

game, the computer opponent made choices using the 'predictive' strategy. This strategy is the

same as the strategy used in ref. [39], algorithm 2 in ref. [45], and competitor 1 in ref. [44]. Briefly, the

computer had knowledge of the mouse's choice and outcome history since the start of the

session. The computer would calculate 9 conditional probabilities that the mouse would choose

left given sequences of the preceding N choices (N = 0 – 4) and N choice-outcome

combinations (N = 1 – 4). A binomial test was then used to test each of these conditional

probabilities against the null hypothesis that the mouse would choose left with a probability of

0.5. If none of the null hypotheses was rejected, the computer would randomly choose between

left or right. If one or more hypotheses were rejected, the computer would choose the side with

the statistically significant conditional probability that was furthest away from 0.5.

Behavioral training – matching pennies against a computer opponent with shifting

strategies. Mice were trained to play the standard version of the matching pennies game; when

a mouse achieved a reward rate of 35% or greater for 3 consecutive days, it was deemed

competent, and we would test the animal against a computer opponent with shifting strategies.

In this variant, the computer opponent, which we called Tom (after Tom, the strategist who

switches strategy to try to catch the mouse Jerry), can play with 1 of 3 different strategies:

predictive, WSLS (win-stay lose-switch), and alternate. Tom would use the same strategy for a

block of trials, drawn randomly from a uniform distribution ranging between 90 and 110, and

then switch abruptly to the different strategy, without any indication to the mouse. For the

predictive strategy, this is the same as described above, except Tom would calculate the

conditional probabilities based on the mouse's choice and outcome history since the start of the

current block. For the WSLS strategy, Tom would randomly choose left or right for the first trial. If

Tom wins, it will stay on that same option for the subsequent trial. But if Tom loses, it will switch

to the other option for the next trial. Tom will continue with this logic until the end of the block.

For the alternate strategy, Tom would randomly choose left or right for the first trial. On the next

trial, Tom will always choose the other option, and continue to alternate between the two options

until the end of the block. At the start of a session, Tom began with the predictive strategy for the

first block, then a second strategy (WSLS or alternate) for the second block, the predictive

strategy for the third block, the same second strategy for the fourth block, and so on. Hence,

Tom switches either between predictive and WSLS or predictive and alternate. The session

continued until either the mouse misses for 10 consecutive trials, or if it made 1000 responses.

Analysis of behavior data. We included all sessions when the mouse responded for at least

50 trials for each block. For the standard version of matching pennies, we had 139 sessions

from 16 $Scn2a^{+/+}$ animals (7 males, 9 females) and 181 sessions from 21 $Scn2a^{+/-}$ animals (12

males, 9 females). For the matching pennies against a computer opponent with shifting between

predictive and WSLS strategies, we had 130 sessions from 38 $Scn2a^{+/+}$ mice (16 males, 22

females) and 108 sessions from 25 $Scn2a^{+/-}$ mice (15 males, 10 females). For the matching

pennies against a computer opponent with shifting between predictive and alternate strategies,

we had 60 sessions from 15 $Scn2a^{+/+}$ mice (7 males, 8 females) and 38 sessions from 8

$Scn2a^{+/-}$ mice (4 males, 4 females).

The behavior data, including animal's choices and outcomes, lick times, and computer's

strategy, were stored by the Presentation software as text log files. The log files were processed

with custom-written code in MATLAB. To compute the probability of switch P(switch), we would

sum the number of trials in which the choice was different than the prior choice (i.e., animal

selected right and the last choice was left, or animal selected left and the last choice was right),

and divide by the total number of trials. To compute the probability of reward P(reward), we

would sum the number of trials in which the animal received a water reward (i.e. animal selected

right and the computer selected right, or animal selected left and the computer selected left),

and divide by the total number of trials. For each session we would calculate P(switch) and

P(reward) separately using trials in which computer strategy was predictive or WSLS, or

predictive or alternate.

Two-photon microscopy. For multi-plane imaging of the spontaneous dendritic calcium

transients, the experiments were performed with a Movable Objective Microscope (MOM, Sutter

Instrument) equipped with a resonant-galvo scanner (Rapid Multi Region Scanner, Vidrio

Technologies or Resonant Scan Box, Sutter Instrument). We used a water-immersion 20x

objective (XLUMPLFLN, 20X/0.95 N.A., Olympus), with an electrically tunable lens (EL-3-10;

Optotune) placed just above the back aperture of the objective. Scanimage 2020 software was

used to control the microscope for image acquisition. The excitation source was either a tunable

Ti:Sapphire femtosecond laser (Chameleon Ultra II, Coherent) or a fixed-wavelength ultrafast

laser (Axon 920-2 TPC, Coherent). The excitation wavelength was 920 nm, and the emission

was collected behind a 475 – 550 nm bandpass filter. The intensity of laser was 120 – 180 mW

after the objective. Time-lapse images were acquired at a resolution of 512 x 512 pixels. The

ETL was set such that 0 V corresponded to the plane of the proximal trunk, and two different

voltages were used to rapidly switch between the 3 different focal planes. The voltages related

to depth through a linear relationship and was calibrated. The voltages were sent using the fast

focus control module in ScanImage. The field of view was slightly larger at the plane of the

apical tuft, compared to the plane of the proximal trunk. The frame rate was 7.93 Hz using

bidirectional scanning. Initially the mouse was habituated in the two-photon microscope for 3

days. The habituation time increased day by day, from 10 minutes to 40 minutes. Prior to the

time-lapse imaging, z-stack images were acquired from the pial surface to -500 μm with a step

size of 5 μm. The z-stack images were used to visualize the dendritic tree of a pyramidal

neuron, such that we may select specific depths for the time-lapse imaging. For PT neurons, we

would choose one depth each within the range for apical tuft (-50 to -150 μm), apical trunk (-250

to -400 μm), and proximal dendrite (<-400 μm). For IT neurons, we would choose one depth

each within the range for apical tuft (-50 to -150 μm), apical trunk (-150 to -350 μm), and

proximal dendrite (<-350 μm). We would acquire time-lapse images from the three depths for 30

minutes.

For imaging during behavior, we used the same imaging system with the behavioral apparatus

set inside the two-photon microscope. There were several other differences. We used a water-

immersion 25x objective (N25X-APO-MP, NIKON). The frame rate was 30.03 Hz using

bidirectional scanning. The intensity of laser was 100 – 140 mW after the objective. We imaged

dendrites in a focal plane that lies between a depth of -50 and -150 μm. To synchronize

behavioral and imaging data, a TTL pulse was sent by the Presentation software at the

beginning of each trial from the USB-201 board of the behavior system. The imaging system

used the TTL pulse as an external trigger to initiate the acquisition of a new imaging file.

For each session, we would initially acquire baseline images for 30 seconds when the animal

has not yet started the task. Then we would acquire time-lapse images, triggered by the

behavioral setup as the task progressed, until the task was completed.

Preprocessing of the multi-plane imaging data. For multi-plane imaging of the spontaneous

dendritic calcium transients, we made three 10-minute recordings consecutively, which together

form the 30-minute session. Each recording generated a multi-page .tiff file saved by

ScanImage. The file contains image frames for the three different focal planes interleaved. We

would extract the image frames corresponding to each focal plane, and then concatenate to

produce the time-lapse images for a focal plane spanning 30 minutes.

We processed with NoRMCorre in MATLAB to correct for non-rigid translational motion. Next,

regions of interest (ROIs) were selected from the motion-corrected time-lapse images, by

comparing with the initial z-stack images of the individual neuron. The software to draw ROIs

was a GUI previously written in the lab. The criteria for ROI selection were as follows: first, the

dendrite was not overlapping with any other dendrites; second, the dendrite belonged to the

target neuron; third, there were event-like transients with rapid onset with amplitude of more

than ~0.1 and exponential decay during the session. To account for the influence of the neuropil,

we calculated the fluorescence signal for each ROI as follows:

$$F(t) = F_{Raw}(t) - cF_{neuropil}(t)$$

716   where $F_{raw}(t)$ is the summed pixel value within the ROI at time t, c is a correction factor set to

717   0.4, and $F_{neuropil}(t)$ is the summed pixel value in the neuropil area, defined as an annulus shaped

718   area centered at the centroid location of the ROI with inner and outer diameters of $2r$ and $3r$,

719   where r is the radius if we assume the ROI area is a circle. The fractional change in

720   fluorescence $\Delta F/F(t)$ was calculated as follows:

$$\frac{\Delta F}{F}(t) = \frac{F(t) - F_0(t)}{F_0(t)}$$

where $F_0(t)$ is the baseline fluorescence, which was the 10^{th} percentile of $F(t)$ within a 2-min

running window centered around time t.

Analysis of the multi-plane imaging data. For PT neurons, we imaged 23 cells in 6 $Scn2a^{+/+}$

mice (5 males, 1 female) and 31 cells in 6 $Scn2a^{+/-}$ mice (3 males, 4 females For IT neurons, we

imaged 12 cells in 7 $Scn2a^{+/+}$ mice (4 males, 3 females) and 13 cells in 5 $Scn2a^{+/-}$ mice (1 male,

4 females). To infer the calcium events from $\Delta F/F(t)$, we used an algorithm called Online Active

Set method to Infer Spikes, or OASIS [36]. We estimated the coupling between different dendritic

compartments using conditional probability. To calculate conditional probability, for example the

probability of event in compartment B given an event in compartment A, we would first identify

all events in compartment A. For each event, we examined a window of ±3 imaging frames (~

±0.38 s) around its occurrence to check for at least one event in compartment B. If a

corresponding event was present, we recorded a 1; otherwise, we recorded a 0. The conditional

probability was then calculated by averaging the 1s and 0s across all events in compartment A.

To calculate the event rate, we divided the total number of calcium events by the duration of the

imaging session.

Preprocessing of the behavioral imaging data. For imaging during behavior, acquisition of an

image file was triggered by a TTL pulse sent by the Presentation software at the start of each

trial. Therefore, each trial generated a multi-page .tiff file saved by ScanImage. To analyze the

data, we first concatenated the image files in batches of 100 trials to produce several large

image files. Then we generated a template for motion correction by calculating the mean

projection using the initial baseline images acquired before matching pennies. Based on this

template image, we processed all the large image files with NoRMCorre in MATLAB to correct

for non-rigid translational motion. Next, regions of interest (ROIs) were selected from the

motion-corrected first large image file. The criteria for ROI selection were as follows: first, the

dendrite was not overlapping with any other dendrites; second, there were event-like transients

with rapid onset with amplitude of more than ~0.1 and exponential decay. Then, the masks for

the ROIs were applied to all other large image files to extract the fluorescence signals. We

calculated the fractional change in fluorescence *ΔF/F(t)* using the same steps, except did not

correct for the neuropil influence (i.e., c = 0).

Analysis of the behavioral imaging data. For imaging during behavior, we had 18 sessions

from 4 *PlexinD1-2A-CreER$^{+/-}$::Scn2a$^{+/+}$* animals (2 males, 2 females), and 15 sessions from 4

PlexinD1-2A-CreER$^{+/-}$::Scn2a$^{+/-}$ animals (4 males). Behavior data were processed with a custom

written script in MATLAB. To plot trial-averaged activity, we aligned *ΔF/F(t)* by the time of cue

onset, and generated peri-stimulus time histogram by averaging across all trials of the same

type (i.e., left & no reward, left & reward, right & no reward, or right & reward). To assess task-

related activity, for each ROI, we would compare the trial-by-trial *ΔF/F* between the cue period (t

= -0.5 – 0.5 s) and baseline (t = -2 – -1 s) or between the outcome period (t = 1 – 2 s) and

baseline (t = -2 – -1 s). To quantify how dendritic calcium transients may relate to choice,

outcome and strategy, we constructed a multiple linear regression model:

$$
\frac{\Delta F}{F}(t) = b_0 + b_1 C_n + b_2 R_n + b_3 C_n R_n + b_4 C_{n-1} + b_5 R_{n-1} + b_6 C_{n-1} R_{n-1} + b_7 S + b_8 S C_n + b_9 S R_n \\
+ \varepsilon(t)
$$

where *ΔF/F(t)* is the fractional change in fluorescence at time *t* relative to cue onset in trial *n*, *C$_n$*

and *C$_{n-1}$* are the choices made by the mouse on the current trial and the previous trial, *R$_n$* and

R$_{n-1}$ are the outcomes received by the mouse on the current trial and the previous trial, *S* is the

computer strategy, *b$_0$, b$_1$,…, b$_9$* are the regression coefficients and $\varepsilon(t)$ is the error variable.

Choices were coded as -1 for left choice and 1 for right choice. Outcomes were coded as 0 for

no reward and 1 for reward. Computer strategy was coded as 1 for predictive and -1 for WSLS.

The regression coefficients were calculated by fitting the model using the MATLAB function *fitlm*

for each imaging session. We fitted for the time period from -2 s before the cue onset to 6 s after

the cue onset, with step size of 25 ms. To collate the results across all sessions, for each

predicting variable, we calculated the fraction of dendritic ROIs in which the regression

coefficient was significantly different from zero ($P < 0.01$).

Histology. After the completion of imaging and behavioral studies, mice were perfused with ice-

cold PBS and paraformaldehyde solution (PFA, 4% v/v in PBS). The brains were extracted and

fixed in 4% PFA at 4°C. The brain was cut into coronal sections at a thickness of 50 μm using a

vibratome (#VT1000S, Leica). The brain sections were mounted on slides with glass coverslips.

If the brains were to be stored beyond 24 hours before sectioning, PFA was replaced with PBS.

For fluorescence imaging of the sections, a confocal microscope (LSM880, Zeiss) equipped with

a 10x/0.45 N.A. water objective was used. Images were acquired at 512 x 512 pixels.

Statistics. All statistics were conducted within MATLAB and R studio. For comparisons across

different groups, a mixed effects model was used with the lme4 package in R. For conditional

probability, we constructed a model with fixed effects including genotype, compartment pairing,

cell type, and all higher order interactions, along with a random effects term (intercept) for

nested ROI per cell per mouse. For event rate, we constructed a model with fixed effects

including genotype, compartment, cell type, and all higher order interactions, along with a

random effects term (intercept) for nested ROI per cell per mouse. The specifics for the other

statistical tests performed are indicated in the main text and the figure captions.

REFERENCES

1. De Rubeis, S., He, X., Goldberg, A.P., Poultney, C.S., Samocha, K., Cicek, A.E., Kou, Y., Liu, L., Fromer, M., Walker, S., et al. (2014). Synaptic, transcriptional and chromatin genes disrupted in autism. Nature *515*, 209-215. 10.1038/nature13772.
2. Sanders, S.J., Murtha, M.T., Gupta, A.R., Murdoch, J.D., Raubeson, M.J., Willsey, A.J., Ercan-Sencicek, A.G., DiLullo, N.M., Parikshak, N.N., Stein, J.L., et al. (2012). De novo mutations revealed by whole-exome sequencing are strongly associated with autism. Nature *485*, 237-241. 10.1038/nature10945.
3. Satterstrom, F.K., Kosmicki, J.A., Wang, J., Breen, M.S., De Rubeis, S., An, J.Y., Peng, M., Collins, R., Grove, J., Klei, L., et al. (2020). Large-Scale Exome Sequencing Study Implicates Both Developmental and Functional Changes in the Neurobiology of Autism. Cell *180*, 568-584 e523. 10.1016/j.cell.2019.12.036.
4. Willsey, A.J., Sanders, S.J., Li, M., Dong, S., Tebbenkamp, A.T., Muhle, R.A., Reilly, S.K., Lin, L., Fertuzinhos, S., Miller, J.A., et al. (2013). Coexpression networks implicate human midfetal deep cortical projection neurons in the pathogenesis of autism. Cell *155*, 997-1007. 10.1016/j.cell.2013.10.020.
5. Nelson, S.B., and Valakh, V. (2015). Excitatory/Inhibitory Balance and Circuit Homeostasis in Autism Spectrum Disorders. Neuron *87*, 684-698. 10.1016/j.neuron.2015.07.033.
6. Nelson, A.D., and Bender, K.J. (2021). Dendritic Integration Dysfunction in Neurodevelopmental Disorders. Dev Neurosci *43*, 201-221. 10.1159/000516657.
7. Lazaro, M.T., Taxidis, J., Shuman, T., Bachmutsky, I., Ikrar, T., Santos, R., Marcello, G.M., Mylavarapu, A., Chandra, S., Foreman, A., et al. (2019). Reduced Prefrontal Synaptic Connectivity and Disturbed Oscillatory Population Dynamics in the CNTNAP2 Model of Autism. Cell Rep *27*, 2567-2578 e2566. 10.1016/j.celrep.2019.05.006.
8. Wang, X., McCoy, P.A., Rodriguiz, R.M., Pan, Y., Je, H.S., Roberts, A.C., Kim, C.J., Berrios, J., Colvin, J.S., Bousquet-Moore, D., et al. (2011). Synaptic dysfunction and abnormal behaviors in mice lacking major isoforms of Shank3. Hum Mol Genet *20*, 3093-3108. 10.1093/hmg/ddr212.
9. Lai, K.O., and Ip, N.Y. (2013). Structural plasticity of dendritic spines: the underlying mechanisms and its dysregulation in brain disorders. Biochim Biophys Acta *1832*, 2257-2263. 10.1016/j.bbadis.2013.08.012.
10. Phillips, M., and Pozzo-Miller, L. (2015). Dendritic spine dysgenesis in autism related disorders. Neuroscience Letters *601*, 30-40. 10.1016/j.neulet.2015.01.011.
11. Sanders, S.J., Campbell, A.J., Cottrell, J.R., Moller, R.S., Wagner, F.F., Auldridge, A.L., Bernier, R.A., Catterall, W.A., Chung, W.K., Empfield, J.R., et al. (2018). Progress in Understanding and Treating SCN2A-Mediated Disorders. Trends Neurosci *41*, 442-456. 10.1016/j.tins.2018.03.011.
12. Begemann, A., Acuna, M.A., Zweier, M., Vincent, M., Steindl, K., Bachmann-Gagescu, R., Hackenberg, A., Abela, L., Plecko, B., Kroell-Seger, J., et al. (2019). Further corroboration of distinct functional features in SCN2A variants causing intellectual disability or epileptic phenotypes. Mol Med *25*, 6. 10.1186/s10020-019-0073-6.

13. Ben-Shalom, R., Keeshen, C.M., Berrios, K.N., An, J.Y., Sanders, S.J., and Bender, K.J.
(2017). Opposing Effects on NaV1.2 Function Underlie Differences Between SCN2A
Variants Observed in Individuals With Autism Spectrum Disorder or Infantile Seizures.
Biol Psychiatry 82, 224-232. 10.1016/j.biopsych.2017.01.009.

14. Spratt, P.W.E., Ben-Shalom, R., Keeshen, C.M., Burke, K.J., Jr., Clarkson, R.L., Sanders, S.J.,
and Bender, K.J. (2019). The Autism-Associated Gene Scn2a Contributes to Dendritic
Excitability and Synaptic Function in the Prefrontal Cortex. Neuron 103, 673-685 e675.
10.1016/j.neuron.2019.05.037.

15. Nelson, A.D., Catalfio, A.M., Gupta, J.P., Min, L., Caballero-Floran, R.N., Dean, K.P., Elvira,
C.C., Derderian, K.D., Kyoung, H., Sahagun, A., et al. (2024). Physical and functional
convergence of the autism risk genes Scn2a and Ank2 in neocortical pyramidal cell
dendrites. Neuron 112, 1133-1149 e1136. 10.1016/j.neuron.2024.01.003.

16. Chen, T.W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R.,
Kerr, R.A., Orger, M.B., Jayaraman, V., et al. (2013). Ultrasensitive fluorescent proteins for
imaging neuronal activity. Nature 499, 295-300. 10.1038/nature12354.

17. Jia, H., Rochefort, N.L., Chen, X., and Konnerth, A. (2010). Dendritic organization of
sensory input to cortical neurons in vivo. Nature 464, 1307-1312. 10.1038/nature08947.

18. Wilson, D.E., Whitney, D.E., Scholl, B., and Fitzpatrick, D. (2016). Orientation selectivity
and the functional clustering of synaptic inputs in primary visual cortex. Nat Neurosci 19,
1003-1009. 10.1038/nn.4323.

19. Otor, Y., Achvat, S., Cermak, N., Benisty, H., Abboud, M., Barak, O., Schiller, Y., Poleg-
Polsky, A., and Schiller, J. (2022). Dynamic compartmental computations in tuft dendrites
of layer 5 neurons during motor behavior. Science 376, 267-275.
10.1126/science.abn1421.

20. O'Hare, J.K., Gonzalez, K.C., Herrlinger, S.A., Hirabayashi, Y., Hewitt, V.L., Blockus, H.,
Szoboszlay, M., Rolotti, S.V., Geiller, T.C., Negrean, A., et al. (2022). Compartment-specific
tuning of dendritic feature selectivity by intracellular Ca(2+) release. Science 375,
eabm1670. 10.1126/science.abm1670.

21. Kerlin, A., Boaz, M., Flickinger, D., MacLennan, B.J., Dean, M.B., Davis, C., Spruston, N.,
and Svoboda, K. (2019). Functional clustering of dendritic activity during decision-
making. eLife 8. 10.7554/eLife.46966.

22. Lacefield, C.O., Pnevmatikakis, E.A., Paninski, L., and Bruno, R.M. (2019). Reinforcement
Learning Recruits Somata and Apical Dendrites across Layers of Primary Sensory Cortex.
Cell Rep 26, 2000-2008 e2002. 10.1016/j.celrep.2019.01.093.

23. Xu, N.L., Harnett, M.T., Williams, S.R., Huber, D., O'Connor, D.H., Svoboda, K., and Magee,
J.C. (2012). Nonlinear dendritic integration of sensory and motor input during an active
sensing task. Nature 492, 247-251. 10.1038/nature11601.

24. Takahashi, N., Ebner, C., Sigl-Glockner, J., Moberg, S., Nierwetberg, S., and Larkum, M.E.
(2020). Active dendritic currents gate descending cortical outputs in perception. Nat
Neurosci 23, 1277-1285. 10.1038/s41593-020-0677-8.

25. Takahashi, N., Oertner, T.G., Hegemann, P., and Larkum, M.E. (2016). Active cortical
dendrites modulate perception. Science 354, 1587-1590. 10.1126/science.aah6066.

26.   Beaulieu-Laroche, L., Toloza, E.H.S., Brown, N.J., and Harnett, M.T. (2019). Widespread
and Highly Correlated Somato-dendritic Activity in Cortical Layer 5 Neurons. Neuron *103*,
235-241 e234. 10.1016/j.neuron.2019.05.014.

27.   Sheffield, M.E., and Dombeck, D.A. (2015). Calcium transient prevalence across the
dendritic arbour predicts place field properties. Nature *517*, 200-204.
10.1038/nature13871.

28.   Baker, A., Kalmbach, B., Morishima, M., Kim, J., Juavinett, A., Li, N., and Dembrow, N.
(2018). Specialized Subpopulations of Deep-Layer Pyramidal Neurons in the Neocortex:
Bridging Cellular Properties to Functional Consequences. J Neurosci *38*, 5441-5455.
10.1523/JNEUROSCI.0150-18.2018.

29.   Anastasiades, P.G., and Carter, A.G. (2021). Circuit organization of the rodent medial
prefrontal cortex. Trends Neurosci *44*, 550-563. 10.1016/j.tins.2021.03.006.

30.   Gao, L., Liu, S., Gou, L., Hu, Y., Liu, Y., Deng, L., Ma, D., Wang, H., Yang, Q., Chen, Z., et al.
(2022). Single-neuron projectome of mouse prefrontal cortex. Nat Neurosci *25*, 515-529.
10.1038/s41593-022-01041-5.

31.   Shepherd, G.M. (2013). Corticostriatal connectivity and its role in disease. Nat Rev
Neurosci *14*, 278-291. 10.1038/nrn3469.

32.   Shao, L.X., Liao, C., Davoudian, P.A., Savalia, N.K., Jiang, Q., Wojtasiewicz, C., Tan, D., Liu,
R.J., Woodburn, S.C., Bilash, O.M., et al. (2025). Psilocybin's lasting action requires
pyramidal cell types and 5-HT2A receptors. Nature, in press.

33.   Marcassa, G., Dascenco, D., Lorente-Echeverria, B., Daaboul, D., Vandensteen, J., Leysen,
E., Baltussen, L., Howden, A.J.M., and de Wit, J. (2025). Synaptic signatures and disease
vulnerabilities of layer 5 pyramidal neurons. Nat Commun *16*, 228. 10.1038/s41467-024-
55470-w.

34.   Grewe, B.F., Voigt, F.F., van 't Hoff, M., and Helmchen, F. (2011). Fast two-layer two-
photon imaging of neuronal cell populations using an electrically tunable lens. Biomed
Opt Express *2*, 2035-2046. 10.1364/BOE.2.002035.

35.   Ali, F., and Kwan, A.C. (2020). Interpreting in vivo calcium signals from neuronal cell
bodies, axons, and dendrites: a review. Neurophotonics *7*, 011402.
10.1117/1.NPh.7.1.011402.

36.   Friedrich, J., Zhou, P., and Paninski, L. (2017). Fast online deconvolution of calcium
imaging data. PLoS Comput Biol *13*, e1005423. 10.1371/journal.pcbi.1005423.

37.   Siniscalchi, M.J., Phoumthipphavong, V., Ali, F., Lozano, M., and Kwan, A.C. (2016). Fast
and slow transitions in frontal ensemble activity during flexible sensorimotor behavior.
Nat Neurosci *19*, 1234-1242. 10.1038/nn.4342.

38.   Schamiloglu, S., Wu, H., Zhou, M., Kwan, A.C., and Bender, K.J. (2023). Dynamic Foraging
Behavior Performance Is Not Affected by Scn2a Haploinsufficiency. eneuro *10*.
10.1523/eneuro.0367-23.2023.

39.   Wang, H., Ortega, H.K., Atilgan, H., Murphy, C.E., and Kwan, A.C. (2022). Pupil Correlates
of Decision Variables in Mice Playing a Competitive Mixed-Strategy Game. eNeuro *9*.
10.1523/ENEURO.0457-21.2022.

40.   Wang, H., Ortega, H.K., Kelly, E.B., Indajang, J., Savalia, N.K., Glaeser-Khan, S., Feng, J., Li,
Y., Kaye, A.P., and Kwan, A.C. (2025). Frontal noradrenergic and cholinergic transients

exhibit distinct spatiotemporal dynamics during competitive decision-making. Sci Adv
11, eadr9916. 10.1126/sciadv.adr9916.

41. Camerer, C. (2003). Behavioral Game Theory: Experiments in Strategic Interaction
(Princeton University Press).

42. Murphy, C.E., Wang, H., Ortega, H.K., Kwan, A.C., and Atilgan, H. (2024). Change point
estimation by the mouse medial frontal cortex during probabilistic reward learning.
eLife. 10.7554/eLife.103001.1.sa4.

43. Ito, M., and Doya, K. (2015). Parallel Representation of Value-Based and Finite State-
Based Strategies in the Ventral and Dorsal Striatum. PLoS Comput Biol *11*, e1004540.
10.1371/journal.pcbi.1004540.

44. Tervo, D.G., Proskurin, M., Manakov, M., Kabra, M., Vollmer, A., Branson, K., and
Karpova, A.Y. (2014). Behavioral variability through stochastic choice and its gating by
anterior cingulate cortex. Cell *159*, 21-32. 10.1016/j.cell.2014.08.037.

45. Lee, D., Conroy, M.L., McGreevy, B.P., and Barraclough, D.J. (2004). Reinforcement
learning and decision making in monkeys during a competitive game. Brain Res Cogn
Brain Res *22*, 45-58. 10.1016/j.cogbrainres.2004.07.007.

46. Matho, K.S., Huilgol, D., Galbavy, W., He, M., Kim, G., An, X., Lu, J., Wu, P., Di Bella, D.J.,
Shetty, A.S., et al. (2021). Genetic dissection of the glutamatergic neuron system in
cerebral cortex. Nature *598*, 182-187. 10.1038/s41586-021-03955-9.

47. Seo, H., and Lee, D. (2007). Temporal filtering of reward signals in the dorsal anterior
cingulate cortex during a mixed-strategy game. J Neurosci *27*, 8366-8377.
10.1523/JNEUROSCI.2369-07.2007.

48. Sul, J.H., Kim, H., Huh, N., Lee, D., and Jung, M.W. (2010). Distinct roles of rodent
orbitofrontal and medial prefrontal cortex in decision making. Neuron *66*, 449-460.
10.1016/j.neuron.2010.03.033.

49. Siniscalchi, M.J., Wang, H., and Kwan, A.C. (2019). Enhanced Population Coding for
Rewarded Choices in the Medial Frontal Cortex of the Mouse. Cereb Cortex *29*, 4090-
4106. 10.1093/cercor/bhy292.

50. Helmchen, F., Svoboda, K., Denk, W., and Tank, D.W. (1999). In vivo dendritic calcium
dynamics in deep-layer cortical pyramidal neurons. Nat Neurosci *2*, 989-996.
10.1038/14788.

51. Marti Mengual, U., Wybo, W.A.M., Spierenburg, L.J.E., Santello, M., Senn, W., and
Nevian, T. (2020). Efficient Low-Pass Dendro-Somatic Coupling in the Apical Dendrite of
Layer 5 Pyramidal Neurons in the Anterior Cingulate Cortex. J Neurosci *40*, 8799-8815.
10.1523/JNEUROSCI.3028-19.2020.

52. Waters, J., and Helmchen, F. (2004). Boosting of action potential backpropagation by
neocortical network activity in vivo. J Neurosci *24*, 11127-11136.
10.1523/JNEUROSCI.2933-04.2004.

53. Murayama, M., and Larkum, M.E. (2009). Enhanced dendritic activity in awake rats. Proc
Natl Acad Sci U S A *106*, 20482-20486. 10.1073/pnas.0910379106.

54. Francioni, V., Padamsey, Z., and Rochefort, N.L. (2019). High and asymmetric somato-
dendritic coupling of V1 layer 5 neurons independent of visual stimulation and
locomotion. Elife *8*. 10.7554/eLife.49145.

55.   Spratt, P.W.E., Alexander, R.P.D., Ben-Shalom, R., Sahagun, A., Kyoung, H., Keeshen, C.M.,
968   Sanders, S.J., and Bender, K.J. (2021). Paradoxical hyperexcitability from Na(V)1.2 sodium
969   channel loss in neocortical pyramidal cells. Cell Rep *36*, 109483.
970   10.1016/j.celrep.2021.109483.

971   56.   Zhang, J., Chen, X., Eaton, M., Wu, J., Ma, Z., Lai, S., Park, A., Ahmad, T.S., Que, Z., Lee,
972   J.H., et al. (2021). Severe deficiency of the voltage-gated sodium channel Na(V)1.2
973   elevates neuronal excitability in adult mice. Cell Rep *36*, 109495.
974   10.1016/j.celrep.2021.109495.

975   57.   Schreiner, D.C., Cazares, C., Renteria, R., and Gremel, C.M. (2022). Information normally
976   considered task-irrelevant drives decision-making and affects premotor circuit
977   recruitment. Nat Commun *13*, 2134. 10.1038/s41467-022-29807-2.

978   58.   Musall, S., Sun, X.R., Mohan, H., An, X., Gluf, S., Li, S.J., Drewes, R., Cravo, E., Lenzi, I., Yin,
979   C., et al. (2023). Pyramidal cell types drive functionally distinct cortical activity patterns
980   during decision-making. Nat Neurosci *26*, 495-505. 10.1038/s41593-022-01245-9.

981   59.   Li, N., Chen, T.W., Guo, Z.V., Gerfen, C.R., and Svoboda, K. (2015). A motor cortex circuit
982   for motor planning and movement. Nature *519*, 51-56. 10.1038/nature14178.

983   60.   Lena, I., and Mantegazza, M. (2019). Na(V)1.2 haploinsufficiency in Scn2a knock-out
984   mice causes an autistic-like phenotype attenuated with age. Sci Rep *9*, 12886.
985   10.1038/s41598-019-49392-7.

986   61.   Tatsukawa, T., Raveau, M., Ogiwara, I., Hattori, S., Miyamoto, H., Mazaki, E., Itohara, S.,
987   Miyakawa, T., Montal, M., and Yamakawa, K. (2019). Scn2a haploinsufficient mice display
988   a spectrum of phenotypes affecting anxiety, sociability, memory flexibility and ampakine
989   CX516 rescues their hyperactivity. Mol Autism *10*, 15. 10.1186/s13229-019-0265-5.

990   62.   Shin, W., Kweon, H., Kang, R., Kim, D., Kim, K., Kang, M., Kim, S.Y., Hwang, S.N., Kim, J.Y.,
991   Yang, E., et al. (2019). Scn2a Haploinsufficiency in Mice Suppresses Hippocampal
992   Neuronal Excitability, Excitatory Synaptic Drive, and Long-Term Potentiation, and Spatial
993   Learning and Memory. Front Mol Neurosci *12*, 145. 10.3389/fnmol.2019.00145.

994   63.   Weinschutz Mendes, H., Neelakantan, U., Liu, Y., Fitzpatrick, S.E., Chen, T., Wu, W., Pruitt,
995   A., Jin, D.S., Jamadagni, P., Carlson, M., et al. (2023). High-throughput functional analysis
996   of autism genes in zebrafish identifies convergence in dopaminergic and neuroimmune
997   pathways. Cell Rep *42*, 112243. 10.1016/j.celrep.2023.112243.

998   64.   Kastner, D.B., Williams, G., Holobetz, C., Romano, J.P., and Dayan, P. (2024). The choice-
999   wide behavioral association study: data-driven identification of interpretable behavioral
components. bioRxiv. 10.1101/2024.02.26.582115.

65.   Middleton, S.J., Kneller, E.M., Chen, S., Ogiwara, I., Montal, M., Yamakawa, K., and
McHugh, T.J. (2018). Altered hippocampal replay is associated with memory impairment
in mice heterozygous for the Scn2a gene. Nat Neurosci *21*, 996-1003. 10.1038/s41593-
018-0163-8.

66.   Wang, C., Derderian, K.D., Hamada, E., Zhou, X., Nelson, A.D., Kyoung, H., Ahituv, N.,
Bouvier, G., and Bender, K.J. (2024). Impaired cerebellar plasticity hypersensitizes
sensory reflexes in SCN2A-associated ASD. Neuron *112*, 1444-1455 e1445.
10.1016/j.neuron.2024.01.029.

67.   Lee, D. (2008). Game theory and neural basis of social decision making. Nat Neurosci *11*,
404-409. 10.1038/nn2065.

68. Wang, H., and Kwan, A.C. (2023). Competitive and cooperative games for probing the
neural basis of social decision-making in animals. Neurosci Biobehav Rev *149*, 105158.
10.1016/j.neubiorev.2023.105158.
69. Izuma, K., Matsumoto, K., Camerer, C.F., and Adolphs, R. (2011). Insensitivity to social
reputation in autism. Proc Natl Acad Sci U S A *108*, 17302-17307.
10.1073/pnas.1107038108.
70. Yoshida, W., Dziobek, I., Kliemann, D., Heekeren, H.R., Friston, K.J., and Dolan, R.J.
(2010). Cooperation and heterogeneity of the autistic mind. J Neurosci *30*, 8815-8818.
10.1523/JNEUROSCI.0400-10.2010.
71. Philippe, R., Janet, R., Khalvati, K., Rao, R.P.N., Lee, D., and Dreher, J.C. (2024).
Neurocomputational mechanisms involved in adaptation to fluctuating intentions of
others. Nat Commun *15*, 3189. 10.1038/s41467-024-47491-2.
72. Planells-Cases, R., Caprini, M., Zhang, J., Rockenstein, E.M., Rivera, R.R., Murre, C.,
Masliah, E., and Montal, M. (2000). Neuronal death and perinatal lethality in voltage-
gated sodium channel alpha(II)-deficient mice. Biophys J *78*, 2878-2891. 10.1016/S0006-
3495(00)76829-9.
1027