1 "**Recurrent Extinction of Resistance Mutations Leads to Convergent Multidrug Resistance in**
2 **Sequential Antibiotic Treatment**"
3
4

5 Adam Lyon[1], Muhammed Sadik Yildiz[1], Samuel Cooke[2], Scott H. Saunders[3], Adam Rosenthal[2], Erdal
6 Toprak[1,3,4]
7
[1] Department of Pharmacology, The University of Texas Southwestern Medical Center, Dallas, TX
75390, USA
[2] Department of Microbiology and Immunology, University of North Carolina, Chapel Hill, NC, USA
[3] Lyda Hill Department of Bioinformatics, The University of Texas Southwestern Medical Center,
Dallas, TX 75390, USA
[4] Lead contact (Erdal.Toprak@utsouthwestern.edu)

Summary
16

To overcome the antibiotic resistance problem, exploiting reproducible evolutionary tradeoffs is
considered key for designing evolution-proof antibiotic therapies. Yet the predictability of resistance
evolution has been viewed as limited, given the often idiosyncratic genetic trajectories observed in
laboratory evolution experiments. To address this, we partially mimicked clinical antibiotic
pharmacodynamics by imposing strong selection and evolved *Escherichia coli* under single or sequential
antibiotic treatment. Under single-antibiotic selection, endpoint resistance, persistence, and tolerance
phenotypes were reproducible, but populations evolving in parallel frequently followed divergent
genetic trajectories. Remarkably, sequential antibiotic use redirected these divergent paths toward
genotypic and phenotypic convergence, driven by extinction of resistance-conferring mutations when
switching to the next effective antibiotic. Single-cell RNA sequencing demonstrated that evolved
cultures contain cells occupying distinct metabolic niches and include more cells in states with lower
translational activity and higher expression of toxin-antitoxin genes. These findings provide
evolutionary insights to inform clinically effective antibiotic treatment strategies that employ sequential
treatment.

48 **<u>INTRODUCTION</u>**

49 Antibiotic resistance is a global health crisis causing over one million deaths annually[1,2]. With limited
50 new antibiotics in development, evidence-based antibiotic choices are essential to prolong the efficacy

of current and future antibiotics. Patients requiring prophylaxis or those with serious infections undergo multiple rounds of antibiotic treatment. Unfortunately, these complex treatment regimens are often established haphazardly with little guidance for clinicians. Repeated treatment failures can lead to multi-drug resistant (MDR) infections, which are associated with higher morbidity and mortality[3]. Therefore, antibiotic choices must consider both clinical efficacy and the evolutionary consequences of treatment failures to mitigate these risks.

Laboratory evolution experiments have revealed many insights about canonical resistance mechanisms such as reduced drug uptake, increased efflux, and drug inactivation. However, we now know there are many other non-canonical antibiotic survival strategies, and these conventional experimental evolution protocols may not capture these non-canonical mechanisms because they do not account for the pharmacological complexity of drug treatment in patients[4-6]. In clinical settings, antibiotic concentrations routinely peak and trough, consistently remaining above the minimum inhibitory concentration (MIC), a clinical metric used to determine susceptibility of a pathogen to a given antibiotic (Figure 1a)[7].

Antibiotic persistence and tolerance are two non-canonical bacterial survival mechanisms about which we have limited knowledge[8-10]. In antibiotic tolerance (Supplementary Figure 1), the entire bacterial population survives longer under antibiotic exposure due to a slower killing rate, despite having the same MIC as antibiotic-sensitive bacteria[11]. Persistence, by contrast, arises from a subpopulation of bacteria that survives antibiotic exposure by dying more slowly, leading to a biphasic killing curve (Supplementary Figure 1). These slower killing rates pose significant challenges for treatments aiming to completely eradicate infections. However, our understanding of tolerance and persistence remains limited, as they are difficult to study[12]. Moreover, their true prevalence in clinical settings is unknown because they are often miscategorized as antibiotic susceptible due to indistinguishable MIC values.

To study the evolution of resistance, persistence, and tolerance under the selection of three clinically important antibiotic compounds, we utilized a laboratory evolution protocol that partially mimics antibiotic pharmacodynamics in clinical treatments by using strong selection while maintaining population diversity. Using this protocol, we evolved *Escherichia coli* populations with single or multiple, sequential antibiotics until they achieved sustained survival. Although phenotypic outcomes were similar, selection with single or sequential antibiotics drove distinct and reproducible differences in the associated mutational trajectories. We found that bacteria employ diverse resistance, persistence, and tolerance mechanisms, specific to each antibiotic, to evade strong killing effects induced by antibiotics. Most strikingly, sequential antibiotic use shifted evolutionary patterns from divergence to convergence, driven by recurrent extinction of resistance-conferring mutations. Despite convergent genetic trajectories in populations evolved under sequential antibiotic use, single-cell RNA sequencing of selected cultures revealed both unique and shared gene expression patterns across distinct single-cell clusters.

RESULTS

We used three antibiotic compounds throughout this work: levofloxacin, amikacin, and cefepime. Each compound represents a separate antibiotic class, and they were chosen due to their frequent empirical and prophylactic use in clinics, particularly in pediatric stem cell transplant patients[13-15]. Levofloxacin is a fluoroquinolone that targets DNA synthesis through DNA gyrase inhibition[16]. Amikacin is an aminoglycoside that binds the 30s ribosomal subunit causing mistranslation which in turn kills bacteria through proteotoxic stress[17,18]. Cefepime is a cephalosporin, a beta lactam antibiotic, that disrupts a key step in cell wall synthesis[19].

To quantitatively evaluate the evolutionary consequences of prolonged single antibiotic treatment, we evolved ten replicate isogenic cultures of a laboratory strain of *E. coli* (K-12, MG1655, abbreviated here as MG) to levofloxacin, amikacin, or cefepime (30 cultures in total). In addition, four untreated isogenic MG cultures were passaged daily without any antibiotic exposure as a control.

Our evolution protocol was designed to reveal clinically relevant bacterial survival strategies and associated genetic changes. Our protocol involves daily exposures of an antibiotic at a dose corresponding to its reported peak serum concentration in human patients, with durations optimized to achieve approximately 99.9% initial killing of the sensitive parent bacteria (Figure 1b). Here, the doses we used were 8.1 µg/ml of levofloxacin, 69 µg/ml of amikacin, and 129 µg/ml of cefepime[20-22]. These doses are higher than the MIC values of the parent strain (Levofloxacin: ~0.11 µg/ml; Amikacin: ~47.08 µg/ml; Cefepime: ~0.18 µg/ml, Figure 2b-d, Supplemental Figure 2).

Throughout the evolution experiments, we sampled the populations daily to quantify bacterial survival by counting colony forming units (CFUs) before and after antibiotic exposure. Antibiotics were removed by washing, and the surviving bacterial populations were transferred to antibiotic-free growth media (LB: Lysogeny Broth, all experiments were performed at 37 °C) for overnight growth. The next day, a modest 10-fold dilution and outgrowth primed the population for the next antibiotic exposure to avoid significant population bottlenecks and loss of diversity which are common in laboratory evolution protocols[23,24] (Figure 1c). This protocol was repeated over several weeks until bacterial populations largely survived (>30 percent) daily antibiotic exposure as measured by percent survival calculated from CFU counts before and after treatment.

The initial killing was strong for each antibiotic. Average survival on Day 1 to levofloxacin, amikacin, and cefepime was 0.029 ± 0.02 %, 0.058 ± 0.11 %, and 0.527 ± 0.49%, respectively (mean $\pm$ standard deviation; n = 10; Figure 1d-f). As the daily selection cycles continued, levofloxacin and amikacin treatments led to survival trajectories that rapidly increased with a sigmoidal shape toward robust average survivals of 97.2 ± 51.8% and 163.7 ± 116.9% (mean $\pm$ standard deviation) after 16 and 14 treatment days, respectively (Figure 1d-e). Cefepime treatment also ultimately led to a treatment failure equivalent. However, unlike survival trajectories for populations evolved under the selection of levofloxacin or amikacin where the trajectories had sigmoidal shapes, those observed in cefepime treatment gradually increased toward a muted survival with a mean survival of 33.8 ± 25.1% despite 20 treatment days (Figure 1f). There was also more daily variation in survival among the individual trajectories relative to what was observed in experiments under levofloxacin or amikacin selection.

Amikacin resistance and collateral sensitivity evolve through *fusA* and *trkH* mutations.

Next, we assessed if increased bacterial survival was linked to known antibiotic resistance mechanisms. We generated dose response curves for each culture of evolved and parent cohorts in all three antibiotics to quantify cross resistance or collateral sensitivity (Figure 2a). Both MIC and inhibitory concentrations that reduced bacterial growth by half (IC_{50}) were determined (Supplemental Figure 2, Methods). We then employed population Illumina whole genome sequencing (Methods) of bulk cultures, both evolved and parent, to identify mutations and their frequencies within and among the replicate evolved populations.

Cultures evolved under amikacin selection (MG^{AMI}) gained highly significant resistance toward amikacin with the average IC_{50} increasing ~12.7-fold (Figure 2b, p = $7.8x10^{-5}$, student's t-test). Modest but statistically significant collateral sensitivities[5,25,26] (reductions in IC_{50}) were observed toward levofloxacin (Figure 2c, p = 0.012) and cefepime (Figure 2d, p = 0.037).

All ten MG^{AMI} cultures strikingly shared a single high frequency mutation of *fusA* (P610L), a gene which encodes ribosomal elongation factor G that processes the translocation of the ribosome during translation and has been associated with aminoglycoside resistance[5,25]. Additionally, *trkH*, a gene encoding a potassium ion channel[27,28], was a common mutational target. Eight of ten cultures had at least one of eight unique TrkH mutations: L80Q, T20P, G156C, V155E, S105Y, Q159H, L185Q, or a 9-nucleotide deletion causing a frameshift after codon 385. TrKH L80Q was the most common mutation and present in five of ten cultures (Figure 2f). TrkH mutations in aminoglycoside resistant bacteria and

145 their association with collateral sensitivity were previously reported by several groups including
146 ours[25,29]. Additional disparate mutations were observed among the replicate cultures (Supplemental
147 Figure 5a).

Levofloxacin resistance is associated with *gyrA* and *rpoB* mutations.

Cultures evolved under levofloxacin selection (MG[LEV]) acquired resistance toward levofloxacin with the
average IC_{50} increasing ~37.5-fold (Figure 2c, p = $2.4x10^{-5}$, student's t-test). No significant change in
amikacin susceptibility was observed in MG[LEV] populations (Figure 2b). However, mild cross resistance
to cefepime was observed (Figure 2d, p = $7.0x10^{-4}$).

All MG[LEV] cultures had a mutation of *gyrA* which encodes DNA gyrase, a target of levofloxacin,
consistent with previous reports[30]. Five unique GyrA mutations were observed: G81D, S83L, S83W,
A119E, and D87Y, ordered by highest occurrence (Figure 2g). In addition to GyrA mutations, six
MG[LEV] cultures had a synonymous mutation of isocitrate dehydrogenase[31-33] (*icd*: Icd H366H, cac→
cat), and nine cultures had a mutation of RNA polymerase beta subunit[34] (*rpoB*: RpoB H1237L, indel,
D1064N, D1064G, E562D, and M129R). Culture 2 had both RpoB D1064N and the *rpoB* indel with
different frequencies (Figure 2g). Although *rpoB* mutations have been reported following ciprofloxacin
selection, the mechanism underlying this association remains unknown[34]. Interestingly, we have not
observed any efflux related mutations in MG[LEV] cultures even though such mutations are common in
cultures evolved in the presence of gradually increased levofloxacin or ciprofloxacin concentrations[5].

Evolution with cefepime promotes tolerance and persistence, with divergent genetic changes.

To our surprise, cultures evolved under cefepime selection (MG[CEF]) failed to produce robust resistance
to cefepime, despite surviving treatment at cefepime doses almost a thousand-fold higher than the MIC.
The IC_{50} of cefepime rose only modestly (<2-fold) (p= 0.014) (Figure 2d). Additionally, no significant
cross resistance was observed toward amikacin (Figure 2b, p = 0.288) or levofloxacin (Figure 2c, p =
0.324).

We next asked whether bacterial survival in the presence of an extremely high cefepime concentration
was due to a noncanonical survival mechanism such as antibiotic tolerance or persistence. Minimum
duration for killing (MDK) measurements were previously proposed to identify and quantitatively
differentiate these phenotypes (Supplementary Figure 1)[11]. To perform these measurements, cultures
undergoing a continuous high-dose antibiotic treatment are periodically sampled for survival (measured
by CFU counts) to determine the surviving fraction relative to the untreated cultures. Over 7 hours, we
monitored the killing effects experienced by the parent strain (MG) and cefepime evolved (MG[CEF])
cultures (Figure 2e). Three MG[CEF] cultures (numbers 1, 3, and 9) exhibited a biphasic shift in the killing
effect at 1 hour in response to the treatment indicative of a persistence phenotype (Figure 2e, pink lines).
A cohort of five MG[CEF] cultures (numbers 2, 4-6, 10) exhibited a characteristically tolerant phenotype
(Figure 2e, teal lines) with a steady and slower killing effect relative to the parent strain. Two MG[CEF]
cultures (numbers 7 and 8) have a biphasic switch at the 1-hour time point indicative of persistence, but
the survivor fraction was much larger than the previously described persistent cohort. With one-hour
survivor fractions of ~10^{-2}, these cultures appear to have a larger persistent subpopulation or an
intermediate phenotype with characteristics of both persistence and tolerance (Figure 2e, maroon lines).

To compare our persistent and tolerant populations to bona fide resistance, we evolved cefepime
resistant *E. coli* populations (MG[CEF-R]) using a conventional laboratory evolution protocol and evolved
six replicate isogenic cultures of MG under cefepime selection. We grew the cultures in a dilution series
of cefepime concentrations, and each day, for 15 days, we propagated the cultures surviving in the
highest concentrations allowing growth into increasingly higher concentrations of antibiotic
(Supplemental Figure 3a). This resulted in significant resistance to cefepime with the average IC_{50} rising
~227-fold (p = 0.027) (Supplemental Figure 3d). MG[CEF-R] cultures also evolved cross resistance to

levofloxacin (Supplemental Figure 3c), but limited change in amikacin susceptibility (Supplemental
Figure 3b).

We also tested whether this protocol-related effect could be generalized beyond our laboratory strain and
if the outcomes could be considered clinically relevant. Previously in our lab, a clinical, non-pathogenic
E. coli isolate (PEc) was genetically barcoded (PbEc, abbreviated here as Pb) and used to evaluate the
evolution of resistance and persistence to cefepime in mice[5]. In that study, *in vitro* evolution assays
using the conventional protocol were also conducted and produced bona fide cefepime resistance (Pb^{CEF-R}
). We evolved six cultures of Pb under the selection of cefepime using our present protocol for 15 days
(Pb^{CEF}). Like its MG^{CEF} counterpart, we found that the survival trajectory of Pb^{CEF} was gradual toward
muted survival (Supplemental Fig 4a), and ultimately there was no increase in cefepime resistance
(Supplemental Fig 4d). No changes in IC$_{50}$ were observed for Pb^{CEF} populations against amikacin or
levofloxacin (Supplemental Figure 4b-c). Instead, MDK assays revealed distinct tolerant and persistent
cohorts among evolved populations (Pb^{CEF}, Supplemental Fig 4e).

Genetic trajectories of cultures evolved under cefepime selection (MG^{CEF}) were highly divergent,
resulting in numerous unique mutations and idiosyncratic genetic changes (Figure 2h). Mutations in
MG^{CEF} cultures were predominantly found at low frequencies (~10-25%, Supplemental Figure 5c) and
only a few mutations were present in more than one culture. Two genes, *prs* and *phoQ*, were common
mutational targets among subsets of MG^{CEF} cultures. *prs*, encoding PRPP synthase (ribose-phosphate
pyrophosphokinase), had five unique mutations (A114V, R79C, A76V, A11S, and I151T) present in
five of the ten MG^{CEF} cultures, with no single mutation occurring in more than two cultures (Figure 2h).
Likewise, *phoQ*, encoding a transmembrane histidine kinase, had four unique mutations (G39C, an
indel, and two different mobile insertion elements) each present in only one MG^{CEF} culture (Figure 2h).
Mutations of *prs* correspond to MG^{CEF} cultures observed as having a tolerant phenotype in the MDK
assay (Figure 2e, teal), consistent with previous reports[35]. No pattern between mutations of *phoQ* and a
survival phenotype (e.g. persistence) was apparent.

In contrast, cefepime resistant MG^{CEF-R} populations had convergent evolution toward mutations in genes
known to be associated with beta lactam resistance including peptidoglycan transpeptidase[36] (*ftsI*) and
with multidrug resistance including outer membrane porin C[37] (*ompC*), an *ompC* repressor, *envZ*[36,38],
efflux pump gene *acrB* of the *acrAB* operon, its repressor *acrR*[36,39], and the *marR* repressor of the
multiple antibiotic resistance operon(*marRAB*)[40,41] (Supplemental Figure 3e). These differences in
mutational outcomes relative to evolution protocol were also observed in evolved clinical isolate Pb^{CEF}
and Pb^{CEF-R} (Supplemental Figure 4f).

Rationally designed sequential antibiotic treatment results in multidrug survival by resistance,
tolerance, and persistence.

To design a rational sequence for multiple-antibiotic therapy, we analyzed the outcomes of single-
antibiotic treatments. MG^{LEV} cultures remained sensitive to amikacin but had elevated resistance to
cefepime, leaving amikacin as a viable continuation for this cohort (Figure 2b, d). Meanwhile, MG^{AMI}
cultures had small but statistically significant collateral effects on sensitivity to levofloxacin and
cefepime, leaving both antibiotics as viable options (Figure 2c-d). Previous studies have reported on the
collateral effects of aminoglycoside resistance toward beta lactam susceptibility[20,25], so we hypothesized
that using amikacin to treat MG^{LEV} cultures would further sensitize them to cefepime as we observed in
MG^{AMI} cultures and resensitize them to levofloxacin .

Based on this rationale, we evolved the MG^{LEV} cultures under the selection of amikacin (MG^{LEV,AMI})
daily until there was sustained survival, and then we evolved those cultures under the selection of
cefepime (MG^{LEV,AMI,CEF}) toward the same endpoint (Figure 3a). The evolution protocol, dosing, and
durations were the same as their single antibiotic treatment counterparts (Figure 1).

As anticipated, amikacin caused strong initial killing, with average survival on Day 1 (the 17th cumulative treatment day) of $0.356 \pm 0.858\%$ (Figure 3a). However, variability in killing across replicates increased compared to Day 1 of the preceding levofloxacin treatment. This was justified as the terminal MG^{LEV} cultures entered the new treatment with greater genetic diversity than the isogenic parent strain (MG) in our initial experiments (MG to MG^{LEV}). The survival trajectories increased rapidly in a sigmoidal pattern (Figure 3a), leading to robust survival. After 15 days of amikacin evolution (31 days in total), the $MG^{LEV,AMI}$ cultures had an average survival of 102.25 ± 56.44 %, so we initiated the cefepime treatment.

The strong killing effect was restored by cefepime on Day 1 (32^{nd} cumulative treatment day) with an average survival of 0.271 ± 0.292 %, and the variability of the killing effect was high (Figure 3a). Like its single antibiotic treatment counterpart (Figure 1f), the survival trajectories through cefepime were gradual and toward an average modest survival of 39.77 ± 34.73 % even after 50 cefepime treatment days (81 days in total).

Evolved bacterial cultures from the final days of each segment of the sequential antibiotic evolution were evaluated for resistance to all three antibiotics (Figure 3b-d, Supplemental Figure 7). Following the amikacin segment of the sequential evolution, $MG^{LEV,AMI}$ cultures became significantly more resistant to amikacin, an ~8-fold change ($p = 4 \times 10^{-6}$) relative to MG^{LEV} cultures (Figure 3c). $MG^{LEV,AMI}$ cultures were also moderately resensitized to levofloxacin relative to MG^{LEV} cultures (~2-fold, $p = 5 \times 10^{-3}$), but $MG^{LEV,AMI}$ cultures remained significantly more levofloxacin resistant than parent MG cultures ($p = 2 \times 10^{-3}$, Figure 3b). As we hypothesized, evolution under the selection of amikacin (MG^{LEV} to $MG^{LEV,AMI}$) reversed the cross resistance to cefepime observed in MG^{LEV}, resulting in a reduction in the average IC_{50} from 0.082 ± 0.024 µg/mL in MG^{LEV} to 0.052 ± 0.026 µg/ml in $MG^{LEV,AMI}$ ($p = 8 \times 10^{-3}$). This IC_{50} value was not significantly higher than the cefepime IC_{50} in the parent MG cultures (Figure 3d).

After evolution under the selection of cefepime ($MG^{LEV,AMI}$ to $MG^{LEV,AMI,CEF}$), levofloxacin resistance did not change significantly relative to $MG^{LEV,AMI}$ (Figure 3b, $p = 0.3$). However, to our surprise, $MG^{LEV,AMI,CEF}$ was significantly resensitized to amikacin relative to $MG^{LEV,AMI}$ where the average amikacin IC_{50} dropped to 9.42 ± 6.15 µg/ml from 26.23 ± 7.86 µg/ml ($p = 10^{-4}$) (Figure 3c). This phenotypic change was not observed in the single antibiotic treatment counterpart (MG^{CEF}). Ultimately, $MG^{LEV,AMI,CEF}$ had no change in cefepime resistance relative to $MG^{LEV,AMI}$ (Figure 3d, $p = 0.395$). Rather, MDK assays with $MG^{LEV,AMI,CEF}$ cultures again revealed that the increased survival was associated with persistence and tolerance phenotypes (Figure 3e) similar to MG^{CEF} populations. Two $MG^{LEV,AMI,CEF}$ cultures (culture 2 and 5) were characteristically tolerant (Figure 3e, teal lines). Two cultures (numbers 3 and 8) had an intermediate phenotype (Figure 3e, maroon lines). The remaining six cultures (numbers 1, 4, 6, 7, 9, and 10) were characteristically persistent (Figure 3e, pink lines) with a biphasic killing curve. The shift occurred at survivor fractions between 10^{-2} and 10^{-3} indicating relatively large persistent subpopulations (Figure 3e). The mix of persistence and tolerance phenotypes was similar to what was observed in MG^{CEF} cultures (Figure 2e) and Pb^{CEF} cultures (Supplemental 4e).

Sequential antibiotic treatment drives unique and convergent evolutionary trajectories

Next, we asked how the evolutionary context and history impacted mutational routes toward survival by Illumina whole genome sequencing of frozen samples isolated from evolved cultured at different time points (Figure 4). When we tracked the five GyrA mutations of MG^{LEV} through the subsequent evolutions under the selection of amikacin and cefepime, we observed that GyrA S83L had an advantage in the amikacin treatment regime where it was either fixed or it replaced the previously dominant GyrA mutations (Figure 4a, d, Supplemental Figure 8a). Strikingly, by day 34, S83L was the dominant GyrA mutation across all cultures and all other GyrA mutations went extinct. Mutations of RpoB on the other hand were also selected against during amikacin treatment and were driven to extinction (Supplemental 8b). In all but culture number 2, RpoB mutations became nearly extinct by the conclusion of amikacin

treatment (day 31). In culture 2, the *rpoB* indel survived amikacin treatment, but subsequently was
selected against through cefepime treatment.

Like MG^{AMI}, MG^{LEV,AMI} cultures acquired mutations in *fusA* and *trkH* genes, yet the number and type of
mutations differed. Where MG^{AMI} cultures had only the FusA P610L mutation, MG^{LEV,AMI} cultures
carried 11 unique FusA mutations (including P610L). This difference was most likely due to greater
initial population diversity of MG^{LEV} cultures before amikacin selection started. Interestingly, all FusA
mutations failed to maintain high frequencies and were subsequently driven to extinction by the final
cefepime treatment day (Figure 4a, Figure 4d, Supplemental Figure 8c), indicating a potential fitness
cost for FusA mutations under cefepime treatment.

In the case of *trkH*, MG^{AMI} cultures had eight different TrkH mutations, while MG^{LEV,AMI} cultures had
only three: T20I, L80Q, and L185Q. MG^{LEV,AMI} cultures possessed high frequency TrkH L185Q
mutations which was the dominant allele, a stark contrast to just one MG^{AMI} culture having the TrkH
L185Q mutation at a low frequency (0.11). TrkH L185Q quickly fixed within MG^{LEV,AMI} cultures
throughout amikacin treatment and maintained in the subsequent cefepime treatment (Figure 4a,
Supplemental Figure 8d).

The evolution of survival under the selection of cefepime in MG^{LEV,AMI,CEF} cultures was markedly
convergent. In contrast to the genetically divergent evolution observed in MG^{CEF}, each of the ten
MG^{LEV,AMI,CEF} cultures acquired nearly identical mutations among a set of nine genes at various times
(Figure 4b-d, Supplemental Figure 8e). Among the nine mutations were GyrA S83L and TrkH L185Q,
which were inherited from previous selection regimens under levofloxacin and amikacin and survived
extinction events during subsequent treatment regime changes to amikacin and cefepime, respectively
(Figure 4d). The additional mutations emerged with a variable cadence across individual cultures
(Figure 4b). Concurrent to TrkH L185Q, a single nucleotide deletion in *glvC*, a poorly characterized
putative phosphoenolpyruvate (PEP)-dependent carbohydrate transport system (PTS) component[42,43],
evolved with nearly identical frequencies indicating a common clade and positive selection for the
mutated *glvC* (Figure 4d). The deletion occurred at nucleotide 1042 of 1072 generating a downstream
stop codon that likely truncates the protein by 17 amino acids.

Next emerged two other concurrent mutations in *selB* and intergenically between *fimB* and *fimE*
(fimB/E). *selB* (Figure 4b-d), encoding a translation factor involved in incorporation of selenocysteine
into proteins[44], had a 2-nucleotide insertion that created a frame shift at amino acid 494 and an early stop
codon at 526, possibly truncating what is typically a 614 amino acid protein. *fimE*, a regulatory gene
that, with *fimB,* controls the production of type 1 fimbriae[45], was likely lost following an IS5 mobile
insertion element inserting between the *fimE* promoter and its transcription start site[46]. This mutation
pair was observed in all cultures beside number 2, and either emerged with TrkH L185Q and the
mutated *glvC* or after.

All MG^{LEV,AMI,CEF} cultures subsequently had a 7-nucleotide long deletion in *rpoZ*, which encodes the
omega subunit of RNA polymerase (Figure 4b-d). The deletion occurs relatively early in the gene (68-74
of 276 nucleotides) and introduces an early stop codon after just 29 amino acids, a severe truncation of
what is typically a 91 amino acid protein. In all cultures except culture 2, a mutation was observed in
hipA (HipA P86L) (Figure 4b-d), a gene named for its association with a highly persistent phenotype[47].
Additionally, mutations of *ftsH*, a gene encoding a membrane-bound protease involved in stress
response[48,49], and of *gatA*, a known galactitol-specific component of the PTS system[50,51] were observed
in all cultures besides numbers 2, 9, and 10. The *ftsH* gene incurs an in-phase deletion of 12 nucleotides
removing codons 261-264 of 644 within its AAA+ domain. *gatA*, meanwhile is disrupted by the mobile
insertion element IS5 following its 69[th] amino acid (Figure 4b-d). By the final day, many of the
discussed mutations had reached high frequencies, and those not fixed shared similar frequencies with
others, indicating their concurrence within a common clade (Figure 4b-c).

The shift from divergent evolution under the selection cefepime as a single treatment in MG^{CEF} toward convergent evolution when cefepime was used a sequential treatment in MG^{LEV,AMI,CEF} was striking and not previously reported to best of our knowledge (Figure 4d). These shifts are illustrated in Figure 4d where bacterial populations evolve under the selection of an antibiotic (color) through various mutational options (arrows) given the single vs sequential context of the evolution. Many mutations within evolved populations are driven to extinction within subsequent antibiotic exposures (Supplemental Figure 9).

Individual convergent mutations are insufficient to produce the observed evolved population level persistence.

To evaluate the impact of the commonly observed mutations in MG^{LEV,AMI,CEF} cultures, we introduced each mutation into the wild-type, parental MG1655 (MG) genome (Methods)[52]. The single mutation strains produced from this mutagenesis are referred to going forward as their parent strain background (MG) with the mutated gene in superscript (e.g., MG^{gatA}, MG^{glvC}).

To evaluate the phenotypic changes in the mutant strains, we measured changes in their exponential growth rate, susceptibility to antibiotics (IC$_{50}$, MIC), and survival by persistence or tolerance in the presence of cefepime (MDK assay, Supplementary Figure 2). The parent strain (MG) had a mean doubling time of 20.07 ± 0.80 minutes (Figure 5a). Mutant strains had limited growth defects. The greatest effects were observed for MG^{gatA} (21.17 ± 0.92 minutes, p = 0.0501), MG^{selB} (21.62 ± 0.41 minutes, p = 0.0017), MG^{rpoZ} (21.91 ± 0.88 minutes, p = 0.0036), and MG^{fimE} (21.66 ± 0.57, p = 0.0026).

We tested resistance of the individual mutant strains to levofloxacin, amikacin, and cefepime. As before, IC$_{50}$ and MIC were estimated from growth data following overnight growth at 37°C in a range of antibiotic concentrations. Mutants MG^{gatA}, MG^{glvC}, MG^{rpoZ}, and MG^{fimE} did not have any resistance change to any of the three antibiotics (Figure 5b-d). Modest levofloxacin resistance was observed for MG^{hipA} (p=0.014) and MG^{selB} (p=0.046) (Figure 5c). MG^{selB} also had slightly elevated resistance to amikacin (p=0.002). MG^{ftsH} was mildly sensitized to amikacin (p=0.04) (Figure 5b). No changes in cefepime resistance were observed for any of the mutants we engineered.

Next, we used the MDK assay to identify the impact of the mutations on survival by tolerance and persistence. The parent strain (MG) is quickly killed by the effects of cefepime in the first hour with a surviving fraction just above 10^{-4} (Figure 5e). Many of the mutant strains were killed at the same rate as parent MG through each timepoint. Only MG^{hipA} was less susceptible through 1 hour with a surviving fraction of 10^{-3}. The rate of killing slowed considerably for MG^{hipA} relative to MG and the other mutants and then continued to have roughly 10-fold greater survival at every time point. Ultimately after 7 hours, MG^{hipA} had a surviving fraction above 10^{-5}, while the other cultures were at or below 10^{-6}. The biphasic shift in its killing is indicative of persistence, in agreement with previous observations of the phenotype in *hipA* mutants.

The relatively high frequencies and penetrance of the other mutations among the replicate MG^{LEV,AMI,CEF} populations suggest a positive selection in the presence of cefepime. This observation strongly indicates that their contribution to survival and fitness is either epistatic in nature or acts in support of the HipA mutation-led phenotype. Indeed, when the survival of MG^{hipA} is directly compared to that of a diverse, evolved culture, MG^{LEV,AMI,CEF}-7, the presence of additional mutations is likely to be associated with enhanced survival. A similar pattern was observed in MG^{prs}, a strain engineered with the Prs A114V mutation, and its corresponding diverse, evolved culture, MG^{CEF}-5 (Supplemental Figure 10d).

In summary, *hipA* and *prs* mutations display persistence and tolerance phenotypes, respectively. However, evolved cultures carrying the exact same mutations exhibit stronger persistence or tolerance phenotypes, suggesting the presence of epistatic interactions between *hipA* and *prs* mutations and the other mutations observed in the evolved populations.

Single cell RNA sequencing of evolved cultures reveals metabolic subpopulations and hipA overexpression

To determine how the diverse, accumulated mutations influence bacterial community composition, we used single-cell RNA sequencing (scRNAseq) to characterize changes in the gene expression landscape resulting from sequential antibiotic exposure. Samples were collected from exponentially grown cultures at the same optical density ($OD_{600} = 1.0$) of the parent strain and two cultures that accumulated largely overlapping mutations: MG^{LEV,AMI,CEF} -4 and MG^{LEV,AMI,CEF}-7. Using ProBac-seq[53], a probe based bacterial scRNAseq method, we analyzed a total of 48,883 single cells from duplicate samples of these three cultures (Supplementary Table 3).

Using dimensionality reduction and graph-based clustering, cells were grouped into 9 unique clusters (Figure 6a). While most cell clusters included at least a few cells from each culture, there were notable instances in which specific clusters were predominantly composed of cells from distinct cultures (Figure 6b, c). The majority of cells in clusters 1 and 2 were of parental origin (>67% and >70% parent cells in each cluster, respectively). Gene-set enrichment analysis (GSEA) identifies translation-related processes and expression of ribosomal subunits as highly enriched in the parental-dominated clusters (Supplementary Table 3, Supplementary Figure 11) These same gene sets were enriched when comparing the wildtype to either culture without cluster-based analysis (Figure 6d, e and supplementary table 4). The signature of lower translational activity in the evolved populations is consistent with the common observation that antibiotic persistence is associated with lower activity levels[54]. For example, expression of the 50S ribosomal subunit L10 (*rplJ*) is higher in parent cultures, and significantly upregulated in clusters 1 and 2 (Figure 6 d-f, Supplementary Tables).

Cells from each of the evolved cultures were commonly and significantly enriched in transcripts for the superpathway of glycolysis, pyruvate dehydrogenase, TCA, and glyoxylate bypass, with several glyoxylate shunt genes overexpressed in these cultures (shared genes in Figure 6d, e and Supplementary Table 4). Culture MG^{LEV,AMI,CEF}-4 comprised the majority of cells in cluster 9 (>72%), which itself was enriched in this superpathway (Figure 6c, Supplementary Figure 12, Supplementary Table 4). Similarly, culture MG^{LEV,AMI,CEF}-7 comprised over 80% of cluster 4 cells, which were enriched in this superpathway (Figure 6c, Supplementary Figure 12, Supplementary Table 4).

In summary, single cell analysis demonstrates that the evolved cultures have cells occupying distinct metabolic niches and contain fewer cells in states that are consistent with high translation activity. This pattern is in line with changes that would be expected from mutations that affect toxin-antitoxin genes including *hipA* identified here. Expression of *hipA* itself was significantly amplified in MG^{LEV,AMI,CEF}-4, while in MG^{LEV,AMI,CEF}-7 too few individual cells were profiled as having significant overexpression of *hipA* (Figure 6 d-e). The distribution of *hipA* expression shifted strongly toward clusters 4 and 9, key clusters within the evolved lineages (Figure 6g). The observation mirrors the subpopulation survival effect which is characteristic of persistence.

DISCUSSION

Predicting how bacteria survive antibiotic treatments is crucial for designing 'evolution-proof' antibiotics and treatment strategies. Laboratory evolution experiments, whether conducted using *in vitro* or *in vivo* models, have been instrumental in advancing our understanding of antibiotic resistance evolution[55,56]. In this study, we demonstrate that treatment protocol (i.e. high antibiotic peak doses observed in human patients) and context (i.e. whether an antibiotic was administered as a single exposure or in sequence), had consequential effects on phenotypic bacterial survival strategies and the mutational routes toward bacterial resistance.

With high-dose, clinically relevant treatments, we demonstrated that use of levofloxacin and amikacin led to canonical resistance evolution while use of cefepime led to persistence and tolerance evolution, and these phenotypic outcomes were independent of the single or sequential context of the treatments

(Figures 2, 3). Likewise, the outcomes of persistence and tolerance were independent of the population
structure (isogenic or polyclonal) and bacterial strain (lab strain or clinical isolate) (Figure 2, 3 and
Supplemental Figure 4).

We also observed context-specific evolution of collateral antibiotic sensitivity[5,25,26]. Amikacin as a
single treatment had only minor collateral effects on sensitivity toward levofloxacin and cefepime
(Figure 2c-d), but amikacin as a sequential treatment following levofloxacin substantially reversed the
levofloxacin-associated cross resistance toward cefepime (Figure 3b,d). A context-specific evolutionary
outcome was also observed with cefepime treatment where cefepime as a single treatment (MG^{CEF}) had
no effect on cross-resistance to amikacin (Figure 2b), but when used as a subsequent treatment following
amikacin, cefepime reduced amikacin resistance of evolved populations (Figure 3c) by selecting against
FusA mutations (Figure 4a-ii, Supplementary Figure 8c).

We observed that sequential antibiotic treatment can shift evolutionary trajectories from genetically
divergent to convergent. In such scenarios, evolutionary constraints such as epistasis between mutations
or fitness costs of mutations in the presence of different antibiotics may provide bacterial populations
with fewer viable mutational routes toward resistance in subsequent treatments creating an opportunity
for rational treatment design.

When we tested the individual convergent mutations selected under sequential antibiotic treatment for
their contributions to survival, we observed that only the HipA mutation resulted in persistent phenotype
(Figure 5e). HipA is a kinase that activates persistence and the stringent response by driving the
accumulation of the alarmone ppGpp. Further, HipA dimerizes and forms a complex with a HipB dimer
to repress the *hipBA* operon, and mutations have been shown to weaken the interaction, allowing greater
free HipA and less transcriptional autoregulation of the operon[57-61]. While MG^{hipA} was the only mutant
strain with increased survival by persistence, the degree of its survival throughout the MDK assay was
lower relative to the survival observed in MDK assays of evolved MG^{LEV,AMI,CEF} (Supplementary Figure
10). We hypothesize that the additional convergent mutants contribute to survival through epistatic
interactions among them. Among them, mutations in *ftsH* and *rpoZ* may have the clearest connection to
the persistent state associated with the HipA mutant.

FtsH was previously shown to degrade stress response regulators to allow bacteria to return to a
prestressed state[48,49], and degrade the LpxC and KdtA enzymes that are involved in the early steps of
LPS biosynthesis[62]. Interestingly, the FtsH-mediated degradation of LpxC is modulated by ppGpp[63]. The
4 codon deletion within the AAA+ domain of FtsH observed here may alter ATP binding or protease
efficiency as a way of adapting to prolonged stress or to compensate or potentiate the HipA persistence
phenotype[64].

RpoZ, the non-essential omega subunit of RNA polymerase, interacts with other RNA polymerase
subunits which creates a binding interface for ppGpp that contributes to the propagation of the stringent
response and persistence phenotype[65,66]. The proposed connection to HipA is further supported in the
observed cadence of mutation emergence, where mutations of FtsH and RpoZ always coincide or
succeed the mutation of HipA throughout evolution under sequential antibiotic treatment.

The results of our single cell RNA sequencing analysis reveal information about transcriptional
complexity within the evolved populations. Clustered cells with widespread expression changes of
metabolic and translation programs highlight additional opportunities to characterize bacterial
susceptibility following distinct treatment regimes. It remains unclear the exact mechanistic role for the
transcriptional shifts in superpathway genes, but it may likely lead to metabolic shifts affecting growth
rates and cell states which contribute to antibiotic survival. Our analysis revealed reduced expression of
ribosomal genes and overexpression of *hipA* in subpopulations of evolved cells consistent with the
biphasic killing effect observed in MDK assays, despite the high frequencies of the mutation.
(Supplemental Figure 10d). Akin to the allele-specific susceptibility and survival we observed, we

hypothesize that some bacterial subpopulations differentially expressing distinct gene sets will have
greater survival during antibiotic treatment.

This work offers an initial roadmap for identifying sequential antibiotic treatments that promote
favorable mutational trajectories in the face of multiple treatment failures. Our ability to predict
mutational routes toward resistance, persistence, and tolerance may improve antibiotic choice based on
known collateral phenotypic effects. Further, as shown here, sequential treatments can shift evolution
from divergent to convergent routes where bacteria may have fewer mutational options to surviving
subsequent treatments, marked by extinction events and variable allelic diversity. In pursuit of rational
treatment design and improved clinically efficacy, future investigations of this kind could be performed
experimentally, as was done here, with more antibiotics and sequences, or rather by sequencing clinical
isolates following sequential antibiotic treatment courses.

483
484 **Materials and Methods**

485
Bacterial strains
Lab strain *Escherichia coli* (MG1655), a genomically barcoded, clinical isolate *Escherichia coli*
(PbEc)[56], and mutant MG1655 clones generated by oligo recombineering were routinely cultured in
lysogeny broth (LB) medium (RPI) aerobically (shaking at 230 rpm) at 37°C. When applicable, growth
was determined by spectrometry (OD_{600}), and survival was determined by enumeration of cultured
colony forming units (CFU) on LB agar plates (RPI). Plate-based assays (MIC) were performed in LB
medium with aerobic growth (shaking at 400 rpm) at 37°C and controlled humidity (80%).

493
494
495 **Antibiotic Stock Preparation**
496 Concentrated stocks of antibiotics levofloxacin (Alfa Aesar) and cefepime (Apotex) were prepared in
497 water, filter sterilized, and stored at -20°C for single use, each 10mg/mL. Amikacin sulfate (Heritage)
498 was diluted as needed from purchased liquid stock.

499
Colony forming unit enumeration
Multiple assays relied on the enumeration of colony forming units. To perform these measurements,
each well of 96 well plates was filled with 180 µL of phosphate buffer saline (PBS, Sigma) solution. 20
µL samples of cultures were then serially diluted (1:10) across the rows of 96 well plate up to a 10^{-6}
dilution. 20 µL of each dilution was plated on LB agar plates in lines, and agar plates were incubated
overnight (37°C). The following day, individually colonies were counted from the plated dilutions with
the maximum countable colonies. When necessary, cultures or aliquots were centrifuged and washed
with PBS to remove antibiotic prior to diluting and plating.

508
Evolution Assays
The evolution assay developed presently was repeated daily beginning with a 1:10 dilution of overnight
cultures into 10 mL of antibiotic-free LB media. Diluted cultures were grown for 1 hour. Prior to
treatment, a 20 µL sample was added to the 96 well plates prepared for CFU enumeration. The
remaining culture was dosed with antibiotic and returned to shaking incubation. Diluting and plating
related to CFU enumeration was performed during the incubation period. Incubation times were
previously optimized to achieve 99.9% killing on the initial day. After the necessary incubation time (1
hour for levofloxacin and amikacin; 2 hours for cefepime), bacteria were pelleted by centrifugation
(6000g, 5 minutes) and washed twice with PBS to remove the antibiotic. The cleaned pellets were
resuspended with the equivalent volume of antibiotic-free LB media, and a post-treatment 20µL sample
of the culture was collected for CFU enumeration. Cultures were promptly returned to shaking and
incubation for overnight growth. The protocol was repeated daily until an increased sustained survival

was observed. Percent survival was calculated by dividing post-treatment CFU counts by pre-treatment
CFU counts. Sequential antibiotic treatments immediately followed the previous antibiotic treatment.
523
For the conventional laboratory evolution protocol, overnight cultures were diluted in antibiotic-free LB
media (1:100). Antibiotic was serially diluted (1:2) across 5 mL LB media in 15 mL conical tubes to
prepare a range of concentrations surrounding a predetermined MIC (MIC/4, MIC/2, MIC, 2xMIC,
4xMIC). Parental strain bacteria from the overnight growth were added (1:500) to tubes containing
antibiotic and then grown aerobically overnight as described previously. The following day, the tube
with the highest antibiotic concentration that allowed for robust growth was used to inoculate (1:500)
freshly prepared conical tubes with an increased range of antibiotic concentrations relative to the prior
day, centered around the passaged concentration (e.g., MIC/2, MIC, 2xMIC, 4xMIC, 8xMIC). This was
repeated daily.
533
534   Routinely, bacteria were collected and saved as a glycerol stock by combining 500 µL of bacteria
535   culture with 500 µL 50% glycerol.
536
Resistance Determination
In 96 well plates, twofold dilutions series of antibiotics were prepared in 100 µL of LB media. Then,
100 µL of diluted bacteria cultures (OD_{600}: 0.01) was added. Each plate included "Media Only" wells to
measure background OD signal and "Cells Only" wells to assess maximum growth in the absence of
antibiotic. Plates were incubated overnight at 37°C, shaking at 400 rpm, in a humidified (80%) incubator
(Infors HT). Following incubation, OD_{600} was measured (Biotek Epoch2 plate reader), and raw OD
values were exported and processed using a custom Python script.
544
First, a global or plate-specific median background was computed from the "Media Only" wells. This
median value was subtracted from the raw data to correct for background signal. Next, each strain–
antibiotic dataset was normalized by dividing the background-corrected OD values by the median OD of
its corresponding "Cells Only" wells, yielding fractional growth that ranged from near 0 (complete
inhibition) to 1 (no inhibition). Using these normalized data, a two-parameter logistic (2PL) curve was
fit to each strain–antibiotic combination. Specifically, the present function was employed to model
growth

$$y(x) = \frac{1}{1 + \left(\frac{x}{IC_{50}}\right)^h}$$

552
553   where x is the antibiotic concentration, IC_{50} is the concentration at which growth is half the no-antibiotic
554   control, and h is the Hill slope reflecting the steepness of the inhibition curve. Experiments were
555   conducted with technical triplicates across 12 antibiotic concentrations, ensuring sufficient data points
556   across multiple conditions for robust curve fitting.
557
Parameter optimization was performed via SciPy's least_squares routine with a robust "dogbox"
algorithm and "soft_l1" loss, which reduces sensitivity to outliers. The minimum inhibitory
concentration (MIC) was then defined as the concentration at which growth reached 5% of the no-drug
control (i.e., 95% inhibition). This value was obtained by numerically solving the fitted 2PL function at
y=0.05. To quantify uncertainty in IC_{50} and MIC estimates, a bootstrap analysis was performed by
resampling each dose–response dataset 1,000 times with replacement and re-fitting the curve. The 2.5[th]
and 97.5[th] percentiles of these bootstrap distributions defined 95% confidence intervals, providing robust
estimates of parameter variability. When the model-predicted IC_{50} or MIC exceeded the maximum
tested antibiotic concentration, an "insufficient drug" flag was assigned, indicating that the true

inhibitory concentration likely lay beyond the tested range, and excluded from further analysis. All
original code is available in this paper's supplemental information.
569
Minimum duration of killing (MDK) assay

Overnight *E. coli* cultures were diluted 1:10 in antibiotic-free LB media and grown aerobically for 1
hour. Initial CFUs were measured as described previously. The cultures were then exposed to 129
µg/mL of cefepime for 7 hours. At hours 1, 2, 3, 5, and 7, 500 µL aliquots were collected in 1.5 mL
microfuge tubes, centrifuged, and washed twice with PBS to remove the antibiotic. Pellets were
resuspended in 500 µL PBS. At hours 5 and 7, final pellets were resuspended in 50 µL PBS, 20 µL of
which was plated and 20uL was diluted further, to increase the limits of detection. CFUs were
enumerated, and survivor fraction was calculated by dividing the ongoing counts by the initial CFU
measurement.

Bacterial population sequencing

Bacteria cell pellets of antibiotic-evolved, passaged untreated control, and parental cultures were
prepared from overnight cultures and sent to Seqcenter (Pittsburgh, PA) for DNA extraction
(ZymoBIOMICS) and whole genome sequencing (Illumina). Samples were sequences using 150 base
pair paired-end reads. For evolved cultures, approximately 2 giga base pairs of data per sample was
received yielding coverage of least 100x throughout the genome. Sequencing data was processed using
breseq (Barrick lab), comparing reads to a MG1655 reference genome [67]. Alleles present in the
passaged, untreated controls were tracked but were removed from the dataset of evolved populations.
Likewise, alleles present in the parental strain relative to the reference genome were subtracted.
Remaining mutations were filtered to include only those which exceeded 10% at any timepoint and
these alleles were traced across timepoints. Commonly mutated genes were presented for each single
antibiotic evolution with full mutation sets presented as supplemental figures. The full breadth of
mutations within the cohorts of evolved cultures following single and sequential evolutions are shown in
Supplemental Figure 9, with the exlusion of MG^{LEV,AMI,CEF} cultures 2, 3, and 8 which had exceedingly
numerous mutations associated with a hypermutator mutation of *mutL*. Further, the *mutL* mutation was
first identified in culture number 2 during the amikacin regime. Later, on day 81, *mutL* was observed in
cultures 3 and 8 along with other mutations present in culture 2, making it difficult to rule out a possible
contamination of those cultures sometime between days 63 and 81.

Oligo recombineering

Chromosomal mutations were made using oligo recombineering, and gene deletions were made using
oligo recombineering followed by Bxb-1-Integrase-Targeting (ORBIT). These methods each incorporate
mutations encoded on transformed oligonucleotides through homologous recombination during DNA
replication, and require elements encoded on a helper plasmid[52]. Parental MG1655 bacteria were made
electrocompetent and then transformed with the pHelper_Ec1_V1_gentR (pHelper-V1) for oligo
recombineering or with pHelper_noMutL_V2_gentR (pHelper-V2) for ORBIT. Selection was made
possible by gentamicin. pHelper-V1 expresses single stranded DNA annealing proteins (SSAP),
CspRecT, and a dominant negative MutL, which improves editing efficiency by suppressing mismatch
repair[68]. Sucrose counterselection is made possible by *sacB* gene encoded on the plasmid. The
expression is driven by the induction of the XylS/m-toluic acid system. pHelper-V2 differs in that it
does not include the dominant negative MutL. Transformed bacteria were induced with toluic acid for 1
hour prior to storage at -80°C.
611
For simple oligo recombineering, 90-nucleotide long oligonucleotides were designed with the desired
mutations at the center flanked by homology regions within the gene, and each was ordered from
Integrated DNA Technologies (IDT) (Table1). Oligonucleotides were prepared and diluted to 25 µM in
water. 50 µL of MG1655-pHelper-V1 was mixed with 2 µL of diluted oligonucleotide, electroporated

(1.8mV), and recovered in LB media for 1 hour at 37°C. Recovered bacteria were steaked on LB agar
for single colonies and incubated overnight at 37°C.
618
619 Individual colonies were picked into 100 µL PBS in 96 well plates for screening by polymerase chain
620 reaction (PCR). Throughout screening, colonies were routinely substreaked on LB-sucrose (7.5%) plates
621 to drive the removal of the pHelper-V1 plasmid. Plasmid loss was confirmed by plating on LB-
622 gentamycin (15 µg/mL) plates.
623

Preparation of electrocompetent cells
625 MG1655 cultures were made electrocompetent prior to transformations with plasmids pHelper-V1 and
626 pHelper-V2, and subsequently MG1655-pHelper-V1 and MG1655-pHelper-V2 were made
627 electrocompetent prior to transformations with oligonucleotides. Each approach varied slightly.
628
Beginning with plasmid transformations, filtered water was aliquoted in 50 mL conical tubes and
refrigerated overnight. MG1655 was cultured overnight in 10 mL LB media. The following day, a 1:100
dilution of the overnight was performed into a fresh 10 mL LB and grown for 2 hours. During this time,
the water aliquots, empty 15 mL conical tubes, and microfuge tubes were placed on ice. After 2 hours of
growth, the culture was split into two, 15 mL tubes. These cultures were centrifuged at 4000g, 2°C, for
10 minutes. The pellets were washed twice with 10 mL of cold water. The final supernatant was fully
removed by pipetting. Pellets were then resuspended in 1 mL of cold water, combined, and then split
into 2 microfuge tubes. The microfuge tubes were centrifuged for 10 minutes, and each pellet was
resuspended in 100 µL of cold water. In separate tubes, 100 µL of the culture was combined with 20 ng
of either plasmid and each was electroporated (2.5 kV). Immediately, 1 mL of prewarmed recovery
media (NEB) was added and moved to a 15 mL conical tube where it recovered for 1 hour at 37°C in the
shaking incubator. Following its recovery, 200 µL of each culture was plated on LB agar plates
containing 15 µg/mL gentamycin for positive selection of transformants. A non-transformed culture was
used as a control. After overnight incubation, single colonies were picked and grown overnight in liquid
culture with gentamycin (15 µg/mL).
644
The following day, the cultures, transformed MG1655-pHelper-V1 and MG1655-pHelper-V2 were
induced and made competent again. Filtered water and 10% glycerol had been refrigerated overnight.
From the overnight cultures, a 1:1000 dilution was made into 200 mL of LB with gentamycin and grown
for 3-4 hours to ~ 0.3 OD. While growing, reagents and tubes were placed in ice. When the cultures
reached 0.3 OD, the cultures were induced with 1 mM toluic acid (added from a 1000x stock solution)
and incubated for 30 minutes. Following incubation, the cultures were removed and placed in an ice bath
with occasional swirling to stop their growth. Then 50 mL aliquots of the cultures were transferred to
iced conical tubes. These were centrifuged at 5000g for 10 minutes at 2°C. The supernatant was poured
off, and pellets were resuspended with 45 mL cold water. Resuspended pellets were centrifuged, and
pellets were resuspended with 10% glycerol. This was repeated twice more, but the final pellets were
resuspended in the residual glycerol. Cultures of the same strain were pooled and then aliquoted (100
µL) into small microfuge tubes. These were chilled on dry ice for 5-10 minutes before being stored at -
80°C. These resulting stocks were competent and induced to allow for the transformation and
incorporation of targeting oligonucleotides.
659
Cultures transformed with oligonucleotides were plated on non-selective plates, and thus colonies
required screening by PCR and then confirmation by Sanger Sequencing (Eurofins). For each mutant
strain prepared, full-length and mutant-specific PCR primers were designed (Table 2). The mutant-
specific primers relied on the small differences in sequences and were paired with the full-length primer
in the opposite direction. PCR reactions were performed using GoTaq polymerase (Promega), the
appropriate primer set, water, and a 1 µL inoculation of the colony-PBS suspension. Prior to screening,

the melting temperatures for reactions using mutant specific primers were optimized by gradient PCR of
parent strain and a pool of presumptive mutant colonies. Then, batches of individual colonies were
screened by mutant-specific PCR. Mutant PCR reaction products were routinely run on 1% agarose gels
with ethidium bromide and imaged under ultraviolet light (Syngene G:BOX Chemi XT4 with GeneSys
software). Positive colonies were then PCR amplified using full-length primer pairs. These full-length
PCR products were cleaned (Nucleospin Gel and PCR cleanup kit) and submitted for Sanger sequencing
(Eurofins).
673
Analysis of the sequencing results often revealed mixed colonies with peaks representing mutant and
wild-type nucleotides. When this occurred, positive colonies were substreaked (LB-sucrose), and
rescreened. After several iterations, Sanger sequencing results were pure, and the mutational status was
confirmed by sequencing further subclones for purity. For the larger mutations made by oligo
recombineering, the efficiency was lower and required a pooling approach to minimize reagent use and
increase throughput. When mutants were confirmed and purity achieved, single colonies were picked
and grown in LB overnight and then saved as stocks in 25% glycerol.
681
Bacterial growth assay
Strains were cultured overnight from glycerol stocks in 10 mL LB media. The following day, a 96 well
plate was prepared with 100 µL LB media in each well. Overnight cultures were diluted in fresh media
to an OD_{600} of 0.1. To media-only wells, an additional 100 µL LB media was added. To culture wells,
with six replicates of each strain, 100 µL of diluted culture was added. The plate was promptly
incubated in Biotek Epoch2 plate reader and grown for 20 hours at 37°C with shaking. Every 2 minutes,
OD measurements were made. Growth curves were analyzed using a custom Python script. OD readings
were first filtered to values between 0.02 and 0.08 to capture the log growth phase. Linear regressions
were performed on the log transformed OD values within this range as a function of time. The resulting
slope was converted to a natural log-based growth rate. Doubling time was then calculated using the
formula
693

$$\frac{\ln(2)}{growth\ rate}$$

694
695 and reported in minutes. The scipy.stats module was used to calculate means, standard deviations, and to
696 perform t-tests of doubling times.
697
Cell fixation for single cell RNA sequencing
Overnight cultures of strains were inoculated (100uL in 10mL) from frozen stocks, avoiding population
bottlenecking. Cultures were diluted to 1 OD, and 1 mL of culture was fixed with 67µL of
paraformaldehyde through a 30 minute incubation. Paraformaldehyde was removed by centrifugation (5
min, 6000 rcf). Pellet was resuspended with 1 mL of 0.2x SSC Buffer (Invitrogen). Buffer was removed
by centrifugation, the pellet was resuspended in 350 µL of MAAM (4 parts methanol, 1 part acetic acid),
and briefly stored at -20°C.
705
Single cell RNA sequencing
Fixed cells were processed using standard ProBac-seq protocols as follows: 100 ul of cells in MAAM
solution were centrifuged at 6000 rcf and washed 3 times in 300 ul PBS solution. After PBS washes
cells were permeabilized by incubation for 30 minutes with 250 units of lysozyme (ReadyLyse, LGC
BioSearch) in 100ul PBS. Cells were centrifuged again at 6000 rcf and washed with 300ul 1x PBS with
0.1% tween solution. Cells were again centrifuged and washed in PBS to remove the detergent and
resuspended in Probe Binding Buffer (100 µl of probe binding buffer consisting of 5 × SSC, 30%
formamide, 9 mM citric acid at pH 6.0, 0.1% Tween 20, 50 ug/ml heparin and 10% low molecular

weight dextran sulfate). Cells were incubated for 30 minutes at 50°C before 25 ul of probes were added at a concentration of 1,200 ng/ul, and the mixture was left to incubate overnight at 50°C with shaking at 300 RPM. Following incubation cells were washed 7 times in probe-wash solution at 50°C ($5\,×\,$SSC, 30% formamide, $9\,$mM citric acid pH 6.0, 0.1% Tween 20 and $50\,$ug/ml heparin) and finally washed twice in room temperature PBS.

Single-cell microfluidic encapsulation was done using a 10X Genomics chip-G and a Chromium Controller with the Chromium Single Cell 3′ Reagents Kit (v3 chemistry) as described by 10X Genomics using a protocol modified to achieve bacterial scRNA-seq described previously [69,70]. For encapsulation a master mix containing the following reagents was prepared: $33\,$μl of 4X ddPCR Multiplex Supermix (BioRad), $4\,$μl of in-drop PCR primer ($10\,$μM), $2.4\,$μl additive A (10X Genomics) and $26.8\,$μl dH$_2$O. Prepared cell samples were diluted to 1,000 cells μl^{-1} and loaded as a total of 16,000 cells per condition to achieve a targeted cell recovery of 10,000 cells. After encapsulation each sample was transferred to an individual PCR tube and cycled 6 times using the following thermocycler program: $94\,$°C for $5\,$min (before cycling), $94\,$°C for $30\,$s followed by $50\,$°C for $30\,$s then $65\,$°C for $30\,$s for 6 cycles and a final hold at $12\,$°C.

Libraries for NGS sequencing were completed using two additional PCR reactions as detailed previously [69,70] and sequenced using an Element AVITI 150 cycle kit using paired end sequencing of 8 bp for the i7 index, 28bp for Read 1 and the remaining cycles for Read 2.

Reads were trimmed and the probe-based UMI was upended in place of the traditional 10X-bead provided UMI using cutadapt in order to allow for true molecular identification, as detailed previously [69,70]. Samples were mapped and aggregated using the 10X cellRanger pipeline. Analysis of single cell libraries was done using the R-based Seurat pipeline [71]. Standard parameters in Seurat were used for data normalization and to identify the variable features. Louvain clustering was used with resolution set to 0.7 using Seurat FindClusters function. We tested a range of resolutions from 0.5 to 1.5 and find that the main clustering features presented in the text remain across this range of clustering parameters. The Seurat function FindMarkers was used and a log-fold change cutoff of 0.5 chosen to highlight highly differential genes in the main-text volcano plots and a log-fold change cutoff of 0.25 for the supplementary volcano plots in the Supplementary Figures. The full output for all differentially expressed genes is shown in the DGE Supplementary Tables.

Quantification and Statistical Analysis

Statistical analyses of MIC data were performed using Python (v3.10) with the scipy.stats package. Means and standard deviations of individual cultures within cohorts were calculated. Student's t-test was used, assuming a normal distribution, to compare the means of relevant groups, treated vs parent and among segments of the sequential antibiotic treatments. Results are reported as mean ± standard deviation with their corresponding p-values. Significance p-value thresholds were noted with asterisks for $p<0.05$ (*), $p<0.01$ (**), $p<0.001$ (***), and $p<0.0001$(****). Minimum duration of killing (MDK) assay data was reported as median ± median absolute deviation.

<u>ACKNOWLEDGEMENTS</u>

This study was supported by the Advanced Research Projects Agency for Health (ARPA-H) (1AYSAX000005-01), NIH grant R01GM125748, UTSW High Risk High Impact Award, and Welch Foundation I-2082-20240404.

Figure 1. Evolution protocol rationale, scheme, and survival outcomes of single antibiotic evolutions. a,
Illustration of the pharmacokinetics in a clinical antibiotic regimen with peak and trough antibiotic concentrations remaining above the minimum inhibitory concentration (MIC). **b,** Laboratory adaptation of clinical parameters for evolution experiments featuring daily antibiotic exposures at peak concentrations with adjustable exposure durations. c**,** Complete evolution protocol workflow. CFU counts are measured before and after antibiotic addition and removal, followed by overnight growth in antibiotic-free media and a 1:10 dilution after overnight growth. d-f**,** Survival trajectories of bacterial cultures evolved in the presence of amikacin, levofloxacin, and cefepime, respectively. Gray lines represent individual replicate cultures, while colored lines represent the mean survival over time for amikacin (blue), levofloxacin (red), cefepime (green).

816
817
818

Figure 2. Phenotypic and genotypic outcomes of evolutions to single antibiotics. a, representative IC_{50} and MIC determination from a dose-response curve. b-d, bar plots of IC_{50} values in antibiotics amikacin, levofloxacin, and cefepime, respectively for parental and evolved cohorts (MG: gray; MG^{AMI}: blue; MG^{CEF}: green; MG^{LEV}: red). Bar height corresponds to the mean IC_{50}, and error bars reflect the standard deviation. Individual IC_{50} values for replicate cultures are shown as dots. Statistical significance (p-value, student's t-test) is denoted by asterisk (* <0.05; **<0.01, *** < 0.001, **** < 0.0001), a convention used in subsequent figures. e, MDK assay for cefepime-evolved cultures (MG^{CEF}) showing the median survivor fraction plotted over time, with error bars representing the median absolute deviation (MAD). Line colors differentiate the parental strain (MG: black) and evolved cultures that are characteristically tolerant (teal), persistent (pink), or of an intermediate phenotype (maroon). f-h, heatmaps of selected mutations and their frequencies among individual cultures: f, MG^{AMI} (blue); g, MG^{LEV} (red); g, MG^{CEF} (green).

Figure 3. Survival and phenotypic outcomes of sequential antibiotic treatment. a, survival trajectories of bacteria
cultures sequentially evolved in the presence of levofloxacin (red), amikacin (blue), and cefepime (green). Gray lines
represent individual cultures, and colored lines indicate the mean survival through the respective antibiotic: levofloxacin
(red), amikacin (blue), cefepime (green). b-d, bar plots of IC_{50} values in antibiotics levofloxacin, amikacin, and cefepime,
respectively, for parental and evolved cohorts (MG: gray; MG^{LEV}: red; $MG^{LEV,AMI}$: blue; $MG^{LEV,AMI,CEF}$: green). Bar height
corresponds to the mean IC_{50}, and error bars reflect the standard deviation. Individual IC_{50} values for replicate cultures are
shown as dots. e, MDK assay for sequentially evolved cultures ($MG^{LEV,AMI,CEF}$) showing the median survivor fraction plotted
over time, with error bars representing the median absolute deviation (MAD). Line colors differentiate the parental strain
(MG: black) and evolved cultures that are characteristically tolerant (teal), persistent (pink), or of a mixed phenotype
(maroon).

Figure 4. Genotypic outcomes of sequential antibiotic evolution. a, mutation trajectories in Culture 4 of *gyrA* (i), *fusA* (ii), *trkH* (iii) and within the convergent clade mutations (iv). b, emergence and cadence of clade mutations within individual cultures. c, heatmap of final frequencies of convergent clade mutations across replicate cultures. d, cartoon depicting the observed impact of single vs. sequential treatment context on divergent and convergent mutation trajectories (arrows) under the selection of levofloxacin (red) amikacin (blue) and cefepime (green). Arrows are noted with mutations.

Figure 5. Phenotypic effects of individual convergent clade mutations.
a, barplot of doubling times of mutant and parent (MG) strains show mean values of six replicates (dots) with error bar indicating the standard deviation. b-c, IC_{50} of individual mutants in each antibiotic, where bar height shows the mean value for three replicates (dots) and error bar indicates standard deviation. d, MDK of mutants with median survival plotted for three replicates and error bar is the median absolute deviation.

Figure 6. a, UMAP two-dimensional representation of the 9 cell clusters identified in the combined dataset of 48,883 cells. b, UMAP showing cells in either the parental culture (top) or evolved cohorts (middle and bottom). c, Stacked bar chart showing the percent community composition of each cell cluster. d,e, Volcano plot of genes expressed in the parent or MG^{Lev,Ami,Cef}-4 (d) and MG^{Lev,Ami,Cef}-7 (e). Genes shared between the two evolved strains are denoted by solid markers. Adjusted P values correspond to two-sided Wilcoxon rank-sum test with Bonferroni correction. **f**. Distribution of the 50S ribosomal subunit *rplJ*. **g**. *hipA* is distributed away from clusters 1 and 2 and predominantly within clusters 9 and 4.

Line numbers: 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75

Supplemental Figure 1. Differences among survival strategies in time kill assay. Illustration of the MDK curves expected for bacteria which are resistant (red), tolerant (teal), persistent (pink), and susceptible (black). Differences include shifts in the minimum duration for killing 99.9% of bacteria ($MDK_{99.9}$) and the biphasic nature of the MDK curve in persistent populations.

Supplemental Figure 2. IC$_{50}$ and MIC values of individual evolved cultures. a, Average IC$_{50}$ values (error bars: standard deviation) for individual cultures of parental strain (MG, gray) and cultures evolved in the presence of amikacin (MG^{AMI}: blue), cefepime (MG^{CEF}: green), and levofloxacin (MG^{LEV}: red). b, Average MIC values (error bars: standard deviation) for individual cultures of parental strain (MG, gray) and cultures evolved in the presence of amikacin (MG^{AMI}: blue), cefepime (MG^{CEF}: green), and levofloxacin (MG^{LEV}: red).

Supplemental Figure 3. Phenotypic and genotypic outcome of cefepime resistance evolution protocol in lab strain. a, Maximum allowable concertation for growth over time in conventional evolution in the presence of cefepime. Plotted line represents the mean concentration and error bars represent the standard deviation. b-d, bar plots of IC_{50} values in antibiotics amikacin, levofloxacin, and cefepime, respectively for parental and cefepime-resistance evolved cohorts (MG: gray; MG^{CEF-R}: purple). Bar height corresponds to the mean IC_{50}, and error bars reflect the standard deviation. Individual IC_{50} values for replicate cultures are shown as dots. e, heatmap of selected mutations and their frequencies among individual cultures of MG^{CEF-R}.

Supplemental Figure 4. Phenotypic and genotypic outcomes of cefepime evolution in clinical strain. a, Survival trajectories of clinical bacterial cultures evolved in the presence cefepime. Gray lines represent individual replicate cultures, and the mean survival is shown in green. b-d, bar plots of IC_{50} values in antibiotics amikacin, levofloxacin, and cefepime, respectively, for unevolved clinical strain (Pb: gray), cefepime-evolved clinical strain using our evolution protocol (Pb^{CEF}: green) and cefepime-evolved clinical strain using the conventional evolution protocol (Pb^{CEF-R}). height corresponds to the mean IC_{50}, and error bars reflect the standard deviation. Individual IC_{50} values for replicate cultures are shown as dots. e, MDK assay for cefepime-evolved clinical cultures in our evolution protocol (Pb^{CEF}) showing the median survivor fraction plotted over time, with error bars representing the median absolute deviation (MAD). Line colors differentiate the parental

strain (Pb: black) and evolved cultures that are characteristically tolerant (teal) and persistent (pink). f, heatmap of mutations and their frequencies among individual cultures of Pb^{CEF} and Pb^{CEF-R}.

Supplemental Figure 5. Extended genotypic outcomes of single antibiotic evolutions. Heatmaps of extended mutations and their frequencies among individual cultures: a, MG^{AMI} (blue); b, MG^{CEF} (green); c, MG^{LEV} (red).

Supplemental Figure 6. Phenotypic and genetic outcome following untreated passage of MG1655. a-c, IC$_{50}$ values of parent and untreated, passages control populations in each antibiotic, amikacin (a), cefepime (b), levofloxacin (c). d, heatmap of mutational changes throughout untreated passages at key timepoints coinciding with sequential antibiotic treatment.

Supplemental Figure 7. IC$_{50}$ and MIC values of sequentially evolved cultures. a, Average IC$_{50}$ values (error bars: standard deviation) for individual cultures of parental strain (MG, gray) and cultures sequentially evolved in the presence of levofloxacin (MG^{LEV}: red), then amikacin (MG^{LEV,AMI}: blue), then cefepime (MG^{LEV,AMI,CEF}: green), and b, Average MIC values (error bars: standard deviation) for individual cultures of parental strain (MG, gray) and cultures sequentially evolved in the presence of levofloxacin (MG^{LEV}: red), then amikacin (MG^{LEV,AM}: blue), then cefepime (MG^{LEV,AMI,CEF}: green).

Supplemental Figure 8. Selected mutation trajectories within each culture. Emergence and fates of selected mutations throughout sequential antibiotic treatment. Columns a-e depict, respectively, mutations of *gyrA*, *rpoB*, *icd*, *fusA*, *trkH*, and those found to evolve convergently.

Supplemental Figure 9. Comprehensive Mutation Frequency Heatmap. All observed mutations among MG^{LEV}, MG^{AMI}, MG^{LEV,AMI}, MG^{LEV,AMI,CEF}, MG^{CEF} cohorts (left to right) and their frequencies (color bar, right).

Supplemental Figure 10. Extended individual mutation characterization. a-c, MIC for individual mutants in each antibiotic, where bar height shows the mean value for three replicates (dots) and error bar indicates standard deviation. d, MDK of mutant strains and evolved cultures with median survival plotted for three replicates and error bar is the median absolute deviation.

Supplemental Figure 11. Volcano plots of genes differentially expressed in cluster 1 (top) and cluster 2 (bottom) compared to all other clusters in the combined dataset. Genes involved in protein translation that are overexpressed in these clusters are colored in red. Fimbriae related genes downregulated in these two clusters are colored black. For a full list of differentially regulated genes in each gene-set and P-values please refer to supplementary table 3.

Supplemental Figure 12. Volcano plots of genes differentially expressed in cluster 4 (top) and cluster 9 (bottom) compared to all other clusters in the combined dataset. Genes involved in purine nucleotide binding that are overexpressed in these clusters are colored in red. Genes in the superpathway of glycolysis, pyruvate dehydrogenase, TCA and Glyoxylate bypass that are overexpressed in these two clusters are colored in purple. For a full list of differentially regulated genes in each gene-set and P-values please refer to supplementary table 3.

Target Gene	Observed Mutation	Modification Made	Oligo nucleotide
gatA	IS5 insertion within gene	added duplicate stop codons at insert site to truncate protein	ATGAACTTTATTTGTTGGCCTTAACAGATAAATGGCTGACGATTATTACTTAGCATGAATCGCCTCACAATGCGGTATTGCAATAGCGTG
glvC	1-nucleotide deletion	deleted nucleotide	CCATGTTGCCCACCACACCAAAGAGATACATTACGGTCGACATTGGGCCGCCAGCACGGCGTGTACCGCAAACAGCAACGGTGAAATGAA
hipA	SNP	alter nucleotide	TACGTGACCGGATCGTTAAACGTTATCATGCCAAATCCAGACAACTGTTTGATTTATTGTCAGAAATAGGGCGAGACAGCGTTGGTGCCG
selB	2-nucleotide insertion	insert nucleotides	CACCACGGTTCGTCACCAAACAGTGGCTCTGCTTTTTGCCAGATGTGGCCTGCTGCTCTTCGCTGAAGCCCGCTTTGTGATCTGGCAGAT
rpoZ	7-nucleotide deletion	delete nucleotides	ACGCTGTAGAGAAAATTGGTAACCGTTTTGACCTGGTACTGGTCGTCGCGCTCGTCAGATGCAGGTAGGCGGAAAGGATCCGCTGGTACC
ftsH	12-nucleotide deletion	delete nucleotides	CAGCATCTGGTTCAGAGTCTGTTCACGTTCATCGTGACCACCGCCACGCTGGCGGCCTACGGCGTCGATTTCATCGATAAAGATGATGCA
fimE	IS5 insertion between promoter and transcription start	full gene deletion	AACTTTATTATCAATAAGTTAAATCAATGATCCTGACGACGGAGACCGCCGTCGTCGACAAGCCCACAATAGTAGTACTCCTAACTGAGA

Supplementary Table 1. Oligonucleotides for recobineering of individual mutants. Detail of target genes, the mutations observed in each, the approach for modification, and the oligonucleotide used in transformation.

Gene	Primer Name	Full-length, Mutant-Specific, Both	Primer Direction	Sequence
glvC	fwd_glvC_full	Both	FWD	CATCTGCAAGAGTTCAGTTTG
glvC	glvC_mt_rv	Mutant-Specific	REV	ACATTACGGTCGACATTGG
glvC	rev_glvC_full	Full-Length	REV	AGTAGAGTTTCACTTCCGCATC
gatA	fwd_gatA_full	Full-Length	FWD	AGGACGGTATATGACTAACCTG
gatA	fwd_gatA_stops	Mutant-Specific	FWD	GAGGCGATTCATGCTAAGTAAT
gatA	rev_gatA_full	Both	REV	GCAAGCGACAATAATCTTG
selB	fwd_selB_full	Full-Length	FWD	GTTATAGCTTGTTGAATGCG
selB	fwd_selB_mut	Mutant-Specific	FWD	AGAGCAGCAGGCCACA
selB	rev_selB_full	Both	REV	GTTGAACCACACTCCTGAT
hipA	fwd_hipA_full	Full-Length	FWD	TCACTTGGATGAACAACCAG
hipA	hipA_mt_fwd	Mutant-Specific	FWD	ATGCCAAATCCAGACAACT
hipA	rev_hipA_full	Both	REV	TTTAATGATGTGCGTCGTC
rpoZ	fwd_rpoZ_full	Full-Length	FWD	GAATGATGACTTCGATACCG
rpoZ	fwd_rpoZ_span	Mutant-Specific	FWD	TACTGGTCGTCGCGCT
rpoZ	rev_rpoZ_full	Both	REV	GACGCTTGATTTGGTCTTC
ftsH	fwd_ftsH_full	Both	FWD	CAGAAACTCGGCGGTAAG
ftsH	rev_ftsH_full	Full-Length	REV	ATGTGAACTTTCAGGATCTGC
ftsH	rev_ftsHm_del	Mutant-Specific	REV	GACCACCGCCACGCT
fimE	fwd_fimE_full	Full-Length	FWD	GACTGATAGCCACATCACTC
fimE	rev_fimE_del	Both	REV	CGTGCTTTCCTCTATGAGTC
p57	pDel_seq_chlor_fwd	Mutant-Specific	FWD	GCGTTACATCCCTGGCTTGTTG

Supplementary Table 2. PCR Primers used for screening and confirming oligorecombineering mutations.
Details of target PCR products, primer names, product type, primer direction, and sequnce.

Sample name	Cell numbers
MG-3 replicate 1	7827
MG-3 replicate 2	10774
MG^{LEV,AMI,CEF}-4 replicate 1	4764
MG^{LEV,AMI,CEF}-4 replicate 2	2764
MG^{LEV,AMI,CEF}-7 replicate 1	9619
MG^{LEV,AMI,CEF}-7 replicate 2	13135
TOTAL CELLS	48883

Supplementary Table 3. scRNA sequencing samples statistics. Samples (top) and clusters (bottom) and the number of cells representing each group

Cluster name	Cell numbers
1	9335
2	7034
3	6008
4	5989
5	5092
6	4967
7	4961
8	3052
9	2445
TOTAL CELLS	48883

References

1. O'Neill, J. (2016). Tackling drug-resistant infections globally: final report and recommendations. Review on Antimicrobial Resistance.

2. Naghavi, M., Vollset, S., Ikuta, K., Swetschinski, L., Gray, A., Wool, E., and Aguilar, G. (2024). Global burden of bacterial antimicrobial resistance 1990-2021: a systematic analysis with forecasts to 2050. Lancet *404*, 1199-1226. 10.1016/s0140-6736(24)01867-1.

3. Park, K.H., Jung, Y.J., Lee, H.J., Kim, H.J., Maeng, C.H., Baek, S.K., Han, J.J., Jeon, W., Kim, D.Y., Lee, Y.M., and Lee, M.S. (2024). Impact of multidrug resistance on outcomes in hematologic cancer patients with bacterial bloodstream infections. Scientific Reports *14*, 15622. 10.1038/s41598-024-66524-w.

4. Munck, C., Gumpert, H.K., Wallin, A.I., Wang, H.H., and Sommer, M.O. (2014). Prediction of resistance development against drug combinations by collateral responses to component drugs. Sci Transl Med *6*, 262ra156. 10.1126/scitranslmed.3009940.

5. Oz, T., Guvenek, A., Yildiz, S., Karaboga, E., Tamer, Y.T., Mumcuyan, N., Ozan, V.B., Senturk, G.H., Cokol, M., Yeh, P., and Toprak, E. (2014). Strength of selection pressure is an important parameter contributing to the complexity of antibiotic resistance evolution. Molecular biology and evolution *31*, 2387-2401. 10.1093/molbev/msu191.

6. Harmand, N., Gallet, R., Martin, G., and Lenormand, T. (2018). Evolution of bacteria specialization along an antibiotic dose gradient. Evol Lett *2*, 221-232. 10.1002/evl3.52.

7. Eyler, R.F., and Shvets, K. (2019). Clinical Pharmacology of Antibiotics. Clin J Am Soc Nephrol *14*, 1080-1090. 10.2215/cjn.08140718.

8. Balaban, N.Q., Helaine, S., Lewis, K., Ackermann, M., Aldridge, B., Andersson, D.I., Brynildsen, M.P., Bumann, D., Camilli, A., Collins, J.J., et al. (2019). Definitions and

guidelines for research on antibiotic persistence. Nature reviews. Microbiology *17*, 441-448. 10.1038/s41579-019-0196-3.

9. Michiels, J.E., Van den Bergh, B., Verstraeten, N., and Michiels, J. (2016). Molecular mechanisms and clinical implications of bacterial persistence. Drug resistance updates : reviews and commentaries in antimicrobial and anticancer chemotherapy *29*, 76-89. 10.1016/j.drup.2016.10.002.

10. Ronneau, S., Hill, P.W., and Helaine, S. (2021). Antibiotic persistence and tolerance: not just one and the same. Current opinion in microbiology *64*, 76-81. 10.1016/j.mib.2021.09.017.

11. Brauner, A., Fridman, O., Gefen, O., and Balaban, N.Q. (2016). Distinguishing between resistance, tolerance and persistence to antibiotic treatment. Nat Rev Microbiol *14*, 320-330. 10.1038/nrmicro.2016.34.

12. Helaine, S., Conlon, B.P., Davis, K.M., and Russell, D.G. (2024). Host stress drives tolerance and persistence: The bane of anti-microbial therapeutics. Cell Host Microbe *32*, 852-862. 10.1016/j.chom.2024.04.019.

13. Alexander, S., Fisher, B.T., Gaur, A.H., Dvorak, C.C., Villa Luna, D., Dang, H., Chen, L., Green, M., Nieder, M.L., Fisher, B., et al. (2018). Effect of Levofloxacin Prophylaxis on Bacteremia in Children With Acute Leukemia or Undergoing Hematopoietic Stem Cell Transplantation: A Randomized Clinical Trial. JAMA *320*, 995-1004. 10.1001/jama.2018.12512.

14. Lehrnbecher, T., Fisher, B.T., Phillips, B., Alexander, S., Ammann, R.A., Beauchemin, M., Carlesse, F., Castagnola, E., Davis, B.L., Dupuis, L.L., et al. (2020). Guideline for Antibacterial Prophylaxis Administration in Pediatric Cancer and Hematopoietic Stem Cell Transplantation. Clin Infect Dis *71*, 226-236. 10.1093/cid/ciz1082.

15. Lehrnbecher, T., Robinson, P., Fisher, B., Alexander, S., Ammann, R.A., Beauchemin, M., Carlesse, F., Groll, A.H., Haeusler, G.M., Santolaya, M., et al. (2017). Guideline for the Management of Fever and Neutropenia in Children With Cancer and Hematopoietic Stem-Cell Transplantation Recipients: 2017 Update. J Clin Oncol *35*, 2082-2094. 10.1200/JCO.2016.71.7017.

16. Kherroubi, L., Bacon, J., and Rahman, K.M. (2024). Navigating fluoroquinolone resistance in Gram-negative bacteria: a comprehensive evaluation. JAC Antimicrob Resist *6*, dlae127. 10.1093/jacamr/dlae127.

17. Serio, A.W., Keepers, T., Andrews, L., and Krause, K.M. (2018). Aminoglycoside Revival: Review of a Historically Important Class of Antimicrobials Undergoing Rejuvenation. EcoSal Plus *8*. 10.1128/ecosalplus.ESP-0002-2018.

18. Rosenberg, C.R., Fang, X., and Allison, K.R. (2020). Potentiating aminoglycoside antibiotics to reduce their toxic side effects. PLoS One *15*, e0237948. 10.1371/journal.pone.0237948.

19. Bush, K., and Bradford, P.A. (2016). β-Lactams and β-Lactamase Inhibitors: An Overview. Cold Spring Harb Perspect Med *6*. 10.1101/cshperspect.a025247.

20. Monogue, M.L., and Nicolau, D.P. (2018). Antibacterial activity of ceftolozane/tazobactam alone and in combination with other antimicrobial agents against MDR Pseudomonas aeruginosa. J Antimicrob Chemother *73*, 942-952. 10.1093/jac/dkx483.

21. Kiser, T.H., Hoody, D.W., Obritsch, M.D., Wegzyn, C.O., Bauling, P.C., and Fish, D.N. (2006). Levofloxacin pharmacokinetics and pharmacodynamics in patients with severe burn injury. Antimicrob Agents Chemother *50*, 1937-1945. 10.1128/aac.01466-05.

22. Cao, G., Zhang, J., Wu, X., Yu, J., Chen, Y., Ye, X., Zhu, D., Zhang, Y., Guo, B., and Shi, Y. (2013). Pharmacokinetics and pharmacodynamics of levofloxacin injection in healthy Chinese volunteers and dosing regimen optimization. J Clin Pharm Ther *38*, 394-400. 10.1111/jcpt.12074.

23. Windels, E.M., Fox, R., Yerramsetty, K., Krouse, K., Wenseleers, T., Swinnen, J., Matthay, P., Verstraete, L., Wilmaerts, D., Van den Bergh, B., and Michiels, J. (2021). Population Bottlenecks Strongly Affect the Evolutionary Dynamics of Antibiotic Persistence. Molecular biology and evolution *38*, 3345-3357. 10.1093/molbev/msab107.

24. Mahrt, N., Tietze, A., Künzel, S., Franzenburg, S., Barbosa, C., Jansen, G., and Schulenburg, H. (2021). Bottleneck size and selection level reproducibly impact evolution of antibiotic resistance. Nat Ecol Evol *5*, 1233-1242. 10.1038/s41559-021-01511-2.

25. Lazar, V., Pal Singh, G., Spohn, R., Nagy, I., Horvath, B., Hrtyan, M., Busa-Fekete, R., Bogos, B., Mehi, O., Csorgo, B., et al. (2013). Bacterial evolution of antibiotic hypersensitivity. Mol Syst Biol *9*, 700.

26. Imamovic, L., and Sommer, M.O. (2013). Use of collateral sensitivity networks to design drug cycling protocols that avoid resistance development. Sci Transl Med *5*, 204ra132. 10.1126/scitranslmed.3006609.

27. Cao, Y., Pan, Y., Huang, H., Jin, X., Levin, E.J., Kloss, B., and Zhou, M. (2013). Gating of the TrkH ion channel by its associated RCK protein TrkA. Nature *496*, 317-322. 10.1038/nature12056.

28. Cao, Y., Jin, X., Huang, H., Derebe, M.G., Levin, E.J., Kabaleeswaran, V., Pan, Y., Punta, M., Love, J., Weng, J., et al. (2011). Crystal structure of a potassium ion transporter, TrkH. Nature *471*, 336-340.

29. Liakopoulos, A., Aulin, L.B.S., Buffoni, M., Fragkiskou, E., van Hasselt, J.G.C., and Rozen, D.E. (2022). Allele-specific collateral and fitness effects determine the dynamics of fluoroquinolone resistance evolution. Proc Natl Acad Sci U S A *119*, e2121768119. 10.1073/pnas.2121768119.

30. Yoshida, H., Bogaki, M., Nakamura, M., and Nakamura, S. (1990). Quinolone resistance-determining region in the DNA gyrase gyrA gene of Escherichia coli. Antimicrob Agents Chemother *34*, 1271-1272. 10.1128/aac.34.6.1271.

31. Nandy, P., Chib, S., and Seshasayee, A. (2020). A Mutant RNA Polymerase Activates the General Stress Response, Enabling Escherichia coli Adaptation to Late Prolonged Stationary Phase. mSphere *5*. 10.1128/mSphere.00092-20.

32. Fraebel, D.T., Mickalide, H., Schnitkey, D., Merritt, J., Kuhlman, T.E., and Kuehn, S. (2017). Environment determines evolutionary trajectory in a constrained phenotypic space. Elife *6*. 10.7554/eLife.24669.

33. Boël, G., Letso, R., Neely, H., Price, W.N., Wong, K.H., Su, M., Luff, J., Valecha, M., Everett, J.K., Acton, T.B., et al. (2016). Codon influence on protein expression in E. coli correlates with mRNA levels. Nature *529*, 358-363. 10.1038/nature16509.

34. Pietsch, F., Bergman, J.M., Brandis, G., Marcusson, L.L., Zorzet, A., Huseby, D.L., and Hughes, D. (2017). Ciprofloxacin selects for RNA polymerase mutations with pleiotropic antibiotic resistance effects. J Antimicrob Chemother *72*, 75-84. 10.1093/jac/dkw364.

35. Fridman, O., Goldberg, A., Ronin, I., Shoresh, N., and Balaban, N.Q. (2014). Optimization of lag time underlies antibiotic tolerance in evolved bacterial populations. Nature *513*, 418-421. 10.1038/nature13469.

36. Adler, M., Anjum, M., Andersson, D.I., and Sandegren, L. (2016). Combinations of mutations in envZ, ftsI, mrdA, acrB and acrR can cause high-level carbapenem resistance in Escherichia coli. J Antimicrob Chemother *71*, 1188-1198. 10.1093/jac/dkv475.

37. Gerken, H., Shetty, D., Kern, B., Kenney, L.J., and Misra, R. (2024). Effects of pleiotropic ompR and envZ alleles of Escherichia coli on envelope stress and antibiotic sensitivity. J Bacteriol *206*, e0017224. 10.1128/jb.00172-24.

38. Zhou, G., Wang, Q., Wang, Y., Wen, X., Peng, H., Peng, R., Shi, Q., Xie, X., and Li, L. (2023). Outer Membrane Porins Contribute to Antimicrobial Resistance in Gram-Negative Bacteria. Microorganisms *11*. 10.3390/microorganisms11071690.

39. Ma, D., Alberti, M., Lynch, C., Nikaido, H., and Hearst, J.E. (1996). The local repressor AcrR plays a modulating role in the regulation of acrAB genes of Escherichia coli by global stress signals. Mol Microbiol *19*, 101-112. 10.1046/j.1365-2958.1996.357881.x.

40. Cohen, S.P., Hächler, H., and Levy, S.B. (1993). Genetic and functional analysis of the multiple antibiotic resistance (mar) locus in Escherichia coli. J Bacteriol *175*, 1484-1492. 10.1128/jb.175.5.1484-1492.1993.

41. Okusu, H., Ma, D., and Nikaido, H. (1996). AcrAB efflux pump plays a major role in the antibiotic resistance phenotype of Escherichia coli multiple-antibiotic-resistance (Mar) mutants. J Bacteriol *178*, 306-308. 10.1128/jb.178.1.306-308.1996.

42. Reizer, J., Michotey, V., Reizer, A., and Saier, M.H., Jr. (1994). Novel phosphotransferase system genes revealed by bacterial genome analysis: unique, putative fructose- and glucoside-specific systems. Protein Sci *3*, 440-450. 10.1002/pro.5560030309.

43. Homma, K., Fukuchi, S., Kawabata, T., Ota, M., and Nishikawa, K. (2002). A systematic investigation identifies a significant number of probable pseudogenes in the Escherichia coli genome. Gene *294*, 25-33. 10.1016/s0378-1119(02)00794-1.

44. Forchhammer, K., Leinfelder, W., and Böck, A. (1989). Identification of a novel translation factor necessary for the incorporation of selenocysteine into protein. Nature *342*, 453-456. 10.1038/342453a0.

45. Abraham, J.M., Freitag, C.S., Clements, J.R., and Eisenstein, B.I. (1985). An invertible element of DNA controls phase variation of type 1 fimbriae of Escherichia coli. Proc Natl Acad Sci U S A *82*, 5724-5727. 10.1073/pnas.82.17.5724.

46. Stentebjerg-Olesen, B., Chakraborty, T., and Klemm, P. (2000). FimE-catalyzed off-to-on inversion of the type 1 fimbrial phase switch and insertion sequence recruitment in an Escherichia coli K-12 fimB strain. FEMS Microbiol Lett *182*, 319-325. 10.1111/j.1574-6968.2000.tb08915.x.

47. Moyed, H.S., and Bertrand, K.P. (1983). hipA, a newly recognized gene of Escherichia coli K-12 that affects frequency of persistence after inhibition of murein synthesis. J Bacteriol *155*, 768-775. 10.1128/jb.155.2.768-775.1983.

48. Tomoyasu, T., Gamer, J., Bukau, B., Kanemori, M., Mori, H., Rutman, A.J., Oppenheim, A.B., Yura, T., Yamanaka, K., Niki, H., and et al. (1995). Escherichia coli FtsH is a membrane-bound, ATP-dependent protease which degrades the heat-shock transcription factor sigma 32. Embo j *14*, 2551-2560. 10.1002/j.1460-2075.1995.tb07253.x.

49. Griffith, K.L., Shah, I.M., and Wolf, R.E., Jr. (2004). Proteolytic degradation of Escherichia coli transcription activators SoxS and MarA as the mechanism for reversing the induction of the superoxide (SoxRS) and multiple antibiotic resistance (Mar) regulons. Mol Microbiol *51*, 1801-1816. 10.1046/j.1365-2958.2003.03952.x.

50. Volpon, L., Young, C.R., Matte, A., and Gehring, K. (2006). NMR structure of the enzyme GatB of the galactitol-specific phosphoenolpyruvate-dependent phosphotransferase system and its interaction with GatA. Protein Sci *15*, 2435-2441. 10.1110/ps.062337406.

51. Yang, J., Peng, Z., Ji, X., Zhang, J., and Du, G. (2022). Galactitol Transport Factor GatA Relieves ATP Supply Restriction to Enhance Acid Tolerance of Escherichia coli in the Two-Stage Fermentation Production of D-Lactate. Fermentation *8*.

52. Saunders, S.H., and Ahmed, A.M. (2024). ORBIT for E. coli: kilobase-scale oligonucleotide recombineering at high throughput and high efficiency. Nucleic Acids Research *52*, e43-e43. 10.1093/nar/gkae227.

53. McNulty, R., Sritharan, D., Pahng, S.H., Meisch, J.P., Liu, S., Brennan, M.A., Saxer, G., Hormoz, S., and Rosenthal, A.Z. (2023). Probe-based bacterial single-cell RNA sequencing predicts toxin regulation. Nature Microbiology *8*, 934-945. 10.1038/s41564-023-01348-4.

54. Wood, W.N., Mohler, K., Rinehart, J., and Ibba, M. (2021). Deacylated tRNA Accumulation Is a Trigger for Bacterial Antibiotic Persistence Independent of the Stringent Response. mBio *12*, 10.1128/mbio.01132-01121. doi:10.1128/mbio.01132-21.

55. Toprak, E., Veres, A., Michel, J.B., Chait, R., Hartl, D.L., and Kishony, R. (2011). Evolutionary paths to antibiotic resistance under dynamically sustained drug selection. Nat Genet *44*, 101-105. 10.1038/ng.1034.

56. Rodrigues, M., Sabaeifard, P., Yildiz, M.S., Lyon, A., Coughlin, L., Ahmed, S., Poulides, N., Toprak, A.C., Behrendt, C., Wang, X., et al. (2024). Susceptible bacteria can survive antibiotic treatment in the mammalian gastrointestinal tract without evolving resistance. Cell Host Microbe *32*, 396-410.e396. 10.1016/j.chom.2024.01.012.

57. Korch, S.B., Henderson, T.A., and Hill, T.M. (2003). Characterization of the hipA7 allele of Escherichia coli and evidence that high persistence is governed by (p)ppGpp synthesis. Mol Microbiol *50*, 1199-1213. 10.1046/j.1365-2958.2003.03779.x.

58. Germain, E., Castro-Roa, D., Zenkin, N., and Gerdes, K. (2013). Molecular mechanism of bacterial persistence by HipA. Mol Cell *52*, 248-254. 10.1016/j.molcel.2013.08.045.

59. Kaspy, I., Rotem, E., Weiss, N., Ronin, I., Balaban, N.Q., and Glaser, G. (2013). HipA-mediated antibiotic persistence via phosphorylation of the glutamyl-tRNA-synthetase. Nat Commun *4*, 3001. 10.1038/ncomms4001.

60. Schumacher, M.A., Balani, P., Min, J., Chinnam, N.B., Hansen, S., Vulić, M., Lewis, K., and Brennan, R.G. (2015). HipBA-promoter structures reveal the basis of heritable multidrug tolerance. Nature *524*, 59-64. 10.1038/nature14662.

61. Semanjski, M., Germain, E., Bratl, K., Kiessling, A., Gerdes, K., and Macek, B. (2018). The kinases HipA and HipA7 phosphorylate different substrate pools in Escherichia coli to promote multidrug tolerance. Sci Signal *11*. 10.1126/scisignal.aat5750.

62. Narberhaus, F., Obrist, M., Führer, F., and Langklotz, S. (2009). Degradation of cytoplasmic substrates by FtsH, a membrane-anchored protease with many talents. Res Microbiol *160*, 652-659. 10.1016/j.resmic.2009.08.011.

63. Schäkermann, M., Langklotz, S., and Narberhaus, F. (2013). FtsH-mediated coordination of lipopolysaccharide biosynthesis in Escherichia coli correlates with the growth rate and the alarmone (p)ppGpp. J Bacteriol *195*, 1912-1919. 10.1128/jb.02134-12.

64. Langklotz, S., Baumann, U., and Narberhaus, F. (2012). Structure and function of the bacterial AAA protease FtsH. Biochim Biophys Acta *1823*, 40-48. 10.1016/j.bbamcr.2011.08.015.

65. Zuo, Y., Wang, Y., and Steitz, T.A. (2013). The mechanism of E. coli RNA polymerase regulation by ppGpp is suggested by the structure of their complex. Mol Cell *50*, 430-436. 10.1016/j.molcel.2013.03.020.

66. Ross, W., Sanchez-Vazquez, P., Chen, A.Y., Lee, J.H., Burgos, H.L., and Gourse, R.L. (2016). ppGpp Binding to a Site at the RNAP-DksA Interface Accounts for Its Dramatic Effects on Transcription Initiation during the Stringent Response. Mol Cell *62*, 811-823. 10.1016/j.molcel.2016.04.029.

67. Deatherage, D.E., and Barrick, J.E. (2014). Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using breseq. Methods in molecular biology *1151*, 165-188. 10.1007/978-1-4939-0554-6_12.

68. Nyerges, Á., Csörgő, B., Nagy, I., Bálint, B., Bihari, P., Lázár, V., Apjok, G., Umenhoffer, K., Bogos, B., Pósfai, G., and Pál, C. (2016). A highly precise and portable genome engineering method allows comparison of mutational effects across bacterial species. Proc Natl Acad Sci U S A *113*, 2502-2507. 10.1073/pnas.1520040113.

69. Samanta, P., Cooke, S.F., McNulty, R., Hormoz, S., and Rosenthal, A. (2024). ProBac-seq, a bacterial single-cell RNA sequencing methodology using droplet microfluidics and large oligonucleotide probe sets. Nature protocols *19*, 2939-2966. 10.1038/s41596-024-01002-1.

70. McNulty, R., Sritharan, D., Pahng, S.H., Meisch, J.P., Liu, S., Brennan, M.A., Saxer, G., Hormoz, S., and Rosenthal, A.Z. (2023). Probe-based bacterial single-cell RNA sequencing predicts toxin regulation. Nat Microbiol *8*, 934-945. 10.1038/s41564-023-01348-4.

71. Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y., Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell Data. Cell *177*, 1888-1902 e1821. 10.1016/j.cell.2019.05.031.