Multiple cis-regulatory elements collaborate to control *mdka*
expression in telencephalic neural stem of adult zebrafish during
constitutive and regenerative neurogenesis

Jincan Chen[1], Masanari Takamiya[1], Agnes Hendriks[1], Tanja Beil[1], Csilla Varnai[2]
Nicolas Diotel[3] and Sepand Rastegar[1,*]

1 Institute of Biological and Chemical Systems-Biological Information Processing
(IBCS-BIP), Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany
2 Department of Cancer and Genomics Sciences, University of Birmingham,
Birmingham, United Kingdom
3 Université de La Réunion, INSERM, UMR 1188, Diabète athérothrombose
Thérapies Réunion Océan Indien (DéTROI), Saint-Pierre, La Réunion, France

*Correspondence to:
S. Rastegar, Institute of Biological and Chemical Systems-Biological Information
Processing (IBCS-BIP), Karlsruhe Institute of Technology, Karlsruhe 76344,
Germany. Tel: +49 721 608 22507. E-mail: sepand.rastegar@kit.edu.

Running title: Transcriptional regulation of *mdka* in the nervous system

17
18 **Keywords:** Neurogenesis; Radial glial cells; Regeneration; Zebrafish; *mdka*; Cis-
19 regulatory elements; Gene transcription regulation
20

Abbreviations: CRE: cis-regulatory element; dpf: days post-fertilization; Dm:
dorsomedial of the telencephalon; dpl: days post-lesion; EGFP: enhanced green
fluorescent protein; FISH: fluorescent *in situ* hybridization; hpf: hours post-fertilization;
IHC: immunohistochemistry; kb: kilo bases; PCNA: proliferating cell nuclear antigen;
RGCs: radial glial cells; mRNA: messenger RNA; *mdka*: *midkine a*; NSCs: neural stem
cells; S100β: S100 calcium-binding protein β; *Tg*: transgenic line; TSS: transcription
start site; VZ: ventricular zone;

Abstract

Zebrafish is a powerful animal model for studying nervous system regeneration due to its remarkable regenerative abilities and the availability of diverse molecular tools. After telencephalic brain injury, neural stem cells (NSCs) in the ventricular zone (VZ) become activated, proliferate, and generate new neurons essential for brain repair. However, the molecular mechanisms regulating these processes remain unclear. Here, we investigate the transcriptional regulation of *midkine-a* (*mdka*), a heparin-binding growth factor gene encoding the secreted protein Midkine-a (Mdka), which is upregulated after injury in radial glial cells (RGCs), the bona fide NSCs of the adult zebrafish telencephalon. Using genome-wide bioinformatic analysis, we identified six putative cis-regulatory elements (CREs) associated with *mdka*. Transgenic assays revealed that these CREs coordinate *mdka* expression during both development and regeneration. In the zebrafish embryo, CRE2, CRE3, CRE4, and CRE6 are required for EGFP expression in the nervous system, with CRE3 showing the strongest activity. In the adult telencephalon, CRE2, CRE4, and CRE6 are active in NSCs, with CRE2 best mimicking *mdka* expression at the ventricular zone. Importantly, individual CREs could not fully reproduce endogenous *mdka* expression, especially under regenerative conditions. In contrast, a combined CRE2346 construct closely recapitulated *mdka* expression in both the embryo and adult telencephalon under homeostatic conditions. These results suggest that *mdka* expression is controlled by a modular and cooperative cis-regulatory architecture that enables precise gene regulation during development, telencephalon homeostasis, and regeneration.

Introduction

Understanding the molecular mechanisms underlying tissue and organ regeneration is critical for addressing the growing challenges posed by neurodegenerative diseases, such as dementia, Parkinson's, and Alzheimer's. These diseases place a significant medical and societal burden on an aging population, underscoring the urgent need for effective therapeutic strategies to address these conditions. Zebrafish has emerged as a key vertebrate model for studying regeneration due to its exceptional ability to regenerate tissues throughout both embryonic and adult stages [1-6]. Unlike most mammals, zebrafish can regenerate organs like the heart, fin, retina, spinal cord, and specific brain regions without scarring and obvious disabilities, which makes them invaluable for studying regenerative processes [2]. With over 70% genetic similarity to humans [7] and a vast number of mutants [8], transgenic lines [9-11], and genomic resources such as the DANIO-CODE Data Co-Ordination Center (DCC) (https://danio-code.zfin.org/), zebrafish provide a unique platform to study gene function, regulation, and cellular dynamics during embryonic development, adulthood under homeostatic and regenerative conditions [12-14]. In zebrafish, the adult telencephalon regenerates by activating dormant RGCs in the ventricular zone (VZ), which produce new neurons to replace those lost from injury [15, 16]. Recent transcriptomic studies have identified several genes significantly upregulated following injury [17-20], one of the most notable being *midkine-a* (*mdka*), a heparin-binding growth factor with multiple roles in development, repair and diseases [21-23]. Interestingly, *mdka* expression increases in quiescent RGCs after telencephalic injury, resembling the behaviour of *id1* (*inhibitor of differentiation 1*) [21], a transcriptional regulator downstream of the BMP signalling pathway [24, 25]. Both *mdka* and *id1* are predominantly expressed in quiescent type 1 cells (RGCs) and are absent in type 2 cells corresponding to actively dividing cells [21]. Both type 1 and type 2 cells express glial markers such as S100β and GFAP, while type 2 cells in addition express the proliferation marker PCNA (proliferating cell nuclear antigen). Lineage tracing experiments have demonstrated that type 2 cells differentiate into type 3 neuroblasts, which migrate and mature into specific neuronal types [26, 27].

Beyond the brain, *mdka* has been implicated in the regeneration of other tissues in zebrafish and other vertebrates, including the heart, fin, and retina, suggesting that it

may play a common role in the regenerative process [28-35]. The zebrafish *midkine* family includes two genes: *mdka* and its paralog *midkine b* (*mdkb*), the result of a gene duplication event [36]. However, only *mdka* is significantly upregulated after injury, highlighting its specific role in regenerative responses [21]. This study investigates the transcriptional regulation of *mdka* in the zebrafish telencephalon, focusing on its regulation in NSCs during neurogenesis and regeneration. We aim to uncover the molecular mechanisms behind its transcription by identifying and characterizing cis-regulatory elements (CREs) that control *mdka* expression in both homeostatic and regenerative contexts.

Our findings, using genome-wide bioinformatic approaches and validation by transgenesis, reveal that multiple CREs work together to regulate *mdka* expression. In embryos, CRE2, CRE3, CRE4, and CRE6 are critical for nervous system expression, while in the adult telencephalon, CRE2, CRE4, and CRE6 are essential for expression in NSCs. Notably, none of the CREs alone were sufficient to fully recapitulate *mdka* expression in the embryonic nervous system or adult telencephalon. These findings highlight the complexity of *mdka* regulation, where individual CREs contribute differently to expression patterns depending on molecular and environmental signals. Furthermore, a transgenic EGFP reporter line containing CRE2, CRE3, CRE4, and CRE6 best mimicked endogenous *mdka* expression in both zebrafish embryos and the adult telencephalon under homeostatic and regenerative conditions, emphasizing the role of multiple molecular cues in *mdka* regulation.

Results

Identification and characterization of *mdka* CREs in the zebrafish embryo

To investigate the transcriptional regulation of *mdka*, we conducted a systematic search for CREs controlling *mdka* expression in zebrafish embryos at 24 hours post-fertilization (hpf) and in the ventricular zone of the adult telencephalon under both physiological and injury-induced conditions. Using data from the DANIO-CODE repository [14, 37], we identified six potential CREs located upstream, downstream, or in intergenic regions of the *mdka* coding sequence, as shown in **Figure 1**. These regions were selected based on chromatin accessibility, as assessed by ATAC-seq (Assay for Transposase-Accessible Chromatin with sequencing) across zebrafish developmental stages, from dome to long-pec. We also considered sequence conservation with the carp genome [38], indicated by blue boxes. Based on epigenetic features provided in the DANIO-CODE resource, including DNA methylation and histone modifications, the potential CREs were classified as promoters or enhancers and categorized by their activation state (repressed, quiescent, or activated) and strength (weak or strong). To validate their regulatory activity *in vivo*, each CRE was cloned upstream of a 1.0-kb *gata2* minimal promoter, corresponding to the most truncated basal promoter (P5) described by Meng et al. (1997) [39]. This promoter has been subsequently validated for enhancer assays by Navratilova et al. (2009) and Zhang et al. (2020) [25, 40], and was used here to drive EGFP expression (**Figure 2A**). These constructs were introduced into zebrafish germline cell to generate stable transgenic lines, *Tg(CREx-gata2aPR:EGFP)*, where CREx represents one of the identified CREs (**Figure 2A**). Five of the six CREs (CRE2-6) drove expression in 24 hpf zebrafish embryos, displaying partially overlapping patterns with endogenous *mdka* expression (**Figure 2B-G; B'-G'; Figure S1A, B**). Among them, *Tg(CRE3-gata2aPR:EGFP)* most closely recapitulated *mdka* expression (**Figure 2D, D'** and **Table 1**). In this line, EGFP was detected along the spinal cord, tail/fin fold, and brain (**Figure 2D, D'; Table 1; Figure S1A** shows a representative 24 hpf zebrafish embryo with specific anatomical regions). However, none of the individual CREs fully replicated the endogenous *mdka* expression pattern (**Figure 2; Table 1**). No EGFP expression was detected in the stable lines generated with CRE1 (**Figure 2B, B'**). Several explanations could account for this, including the possibility that CRE1

functions as a silencer. This hypothesis will be briefly discussed in the relevant section.

Additionally, each CRE was tested in combination with three different promoters: the

ctgf promoter [41], the zebrafish synthetic promoter (zsp) [42], and the *mdka* intrinsic

promoter (mdkaPR, this study). EGFP expression in these transient transgenics

closely resembled that observed using the *gata2a* promoter (**Figure S2**), supporting

the conclusion that EGFP expression is primarily driven by the CREs rather than by

the specific promoter used. Given the consistent expression patterns observed with

the *gata2a* promoter, all subsequent characterization of the CREs was carried out

using the *gata2aPR:EGFP* construct.

Characterization of CREs driving *mdka* expression in adult NSCs

In the adult zebrafish telencephalon, EGFP expression was detected in four out of six

transgenic lines generated (**Figure 3C, C'; D, D'; E, E'; G, G'**). No expression was

observed for CRE1 and CRE5 (**Figure 3B, B'; F, F'**). CRE3 drove strong ectopic

EGFP expression in the dorsal pallium (Dp) and lateral pallium (Dl), whereas no

expression was detected in the dorsomedial region (Dm) of the VZ (**Figure 3D, D'**). In

contrast, CRE2, CRE4, and CRE6 directed EGFP expression in the VZ of the

telencephalon, where RGCs reside (**Figure 3C, C'; E, E'; G, G'**, and **Table 2**). To

determine if EGFP transgene expression accurately reflects endogenous *mdka*

expression, we compared it with endogenous *mdka* expression at single-cell resolution

(**Figure 3H-J**). In CRE2, CRE4 and CRE6 transgenic lines, EGFP-positive cells were

present along the VZ with endogenous *mdka*-positive RGCs (**Figure 3H-J''**).

Quantification of confocal stack images showed that roughly half of *mdka*-positive

RGCs co-expressed EGFP in each CRE transgenic lines, with no significant difference

among the three CRE lines (**Figure 3K**). In the CRE6 transgenic line, EGFP

expression was also observed outside of VZ, where endogenous *mdka* is not

expressed, suggesting that CRE6 activity leads to ectopic transgene expression

(**Figure 3J''**, white arrows). Since we previously reported that *mdka* expression is

predominant in quiescent RGCs compared to activated RGCs [21], we next

exanimated whether CRE2, CRE4 and CRE6 drive EGFP expression in similar

manner. To investigate this, we analyzed the expression of the RGC marker S100β

and the proliferation marker PCNA in EGFP-positive cells of each CRE transgenic line

(**Figure 3L**). All CRE transgenic lines exhibited robust EGFP expression, primarily in

quiescent type 1 RGCs marked by S100β+/PCNA−, closely resembling the

endogenous *mdka* expression (**Figure 3M; Figure S1C; Figure S3A-A''', B-B'''**).

Although each CRE captured specific aspects of endogenous *mdka* expression, none

of them alone fully recapitulated its complete spatial pattern in the adult telencephalon.

A similar discrepancy was observed during embryonic development at 24 hpf (**Figure**

2). While *Tg(CRE2-gata2aPR:EGFP)*, *Tg(CRE4-gata2aPR:EGFP)*, and *Tg(CRE6-*

gata2aPR:EGFP) each reflect certain aspects of *mdka* expression, their activity was

incomplete in both spatial distribution and intensity. Interestingly, CREs that were

strongly active in embryonic central nervous system (**Figure 2C**, **C'**) also exhibited

activity in the adult telencephalon (**Figure 3C, E and G**). However, their expression

patterns differed from that of the endogenous *mdka* mRNA, suggesting that distinct

regulatory elements control *mdka* expression at different developmental stages.

These findings indicate that full *mdka* expression likely requires a combination of

multiple CREs, and that these combinations may vary between embryonic and adult

stages, particularly in the adult telencephalon.

Identification of injury-responsive CREs regulating *mdka* expression in the adult

telencephalon

Since *mdka* expression is significantly upregulated during telencephalon regeneration

following brain injury, particularly in type 1 RGCs [21], we subjected all six *mdka* CRE

transgenic lines to telencephalon injury (**Figure 4B-G, B'-G'**). All injury-related

analyses were performed at 5 days post-lesion (dpl), based on prior RNA-sequencing

and RT-qPCR data showing that *mdka* expression peaks at this time point in the

injured hemisphere, with a 1.37-fold increase compared to the uninjured side ($p = 2.68$

$\times 10^{-9}$) [21]. This time point therefore provided an optimal window to evaluate injury-

induced regulatory activity of the CREs. The injury was induced in the right hemisphere

using a 30G syringe needle inserted through the skull [27, 43], while the uninjured

contralateral hemisphere served as an internal control (**Figure 4A**). Following injury,

EGFP reporter expression remained absent in the VZ of the Dm region in transgenic

lines carrying CRE1, CRE3, and CRE5 (**Figure 4B, B', D, D', F,** and **F'**), consistent

with the lack of expression observed under homeostatic conditions (**Figure 3B, B', D,**

D', F, and **F'**). In contrast, transgenic lines with CRE2 (**Figure 4C, C'**), CRE4 (**Figure**

4E, E'), and CRE6 (**Figure 4G, G'**) showed EGFP expression in the Dm of the VZ.

However, for all three CREs, the number of EGFP-positive RGCs in the injured
hemisphere remained largely unchanged compared to the uninjured contralateral side,
with no significant differences between the CREs (**Figure 4H**). Assessment of RGC
stemness, based on the proportion of type 1 RGCs among EGFP-positive cells,
revealed differences between CREs in the injured (ipsilateral) brain hemisphere.
Specifically, CRE2 showed a significantly higher proportion of quiescent type 1 RGCs
compared to CRE6 (**Figure 4I**, $p < 0.01$). In contrast, on the contralateral (uninjured)
side, approximately three-quarters of EGFP-positive RGCs remained quiescent
across all CRE lines, consistent with the proportions observed under homeostatic
conditions (**Figure 3M; Figure 4I**). These results suggest that none of the individual
CREs contain a strongly injury-responsive regulatory element sufficient to upregulate
mdka expression during regeneration. Similar to the regulation observed under
physiological conditions, it is likely that multiple CREs must act in combination to
mediate a full regenerative response.

Cooperative CRE activity enhances faithful transgene expression in embryos,
adult RGCs, and in response to injury

Analysis of individual stable transgenic lines showed that none of the six CREs alone
fully recapitulated the spatial expression pattern of *mdka* throughout embryonic
development and adulthood, under either homeostatic or regenerative conditions. To
investigate potential cooperative interactions among these CREs, we generated a
combined CRE2346 transgenic reporter construct (**Figure 5A**). This construct
integrates CRE3, which showed strong regulatory activity during embryogenesis,
together with CRE2, CRE4, and CRE6, which were effective in the adult
telencephalon. Under homeostatic conditions, the CRE2346 construct drove EGFP
expression patterns that closely mimicked endogenous *mdka* expression in embryos
(**Figure 5B-B''**). In the adult telencephalon (**Figure 5C, D-D''**), CRE2346 mediated
specific EGFP expression in the Dm of the VZ (white arrowheads, **Figure 5D'**),
displaying RGC morphology and co-expression with the RGC marker S100β.
Transgene fidelity assessment revealed that CRE2346 was active in the majority
(median 88%) of *mdka*-positive RGCs, demonstrating significantly higher coverage
than any individual CRE (**Figure 5E**). However, analysis of RGC stemness, assessed

by immunolabeling for PCNA and S100β, showed no significant effect of the CRE
combination (**Figure 5F-F''', G**).

We next investigated how the combination of *mdka* cis-regulatory elements (CREs)
influences transgene expression during regenerative neurogenesis following brain
injury (**Figure 6**). After stab injury, EGFP expression driven by the CRE2346 construct
was observed in the ventricular zone (VZ) of both the injured (ipsilateral) and uninjured
(contralateral) hemispheres, closely mirroring the endogenous *mdka* expression
pattern (**Figure 6A, B-B''**). Notably, injury-induced *mdka* upregulation at the lesion
site, which occurs outside the VZ, was also faithfully reproduced by the CRE2346 line
(**Figure 6C-C''**). Quantification revealed that the number of radial glial cells (RGCs)
expressing EGFP in the injured hemisphere was comparable to that in the
contralateral side across all CRE lines, further reflecting endogenous *mdka* expression
(**Figure 6D**). Moreover, EGFP intensity was significantly increased in the injured
hemisphere in the CRE2346 line ($p < 0.001$), aligning with our prior observations of
injury-induced *mdka* upregulation (**Figure 6E**) [21]. To evaluate the impact of
combining CREs, we compared transgene fidelity and ectopic expression between
CRE2346 and single CRE lines (**Figure 6F-G**). The combination significantly improved
transgene fidelity, with consistent expression levels between ipsilateral and
contralateral hemispheres (**Figure 6F**). Additionally, CRE2346 effectively suppressed
the ectopic, off-target expression seen with CRE4 and CRE6 alone, which had
previously shown high levels of non-specific activity (**Figure 6G**). We then assessed
whether the CRE combination preserved the identity and stemness of transgene-
expressing RGCs. Co-immunolabeling with S100β and PCNA allowed us to
distinguish quiescent (type 1) from activated (type 2) RGCs (**Figure 6H-I'''**). In the
uninjured (contralateral) hemisphere, CRE2346 increased the proportion of EGFP+
quiescent RGCs to 84%, closely approaching the 94% observed for endogenous *mdka*
expression (**Figure 6J**). However, in the injured hemisphere, this fidelity was reduced.
All constructs, including CRE2346, showed a marked decrease in the proportion of
quiescent RGCs (39-69%), significantly lower than the 90% observed with
endogenous *mdka*. This suggests that although the CRE2346 line closely replicates
endogenous *mdka* activity under normal conditions and partially during regeneration,
it likely lacks additional regulatory elements necessary for fully recapitulating *mdka*'s
injury response, particularly for maintaining the quiescent RGC state.

In summary, our findings demonstrate that the CRE2346 combination effectively
drives *mdka*-like EGFP expression in RGCs with high spatial specificity and
responsiveness to injury (**Table 3**). This highlights the necessity of multiple regulatory
elements for precise spatial and temporal control of *mdka* expression under both
physiological and regenerative conditions. Interestingly, all four CREs contain multiple
transcription factor binding sites, including robust AP-1 binding sites within their
evolutionarily conserved regions (**Figure S5**). AP-1 is a well-known regulator of gene
expression during regeneration across various tissues and species, including the
zebrafish heart, fin, and tail [35, 44-48], supporting the potential role of these CREs in
the regenerative response to injury.

Discussion

Our study reveals that *mdka* expression in the zebrafish central nervous system is regulated by a complex and highly context-dependent network of CREs. These elements act in a modular, combinatorial, and sometimes redundant fashion, reflecting a multifaceted regulatory architecture that governs *mdka* transcription across different developmental stages and physiological states. Consistent with previous reports, *mdka* displays dynamic, tissue-specific expression patterns. For example, in the adult zebrafish telencephalon, *mdka* is expressed in RGCs under homeostatic conditions [21], while in the retina, it is absent in resting stem cells but becomes induced in proliferating Müller glia following injury [29, 49]. Interestingly, in the injured telencephalon, *mdka* expression remains restricted to quiescent RGCs, underscoring its tight, tissue-specific transcriptional control [21].

Modular and context-specific regulation of *mdka*

Our transgenic analyses demonstrate that *mdka* expression is orchestrated by multiple CREs, each contributing distinct spatiotemporal aspects to its regulatory program. Some elements, such as CRE5 and CRE3, are active primarily during embryogenesis. For example, CRE3 drives strong expression in the embryonic brain and spinal cord but fails to recapitulate specific adult brain patterns, instead causing ectopic and nonspecific activation (**Table 3**). In contrast, elements like CRE2 are active mainly in the adult telencephalon, particularly in RGCs located in the Dm of the VZ. This modularity is not limited to the brain. In the heart, previous studies showed that a transgene containing *mdka* intronic sequences (*mdka*_e2, overlapping our CRE3, **Table S1**) was active in the developing epicardium but inactive after adult injury [35]. Meanwhile, a downstream element (*mdka*_e2, overlapping CRE6, **Table S1**) was inactive during development but became induced in a subset of adult epicardial cells following injury [35]. These examples reinforce the idea that distinct CREs mediate *mdka* expression in different developmental or physiological contexts.

Combinatorial input cooperative and redundant CRE activity

While individual CREs contribute to specific features of *mdka* expression, none alone were sufficient to fully recapitulate endogenous expression, particularly in response to

injury. For instance, CRE2, CRE4, and CRE6 each drove weak expression in adult
telencephalic RGCs, but none showed injury responsiveness on their own. However,
when combined in a multi-CREs construct, these same elements collectively drove
expression that closely mimicked endogenous *mdka*, including robust injury-induced
activation. This supports a model in which full *mdka* expression depends on the
cooperative action of multiple CREs. Such combinatorial regulation may reflect the
integration of distinct signaling inputs across separate regulatory elements. One
possibility is that different signaling pathways act through specific CREs, with their
additive or synergistic activity required to achieve the proper spatiotemporal
expression profile. In this context, the effects of different CREs may be additive
meaning that multiple enhancers with similar expression patterns together drive
expression to levels that individual elements cannot reach on their own. Alternatively,
or additionally, a single signaling pathway might act on multiple CREs in parallel, with
full activation only occurring when all responsive elements are engaged. This
combinatorial strategy may also contribute to enhanced specificity by limiting enhancer
activity to overlapping spatial domains, thereby reducing ectopic expression. The
presence of conserved AP-1 binding motifs within CRE2, CRE3, CRE4, and CRE6
(**Figure S5A-D**), coupled with the established role of the AP-1 complex in regeneration
[34, 44, 47], supports both scenarios. Importantly, our observations also reveal
redundancy among CREs. Although each element has specific functions, some show
overlapping activity. For instance, despite weak individual responses to injury, CRE2,
CRE4, and CRE6 all target the same VZ region. Furthermore, CRISPR/Cas9-
mediated deletion of a regulatory element in the regenerating fin did not abolish *mdka*
expression, and regeneration proceeded without detectable patterning defects [34].
This suggests that redundant or compensatory enhancer activity may preserve gene
expression when individual elements are disrupted. Such redundancy adds
robustness to the regulatory system, ensuring consistent *mdka* activation even in
fluctuating or damaged environments. Together, our results highlight a regulatory logic
in which enhancer modularity, cooperation, and redundancy work in concert to control
mdka expression with high precision and resilience.

Broader implications for gene regulation and regenerative biology

The modular and distributed regulatory architecture of *mdka* is consistent with classical models of enhancer function. The *even-skipped (eve)* gene in *Drosophila*, for example, is expressed in a seven-stripe pattern during embryogenesis, with each stripe controlled by a separate enhancer integrating local signals [50, 51]. Similarly, in mammals, the neural crest gene *sox10* is regulated by multiple enhancers with overlapping and redundant functions, enabling robust expression across tissues and stages [52, 53]. These cases, like *mdka*, illustrate how enhancer modularity enables nuanced, context-dependent expression and buffering against perturbation. Our *mdka* reporter lines provide a useful system for dissecting transcriptional programs that govern adult neurogenesis and injury-induced gene activation. These tools could be employed to identify upstream regulators of regenerative gene expression and screen for factors that modulate regenerative capacity *in vivo*.

Regulatory complexity vs. simplicity: A comparative perspective

Intriguingly, the complexity of *mdka* regulation contrasts with the simpler enhancer architecture observed for other genes expressed in the same cellular context. For example, *id1*, which is co-expressed with *mdka* in quiescent RGCs under both physiological and regenerative conditions, appears to be controlled by a single enhancer responsive primarily to BMP signaling [24, 25]. Despite their overlapping expression domains, *id1* and *mdka* rely on very different regulatory strategies. This contrast raises important questions: Why do some genes require complex enhancer networks while others are regulated through simpler mechanisms? Comparative studies between genes like *mdka* and *id1* could illuminate how enhancer complexity evolves and functions in the context of regeneration and plasticity.

Limitations and future directions in defining the *mdka* regulatory landscape

Our primary goal was to identify and functionally validate CREs that positively regulate *mdka* expression. Using *in vivo* transgenesis in zebrafish embryos and adults, we discovered several elements that drive EGFP expression in patterns consistent with endogenous *mdka* activity. These findings provide a foundation for understanding how *mdka* is regulated during development and regeneration. However, a key limitation of our approach is that it is optimized to detect enhancer activity and does not capture negative regulatory elements such as silencers. Traditional reporter assays are unable

to distinguish between a lack of enhancer function and active repression. For instance,

CRE1 did not drive detectable EGFP expression on its own, but whether this indicates

silencer activity or simply absence of regulatory function remains unresolved. To

address this, future studies should explore combinatorial testing, such as integrating

CRE1 with active enhancers like CRE2346, to determine if it can restrict or suppress

ectopic expression. Such designs could uncover context-dependent repression that

would remain hidden when elements are tested in isolation. Our data also underscore

the modular nature of the *mdka* regulatory architecture, which involves synergy,

additivity, and redundancy among CREs. Additive effects, where multiple elements

with overlapping activity are needed to achieve full expression, should be

distinguished from redundancy, where one CRE can compensate for the absence of

another. Both features likely contribute to the robustness and precision of *mdka*

expression in both physiological and regenerative contexts. A fuller understanding of

this regulatory logic will require experimental strategies that can reveal both positive

and negative regulatory interactions.

Another limitation relates to the use of fluorescent reporter systems. The relative

stability of EGFP protein may result in prolonged signal even after *mdka* mRNA

expression has subsided, potentially contributing to discrepancies in spatial

localization. To address this, we performed *in situ* hybridization for *egfp* mRNA, which

confirmed both expected expression in the dorsomedial region and ectopic expression

in the parenchyma (data not shown). These findings further support the idea that

certain silencer elements may be absent from our current constructs. Given its lack of

activity in isolation, CRE1 remains a strong candidate for a negative regulatory

element, and its function in combination with other enhancers will be a priority for future

studies. In addition, mRNA-level analysis across multiple transgenic lines will be

essential for identifying those that most accurately recapitulate endogenous *mdka*

expression in radial glial cells. Collectively, these insights highlight the importance of

both enhancer and silencer elements in shaping the spatial and temporal dynamics of

mdka regulation under homeostatic and regenerative conditions.

Materials and methods

Zebrafish husbandry

All zebrafish lines were maintained at the European Zebrafish Resource Center (EZRC). Experiments were performed on 6- to 12-month-old wild-type fish or transgenic reporter lines (**Table S2**). Animal husbandry was carried out in accordance with permit guidelines under §11 TSchG (Regierungspräsidium Karlsruhe, Aktenzeichen 35–9185.64/BH KIT). Fish were kept according to the European Society for Fish Models in Biology and Medicine (EuFishBioMed) guidelines at the Karlsruhe Institute of Technology (KIT) [54], and animal experimentation adhered to German animal protection standards, approved by the Government of Baden-Württemberg (Regierungspräsidium Karlsruhe, Reference Numbers: 35-9185.81/G-214/21 and 35-9185.81/G-215/21). Efforts were made to minimize animal distress and reduce their number.

Plasmids construction

Plasmids for enhancer/promoter reporter assays were generated using Gateway cloning and TOPO/TA cloning strategies. The primers used for PCR amplification of putative CREs, and promoter are listed in **Table S3,** while the coordinates and sizes of the CREs and the promoter are provided in **Table S1**. Most plasmids (**Table S4**) were constructed using commonly available vectors, with detailed protocols for Gateway cloning available as described in Kwan et al., 2007 [55]. The pTol2-mdkaCRE2346-gata2aPR:egfp-pA plasmid was created by combining four *mdka* putative CREs (CRE2, CRE3, CRE4, and CRE6) into the pT2KHGPzGATA2C1 destination vector and was synthesized by GenScript Biotech Corporation.

Microinjection to generate transient and stable transgenesis

One-cell stage embryos were injected with 1 µl of 20 ng/µl Tol2 transposase mRNA, 1 µl of 10 % phenol red, 50 ng/µl plasmid DNA, and nuclease-free water (final volume 10 µl) directly into the yolk through the chorion. After injection, embryos were incubated at 28 °C in E3 medium with methylene blue. At 24 hpf, embryos were sorted based on EGFP fluorescence. Positive F0 founders were identified by EGFP

426 expression, raised to sexual maturity, and outcrossed with wildtype zebrafish. Stable
427 EGFP expression in ~20 % of F1 progeny confirmed successful stable transgenesis.

Fluorescent *in situ* hybridization (FISH) of whole-mount embryos

429 Embryos were collected at the desired developmental stage, and chorions were
430 removed manually or with pronase (1:100 w/v, 28.5 °C). The *mdka* probe (Molecular
431 Instruments Inc.) was used, following the HCR™ RNA-FISH protocol with minor
432 modifications, available on the Molecular Instruments website.

Stab injury of adult telencephalon and preparation of sections

434 The stab injury experiment and telencephalon sections were performed following the
435 protocol described by Schmidt et al. [43].

FISH on adult telencephalon sections and embryos

For experiments using probes from Molecular Instruments Inc., the protocol "HCR™
RNA-FISH protocol for fresh frozen or fixed frozen tissue sections was followed, with
minor modifications. Specifically, for 24 hpf embryos, proteinase K treatment was
conducted at a concentration of 10 µg/ml (diluted from a 1000 × stock to 1 × working
concentration) for 10 min to improve probe penetration. The *mdka* probe used for
fluorescent *in situ* hybridization was purchased from the Molecular Instruments Inc.
with a dilution of 1:500.

Immunohistochemistry (IHC) on adult telencephalon sections

Vibratome sections were blocked with 1 ml blocking buffer for 1 hour at RT. After
removing the buffer, sections were incubated overnight at 4 °C with primary antibodies
in 300 µl blocking buffer. The next day, sections were washed 3 × 10 min with PTW (1
× PBS containing 0.1% Tween-20), then incubated with secondary antibodies (diluted
in PTW) for 2 hours at RT in the dark. DAPI staining (1:2000 in PTW) was performed
for 20 min if needed, followed by 3 × 5 min washes in PTW. Sections were mounted
and imaged. All antibodies used are listed in **Table S5**.

Mounting for imaging

For embryos, dechorionated embryos were embedded in a 1.6 % low melting-point
agarose (Sigma) and 0.016 % Tricaine (Sigma) mixture in E3 medium, oriented within
1 min using a needle or fine hair, left at room temperature for 3 min, cooled at 4 °C for
3 min, supplemented with water drops to prevent evaporation. While for adult
telencephalon sections, fluorescently labelled brain sections were mounted on glass
slides with 0.17 mm coverslips using Aqua Polymount (PolySciences Inc., USA) and
stored at 4 °C in the dark before and after imaging [43].

Confocal imaging

Fluorescently labelled embryos and telencephalon sections were imaged using Leica
TCS SP5 DM5000 and SP8/DLS confocal fluorescence microscope (Leica
Microsystems, Wetzlar, Germany) equipped with 405 nm, 488 nm, 561 nm and 633
nm lasers. Excitation (Ex) wavelengths and emission (Em) ranges were as follows: Ex
488 nm for AlexaFluor 488 (EGFP, Em 492-550 nm), Ex 561 nm for AlexaFluor 561
(PCNA, Em 565-605 nm), Ex 633 nm for AlexaFluor 633 (S100β, Em 650-740 nm),
Ex 633 nm for AlexaFluor 633-labeled *mdka* RNA probe (EM 655-724 nm), and Ex
405 nm for DAPI (Em 409-480 nm). Imaging was performed with sequential scanning
at 16-bit color depth with a 1.0 AU pinhole aperture. Images were acquired at a
resolution of 1024 × 1024-pixel resolution, with a pixel size of approximately 1.55 µm
× 1.55 µm (field of view: ~1.55 mm × 1.55 mm), using a 63 ×/1.20 NA Water Immersion
objective (Leica Microsystems) at a scanning frequency of 400 Hz. Laser intensity,
PMT gain, detector settings, AOTF (Acousto-Optic Tuneable Filters) attenuation and
other imaging settings were kept constant across experimental groups to ensure
comparability.

Image processing and arrangement

Images were processed with consistent exposure times and gain settings across all
series. Z-stack maximum projections were generated and cropping or rotation was
performed in ImageJ/Fiji. For wide-field images in **Figure S2**, deconvolution using the
Iterative Deconvolve 3D and Diffraction PSF 3D plugins were applied to remove out-
of-focus blur and enhance clarity. All figures, including flowcharts, were created and
arranged using Microsoft PowerPoint (version 16.78) or Adobe Illustrator 2022
(version 26.4.1).

Statistical analysis

Three sections per brain were analyzed from at least three individuals (with sample

sizes indicated in the figure legends). Each brain section was treated as a single data

point. All statistical analyses were performed and visualized using R. To determine the

appropriate statistical test, data normality was assessed using the *Shapiro–Wilk* test

with a significance level of $\alpha = 0.05$. For normally distributed data, parametric tests

were used: the *Welch t*-test for comparisons between two samples, and the *Tukey*

HSD test following a significant *ANOVA* ($\alpha = 0.05$) for comparisons involving more

than two groups. For non-normally distributed data, the *Mann–Whitney U* test was

used for unpaired two-sample comparisons, and the *Dunn* test was applied following

a significant *Kruskal–Wallis* rank sum test ($\alpha = 0.05$) for multiple-group comparisons.

Acknowledgements

We would like to sincerely thank Anne Schröck for her invaluable technical support

throughout this project. We are grateful to Luisa Lübke, Gaoqun Zhang, and Barbara

Schornick for their contributions during the early phase of this study, and to Fushun

Chen for his initial support with bioinformatics. We also thank Ferenc Müller and Yavor

Hadzhiev for their helpful discussions and advice on bioinformatics approaches.

Research in Sepand Rastegar's laboratory is supported by the BioInterfaces in

Technology and Medicine and Natural, Artificial, and Cognitive Information Processing

(NACIP) programmes of the Helmholtz Association, as well as by project grants from

the German Research Foundation (DFG GRK2039, STR 439/17-1, RA 3469/5-1) and

the Marie Skłodowska-Curie Initial Training Networks (ITNs) ZenCode and Danio-

Recode. Jincan Chen is supported by a grant from the China Scholarship Council

(CSC), grant number 202106310017. We thank the KIT Publication Fund of the

Karlsruhe Institute of Technology for its support. Open access funding was enabled

and organized by Projekt DEAL.

Conflict of interest

The authors declare no conflict of interest.

Author contributions

SR designed, supervised the experiments, analysed the data, and co-wrote the

manuscript with JC, MT and ND. JC, AH, MT and TB conducted experiments and

analysed data. JC and MT analysed and quantified the data. CV conducted the Hi-C

and 4C data analysis. All authors have reviewed and approved the final version of the

manuscript for publication.

Data availability statement

The data that supports the findings of this study are available in the main text and the

supporting information of this article.

References

1.   Chen, J. C., Sanchez-Iranzo, H., Diotel, N. & Rastegar, S. (2024) Comparative insight into the regenerative mechanisms of the adult brain in zebrafish and mouse: highlighting the importance of the immune system and inflammation in successful regeneration, *Febs Journal.* **291**, 4193-4205.

2.   Marques, I. J., Lupi, E. & Mercader, N. (2019) Model systems for regeneration: zebrafish, *Development.* **146**.

3.   Becker, T. & Becker, C. G. (2022) Regenerative neurogenesis: the integration of developmental physiological and immune signals, *Development.* **149**.

4.   Gemberling, M., Bailey, T. J., Hyde, D. R. & Poss, K. D. (2013) The zebrafish as a model for complex tissue regeneration, *Trends Genet.* **29**, 611-620.

5.   Kizil, C., Kaslin, J., Kroehne, V. & Brand, M. (2012) Adult neurogenesis and brain regeneration in zebrafish, *Dev Neurobiol.* **72**, 429-461.

6.   Alunni, A. & Bally-Cuif, L. (2016) A comparative view of regenerative neurogenesis in vertebrates, *Development.* **143**, 741-753.

7.   Howe, K.Clark, M. D.Torroja, C. F.Torrance, J.Berthelot, C.Muffato, M.Collins, J. E.Humphray, S.McLaren, K.Matthews, L.McLaren, S.Sealy, I.Caccamo, M.Churcher, C.Scott, C.Barrett, J. C.Koch, R.Rauch, G. J.White, S.Chow, W.Kilian, B.Quintais, L. T.Guerra-Assunçao, J. A.Zhou, Y.Gu, Y.Yen, J.Vogel, J. H.Eyre, T.Redmond, S.Banerjee, R.Chi, J. X.Fu, B. Y.Langley, E.Maguire, S. F.Laird, G. K.Lloyd, D.Kenyon, E.Donaldson, S.Sehra, H.Almeida-King, J.Loveland, J.Trevanion, S.Jones, M.Quail, M.Willey, D.Hunt, A.Burton, J.Sims, S.McLay, K.Plumb, B.Davis, J.Clee, C.Oliver, K.Clark, R.Riddle, C.Eliott, D.Threadgold, G.Harden, G.Ware, D.Mortimer, B.Kerry, G.Heath, P.Phillimore, B.Tracey, A.Corby, N.Dunn, M.Johnson, C.Wood, J.Clark, S.Pelan, S.Griffiths, G.Smith, M.Glithero, R.Howden, P.Barker, N.Stevens, C.Harley, J.Holt, K.Panagiotidis, G.Lovell, J.Beasley, H.Henderson, C.Gordon, D.Auger, K.Wright, D.Collins, J.Raisen, C.Dyer, L.Leung, K.Robertson, L.Ambridge, K.Leongamornlert, D.McGuire, S.Gilderthorp, R.Griffiths, C.Manthravadi, D.Nichol, S.Barker, G.Whitehead, S.Kay, M., et al. (2013) The zebrafish reference genome sequence and its relationship to the human genome, *Nature.* **496**, 498-503.

8. Driever, W., SolnicaKrezel, L., Schier, A. F., Neuhauss, S. C. F., Malicki, J., Stemple, D. L., Stainier, D. Y. R., Zwartkruis, F., Abdelilah, S., Rangini, Z., Belak, J. & Boggs, C. (1996) A genetic screen for mutations affecting embryogenesis in zebrafish, *Development.* **123**, 37-46.

9. Ellingsen, S., Laplante, M. A., König, M., Kikuta, H., Furmanek, T., Hoivik, E. A. & Becker, T. S. (2005) Large-scale enhancer detection in the zebrafish genome, *Development.* **132**, 3799-3811.

10. Choe, C. P., Choi, S. Y., Kee, Y., Kim, M. J., Kim, S. H., Lee, Y., Park, H. C. & Ro, H. (2021) Transgenic fluorescent zebrafish lines that have revolutionized biomedical research, *Lab Anim Res.* **37**, 26.

11. Parinov, S., Kondrichin, I., Korzh, V. & Emelyanov, A. (2004) Tol2 transposon-mediated enhancer trap to identify developmentally regulated zebrafish genes in vivo, *Dev Dyn.* **231**, 449-59.

12. Hörtenhuber, M., Mukarram, A. K., Stoiber, M. H., Brown, J. B. & Daub, C. O. (2020) *-DCC: A platform to collect, annotate, and explore a large variety of sequencing experiments, *Gigascience.* **9**.

13. Yang, H., Luan, Y., Liu, T., Lee, H. J., Fang, L., Wang, Y., Wang, X., Zhang, B., Jin, Q., Ang, K. C., Xing, X., Wang, J., Xu, J., Song, F., Sriranga, I., Khunsriraksakul, C., Salameh, T., Li, D., Choudhary, M. N. K., Topczewski, J., Wang, K., Gerhard, G. S., Hardison, R. C., Wang, T., Cheng, K. C. & Yue, F. (2020) A map of cis-regulatory elements and 3D genome structures in zebrafish, *Nature.* **588**, 337-343.

14. Baranasic, D., Hörtenhuber, M., Balwierz, P. J., Zehnder, T., Mukarram, A. K., Nepal, C., Várnai, C., Hadzhiev, Y., Jimenez-Gonzalez, A., Li, N., Wragg, J., D'Orazio, F. M., Relic, D., Pachkov, M., Díaz, N., Hernández-Rodríguez, B., Chen, Z. L., Stoiber, M., Dong, M., Stevens, I., Ross, S. E., Eagle, A., Martin, R., Obasaju, O., Rastegar, S., McGarvey, A. C., Kopp, W., Chambers, E., Wang, D., Kim, H. R., Acemel, R. D., Naranjo, S., Lapinski, M., Chong, V., Mathavan, S., Peers, B., Sauka-Spengler, T., Vingron, M., Carninci, P., Ohler, U., Lacadie, S. A., Burgess, S. M., Winata, C., van Eeden, F., Vaquerizas, J. M., Gómez-Skarmeta, J. L., Onichtchouk, D., Brown, B., Bogdanovic, O., van Nimwegen, E., Westerfield, M., Wardle, F. C., Daub, C. O., Lenhard, B. & Müller, F. (2022) Multiomic atlas with functional stratification and developmental dynamics of zebrafish regulatory elements, *Nat Genet.* **54**, 1037-1050.

15. Diotel, N., Lübke, L., Strähle, U. & Rastegar, S. (2020) Common and distinct features of adult neurogenesis and regeneration in the telencephalon of zebrafish and mammals, *Front Neurosci-Switz.* **14**.

16. Ghaddar, B., Lübke, L., Couret, D., Rastegar, S. & Diotel, N. (2021) Cellular mechanisms participating in brain repair of adult zebrafish and mammals after injury, *Cells-Basel.* **10**.

17. Rodriguez Viales, R., Diotel, N., Ferg, M., Armant, O., Eich, J., Alunni, A., März, M., Bally-Cuif, L., Rastegar, S. & Strahle, U. (2015) The helix-loop-helix protein Id1 controls stem cell proliferation during regenerative neurogenesis in the adult zebrafish telencephalon, *Stem Cells.* **33**, 892-903.

18. Demirci, Y., Cucun, G., Poyraz, Y. K., Mohammed, S., Heger, G., Papatheodorou, I. & Ozhan, G. (2020) Comparative Transcriptome Analysis of the Regenerating Zebrafish Telencephalon Unravels a Resource With Key Pathways During Two Early Stages and Activation of Wnt/beta-Catenin Signaling at the Early Wound Healing Stage, *Front Cell Dev Biol.* **8**, 584604.

19. Gourain, V., Armant, O., Lubke, L., Diotel, N., Rastegar, S. & Strahle, U. (2021) Multi-Dimensional Transcriptome Analysis Reveals Modulation of Cholesterol Metabolism as Highly Integrated Response to Brain Injury, *Front Neurosci.* **15**, 671249.

20. Lange, C., Rost, F., Machate, A., Reinhardt, S., Lesche, M., Weber, A., Kuscha, V., Dahl, A., Rulands, S. & Brand, M. (2020) Single cell sequencing of radial glia progeny reveals the diversity of newborn neurons in the adult zebrafish brain, *Development.* **147**.

21. Lübke, L., Zhang, G. Q., Straehle, U. & Rastegar, S. (2022) *mdka* expression is associated with quiescent neural stem cells during constitutive and reactive neurogenesis in the adult zebrafish telencephalon, *Brain Sci.* **12**.

22. Winkler, C. & Yao, S. (2014) The midkine family of growth factors: diverse roles in nervous system formation and maintenance, *Br J Pharmacol.* **171**, 905-12.

23. Aynacioglu, A. S., Bilir, A. & Tuna, M. Y. (2019) Involvement of midkine in autoimmune and autoinflammatory diseases, *Mod Rheumatol.* **29**, 567-571.

24. Zhang, G. Q., Lübke, L., Chen, F. S., Beil, T., Takamiya, M., Diotel, N., Strähle, U. & Rastegar, S. (2021) Neuron-radial glial cell communication via BMP/Id1 signaling is key to long-term maintenance of the regenerative capacity of the adult zebrafish telencephalon, *Cells-Basel.* **10**.

25. Zhang, G. Q., Ferg, M., Lübke, L., Takamiya, M., Beil, T., Gourain, V., Diotel, N., Strähle, U. & Rastegar, S. (2020) Bone morphogenetic protein signaling regulates Id1-mediated neural stem cell quiescence in the adult zebrafish brain via a phylogenetically conserved enhancer module, *Stem Cells.* **38**, 875-889.

26. Adolf, B., Chapouton, P., Lam, C. S., Topp, S., Tannhäuser, B., Strähle, U., Götz, M. & Bally-Cuif, L. (2006) Conserved and acquired features of adult neurogenesis in the zebrafish telencephalon, *Dev Biol.* **295**, 278-293.

27. März, M., Schmidt, R., Rastegar, S. & Strähle, U. (2011) Regenerative response following stab injury in the adult zebrafish telencephalon, *Dev Dynam.* **240**, 2221-2231.

28. Grivas, D., Gonzalez-Rajal, A. & de la Pompa, J. L. (2021) Midkine-a regulates the formation of a fibrotic scar during zebrafish heart regeneration, *Front Cell Dev Biol.* **9**, 669439.

29. Gramage, E., D'Cruz, T., Taylor, S., Thummel, R. & Hitchcock, P. F. (2015) Midkine-a protein localization in the developing and adult retina of the zebrafish and its function during photoreceptor regeneration, *Plos One.* **10**, e0121789.

30. Ang, N. B., Saera-Vila, A., Walsh, C., Hitchcock, P. F., Kahana, A., Thummel, R. & Nagashima, M. (2020) Midkine-a functions as a universal regulator of proliferation during epimorphic regeneration in adult zebrafish, *Plos One.* **15**, e0232308.

31. Nagashima, M., D'Cruz, T. S., Danku, A. E., Hesse, D., Sifuentes, C., Raymond, P. A. & Hitchcock, P. F. (2020) Midkine-a is required for cell cycle progression of Müller glia during neuronal regeneration in the vertebrate retina, *J Neurosci.* **40**, 1232-1247.

32. Kikuchi-Horie, K., Kawakami, E., Kamata, M., Wada, M., Hu, J. G., Nakagawa, H., Ohara, K., Watabe, K. & Oyanagi, K. (2004) Distinctive expression of midkine in the repair period of rat brain during neurogenesis: immunohistochemical and immunoelectron microscopic observations, *J Neurosci Res.* **75**, 678-87.

33. Lien, C. L., Schebesta, M., Makino, S., Weber, G. J. & Keating, M. T. (2006) Gene
expression analysis of zebrafish heart regeneration, *Plos Biol.* **4**, e260.

34. Thompson, J. D., Ou, J. H., Lee, N., Shin, K., Cigliola, V., Song, L. Y., Crawford,
G. E., Kang, J. S. & Poss, K. D. (2020) Identification and requirements of enhancers
that direct gene expression during zebrafish fin regeneration, *Development.* **147**.

35. Weinberger, M., Simoes, F. C., Gungoosingh, T., Sauka-Spengler, T. & Riley, P. R.
(2024) Distinct epicardial gene regulatory programs drive development and
regeneration of the zebrafish heart, *Dev Cell.* **59**, 351-367.

36. Winkler, C., Schäfer, M., Duschl, J., Schartl, M. & Volff, J. N. (2003) Functional
divergence of two zebrafish midkine growth factors following fish-specific gene
duplication, *Genome Res.* **13**, 1067-1081.

37. Gonzalez, A. J., Baranasic, D. & Müller, F. (2023) Zebrafish regulatory genomic
resources for disease modelling and regeneration, *Dis Model Mech.* **16**.

38. Chen, Z. L., Omori, Y., Koren, S., Shirokiya, T., Kuroda, T., Miyamoto, A., Wada,
H., Fujiyama, A., Toyoda, A., Zhang, S. Y., Wolfsberg, T. G., Kawakami, K., Phillippy,
A. M., Mullikin, J. C., Burgess, S. M. & Prog, N. C. S. (2019) De novo assembly of the
goldfish (*Carassius auratus*) genome and the evolution of genes after whole-genome
duplication, *Sci Adv.* **5**.

39. Meng, A. M., Tang, H., Ong, B. A., Farrell, M. J. & Lin, S. (1997) Promoter analysis
in living zebrafish embryos identifies a *cis*-acting motif required for neuronal
expression of GATA-2, *P Natl Acad Sci USA.* **94**, 6267-6272.

40. Navratilova, P., Fredman, D., Hawkins, T. A., Turner, K., Lenhard, B. & Becker, T.
S. (2009) Systematic human/zebrafish comparative identification of *cis*-regulatory
activity around vertebrate developmental transcription factor genes, *Dev Biol.* **327**,
526-540.

41. Rastegar, S., Hess, I., Dickmeis, T., Nicod, J. C., Ertzer, R., Hadzhiev, Y., Thies,
W. G., Scherer, G. & Strähle, U. (2008) The words of the regulatory code are arranged
in a variable manner in highly conserved enhancers, *Dev Biol.* **318**, 366-377.

42. Begeman, I. J., Emery, B., Kurth, A. & Kang, J. S. (2022) Regeneration and
developmental enhancers are differentially compatible with minimal promoters, *Dev
Biol.* **492**, 47-58.

43. Schmidt, R., Beil, T., Strähle, U. & Rastegar, S. (2014) Stab wound injury of the
zebrafish adult telencephalon: A method to investigate vertebrate brain neurogenesis
and regeneration, *Jove-J Vis Exp*.

44. Beisaw, A., Kuenne, C., Guenther, S., Dallmann, J., Wu, C. C., Bentsen, M.,
Looso, M. & Stainier, D. Y. R. (2020) AP-1 contributes to chromatin accessibility to
promote sarcomere disassembly and cardiomyocyte protrusion during zebrafish heart
regeneration, *Circ Res.* **126**, 1760-1778.

45. Nakamura, M., Yoshida, H., Takahashi, E., Wlizla, M., Takebayashi-Suzuki, K.,
Horb, M. & Suzuki, A. (2020) The AP-1 transcription factor JunB functions in Xenopus
tail regeneration by positively regulating cell proliferation. **522**, 990-995.

46. Wang, W., Hu, C., Zeng, A., Alegre, D., Hu, D., Gotting, K., Ortega Granillo, A.,
Wang, Y., Robb, S., Schnittker, R., Zhang, S., Alegre, D., Li, H., Ross, E., Zhang, N.,
Brunet, A. & Sánchez Alvarado, A. (2020) Changes in regeneration-responsive
enhancers shape regenerative capacities in vertebrates, *Science.* **369**, eaaz3090.

47. Tamaki, T., Yoshida, T., Shibata, E., Nishihara, H., Ochi, H. & Kawakami, A. (2023)
Splashed E-box and AP-1 motifs cooperatively drive regeneration response and shape
regeneration abilities, *Biol Open.* **12**.

48. Dong, Y., Yang, Y., Wang, H., Feng, D., Nist, E., Yapundich, N., Spurlock, B., Craft,
M., Qian, L. & Liu, J. (2024) Single-cell chromatin profiling reveals genetic programs
activating proregenerative states in nonmyocyte cells, *Sci Adv.* **10**, eadk4694.

49. Gramage, E., Li, J. & Hitchcock, P. (2014) The expression and function of midkine
in the vertebrate retina, *Brit J Pharmacol.* **171**, 913-923.

50. Goto, T., Macdonald, P. & Maniatis, T. (1989) Early and late periodic patterns of
even skipped expression are controlled by distinct regulatory elements that respond
to different spatial cues, *Cell.* **57**, 413-422.

51. Berrocal, A., Lammers, N. C., Garcia, H. G. & Eisen, M. B. (2020) Kinetic sculpting
of the seven stripes of the Drosophila even-skipped gene, *Elife.* **9**, e61635.

52. Werner, T., Hammer, A., Wahlbuhl, M., Bösl, M. R. & Wegner, M. (2007) Multiple
conserved regulatory elements with overlapping functions determine Sox10
expression in mouse embryogenesis, *Nucleic Acids Res.* **35**, 6526-38.

53. Betancur, P., Bronner-Fraser, M. & Sauka-Spengler, T. (2010) Genomic code for
Sox10 activation reveals a key regulatory enhancer for cranial neural crest, *Proc Natl*
Acad Sci U S A. **107**, 3570-5.

54. Aleström, P., D'Angelo, L., Midtlyng, P. J., Schorderet, D. F., Schulte-Merker, S.,
Sohm, F. & Warner, S. (2020) Zebrafish: Housing and husbandry recommendations,
Lab Anim-Uk. **54**, 213-224.

55. Kwan, K. M., Fujimoto, E., Grabher, C., Mangum, B. D., Hardy, M. E., Campbell,
D. S., Parant, J. M., Yost, H. J., Kanki, J. P. & Chien, C. B. (2007) The *Tol2*kit: A
multisite gateway-based construction kit for *Tol2* transposon transgenesis constructs,
Dev Dynam. **236**, 3088-3099.

Table 1. Summary of EGFP expression patterns driven by individual CREs in 24 hpf embryos.

	Endogenous	CRE1	CRE2	CRE3	CRE4	CRE5	CRE6
Brain	++	-	+	+	+	+	+
Retina	++	-	+	++	++	+	++
Spinal cord	++	-	+	++	+	+	+
Notochord	+	-	+	-	-	-	-
Fin fold/Tail	++	-	-	++	+	-	+
Somite/Muscle	++	-	+	-	++	-	++

Note: -, no detectable expression; +, weak expression; ++, strong expression.

Table 2. Summary of EGFP expression patterns driven by individual CREs in the adult telencephalon.

	Endogenous	CRE1	CRE2	CRE3	CRE4	CRE5	CRE6
Dm	++	-	++	-	+	-	+
Dl	++	-	+	+	+	-	+
Dp	+	-	-	++	-	-	-
Vd	+	-	+	-	-	-	-
Vv	+	-	+	-	++	-	-

Note: -, no detectable expression; +, weak expression; ++, strong expression. Abbreviations: Dm, dorsomedial pallium; Dp, dorsal pallium; Dl, lateral pallium; VZ, ventricular zone; Vd, dorsal nucleus of ventral telencephalon; Vv, ventral nucleus of ventral telencephalon.

Table 3. Summary of EGFP expression patterns observed in distinct *mdka* CRE transgenic lines.

CRE(s)	Activity summary
CRE2	Adult regulatory sequence; active in physiological conditions; best matching *mdka* expression
CRE3	Predominantly active during embryonic development; overlaps with mdka expression; shows ectopic expression in the adult telencephalon in Dp and Dl.
CRE4	Adult regulatory sequence; active in NSCs; does not show a significant response to injury
CRE6	Adult regulatory sequence; active in NSCs with ectopic activity in neurons; no significant response to injury
CRE2346	Recapitulates *mdka* expression; EGFP reporter overlaps with *mdka* mRNA in both embryo and adult; inducible upon injury

Note: Dp, dorsal pallium; Dl, lateral pallium; NSCs, neural stem cells.

 Figure legends

Figure 1. Identification of potential *mdka* regulatory sequences based on
embryonic developmental ATAC-seq data. Snapshot of UCSC genome browser
tracks displaying ATAC-seq data from different embryonic developmental stages
(Dome, 75%-epiboly, 5-9 somites, Prime-5 and Long-pec) across a 45-kilobase region
(chr7: 38,875,000-38,920,000; danRer10 assembly). Green peak profiles represent
open chromatin regions identified by ATAC-seq. Color-coded horizontal bars indicate
the genomic position of putative regulatory elements based on PADRES (Predicted
ATAC-seq-supported developmental regulatory elements): promoters (1_TssA and
2_TssA2, blue), enhancers (5_EnhA1, red; 7_EnhWk1, orange), and other elements
(e.g., 10_Quies, grey). The grey box highlights six putative *mdka* CREs (CRE1 to
CRE6) identified in this study, while the magenta box marks the putative *mdka*
promoter region. The relative positions of all CREs are shown in reference to the *mdka*
gene and its transcription start site (TSS). PhastCons conservation scores derived
from zebrafish-carp genome alignment highlights conserved non-coding regions
(CNEs) associated with the *mdka* locus.

Figure 2. Stable EGFP transgenic line generated using potential *mdka* CREs. A,
Schematic of the plasmid pTol2-mdkaCREx-gata2aPR:egfp-pA, where CREx
represents one of the six putative CREs (CRE1 to CRE6). **B-G**, Stable EGFP reporter
expression driven by *mdka* CRE1 (B), CRE2 (C), CRE3 (D), CRE4 (E), CRE5 (F),
CRE6 (G) in embryo at 24 hours post-fertilization (hpf). **B'-G'**, Merged images showing
mdka mRNA (magenta) and EGFP (green) expression in the corresponding CRE
transgenic lines. *Tg(CRE3-gata2aPR:EGFP)* embryo exhibited strong co-localization
of *mdka* mRNA and EGFP signals in the spinal cord (yellow arrowheads, D, D') and
fin fold region (white arrowheads, D, D'). *Tg(CRE4-gata2aPR:EGFP)* and *Tg(CRE6-*
gata2aPR:EGFP) embryos showed strong activity in the retina (green arrowheads)
and brain regions (red arrowheads, E, E', G, G'), respectively. *Tg(CRE1-*
gata2aPR:EGFP) showed no detectable EGFP expression (B, B'). Scale bar: 200 μm.
766
Figure 3. The *mdka* transgenic lines, CRE2, CRE4 and CRE6, drive radial glial
EGFP expression in the adult telencephalon during constitutive neurogenesis.
A, Schematic illustrating the orientation of transverse sections through the
telencephalon, with reference to the location of radial glial cells (RGCs). Dm:

dorsomedial pallium; VZ: ventricular zone; RGCs: radial glial cells. **B-G'**, EGFP
reporter expression in the adult telencephalon across various *mdka* CRE transgenic
lines: CRE1 (B-B'. *n* = 3 brains), CRE2 (C-C', *n* = 4), CRE3 (D-D', *n* = 6), CRE4 (E-E',
n = 3), CRE5 (F-F', *n* = 4), and CRE6 (G-G', *n* = 3). The location of VZ is indicated on
the left hemisphere of CRE1 telencephalon, where EGFP expression is absent (B-B').
Rectangles in B-G indicate magnified regions shown in B'-G', respectively. Dashed
lines outline the telencephalon contour (B-G). **H-J''**, Magnified transverse views of the
Dm region showing EGFP (green) and *mdka* mRNA expression (magenta) in three
transgenic lines with radial glial EGFP expression: *Tg(CRE2-gata2aPR:EGFP)*(H-H''),
Tg(CRE4-gata2aPR:EGFP)(I-I''), and *Tg(CRE6-gata2aPR:EGFP)* (J-J''). White
arrowheads highlight RGCs of *mdka* mRNA and EGFP co-localization, reflecting
faithful activity of CRE2, CRE4 and CRE6. Arrows in the panel J'' indicate the ectopic
EGFP expression of CRE6 located outside of VZ. **K**, The fidelity of CRE2, CRE4 and
CRE6 activity was evaluated by quantifying the percentage of EGFP-positive cells
among endogenous *mdka*-positive RGCs with the confocal images as shown the
panels H-J. Results are shown as boxplots with the median values. No significant
difference was observed between different CREs (*ANOVA* test, *p* = 0.84). **L**, S100β+
EGFP cells are classified either as type 1 (quiescent, PCNA-negative) or type 2
(activated, PCNA-positive) based on PCNA expression. **M**, Quantification of type 1
and type 2 RGC population in CRE2, CRE4, and CRE6. The mean ± standard
deviation (error bar) of type 1 RGCs is expressed as a percentage of the total RGC
population. No significant difference was observed between different CREs (*n.s.*;
Kruskal-Wallis test, *p* = 0.97). Representative confocal images corresponding to this
quantification are provided in Supplementary Figure S3A-A''', B-B''' and Figure S4A-
A''', C-C''', showing co-immunostaining for EGFP, S100β, and PCNA. Each data point
represents an individual brain slice. Scale bar: 200 μm (B-G), 25 μm (B'-G', H-J) and
10 μm (H'-H'', I'-I'', J'-J'' and L). *n* = 3 brains (CRE1; B, B'), *n* = 6 (CRE2; C, C', H-H'',
K, M), *n* = 6 (CRE3; D, D'), *n* = 3 (CRE4; E, E', I-I'', K, M), *n* = 4 (CRE5; F, F'), *n* = 3
(CRE6; G, G', J-J', K, M).

Figure 4. During regenerative neurogenesis, *mdka* CRE2, CRE4 and CRE6,
maintain radial glial EGFP expression, though show variability in the injured
brain hemisphere. A, Schematic illustrating the experimental design and orientation
of telencephalic brain sections. Adult zebrafish were subjected to stab injury in the

right hemisphere of the telencephalon, and brains were collected and fixed at 5 days
post-lesion (dpl). **B-G'**, EGFP expression from stable transgenic lines for CRE1 (B,
B'), CRE2 (C, C'), CRE3 (D, D'), CRE4 (E, E'), CRE5 (F, F'), and CRE6 (G, G') is
shown for both contralateral and ipsilateral sides (asterisks indicate ipsilateral/injured
side) of the telencephalon. Rectangles in B-G indicate magnified regions shown in B'-
G', respectively. Dashed lines mark the telencephalon contour (B)**. H,** Quantification
of EGFP-positive and S100β-positive RGCs. The number of EGFP-positive RGCs in
the injured (ipsilateral) hemispheres of CRE2, CRE4, and CRE6 transgenic lines is
shown as a percentage relative to the corresponding contralateral side in boxplots. No
significant differences were observed among CREs (*ANOVA, p* > 0.05). **I,** Stemness
of EGFP-positive RGCs in CRE2, CRE4, and CRE6 was assessed. Relative
proportions of type 1 (quiescent, S100β$^+$/PCNA$^-$) and type 2 (activated,
S100β$^+$/PCNA$^+$) RGCs in contralateral and ipsilateral hemispheres are presented as
stacked bar plots. **While no statistical differences were found among CREs in the
contralateral hemisphere, a significant difference was observed between CRE2 and
CRE6 in the ipsilateral (injured) hemisphere (*p* < 0.01). Representative confocal
images used for quantifications in panels H and I are provided in Supplementary
Figures S3C-D''' and S4B-D''', showing co-immunostaining for EGFP, S100β, and
PCNA. Each data point represents an individual brain slice. Scale bars: 200 µm (B-
G); 25 µm (B'-G'). Sample sizes: *n* = 3 brains (B, B'), *n* = 4 (C, C'), *n* = 5 (D, D'), *n* = 3
(E, E'), *n* = 4 (F, F'), *n* = 3 (G, G'), *n* = 3 for each CRE (H, I).
826
Figure 5. Combination of CREs enhances faithful transgene expression in both
embryos and the adult telencephalon under homeostatic conditions. A,
Schematic of the pTol2-mdkaCRE2346-gata2aPR:egfp-pA construct, in which a
combination of CRE2, CRE3, CRE4, and CRE6 drives EGFP expression initiated by
the *gata2a* promoter region (*gata2aPR*). **B-B''**, Comparison of EGFP transgene (B')
and *mdka* mRNA (B'') expression in 24 hpf embryo. Merged EGFP and *mdka* mRNA
signals (B) show co-localization primarily in the brain and spinal cord (white arrows).
C, Merged EGFP and *mdka* mRNA expression in a transverse section of the adult
telencephalon under homeostatic conditions. The magnified Dm region (dashed
rectangle) is shown in D-D''. **D-D''**, Magnified views of the telencephalic Dm domain
showing merged signals (D), EGFP driven by CRE2346 (green, D'), and endogenous
mdka mRNA (magenta, D''). White arrowheads indicate RGCs in the ventricular zone

(VZ). **E,** Faithfulness of EGFP transgene expression driven by CRE2, CRE4, CRE6

and the combined CRE2346. EGFP expression in VZ of each CRE transgenic line was

compared with endogenous *mdka* mRNA. Data are shown as boxplots representing

the percentage of EGFP$^+$/*mdka*$^+$ double-positive cells relative to total *mdka*$^+$ cells.

Median values are indicated. Significant differences were observed between

CRE2346 and individual CREs (*ANOVA* followed by *Tukey's* test; CRE2346 vs CRE2

[$p < 0.01$], vs CRE4 [$p < 0.01**$], vs CRE6 [$p < 0.05*$]). The data for individual CREs

are identical to that shown in Figure 3K. **F-G**, Stemness analysis of RGCs labeled by

CRE2346-driven EGFP. EGFP-positive cells in the VZ of the telencephalic Dm domain

were categorized as type 1 (quiescent, S100β$^+$/PCNA$^-$) or type 2 (activated,

S100β$^+$/PCNA$^+$) using immunolabeling. **F-F'''**, Transverse telencephalon sections

showing EGFP (F'), PCNA (F''), S100β (F''') and merged image (F), highlighting type

1 and 2 RGCs under homeostatic conditions. **G,** Stemness of CRE2346-driven EGFP-

positive RGCs was assessed and compared to those of individual CREs in a stacked

bar chart. The individual CRE data are reused from Figure 3M. Mean ± standard

deviation of type 1 RGCs (as a proportion of EGFP-positive RGCs) is shown. Each

data point represents an individual brain slice. No significant differences were

observed among different CREs (*n.s.*; *Kruskal-Wallis* test, $p = 0.05$). Scale bars: 200

μm (B-B'' and C); 25 μm (D-D'', F-F'''). Sample sizes: $n = 3$ brains (C, D-D'', E); $n = 5$

brains (F-F''', G).

Figure 6. Combination of CREs enhanced faithful transgene expression in the

adult telencephalon after brain injury. A-C'', Merged images of EGFP expression

driven by CRE2346 (green) and endogenous *mdka* mRNA (magenta) in a transverse

section of the adult telencephalon during regenerative neurogenesis at 5 dpl. The

hemisphere ipsilateral to the injury is marked with an asterisk. **A**, Overview of the

telencephalon. Stippled rectangular regions indicate areas magnified in B-B'' (Dm

domain) and C-C'' (site of injury). **B-B''**, Magnified merged image of CRE2346-driven

EGFP (green, B') and endogenous *mdka* mRNA (magenta, B''). **C-C'**, Magnified

merged view at the site of injury (C), showing upregulated expression of CRE2346-

driven EGFP (green, C') and endogenous *mdka* mRNA (magenta, C''). **D,** The number

of S100β-positive RGCs expressing either CRE transgenes or endogenous *mdka*

mRNA in the hemisphere ipsilateral to the injury did not differ between CRE constructs

and the endogenous gene (*n.s.*; *Kruskal-Wallis* test, $p = 0.11$). The number of
ipsilateral RGCs is normalized to that of contralateral RGCs. **E**, In the CRE2346
transgenic line, EGFP levels in the ipsilateral hemisphere are significantly higher than
in the contralateral side (*Welch t*-test, $p < 0.001$***). Normalized integrated
fluorescence across brain slices is presented in arbitrary units (AU). **F-G**, Faithfulness
of (F) and ectopic (G) EGFP transgene expression driven by either CRE2, CRE4,
CRE6 or CRE2346 in contralateral ("c") and ipsilateral hemispheres ("I") after brain
injury. **F**, EGFP expression in VZ of each CRE transgenic line were compared with
endogenous *mdka* mRNA expression. Results are presented in boxplots showing the
percentage of cells double-positive for EGFP and endogenous *mdka* mRNA relative
to the total number of *Stemness*-positive cells. Significant differences were observed
between CRE2346 and each individual CREs (*Dunn's post hoc* test after *Kruskal–*
Wallis: CRE2346 vs CRE2 [$p < 0.01$**], vs CRE4 [$p < 0.01$**] and vs CRE6 [$p <
0.001$***]). No significant difference was observed between contralateral and
ipsilateral hemispheres in all cases (*n.s.*; *Dunn's post hoc* test after *Kruskal–Wallis*
analysis). **G**, Ectopic EGFP expression outside the VZ of each CRE transgenic line
are shown relative to the number of endogenous *mdka*-positive RGCs. Although no
significant difference was observed between contralateral and ipsilateral hemispheres,
CRE2346 showed significantly lower ectopic transgene expression compared to
CRE4 (Adjusted *p*-values in *Dunn* test following *Kruskal-Wallis* test; $p < 0.001$***) and
CRE6 ($p < 0.001$***). **H-J**, Evaluation of RGC stemness in the CRE2346 line using
PCNA and S100β immunostaining. **H-I'''**, Transverse section of the adult
telencephalon during regenerative neurogenesis at 5 dpl, showing the merged image
(H-I) of CRE2346-driven EGFP (green, H'-I'), PCNA (magenta, H''-I'') and S100β
(blue, H'''-I'''). The white rectangular regions in H-H''' are magnified in I-I''',
respectively. Representative type 1 (S100β$^+$/PCNA$^−$) and type 2 (S100β$^+$/PCNA$^+$)
RGCs are indicated (I-I'''). **J**, Stemness of CRE2346-driven EGFP-positive RGCs
compared to individual CRE lines and endogenous *mdka* mRNA ("*mdka*"). Pairwise
comparisons were performed using *Dunn*'s test following a significant *Kruskal–Wallis*
test (*n.s.* for $p >= 0.05$, * for $p < 0.05$, ** for $p < 0.01$ and *** for $p < 0.001$). The results
for hemispheres contralateral ("C") or ipsilateral ("I") to the injury are plotted adjacent
to each other. The data for CRE2, CRE4 and CRE6 are identical to Figure 4I. Each
data point represents an individual brain slice. Black solid dots indicate outliers. Scale
bars: 200 μm (A, H-H'''); 25 μm (B–B'', C-C'', I–I'''). Sample sizes: $n = 4$ brains (A, B-

B'', C-C''); n = 6 brains (CRE2346; D, H-H''', I-I''', J); n = 3 brains (E, F, G); n = 3 brains

(endogenous expression of *mdka*; D, J).

A pTol2-mdkaCREx-gata2aPR:egfp-pA

A Adult zebrafish — telencephalon — Dm — VZ — RGC

EGFP / unjured

CRE1	CRE2	CRE3	CRE4	CRE5	CRE6

B, C, D, E, F, G (200 µm)
B', C', D', E', F', G' (25 µm)
VZ, Dm labels

EGFP *mdka* mRNA

CRE2	CRE4	CRE6

H, H', H'', I, I', I'', J, J', J''
RGC, VZ, Dm labels (10 mm)

K Transgene fidelity

n.s.

% total *mdka*(+) RGC — 49%, 46%, 58%
CRE 2, 4, 6

L RGC classification

merged | EGFP | S100β | nucleus / PCNA

type 1 (quiescent) PCNA−
type 2 (activated) PCNA+
S100β+ (10 mm)

M

n.s.

% total EGFP(+) RGC
type 2 / type 1
79 ±8, 79 ±9, 79 ±6
CRE 2, 4, 6

A pTol2-mdkaCRE2346-gata2aPR:egfp-pA

24 hpf embryo

Adult telencephalon (faithfullness/homeostatic condition)

B merge

B' EGFP

B'' *mdka* mRNA

C merge

D merge

D' EGFP

D'' *mdka* mRNA

E transgene fidelity

Adult telencephalon (RGC stemness/homeostatic condition)

F merge

F' EGFP

F'' PCNA

F''' S100β

G