Histone H3 tail charge patterns govern nucleosome condensate formation and dynamics

Erin F. Hammonds[1], Anurag Singh[2], Krishna K. Suresh[3], Sean Yang[3], Sarah S. Meidl Zahorodny[1], Ritika Gupta[2], Davit A. Potoyan[3]*, Priya R. Banerjee[2]*, Emma A. Morrison[1]*

[1]Department of Biochemistry, Medical College of Wisconsin, 8701 Watertown Plank Rd., Milwaukee, WI 53226
[2]Department of Physics, The State University of New York at Buffalo, Buffalo, NY 14260
[3]Department of Chemistry, Iowa State University, Ames, IA 50011
*Corresponding: emorrison@mcw.edu, prbanerj@buffalo.edu, potoyan@iastate.edu

ABSTRACT

Emerging models of nuclear organization suggest that chromatin forms functionally distinct microenvironments through phase separation. As chromatin architecture is organized at the level of the nucleosome and regulated by histone post-translational modifications, we investigated how these known regulatory mechanisms influence nucleosome phase behavior. By systematically altering charge distribution within the H3 tail, we found that the terminal and central regions modulate the phase boundary and tune nucleosome condensate viscosity differentially, as revealed by microscopy-based assays, microrheology, and simulations. Nuclear magnetic resonance relaxation experiments revealed that H3 tails remain dynamically mobile within condensates, and their mobility correlates with condensate viscosity. These results demonstrate that the number, identity, and spatial arrangement of basic residues in the H3 tail critically regulate nucleosome phase separation. Our findings support a model in which nucleosomes, through their intrinsic properties and modifications, actively shape the local chromatin microenvironment—providing new insight into the histone language in chromatin condensates.

INTRODUCTION

Dynamic chromatin organization is being re-examined through the lens of phase separation (1–6). Current models invoke phase separation and biomolecular condensates in compartmentalization and the formation of localized environments that are functionally distinct within the nucleus, which dynamically change in response to stimuli (2, 3, 5–11). Chromatin regions ranging from transcriptionally silenced heterochromatin to transcriptionally active super-enhancers form phase-separated regions (12, 13), and condensates with such diverse functions have varied physicochemical and material properties (9, 14, 15). It is appealing to consider a model wherein the nucleus exists as a collection of phase-separated regions, each enriched in a distinct set of chromatin, epigenetic modifications, proteins, and RNA that determine the local environment and inform the biological activity within.

The nucleus is rich in multivalent biopolymers capable of undergoing phase separation, and chromatin is no exception with its histone and DNA components. Chromatin can undergo phase separation in the absence of chromatin-interacting proteins, and this holds true even down to the level of the nucleosome core particle (NCP), the basic subunit of chromatin (7, 16–21). Recent reports have also shown the effect of nucleosome spacing (length of linker DNA) on nucleosome mobility and thermodynamic stability of chromatin condensates (22, 23). Within the NCP, ~147-bp DNA wraps around a folded histone core, and the intrinsically disordered termini, or tails, of the histones protrude. The histone tails are enriched in basic residues—the histone tails range from 11-36 residues in length and are 31-42% lysines and arginines while only 0-8% aspartates and glutamates. The histone tails form both intra- and inter-nucleosomal interactions with DNA within the context of nucleosomes and nucleosome arrays, and the H3 and H4 tails dominate inter-nucleosomal interactions (24–33). Chromatin phase separation is driven, at least in part, by inter-nucleosomal interactions between these highly basic histone tails and DNA (7, 16, 21). In support, removal of the H3 or H4 tails individually or all histone tails simultaneously abrogates phase separation (7, 16). Furthermore, neutralizing mutations of the H4 tail basic patch abrogate phase separation while neutralization of the H2A/H2B acidic patch does not, supporting the importance of histone tail-DNA interactions over histone tail-histone core interactions in phase separation (7). Here, we hypothesize that site-specific mutations of basic residues within histone tails alter histone–DNA electrostatic interactions, thereby modulating the condensate properties of NCPs. Because the emergent properties of chromatin subcompartments arise from the underlying physical characteristics of their constituent chromatin units, it is essential to elucidate how such molecular-level perturbations influence the material properties of nucleosome condensates.

Post-translational modifications (PTMs) on histone proteins are epigenomic marks that regulate chromatin organization, including via phase separation (34). Although histone PTMs can regulate chromatin condensates indirectly through interactions with proteins including HP1α, linker histone, and BRD4, they also directly regulate chromatin (7, 35, 36). Charge-modulating PTMs of histone tail residues, including acylation of lysine and citrullination of arginine, alter chromatin compaction and are poised to disrupt the multivalent electrostatic interactions between histone tails and DNA that drive nucleosome phase separation (37, 38). In initial investigations into the direct effects of histone PTMs, non-specific hyperacetylation of histones by p300 led to the dissolution of nucleosome array condensates (7). While both lysine and arginine carry a single positive charge at neutral pH, they differ in their interactions with DNA. Arginines are generally thought to form stronger interactions with DNA due to more favorable enthalpic contributions with the phosphate backbone, a greater hydrogen-bonding capacity, a greater propensity for cation-π interactions, and a lower energetic cost of desolvation (39–41). Greater compaction and condensation are observed with arginine-rich than lysine-rich peptides (42–47). The more favorable interaction of arginines with DNA is also observed in simulations of NCPs and nucleosomes (32, 48, 49). What remains unclear is how the position-

specific contributions of lysine and arginine residues within histone tails, and their neutralization
by PTMs, influence nucleosome condensate formation and dynamics.
Here, we begin to systematically dissect the influence of lysines and arginines and their
charge-neutralizing modifications by investigating the role of basic residues of the H3 tail, the
longest of the core histone tails (24), on nucleosome phase separation. In the case of the H3
tail, six lysines and four arginines are distributed along its length with the potential to contribute
to phase separation via inter-nucleosomal interactions with DNA. We investigate the position-
dependent effect of perturbing lysines or arginines in the terminal or central regions on
nucleosome phase separation. Basic residues are perturbed in these regions via neutralizing
mutations to glutamine as mimetics for acetylation and citrullination, or by converting all basic
residues to a single type (i.e., either lysines or arginines). Using microscopy-based techniques
in conjunction with simulations, we demonstrate that the number and position of lysine and
arginine neutralizations within the H3 tail determine nucleosome phase boundaries and the
viscosity of their condensates. NMR dynamics experiments show minimal differences in
picosecond-to-nanosecond timescale dynamics of the H3 tail within condensates and display a
correlation between condensate viscosity and H3 tail dynamics. Together, these data contribute
to a model wherein the intrinsic properties of nucleosomes and their condensates are involved
in defining the local chromatin microenvironment, which can be tuned via position-specific
charge-modulating histone PTMs.
99
100
101 **MATERIALS AND METHODS**
102
Expression and purification of full-length histones. Full-length human histone sequences
were expressed and purified as previously described (16, 50). These protein sequences
correspond to UniProt sequences P0C0S8, P62807, Q71DI3 (with C110A), and P62805. Briefly,
each histone sequence was expressed from pET3a vectors using T7 Express lysY competent
E. coli (New England Biolabs) or Novagen Rosetta 2(DE3) pLysS competent *E. coli* (Millipore
Sigma). Mutants were generated via Q5 Site-Directed Mutagenesis Kit (New England Biolabs).
The histone H3 constructs used in this study were: unmodified, R2Q, R8Q, R2/8Q (2xR-
$Q_{terminal}$), R17/26Q (2xR-$Q_{central}$), R2/8/17/26Q (4xR-Q), K4/9Q (2xK-$Q_{terminal}$), K14/18/23/27Q
(4xK-$Q_{central}$), K4/9/18/27Q (4xK-Q_{even}), K4/9/14/18/23/27Q (6xK-Q), R2/8/17/26K (All-K),
K4/9/14/18/23/27R (All-R), and tailless [here, referring to H3(28-135)]. Briefly, histones were
extracted from inclusion bodies and purified in Urea Buffer (7 M urea, 20 mM sodium acetate pH
5.2, 100 mM NaCl, 5 mM β-ME, 1 mM EDTA) sequentially over Q-Sepharose Fast Flow resin
(GE Healthcare) and SP-Sepharose Fast Flow resin (GE Healthcare) or Proteindex IEX-CM
Agarose 6 Fast Flow resin (Marvelgent Biosciences). Pure histones were dialyzed into ddH$_2$O
with 1 mM β-ME and stored lyophilized at -20°C until needed for nucleosome reconstitution. ESI
mass spectrometry (Finnigan LTQ, Thermo Electron Corporation, San Jose, CA) was used for
histone quality assessment based on molecular weight and used to ensure no unintended
modifications, such as carbamylation.
121
Amplification and purification of 147 bp Widom 601 DNA. DNA used in both NCP
reconstitution and peptide-DNA assays was obtained using a plasmid with 32 repeats of the
147-bp Widom 601 sequence (ATCGAGAATC CCGGTGCCGA GGCCGCTCAA
TTGGTCGTAG ACAGCTCTAG CACCGCTTAA ACGCACGTAC GCGCTGTCCC
CCGCGTTTTA ACCGCCAAGG GGATTACTCC CTAGTCTCCA GGCACGTGTC
AGATATATAC ATCCGAT). Bacterial amplification and purification of 147-bp DNA was
performed as previously described (16, 50). Following transformation using NEB 5-alpha
Competent *E. coli* (New England Biolabs), the bacteria were grown in 2xYT media pH 7.5. The
plasmid was purified using an alkaline lysis method and phenol-chloroform extractions. The

147-bp fragments were released via digestion by EcoRV-HF enzyme (New England Biolabs)
and purified from the vector using PEG precipitation and HiTrap DEAE Fast Flow Columns (GE
Healthcare) via FPLC (Bio-Rad). DNA purity was assessed using 5% native-PAGE gel
electrophoresis. The 147-bp DNA fragments were stored following ethanol precipitation at -
20°C.

Nucleosome core particle reconstitution. NCP reconstitutions were performed largely as
previously described (16, 51). Briefly, histones H2A, H2B, H3, and H4 are individually
resuspended at 2 mg/mL in Unfolding Buffer (6 M guanidine-HCl, 20 mM Tris pH 7.5, 5 mM
DTT). Most histone complexes were refolded as an H2A/H2B dimer and H3/H4 tetramer, except
for complexes with K4/9/14/18/23/27R-H3 and tailless-H3, which were refolded as an
H2A/H2B/H3/H4 octamer. All complexes were refolded at 1 mg/mL histone via dialysis into a
High Salt Buffer (2 M KCl, 20 mM Tris pH 7.5, 1 mM EDTA, 0.5 mM benzamidine, 1 mM DTT).
Each histone complex was purified via FPLC using a HiLoad 16/600 Superdex 200 column (GE
Healthcare), and purity was assessed using 18% SDS-PAGE gel electrophoresis. Prior to the
final NCP reconstitution, purified 147-bp Widom 601 DNA was resuspended in High Salt Buffer
at a final concentration of 8-11 µM. The final NCP reconstitution was performed at a molar ratio
of 1:1:2.2 or 1:1:2 in terms of 147-bp Widom 601 DNA:H3/H4 tetramer:H2A/H2B dimer. For
samples with K4/9/14/18/23/27R-H3 and tailless-H3, the final NCP reconstitution was performed
at 1:1:0.2 molar ratio of 147-bp Widom 601 DNA:H2A/H2B/H3/H4 octamer:H2A/H2B dimer. The
NCP reconstitution was performed via slow desalting with an exponential gradient followed by
overnight dialysis into No Salt Buffer (20 mM Tris pH 7.5, 1 mM EDTA, 0.5 mM benzamidine, 1
mM DTT). NCPs were purified via a 10-40% sucrose gradient (BioComp Instruments). NCP
purity was assessed via sucrose gradient profile (absorbance at 260 nm) and 5% native-PAGE.
Additionally, histone integrity was assessed using 18% SDS-PAGE prior to use in experiments.

Expression and purification of H3 tail peptides. Unmodified H3(1-44) was used as an
extended histone tail peptide sequence. The following H3(1-44) mutants were also used:
R2/8Q, R17/26Q, R2/8/17/26Q, K4/9Q, K14/18/23/27Q, K4/9/18/27Q, K4/9/14/18/23/27Q,
R2/8/17/26K, and K4/9/14/18/23/27R. Mutant H3 tail peptides were generated using Q5 Site-
Directed Mutagenesis Kit (New England Biolabs). Peptides were expressed and purified as
previously described (16). Briefly, each peptide was expressed from a pET3a vector in
BL21(DE3) competent *E. coli* (New England Biolabs). Following cell lysis, the H3 tail peptides in
the soluble portion were purified using Q-Sepharose Fast Flow resin (GE Healthcare) and a
Source 15S column (GE Healthcare) via FPLC (Bio-Rad). Following lyophilization, peptides
were purified in S30 Buffer (50mM KPi pH 7, 50mM KCl, 0.5mM EDTA) via FPLC using a
HiLoad 16/600 Superdex 30 pg column (GE Healthcare). Peptides were further purified using a
PROTO 300 C18 column (Higgins Analytical) via HPLC (Shimadzu). ESI mass Spectrometry
(Finnigan LTQ, Thermo Electron Corporation, San Jose, CA) was used to confirm intact masses
for H3 tail peptides. Peptides were lyophilized and stored at -20°C prior to experiments.

Preparation of plates for phase separation assays. Prior to phase separation assays, each
multiwell plate was washed with 1% Hellmanex at 30°C for 40 min, washed extensively with
autoclaved ddH$_2$O, and left to air dry. Each plate was then treated with Sigmacote (Sigma-
Aldrich) for 10 min, washed extensively with water, and left to air dry.

NCP stock preparation for phase separation assays. NCP stocks were prepared in MOPS
Buffer [20 mM MOPS pH 7 (8 mM NaOH to pH), 0.1 mM EDTA] using a 10 kDa MWCO
centrifugal filtration unit (Millipore Sigma) and were stored at 4°C for a minimum of 12 hours
before checking stock concentration. To determine stock concentration, a secondary NCP stock
was made into MOPS Buffer. NCP concentration was determined spectrophotometrically from a

1:100 dilution of the secondary stock into 2 M KCl using a NanoDrop (Thermo Fisher).
Concentration was calculated using the 147-bp Widom 601 DNA extinction coefficient at 260 nm
(ε_{260} = 2,312,300.9 M^{-1}cm^{-1}). The primary stock concentration was then back-calculated.
185
Turbidity assay. All components [NCP, H3(1-44), 601 DNA, MgCl$_2$ stock, and KCl stock] were
prepared in MOPS buffer [20 mM MOPS pH 7 (with 8 mM NaOH to pH), 0.1 mM EDTA] and
warmed to room temperature before conducting each assay. Final buffer conditions were 20
mM MOPS pH 7 (8 mM NaOH to pH), 0.1 mM EDTA, 150 mM KCl, and 2 mM MgCl$_2$. Note that
the addition of 2 mM MgCl$_2$ is distinct from our previously reported phase separation assays
(16). Most assays were run in µClear 96-Well Half Area Microplates (Greiner Bio-One) sealed
with Optical Adhesive Film (Thermo Fisher Scientific). Absorbance measurements were
recorded using a FlexStation 3 plate reader (Molecular Devices) at 600 nm (A$_{600}$) for 60 minutes
at one-minute intervals. Data were recorded with a set temperature of 25 °C, which had an
actual experimental range of 24.9-25.6 °C (average with standard deviation of 25.3 $\pm$ 0.1 °C for
NCP samples). The maximum A$_{600}$ value measured within the first 10 minutes of the time course
was used in subsequent analyses. Data were plotted using Igor Pro (Wavemetrics) and R
R Statistical Software (v4.2.2; R Core Team 2022).
For assays with NCP, components were mixed in the order of MOPS buffer, KCl, MgCl$_2$,
and with the final addition of concentrated NCP stock to initiate the reaction. Experiments were
conducted at NCP concentrations of 0, 10, 25, 50, 100, 200, and 300 µM. A total of three
replicates were collected for each sample, with varied NCP concentration order between
replicates. Assays were conducted at a final volume of 36.5 µL in µClear 96-Well Half Area
Microplates (Greiner Bio-One) or at a final volume of 75 µL (for matched sample height) in
µClear 96-Well Full Area Microplates (Greiner Bio-One).
For H3 tail and DNA, components were mixed in the order of MOPS buffer, KCl, MgCl$_2$,
147 bp 601 DNA, and with the final addition of concentrated H3(1-44) stock to initiate the
reaction. Assays were performed at 50 µM H3 tail peptide with increasing DNA concentrations
(0, 0.625, 1.25, 2.5, 5, 10, 12.5, 15, 25 µM) and a final volume of 20 µL in each well.
210
Brightfield microscopy. Images were acquired using Cytation10 Confocal Imaging Reader
(Agilent) in brightfield mode with a 60x objective, collecting from -30 µm to 200 µm in 2 µm
steps. Samples were prepared in a similar manner to those for turbidity assays. All components
[NCP, H3(1-44), 601 DNA, MgCl$_2$ stock, and KCl stock] were prepared in MOPS buffer [20 mM
MOPS pH 7 (with 8 mM NaOH to pH), 0.1 mM EDTA] and warmed to room temperature before
conducting each assay. Final assay buffer conditions were 20 mM MOPS pH 7 (8 mM NaOH to
pH), 0.1 mM EDTA, 150 mM KCl, and 2 mM MgCl$_2$. Assays used µClear 96-Well Half Area
Microplates (Greiner Bio-One) sealed with Optical Adhesive Film (Thermo Fisher Scientific).
Final experimental well volumes were 20 or 36.5 µL. The microscopy images were collected at
room temperature with an experimental range of 27.4-28.2 °C for peptide-DNA assays and
27.4-29.0 °C (average with standard deviation of 28.0 $\pm$ 0.3 °C) for NCP assays.
The NCP was pipetted and imaged in two ways in order to capture faster-settling
condensates. The first method pipetted concentrations 0 to 300 µM in order and imaged from 0
to 300 µM. The second method pipetted concentrations 0 to 300uM in order and imaged from
300 to 0 µM. NCP microscopy replicates were conducted with n = 4 for unmodified and n = 3
for all mutants, which includes two biological replicates for all constructs except the two single
mutants (R2Q and R8Q). Additional concentrations were filled in at 5 or 20 µL volumes in
µClear 384-well small volume LoBase or 96-Well Half Area Microplates (Greiner Bio-One),
collected in triplicate with a single biological replicate. At 300 µM NCP, phase-separated
samples of R2/8/17/26Q-H3-NCP separated into two distinct layers by the time automated data
collection began, with the lack of visible droplets falsely suggesting a lack of phase separation.

Thus, for this mutant at 300 µM, an additional single-well experiment was collected to shorten
the experiment dead time as much as possible. This single-well imaging is used in **Figure 1D**.
Images for figures were made using Gen5 (Agilent). Images were further analyzed using
ImageJ version 2.1.0/1.54f. In an attempt to objectively define the presence of droplets in
brightfield images, a pseudo-coefficient of variation (CV) was calculated. The total pixel intensity
values of a standard deviation z-projection and an average intensity z-projection (from 0 to 200
µm) were determined for each NCP concentration. The reported CV was taken as the ratio of
the intensities of the standard deviation projection and the average intensity projection,
multiplied by 100. The CV values were averaged across replicates and plotted (**Supplementary
Figure 4A**) using R Statistical Software (v4.2.2; R Core Team 2022). A CV cutoff of 5.0 was
used to delineate phase separation. These values agree with a visual assessment of droplets.
For H3 tail peptide and DNA, only one set of microscopy images was obtained, which were
visually assessed for droplets.
245
Fluorescence microscopy assay. For fluorescence imaging of NCP samples, YOYO-1 dye
(Invitrogen) was added to NCP stocks at a 1:100 molar ratio with a minimum incubation time of
16 hours prior to inducing phase separation and imaging. Samples were prepared at 200 µM
NCP in a final volume of 4 µL. Final buffer conditions were 20 mM MOPS pH 7 (8 mM NaOH to
pH), 0.1 mM EDTA, 150 mM KCl, and 2 mM $MgCl_2$. Samples were imaged at room temperature
in µClear 384-well small volume LoBase microplates (Greiner Bio-One) sealed with optical
adhesive film (Thermo Fisher Scientific). Microscopy was performed on a Nikon Eclipse Ti2-E
inverted spinning disc microscope equipped with a Yokogawa confocal scanner unit (CSU-W1),
4 laser lines (405/488/568/633), a Hamamatsu ORCA-Flash4.0 V3 sCMOS Camera system with
an 82% quantum efficiency chip and a water immersion dispenser. The system is run using the
NIS-Elements Advanced Research package software (version X). DIC and confocal images
were collected with a z-step size of 2 µm using a 60x water objective (SR Plan Apo IR). The
field of view for the camera is 2048 x 2044 pixels with resolution 0.16 µm/pixel for the 60x water
objective. Images were processed using Fiji and ImageJ version 2.1.0/1.54f.
260
NCP sample recovery, quality assessment, and re-use. Following assays with non-
fluorescently labeled NCPs, samples were pooled and diluted to reduce salt and dissipate NCP
phase separation. Both pre-assay and post-assay NCP samples were run on 5% native-PAGE
and 18% SDS-PAGE. Samples were then concentrated and exchanged into MOPS buffer using
a 10 kDa MWCO centrifugal filtration unit (Millipore Sigma). Following gel confirmation and
exchange, the samples were reused in replicate phase separation assays as done previously
(16).
268
Video particle tracking (VPT) microrheology of NCP samples. VPT measurements were
performed on phase-separated samples of NCP variants prepared at a final NCP concentration
of 200 µM in 20 mM MOPS pH 7 (8 mM NaOH to pH), 0.1 mM EDTA, 150 mM KCl, and 2 mM
$MgCl_2$. Yellow-green carboxylate-modified polystyrene beads of 1 µm size were mixed during
sample preparation. These beads were confirmed to partition into the dense phases of phase-
separated NCP samples (**Supplementary Video 1**) and were used as probe particles for VPT
measurements. The samples of volume 5 or 10 µL were sandwiched between a glass slide and
coverslip treated with 5% w/v PEG in 5% ethanol. The samples were surrounded by mineral oil
to prevent evaporation. The prepared sample sandwich was placed on a temperature-controlled
stage (Instec) attached to a Zeiss Primovert inverted microscope. Temperature was controlled
using the thermal stage, while the actual temperature was determined using a thermocouple.
The samples were allowed to fuse before imaging and were rested on the microscope for 30-60
minutes before data collection. The bead motion inside the NCP dense phases was captured at
24 °C in the form of videos using a 40x air objective lens and Teledyne FLIR Blackfly S USB3

CMOS camera for 1000 frames with 100 ms exposure time. The acquired videos were
processed using Fiji (version 1.54f) (52). The probe particles were tracked to get their
trajectories using the Trackmate (53) software plugin in Fiji. During tracking, intensity and quality
filters were used to avoid tracking aggregated particles.
The analysis routine for estimating the mean square displacement (MSDs) from particle
trajectories is detailed in our previous work (54, 55). In brief, the center of the mass trajectory of
the bead motion was utilized to correct the bead trajectories for any possible drift during video
acquisition. The center of the mass trajectory was estimated from the velocities of the bead
using

$$X_{COM}(k) = X_0 + \sum_{j=0}^{k} \frac{1}{N_j} \sum_{i=1}^{N} v_{i,j} \tag{1}$$

292 where X_{COM} is the center of mass vector calculated for k_{th} frame, j is the individual frame, and
293 N_j is the number of particles in the frame j. X_0 is the center of the mass vector of the first frame.
294 After correction, the obtained position vector of the particle R can be used to estimate the MSDs
295 using

$$MSD(\tau) = \langle R(t + \tau) - R(t) \rangle_{t,N} \tag{2}$$

296 where τ is the lag time. We used our custom Python scripts, also deposited and can be found
297 on GitHub (https://github.com/BanerjeeLab-repertoire/Material-properties), to compute MSD
298 across various lag times τ. The ensemble average MSD was then determined based on data
299 from approximately 20–50 individual beads. The MSD was then fitted (54, 55) using
300

$$MSD(\tau) = 4D\tau^\alpha + N \tag{3}$$

301 to obtain the diffusion coefficient D of the beads. Here, α is the diffusivity parameter which
302 signifies the nature of bead diffusion inside the dense phases of NCP samples. The ensemble-
303 average MSDs for which the fitting yielded $\alpha=1$ were used to estimate the viscosity η using the
304 Stokes-Einstein equation (54, 55)

$$\eta = \frac{k_B T}{6\pi DR} \tag{4}$$

where κ_B is the Boltzmann constant, R is the radius of the particles, D is the diffusion coefficient
of beads obtained by fitting the ensemble-average MSDs to Eq. (3) and T is the temperature of
the sample.
The videos were recorded at 9-10 randomly chosen positions of the dense phases of
NCP samples, and data were collected on three technical replicates for each NCP mutant.
Statistical analyses were performed using R Statistical Software (v4.2.2; R Core Team 2022).
To determine variance within viscosity measurements, a one-way ANOVA was initially used.
Measurements obtained were determined to be significant with a p-value of 1.11E-64 (n=3). To
determine which NCP viscosity measurements are significant to one another, a post-hoc
Tukey's test was used where a p-value of < 0.05 was deemed significant for the pairwise
comparisons between the NCPs. Data were plotted using R.
316
NMR sample preparation. Purified NCPs reconstituted with [15]N-H3 (Unmodified, 4xR-Q, All-K,
or All-R) were exchanged into MOPS buffer [20 mM MOPS pH 7 (with 8 mM NaOH to pH), 0.1
mM EDTA] using consecutive rounds of concentration and dilution in 10k MWCO centrifugal
filtration devices (Millipore Sigma). NCP stock concentration was determined
spectrophotometrically via absorbance at 260 nm after diluting into 2 M KCl.

Phase separation was induced by diluting to 400 μM NCP in final solution conditions of
18 mM MOPS pH 7 (with 7.2 mM NaOH to pH), 0.09 mM EDTA, 150 mM KCl, 2 mM MgCl$_2$, and
10% D$_2$O. The sample was transferred to a 1.7 mm NMR tube (Bruker) and left to separate into
two layers overnight. The top (light phase) layer was then transferred to a second 1.7 mm NMR
tube. Care was taken to ensure the NMR coil volume was filled with either dense or light phase.
327
NMR data collection. Data were collected on a Bruker Avance III HD 600 MHz spectrometer
with a 1.7-mm cryogenic probe at 298 K and running Topspin 3.5pl7. For chemical shift
analysis, ^{1}H-^{15}N HSQC spectra were collected with 16 scans, 1024 (^{1}H) x 300 (^{15}N) total points,
acquisition times of 51 ms (^{1}H) and 75 ms (^{15}N), and spectra widths of ~17 ppm (^{1}H) and ~33
ppm (^{15}N). All nuclear spin relaxation experiments were collected with 1600 (^{1}H) x 228 (^{15}N) total
points, acquisition times of 102 ms (^{1}H) and 84 ms (^{15}N), and spectra widths of 13 ppm (^{1}H) and
22 ppm (^{15}N). {^{1}H}-^{15}N steady-state heteronuclear nuclear Overhauser effect (hnNOE) data
were collected using the Bruker pulse sequence hsqcnoef3gpsi with the reference and
saturated spectra interleaved. Experiments were collected with 32 (for dense phase samples),
72 (for 4xR-Q-H3 light phase sample), 88 (for All-R light phase), or 96 (for unmodified-H3 and
All-K-H3 light phase samples) scans and an interscan delay of 5 s. The dense phase hnNOE
experiments were collected at the start of the experimental queue and again 13-18 days later to
check for sample aging. The data were unchanged within experimental error, and data from the
initial hnNOE experiment were used in figures. Longitudinal (R1) ^{15}N relaxation experiments
were collected with total relaxation loop lengths of 0.033 s, 0.195 s, 0.390 s, 0.585 (x2), 0.813 s,
1.301 s, and 1.951 s in randomized order. Experiments were collected with 24 (for dense phase
samples), 32 (for R2/8/17/26Q-H3 light phase sample), or 40 (for unmodified-H3 and
R2/8/17/26K-H3 light phase samples) scans and an interscan delay of 2 s. Transverse (R2) ^{15}N
relaxation experiments were collected with an effective CPMG field of 500 Hz and total
relaxation CPMG loop lengths of 0.0044 s, 0.013 s, 0.026 s, 0.044 s (x2), 0.062 s, 0.084 s, and
0.110 s. Experiments included temperature compensation blocks and were collected in
randomized order, with the same number of scans as for R1, and with an interscan delay of 2.5
s. ^{1}H-^{15}N HSQC spectra were collected between data series to confirm sample integrity. For
^{15}N-All-R-H3-NCP, 1D versions of the R1 and R2 experiments were collected with 256 scans at
the same delay times as for the 2D versions.
353
NMR data analysis. NMRPipe and CcpNmr Analysis 2.5.2 were used within NMRbox for
processing and analysis, respectively, of NMR data (56–58). Data were plotted using Igor Pro
(Wavemetrics). Amide assignments for NCPs with unmodified H3 are from BMRB entry 50806.
These assignments were transferred to H3 mutants via inspection and comparison to previously
published spectra (59). Chemical shift differences ($\Delta\delta$) between solution phases were calculated
from the differences in amide proton ($\Delta\delta_H$) and amide nitrogen ($\Delta\delta_N$) chemical shifts using:
$\Delta\delta = \sqrt{(\Delta\delta_H)^2 + (0.154\Delta\delta_N)^2}$ (60). Spectra were processed using a cosine-squared window
function for both dimensions, and each dimension was doubled in size by zero-filling twice and
rounding to the nearest power of two. HnNOE values for each residue were calculated as the
peak height ratio between the saturated and reference spectra, and errors were calculated via
standard error propagation of the spectral noise as determined by CcpNmr Analysis 2.5.2. R1
and R2 relaxation rates were determined by fitting peak heights from the relaxation series to a
single-exponential decay without offset. Errors were determined from the covariance matrix
within CcpNmr Analysis 2.5.2. As discussed for previous analyses, there is the caveat resulting
from the low chemical shift dispersion of IDRs that relaxation rates can be convolution by
nearby peaks in dense spectral regions (59). For the ^{1}H-1D versions of R1 and R2 experiments,
the amide backbone region was integrated in CcpNMR Analysis 3.3.2.1 and fit using Igor Pro

(Wavemetrics) to a single-exponential decay without offset where the reported error is the
estimated standard deviation of the fit coefficient.
373
Coarse-grained molecular simulations of H3 tail with DNA fragments. We employed a one-
bead hydrophobicity scale (HPS) model to calculate the phase diagram and viscosity of H3 tail-
DNA co-condensates. The H3 tail sequence is the same as the peptide tested experimentally
(ARTKQTARKSTGGKAPRKQLATKAARKSAPATGGVKKPHRYRPG). A 20-thymine (T) strand
complementary base-paired with adenine (A) was used as the DNA sequence. The two-
component system comprises double-stranded DNA and H3 peptides with 40 and 44 beads,
respectively. The beads in the DNA chain represent adenine-base (A) and thymine-base (T)
nucleotides, and the beads in the peptide represent amino acids in H3(1-44). The non-bonded
interactions between residues i, j are modeled via the Ashbaugh-Hatch potential, and salt-
screened electrostatic interactions are modeled via the Debye-Hückel potential:

$$U_{ij}^{AH}(r) = \begin{cases} U^{LJ}(r) - \lambda_{ij}U^{LJ}(r_c) + \epsilon(1 - \lambda_{ij}), & r \le 2^{1/6}\sigma_{ij} \\ \lambda_{ij}[U^{LJ}(r) - U^{LJ}(r_c), & 2^{1/6}\sigma_{ij} < r \le r_c \\ 0, & r > r_c \end{cases}$$

$$U_{ij}^{DH}(r) = \frac{q_i q_j e^2}{4\pi\epsilon}\frac{\exp(-\kappa r)}{r}$$

where ϵ = 0.8368 kJ/mol, r_c = 4 nm, and U^{LJ} is the Lennard-Jones potential $U^{LJ}(r) = 4\epsilon[(\frac{\sigma_{ij}}{r})^{12} -$
$(\frac{\sigma_{ij}}{r})^6 - \sigma_{ij} = (\sigma_i + \sigma_j)/2$ and $\lambda_{ij} = (\lambda_i + \lambda_j)/2$ are arithmetic averages of monomer size and
hydrophobicity value, respectively. The pair potential with $\lambda = 0$ consists of only the repulsive
term equivalent to the Weeks-Chandler-Andersen functional form. We use σ values of amino
acids from van der Waals volumes and λ values of amino acids from the recently proposed
CALVADOS 2 parameters (61). In the Debye-Hückel potential, q is the charge number of
particles and $\kappa^{-1} = \sqrt{\varepsilon k_B T/2e^2 I}$ is the Debye length. In our simulations, the ionic strength of the
electrolyte I = 0.17 M. The temperature-dependent dielectric constant used in our simulations
has the following empirical relation: $\frac{\varepsilon(T)}{\varepsilon_0} = \frac{5321}{T} + 233.76 - 0.9297T + 1.41710 \times 10^{-3}T^2 -$
$8.29210 \times 10^{-7}T^3$. All coarse-grained beads are connected via bonded interactions modeled by
harmonic potentials $U^{bond}(r) = \frac{k}{2}(r - r_0)^2$ where k = 1000 kJ/mol/nm^2, r_0 = 0.5 nm for DNA, and
r_0 = 0.38 nm for peptide. In dsDNA, there are additional base pair interactions and stacking
interactions between adjacent base-pair groups. The base pair interaction is represented by a
repulsive Morse potential $U_{bp} = Dexp(-\alpha(r - l_0))$ with D = 472 kJ/mol, α = 2.876 nm^{-1} and
$l_0 = 1.243$nm. The stacking interaction is modeled using a Lennard-Jones potential,
characterized by ϵ = 49.2 kJ/mol, σ = 0.2938 nm, and a cut-off distance r_c = 5.65 nm. We have
applied OpenMM (v7.7) for H3 tail-DNA condensate simulations.
We performed coexistence simulations in a slab geometry at different T for
characterizing their phase diagrams. The number of dsDNA was set at N_{DNA} = 180. The number
of H3 tail peptides was set around N_{H3} = 520 – 720. These simulations were carried out at each
T for a duration of 5 × 10^7 – 7 × 10^7 steps, and the dense phase and dilute phase
concentrations were computed based on the last 2 × 10^7 steps of the simulation trajectory. We
estimated the critical temperatures by fitting the phase diagram using the law of coexistence
densities and the critical concentrations by assuming the law of rectilinear diameter holds.
We ran condensate simulations in a cubic box for viscosity measurement. The number
of dsDNA was set around N_{DNA} = 84-180. The number of H3 tail peptides was set around N_{H3} =
254 – 426 to keep the approximate charge neutrality of the co-condensates. The total number of
the CG monomers was kept at about 2 × 10^4 to eliminate the finite-size effect. The initial length

of the cubic box was set at 40 nm, which is much larger than the equilibrium size. The box was
then shrunk in the subsequent NPT simulations. After energy minimization, we used the NPT
ensemble with a timestep of 0.01 ps, to equilibrate all systems using an additional 5×10^7 steps.
Finally, we fixed the box size as the average volume in NPT simulations and carried out
additional NVT simulations for 1×10^7 steps.
418
419
420   **RESULTS**
421
Design of H3 tail basic residue mutants to probe nucleosome phase separation.
We set out to investigate the role of charge neutralization of H3 tail basic residues and
the distribution of these neutralizations along the tail on nucleosome phase separation. Charge-
modulation of the histone tails is expected to perturb the balance between multivalent intra- and
inter-nucleosomal tail-DNA interactions (24) and, in turn, phase separation propensity and
condensate properties. Furthermore, the most N-terminal basic residues in the tail are
anticipated to have greater potential to form inter-nucleosomal interactions and facilitate phase
separation than the basic residues in the center of the tail sequence.
Here, we focus on six lysine and four arginine residues within the H3 tail that can
promote phase separation through inter-nucleosomal electrostatic interactions with DNA. To
examine the position-dependent effects of perturbing these basic residues, we divided the H3
tail into three regions: the **'terminal' region**, comprising residues preceding the first TGG motif;
the **'hinge' region**, corresponding to the uncharged segment that contains the second TGG
motif and precedes the histone core; and the **'central' region**, spanning between the 'terminal'
and 'hinge' regions (**Supplementary Fig. 1**). Neutralizing mutations of basic residues to
glutamine in the terminal and central regions were introduced as PTM mimetics of arginine
citrullination and lysine acetylation. We designed H3 mutants (**Supplementary Figure 1**) to
separately neutralize arginines and lysines within the terminal region (R2/8Q, referred to as
2xR-Q$_{terminal}$, and K4/9Q, referred to as 2xK-Q$_{terminal}$) and central region (R17/26Q, referred to as
2xR-Q$_{central}$, and K14/18/23/27Q, referred to as 4xK-Q$_{central}$). Additionally, we designed H3
mutants with neutralization of the two terminal region arginines individually (R2Q and R8Q), all
four arginine residues (R2/8/17/26Q, referred to as 4xR-Q), the four lysine residues adjacent to
arginine residues (K4/9/18/27Q, referred to as 4xK-Q$_{even}$), and all six lysine residues across the
terminal and central regions (K4/9/14/18/23/27Q, referred to as 6xK-Q). To test the importance
of H3 tail basic residue identity, we also designed two mutants to convert all basic residues
across the terminal and central regions of the H3 tail to only one type: R2/8/17/26K-H3 (All-K-
H3) and K4/9/14/18/23/27R-H3 (All-R-H3).
449
Sequence modifications to the H3 tail modulate tail-DNA condensate formation.
To probe the effect of H3 tail modifications, we started with a simplified two-component
system containing purified H3 tail peptide [residues 1-44, H3(1-44)] and the 147-bp Widom 601
DNA sequence (**Figure 1A-B**). Previously, we demonstrated that the phase separation behavior
of H3 tail peptide with DNA is similar to the reentrant condensation observed for other systems
with multivalent interactions between nucleic acids and arginine- and lysine-rich intrinsically
disordered peptides (16, 44, 45, 62–64). Using microscopy and turbidity measurements, we
determined the phase boundary for this two-component system and compared it to the
estimated charge neutralization ratio for H3(1-44) mutants and 147-bp Widom DNA, which is
predicted to display the maximum turbidity if the system follows complex coacervation (**Figure
1A, 1C, 1D**). Overall, increasing the number of neutralizations narrows the condensate
formation regime with a subtle shift in maximum turbidity to lower ratios of DNA to peptide
(**Figure 1C-D, Supplementary Figure 2)**. Phase separation near the estimated charge
neutralization ratio is seen for all H3 tail peptides except with 4xR-Q and 6xK-Q mutations, for

which no phase separation is observed. In contrast to 4xR-Q-H3(1-44), the two 4xK-Q mutants
phase separate with DNA, supporting a differential effect between modifications to arginine and
lysine. The similar effect of 2xR-Q$_{terminal}$ and 2xR-Q$_{central}$ mutations suggests no clear position
dependence within this simplified two-component system, although a mild narrowing of the
phase boundary is observed between the two 4xK-Q mutants. Unmodified-, All-K-, and All-R-
H3(1-44) have the same net charge and display similar phase separation, but All-R-H3(1-44) is
distinct in showing a broader transition via turbidity (**Figure 1C-D**, **Supplementary Figure 2A**).
These results suggest a differential impact between arginines and lysines in driving condensate
formation via H3 tail-DNA interactions, as well as a role for neutralization modifications in
regulation of these condensates.

Figure 1. Phase separation of H3 tail peptide with DNA is modulated by perturbation of basic residues. A. Mutants of H3(1-44) are summarized and shown explicitly in **Supplementary Figure 1**. The net charge and estimated charge-neutral molar ratio of peptide:DNA are tabulated for each mutant. **B.** Sequence of H3 peptide, H3(1-44), which includes the H3 tail. Arginine and lysine residues are highlighted in shades of dark and light blue, respectively. **C.** Turbidity (absorbance at 600 nm) values (average of n=3) are shown in greyscale for unmodified and mutant versions of H3(1-44) at 50 µM with increasing amounts of 147-bp Widom 601 DNA, shown as molar ratio. Turbidity is plotted as a heatmap ranging from 0.00 (white) to 0.27 (black). Values greater than 0.03 are outlined with a black square to indicate light scattering due to droplet formation. Data were recorded with a set temperature of 25 °C, which had an actual experimental range of 24.9-25.6 °C. **D.** Brightfield microscopy images were collected at the same molar ratios as in (**C**). The scale bar is 25 µm. Images are outlined in black to indicate droplets. See **Supplementary Figure 2** for an alternative representation of turbidity data and larger images. Microscopy images were collected at room temperature with an experimental range of 27.4-28.2 °C. Construct identity in (**C**) and (**D**) aligns with the table in (**A**).

490
491
Nucleosome core particle phase boundary shifts with basic residue neutralization in a position-dependent manner.

493   We next asked whether this series of H3 modifications influences phase separation within the context of the nucleosome, where the H3 tails are only two of ten tails and are tethered to the core in close proximity to DNA. In this situation, we predicted a positional dependence of basic residue neutralization. To probe the effect of charge-neutralizing mutations of H3 tail basic residues on NCP phase separation (**Figure 2A-B**), we started by determining the saturation concentration (c_{sat}) of each mutant-H3-NCP using turbidity and brightfield microscopy. These assays were conducted near room temperature (25.3 $\pm$ 0.1 °C and 28.0 $\pm$ 0.3 °C, respectively) and at physiologically relevant salt concentrations of 150 mM KCl and 2 mM MgCl$_2$, for a range of discrete NCP concentrations from 0 to 300 µM.

503   Unmodified-NCP was used as the point of comparison for all mutants. Droplets were observed via brightfield microscopy at unmodified-NCP concentrations of 50 µM and greater, which is also in agreement with sample turbidity (**Figure 2C-D**, **Supplementary Figures 3-4**, **Supplementary Tables 1-2**). Note that this boundary is shifted to lower concentrations from our previous publication due to the inclusion of MgCl$_2$ (16). In agreement with previous results acquired in the absence of MgCl$_2$ (16), joint removal of the terminal and central regions of the H3 tail (referred to as tailless-H3-NCP) abrogates phase separation at concentrations up to at least 300 µM (**Figure 2C-D**). We interpret these results to support the importance of the H3 tail in NCP phase separation under the experimental conditions tested.

512   A comparison of arginine and lysine mutants provides insight into the influence of charge neutralization of H3 tail basic residues on NCP phase separation as a function of identity, number, and position (**Figure 2C-D**, **Supplementary Figures 3-4**). When comparing arginine neutralizations, a dependence on number and position is observed: 2xR-Q$_{central}$-H3-NCP has similar c_{sat} (50 µM NCP) as compared to unmodified-NCP while 2xR-Q$_{terminal}$ and 4xR-Q shift the c_{sat} to higher concentrations (100 and 200 µM NCP, respectively). To further break down the effect of 'terminal' region arginines, we individually neutralized R2 and R8. R2Q produces a more pronounced shift in c_{sat} than R8Q (100 and 75 µM NCP, respectively) and is similar to the effect of the double neutralization. Lysine neutralizations exhibit a dependence on number: 2xK-Q$_{terminal}$ shifts c_{sat} to 100 µM NCP while 4xK-Q$_{central}$ and 4xK-Q$_{even}$ shift c_{sat} to 200-300 µM NCP. The effects on c_{sat} are similar whether neutralizing two arginines or two lysines in the 'terminal' region; the effects of 'central' region neutralizations are confounded by the difference in the number of arginines and lysines (i.e., 2 vs. 4) in the 'central' region. When all six H3 tail lysines are neutralized (6xK-Q), we do not observe phase separation at the concentrations tested. A comparison of the conversion of all basic residues across the terminal and central regions of the H3 tail to only one type suggests that c_{sat} decreases to 15 and 35 µM for All-K and All-R, respectively (**Figure 2C-D**, **Supplementary Figures 3-4**), the only mutants tested to decrease c_{sat}.

530   Together, these data suggest that the type (i.e., arginine vs. lysine), number (i.e., two vs. four vs. six), and distribution (i.e., terminal vs. central) of basic residues neutralized play a role in defining nucleosome phase separation propensity. Results obtained within the NCP context exhibit trends similar to those observed in the simplified peptide–DNA system, but with an added dependence on residue position and a reduced sensitivity to residue number. It is somewhat surprising that the peptide–DNA model captures the overall trends of the more complex NCP system so well, given the absence of the other histone tails. Differential contributions of arginine and lysine to NCP phase separation propensity support that a mixture of arginine and lysine residues may be important for reasons beyond their distinct modifications and recognition by reader proteins.

Figure 2. Nucleosome phase boundary shifts with perturbation of basic residues. A. Histone H3 tail amino acid sequence that protrudes from the nucleosome core. Arginine and lysine residues are highlighted in dark and light blue shades, respectively. **B.** NCP with basic residues of H3 tail colored according to (**A**). NCP conformation is from a molecular dynamics simulation end state (65). A cartoon illustrates that, while inter-nucleosomal tail-DNA interactions are a main driver of nucleosome phase separation, both intra- and inter-nucleosomal interactions are presumed to exist. **C.** Turbidity (absorbance at 600 nm) values (average of n=3) are shown in greyscale for NCP with unmodified or mutant H3, as labeled, and as a function of NCP concentration. Turbidity is plotted as a heatmap ranging from 0.01 (white) to 1.1 (black). Values greater than 0.1 are outlined with a black square to indicate light scattering due to droplet formation. Data were recorded with a set temperature of 25 °C, which had an actual experimental range of 24.9-25.6 °C. **D.** Brightfield microscopy images were collected for the same NCP concentrations as in (**C**), along with additional concentrations to improve c_{sat} determination. The scale bar is 50 μm. A coefficient of variation value (see Methods) was determined for each image. Values greater than 5.0 are considered significant and are outlined in blue to indicate droplets. Full field-of-view images are shown in **Supplementary Figure 4**. Microscopy images were collected at room temperature with an experimental range of 27.4-29.0 °C.

Basic residues within the histone H3 tail govern nucleosome condensate material properties.

Next, we sought to understand how the neutralization of basic residues within the H3 tail influences the microenvironment of nucleosome condensates. Using temperature-controlled video particle tracking (VPT) microrheology (**Figure 3A**), we determined the viscosity of NCP condensates. Overall, neutralization of basic residues decreases the viscosity of NCP

condensates, and the degree of neutralization correlates with the extent of enhanced
condensate fluidity (**Figure 3B-C**, **Supplementary Tables 3-4**). We observe a decrease in
viscosity when we increase the number of lysine neutralizations (compare unmodified at $110 \pm$
10 mPa s to 2xK-$Q_{terminal}$- and 4xK-Q_{even}-H3-NCP at 32 ± 8 and 13 ± 1 mPa s, respectively) and
arginine neutralizations (compare unmodified to R2Q, R8Q, 2xR-$Q_{terminal}$-, 2xR-$Q_{central}$-, and 4xR-
Q-H3-NCP at 39 ± 2 mPa s, 51 ± 4 mPa s, 26 ± 2 mPa s, 43 ± 5 mPa s, and 2.4 ± 0.5 mPa s,
respectively). Neutralization of all four arginine residues within the H3 tail (4xR-Q-H3-NCP)
leads to the lowest measured viscosity among the samples tested. Similar to c_{sat} (**Figure 2D**),
there appears to be a dependence on position. Specifically, a more pronounced decrease in
viscosity is observed for R2Q as compared to R8Q and with neutralization of two arginines in
the 'terminal' region as compared to the 'central' region (compare 2xR-$Q_{terminal}$- and 2xR-$Q_{central}$-
H3-NCP). Furthermore, we observe that the type of residue neutralized influences NCP
condensate viscosity in some samples (compare 4xR-Q- and 4xK-Q_{even}-H3-NCP). Interestingly,
All-K-H3-NCP has decreased condensate viscosity (84 ± 4 mPa s) as compared to unmodified-
NCP while All-R-H3-NCP is the one modification tested that increases the condensate viscosity
(250 ± 30 mPa s). Collectively, the data reveal that modifications to basic residues enable
tuning of NCP condensate viscosity across nearly two orders of magnitude. Notably, a global
linear correlation exists between viscosity and c_{sat} on a natural logarithm scale with two
exceptions: All-K- and 4xR-Q-H3-NCP (**Figure 3D**). These differences may stem from a
complex interplay between hydrodynamic effects and sequence-encoded thermodynamic
interactions in these systems (66).
587

Figure 3. Nucleosome condensate viscosity is tuned by H3 tail charge patterns. A. Schematic representation of the video particle tracking (VPT) microrheology employed to determine the viscosity of condensates formed by unmodified- and mutant-H3-NCPs. The motion of 1-µm yellow-green fluorescent carboxylate-modified beads embedded inside condensates was tracked. Individual trajectories of the tracked beads were analyzed to estimate the mean squared displacement (MSD) of the beads. **B.** Ensemble-average MSD profiles of the tracked beads in condensates of unmodified- and mutant-H3-NCPs at 24 °C. A shift towards lower MSDs depicts slower diffusivity of the beads inside the dense phases of the NCP mutants. **C.** Viscosities of the condensates formed by unmodified- and mutant-H3-NCPs were estimated from the fit of the MSDs to the generalized Stokes–Einstein relation in the diffusive regime, using the power-law scaling of the form $MSD = 4D\tau^{\alpha}$ (Methods section). In the box plot, the black center horizontal line marks the median, the grey box marks the interquartile range (IQR) from the 25^{th} to 75^{th} percentile, and the grey vertical lines mark the minimum (25^{th} percentile − 1.5*IQR) and maximum (75^{th} percentile +1.5*IQR). All viscosities are significantly different from unmodified NCP condensates as determined via ANOVA with Tukey post-hoc analysis. Additional significance comparisons are reported in **Supplementary Table 4**. **D.** Inverse correlation between the natural log of the viscosity and c_{sat} (as determined from brightfield microscopy in **Figure 2D**) of unmodified- and mutant-H3-NCP. The dashed blue line is a guide to the eye for the inverse correlation. The figure contains data from measurements at 9-10 randomly chosen positions of the dense phases over three independently prepared sample repeats. Data symbols are consistent across the figure.

Phase behavior and viscosity of H3 tail-DNA condensates from coarse-grained simulations.

The thermodynamic and material properties of H3 tail-DNA condensates were investigated using coarse-grained simulations (**Figure 4A-B**). To conduct the simulations, each amino acid of the histone tail, the same H3(1-44) peptide sequence as used experimentally (**Figure 1B**), and each nucleotide of a 20-bp poly(dT:dA) double-stranded DNA were coarse-grained into single beads. For the tail peptide, we use an established single-bead model called HPS-CALVADOS (61) in which the effect of residue identity on intermolecular attraction is encoded through the λ-parameter that quantifies the hydropathy scale of residues. In this parametrization, arginine has a substantially higher λ than lysine, reflecting its experimentally established stronger capability to engage in hydrogen-bonding, cation–π, and electrostatic interactions in crowded or partially dehydrated environments. As a result, arginine-rich sequences exhibit markedly stronger effective cohesion than their lysine-rich counterparts.

The phase diagram (**Figure 4C**) shows broad agreement with the experimental results obtained with NCPs (**Figure 2C-D**). The All-R H3 tail mutant displays higher critical temperatures compared to the unmodified and all other mutants. With increasing arginine neutralization (compare unmodified, 2xR-Q$_{terminal}$, and 4xR-Q), both the critical temperature and concentration decrease. The higher critical temperature of 2xR-Q$_{central}$ compared to 2xR-Q$_{terminal}$ suggests a positional effect, where the location of arginine neutralizations within the histone tail influences phase separation behavior and condensate stability. A comparison of the substitution of all arginines with either lysines (All-K) or glutamines (4xR-Q) shows a significant increase in the critical temperature for All-K, along with a broader phase separation boundary (i.e., the condensate concentration is higher at 300 K). The 4xR-Q mutant and its lysine corollary, 4xK-Q$_{even}$, exhibit very similar values of critical temperatures as indicated by the fitting results. The 4xR-Q mutant exhibits the narrowest phase separation boundary compared to all other mutants and the unmodified peptide. The 6xK-Q mutant is identified by the coarse-grained model as having the lowest critical temperature, and under our experimental conditions, this mutant failed to form condensates either in the context of peptide-DNA systems or NCP (**Figure 1C-D, Figure 2C-D**).

Next, we quantified viscosities from coarse-grained simulations of the H3 tail–DNA system using the same sequence variants examined experimentally. In coarse-grained models,

we have previously demonstrated that the microscopic origin of viscosity arises from the
association and dissociation kinetics of sticky, positively charged residues (primarily arginine
and lysine) at the DNA interface (55, 67). When these residues are clustered, their interactions
with DNA become cooperative, increasing the effective dissociation energy. As a result,
viscosity grows nonlinearly with the local density of high-affinity residues. The computational
trends we obtain closely parallel the experimental measurements on NCPs (**Supplementary**
Figure 5) and are fully consistent with this microscopic mechanism. All variants except All-R
show reduced viscosity relative to unmodified H3, whereas All-R displays a dramatically
elevated viscosity. That is because All-R represents a highly cohesive sequence: it contains
nearly twice as many arginines as unmodified-H3(1-44), and the patterning of these residues is
dramatically altered, creating several arginine repeats. No other constructs contain arginine
dipeptides and at most one short window with two arginine residues in three consecutive
positions. In contrast, All-R contains three arginine dipeptides and eight distinct 3-residue
windows enriched in arginine. This dense, patterned clustering of strongly attractive residues
produces a nonlinear amplification of cohesive interactions, leading to slower dissociation from
DNA and therefore substantially higher viscosity. Within the remaining mutants, the 2xR-Q$_{terminal}$
construct exhibits lower viscosity than the 2xR-Q$_{central}$ mutant, and the 4xR-Q mutant shows the
lowest viscosity of all. These results also support an inverse relationship between viscosity and
c$_{sat}$ across the studied constructs (**Figure 4D**, **Supplementary Table 5**), suggesting that both
equilibrium stability and condensate dynamics are governed by the same underlying pattern of
sequence-encoded interactions.
662

663
Figure 4: Phase behavior and viscosity of H3 tail-DNA condensates of unmodified and mutants
from coarse-grained MD simulations. A. Snapshots of 4xR-Q-H3(1-44) (blue) with DNA (red)
condensates at 250 K (upper) and 300 K (lower) from coarse-grained simulations, highlighting the
transition between condensed and dilute phases. **B.** Snapshot from viscosity calculation simulations
performed at 300 K, where the simulation box is systematically shrunk to achieve the condensed-phase
density. **C.** Phase diagrams of unmodified- and mutant-H3 tail peptide with DNA, showcasing the critical
temperature and concentrations associated with phase transitions. Temperature (T) is plotted as a ratio,

relative to the critical temperature of unmodified-NCP ($T_{c, unmodified}$). **D.** Calculated viscosity as a function of
saturation concentration (c_{sat}) derived from the phase diagram analysis of the unmodified- and mutant-H3
tail peptides. The dashed blue line is a guide to the eye for the inverse correlation. Symbol legend in (**D**)
corresponds to (**C**) and (**D**).
675
676
H3 tails remain highly mobile in nucleosome condensates.
As histone tail dynamics are proposed to be important in the histone language (59, 68,
69), we next investigated how these functional protein motions are perturbed by the condensate
environment. While the condensate environment severely restricts local motions in some
systems (70–72), several condensate systems have been reported to display high mobility of
intrinsically disordered regions at the molecular level (73, 74). The measured viscosities of the
NCP condensates (**Figure 3C**, **Supplementary Table 3**) are at the lower end of the values
previously reported for biomolecular condensates (75), suggesting that the condensates are
sufficiently fluid to permit experimental measurement of tail dynamics.
We used solution NMR spectroscopy to investigate the mobility of the H3 tails within
NCP condensates. In these samples, only the H3 component of the NCP is [15]N-labeled, and
only the tail is visible by solution NMR with this labeling approach due to the large size of the
complex (32, 76). Notably, backbone resonances were still observable for the nucleosomal H3
tails within the dense phase of a phase-separated sample, supporting that the tails retain a high
level of mobility within condensates (**Supplementary Figure 6A**, left). However, resonances for
residues towards the ends of the tail (see T3, K4, and V35) broaden almost beyond detection.
Chemical shift differences ($\Delta\delta$) determined from [1]H-[15]N HSQC spectra were minimal between
the dense and light phases (**Supplementary Figure 6C**). This is commonly observed for IDPs
within condensates (e.g., see (70, 71, 73)). The visibility of most H3 resonances in the dense
phase spectra, along with the minimal changes in chemical environment, supports a model
wherein the H3 tails remain disordered and bound to DNA (presumably via a mixture of intra-
and inter-nucleosomal interactions) in a dynamic ensemble of conformations.
We employed NMR spin relaxation experiments to gain insight into nucleosomal H3 tail
dynamics on the ps-ns timescale. On average, [15]N-R1 values increased by 3% while [15]N-R2
values increased by 71% in the dense phase as compared to the light phase (**Figure 5A-B**:
open vs. closed circles, **Supplementary Figure 7**: grey vs. black). The hnNOE profiles for the
two phases are the same within the experimental error (**Figure 5C**, **Supplementary Figure 8**),
with the caveat that the signal quality is poor for [15]N-unmodified-H3-NCP samples. Thus, while
there are only minor, if any, differences between the two phases for [15]N-R1 and hnNOE, the
[15]N-R2 values exhibit a strong dependence on phase (**Figure 5A-C**, **Supplementary Figures**
7-8). The R2 profile for residues 26-29 stands out as distinct between the two phases. The
difference in R2 profile shape for R26, K27, S28, and A29 suggests the possibility of interesting
conformational dynamics changes at the boundary between the 'central' and 'hinge' regions
between the two phases, which we speculate could originate from differences between intra-
and inter-nucleosomal interactions.
The data support that the H3 tails remain mobile on the ps-ns timescale within
nucleosome condensates, while the increased R2 values could reflect i) the increase in
viscosity, ii) a shift in the equilibrium between intra- and inter-nucleosomal tail-DNA interactions
toward inter-nucleosomal interactions within the dense phase, or iii) a combination of the two.
Previous in-cell NMR studies have provided a framework for understanding these data. Similar
to the dense phase of a phase-separated sample, the higher viscosity within a cell is expected
to result in decreased R1 values and increased R2 values, but an exaggerated effect is
observed experimentally for R2 (77, 78). These in-cell observations can be explained by
transient interactions with large biomolecules or complexes (77, 78). The minimal differences in
R1 that we observe between the light and dense phases of unmodified NCP suggest that the

large differences in R2 reflect an increase in inter-nucleosomal interactions within the dense phase. This is consistent with the known role of tail-DNA interactions in nucleosome condensates (7, 16, 21). However, we cannot rule out the possibility that the large differences in R2 could be due to R_{ex} contributions on the ms timescale.

Figure 5. NMR fast-timescale dynamics indicate highly mobile H3 tails within nucleosome condensates. Plots are shown for [15]N-R1 rates (**A**), [15]N-R2 rates (**B**), and hnNOE values (**C**) as a function of residue for unmodified- (black circles), All-K- (blue squares), 4xR-Q- (red triangles), and All-R- (green diamonds) H3-NCP. Data were collected on the light phase (open symbols) and dense phase (closed symbols) of NCP samples prepared in 20 mM MOPS pH 7, 150 mM KCl, 2 mM MgCl$_2$, 0.1 mM EDTA, and 10% D$_2$O. Error bars were determined via the covariance matrix in fitting R1 and R2 decay curves

and represent standard error propagation of the spectral noise for hnNOE values. Residues without data
for any sample are colored grey in the x-axis labels. Data were collected at 600 MHz and 298 K. **D.**
Correlation between dense phase viscosity measured via microrheology and average ^{15}N-R2 rates. For
viscosity, the average and standard deviation of experimental replicates are shown. For R2, the average
and standard deviation across residues plotted in (**B**) are shown. For All-R, the exponential decay and fit
error from the ^{1}H-1D is also shown (open symbol, see Methods and Supplementary Figure 9). The All-R
R2 is likely an underestimate of the true average since data are only from the few most mobile residues.
741
742
H3 tail basic residues govern tail mobility within NCP condensates.
We next investigated nucleosomal H3 tail mobility within a subset of the mutants toward
the extremes of condensate changes in our NCP system: 4xR-Q, All-K, and All-R. ^{1}H-^{15}N
spectral quality (in terms of sensitivity) was improved for All-K and 4xR-Q as compared to
unmodified (**Supplementary Figure 6A-B**). Chemical shift differences ($\Delta\delta$) determined between
the dense and dilute phases from ^{1}H-^{15}N HSQC spectra were equal to (for All-K) or smaller than
(for 4xR-Q) those observed for unmodified (**Supplementary Figure 6C**). In contrast, spectral
quality was poor for All-R, especially for the dense phase (**Supplementary Figure 6A-B**): in ^{1}H-
^{15}N HSQC spectra, signal is largely only observed for the two TGG motifs and the extended
'hinge' region around the second TGG motif, the H3 tail regions known to be most dynamic (76).
NMR analyses are reported below for All-R with the caveat that the signal quality of these
samples is very poor, and the data should be treated as qualitative.
We repeated NMR spin relaxation experiments with the mutants to gain insight into their
effect on ps-ns nucleosomal H3 tail dynamics. Within each phase of All-K and 4xR-Q, ^{15}N-R1
values show a mild increase while ^{15}N-R2 and hnNOE values decrease as compared to
unmodified-NCP (**Figure 5A-C**, **Supplementary Figures 7-8**). A stronger effect is observed for
4xR-Q. The increased mobility of the H3 tail within 4xR-Q-H3-NCP was expected in the light
phase due to a previous study under non-phase-separating conditions (59) and is consistent
with arginine-DNA interactions constraining the mobility of the tails. The increased mobility of
the All-K-H3 tail (albeit not as high as for 4xR-Q) is consistent with arginines forming stronger
interactions with DNA than lysines. As with unmodified, differences in R1 and hnNOE between
the phases are minor for All-K and 4xR-Q, and differences in dense phase ^{15}N-R2 values are
most pertinent for understanding changes in tail mobility within NCP condensates. As discussed
above with unmodified-NCP, R2 profiles from the dense phase likely reflect changes in inter-
nucleosomal interactions in addition to changes in viscosity. The stepwise decrease in dense
phase R2 values when moving from unmodified to All-K to 4xR-Q (**Figure 5B**) could be a
combination of the increased mobility of the tail (with weaker lysine-DNA interactions or arginine
neutralization) and a decrease in inter-nucleosomal interactions (or a related change in the
inter-nucleosomal interaction configuration). The minimal differences in R2 between the two
phases of 4xR-Q could indicate that there are only minor differences in tail-DNA interaction
configurations between the two phases. Minimal data are available for All-R, and while the data
only report on the most dynamic residues, they hint at interesting trends for this mutant: i)
hnNOEs from the light phase 'hinge' region are elevated and suggest that the high arginine
content decreases ps-ns dynamics of the tail (**Figure 5C**), and ii) differences in R2 between the
light and dense phases align with the trends observed for unmodified and All-K (**Figure 5B**,
Supplementary Figures 7-9).
Together, these NMR data support mobility of the H3 tails on the ps-ns timescale within
NCP condensates. Neutralizing arginines or substituting in a weaker DNA-interactor (i.e., lysine)
increases the mobility of the H3 tails, while swapping arginine in for lysine has the opposite
effect. The mutants tested via NMR show a correlation between average R2 values and
viscosity (**Figure 5D**). Note that the average R2 for All-R is from only the most mobile residues
and thus assumed to be an underestimate of the true average across the tail. Differences in R2

between the mutants and their two phases may report on differences in intra- vs. inter-
nucleosomal interactions and the interaction network within condensates.
787
788
DISCUSSION
Here, we show that nucleosome phase separation propensity and properties are
influenced by the number, identity, and distribution of basic residues and their modifications
within the H3 tail. We tested the effects of H3 tail arginine and lysine neutralization on the phase
boundary and material properties of NCP condensates. Overall, neutralizations within the H3 tail
decrease the phase separation propensity of the NCP and the corresponding viscosity of their
condensates. Neutralizing arginines in the 'terminal' region of the tail has a stronger effect than
in the 'central' region of the tail, supporting a positional dependence where residues that can
reach further to bridge nucleosomes play a larger role in phase separation. In general, arginines
within the H3 tail have a stronger contribution to phase separation than lysines. Shifting the H3
tail content towards lysines decreases condensate viscosity while shifting towards arginine is
the one modification tested that leads to an increase in viscosity. The H3 tails remain highly
mobile within nucleosome condensates unless arginine content is increased, and tail mobility
correlates with condensate viscosity. We interpret our data in terms of tail-DNA interactions but
acknowledge that we cannot rule out contributions from histone-histone interactions. The
findings presented here highlight the intrinsic material properties of chromatin at the mesoscale.
However, the transition from mesoscale condensates to microscale chromatin domains likely
involves additional layers of regulation and spatial organization that merit future investigation.
Together, our data support that modifying H3 tail basic residues has a differential direct effect
on nucleosome phase separation (as a function of tail charge patterns), with implications for the
direct effect of charge-modulating histone PTMs on chromatin organization via phase
separation.
Here, the systematic perturbation of H3 tail basic residues begins to address the role of
charge patterning in the histone tails in regulating chromatin phase separation. Charge
patterning in IDRs is important in charge-driven phase separation (79) and is relevant within the
nucleus. Charge blockiness of IDRs is important in controlling which transcriptional regulators
get incorporated into MED1 Mediator condensates (80). A recent study investigated the charge
patterning of the linker histone H1 C-terminal tail and its role in condensates formed with DNA
(81). The C-terminal tails of linker histones are well-mixed sequences that are lysine-rich and
generally arginine-poor. In the context of the H1 C-terminal tail, higher arginine as compared to
lysine content increases phase separation propensity and decreases diffusion within the
condensate (in terms of FRAP recovery). This is consistent with the higher viscosity of All-R-H3-
NCP as compared to the unmodified-H3 sequence containing a mixture of lysine and arginine.
The H3 tails retain a high degree of mobility within NCP condensates. Modifying arginine
residues via neutralization or swapping them out for a weaker DNA-interactor (i.e., lysine) has a
larger effect than the condensate environment on ps-ns tail dynamics as observed via hnNOE.
This highlights that the NCP condensate environment minimally perturbs tail mobility and also
emphasizes the role of dynamics in the histone language (59, 68, 69). ^{15}N-R2 values likely
reflect differences in intra- vs. inter-nucleosomal interactions and network configuration within
the condensate as modulated by basic residue modifications. Biological implications of retained
tail mobility include that the tails are likely not occluded by the condensate environment from
interacting with regulatory proteins in, for example, a transcriptionally active condensate.
Perhaps other regulatory proteins within these nucleosome condensates are also highly mobile,
but this remains to be tested. There is precedent for the high H3 tail mobility observed in NCP
condensates. Within condensates of the H1 C-terminal tail with prothymosin-⍺, long-range chain
reconfiguration of prothymosin-⍺ is only slowed by a factor of three while the viscosity of the
condensate is 300x greater as compared to the light phase (74). Additionally, the germ-granule

protein Ddx4 forms a condensate that is ~50x more concentrated than the light phase but with
similar hnNOE values (73). This contrasts with other systems that exhibit slowed dynamics in
condensates. For example, hnRNPA2, the low-complexity domain of FUS, and an elastin-like
polypeptide show larger changes in ^{15}N-R1 and hnNOE values that are in line with a greater
restriction in local motions (70–72).

Our results add to the emerging model of phase separation driving chromatin
organization and regulation. Various nuclear factors are poised to tune condensate formation
and physicochemical properties. It is anticipated that condensate viscosity influences the
biological function of each chromatin subcompartment, with less viscous condensates
associated with more active chromatin and vice versa (9, 14, 15). We propose that charge-
neutralizing PTMs, such as lysine acetylation and arginine citrullination, tune phase separation
to lower viscosity and allow for greater access to chromatin (**Figure 6**). The differences we
observe in the phase separation propensity, condensate viscosity, and tail mobility within
nucleosome condensates with charge-modulating mutations within the H3 tail support that the
histone language (i.e., the positions and the types of modifications) mediates accessibility and
biological function in part by tuning the biophysical properties of condensates. Furthermore,
these data suggest that histone PTMs may largely tune chromatin phase separation in one
direction, to lower viscosity, because many neutralize basic residues or add negative charge. In
order to tune chromatin condensates to higher viscosity, additional contributions from linker
DNA length or condensate-promoting proteins such as linker histone H1 and HP1◌ are
necessary (7, 22, 23, 82, 83). Variations in histone sequence (histone variants) are also poised
to influence phase separation, especially by altering the composition and relative prevalence of
arginine and lysine (as seen for linker histone (81)). It remains an open question whether other
interactions within chromatin influence phase separation (e.g., DNA sequence and methylation,
the histone core, and tail-tail interactions). The environment and physicochemical properties of
each phase-separated region of chromatin are anticipated to be determined by a combination of
all nuclear factors present—chromatin, chromatin-interacting proteins, and epigenomic
modifications.
864

865
866 **Figure 6. Histone tail charge patterning via modifications modulates chromatin condensates.** In
867 this model, charge-neutralizing PTMs, such as lysine acetylation and arginine citrullination, tune phase
868 separation to lower viscosity and allow for greater access to chromatin. The differences observed in the
869 viscosity and tail mobility of these nucleosome condensates suggest the histone language tunes
870 accessibility and biological function in part by tuning the material properties of condensates.
871

DATA AVAILABILITY

The data underlying this article are available in the manuscript, the supplementary materials, and upon reasonable request to the appropriate corresponding author. Unique plasmids are available from Addgene (plasmid # 240634) or upon request. The relaxation datasets presented in this study are deposited in the Biological Magnetic Resonance Data Bank (BMRB) with deposition numbers 53057, 53058, 53059, and 53458. All analysis and calculations for VPT were done using Python scripts from previous publications (54, 55) that are available on GitHub (https://github.com/BanerjeeLab-repertoire/Material-properties).

SUPPLEMENTARY DATA

Supplementary data are available at *NAR* Online.

ACKNOWLEDGMENTS

Thanks to Drs. Catherine Musselman, Karolin Luger, and Michael Poirier for gifts of histone plasmids.

AUTHOR CONTRIBUTIONS

EM conceived of the initial project. EH, EM, DP, and PB designed the studies. EM, DP, and PB acquired funding and supervised the research. EH and SMZ prepared samples. EH, AS, KS, SY, RG, and EM performed the research and analyzed the data. EH, AS, KS, and EM visualized the data. EH and EM wrote the manuscript with contributions from AS, KS, DP, and PB.

FUNDING

This work was supported by the National Institutes of Health [grant R35GM142594 to E.A.M, R35GM138186 to P.R.B, R35GM138243 to D.A.P., S10OD020000 to the MCW NMR facility]. This study made use of NMRbox: National Center for Biomolecular NMR Data Processing and Analysis, a Biomedical Technology Research Resource (BTRR), which is supported by the National Institutes of Health [grant P41GM111135]. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

CONFLICT OF INTEREST

P.R.B. is a member of the Biophysics Reviews (AIP Publishing) editorial board. This affiliation did not influence the work reported here. All other authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

MATERIALS & CORRESPONDENCE

Correspondence on microrheology measurements should be addressed to Priya Banerjee at prbanerj@buffalo.edu. Correspondence on simulations should be addressed to David Potoyan at potoyan@iastate.edu. Other correspondence and material requests should be addressed to Emma Morrison at emorrison@mcw.edu.

REFERENCES

1. Fujishiro,S., Sasai,M. and Maeshima,K. (2025) Chromatin domains in the cell: Phase separation and condensation. *Curr. Opin. Struct. Biol.*, **91**, 103006.

2. Rippe,K. (2022) Liquid–Liquid Phase Separation in Chromatin. *Csh Perspect Biol*, **14**, a040683.

3. Sabari,B.R., Dall'Agnese,A. and Young,R.A. (2020) Biomolecular Condensates in the Nucleus. *Trends Biochem Sci*, **45**, 961–977.

4. Shin,Y., Chang,Y.-C., Lee,D.S.W., Berry,J., Sanders,D.W., Ronceray,P., Wingreen,N.S., Haataja,M. and Brangwynne,C.P. (2018) Liquid Nuclear Condensates Mechanically Sense and Restructure the Genome. *Cell*, **175**, 1481-1491.e13.

5. Hildebrand,E.M. and Dekker,J. (2020) Mechanisms and Functions of Chromosome Compartmentalization. *Trends Biochem. Sci.*, **45**, 385–396.

6. Narlikar,G.J. (2020) Phase-separation in chromatin organization. *J Biosciences*, **45**, 5.

7. Gibson,B.A., Doolittle,L.K., Schneider,M.W.G., Jensen,L.E., Gamarra,N., Henry,L., Gerlich,D.W., Redding,S. and Rosen,M.K. (2019) Organization of Chromatin by Intrinsic and Regulated Phase Separation. *Cell*, **179**, 470-484.e21.

8. Pei,G., Lyons,H., Li,P. and Sabari,B.R. (2024) Transcription regulation by biomolecular condensates. *Nat. Rev. Mol. Cell Biol.*, 10.1038/s41580-024-00789-x.

9. Erdel,F. (2020) Biophysical mechanisms of chromatin patterning. *Curr Opin Genet Dev*, **61**, 62–68.

10. Bhat,P., Honson,D. and Guttman,M. (2021) Nuclear compartmentalization as a mechanism of quantitative control of gene expression. *Nat Rev Mol Cell Bio*, **22**, 653–670.

11. Li,X., Liu,C., Lei,Z., Chen,H. and Wang,L. (2024) Phase-separated chromatin compartments: Orchestrating gene expression through condensation. *Cell Insight*, **3**, 100213.

12. Sabari,B.R., Dall'Agnese,A., Boija,A., Klein,I.A., Coffey,E.L., Shrinivas,K., Abraham,B.J., Hannett,N.M., Zamudio,A.V., Manteiga,J.C., *et al.* (2018) Coactivator condensation at super-enhancers links phase separation and gene control. *Science*, **361**, eaar3958.

13. Strom,A.R., Emelyanov,A.V., Mir,M., Fyodorov,D.V., Darzacq,X. and Karpen,G.H. (2017) Phase
separation drives heterochromatin domain formation. *Nature*, **547**, 241–245.

14. Larson,A.G. and Narlikar,G.J. (2018) The Role of Phase Separation in Heterochromatin Formation,
Function, and Regulation. *Biochemistry*, **57**, 2540–2548.

15. Eshghi,I., Eaton,J.A. and Zidovska,A. (2021) Interphase Chromatin Undergoes a Local Sol-Gel
Transition upon Cell Differentiation. *Phys. Rev. Lett.*, **126**, 228101.

16. Hammonds,E.F., Harwig,M.C., Paintsil,E.A., Tillison,E.A., Hill,R.B. and Morrison,E.A. (2022)
Histone H3 and H4 tails play an important role in nucleosome phase separation. *Biophys. Chem.*, **283**,
106767.

17. Gibson,B.A., Blaukopf,C., Lou,T., Chen,L., Doolittle,L.K., Finkelstein,I., Narlikar,G.J., Gerlich,D.W.
and Rosen,M.K. (2023) In diverse conditions, intrinsic chromatin condensates have liquid-like
material properties. *Proc National Acad Sci*, **120**.

18. Strickfaden,H., Tolsma,T.O., Sharma,A., Underhill,D.A., Hansen,J.C. and Hendzel,M.J. (2020)
Condensed Chromatin Behaves like a Solid on the Mesoscale In Vitro and in Living Cells. *Cell*, **183**,
1772-1784.e13.

19. Chen,Q., Zhao,L., Soman,A., Arkhipova,A.Y., Li,J., Li,H., Chen,Y., Shi,X. and Nordenskiöld,L.
(2022) Chromatin Liquid–Liquid Phase Separation (LLPS) Is Regulated by Ionic Conditions and Fiber
Length. *Cells*, **11**, 3145.

20. Bertin,A., Mangenot,S., Renouard,M., Durand,D. and Livolant,F. (2007) Structure and Phase
Diagram of Nucleosome Core Particles Aggregated by Multivalent Cations. *Biophys J*, **93**, 3652–
3663.

21. Farr,S.E., Woods,E.J., Joseph,J.A., Garaizar,A. and Collepardo-Guevara,R. (2021) Nucleosome
plasticity is a critical element of chromatin liquid–liquid phase separation and multivalent nucleosome
interactions. *Nat Commun*, **12**, 2883.

22. Zhou,H., Huertas,J., Maristany,M.J., Russell,K., Hwang,J.H., Yao,R., Hutchings,J., Shiozaki,M.,
Zhao,X., Doolittle,L.K., *et al.* (2025) Multi-scale structure of chromatin condensates rationalizes
phase separation and material properties. *bioRxiv*, 10.1101/2025.01.17.633609.

23. Chen,L., Maristany,M.J., Farr,S.E., Luo,J., Gibson,B.A., Doolittle,L.K., Espinosa,J.R., Huertas,J.,
Redding,S., Collepardo-Guevara,R., *et al.* (2025) Nucleosome spacing can fine-tune higher-order
chromatin assembly. *Nat. Commun.*, **16**, 6315.

24. Pepenella,S., Murphy,K.J. and Hayes,J.J. (2013) Intra- and inter-nucleosome interactions of the core
histone tail domains in higher-order chromatin structure. *Chromosoma*, **123**, 3–13.

25. Gordon,F., Luger,K. and Hansen,J.C. (2005) The Core Histone N-terminal Tail Domains Function
Independently and Additively during Salt-dependent Oligomerization of Nucleosomal Arrays*. *J Biol
Chem*, **280**, 33701–33706.

26. Kan,P.Y., Lu,X., Hansen,J.C. and Hayes,J.J. (2007) The H3 Tail Domain Participates in Multiple
Interactions during Folding and Self-Association of Nucleosome Arrays. *Molecular and cellular*
biology, **27**, 2084–2091.

27. Kan,P.-Y., Caterino,T.L. and Hayes,J.J. (2009) The H4 tail domain participates in intra- and
internucleosome interactions with protein and DNA during folding and oligomerization of nucleosome
arrays. *Molecular and cellular biology*, **29**, 538–546.

28. Zheng,C., Lu,X., Hansen,J.C. and Hayes,J.J. (2005) Salt-dependent Intra- and Internucleosomal
Interactions of the H3 Tail Domain in a Model Oligonucleosomal Array. *The Journal of biological*
chemistry, **280**, 33552–33557.

29. Schwarz,P.M., Felthauser,A., Fletcher,T.M. and Hansen,J.C. (1996) Reversible Oligonucleosome
Self-Association: Dependence on Divalent Cations and Core Histone Tail Domains †. *Biochemistry-*
us, **35**, 4009–4015.

30. Bertin,A., Renouard,M., Pedersen,J.S., Livolant,F. and Durand,D. (2007) H3 and H4 Histone Tails
Play a Central Role in the Interactions of Recombinant NCPs. *Biophys J*, **92**, 2633–2645.

31. Berezhnoy,N.V., Liu,Y., Allahverdi,A., Yang,R., Su,C.-J., Liu,C.-F., Korolev,N. and Nordenskiöld,L.
(2016) The Influence of Ionic Environment and Histone Tails on Columnar Order of Nucleosome
Core Particles. *Biophysical Journal*, **110**, 1720–1731.

32. Morrison,E.A., Bowerman,S., Sylvers,K.L., Wereszczynski,J. and Musselman,C.A. (2018) The
conformation of the histone H3 tail inhibits association of the BPTF PHD finger with the nucleosome.
Elife, **7**, e31481.

33. Moore,S.C. and Ausió,J. (1997) Major Role of the Histones H3-H4 in the Folding of the Chromatin
Fiber. *Biochem Bioph Res Co*, **230**, 136–139.

34. Millán-Zambrano,G., Burton,A., Bannister,A.J. and Schneider,R. (2022) Histone post-translational
modifications — cause and consequence of genome function. *Nat. Rev. Genet.*, **23**, 563–580.

35. Wang,L., Gao,Y., Zheng,X., Liu,C., Dong,S., Li,R., Zhang,G., Wei,Y., Qu,H., Li,Y., *et al.* (2019)
Histone Modifications Regulate Chromatin Compartmentalization by Contributing to a Phase
Separation Mechanism. *Molecular cell*, **76**, 646-659.e6.

36. Larson,A.G., Elnatan,D., Keenen,M.M., Trnka,M.J., Johnston,J.B., Burlingame,A.L., Agard,D.A.,
Redding,S. and Narlikar,G.J. (2017) Liquid droplet formation by HP1α suggests a role for phase
separation in heterochromatin. *Nature*, **547**, 236–240.

37. Nitsch,S., Shahidian,L.Z. and Schneider,R. (2021) Histone acylations and chromatin dynamics:
concepts, challenges, and links to metabolism. *Embo Rep*, **22**, e52774.

38. Fuhrmann,J., Clancy,K.W. and Thompson,P.R. (2015) Chemical Biology of Protein Arginine
Modifications in Epigenetic Regulation. *Chem Rev*, **115**, 5413–5461.

39. Mascotti,D.P. and Lohman,T.M. (1997) Thermodynamics of Oligoarginines Binding to RNA and
DNA †. *Biochemistry-us*, **36**, 7272–7279.

40. Wintjens,R., Liévin,J., Rooman,M. and Buisine,E. (2000) Contribution of cation-π interactions to the
stability of protein-DNA complexes11Edited by J. Thornton. *J Mol Biol*, **302**, 393–408.

41. Rohs,R., West,S.M., Sosinsky,A., Liu,P., Mann,R.S. and Honig,B. (2009) The role of DNA shape in
protein-DNA recognition. *Nature*, **461**, 1248–1253.

42. DeRouchey,J., Hoover,B. and Rau,D.C. (2013) A Comparison of DNA Compaction by Arginine and
Lysine Peptides: A Physical Basis for Arginine Rich Protamines. *Biochemistry-us*, **52**, 3000–3009.

43. Leng,M. and Felsenfeld,G. (1966) The preferential interactions of polylysine and polyarginine with
specific base sequences in DNA. *Proc. Natl. Acad. Sci.*, **56**, 1325–1332.

44. Lenton,S., Hervø-Hansen,S., Popov,A.M., Tully,M.D., Lund,M. and Skepö,M. (2021) Impact of
Arginine–Phosphate Interactions on the Reentrant Condensation of Disordered Proteins.
Biomacromolecules, 10.1021/acs.biomac.0c01765.

45. Alshareedah,I., Kaur,T., Ngo,J., Seppala,H., Kounatse,L.-A.D., Wang,W., Moosa,M.M. and
Banerjee,P.R. (2019) Interplay between Short-Range Attraction and Long-Range Repulsion Controls
Reentrant Liquid Condensation of Ribonucleoprotein–RNA Complexes. *J Am Chem Soc*, **141**, 14593–
14602.

46. Boeynaems,S., Holehouse,A.S., Weinhardt,V., Kovacs,D., Lindt,J.V., Larabell,C., Bosch,L.V.D.,
Das,R., Tompa,P.S., Pappu,R.V., *et al.* (2019) Spontaneous driving forces give rise to protein–RNA
condensates with coexisting phases and complex material properties. *Proc National Acad Sci*, **116**,
201821038.

47. Fisher,R.S. and Elbaum-Garfinkle,S. (2020) Tunable multiphase dynamics of arginine and lysine
liquid condensates. *Nat Commun*, **11**, 4628.

48. Li,Z. and Kono,H. (2016) Distinct Roles of Histone H3 and H2A Tails in Nucleosome Stability.
Scientific reports, **6**, srep31437.

49. Peng,Y., Li,S., Onufriev,A., Landsman,D. and Panchenko,A.R. (2021) Binding of regulatory proteins
to nucleosomes is modulated by dynamic histone tails. *Nat Commun*, **12**, 5280.

50. Paintsil,E.A. and Morrison,E.A. (2022) DNA-Protein Interactions, Methods and Protocols. *Methods
Mol Biology*, **2599**, 163–175.

51. Hammonds,E.F. and Morrison,E.A. (2022) DNA-Protein Interactions, Methods and Protocols.
Methods Mol. Biol., **2599**, 177–190.

52. Schindelin,J., Arganda-Carreras,I., Frise,E., Kaynig,V., Longair,M., Pietzsch,T., Preibisch,S.,
Rueden,C., Saalfeld,S., Schmid,B., *et al.* (2012) Fiji: an open-source platform for biological-image
analysis. *Nat Methods*, **9**, 676–682.

53. Tinevez,J.-Y., Perry,N., Schindelin,J., Hoopes,G.M., Reynolds,G.D., Laplantine,E., Bednarek,S.Y.,
Shorte,S.L. and Eliceiri,K.W. (2017) TrackMate: An open and extensible platform for single-particle
tracking. *Methods*, **115**, 80–90.

54. Alshareedah,I., Moosa,M.M., Pham,M., Potoyan,D.A. and Banerjee,P.R. (2021) Programmable
viscoelasticity in protein-RNA condensates with disordered sticker-spacer polypeptides. *Nat.*
Commun., **12**, 6620.

55. Alshareedah,I., Singh,A., Yang,S., Ramachandran,V., Quinn,A., Potoyan,D.A. and Banerjee,P.R.
(2024) Determinants of viscoelasticity and flow activation energy in biomolecular condensates. *Sci.*
Adv., **10**, eadi6539.

56. Maciejewski,M.W., Schuyler,A.D., Gryk,M.R., Moraru,I.I., Romero,P.R., Ulrich,E.L.,
Eghbalnia,H.R., Livny,M., Delaglio,F. and Hoch,J.C. (2017) NMRbox: A Resource for Biomolecular
NMR Computation. *Biophys J*, **112**, 1529–1534.

57. Delaglio,F., Grzesiek,S., Vuister,G.W., Zhu,G., Pfeifer,J. and Bax,A. (1995) NMRPipe: a
multidimensional spectral processing system based on UNIX pipes. *Journal of biomolecular NMR*, **6**,
277–293.

58. Vranken,W.F., Boucher,W., Stevens,T.J., Fogh,R.H., Pajon,A., Llinas,M., Ulrich,E.L., Markley,J.L.,
Ionides,J. and Laue,E.D. (2005) The CCPN data model for NMR spectroscopy: development of a
software pipeline. *Proteins*, **59**, 687–696.

59. Jennings,C.E., Zoss,C.J. and Morrison,E.A. (2023) Arginine anchor points govern H3 tail dynamics.
Frontiers Mol Biosci, **10**, 1150400.

60. Tugarinov,V. and Kay,L.E. (2003) Quantitative NMR studies of high molecular weight proteins:
application to domain orientation and ligand binding in the 723 residue enzyme malate synthase G.
Journal of molecular biology, **327**, 1121–1133.

61. Cao,F., Bülow,S. von, Tesei,G. and Lindorff☐Larsen,K. (2024) A coarse☐grained model for
disordered and multi☐domain proteins. *Protein Sci.*, **33**, e5172.

62. Banerjee,P.R., Milin,A.N., Moosa,M.M., Onuchic,P.L. and Deniz,A.A. (2017) Reentrant Phase
Transition Drives Dynamic Substructure Formation in Ribonucleoprotein Droplets. *Angewandte*
Chemie Int Ed, **56**, 11354–11359.

63. Teif,V.B. and Bohinc,K. (2011) Condensed DNA: Condensing the concepts. *Prog Biophysics Mol*
Biology, **105**, 208–222.

64. Nguyen,T.T. and Shklovskii,B.I. (2001) Complexation of DNA with positive spheres: Phase diagram
of charge inversion and reentrant condensation. *J Chem Phys*, **115**, 7298–7308.

65. Morrison,E.A., Baweja,L., Poirier,M.G., Wereszczynski,J. and Musselman,C.A. (2021) Nucleosome
composition regulates the histone H3 tail conformational ensemble and accessibility. *Nucleic Acids*
Res, **49**, gkab246-.

66. Alshareedah,I., Borcherds,W.M., Cohen,S.R., Singh,A., Posey,A.E., Farag,M., Bremer,A.,
Strout,G.W., Tomares,D.T., Pappu,R.V., *et al.* (2024) Sequence-specific interactions determine
viscoelasticity and ageing dynamics of protein condensates. *Nat. Phys.*, **20**, 1482–1491.

67. Yang,S., Banerjee,P.R. and Potoyan,D.A. (2024) Microscopic Origins of Flow Activation Energy in
Biomolecular Condensates. *J. Phys. Chem. B*, **128**, 12348–12357.

68. Tsunaka,Y., Furukawa,A. and Nishimura,Y. (2022) Histone tail network and modulation in a
nucleosome. *Curr Opin Struc Biol*, **75**, 102436.

69. Ghoneim,M., Fuchs,H.A. and Musselman,C.A. (2021) Histone Tail Conformations: A Fuzzy Affair
with DNA. *Trends Biochem Sci*, 10.1016/j.tibs.2020.12.012.

70. Ryan,V.H., Dignon,G.L., Zerze,G.H., Chabata,C.V., Silva,R., Conicella,A.E., Amaya,J., Burke,K.A.,
Mittal,J. and Fawzi,N.L. (2018) Mechanistic View of hnRNPA2 Low-Complexity Domain Structure,
Interactions, and Phase Separation Altered by Mutation and Arginine Methylation. *Molecular cell*, **69**,
465-479.e7.

71. Burke,K.A., Janke,A.M., Rhine,C.L. and Fawzi,N.L. (2015) Residue-by-Residue View of In Vitro
FUS Granules that Bind the C-Terminal Domain of RNA Polymerase II. *Molecular cell*, **60**, 231–241.

72. Reichheld,S.E., Muiznieks,L.D., Keeley,F.W. and Sharpe,S. (2017) Direct observation of structure
and dynamics during phase separation of an elastomeric protein. *Proc. Natl. Acad. Sci.*, **114**, E4408–
E4415.

73. Brady,J.P., Farber,P.J., Sekhar,A., Lin,Y.-H., Huang,R., Bah,A., Nott,T.J., Chan,H.S., Baldwin,A.J.,
Forman-Kay,J.D., *et al.* (2017) Structural and hydrodynamic properties of an intrinsically disordered
region of a germ cell-specific protein on phase separation. *Proc. Natl. Acad. Sci.*, **114**, E8194–E8203.

74. Galvanetto,N., Ivanović,M.T., Chowdhury,A., Sottini,A., Nüesch,M.F., Nettels,D., Best,R.B. and
Schuler,B. (2023) Extreme dynamics in a biomolecular condensate. *Nature*, **619**, 876–883.

75. Sanfeliu-Cerdán,N. and Krieg,M. (2025) The mechanobiology of biomolecular condensates. *Biophys.*
Rev., **6**, 011310.

76. Stützer,A., Liokatis,S., Kiesel,A., Schwarzer,D., Sprangers,R., Söding,J., Selenko,P. and Fischle,W.
(2016) Modulations of DNA Contacts by Linker Histones and Post-translational Modifications
Determine the Mobility and Modifiability of Nucleosomal H3 Tails. *Molecular cell*, **61**, 247–259.

77. Ye,Y., Liu,X., Zhang,Z., Wu,Q., Jiang,B., Jiang,L., Zhang,X., Liu,M., Pielak,G.J. and Li,C. (2013)
19F NMR Spectroscopy as a Probe of Cytoplasmic Viscosity and Weak Protein Interactions in Living
Cells. *Chem. A Eur. J.*, **19**, 12705–12710.

78. Leeb,S., Yang,F., Oliveberg,M. and Danielsson,J. (2020) Connecting Longitudinal and Transverse
Relaxation Rates in Live-Cell NMR. *J. Phys. Chem. B*, **124**, 10698–10707.

79. Chang,L.-W., Lytle,T.K., Radhakrishna,M., Madinya,J.J., Vélez,J., Sing,C.E. and Perry,S.L. (2017)
Sequence and entropy-based control of complex coacervates. *Nat. Commun.*, **8**, 1273.

80. Lyons,H., Veettil,R.T., Pradhan,P., Fornero,C., Cruz,N.D.L., Ito,K., Eppert,M., Roeder,R.G. and
Sabari,B.R. (2023) Functional partitioning of transcriptional regulators by patterned charge blocks.
Cell, **186**, 327-345.e28.

81. Watson,M., Sabirova,D., Hardy,M.C., Pan,Y., Carpentier,D.C.J., Yates,H., Wright,C.J., Chan,W.H.,
Destan,E. and Stott,K. (2024) A DNA condensation code for linker histones. *Proc. Natl. Acad. Sci.*,
121.

82. Keenen,M.M., Brown,D., Brennan,L.D., Renger,R., Khoo,H., Carlson,C.R., Huang,B., Grill,S.W.,
Narlikar,G.J. and Redding,S. (2021) HP1 proteins compact DNA into mechanically and positionally
stable phase separated domains. *Elife*, **10**, e64563.

83. Deshpande,P., Prentice,E., Ceballos,A.V., Casaccia,P. and Elbaum-Garfinkle,S. (2024) Epigenetic
marks uniquely tune the material properties of HP1α condensates. *Biophys. J.*, **123**, 1508–1518.

1142