Deconvoluting single-cell transcriptomics reveals cellular programs

regulated by cell-cell communication in colorectal cancer

Han Zhang[1], Binfeng Lu[3], Aodong Qiu[1], Gregory F. Cooper[1], Anwaar Saeed[2], John W. Paisley[4],

Xinghua Lu[1], Lujia Chen[1]*

[1]Department of Biomedical Informatics, University of Pittsburgh, Pittsburgh, PA, 15206, USA

[2]UPMC Hillman Cancer Center, Pittsburgh, PA 15232, USA

[3]Center for Discovery and Innovation, Hackensack Meridian Health, Nutley, NJ 07110, USA

[4]Columbia University, New York, NY 10027, USA

*Correspondence: luc17@pitt.edu

Conflicts of Interest Statement:
Lujia Chen and Xinghua Lu report leadership roles with DeepRx. Anwaar Saeed reports reports a leadership role

with Autem therapeutics, Exelixis, KAHR medical and Bristol-Myers Squibb; consulting or advisory board role

with AstraZeneca, Bristol-Myers Squibb, Merck, Exelixis, Pfizer, Xilio therapeutics, Taiho, Amgen, Autem

therapeutics, KAHR medical, and Daiichi Sankyo; institutional research funding from AstraZeneca, Bristol-Myers

Squibb, Merck, Clovis, Exelixis, Actuate therapeutics, Incyte Corporation, Daiichi Sankyo, Five prime therapeutics,

Amgen, Innovent biologics, Dragonfly therapeutics, Oxford Biotherapeutics, Arcus therapeutics, and KAHR

medical; and participation as a data safety monitoring board chair for Arcus therapeutics. The remaining authors

have no relevant financial interests to disclose.

20
Abstract

Cells within a tissue microenvironment communicate through intricate cell-cell communication

(CCC) networks. In this meta-analysis of eight single-cell cohorts encompassing 153 patients

and 279 samples, we advance the understanding of CCC networks in colorectal cancers through

a novel analytical framework. Employing hierarchical topic modeling, we identify gene

expression modules (GEMs) that mirror single-cell signaling states, crucial for deciphering the

complexity of intercellular interactions. By applying causal discovery methods, we

systematically uncover GEMs likely regulated by ligand-receptor signaling and cross-cell-type

communication. This analysis reveals cross-cell-type CCC programs, marked by highly

correlated GEMs across various cell types, shedding light on the intricate CCC networks within

the tumor microenvironment. Spatial transcriptomics further validate these findings by

demonstrating the co-localization of GEMs within CCC programs in distinct spatial domains,

emphasizing the spatial dynamics of tumor intercellular communication. Our interactive website

(http://44.192.10.166:3838/) and analytical framework equip researchers with powerful tools to

explore these complex mechanisms, potentially uncovering novel drug targets and refining

strategies for precision immunotherapies. This comprehensive study not only presents a detailed

catalog of CCC networks driven by ligand-receptor interactions in colorectal cancer but also

highlights the significance of integrating multi-sample and patient data to unravel the molecular

underpinnings of cancer communication pathways.

Introduction

Colorectal cancer (CRC) is the 3[rd] most common cancer and the 2[nd] cause of cancer death [1]. The discovery of the consensus molecular subtypes (CMS) among CRC tumors reflects the heterogeneity of composition and functional states of cells in the tumor microenvironments (TME), which in turn may underlie diverse behaviors of disease progression and responses to anticancer treatments by tumors. For example, except for those with deficient mismatch repair (dMMR), most CRCs do not respond to anti-PD-1 immune therapy [2]. Such a relationship between genomic pattern and clinical response is the result of cell-cell communications (CCC) within the TME, through which the downstream signaling pathways are mediated and regulated [3,4]. Therefore, a comprehensive understanding of the cell states and communication network in the TME are vital for uncovering the heterogeneous mechanisms of tumor growth, immune evasion and treatment resistance, which are essential for precision oncology.

Modeling CCC requires inferring the cellular states of individual cells, identifying synchronous changes in the cell-state among various cell types that may arise due to CCC, and investigating the molecular mechanism underlying CCC [5-8]. Single-cell RNA sequencing (single-cell RNA-seq) is a powerful technology for profiling the transcriptomes of individual cells. As gene expressions are tightly regulated in a cell, the co-expression patterns of genes serve as proxies of the states of signaling pathways in a cell, which can be discovered by deconvoluting scRNA-seq data. Furthermore, the emergence of spatial transcriptomic technology provides additional spatial relationships between cells within their native spatial context [9,10]. Thus, integrating scRNA-seq with spatial transcriptome provides the spatial context for interactions among different cell populations [11] and enhances the accuracy of modeling CCC.

Single-cell transcriptomes can be used to investigate cellular states at different levels. At the whole transcriptome level, cells sharing similar overall transcriptomic profiles can be discovered as discrete cell clusters using state-of-the-art clustering techniques [12-15]. Although these approaches are good at classifying different cell populations, it does not allow examination of transcriptomic programs within individual cells that underpin the biological functions of these cells [16]. To discover and represent transcriptomic programs within each cell, different statistical- and factorization- based models, e.g., non-negative matrix factorization (NMF) and latent Dirichlet allocation (LDA), have been used to deconvolute transcriptional programs by capturing the co-expression pattern of certain cell subsets [17-20]. However, LDA models impose strong assumption of the independence among gene programs while NMF models do not explicitly identify dependencies among programs. Moreover, these methodologies fail to capture the well-established phenomenon that cells belonging to a cell type, such as T and NK (TNK) cells, exhibit highly coordinated transcriptomic programs. As a result, both approaches inadequately represent the nested nature of gene expression modules (GEM) that are observed in biology. These GEMs are more accurately represented as a hierarchical structure, wherein transcriptomic programs shared by cells across different cell subtypes, such as CD8+ T cells and CD4+ T cells, are positioned close to the root of the hierarchy. In contrast, transcriptomic programs underlying the differentiation of specialized cells are positioned as leaves. The methods of capturing such hierarchical relationships of transcriptomic programs remain relatively underdeveloped.

Transduction of signals across cells involves ligand-receptor interactions. Different methods have been developed to find potential ligand-receptor-mediated intercellular communication, e.g.,

CellChat, CellPhoneDBv2.0 [6,21,22]. However, those methods are mainly based on the expression

and enrichment of ligand-receptor pairs in specific cell clusters but do not explicitly model the

impact of the signals transmitted by a ligand-receptor pair, e.g., genes regulated by PD-L1-PD-1

signaling. Furthermore, few methods exist for modeling coordinated cellular states of different

cells within a tumor, e.g., certain subtype of myeloid cells and subtype of stromal cells tend to

co-exist in tumors, or inferring whether the changed state of one type of cell directly influences

the state of other types of cells.

To solve these challenges, we propose a novel framework for integrating bulk, single-cell, and

spatial RNA-seq data to depict a comprehensive landscape of CRC TMEs (**Fig. 1**). We employed

the nested Hierarchical Dirichlet process (nHDP) model to deconvolute a large collection of

single-cell transcriptomes from 8 cohorts with over 153 CRC patients and over 626,858 single-

cells (**Fig. 1A, Extended Data Table. 1 & Fig. 1**). nHDP organizes the discovered gene

expression modules (GEMs) in a hierarchy, enabling it to separate GEMs (root node)

representing general transcriptomic processes from those reflecting specific functional state of

cells (leave node) (**Fig. 1B & 1C**). We then systematically investigated whether the expression

of a GEM was regulated by ligand-receptor signaling (**Fig. 1D**) using the hill function that is

used in pharmacology to measure the binding affinity between ligand and receptor. By studying

the coordinated expression of GEMs, we systematically investigated the statistically significant

intercellular communications across cell types (**Fig. 1E**) with spatial transcriptome and then

extended our analysis to discover patterns of intercellular communication programs (GEM

programs) shared within subpopulations of tumors (**Fig. 1F**). Employing techniques of causal

analyses, we further searched for ligand-receptor pairs that contribute to mediate signal

transduction of a CCC and further constructed a CCC network among cells expressing

coordinated GEMs as well as biomarker discovery (**Fig. 1G**). We envision this comprehensive

framework enables researchers to study complex mechanisms underlying distinct TMEs,

investigate different immune evasion mechanisms, discover potential drug targets, and guide

precision immune therapies.

115
Fig. 1: General design of the study. **A.** Merged CRC single-cell RNA-seq data from 8 cohorts.

B. Diagram of the nHDP model using the expression of genes (a *cell-by-gene* matrix) in TNK

cells as input. The transcriptome of one cell can be represented by a unique mixture of GEMs in

the nHDP tree. The nHDP model outputs two matrices: a *gene-by-GEM* matrix reflecting the

gene composition of a GEM and a *cell-by-GEM* matrix representing the GEM compositions of

each cell. **C.** nHDP discovered the tree-structured organization of GEMs. Each node in the tree

represents a GEM. A pie chart represents the percentage of each cell type the GEM is expressed.

D. Illustration of modeling the relationship between ligand-receptor (LR) and its downstream

GEMs. **E.** Modeling cell-cell communication by combining single-cell and spatial transcriptomes.

F. Learning patterns of coordinated expression of GEMs across cell types to discover distinct

GEM programs and associated tumor microenvironments. **G.** Downstream discoveries enabled

by deconvoluting single-cell transcriptomes by nHDP: **1)** GEMs highly associated with the

signal of LR. **2)** Subnetwork of intercellular communication for GEMs and LRs of interest. **3)**

GEM as potential novel biomarker for therapies.

Result

A comprehensive atlas of GEMs expressed by cells in CRC tumor microenvironments.

We collected single-cell RNA-seq data from 8 independent CRC cohorts. Then, we separated cells into 5 major cell types: TNK: mainly including T and natural killing (NK) cells (N=246,560); Myeloid: myeloid cells, including monocytes, macrophages, neutrophils, and dendritic cells (N=94,663); PlasmaB: including B and plasma cells (N=96,637); Stroma: stromal cells, including endothelial, fibroblast and myofibroblast cells (N=69,128); Epi: epithelial cells (N=119,870) (**Extended Data Table. 1**). The merged cells for each cell category were clustered and labeled using Seurat pipeline and canonical marker genes. After the integration and batch removal, we identified 18 clusters in TNK cells (**Fig. 2A**)**,** 18 cell populations in both Myeloid cells and PlasmaB cells and 17 cell groups in Stroma cells (**Extended Data Fig. 1**). We also showed that the distributions of datasets are well maintained in all clusters in TNK cells (**Fig. 2B**), suggesting the batch effects were regressed out; thus, the pooled TNK cells reflect the landscape of tumor-infiltrating TNK cells in CRC.

To model the tree structure of the TNK gene expression modules (GEMs), we utilized the nested hierarchical Dirichlet process (nHDP) model to identify a series of nested gene expression modules (GEMs) [23], where a GEM is a probability distribution over the gene space that captures the coordinated expression pattern of a set of genes in a subpopulation of cells. The nHDP model organizes inferred GEMs in a hierarchical tree structure, and we set branching factors of 5, 4, and 3 at 1^{st}, 2^{nd}, and 3^{rd} layers, respectively, leading to a total of 85 nodes to capture GEMs with different levels of granularity and specificity. **Fig. 2C** shows the tree organization of the GEMs learned using TNK cells as input. The nHDP model populated the nodes in the hierarchy through

a data-driven approach. Each node in the tree represents a GEM. The ones close to the root

(TNK GEMs 1-5 in **Fig. 2C**) represent GEMs that are more broadly expressed in all T, innate

lymphoid cells (ILC), mucosal-associated invariant T cells (MAIT) and NK cells, which can be

used to define major subtypes of T and NK (TNK) cells. The ones close to the leaves represent

GEMs specifically expressed in more differentiated cells. In the TNK GEM tree, only 43 TNK

GEMs were identified as biologically meaningful and represented as filled nodes. The nodes

identified as biologically meaningless were shown as empty nodes without color.

We systematically inspected the TNK GEMs reflecting TNK transcriptomic processes. The pie

chart within each GEM illustrates the proportion of cells expressing that specific GEM across

various T cell subtypes, including $CD8^+$ T cells, $CD4^+$ T cells, ILC, MAIT, NK cells, regulatory

T cells (Treg) and Proliferation T cells. It is interesting to note that major branches of the nHDP

hierarchy tree, as defined by GEMs 1-4, represent major TNK subtypes, including CD8/NK,

CD4, Treg, and proliferation T cells (**Fig. 2C**). Projecting the expression of GEMs onto the

UMAP of TNK cells (**Fig. 2D**), we noticed that certain GEMs were concentrated in a particular

cell cluster or a combination of clusters, representing a characteristic transcriptomic program that

is active in the cell cluster. For example, TNK50 is concentrated in cells belonging to cluster #4,

which consists of T helper (Th) cells expressing CXCL13; TNK35 is mainly expressed in T cell

cluster #6, a subset of T_{reg}. Conversely, there are also GEMs shared across multiple cell clusters,

suggesting a transcriptomic program that is shared across multiple subtypes of T cells. For

example, TNK26 is expressed in a subset of $CXCL13^+CD4^+$ T cells (cluster #4), $MKI67^+$

proliferative T cells (cluster #10), and $CXCL13^+CD8^+$ T cells (cluster #11). TNK42 is expressed

in MAIT cells (cluster #9) and ILC (cluster #17). Finally, TNK12 is expressed broadly among

Treg, Th, and proliferating T cells. These results indicate that our model can capture

transcriptomic programs that are both shared and specialized by cells belonging to different cell

subtypes. To test the performance of our model, we also compared the distributions of the

transcriptomic programs that were found by our model and NMF [24]. Our result demonstrates that

the nHDP excels in capturing more specific cellular programs shared across multiple cell clusters

when compared to NMF as shown in **Extended Data Fig. 2**. For instance, TNK15 captures a

cellular program of greater specificity, shared among cells from 4 TNK cell clusters, surpassing

the performance of NMF5. NMF5 captures a more generalized cellular program that combines

programs represented by TNK15, TNK42, and TNK51. Similarly, NMF58 captures the cellular

programs combining programs represented by TNK17, TNK29, and TNK50.

To make the GEMs more biologically explainable, we depict the distinctive gene composition

profile of each GEM (top 20) through a word cloud, and the top three associated transcription

factors identified by transcription factor analysis using ChEA3 [25] were illustrated as black dots in

Fig. 2D. It is interesting to note that TNK26 is enriched with genes characteristic of exhausted T

cells including PDCD1, CXCL13 and TIGIT. Based on the transcription factor analysis,

NFATC2 regulated the top 50 genes associated with TNK26. Several studies have shown that the

transcription factor NFAT is involved in the exhaustion process of activated CD8$^+$ T cells [26]. In

addition, the top genes associated with TNK26 and TNK50 are both regulated by transcription

factor ZNF831, which has been shown to inhibit the functional activity of effector cells through

STAT pathway [27]. This indicates the potential existence of a signal leading to the regulation of T

cell exhaustion. We further examined the distribution of GEMs in different TNK clusters (**Fig.**

2E), unveiling which GEMs are broadly shared among cells assigned to multiple clusters and

which GEMs are specific to a particular cell cluster. **Extended Data Fig. 3** displays the detailed

comparison between GEMs and TNK cell clusters by measuring the correlation between top 50

genes of a GEM and highly variable genes in each TNK cell subtype. Furthermore, we

performed pathway enrichment analyses in Reactome to investigate the functions of top genes

associated with GEMs (**Fig. 2F**). Interestingly, several TNK GEMs, including TNK6, TNK26,

TNK35, and TNK41, are associated with PD-1 signaling pathway. The dot plots showing the

distribution of GEMs in traditional cell clusters and the enriched signaling pathways for other

cell types (PlasmaB, Myeloid, Stromal) are listed in **Extended Data Fig. 4 & 5**. The detailed

annotation of each GEM, including the UMAP, cell composition, and top associated genes, is

listed in our interactive website. Besides, the enriched MsigDB gene sets (H & C1-C8) for each

GEM, derived from the hypergeometric analysis, is shown in **Extended Data Table. 7**.

In summary, by deconvoluting the transcriptomes of a large number of single cells using nHDP,

our approach revealed GEMs that represent transcriptomic programs within or across traditional

cell subtypes, which reflect certain signaling events within these cells.

214
Fig. 2: Deconvoluting single-cell transcriptomics to discover GEMs reflecting the state of

signaling pathways. **A.** UMAP plot of 246,560 merged TNK cells from 8 CRC cohorts. **B.** The

proportion plot reflects the origins of cells in each cell cluster. **C.** The hierarchical tree structure

of TNK GEMs was learned from the nHDP model, which was trained with merged TNK cells.

The nHDP model has three levels. The first level contains GEMs from 1-5. The second level

contains GEMs from 6-25. The third level contains GEMs from 26-85. Each color represents a

subtype of T cell. The unused GEMs are colored grey. **D.** The UMAP plot, word cloud plot, and

top associated TFs of selected TNK GEMs. **E.** The functional involvement of TNK GEMs within

each cell cluster. **F.** The enriched signaling pathways associated with TNK GEMs.

Discovering potential regulatory relationships between ligand-receptor signaling and

GEMs

We set out to examine if the expression of a GEM in cells is potentially regulated by certain

ligand-receptor signaling by testing the hypothesis posited in **Fig. 3A**: If a ligand-receptor pair

regulates the expression of the member genes of a GEM, the overall expression value of the

GEM can be modeled with a sigmoid dose-response model with respect to ligand-receptor pair.

We modeled the relationship of the expression of ligand, receptor, and GEM at the tumor sample

level. We first generated pseudo-bulk RNA-seq (a *tumor-by-GEM* matrix) from the single-cell

RNA-seq (a *cell-by-GEM* matrix) for tumor samples. Then we estimated the concentration of

ligand and receptor proteins based on the pseudo-bulk expression (see Methods) of the ligand

($[L]$) and receptor ($[R]$) for each of the 151 tumors. Additionally, we approximated the

concentrate of protein complex as the product of ligand and receptor: $[LR] \propto [L]*[R]$ [5]. Pseudo-

bulk expression of GEMs was estimated using the GSVA analysis. Subsequently, we conducted

a systematic search against all GEMs with known LR pairs from NicheNet database and retained

the LR pairs that could predict the expression of the GEM with statistical significance (**Extended**

Data Table. 2).

We investigated the GEMs that are strongly associated with immune checkpoints programmed

cell death protein 1 (PD-1, encoded by *PDCD1*) and cytotoxic T lymphocyte-associated protein

4 (*CTLA4*) signaling (**Fig. 3B**). For the PD-1 signaling, the top two TNK GEMs that could be

predicted by approximated [*LR*] with the strongest non-linear correlation were TNK26 and

TNK50, with Spearman's correlation coefficients between prediction and observation being

ρ =0.76, *p*<2.2e-16 and ρ =0.68, *p*<2.2e-16, respectively. The UMAPs show that the

distributions of *PDCD1* and the two GEMs are highly correlated at the single-cell level (**Fig. 3B-**

C). The results also suggest potential PD-L1-PD-1 signaling is associated with distinct GEMs in

different T cell subtypes. Specifically, [*CD274*]*[*PDCD1*] is associated with TNK50 in

CXCL13$^+$CD4$^+$ T cells, whereas [*CD274*]*[*PDCD1*] is associated TNK26 mainly in exhausted

CD8$^+$ T cells. Importantly, the approximated ligand-receptor complex [*LR*] is more predictive of

the expression of these two GEMs when compared with only using the receptor expression [R]

alone. Similarly, the results show that the expression values of TNK35 can be predicted by

approximated *CD86* and *CTLA4* complex with Spearman's ρ =0.59, *p*<2.2e-16. The expression

of *CTLA4* is also highly overlapped with the expression of TNK35 in Treg cells (**Fig. 3B-C**),

suggesting the formation of this complex at these cells is associated with expression of the GEM

in a non-linear dose-response fashion. In summary, our results show that identifying GEMs

enabled the investigation of the functional impact of potential ligand-receptor signaling. We

visualized a subset of ligand-receptor pairs and TNK GEMs that exhibited statistically significant

nonlinear dose-response relationships (**Extended Data Fig. 6 & Table. 2**).

The strong non-linear dose-response relationship between a ligand-receptor pair and a GEM (e.g.,

CD274-PDCD1 and TNK26) suggests the hypothesis that the expression of some members of a

GEM may be regulated by the ligand-receptor interaction. Thus, PD-1 signaling pathway is

likely to be more active in tumors expressing a high level of TNK26, and thereby, these tumors

might be more susceptive to anti-PD-1 treatment. To test the association of the expression level

of TNK26 and response to anti-PD-1 related treatment, we collected clinical and transcriptomic

data from Chen et al.'s CRC cohort treated with anti-PD-1 plus chemotherapy [4], which involves

20 CRC patients. The transcriptome and clinical responses to the treatment were measured pre-,

on-, and post-treatment. The patients were classified as CR (complete response), PR (partial

response), and SD (stable disease), based on the classification provided in the study. We

compared the expression level of GEMs learned from nHDP model in these three groups using

the pre-treatment transcriptome to identify discriminative GEM features. The analysis indicates a

higher TNK26 expression associated with CR group compared to the PR and SD group (**Fig. 3D**).

This association was further tested across one established CRC studies involving patients treated

with the combination of carbozantinib (a small molecule inhibitor of tyrosine kinases including

VEGFR2) and durvalumab (an immune checkpoint inhibitor targeting PD-L1) [28] and three

established melanoma studies [29-31] involving patients treated with anti-PD-1 antibodies. We

performed the wilcoxon test to identify discriminative GEMs. Consistently, TNK26 expression is

significantly higher in the responder group compared to the non-responder group (**Extended**

Data Fig. 7A & 7C). Moreover, by comparing the peak of TNK26 expression distribution before

and after treatment (**Fig. 3D**), we observed an apparent decrease following the anti-PD-1 therapy,

suggesting that TNK26 may be a downstream target of PD-1 signaling in the CR group. This

discovery is aligned with the statement from the original paper that CR patients experience a

transition from a hot to a cold immune state driven by effective tumor clearance. The results

support the notion that the expression level of TNK26 reflects the involvement of the PD-1

signaling in tumors, and thus can potentially be used as a novel biomarker for responsiveness to

the anti-PD-1 therapy. Regarding TNK50, a significantly higher expression in responder group

than in non-responder group is observed exclusively in the melanoma dataset. In the CAMILLA

trial, even though TNK50 is shown to be correlated with TNK26, TNK50 did not reach statistical

significance, likely due to the small number of responders with one apparent outlier, as depicted

in **Extended Data Fig. 7B**. More examples of GEMs showing significant difference between

responder group and non-responder group in CAMILLA trial and melanoma cohorts are listed in

Extended Data Fig. 7A & 7C, respectively.

Fig. 3: Discovering ligand-receptors strongly associated with specific pathways regulating the expression of GEMs. A. Illustration of GEMs reflective of downstream transcriptome mediated by biochemical signaling through ligand-receptor binding. The significant association between a GEM and a ligand-receptor pair needs to satisfy the sigmoid fitting. **B.** The overlap of cells expressing receptor and GEMs of interest. Left: the overlap of *PDCD1*, TNK26 and TNK50. Right: the overlap of *CTLA4* and TNK35. **C.** From left to right, the plot of sigmoid fitting between GEM and ligand-receptor pair with *p*-value and residual standard error (RSE), UMAP

plot of the receptor gene, and UMAP plot of the GEM. **D.** Comparison of the expression levels

of TNK26, TNK50 and TNK35 in the CR (complete response), PR (partial response) and SD

(stable disease) groups before (pre-) and during (on-) anti-PD-1 immunotherapy treatment.

The CCC between SPP1-macrophage and cancer-associated fibroblast through ligand-

receptor *SPP1-ITGAV*

In addition to the cell-cell communications of TNK cells in the TME, the significance of

crosstalk between macrophages and cancer-associated fibroblasts (CAFs) has garnered

increasing attention in recent studies [32-34]. Motivated by this, we embarked on an investigation

into this crucial aspect. After searching for highly correlated GEMs that are expressed in stromal

cells and myeloid cells, our analysis led to a pair of GEMs, Mye11-Stroma11, which exhibited

strong Spearman correlation ($\rho = 0.35$, p = 9.2e-06) at the pseudo-bulk RNA level (**Fig. 4A**). The

results led to a hypothesis that myeloid cells expressing Mye11 may communicate with CAF

cells expressing Stroma11. Based on the annotation of GEM distribution, Mye11 is expressed in

a separate subpopulation of M2 macrophage, Stroma11 is expressed in a subpopulation of tumor

promoting CAF. After searching for the *LR* potentially mediating such communication, we

identified a LR pair *SPP1/ITGAV*, which predict the expression of Stroma11 (**Fig. 4A**) in a

nonlinear dose-response fashion (p-value of sigmoid fitting being 0.0004). Examining the

distribution of the ligands and receptor genes and the above GEMs in UMAPs showed that *SPP1*

and Mye11 were co-expressed in a subpopulation of M2 macrophage subpopulation (**Fig. 4B**).

ITGAV, which encodes subunits of integrins, is widely expressed in stromal cells, including both

fibroblasts and endothelial cells. This suggests that integrin signaling may have distinct

downstream effects in different stromal subpopulations, with Stroma11 being one of them. This

claim is supported by the existence of other *GEM X-LR-GEM Y* triplet communications guided

through *ITGAV* (**Extended Data Table. 6**).

To reduce the false positive discoveries, we require the *GEM X-LR-GEM Y* triplet to pass both

conditional independence test and spatial colocalization test. Based on the conditional

independence test, Mye11 and Stroma11 become independent when conditioned on the presence

of *SPP1-ITGAV*, indicating a potential causal relationship. We further tested this triplet using

two spatial transcriptomic cohorts (metastatic CRC patients from Wu et al [35] and CRC patients

from Valdeolivas et al [36]) by analyzing the co-localization of the ligands, receptors, GEMs and

their interactions in the spatial domains. The HE slides in these two spatial cohorts are provided

in **Extended Data Fig. 10**. Using a CRC tumor section (SN048_A121573_Rep2) as an example,

the spatial expression of ligand (*SPP1*), receptors (*ITGAV*) and GEMs (Mye11 and Stroma11)

were highly correlated (**Fig. 4D**), indicating that cells expressing these genes and GEMs likely

interact within spatial domains. The spatial correlation of Mye11-Stroma11, Mye11-*SPP1*,

Stroma11-*ITGAV*, and Stroma11-*SPP1-ITGAV* were 0.78, 0.75, 0.41, and 0.51, respectively. The

p-values associated with the spatial correlations shown above are less than 1e-22 (See **Extended**

Data Table. 6). The spatial colocalization of this communication (Mye11-*SPP1/ITGAV*-

Stroma11) was observed across multiple spatial slides (14 slides). These results support the

hypothesis that SPP1+ macrophages communicate with CAFs via the *SPP1-ITGAV* ligand-

receptor pair.

The CCC between macrophage and endothelial cells through ligand-receptor *VEGFA-

*FLT1***

By analyzing the transcriptomic data from the CAMILLA trial involving combinational anti-PD-

L1 immunotherapy, Stroma20 emerged as the most discriminative GEM feature, showing strong

potential for predicting drug response in metastatic CRC patients (**Extended Data Fig. 7A**).

Stroma20 is expressed in a subpopulation of tip cell/angiogenic endothelial cells. The top three

associated TFs are BCL6B, SOX18, and SOX7 and the top enriched signaling pathway

associated with the leading genes of Stroma20 is NOTCH1 signaling. The relationship between

NOTCH signaling and VEGFR has been demonstrated in previous studies [37].

We then searched for the triplet pairs including Stroma20 that passed both the conditional

independence test and spatial colocalization test, and identified one significant communication:

Mye9-*VEGFA-FLT1*-Stroma20. Mye9 is a subpopulation of macrophage and dendritic cells, and

exhibited strong correlation with Stroma20 (Spearman $\rho = 0.44$, p = 2.7e-08) at the pseudo-bulk

RNA level (**Fig. 4A**). UMAPs in **Fig. 4C** show the co-expression of *VEGFA* and Mye9, and the

co-expression of *FLT1* and Stroma20. The spatial expression of ligand (*VEGFA*), receptors

(*FLT1*) and GEMs (Mye9 and Stroma20) were highly correlated, as shown in **Fig. 4E**. Using the

same slide in **Fig. 4D**, the spatial correlations for Mye9-Stroma20, Mye9-*VEGFA*, Stroma20-

FLT1, Stroma20-*VEGFA-FLT1* are 0.69, 0.56, 0.31, and 0.31, respectively. The p-values

associated with the spatial correlations shown above are less than 1e-22 (See **Extended Data**

Table. 6). This communication (Mye9-*VEGFA-FLT1*-Stroma20) is spatially colocalized in

multiple slides (six slides). Our findings provide insights into identifying potential biomarkers

and their associated communications for current and novel treatments that elucidate underlying

biological mechanisms.

The example discussed above demonstrates our ability to identify potential CCC candidates

mediated by the ligand-receptor signaling, supported by the integration of scRNA-seq (co-

expression), pseudo-bulk RNA-seq (correlation and conditional independence test) and spatial

transcriptomic data (spatial colocalization test).

Fig. 4: Modeling intercellular communication. A. Left two: The scatter plot of Mye11-vs-Stroma11 and Mye9-vs-Stroma20 using the pseudo-bulk transcriptome data; Right two: The sigmoid fitting between Stroma11 and ligand-receptor pair *SPP1-ITGAV*, and the sigmoid fitting between Stroma20 and ligand-receptor pair *VEGFA-FLT1* using the pseudo-bulk transcriptome data. B. The UMAP plot of cells expressing *SPP1*, Mye11, *ITGAV* and Stroma11. **C.** The UMAP plot of cells expressing *VEGFA*, Mye9, *FLT1*, and Stroma20. **D**. The spatial distribution plot of *SPP1*, *ITGAV*, *SPP1-ITGAV*, Mye11, Stroma11, and Mye11-Stroma11 using the spatial slide (SN048_A121573_Rep2) as an example. **E.** The spatial distribution plot of *VEGFA*, *FLT1*, *VEFGA-FLT1*, Mye9, Stroma20, and Mye9-Stroma20 using the same slide.

Discovering coordinated CCC programs

We further investigated patterns of coordinated expression of GEMs across different cell types (TNK, B, myeloid, stromal, and epithelial) in CRCs. We represented each tumor in the GEM space (pseudo-bulk expression: *cell-by-GEM* matrix) and performed double clustering of GEMs and tumors. The results (**Fig. 5A**) showed that the expression of many GEMs was highly coordinated. We hypothesized that the coordinated expression of GEMs across multiple types of cells in a TME is likely due to two possible mechanisms: environmental factors that orchestrate the expression of GEMs (confounding) or a CCC network that allows cells to communicate and reach homeostasis. We refer to a set of GEMs across multiple cell categories that displays a highly coordinated expression pattern as a CCC program, and we have identified 9 CCC programs (rows). Based on their GEM composition, clustering tumors revealed 6 sample clusters (columns). A complete list of CCC programs is in **Extended Data Table. 4**.

We further investigate the potential molecular basis of CCC programs. CCC Program 2 consists

of 16 GEMs from different cell categories (**Fig. 5B**). The pathway analysis indicates that GEMs

in CCC Program 2 are significantly enriched of genes involved in cell cycle pathways, such as

the C-MYC signaling pathway, a hallmark pathway of cell proliferation in different cell types [38].

The GEMs belonging to this CCC program include GEMs from TNK, B, myeloid, epithelial and

stromal cells, respectively (**Fig. 5B**), suggesting they are subpopulations of cells in a

proliferation stage from each cell category. Of interest, we noted that TNK9 is enriched in the T-

cell cluster #10, which is labeled as proliferative T cells. Not surprisingly, an epithelial GEM

belonging to this program, Epi9, is broadly active in epithelial cells, reflecting a shared

transcriptomic program involved in cell proliferation by cancer cells. The UMAPs in **Fig. 5C**

indicate that various cell categories expressing these GEMs also express the proliferation marker

gene *MKI67*. We hypothesized that the expression of GEMs of proliferation nature in different

cell types may be associated with signals that promote proliferation of cells, e.g., receptor

tyrosine kinase (RTK) pathways, Wnt/β-catenin pathway, and PI3K-AKT-mTOR pathway. Our

pathway enrichment analysis shows that the top genes associated with most of GEMs in CCC

Program 2 are enriched in the MYC pathway, which is downstream of RTKs. To test the

hypothesis, we investigated the spatial distribution of the GEMs in tumors. The results (**Fig. 5D**)

show the co-localization of GEMs in the CCC Program 2. We calculated the spatial correlations

between MKI67 with respect to B21, Stroma68, and Epi23, using their expression values in

spatial spots. The correlation coefficients are 0.708, 0.762, and 0.804, respectively, providing

evidence for the strong coordinated expression of these GEMs in spatial domains. We further

checked the cell composition of spatial spots expressing these GEMs. The inference of cell

composition in tumor spots (**Extended Data Fig. 8A**) shows that those spots contain a mixture

of cells potentially expressing these GEMs. The compositions of B21, Stroma68 and Epi23,

represented by their top associated genes, are not overlapping (**Extended Data Fig. 8B**). Each of

the three GEMs captures cell-type-specific proliferation signals. The indicates that the diverse

cells expressing GEMs in GEM program 2 may be stimulated by proliferation-related factors

existing in tissue microenvironments.

Moreover, we performed deconvolution analysis on the GEMs in CCC Program 2 utilizing the

single-cell RNA-seq transcriptome data obtained from the peripheral blood sample prior to the

treatment of the regorafenib and nivolumab combination therapy [39]. Our analysis revealed that

the expression levels of GEMs, including TNK9, B18, and Mye21, in the proliferation program

(CCC program 2), exhibited significant differences between responders and non-responders, as

illustrated in **Extended Data Fig. 9**. This observation aligns with the findings reported in the

original paper and highlights the potential of cellular signals related to MKI67 proliferation as

indicative markers for predicting clinical outcomes in patients treated with a combination of

multi-kinase inhibitors and immune checkpoint inhibitors.

We also searched for CCC programs in which coordinated expression of GEMs is likely due to

CCC between the GEMs. As an example, CCC Program 9 includes TNK26, TNK35, TNK50,

Mye72, Mye77, B57, and B6. To investigate whether cells expressing these GEMs communicate

with each other, we searched for LR pairs that were co-expressed with these GEMs, e.g., (*GEM*

X and *L*) and (*R* and *GEM Y*), as a candidate pair of communicating cells, using the pseudo-bulk

RNA-seq data. We found three main LR pairs through which GEMs in the program may

communicate, including *CD274-PDCD1*, *CD86-CTLA4*, and *CXCL13-CXCR5*, which are

closely associated with T-cell immune checkpoint molecules in the tumor microenvironment. We

illustrate the possible communication channels among the cells expressing those GEMs using

directed edges based on the co-expression of *GEM X-L* and *R-GEM Y* (**Fig. 5E**). The expression

level of the ligands and receptors in the cells expressing GEMs in the CCC Program 9 is shown

in **Fig. 5F**. The cells expressing TNK GMEs in CCC Program 9, such as TNK26 and TNK35,

mainly express receptor genes *PDCD1* and *CTLA4*. GEM B57 is concentrated in IgA/G cells

expressing the *CXCR5*, which is the receptor of *CXCL13*. Two myeloid GEMs, including Mye72

and Mye77, concentrated in dendritic cells, express ligand genes *CD274*, *CD80*, and *CD86*.

We then investigated how GEMs in CCC program 9, such as TNK26, communicate with GEMs

in other CCC programs. To reduce false-positive discoveries, we required the communication

triplets (*GEM X–LR–GEM Y*) to pass both the conditional independence test and spatial

colocalization. **Fig. 5G** highlights one of the most significant communications involving TNK26

that meets both criteria: TNK26—*CSF1-CSF1R*—Mye8. The Spearman correlations of TNK26-

Mye8, TNK26-*CSF1*, *CSF1R*-Mye8 in pseudo-bulk are 0.33 (*p*=3.5e-05), 0.39 (p=8.6e-07) and

0.57 (p<2.2e-16). The ligand, receptor, and two GEMs were shown to be colocalized in spatial

domains in 11 slides. **Fig. 5G** uses two slides as examples to show the significant colocalization.

The spatial correlations of TNK26-Mye8, TNK26-*CSF1*, and *CSF1-CSF1R*-Mye8 are 0.85, 0.78,

and 0.60 (upper slide: SN048_A416371_Rep2), and 0.64, 0.32 and 0.39 (lower slide:

SN048_A121573_Rep2), respectively. The p-values associated with the spatial correlations

above are all less than 1e-22 (See **Extended Data Table. 6**). Several studies have demonstrated

the importance of CSF1R signaling in CD8□ T cell exhaustion in cancer types such as breast

cancer [40-42], as well as the potential of CSF1R inhibitors as monotherapy or in combination with

other immunotherapies, including immune checkpoint inhibitors [43]. The strong association of

TNK26 with signaling pathways, such as *PD-L1-PDCD1* and *CSF1-CSF1R,* highlights the

potential of TNK26 and its related interactions as targets for mono/combination immunotherapy.

In summary, the findings presented in this section collectively illustrate the feasibility of

uncovering a unique CCC mechanism within a micro-TME. This is achieved by modeling the

synchronized expression of GEMs, identifying potential LRs responsible for mediating

communication, and providing evidence of spatial colocalization for the components involved in

the identified CCC channel.

Fig. 5: Discovering patterns of intercellular communication channels that define transcriptome programs across different cell types. A. Clustering of GEMs and CRC tumors. Pseudo-bulk profile for each tumor was constructed from single-cell RNA-seq data. Expression values of a GEM were estimated based on the GSVA using the top 50 weighted genes in a GEM. The GEMs (rows) are grouped into 9 CCC programs by clustering analysis, representing patterns of intercellular communication channels. The tumors (columns) are grouped into 6 clusters. **B.** The GEM members of the CCC Program 2. The majority of genes associated with these GEMs

are MYC pathway targets, as indicated by the HALLMARK_MYC_TARGETS in GSEA. **C.**

UMAP of MKI67 and GEMs in CCC program 2 in different cell-type categories. **D.** Spatial

colocalization of proliferation marker MKI67 and GEMs in CCC Program 2. **E.** The potential

CCC network and the ligand-receptors involved in GEMs in CCC Program 9. The red and blue

rectangle in the center represents ligands and receptors respectively. The edge between a

ligand/receptor and a GEM is added if the ligand/receptor is expressed in the cells expressing a

GEM. **F.** The expression level of receptors (blue) and its associated ligands (red) in cells

expressing the GEMs in CCC program 9. TNK and B GEMs in CCC Program 9 mainly express

receptors (*PDCD1, CTLA4, and CXCR5*). Myeloid GEMs mainly express ligands (*CD80, CD86*

and CD274). **G.** The spatial colocalization of ligand, receptor and GEMs: TNK26 -> *CSF1-*

CSF1R-> Mye8 in two spatial slides (SN048_A416371_Rep2 and SN048_A121573_Rep2).

Construction of cell-cell communication subnetworks connected by ligand-receptor

signaling

Discovering a pair of highly correlated GEMs across cell types suggests possible CCC between

the cells expressing the GEMs. Subsequently, we conducted a systematic search and examination

of potential CCC channels and networks. Using CRC pseudo-bulk RNA-seq data, we first

searched for highly correlated GEMs (*GEM X* and *GEM Y*) that are expressed in different cell

types (or clusters) to discover potential CCCs. In addition, we performed a causal analysis to find

potential ligands and receptors that mediate CCCs. To that end, we then search for *LR* pairs such

that *L* is co-expressed with *GEM X* and *R* is co-expressed with *GEM Y.* Finally, we assess

whether conditioning on *LR*, the two marginally correlated GEMs become independent, which is

denoted as *GEM X ⫫ GEM Y | LR*. Conditional independence is an important concept in causal

analysis, and in our setting, if a pair of highly correlated GEMs becomes independent when

conditioning on a LR pair, this LR pair is potentially involved in transmitting signals between

cells expressing *GEM X* and *GEM Y*. Finding a *GEM X-LR-GEM Y* triplet leads to the hypothesis

depicted in **Fig. 6A**. More specifically, a change in the state of the signaling pathway (*X*) in one

cell promotes the release of the ligand and, at the same time, leads to transcriptomic changes

represented by *GEM X*, which serves as a proxy of the pathway (*X*) state. The binding of the

ligand with its receptor in another cell results in a cellular effect represented by the signaling

pathway (*Y*), which leads to the changed gene transcription represented by *GEM Y*. Further

extending such CCC analysis to multiple GEMs and cell types leads to a CCC network (CCCN),

which may be used to understand how different cells reach homeostatic states in a TME through

CCCs. If there are GEMs/LR pairs of interest, a CCC sub-network can be constructed based on

the *GEM X-LR-GEM Y* triplets. We identified and illustrated a CCC sub-network related to

immune checkpoint signaling in **Fig. 6B**. The network consists of T cells expressing TNK GEMs

(TNK26, TNK35 and TNK50) and non-TNK cells expressing GEMs (Stroma37, B44, Mye21,

Mye43, and Mye46), likely transmitted through PD-1, CTLA4 and CXCL13 signaling. In this

network, the communication between cells expressing Mye9 (a combination of dendritic and

macrophage cells) and TNK35 (LAYN$^+$ Effector Treg) can be mediated through *CD274-PDCD1*;

the communication between the cells expressing Mye43 (TAM: tumor-associated macrophage)

and TNK35/TNK26 (PDCD1$^+$CXCL13$^+$ CD4$^+$ and CD8$^+$ T cells) is transmitted through

CD80/86-CTLA4; the communication between Mye26-expressing NLRP3$^+$INHBA$^+$ macrophage

cells and TNK50-expressing CXCL13 producing CD4$^+$ T cells is facilitated via *CD274-PDCD1*;

the communication between TNK50 and B44-expressing follicular B helper cell is transmitted

through *CXCL13-CXCR5* which is consistent with the current literature [44]. Moreover, we also

discovered the communication between a specific subset of mCAF and effector Treg. The

communication between Stroma37 expressing MMP1$^+$CXCL5$^+$mCAF cells and TNK35

expressing LAYN1$^+$Treg cells is transmitted through *CD274-PDCD1*. *MMP1* ranks the top 3rd

gene associated with Stroma37 and has been shown to play an important role in CAF-induced

protection and lead to therapy resistance [45]. The communication between stromal cells and Tregs

has been validated in several studies [46]. Besides, TNK51 expressing Th1 cells communicate with

myeloid GEMs (Mye8 and Mye42) expressing suppressor-like macrophage (M2) through *TNF-*

TNFRSF1B [47,48]. The above results demonstrate that re-constructing CCC networks provides

insights into potential molecular mechanisms of CCCN underlying immune evasion mechanisms

of tumors.

NK cells are a promising alternative platform for cellular immunotherapy, and clinical trials of

NK cell immunotherapy have produced many encouraging clinical results [49]. In **Fig. 6C**, a CCC

sub-network illustrates the primary communications between TNK32-expressing NK cells and

specific cellular functions of stromal, myeloid cells, and B cells through *ICAM1/3-LFA-1*, *IL10-*

IL10RA, and *IL15-IL15RB* interactions. Besides, we discovered a highly ranked potential CCC

channel involving Stroma22 expressing CAF, TNK14 expressing Treg and TNK50 expressing

CXCL13$^+$CD4$^+$ T cells through *OX40L-OX40,* which is a costimulatory receptor as shown in **Fig.**

6D. Recent studies have shown the benefit of combining OX40 agonist and PD-1 inhibitor to

improve the immunotherapy outcome [50-52]. In both the CAMILLA trial dataset, where combined

therapy involved a VEGFR2 inhibitor and anti-PD-L1, and the dataset from melanoma patients

treated solely with anti-PD-1 immunotherapy, Stroma22 exhibits a notable difference between

the responder and non-responder groups. (**Extended Data Fig. 7A**). In addition to the

aforementioned findings, we have identified additional *GEM X-LR-GEM Y* triplets as potential

CCCNs in colorectal cancer (CRC). These discoveries offer valuable insights into exploring

intercellular communication within the TME and understanding the key cellular transcriptomic

processes associated with ligand-receptor signaling. The details are listed in **Extended Data**

Table. 3. To enhance the user-friendliness of our framework, we've created an interactive

website where users can explore the annotation of each GEM, identify the top-ranked GEMs

associated with each ligand-receptor pair, and investigate the *GEM X-LR-GEM Y* triplets of

interest (http://44.192.10.166:3838/).

Fig. 6: The newly discovered cellular networks associated with ligand-receptor interactions involved in the immune checkpoint inhibitor therapy. A. The framework of the conditional independence analysis to find the causal relationship between GEM-expressing cells transmitted by ligand-receptor signals. **B.** Subnetwork of intercellular communication related to PDL1/2-PD-1, CD80/CD86-CTLA4, and CXCL13-CXCR5 based on the conditional independence analysis.

572   The detailed annotation of each GEM is listed in the interactive website. **C.** Subnetwork of

573   intercellular communication related to TNK32 which represents the cellular function of NK cells.

574   **D.** Subnetwork of intercellular communication between Stroma22 and GEMs from other cell

575   types through the OX40L-OX40 signaling.

Discussion

In this study, we showed that by deconvoluting the hierarchically organized transcriptomic

programs and closely representing cell differential programs, the nHDP model can identify

GEMs that are potentially regulated through CCC, discover possible CCC channels, and generate

the hypothesis of the molecular mechanisms of CCC underlying heterogeneous tumor

microenvironments of CRC. Detecting and representing the states of cellular signaling systems

in individual cells is a fundamental task in single-cell biology. While there are different

approaches for deconvoluting single-cell transcriptomic components [18,53], our results indicate

that by modeling the hierarchical and compositional relationships among transcriptomic

programs of cells, the nHDP model captures fine-grained transcriptomic signals better than other

non-hierarchical models.

Although single-cell RNA-seq assays offer unparalleled cell-type-specific insights, their practical

use is impeded by several obstacles (e.g. high cost of library preparation and sequencing) that

hinder their broad application in clinical contexts. The evolution of bulk RNA-seq data over the

past decades has yielded a substantial volume of samples conducive to the execution of pattern

and machine learning investigations. The study of CCC, especially the TME associated with

drug treatments, could be further promoted by integrating insights leaned from scRNA-seq data.

Therefore, effective methods for bulk RNA-seq deconvolution are still urgently needed. The

GEMs learned from nHDP could be used as novel signatures for deconvoluting the specific

transcriptomic processes in both bulk RNA-seq data and spatial transcriptomic data. Compared

with two state-of-the-art statistical models, non-negative matrix factorization (NMF) and latent

Dirichlet allocation (LDA), for identifying transcriptomic programs, nHDP model has the

advantage of capturing the hierarchical structure of signaling pathways. GEMs derived from the

nHDP model reflect the cellular states across multiple cell subtypes or specific to a particular cell

subtype. Our recent study [54] showed that GEMs offered a more precise estimation of biological

signals, such as immune cell infiltration, in comparison to the state-of-the-art deconvolution

methods like Cibersort and SCNA [55,56]. This improved accuracy enables the identification of

different lymphocyte infiltration levels.

Identifying highly specialized GEMs enabled us to use the expression level of GEMs as proxies

to represent the state of signaling pathways (unobserved) that regulate their expression. This, in

turn, enabled us to search ligand-receptors that likely transmit extracellular signals to the

pathways inside single cells, generating the hypothesis of how across-cell signaling influences

intracellular signaling through CCC. Our results indicate that the *CD274-PDCD1* ligand-receptor

pair may influence the state of the cellular signaling pathways that regulate the expression of

certain members of TNK26 in exhausted T cells. The hypothesis is supported by multiple lines of

evidence: The non-linear dose-response relationship between the approximated ligand-receptor

complex and the GEM provides support from a biochemical point of view. The significant

difference in TNK26 expression between responding and non-responding tumors treated with

anti-PD-1 agents (both newly developed clinical trials in CRC and melanoma datasets) indicates

that tumors with high TNK26 expression display active PD-L1-PD-1 signaling, rendering them

more susceptible to the treatment. Conversely, tumors with low TNK26 expression exhibit

relatively inactive PD-L1-PD-1 signaling, leading to their reduced responsiveness to the

treatment. Based on these results, we hypothesize that the PD-L1-PD-1 signaling likely regulates

some members of the TNK26. Further investigation of which members are regulated by this

signal, e.g., using principled causal discovery methods, may enable researchers to delineate

mechanisms of PD-1 signaling. Similar studies of the impact of other commonly co-expressed

immune checkpoint proteins (receptors) in tumor-experienced T cells may reveal how different

checkpoint signaling pathways cooperate and lead to immune suppression.

The discovery of a strong correlation of GEMs across cell types indicates that the expression of

GEMs by different cell types is coordinated, and such coordination is likely due to CCC. We

identified coordinated GEM expression patterns, referred to as CCC programs, and their spatial

colocalization provides additional support for intercellular communication networks. We further

noticed the differences in the activities of CCC programs may underlie the heterogeneity of

TMEs, leading to tumor clusters that recapitulate well-appreciated molecular subtyping of CRC.

Further investigation identifies ligand-receptors that likely transmit signals across GEMs in CCC

channels.

We anticipate further investigating CCC channels active in each tumor/subcluster may provide

mechanistic insights into the TME heterogeneity. In our population-wide exploration of CCC, we

observed instances where multiple ligands could communicate with a single receptor, as

illustrated by the TNK32 in **Fig. 6C**. In the causal theory, those ligands are multiple parents of

the receptor. If we examine whether instantiated values of variables in a local network lead to

context-specific independence, only certain parents may contribute to the state of children from

the aspect of an individual tumor sample. For instance, in an individual tumor sample, only one

type of cell among stromal, myeloid, or B cells may be involved in the interaction with TNK32-

expressing cells. Our forthcoming efforts will involve employing the individualized Causal

Bayesian Network (iCBN) model [57] to reveal the individualized CCCN within each tumor

sample. This effort will contribute to a deeper understanding of the mechanisms underlying

heterogeneity. Besides, this study approximates the ligand receptor interaction based on the

product of [L] and [R] due to the feasibility. We realized that this approximation method using

the gene expression is insufficient. However, since gene expression is one type of data available

as a surrogate for ligand/receptor proteins, the method applied in this study is among the widely

used approximation approach by the research community [5,6].

We envision that the framework presented in this study for deconvoluting single-cell

transcriptomic programs to model CCC can be applied to various cancer types. Enhancing the

nHDP model involves refining the initialization process and filtering variable genes, leading to a

more effective capture of relationships within transcriptomic programs. This improvement will

boost the model's ability to detect GEMs regulated by specific transcriptomic programs. Future

methodological developments for detecting fine-grained transcriptomic programs may involve

leveraging recently developed topic modeling models to capture the compositionally co-

expressed genes. Additionally, integrating single-cell RNA-seq with single-cell chromatin

accessibility sequencing (scATAC) could aid in discovering active motifs associated with CCC

networks. In conclusion, this study advances the understanding of CCCs mechanisms underlying

heterogeneous TME and immune evasion mechanisms in CRC. These insights may uncover

novel targets for immunotherapy and guide precision immunotherapy [23].

Online Method

Data Collection and Preprocessing

Single-cell transcriptomics

We collected single-cell RNA-seq data from 8 publicly available colorectal cancer data sources

(**Extended Data Table. 1**) [24,35,58,59]. The total number of patients is 153, of which there are 179

tumor samples and 100 normal samples. We retained cells that met the following criteria: nUMI >

500, nGene > 200, log10GenesPerUMI > 0.8, and mitoRatio < 0.2. To perform gene-level

filtering, we retained only genes expressed in at least 0.1% of the cells. After the quality control

(QC) analysis, we applied the total-count correction (library-size correction) and normalized the

data matrix to 10,000 reads per cell. The annotations of cell major type were obtained from

original publications and used for further analysis. We then divided cells into five major

categories: 246,560 T and natural killer (TNK) cells, 94,663 myeloid cells, 96,637 plasma and B

cells, 69,130 [6]stromal cells, and 119,870 epithelial cells. When performing clustering analysis

within each cell category, the Harmony algorithm [60] was applied to remove the batch effect, cell

clustering was carried out using the Leiden algorithm [61], and the UMAP (Uniform Manifold

Approximation and Projection) algorithm was used to visualize clusters [62]. Seurat pipeline was

used to find subclusters and their top variable genes within TNK, myeloid, plasmaB, stromal,

and epithelial cells, respectively [15,63]. The cell types were annotated with well-known marker

genes: *CD3D, CD8A, FOXP3, IL7R, NKG7, GNLY* for T and NK cells; *CD79A, MS4A1, MZB1*

for B and plasma cells; *CD14, CD68, CD1C, S100A8, C1QA, LYZ* for myeloid cells. The

detailed markers we used were listed in **Extended Data Table. 5**.

Bulk RNA-seq

The bulk RNA-seq cohorts collected in this study mainly serve for discovering potential GEM

features strongly associated with the anti-PD-1 immunotherapy. Bulk RNA-seq data of 20

patients with metastatic CRC, treated with the combination of Cabozantinib plus durvalumab in

the CAMILLA clinical trial and characterized by proficient microsatellite stable (MSS) tumors,

were collected upon request [28]. The number of responders and non-responders is 4 and 16,

respectively. The responders include the CR (complete response) and PR (partial response) and

the non-responders include the PD (progressive disease) and SD (stable disease). Three

melanoma bulk RNA-seq data with response to anti-PD-1 treatment were downloaded from Liu

et al, Riaz et al, and Gide et al for the validation analysis [29-31].

Pseudo-bulk RNA-seq

To facilitate the clustering of GEMs and the searching for ligand-receptor pairs using the CRC

single-cell data, we generated pseudo-bulk RNA-seq results for tumor samples (a *tumor-by-*

matrix). For the tumor samples (n=151) that include both $CD45^-$ and $CD45^+$ cells, UMI counts of

the same gene in each sample were aggregated, resulting in the expression data in raw count

format. We then normalized the raw count expression data to $log_2(TPM+1)$.

Spatial transcriptome

Spatial transcriptomic data of colorectal cancer and its corresponding liver metastasis using 10x

Visium technology were collected from the study by Wu et al [35] and Valeolivas et al [36]. These

two cohorts contain 8 and 16 slides, respectively. As shown in **Extended Data Fig. 10**, these two

cohorts contains 8 slides (Gao_colon1, Gao_colon2, Gao_colon3, Gao_colon4, Gao_liver1,

Gao_liver2, Gao_liver3, and Gao_liver4) and 16 slides (SN048_A121573_Rep1,

SN048_A121573_Rep2, SN048_A416371_Rep1, SN048_A416371_Rep2,

SN84_A120838_Rep1, SN84_A120838_Rep2, SN84_A121573_Rep1, SN84_A121573_Rep2,

SN123_A595688_Rep1, SN123_A798015_Rep1, SN123_A551763_Rep1,

SN123_A938797_Rep1_X, SN124_A551763_Rep2, SN124_A938797_Rep2,

SN124_A595688_Rep2, SN124_A798015_Rep2), respectively. We first used sctransform

method [64] to normalize the raw count in each spot of the slide (a *spot-by-gene* matrix). The mean

of top 50 genes associated with each GEM was then used to estimate the GEM values from

normalized expression data. The deconvoluted spatial transcriptomic data is represented as a

spot-by-GEM matrix. Function RCTD (R package spacexr) [65] was used to infer the cell-type

composition in tumor region.

Identification of GEMs using nHDP

We used the nested hierarchical Dirichlet processes (nHDP) model [54,66] to identify patterns of

GEMs from single-cell RNA-seq data. The flow of the training process is shown in **Fig. 1B &**

1C. We binarized the UMI count of each gene within a cell as the input of the nHDP model. If a

gene is expressed in a cell with non-zero value, it will be assigned a binary value 1; Otherwise, it

will be assigned a binary value of 0. A trained nHDP model returns two matrices: a cell-by-GEM

count matrix, which collects the counts of UMIs assigned to a GEMs in each cell; a GEM-by-

gene matrix, which represents the probability profile of genes in each GEM (**Fig. 1B**). We

constructed a three-layer tree structure with the following branching factors: 5, 4, and 3 for 1^{st},

2^{nd}, and 3^{rd} layers, respectively, as shown in **Fig. 1C**. In this model, the choice of tree structure

(e.g., 5-4-3) was guided by biological priors such as the expected number of major lineages,

ensuring that top level GEMs capture common programs while deeper nodes refine subtype- and

function-specific signals. We systematically evaluated alternative tree structures (3-2-2, 4-3-2, 5-

4-3) and demonstrated that larger and deeper trees yielded consistently lower perplexity and

higher topic coherence, particularly as the training size increased[54]. The switching parameter γ

was set to balance exploration of deeper nodes with probability ranging from 0 to 1, and mini-

batch training with L1 K-means initialization further improved robustness against noise. To

validate reproducibility, we performed simulation studies and tested independent datasets

(PBMC 3K, 8K, 33K) repeatedly, finding consistent GEM structures and transferability across

donors. Together, these analyses demonstrate that the nHDP model is robust to parameter

choices when informed by biological context, and reproducible across independent runs and

datasets.

The expression of GEMs (cell-by-GEM matrix) is added to the Seurat object as new features for

plotting the UMAP of GEMs. The dominant function represented by a GEM is based on the rank

of cell subtype percentage ("Distribution" in **Extended Data Fig. 4**). We used word clouds R

package "wordcloud2" to visualize the probability profile of the top 20 associated genes of a

GEM (**Fig. 1D**). We trained a nHDP model for each cell category (TNK, B, Mye, Stroma, and

Epi) and retained GEMs expressed in over 1% of cells in a category for further analysis.

Functional analysis of GEMs

Transcription factor and enrichment analysis were applied to conduct functional analysis on GEMs. CHEA3 was used to conduct the transcription factor analysis [25]. Reactome, KEGG and GO databases were used to conduct comprehensive pathway analysis using R package "msigdbr" [67]. Besides, we conducted a hypergeometric analysis [68] to annotate GEMs with well-known gene sets (H and C1-C8) listed in MsigDB [69]. The significant enrichment was defined as p-value <= 0.05 & FDR <= 0.05.

Estimation of GEM level in bulk and spatial RNA-seq data

To estimate the expression level of a GEM in bulk/pseudo-bulk/spatial RNA-seq data, we used two methods: 1) gene set variation analysis (GSVA) of the top 50 weighted genes in each GEM [70]. 2) mean expression of the top 50 weighted genes in each GEM.

Fitting non-linear relationship between ligand-receptor pairs and GEMs

The ligand-receptor database was downloaded from Nichenet [6,21,22]. To fit the dose-response curve between the expression of ligand-receptor and GEM value, we model the quantity of LR complex [LR] as the product of expression of L and R based on the fact that $[LR] \propto [L][R]$ using the pseudo-bulk RNA-seq data. We first scaled the GEM value to [0, 1] and then created a 4-parameter log-logistic function as $f(x) = c + \frac{d-c}{1+\exp(b(\log(x)) - \log(e)))}$, where d and c on 0 and infinite dose, e is the point of inflection and b being the hill slope using *drc/v3.0.1* [71]. The coefficients and its associated *p*-values were estimated and adjusted with Benjamini-Hochberg

procedure[22]. Sigmoid fitting with a significant adjusted *p*-value (<0.05) demonstrates a good fit

between the expression of LR complex and the scaled GEM values.

Assessing conditional independence of *GEM X-LR-GEM Y* triplets

To find out the ligand-receptor pairs that potentially mediate CCC between cells across cell

categories expressing two correlated GEMs, conditional independence tests of *GEM X ⫫ GEM Y*

| LR were performed using the *bnlearn/v4.7.1* package in R [72]. We estimated expression status of

each GEM using the GSVA scores (*GSVA/v1.42.0*) [70] based on its top 50 genes. We calculated

all pairwise correlations between GEMs expressed in different cell categories. If a pair of cross-

cell-category GEMs exhibits significant correlation, we further search for LR pairs that rendered

conditional independence (*GEM X ⫫ GEM Y | LR*). *P*-values of conditional independence tests

were adjusted to control the false discovery rate using Benjamini-Hochberg method. We set the

threshold of the conditional correlation between *GEM1* and *GEM2* at an adjusted *p*-value > 0.05

as conditional independent.

We further performed post-hoc proportional tests to remove false positive findings. For each

conditional independent *GEM X—LR—GEM Y* triplet, we investigated the co-expression of *GEM*

X—L in potential signal-originating cells and the co-expression of *R—GEM Y* in potential signal-

receiving cells. To conduct the proportional tests, we first construct a 2 by 2 contingency table to

measure the frequency of co-expression GEM and the tested gene. GEM and gene were

considered presenting if the corresponding values are beyond 0 and not presenting otherwise.

Then the statistics of proportional tests $z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$ were estimated to test if the proportion of

co-expressing cells is significantly higher than the proportion of not co-expressing cells, where $\hat{p}$

is the proportion of cells that co-express GEM and tested gene and p_0 is the proportion of cells

that not co-express GEM and tested gene, n is the total number of cells that express such GEM.

p-value of each post-hoc test was also adjusted using the Benjamini-Hochberg method for false

discovery rate control [73]. Ligand-receptor pairs that passed both conditional independence test

and post-hoc proportional test were considered valid for downstream analysis.

To reduce false positives, we applied multiple criteria to filter GEM pairs. At the pseudo-bulk

level, the correlations of *GEM X–GEM Y*, *GEM X–L*, and *R–GEM Y* were all required to exceed

0.3. In addition, we examined expression at the single-cell level to confirm the presence of

specific molecules (ligand/receptor genes) in GEM-expressing cells, thereby further reducing

false positives. Specifically, we required that more than 10% of cells expressing GEM X also

express ligand gene *L*, and more than 10% of cells expressing *GEM Y* also express receptor gene

R.

Assessing Spatial Colocalization

Spatial colocalization was evaluated by calculating spot-wise spatial correlations. To account for

soluble ligands, we assigned a weight of 1 to the center spot and first-layer neighbors, and a

weight of 0.5 to second-layer neighbors. Because each spatial spot contains a mixture of cells,

we inferred the cell composition within each spot (with the percentages of different cell types

summing to 1) using RCTD [65].

We then multiplied the GEM value by the cell type percentage to estimate the expression of

GEMs for a specific cell type. We calculated five types of correlations: (1) *GEM X-L* (ligand), (2)

GEM X-GEM Y, (3) *R* (receptor)-*GEM Y*, (4) *GEM X-LR*, and (5) *LR-GEM Y*. For *GEM X-L* and

GEM X-GEM Y correlations, we used lagged *GEM X* values derived from the two layers of

neighboring spots. For *GEM X-L* and *R-GEM Y* correlations, we first computed the ligand–

receptor product using the lagged ligand value, and then computed spot-wise correlation between

ligand–receptor product and *GEM X* or *GEM Y*. Statistical significance of each correlation was

assessed using Pearson's correlation test to obtain p-values. Spatial pairs were considered

colocalized only if the correlations of *GEM X-GEM Y*, *GEM X-L* and *LR-GEM Y* all exceed 0.3.

We then multiplied the GEM value by the cell type percentage to estimate GEM expression for a

specific cell type. We calculated five types of correlations: (1) *GEM X–L* (ligand), (2) *GEM X–*

GEM Y, (3) *R* (receptor)–*GEM Y*, (4) *LR–GEM Y*. For *GEM X–L* and *GEM X–GEM Y*

correlations, we used lagged GEM X values derived from two layers of neighboring spots. For

LR–GEM Y correlations, we first computed the ligand–receptor product using the lagged ligand

value, and then calculated spot-wise correlations between the ligand–receptor product and *GEM*

Y. The statistical significance of each correlation was assessed using Pearson's correlation test to

obtain p-values. Spatial pairs were considered colocalized only if the correlations of *GEM X–*

GEM Y, *GEM X–L*, and *LR–GEM Y* all exceeded 0.3.

Identifying programs of GEMs

Hierarchical clustering was performed on a Euclidean distance matrix [74] to identify GEM

programs using the pseudo-bulk expression of GEMs (sample-by-GEM matrix) as input.

Evaluating GEMs as potential biomarkers for anti-PD-1 therapies

We performed GSVA analysis to deconvolute the expression levels of GEMs for each bulk

tumor sample. Samples were divided into the response group if they showed complete response

(CR) or partial response (PR) to anti-PD-1 whereas non-response group if progressive disease

(PD) or stable disease (SD) were observed. We employed the one-sided Wilcoxon rank test [75] to

identify GEMs that exhibited a significant difference in expression levels between these two

groups.

Availability

nHDP model designed for analyzing single-cell transcriptome (scGEM):

https://github.com/hansolo-bioinfo/scGEM

CRC Atlas: https://github.com/hansolo-bioinfo/CRCAtlas

Interactive website: http://44.192.10.166:3838/

Ethics

No animal or human subjects is involved in this study.

Acknowledgments

This research was supported in part by The National Library of Medicine, National Human

Genome Research Institute, and National Cancer Institute at the National Institutes of Health

864   grants R00LM013089, R01HG014023, R01LM012011 and R01CA254274. The Department

865   specifically disclaims responsibility for any analysis, interpretations or conclusions.

Reference

1   Xi, Y. & Xu, P. Global colorectal cancer burden in 2020 and projections to 2040. *Transl*
Oncol **14**, 101174, doi:10.1016/j.tranon.2021.101174 (2021).
2   Sahin, I. H. *et al.* Immune checkpoint inhibitors for the treatment of MSI-H/MMR-D
colorectal cancer and a perspective on resistance mechanisms. *Br J Cancer* **121**, 809-818,
doi:10.1038/s41416-019-0599-y (2019).
3   Takahashi, N. *et al.* Dynamics of genomic and immune responses during primary
immunotherapy resistance in mismatch repair-deficient tumors. *Cold Spring Harb Mol*
Case Stud **6**, doi:10.1101/mcs.a005678 (2020).
4   Chen, Y. *et al.* Spatiotemporal single-cell analysis decodes cellular dynamics underlying
different responses to immunotherapy in colorectal cancer. *Cancer Cell* **42**, 1268-1285
e1267, doi:10.1016/j.ccell.2024.06.009 (2024).
5   Armingol, E., Officer, A., Harismendy, O. & Lewis, N. E. Deciphering cell-cell
interactions and communication from gene expression. *Nat Rev Genet* **22**, 71-88,
doi:10.1038/s41576-020-00292-x (2021).
6   Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat*
Commun **12**, 1088, doi:10.1038/s41467-021-21246-9 (2021).
7   Giladi, A. *et al.* Dissecting cellular crosstalk by sequencing physically interacting cells.
Nat Biotechnol **38**, 629-637, doi:10.1038/s41587-020-0442-2 (2020).
8   Shao, X., Lu, X., Liao, J., Chen, H. & Fan, X. New avenues for systematically inferring
cell-cell communication: through single-cell transcriptomics data. *Protein Cell* **11**, 866-
880, doi:10.1007/s13238-020-00727-5 (2020).
9   Marx, V. Method of the Year: spatially resolved transcriptomics. *Nat Methods* **18**, 9-14,
doi:10.1038/s41592-020-01033-y (2021).
10   Moses, L. & Pachter, L. Museum of spatial transcriptomics. *Nat Methods* **19**, 534-546,
doi:10.1038/s41592-022-01409-2 (2022).
11   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol*
36, 411-420, doi:10.1038/nbt.4096 (2018).
12   Luecken, M. D. & Theis, F. J. Current best practices in single-cell RNA-seq analysis: a
tutorial. *Mol Syst Biol* **15**, e8746, doi:10.15252/msb.20188746 (2019).
13   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression
data analysis. *Genome Biol* **19**, 15, doi:10.1186/s13059-017-1382-0 (2018).
14   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902
e1821, doi:10.1016/j.cell.2019.05.031 (2019).
15   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587
e3529, doi:10.1016/j.cell.2021.04.048 (2021).
16   Kotliar, D. *et al.* Identifying gene expression programs of cell-type identity and cellular
activity with single-cell RNA-Seq. *Elife* **8**, doi:10.7554/eLife.43803 (2019).

17   Zhu, X., Ching, T., Pan, X., Weissman, S. M. & Garmire, L. Detecting heterogeneity in
single-cell RNA-Seq data by non-negative matrix factorization. *PeerJ* **5**, e2888,
doi:10.7717/peerj.2888 (2017).
18   Liu, L. *et al.* Deconvolution of single-cell multi-omics layers reveals regulatory
heterogeneity. *Nat Commun* **10**, 470, doi:10.1038/s41467-018-08205-7 (2019).
19   Venkatasubramanian, M., Chetal, K., Schnell, D. J., Atluri, G. & Salomonis, N.
Resolving single-cell heterogeneity from hundreds of thousands of cells through
sequential hybrid clustering and NMF. *Bioinformatics* **36**, 3773-3780,
doi:10.1093/bioinformatics/btaa201 (2020).
20   Shao, C. & Hofer, T. Robust classification of single-cell transcriptome data by
nonnegative matrix factorization. *Bioinformatics* **33**, 235-242,
doi:10.1093/bioinformatics/btw607 (2017).
21   Efremova, M., Vento-Tormo, M., Teichmann, S. A. & Vento-Tormo, R. CellPhoneDB:
inferring cell-cell communication from combined expression of multi-subunit ligand-
receptor complexes. *Nat Protoc* **15**, 1484-1506, doi:10.1038/s41596-020-0292-x (2020).
22   Browaeys, R., Saelens, W. & Saeys, Y. NicheNet: modeling intercellular communication
by linking ligands to target genes. *Nat Methods* **17**, 159-162, doi:10.1038/s41592-019-
0667-5 (2020).
23   Chen, X. *et al.* An individualized causal framework for learning intercellular
communication networks that define microenvironments of individual tumors. *PLoS*
computational biology **18**, e1010761, doi:10.1371/journal.pcbi.1010761 (2022).
24   Pelka, K. *et al.* Spatially organized multicellular immune hubs in human colorectal
cancer. *Cell* **184**, 4734-4752 e4720, doi:10.1016/j.cell.2021.08.003 (2021).
25   Keenan, A. B. *et al.* ChEA3: transcription factor enrichment analysis by orthogonal
omics integration. *Nucleic Acids Res* **47**, W212-W224, doi:10.1093/nar/gkz446 (2019).
26   Martinez, G. J. *et al.* The transcription factor NFAT promotes exhaustion of activated
CD8(+) T cells. *Immunity* **42**, 265-278, doi:10.1016/j.immuni.2015.01.006 (2015).
27   Fan, J. *et al.* Zinc finger protein 831 promotes apoptosis and enhances chemosensitivity
in breast cancer by acting as a novel transcriptional repressor targeting the STAT3/Bcl2
signaling pathway. *Genes Dis* **11**, 430-448, doi:10.1016/j.gendis.2022.11.023 (2024).
28   Saeed, A. *et al.* Cabozantinib plus durvalumab in advanced gastroesophageal cancer and
other gastrointestinal malignancies: Phase Ib CAMILLA trial results. *Cell Rep Med* **4**,
100916, doi:10.1016/j.xcrm.2023.100916 (2023).
29   Liu, D. *et al.* Integrative molecular and clinical modeling of clinical outcomes to PD1
blockade in patients with metastatic melanoma. *Nat Med* **25**, 1916-1927,
doi:10.1038/s41591-019-0654-5 (2019).
30   Riaz, N. *et al.* Tumor and Microenvironment Evolution during Immunotherapy with
Nivolumab. *Cell* **171**, 934-949 e916, doi:10.1016/j.cell.2017.09.028 (2017).
31   Gide, T. N. *et al.* Distinct Immune Cell Populations Define Response to Anti-PD-1
Monotherapy and Anti-PD-1/Anti-CTLA-4 Combined Therapy. *Cancer Cell* **35**, 238-255
e236, doi:10.1016/j.ccell.2019.01.003 (2019).
32   Hoeft, K. *et al.* Platelet-instructed SPP1(+) macrophages drive myofibroblast activation
in fibrosis in a CXCL4-dependent manner. *Cell Rep* **42**, 112131,
doi:10.1016/j.celrep.2023.112131 (2023).

33   Qi, J. *et al.* Single-cell and spatial analysis reveal interaction of FAP(+) fibroblasts and
SPP1(+) macrophages in colorectal cancer. *Nat Commun* **13**, 1742, doi:10.1038/s41467-
022-29366-6 (2022).
34   Mao, X. *et al.* Crosstalk between cancer-associated fibroblasts and immune cells in the
tumor microenvironment: new findings and future perspectives. *Mol Cancer* **20**, 131,
doi:10.1186/s12943-021-01428-1 (2021).
35   Wu, Y. *et al.* Spatiotemporal Immune Landscape of Colorectal Cancer Liver Metastasis
at Single-Cell Level. *Cancer Discov* **12**, 134-153, doi:10.1158/2159-8290.CD-21-0316
(2022).
36   Valdeolivas, A. *et al.* Profiling the heterogeneity of colorectal cancer consensus
molecular subtypes using spatial transcriptomics. *NPJ Precis Oncol* **8**, 10,
doi:10.1038/s41698-023-00488-4 (2024).
37   Tanaka, M. *et al.* BCL6B Contributes to Ocular Vascular Diseases via Notch Signal
Silencing. *Arterioscler Thromb Vasc Biol* **43**, 927-942,
doi:10.1161/ATVBAHA.123.318987 (2023).
38   Wang, H. *et al.* c-Myc depletion inhibits proliferation of human tumor cells at various
stages of the cell cycle. *Oncogene* **27**, 1905-1915, doi:10.1038/sj.onc.1210823 (2008).
39   Kim, H. D. *et al.* Regorafenib plus nivolumab in unresectable hepatocellular carcinoma:
the phase 2 RENOBATE trial. *Nat Med* **30**, 699-707, doi:10.1038/s41591-024-02824-y
(2024).
40   Sato, T. *et al.* Sustained inhibition of CSF1R signaling augments antitumor immunity
through inhibiting tumor-associated macrophages. *JCI Insight* **10**,
doi:10.1172/jci.insight.178146 (2025).
41   Neubert, N. J. *et al.* T cell-induced CSF1 promotes melanoma resistance to PD1 blockade.
Sci Transl Med **10**, doi:10.1126/scitranslmed.aan3311 (2018).
42   Riaz, N. *et al.* Prognostic Significance of CSF-1R Expression in Early Invasive Breast
Cancer. *Cancers (Basel)* **13**, doi:10.3390/cancers13225769 (2021).
43   Wen, J., Wang, S., Guo, R. & Liu, D. CSF1R inhibitors are emerging immunotherapeutic
drugs for cancer treatment. *Eur J Med Chem* **245**, 114884,
doi:10.1016/j.ejmech.2022.114884 (2023).
44   Rao, D. A. T Cells That Help B Cells in Chronically Inflamed Tissues. *Front Immunol* **9**,
1924, doi:10.3389/fimmu.2018.01924 (2018).
45   Cui, Q. *et al.* Upregulating MMP-1 in carcinoma-associated fibroblasts reduces the
efficacy of Taxotere on breast cancer synergized by Collagen IV. *Oncol Lett* **16**, 3537-
3544, doi:10.3892/ol.2018.9092 (2018).
46   Sun, R. *et al.* Amphiregulin couples IL1RL1(+) regulatory T cells and cancer-associated
fibroblasts to impede antitumor immunity. *Sci Adv* **9**, eadd7399,
doi:10.1126/sciadv.add7399 (2023).
47   Wilk, A. J., Shalek, A. K., Holmes, S. & Blish, C. A. Comparative analysis of cell-cell
communication at single-cell resolution. *Nat Biotechnol*, doi:10.1038/s41587-023-01782-
z (2023).
48   Nonaka, K. *et al.* Th1 polarization in the tumor microenvironment upregulates the
myeloid-derived suppressor-like function of macrophages. *Cell Immunol* **369**, 104437,
doi:10.1016/j.cellimm.2021.104437 (2021).
49   Chu, J. *et al.* Natural killer cells: a promising immunotherapy for cancer. *J Transl Med* **20**,
240, doi:10.1186/s12967-022-03437-0 (2022).

50   van der Sluis, T. C. *et al.* OX40 agonism enhances PD-L1 checkpoint blockade by
shifting the cytotoxic T cell differentiation spectrum. *Cell Rep Med* **4**, 100939,
doi:10.1016/j.xcrm.2023.100939 (2023).
51   Ma, Y. *et al.* Combination of PD-1 Inhibitor and OX40 Agonist Induces Tumor Rejection
and Immune Memory in Mouse Models of Pancreatic Cancer. *Gastroenterology* **159**,
306-319 e312, doi:10.1053/j.gastro.2020.03.018 (2020).
52   Lao, J. *et al.* OX40 enhances T cell immune response to PD-1 blockade therapy in non-
small cell lung cancer. *Int Immunopharmacol* **108**, 108813,
doi:10.1016/j.intimp.2022.108813 (2022).
53   Im, Y. & Kim, Y. A Comprehensive Overview of RNA Deconvolution Methods and
Their Application. *Mol Cells* **46**, 99-105, doi:10.14348/molcells.2023.2178 (2023).
54   Zhang, H., Lu, X., Lu, B. & Chen, L. scGEM: Unveiling the Nested Tree-Structured
Gene Co-Expressing Modules in Single Cell Transcriptome Data. *Cancers (Basel)* **15**,
doi:10.3390/cancers15174277 (2023).
55   Davoli, T., Uno, H., Wooten, E. C. & Elledge, S. J. Tumor aneuploidy correlates with
markers of immune evasion and with reduced response to immunotherapy. *Science* **355**,
doi:10.1126/science.aaf8399 (2017).
56   Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression profiles.
Nat Methods **12**, 453-457, doi:10.1038/nmeth.3337 (2015).
57   Liu, Z. *et al.* A Novel Bayesian Framework Infers Driver Activation States and Reveals
Pathway-Oriented Molecular Subtypes in Head and Neck Cancer. *Cancers (Basel)* **14**,
doi:10.3390/cancers14194825 (2022).
58   Joanito, I. *et al.* Single-cell and bulk transcriptome sequencing identifies two epithelial
tumor cell states and refines the consensus molecular classification of colorectal cancer.
Nat Genet **54**, 963-975, doi:10.1038/s41588-022-01100-4 (2022).
59   Liu, Y. *et al.* Immune phenotypic linkage between colorectal cancer and liver metastasis.
Cancer Cell **40**, 424-437 e425, doi:10.1016/j.ccell.2022.02.013 (2022).
60   Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat Methods* **16**, 1289-1296, doi:10.1038/s41592-019-0619-0 (2019).
61   Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-
connected communities. *Sci Rep* **9**, 5233, doi:10.1038/s41598-019-41695-z (2019).
62   Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP.
Nat Biotechnol, doi:10.1038/nbt.4314 (2018).
63   Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of
single-cell gene expression data. *Nat Biotechnol* **33**, 495-502, doi:10.1038/nbt.3192
(2015).
64   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-
seq data using regularized negative binomial regression. *Genome Biol* **20**, 296,
doi:10.1186/s13059-019-1874-1 (2019).
65   Cable, D. M. *et al.* Robust decomposition of cell type mixtures in spatial transcriptomics.
Nat Biotechnol **40**, 517-526, doi:10.1038/s41587-021-00830-w (2022).
66   Paisley, J., Wang, C., Blei, D. M. & Jordan, M. I. Nested Hierarchical Dirichlet Processes.
IEEE Trans Pattern Anal Mach Intell **37**, 256-270, doi:10.1109/TPAMI.2014.2318728
(2015).
67   Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**,
1739-1740, doi:10.1093/bioinformatics/btr260 (2011).

68   Cao, J. & Zhang, S. A Bayesian extension of the hypergeometric test for functional
enrichment analysis. *Biometrics* **70**, 84-94, doi:10.1111/biom.12122 (2014).
69   Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set
collection. *Cell Syst* **1**, 417-425, doi:10.1016/j.cels.2015.12.004 (2015).
70   Hanzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for
microarray and RNA-seq data. *BMC Bioinformatics* **14**, 7, doi:10.1186/1471-2105-14-7
(2013).
71   Ritz, C., Baty, F., Streibig, J. C. & Gerhard, D. Dose-Response Analysis Using R. *PLoS*
One **10**, e0146021, doi:10.1371/journal.pone.0146021 (2015).
72   M, S. Learning Bayesian Networks with the bnlearn R Package. *Journal of Statistical*
Software **35**, 1-22, doi:10.18637/jss (2010).
73   Haynes, W. Benjamini–Hochberg Method. *Benjamini–Hochberg Method. In: Dubitzky,*
W., Wolkenhauer, O., Cho, KH., Yokota, H. (eds) Encyclopedia of Systems Biology.
Springer, New York, NY., doi:https://doi.org/10.1007/978-1-4419-9863-7_1215 (2013).
74   Murtagh, F. Algorithms for hierarchical clustering: an overview. . *Wiley Interdisciplinary*
Reviews: Data Mining and Knowledge Discovery **2**, 86-97 (2012).
75   Forrester, J. C. & Ury, H. K. The Signed-Rank (Wilcoxon) test in the rapid analysis of
biological data. *Lancet* **1**, 239-241, doi:10.1016/s0140-6736(69)91245-8 (1969).

1060

1061

1062

1063

1064

1065

1066

1067

1068

1069

1070

1071

1072 **Supplementary Figures**

1073

1074 **Supp Fig. 1: UMAP of B, Myeloid and Stromal cells. A.** The cell clusters for B and Plasma

1075 cells. **B.** The cell clusters for myeloid cells. **C.** The cell clusters for stromal cells.

1076

1077

1078 **Supp Fig. 2: Comparison between GEMs learned from nHDP and GEMs learned from**

1079 **NMF. A.** The cells expressing NMF5 are the combination of cells expressing TNK15, TNK42

1080 and TNK51. **B.** The cells expressing NMF58 are the combination of cells expressing TNK17,

1081 TNK29, and TNK50.

1082

1083

1084

1085

1086

1087

1088

1089

Supp Fig. 3: Cosine similarity between top50 GEM associated genes and highly expressed marker genes in T cell subtypes. A. Heatmap of cosine similarities between differentially expressed genes of T cell subtype and top 50 genes of all 85 T cell GEMs learned from nHDP model. GEMs are labeled with respect to their tree levels (L1, L2, and L3) and major T cell subtypes including CD4[+], CD8[+], ILC, MAIT, NK, Treg and proliferation T cells. **B.** The heatmap of cosine similarity for significant T cell GEMs summarized in **Fig. 2E** are shown.

1090

1091

1092

1093

1094

1095

1096

1097

1098

1099

1100

Supp Fig. 4: The dot plot comparing the similarity between GEMs and traditional cell

clusters. A. The dot plot for B and Plasma cells. **B.** The dot plot for myeloid cells. **C.** The dot

plot for stromal cells. **D.** The dot plot for TNK cells. The color and size of the dot reflect the

average expression and percentage of overlapped cells, respectively.

1101

1102

1103

1104

1105

1106

Supp Fig. 5: The enriched signaling pathways. A. The enriched signaling pathway for myeloid

cells. **B.** The enriched signaling pathway for B and plasma cells. **C.** The enriched signaling

pathway for stromal cells.

1110
1111
1112 **Supp Fig. 6: The summarization of top ligand-receptor pairs significantly associated with**

1113 **TNK GEMs**. The residual standard error (RSE) of the sigmoid fitting model against ligand-

1114 receptor expression production and GEM signals are shown in gradient color. Ligand-receptor

1115 pairs with significant p-value of sigmoid model are displayed.

1116

Supp Fig. 7: Example of GEMs significantly differentially expressed in responder group

and non-responder group. A. Example of GEMs significantly differentially expressed in

responder group and non-responder group treated with Cabozantinib plus durvalumab

(CAMILLA trial). **B.** The correlation between TNK26 and TNK50 in CAMILLA cohort. **C.**

Example of GEMs significantly differentially expressed in responder group and non-responder

group treated with solo anti-PD-1 immunotherapy (three independent melanoma datasets).

1123

A

Inference of cell-type composition in tumor region

🔴🟢 Fibroblast 🟡🔵 Lymphocyte 🟠🔵🟣 Epithelial cell

B

	B21	Stroma68	Epi23
B21	1	0	0
Stroma68	0	1	0.02
Epi23	0	0.02	1

1124
1125 **Supp Fig. 8: Exploring the uniqueness of three GEMs in CCC program 2. A.** The inference

 of cell-type composition in tumor region of **Fig. 5D**. **B.** The cosine similarity between B21,

 Stroma68 and Epi23 using the expression level of the top 50 associated genes.

1130 **Suppl Fig. 9: Example of GEMs in the proliferation program (CCC program2), which are**

1131 **significantly differentially expressed in responders and non-responders treated with**

1132 **regorafenib and nivolumab combination therapy.**

1133

1134

Suppl Fig. 10: HE slides in the two publicly available spatial transcriptomics cohorts used

in this study.

1135

1136

1137