Deconvoluting single-cell transcriptomics reveals cellular programs regulated by cell-cell communication in colorectal cancer

Han Zhang[1], Binfeng Lu[3], Gregory F. Cooper[1], Anwaar Saeed[2], John W. Paisley[4], Xinghua Lu[1], Lujia Chen[1]*

[1]Department of Biomedical Informatics, University of Pittsburgh, Pittsburgh, PA, 15206, USA

[2]UPMC Hillman Cancer Center, Pittsburgh, PA 15232, USA

[3]Center for Discovery and Innovation, Hackensack Meridian Health, Nutley, NJ 07110, USA

[4]Columbia University, New York, NY 10027, USA

*Correspondence: luc17@pitt.edu

Conflicts of Interest Statement:
Anwaar Saeed reports a leadership role with Autem therapeutics, Exelixis, KAHR medical and Bristol-Myers Squibb; consulting or advisory board role with AstraZeneca, Bristol-Myers Squibb, Merck, Exelixis, Pfizer, Xilio therapeutics, Taiho, Amgen, Autem therapeutics, KAHR medical, and Daiichi Sankyo; institutional research funding from AstraZeneca, Bristol-Myers Squibb, Merck, Clovis, Exelixis, Actuate therapeutics, Incyte Corporation, Daiichi Sankyo, Five prime therapeutics, Amgen, Innovent biologics, Dragonfly therapeutics, Oxford Biotherapeutics, Arcus therapeutics, and KAHR medical; and participation as a data safety monitoring board chair for Arcus therapeutics. The remaining authors have no relevant financial interests to disclose.

Abstract

Cells within a tissue microenvironment communicate through intricate cell-cell communication (CCC) networks. In this meta-analysis of eight single-cell cohorts encompassing 153 patients and 279 samples, we advance the understanding of CCC networks in colorectal cancers through a novel analytical framework. Employing hierarchical language modeling, we identify gene expression modules (GEMs) that mirror single-cell signaling states, crucial for deciphering the complexity of intercellular interactions. By applying causal discovery methods, we systematically uncover GEMs likely regulated by ligand-receptor signaling and cross-cell-type communication. This analysis reveals cross-cell-type CCC programs, marked by highly correlated GEMs across various

cell types, shedding light on the intricate CCC networks within the tumor microenvironment.

Spatial transcriptomics further validate these findings by demonstrating the co-localization of

GEMs within CCC programs in distinct spatial domains, emphasizing the spatial dynamics of

tumor intercellular communication. Our interactive website (http://44.192.10.166:3838/) and

analytical framework equip researchers with powerful tools to explore these complex mechanisms,

potentially uncovering novel drug targets and refining strategies for precision immunotherapies.

This comprehensive study not only presents a detailed catalog of CCC networks driven by ligand-

receptor interactions in colorectal cancer but also highlights the significance of integrating multi-

sample and patient data to unravel the molecular underpinnings of cancer communication

pathways.

Introduction

Colorectal cancer (CRC) is the 3[rd] most common cancer and the 2[nd] cause of cancer death[1]. The discovery of the consensus molecular subtypes (CMS) among CRC tumors reflects the heterogeneity of composition and functional states of cells in the tumor microenvironments (TME), which in turn may underlie diverse behaviors of disease progression and responses to anticancer treatments by tumors. For example, except for those with deficient mismatch repair (dMMR), most CRCs do not respond to anti-PD-1 immune therapy [2]. Such relationship between genomic pattern and clinical response is the result of cell-cell communications (CCC) within in the TME, through which the downstream signaling pathways are mediated and regulated. Therefore, the comprehensive understanding of the cell states and communication network in the TME are vital for uncovering the heterogeneous mechanisms of tumor growth, immune evasion and treatment resistance, which are essential for precision oncology.

Modeling CCC requires inferring the cellular states of individual cells, identifying synchronous changes in the cell-state among various cell types that may arise due to CCC, and investigating the molecular mechanism underlying CCC[3-6]. Single-cell RNA sequencing (scRNA-seq) is a powerful technology for profiling the transcriptomes of individual cells. As gene expressions are tightly regulated in a cell, the co-expression patterns of genes serve as proxies of the states of signaling pathways in a cell, which can be discovered by deconvoluting scRNA-seq data. Furthermore, the emergence of spatial transcriptomic technology provides additional spatial relationships between cells within their native spatial context [7,8]. Thus, integrating scRNA-seq with spatial transcriptome provides the spatial context for interactions among different cell populations [9] and enhances the accuracy of modeling CCC.

Single-cell transcriptomes can be used to investigate cellular states at different levels. At the whole

transcriptome level, cells sharing similar overall transcriptomic profiles can be discovered as

discrete cell clusters using state-of-the-art clustering techniques [10-13]. Although these approaches

are good at classifying different cell populations, it does not allow examination of transcriptomic

programs within individual cells that underpin the biological functions of these cells [14]. To discover

and represent transcriptomic programs within each cell, different statistical- and factorization-

based models, e.g., non-negative matrix factorization (NMF) and latent Dirichlet allocation (LDA),

have been used to deconvolute transcriptional programs by capturing the co-expression pattern of

certain cell subsets [15-18]. However, these models often have strong assumption of the independence

among gene programs. Moreover, these methodologies fail to capture the well-established

phenomenon that cells belonging to a lineage exhibit highly coordinated transcriptomic programs.

These programs are more accurately represented as a hierarchical structure, wherein transcriptomic

programs shared by numerous cells within a lineage are positioned close to the root of the hierarchy.

In contrast, transcriptomic programs underlying the differentiation of specialized cells are

positioned as leaves. The methods of capturing such hierarchical relationships of transcriptomic

programs remain relatively underdeveloped.

Transduction of signals across cells involves ligand-receptor interactions. Different methods have

been developed to find potential ligand-receptor-mediated intercellular communication, e.g.,

CellChat, CellPhoneDBv2.0, and Nichenet [4,19,20]. However, those methods are mainly based on

the expression and enrichment of ligand-receptor pairs in specific cell clusters but do not explicitly

model the impact of the signals transmitted by a ligand-receptor pair, e.g., genes regulated by PD-

L1/PD-1 signaling. Furthermore, few methods exist for modeling coordinated cellular states of

different cells within a tumor, e.g., certain subtype of myeloid cells and subtype of stromal cells

tend to co-exist in tumors, or inferring whether the changed state of one type of cell directly

influences the state of other types of cells.

To solve these challenges, we propose a novel framework for integrating bulk, single-cell, and

spatial RNA-seq data to depict a comprehensive landscape of CRC TMEs (**Fig. 1**). We employed

the nested Hierarchical Dirichlet process (nHDP) model to deconvolute a large collection of single-

cell transcriptomes from 8 cohorts with over 153 CRC patients and over 626,858 single-cells (**Fig.**

1A, Extended Data Table 1 & Fig. 1). nHDP organizes the discovered gene expression modules

(GEMs) in a hierarchy, enabling it to separate GEMs (transcriptomic programs) representing major

cell lineage markers from those reflecting the functional state of cells (**Fig. 1B-C**). We then

systematically investigated whether the expression of a GEM was regulated by ligand-receptor

signaling (**Fig. 1D**) using the hill function that is used in pharmacology to measure the binding

affinity between ligand and receptor. By studying the coordinated expression of GEMs, we

systematically investigated the statistically significant intercellular communications across cell

types (**Fig. 1E**) and discovered patterns of intercellular communication programs (GEM programs)

shared within subpopulations of tumors (**Fig. 1F**). Employing techniques of causal analyses, we

further searched for ligand-receptor pairs that contribute to mediate signal transduction of a CCC

and further constructed a CCC network among cells expressing coordinated GEMs (**Fig. 1G**). We

envision this comprehensive framework enables researchers to study complex mechanisms

underlying distinct TMEs, investigate different immune evasion mechanisms, discover potential

drug targets, and guide precision immune therapies.

Fig. 1: General design of the study. **A.** Merged CRC single-cell RNAseq data from 8 cohorts. **B.** Diagram of the nHDP model using the expression of genes (a *cell-by-gene* matrix) in TNK cells as input. The transcriptome of one cell can be represented by a unique mixture of GEMs in the nHDP tree. The nHDP model outputs two matrices: a *gene-by-GEM* matrix reflecting the gene

composition of a GEM and a *cell-by-GEM* matrix representing the GEM compositions of each cell.

C. nHDP discovered the tree-structured organization of GEMs. Each node in the tree represents a

GEM. A pie chart represents the percentage of each cell type the GEM is expressed. **D.** Illustration

of modeling the relationship between ligand-receptor (LR) and its downstream GEMs. **E.**

Modeling cell-cell communication by combining single-cell and spatial transcriptomes. **F.**

Learning patterns of coordinated expression of GEMs across cell types to discover distinct GEM

programs and associated tumor microenvironments. **G.** Downstream discoveries enabled by

deconvoluting single-cell transcriptomes by nHDP: **1)** GEMs highly associated with the signal of

LR. **2)** Subnetwork of intercellular communication for GEMs and LRs of interest. **3)** GEM as

potential novel biomarker for therapies.

Result

A comprehensive atlas of GEMs expressed by cells in CRC tumor microenvironments.

We collected single-cell RNA-seq data from 8 independent CRC cohorts. Then, we separated cells into 5 major lineages: TNK: mainly including T and natural killing (NK) cells (N=246,560); Myeloid: myeloid cells, including monocytes, macrophages, neutrophils, and dendritic cells (N=94,663); PlasmaB: including B and plasma cells (N=96,637); Stroma: stromal cells, including endothelial, fibroblast and myofibroblast cells (N=69,128); Epi: epithelial cells (N=119,870) (**Extended Data Table 1**). The merged cells for each cell category were clustered and labeled using Seurat pipeline and canonical marker genes. After the integration and batch removal, we identified 18 clusters in TNK cells (**Fig. 2A**), 18 cell populations in both Myeloid cells and PlasmaB cells and 17 cell groups in Stroma cells (**Extended Data Fig. 1**). We also showed that the distributions of datasets are well maintained in all clusters in TNK cells (**Fig. 2B**), suggesting the batch effects were regressed out; thus, the pooled TNK cells reflect the landscape of tumor-infiltrating TNK cells in CRC.

To model the tree structure of the TNK gene expression modules (GEMs), we utilized the nested hierarchical Dirichlet process (nHDP) model to identify a series of nested gene expression modules (GEMs) [21], where a GEM is a probability distribution over the gene space that captures the coordinated expression pattern of a set of genes in a subpopulation of cells. The nHDP model organizes inferred GEMs in a hierarchical tree structure, and we set branching factors of 5, 4, and 3 at 1^{st}, 2^{nd}, and 3^{rd} layers, respectively, leading to a total of 85 nodes to capture GEMs with different levels of granularity and specificity. **Fig. 2C** shows the tree organization of the GEMs learned using TNK cells as input. The nHDP model populated the nodes in the hierarchy through

a data-driven approach. Each node in the tree represents a GEM. The ones close to the root (TNK

GEMs 1-5 in **Fig. 2C**) represent GEMs that are more broadly expressed in all T, innate lymphoid

cells (ILC), mucosal-associated invariant T cells (MAIT) and NK cells, which can be used to

define major lineage of T and NK (TNK) cells. The ones close to the leaves represent GEMs

specifically expressed in more differentiated cells. In the TNK GEM tree, only 43 TNK GEMs

were identified as biologically meaningful and represented as filled nodes. The nodes identified as

biologically meaningless were shown as empty nodes without color.

We systematically inspected the TNK GEMs reflecting TNK transcriptomic processes. The pie

chart within each GEM illustrates the proportion of cells expressing that specific GEM across

various T cell subtypes, including $CD8^+$ T cells, $CD4^+$ T cells, ILC, MAIT, NK cells, regulatory

T cells (Treg) and Proliferation T cells. It is interesting to note that major branches of the nHDP

hierarchy tree, as defined by GEMs 1-4, represent the major TNK lineages, including CD8/NK,

CD4, Treg, and proliferation T cells (**Fig. 2C**). Projecting the expression of GEMs onto the UMAP

of TNK cells (**Fig. 2D**), we noticed that certain GEMs were concentrated in a particular cell cluster,

representing a characteristic transcriptomic program active in the cell cluster. For example, TNK50

is concentrated in cells belonging to cluster #4, which consists of T helper (Th) cells expressing

CXCL13; TNK35 is mainly expressed in T cell cluster #6, a subset of T_{reg}. Conversely, there are

also GEMs shared across multiple cell clusters, suggesting a transcriptomic program that is shared

across multiple subtypes of T cells. For example, TNK26 is expressed in a subset of

$CXCL13^+CD4^+$ T cells (cluster #4), $MKI67^+$ proliferative T cells (cluster #10), and

$CXCL13^+CD8^+$ T cells (cluster #11). TNK42 is expressed in MAIT cells (cluster #9) and ILC

(cluster #17). Finally, TNK12 is expressed broadly among Treg, Th, and proliferating T cells.

These results indicate that our model can capture transcriptomic programs that are both shared and

specialized by cells belonging to different lineages. To test the performance of our model, we also

compared the distributions of the transcriptomic programs that were found by our model and NMF

[22]. Our result demonstrates that the nHDP excels in capturing more specific cellular programs

shared across multiple cell clusters when compared to NMF as shown in **Extended Data Fig. 2**.

For instance, TNK15 captures a cellular program of greater specificity, shared among cells from

4 TNK cell clusters, surpassing the performance of NMF5. NMF5 captures a more generalized

cellular program that combines programs represented by TNK15, TNK42, and TNK51. Similarly,

NMF58 captures the cellular programs combining programs represented by TNK17, TNK29, and

TNK50.

To make the GEMs more biologically explainable, we depict the distinctive gene composition

profile of each GEM (top 20) through a word cloud, and the top three associated transcription

factors identified by transcription factor analysis using ChEA3 [23] were illustrated as black dots in

Fig. 2D. It is interesting to note that TNK26 is enriched with genes characteristic of exhausted T

cells including PDCD1, CXCL13 and TIGIT. Based on the transcription factor analysis, NFATC2

regulated the top 50 genes associated with TNK26. Several studies have shown that the

transcription factor NFAT is involved in the exhaustion process of activated $CD8^+$ T cells [24]. In

addition, the top genes associated with TNK26 and TNK50 are both regulated by transcription

factor ZNF831, which has been shown to inhibit the functional activity of effector cells through

STAT pathway [25]. This indicates the potential existence of a signal leading to the regulation of T

cell exhaustion. We further examined the distribution of GEMs in different TNK clusters (**Fig.**

2E), unveiling which GEMs are broadly shared among cells assigned to multiple clusters and

which GEMs are specific to a particular cell cluster. **Extended Data Fig.3** displays the detailed

comparison between GEMs and TNK cell clusters by measuring the correlation between top 50

genes of a GEM and highly variable genes in each TNK cell subtype. Furthermore, we performed

pathway enrichment analyses in Reactome to investigate the functions of top genes associated with

GEMs (**Fig. 2F**). Interestingly, several TNK GEMs, including TNK6, TNK26, TNK35, and

TNK41, are associated with PD-1 signaling pathway. The dot plots showing the distribution of

GEMs in traditional cell clusters and the enriched signaling pathways for other cell types (PlasmaB,

Myeloid, Stromal) are listed in **Extended Data Fig. 4 & 5**. The detailed annotation of each GEM,

including the UMAP, cell composition, and top associated genes, is listed in our interactive website.

In summary, by deconvoluting the transcriptomes of a large number of single cells using nHDP,

our approach revealed GEMs that represent transcriptomic programs within or across traditional

cell subtypes, which reflect certain signaling events within these cells.

Fig. 2: Deconvoluting single-cell transcriptomics to discover GEMs reflecting the state of signaling pathways. **A.** UMAP plot of 246,560 merged TNK cells from 8 CRC cohorts. **B.** The proportion plot reflects the origins of cells in each cell cluster. **C.** The hierarchical tree structure of TNK GEMs was learned from the nHDP model, which was trained with merged TNK cells. The nHDP model has three levels. The first level contains GEMs from 1-5. The second level

contains GEMs from 6-25. The third level contains GEMs from 26-85. Each color represents a

subtype of T cell. The unused GEMs are colored grey. **D.** The UMAP plot, word cloud plot, and

top associated TFs of selected TNK GEMs. **E.** The functional involvement of TNK GEMs within

each cell cluster. **F.** The enriched signaling pathways associated with TNK GEMs.

Discovering potential regulatory relationships between ligand-receptor signaling and GEMs

We set out to examine if the expression of a GEM in cells is potentially regulated by certain ligand-

receptor signaling by testing the hypothesis posited in **Fig. 3A**: If a ligand-receptor pair regulates

the expression of the member genes of a GEM, the overall expression value of the GEM can be

modeled with a sigmoid dose-response model with respect to ligand-receptor pair. We modeled

the relationship of the expression of ligand, receptor, and GEM at the tumor sample level. We first

generated pseudo-bulk RNAseq (a *tumor-by-GEM* matrix) from the scRNAseq (a *cell-by-GEM*

matrix) for tumor samples. Then we estimated the concentration of ligand and receptor proteins

based on the pseudo-bulk expression (see Methods) of the ligand ($[L]$) and receptor ($[R]$) for each

of the 151 tumors. Additionally, we approximated the concentrate of protein complex as the

product of ligand and receptor: $[LR] \propto [L]*[R]$ [3]. Pseudo-bulk expression of GEMs was estimated

using the GSVA analysis. Subsequently, we conducted a systematic search against all GEMs with

known LR pairs from NicheNet database and retained the LR pairs that could predict the

expression of the GEM with statistical significance (**Extended Data Table 2**).

We investigated the GEMs that are strongly associated with immune checkpoints programmed cell

death protein 1 (PD-1, encoded by *PDCD1*) and cytotoxic T lymphocyte-associated protein 4

(*CTLA4*) signaling (**Fig. 3B**). For the PD-1 signaling, the top two TNK GEMs that could be

predicted by approximated [LR] with the strongest non-linear correlation were TNK26 and TNK50,

with Pearson's correlation coefficients between prediction and observation being $\rho = 0.782$, $p =$

7.2831e-9 and $\rho = 0.685$, $p = 6.8891$e-07, respectively. The UMAPs show that the distributions of

PDCD1 and the two GEMs are highly correlated at the single-cell level (**Fig. 3B-C**). The results

also suggest potential PD-L1/PD-1 signaling is associated with distinct GEMs in different T cell

subtypes. Specifically, [*CD274*]*[*PDCD1*] is associated with TNK50 in CXCL13$^+$CD4$^+$ T cells,

whereas [*CD274*]*[*PDCD1*] is associated TNK26 mainly in exhausted CD8$^+$ T cells. Importantly,

the approximated ligand-receptor complex [LR] is more predictive of the expression of these two

GEMs when compared with only using the receptor expression [R] alone. Similarly, the results

show that the expression values of TNK35 can be predicted by approximated *CD86* and *CTLA4*

complex, and the expression of *CTLA4* is highly correlated with the expression of TNK35 in Treg

cells (**Fig. 3B-C**), suggesting the formation of this complex at these cells is associated with

expression of the GEM in a non-linear dose-response fashion. In summary, our results show that

identifying GEMs enabled the investigation of the functional impact of potential ligand-receptor

signaling. We visualized a subset of ligand-receptor pairs and TNK GEMs that exhibited

statistically significant nonlinear dose-response relationships (**Fig. 3D, Extended Data Table 2**).

The strong non-linear dose-response relationship between a ligand-receptor pair and a GEM (e.g.,

CD274-PDCD1 and TNK26) suggests the hypothesis that the expression of some members of a

GEM may be regulated by the ligand-receptor interaction. Thus, PD-1 signaling pathway may be

more active in tumors expressing a high level of TNK26, and thereby, these tumors might be more

susceptive to anti-PD-1 treatment. To test the association of the expression level of TNK26 and

response to anti-PD-1 related treatment. We collected the clinical and transcriptomic data of CRC

patients from the CAMILLA trial [26]. This study involves metastatic CRC patients with proficient

microsatellite stable (MSS) tumors. The patients received treatment with the combination of

carbozantinib (a small molecule inhibitor of tyrosine kinases including VEGFR2) and durvalumab

(an immune checkpoint inhibitor targeting PD-L1). The transcriptome and clinical responses to

the combined treatment were measured. We separated the patients into responder and non-

responder groups, according to designation by the studies, and we compared the express level of

GEMs learned from nHDP model in these two groups. The analyses indicate a significant elevation

in TNK26 expression in the responder group compared to the non-responder group (**Fig. 3E**). This

association was further tested across three established melanoma studies [27] involving patients

treated with anti-PD-1 antibodies. Consistently, TNK26 expression is significantly higher in the

responder group compared to the non-responder group (**Extended Data Fig. 6C**). The results

support the notion that the expression level of TNK26 reflects the involvement of the PD-1

signaling in the tumors, and thus can potentially be used as a novel biomarker for responsiveness

to the anti-PD-1 therapy. Regarding TNK50, a significant difference is observed exclusively in the

melanoma dataset. The TNK50 expression is significantly higher in responder group than in non-

responder group. However, in CAMILLA trial, even though TNK50 is shown to be correlated with

TNK26, TNK50 did not reach statistical significance, likely due to the small number of responders

with one apparent outlier, as depicted in **Extended Data Fig.6B**. More examples of GEMs

showing significant difference between responder group and non-responder groups in CAMILLA

trial and melanoma cohorts are listed in **Extended Data Fig. 6A & 6C**, respectively.

Fig. 3: Discovering ligand-receptors strongly associated with specific pathways regulating the expression of GEMs. A. Illustration of GEMs reflective of downstream transcriptome mediated by biochemical signaling through ligand-receptor binding. The significant association between a GEM and a ligand-receptor needs to satisfy the sigmoid fitting. **B.** The overlap of cells expressing receptor and GEMs of interest. Left: the overlap of *PDCD1*, TNK26 and TNK50. Right: the overlap of *CTLA4* and TNK35. **C.** From left to right, the plot of sigmoid fitting between GEM

and ligand-receptor pair with *p*-value and residual standard error (RSE), UMAP plot of the receptor

gene, and UMAP plot of the GEM. **D.** The summarization of top ligand-receptor pairs significantly

associated with TNK GEMs. **E.** Comparison of the expression level of TNK26 and TNK50

between the responder group and non-responder group treated with Cabozantinib plus durvalumab.

Carbozantinib is a small molecule inhibitor of tyrosine kinases including VEGFR2 and

durvalumab is an immune checkpoint inhibitor of PD-L1. The p-value was calculated using one-

sided Wilcoxon test.

CCC between cancer-associated fibroblast and SPP1-macrophage through ligand-receptor

pairs *FN1-ITGAV* and *SPP1-ITGAV*

The significance of crosstalk between macrophages and cancer-associated fibroblasts (CAFs) has

garnered attention in recent studies [28-30]. Motivated by this, we embarked on an investigation into

this crucial aspect. After searching for highly correlated GEMs that are expressed in stromal cells

and myeloid cells, our analysis led to a pair of GEMs, Mye59 and Stroma14, which exhibited the

strongest correlation (Pearson ρ=0.78) at the pseudo-bulk RNA level. A further search of myeloid

GEMs correlated to Stroma14 led to the additional discovery of Mye43 (Pearson ρ=0.58) (**Fig. 4A**

top row). The results led to a hypothesis that myeloid cells expressing Mye59 or Mye43 may

communicate with stromal cells expressing Stroma14. Based on the annotation of GEM

distribution, Mye59 and Mye43 are expressed in a separate subpopulation of M2 macrophage, and

Stroma14 is expressed in a subpopulation of FAP$^+$POSTN$^+$ fibroblast cells defined as matrix CAF

(mCAF)[31]. The search for the *LR* potentially mediating such communication led to *FN1-ITGAV*

and *SPP1-ITGAV* pairs, which predict the expression of Stroma14 (**Fig. 4A** bottom row) in a

nonlinear dose-response fashion. Examining the distribution of the ligands and receptor genes and

the above GEMs in UMAPs showed that *FN1* and Mye59 were co-expressed in a subpopulation

of macrophage cells, and *SPP1* and Mye43 were co-expressed in another distinct macrophage

subpopulation (**Fig. 4B**). While *ITGAV* (encoding a unit of integrin) was widely expressed in

stromal cells, Stroma14 was expressed in a subpopulation of mCAF, suggesting that integrin

signaling may have different downstream effects in distinct subpopulations of stromal cells (**Fig.**

4B). The above results led to the hypothesis: a subpopulation of *ITGAV*-expressing mCAF in CRC

may interact with distinct subpopulations of macrophage cells that express cells *FN1* and *SPP1*,

respectively, and such interactions induce the expression of the Stroma14 in CAFs.

We tested the above hypothesis using spatial transcriptomic data of metastatic CRC patients from

Pelka et al [22], by studying the co-localization of the above ligands, receptors, and GEMs in the

spatial domain and modeling their interactions in spatial domains. In a CRC liver metastatic tumor

section, the spatial expression of ligand (*FN1*, *SPP1*) and receptors (*ITGAV*) and GEMs (Mye59,

Mye43, and Stroma14) were highly correlated (**Fig. 4C**), indicating that cells expressing these

genes and GEMs likely interact in spatial domains. There is a notable difference in the spatial

distributions of Mye59 and Mye43. Particularly, myeloid cells expressing Mye59 concentrate at

tumor-associated fibroblast-rich tissue and the tumor leading edge, interfacing with normal liver

tissue, whereas mey43 is mainly expressed in the tumor core. Examining the distribution of *ITGAV*

and Stroma14 showed that while *ITGAV* was expressed in both tumor and adjacent normal liver

tissues (indicating the presence of stromal cells), Stroma14 is mainly expressed in mCAF in the

tumor area. This indicates that the two CCC channels (Mye59-Stroma14 and Mye43-Stroma14)

may play different signaling roles.

The spatial distribution of the approximated *FN1- ITGAV* complex agrees well with the spatial

distribution of Mye59 and Stroma14, and their distribution is distinguishable from that of *SPP1-*

ITGAV and Mye43 (**Fig. 4C**). We further modeled the relationship between approximated $[LR]$

and expression of Stroma14 in the spatial domains. The approximated *FN1-ITGAV* complex

concentration showed a clear sigmoid dose-response curve ($\rho = 0.6$, $p = 1.4429e\text{-}14$), and the

association between *SPP1-ITGAV* complex and Stroma14 is also statistically significant ($\rho =$

$0.315, p = 0.0007$). The sigmoid fittings between Stroma14 and two ligand-receptor pairs in the

spatial domain are shown in **Fig. 4D**. The above results led to a hypothetic model of the TME

involving the above ligands, receptors, and GEMs (**Fig. 4E**). We hypothesize that myeloid cells

expressing Mye59 (macrophage I) may align with the tumor leading edge as the tumor invades,

whereas myeloid cells expressing Mye43 (macrophage II) are more concentrated within the tumor

core. The interaction between these macrophage cells and mCAF cells leads to the expression of

Stroma14, which is characteristic of mCAF but absent from stromal cells in normal liver tissue.

By deconvoluting cellular signaling activities as GEMs, our approach uncovers the signaling

mechanisms through which various SPP1+ macrophage subpopulations interact with different

CAF subsets in distinct spatial domains. The example discussed above demonstrates our ability to

identify potential CCC candidates transmitted by the ligand-receptor signal, supported by both

scRNA-seq and spatial transcriptomic data.

Fig. 4: Modeling intercellular communication. A. Up panels: The scatter plot of Stroma14-vs-Mye43 and Stroma14-vs-Mye59 using the pseudo-bulk transcriptome data. Lower panels: the sigmoid fitting between Stroma14 and two ligand-receptor pairs (*FN1-ITGAV* and *SPP1-ITGAV*) using the pseudo-bulk transcriptome data. **B.** The UMAP plot of cells expressing *FN1, SPP1, ITGAV*, Mye59, Mye43, and Stroma14, respectively. **C.** The spatial distribution plot of HE, tissue clustering, GEMs, ligands, receptors, and ligand-receptor pairs. **D.** The sigmoid fitting between

Stroma14 and two ligand-receptor pairs (*FN1-ITGAV* and *SPP1-ITGAV*) using spatial

transcriptome data. **E.** Illustration of a hypothetic spatial model for the tumor microenvironment

of the spatial sample.

Discovering coordinated CCC programs

We further investigated patterns of coordinated expression of GEMs across different cell types

(TNK, B, myeloid, stromal, and epithelial) in CRCs. We represented each tumor in the GEM space

(pseudo-bulk expression: *cell-by-GEM* matrix) and performed double clustering of GEMs and

tumors. The results (**Fig. 5A**) showed that the expression of many GEMs was highly coordinated.

We hypothesized that the coordinated expression of GEMs across multiple types of cells in a TME

is likely due to two possible mechanisms: environmental factors that orchestrate the expression of

GEMs (confounding) or a CCC network that allows cells to communicate and reach homeostasis.

We refer to a set of GEMs across multiple cell categories that displays a highly coordinated

expression pattern as a CCC program, and we have identified 9 CCC programs (rows). Based on

their GEM composition, clustering tumors revealed 6 sample clusters (columns). A complete list

of CCC programs is in **Extended Data Table 4**.

We further investigate the potential molecular basis of CCC programs. CCC Program 2 consists

of 16 GEMs from different cell categories (**Fig. 5B**). The pathway analysis indicates that GEMs

in CCC Program 2 are significantly enriched of genes involved in cell cycle pathways, such as the

C-MYC signaling pathway, a hallmark pathway of cell proliferation in different cell types [32]. The

GEMs belonging to this CCC program include GEMs from TNK, B, myeloid, epithelial and

stromal cells, respectively (**Fig. 5B**), suggesting they are subpopulations of cells in a proliferation

stage from each cell category. Of interest, we noted that TNK9 is enriched in the T-cell cluster

#10, which is labeled as proliferative T cells. Not surprisingly, an epithelial GEM belonging to

this program, Epi9, is broadly active in epithelial cells, reflecting a shared transcriptomic program

involved in cell proliferation by cancer cells. The UMAPs in **Fig. 5C** indicate that various cell

categories expressing these GEMs also express the proliferation marker gene *MKI67*. We

hypothesized that the expression of GEMs of proliferation nature in different cell types may be

associated with signals that promote proliferation of cells, e.g., receptor tyrosine kinase (RTK)

pathways, Wnt/β-catenin pathway, and PI3K-AKT-mTOR pathway. Our pathway enrichment

analysis shows that the top genes associated with most of GEMs in CCC Program 2 are enriched

in the MYC pathway, which is downstream of RTKs. To test the hypothesis, we investigated the

spatial distribution of the GEMs in tumors. The results (**Fig. 5D**) show the co-localization of GEMs

in the CCC Program 2. We calculated the spatial correlations between MKI67 with respect to B21,

Stroma68, and Epi23, using their expression values in spatial spots. The correlation coefficients

are 0.708, 0.762, and 0.804, respectively, providing evidence for the strong coordinated expression

of these GEMs in spatial domains. We further checked the cell composition of spatial spots

expressing these GEMs. The inference of cell composition in tumor spots (**Extended Data Fig.**

7A) show that those spots contain a mixture of cells potentially expressing these GEMs. The

compositions of B21, Stroma68 and Epi23, represented by their top associated genes, are not

overlapping (**Extended Data Fig. 7B**). Each of the three GEMs captures cell-type-specific

proliferation signals. The indicates that the diverse cells expressing GEMs in GEM program 2 may

be stimulated by proliferation-related factors existing in tissue microenvironments.

Moreover, we performed deconvolution analysis on the GEMs in CCC Program 2 utilizing the

single-cell RNA-seq transcriptome data obtained from the peripheral blood sample prior to the

treatment of the regorafenib and nivolumab combination therapy[33]. Our analysis revealed that the

expression levels of GEMs, including TNK9, B18, and Mye21, in the proliferation program

(CCC program 2), exhibited significant differences between responders and non-responders, as

illustrated in Extended Data Fig.8. This observation aligns with the findings reported in the

original paper and highlights the potential of cellular signals related to MKI67 proliferation as

indicative markers for predicting clinical outcomes in patients treated with a combination of

multi-kinase inhibitors and immune checkpoint inhibitors.

We also searched for CCC programs in which coordinated expression of GEMs is likely due to

CCC between the GEMs. As an example, CCC Program 9 includes TNK26, TNK35, TNK50,

Mye72, Mye77, B57, and B6. To investigate whether cells expressing these GEMs communicate

with each other, we searched for LR pairs that were co-expressed with these GEMs, e.g., (*GEM X*

and L) and (*R and GEM Y*), as a candidate pair of communicating cells, using the pseudo-bulk

RNA-seq data. We found three main LR pairs through which GEMs in the program may

communicate, including *CD274-PDCD1*, *CD86-CTLA4*, and *CXCL13-CXCR5*, which are closely

associated with T-cell immune checkpoint molecules in the tumor microenvironment. We illustrate

the possible communication channels among the cells expressing those GEMs using directed edges

based on the co-expression of *GEM X-L* and *R-GEM Y* (**Fig. 5E**). The expression level of the

ligands and receptors in the cells expressing GEMs in the CCC Program 9 is shown in **Fig. 5F**.

The cells expressing TNK GMEs in CCC Program 9, such as TNK26 and TNK35, mainly express

receptor genes *PDCD1* and *CTLA4*. GEM B57 is concentrated in IgA/G cells expressing the

CXCR5, which is the receptor of CXCL13. Two myeloid GEMs, including Mye72 and Mye77,

concentrated in dendritic cells, express ligand genes *CD274*, *CD80*, and *CD86*. We then

investigated whether these ligands, receptors, and GEMs are colocalized in spatial domains as

evidence of possible CCC among the cells expressing those GEMs. Indeed, the spatial

transcriptomic data show that *GEM X-LR-GEM Y* triplets of inferred CCC channels, e.g., Mye72—

CD274/PDCD1—TNK50 (Pearson correlations of Mye72-*CD274* and *PDCD1*-TNK50 are 0.059

and 0.545 respectively) and TNK41—*CXCL13/CXCR5*—B6 (Pearson correlations of TNK41-

CXCL13 and CXCR5-B6 are 0.574 and 0.628 respectively), tend to be colocalized in spatial

domains (**Fig. 5G**). The low correlation between *CD274* and Mye72 is due to the sparse

distribution of *CD274* with diffusion in the spatial domain. These results support the hypothesis

that CCC between cells in the spatial domain underlies the observed coordinated expression of

GEMs, i.e., CCC programs.

In summary, the findings presented in this section collectively illustrate the feasibility of

uncovering a unique CCC mechanism within a micro-TME. This is achieved by modeling the

synchronized expression of GEMs, identifying potential LRs responsible for mediating

communication, and providing evidence of spatial colocalization for the components involved in

the identified CCC channel.

Fig. 5: Discovering patterns of intercellular communication channels that define transcriptome programs across different cell types. A. Clustering of GEMs and CRC tumors. Pseudo-bulk profile for each tumor was constructed from scRNA-seq data. Expression values of a GEM were estimated based on the GSVA using the top 50 weighted genes in a GEM. The GEMs (rows) are grouped into 9 CCC programs by clustering analysis, representing patterns of intercellular communication channels. The tumors (columns) are grouped into 6 clusters. **B.** The GEM members of the CCC Program 2. The majority of genes associated with these GEMs are

MYC pathway targets, as indicated by the HALLMARK_MYC_TARGETS in GSEA. **C.** UMAP

of MKI67 and GEMs in CCC program 2 in different cell-type categories. **D.** Spatial colocalization

of proliferation marker MKI67 and GEMs in CCC Program 2. **E.** The potential CCC network and

the ligand-receptors involved in GEMs in CCC Program 9. The red and blue rectangle in the center

represents ligands and receptors respectively. The edge between a ligand/receptor and a GEM is

added if the ligand/receptor is expressed in the cells expressing a GEM. **F.** The expression level of

receptors (blue) and its associated ligands (red) in cells expressing the GEMs in CCC program 9.

TNK and B GEMs in CCC Program 9 mainly express receptors (*PDCD1, CTLA4, and CXCR5*).

Myeloid GEMs mainly express ligands (*CD80, CD86 and CD274*). **G.** The spatial colocalization

of ligand, receptor and GEMs using Mye73 -> *CD274-PDCD1->* TNK50 and TNK26 -> *CXCL13-*

CXCR5 -> B6 as examples.

Construction of cell-cell communication subnetworks connected by ligand-receptor signaling

Discovering a pair of highly correlated GEMs across cell types suggests possible CCC between

the cells expressing the GEMs. Subsequently, we conducted a systematic search and examination

of potential CCC channels and networks. Using CRC pseudo-bulk RNA-seq data, we first searched

for highly correlated GEMs (*GEM X* and *GEM Y*) that are expressed in different cell types (or

clusters) to discover potential CCCs. In addition, we performed a causal analysis to find potential

ligands and receptors that mediate CCCs. To that end, we then search for *LR* pairs such that *L* is

co-expressed with *GEM X* and *R* is co-expressed with *GEM Y*. Finally, we assess whether

conditioning on LR, the two marginally correlated GEMs become independent, which is denoted

as *GEM X* ⊥ *GEM Y| LR*. Conditional independence is an important concept in causal analysis,

and in our setting, if a pair of highly correlated GEMs becomes independent when

conditioning on a LR pair, this LR pair is potentially involved in transmitting signals between

cells expressing *GEM X* and *GEM Y*. Finding a *GEM X-LR-GEM Y* triplet leads to the hypothesis

depicted in **Fig. 6A**. More specifically, a change in the state of the signaling pathway (*X*) in one

cell promotes the release of the ligand and, at the same time, leads to transcriptomic changes

represented by *GEM X*, which serves as a proxy of the pathway (*X*) state. The binding of the ligand

with its receptor in another cell results in a cellular effect represented by the signaling

pathway (*Y),* which leads to the changed gene transcription represented by *GEM Y*. Further

extending such CCC analysis to multiple GEMs and cell types leads to a CCC network (CCCN),

which may be used to understand how different cells reach homeostatic states in a TME through

CCCs. If there are GEMs/LR pairs of interest, a CCC sub-network can be constructed based on

the *GEM X-LR-GEM Y* triplets. We identified and illustrated a CCC sub-network related to

immune checkpoint signaling in **Fig. 6B**. The network consists of T cells expressing TNK GEMs

(TNK26, TNK35 and TNK50) and non-TNK cells expressing GEMs (Stroma37, B44, Mye21,

Mye43, and Mye46), likely transmitted through PD-1, CTLA4 and CXCL13 signaling. In this

network, the communication between cells expressing Mye9 (a combination of dendritic and

macrophage cells) and TNK26 (PDCD1$^+$CXCL13$^+$ CD4$^+$ and CD8$^+$ T cells) can be mediated

through *CD274-PDCD1*; the communication between the cells expressing Mye43 (TAM: tumor-

associated macrophage) and TNK35 (LAYN$^+$ Effector Treg)/TNK26 is transmitted through

CD80/86-CTLA4; the communication between Mye26-expressing NLRP3$^+$INHBA$^+$ macrophage

cells and TNK50-expressing CXCL13 producing CD4$^+$ T cells is facilitated via *CD274-PDCD1*;

the communication between TNK50 and b44-expressing follicular B helper cell is transmitted

through *CXCL13-CXCR5* which is consistent with the current literature [34]. Moreover, we also

discovered the communication between a specific subset of mCAF and effector Treg. The

communication between Stroma37 expressing MMP1$^+$CXCL5$^+$mCAF cells and TNK35

expressing LAYN1$^+$Treg cells is transmitted through *CD274-PDCD1*. *MMP1* ranks the top 3rd

gene associated with Stroma37 and has been shown to play an important role in CAF-induced

protection and lead to therapy resistance [35]. The communication between stromal cells and Tregs

has been validated in several studies [36]. Besides, TNK51 expressing Th1 cells communicate with

myeloid GEMs (Mye8 and Mye42) expressing suppressor-like macrophage (M2) through *TNF-*

TNFRSF1B[37,38]. The above results demonstrate that re-constructing CCC networks provides

insights into potential molecular mechanisms of CCCN underlying immune evasion mechanisms

of tumors.

NK cells are a promising alternative platform for cellular immunotherapy, and clinical trials of NK

cell immunotherapy have produced many encouraging clinical results [39]. In **Fig. 6C**, a CCC sub-

network illustrates the primary communications between TNK32-expressing NK cells and specific

cellular functions of stromal, myeloid cells, and B cells through *ICAM1/3-LFA-1*, *IL10-IL10RA*,

and *IL15-IL15RB* interactions. Besides, we discovered a highly ranked potential CCC channel

involving Stroma22 expressing CAF, TNK14 expressing Treg and TNK50 expressing

CXCL13$^+$CD4$^+$ T cells through *OX40L-OX40,* which is a costimulatory receptor as shown in **Fig.**

6D. Recent studies have shown the benefit of combining OX40 agonist and PD-1 inhibitor to

improve the immunotherapy outcome [40-42]. In both the CAMILLA trial dataset, where combined

therapy involved a VEGFR2 inhibitor and anti-PD-L1, and the dataset from melanoma patients

treated solely with anti-PD1 immunotherapy, Stroma22 exhibits a notable difference between the

responder and non-responder groups. (**Extended Data Fig. 6**). In addition to the aforementioned

findings, we have identified additional *GEM X-LR-GEM Y* triplets as potential CCCNs in

colorectal cancer (CRC). These discoveries offer valuable insights into exploring intercellular

communication within the TME and understanding the key cellular transcriptomic processes

associated with ligand-receptor signaling. The details are listed in **Extended Data Table 3**. To

enhance the user-friendliness of our framework, we've created an interactive website where users

can explore the annotation of each GEM, identify the top-ranked GEMs associated with each

ligand-receptor pair, and investigate the *GEM X-LR-GEM Y* triplets of interest

(http://44.192.10.166:3838/).

Fig. 6: The newly discovered cellular networks associated with ligand-receptor interactions involved in the immune checkpoint inhibitor therapy. A. The framework of the conditional independence analysis to find the causal relationship between GEM-expressing cells transmitted by ligand-receptor signals. **B.** Subnetwork of intercellular communication related to PDL1/2-PD1, CD80/CD86-CTLA4, and CXCL13-CXCR5 based on the conditional independence analysis. The

532   detailed annotation of each GEM is listed in the interactive website. **C.** Subnetwork of intercellular

533   communication related to TNK32 which represents the cellular function of NK cells. **D.**

534   Subnetwork of intercellular communication between Stroma22 and GEMs from other cell types

535   through the OX40L-OX40 signaling.

Discussion

In this study, we showed that by deconvoluting the hierarchically organized transcriptomic

programs and closely representing cell differential programs, the nHDP model can identify GEMs

that are potentially regulated through CCC, discover possible CCC channels, and generate the

hypothesis of the molecular mechanisms of CCC underlying heterogeneous tumor

microenvironments of CRC. Detecting and representing the states of cellular signaling systems in

individual cells is a fundamental task in single-cell biology. While there are different approaches

for deconvoluting single-cell transcriptomic components [16,43], our results indicate that by modeling

the hierarchical and compositional relationships among transcriptomic programs of cells, the

nHDP model captures fine-grained transcriptomic signals better than other non-hierarchical

models.

Although scRNAseq assays offer unparalleled cell-type-specific insights, their practical use is

impeded by several obstacles (e.g. high cost of library preparation and sequencing) that hinder

their broad application in clinical contexts. The evolution of bulk RNA-seq data over the past

decades has yielded a substantial volume of samples conducive to the execution of pattern and

machine learning investigations. The study of CCC, especially the TME associated with drug

treatments, could be further promoted by integrating insights leaned from scRNA-seq data.

Therefore, effective methods for bulk RNA-seq deconvolution are still urgently needed. The

GEMs learned from nHDP could be used as novel signatures for deconvoluting the specific

transcriptomic processes in both bulk RNA-seq data and spatial transcriptomic data. Compared

with two state-of-the-art statistical models, non-negative matrix factorization (NMF) and latent

Dirichlet allocation (LDA), for identifying transcriptomic programs, nHDP model has the

advantage of capturing the hierarchical structure of signaling pathways. GEMs derived from the

nHDP model reflect the cellular states across multiple cell subtypes or specific to a particular cell

subtype. Our recent study [44] showed that GEMs offered a more precise estimation of biological

signals, such as immune cell infiltration, in comparison to the state-of-the-art deconvolution

methods like Cibersort and SCNA [45,46]. This improved accuracy enables the identification of

different lymphocyte infiltration levels.

Identifying highly specialized GEMs enabled us to use the expression level of GEMs as proxies to

represent the state of signaling pathways (unobserved) that regulate their expression. This, in turn,

enabled us to search ligand-receptors that likely transmit extracellular signals to the pathways

inside single cells, generating the hypothesis of how across-cell signaling influences intracellular

signaling through CCC. Our results indicate that the *CD274-PDCD1* ligand-receptor pair may

influence the state of the cellular signaling pathways that regulate the expression of certain

members of TNK26 in exhausted T cells. The hypothesis is supported by multiple lines of evidence:

The non-linear dose-response relationship between the approximated ligand-receptor complex and

the GEM provides support from a biochemical point of view. The significant difference in TNK26

expression between responding and non-responding tumors treated with anti-PD-1 agents (both

newly developed clinical trials in CRC and melanoma datasets) indicates that tumors with high

TNK26 expression display active PD-L1/PD-1 signaling, rendering them more susceptible to the

treatment. Conversely, tumors with low TNK26 expression exhibit relatively inactive PD-L1-PD-

1 signaling, leading to their reduced responsiveness to the treatment. Based on these results, we

hypothesize that the PD-L1/PD-1 signaling likely regulates some members of the TNK26. Further

investigation of which members are regulated by this signal, e.g., using principled causal discovery

methods, may enable researchers to delineate mechanisms of PD-1 signaling. Similar studies of

the impact of other commonly co-expressed immune checkpoint proteins (receptors) in tumor-

experienced T cells may reveal how different checkpoint signaling pathways cooperate and lead

to immune suppression.

The discovery of a strong correlation of GEMs across cell types indicates that the expression of

GEMs by different cell types is coordinated, and such coordination is likely due to CCC. We

identified coordinated GEM expression patterns, referred to as CCC programs, and their spatial

colocalization provides additional support for intercellular communication networks. We further

noticed the differences in the activities of CCC programs may underlie the heterogeneity of TMEs,

leading to tumor clusters that recapitulate well-appreciated molecular subtyping of CRC. Further

investigation identifies ligand-receptors that likely transmit signals across GEMs in CCC channels.

We anticipate further investigating CCC channels active in each tumor/subcluster may provide

mechanistic insights into the TME heterogeneity. In our population-wide exploration of CCC, we

observed instances where multiple ligands could communicate with a single receptor, as illustrated

by the TNK32 in **Fig. 6C**. In the causal theory, those ligands are multiple parents of the receptor.

If we examine whether instantiated values of variables in a local network lead to context-specific

independence, only certain parents may contribute to the state of children from the aspect of an

individual tumor sample. For instance, in an individual tumor sample, only one type of cell among

stromal, myeloid, or B cells may be involved in the interaction with TNK32-expressing cells. Our

forthcoming efforts will involve employing the individualized Causal Bayesian Network (iCBN)

model [47] to reveal the individualized CCCN within each tumor sample. This effort will contribute

to a deeper understanding of the mechanisms underlying heterogeneity.

We envision that the framework presented in this study for deconvoluting single-cell

transcriptomic programs to model CCC can be applied to various cancer types. Enhancing the

nHDP model involves refining the initialization process and filtering variable genes, leading to a

more effective capture of relationships within transcriptomic programs. This improvement will

boost the model's ability to detect GEMs regulated by specific transcriptomic programs. Future

methodological developments for detecting fine-grained transcriptomic programs may involve

leveraging recently developed language models to capture the compositionally co-expressed genes.

Additionally, integrating scRNAseq with single-cell chromatin accessibility sequencing (scATAC)

could aid in discovering active motifs associated with CCC networks. In conclusion, this study

advances the understanding of CCCs mechanisms underlying heterogeneous TME and immune

evasion mechanisms in CRC. These insights may uncover novel target for immunotherapy and

guide precision immunotherapy [21].

Online Method

Data Collection and Preprocessing

Single-cell transcriptomics

We collected single-cell RNA sequencing data from 8 publicly available colorectal cancer data

sources (**Extended Data Table 1**) [22,48-50]. The total number of patients is 153, of which there are

179 and 100 scRNA tumor and normal samples respectively. We retained cells that met the

following criteria: nUMI > 500, nGene > 200, log10GenesPerUMI > 0.8, and mitoRatio < 0.2. To

perform gene-level filtering, we retained only genes expressed in at least 0.1% of the cells. After

the quality control (QC) analysis, we applied the total-count correction (library-size correction)

and normalized the data matrix to 10,000 reads per cell. The annotations of cell major type were

obtained from original publications and used for further analysis. We then divided cells into five

major categories: 246,560 T and natural killer (TNK) cells, 94,663 myeloid cells, 96,637 plasma

and B cells, 69,128 stromal cells, and 119,870 epithelial cells. When performing clustering

analysis within each cell category, the Harmony algorithm [51] was applied to remove the batch

effect, cell clustering was carried out using the Leiden algorithm [52], and the UMAP (Uniform

Manifold Approximation and Projection) algorithm was used to visualize clusters.

Seurat pipeline was used to find subclusters and its top variable genes within TNK, myeloid,

plasmaB, stromal, and epithelial cells, respectively [13,53]. The cell types were annotated with well-

known marker genes: *CD3D, CD8A, FOXP3, IL7R, NKG7, GNLY* for T and NK cells; *CD79A,*

MS4A1, MZB1 for B and plasma cells; *CD14, CD68, CD1C, S100A8, C1QA, LYZ* for myeloid

cells. The detailed markers we used were listed in **Extended Data Table 5**.

Bulk RNA-seq

The TCGA bulk RNA-seq data of colorectal cancer (COADREAD) was downloaded from UCSC

Xena [54]. The total number of tumor samples is 290. Bulk RNA-seq data of 20 patients with

metastatic CRC, treated with the combination of Cabozantinib plus durvalumab in the CAMILLA

clinical trial and characterized by proficient microsatellite stable (MSS) tumors, were collected

upon request [26]. The number of responders and non-responders is 4 and 16, respectively. The bulk

RNA-seq data with PD1 response of melanoma was downloaded from GSE96061, PRJEB23709

and dbGap:phs000452.v3.p1 for validation analysis [55-57]. To facilitate the clustering of tumors and

ligand receptor search using the CRC single-cell data atlas, we generated pseudo-bulk RNAseq

results for tumor samples (a *tumor-by-GEM* matrix). For the tumor samples (n=151) that include

both CD45$^-$ and CD45$^+$ cells, UMI counts of the same gene in each sample were aggregated,

resulting in the expression data in raw count format. We then normalized the raw count expression

data to $\log_2$(TPM+1).

Spatial transcriptomics

Spatial transcriptomic data of colorectal cancer and its corresponding liver metastasis using 10x

Visium technology were collected from the study by Wu et al [49]. We first used sctransform

method[58] to normalize the raw count in each spot of the slide (a *spot-by-gene* matrix). The GSVA

method was then used to estimate the GEM values from normalized expression data. The

deconvoluted spatial transcriptomic data is represented as a *spot-by-GEM* matrix. STdeconvolve

was used to infer the cell-type composition in tumor region [59].

Identification of GEMs using nHDP

We used the nested hierarchical Dirichlet processes (nHDP) model [44,60] to identify patterns of

GEMs from scRNA data. The flow of the training process is shown in **Fig.1B-C**. We binarized the

UMI count of each gene within a cell as the input of the nHDP model. A trained nHDP model

returns two matrices: a cell-by-GEM count matrix, which collects the counts of UMIs assigned to

a GEMs in each cell; a GEM-by-gene matrix, which represents the probability profile of genes in

each GEM (**Fig.1B**). We constructed a three-layer tree structure with the following branching

factors: 5, 4, and 3 for 1st, 2nd, and 3rd layers, respectively, as shown in **Fig.1C**. The expression of

GEMs (cell-by-GEM matrix) is added to the Seurat object as new features for plotting the UMAP

of GEMs. The dominant function represented by a GEM is based on the rank of cell subtype

percentage ("Distribution" in **Extended Data Fig. 4**). We used word clouds R package

"wordcloud2" to visualize the probability profile of the top 20 associated genes of a GEM (**Fig.1D**).

We trained a nHDP model for each cell category (TNK, B, Mye, Stroma, and Epi) and retained

GEMs expressed in over 1% of cells in a category for further analysis.

Functional analysis of GEMs

680   Transcription factor and enrichment analysis are applied to conduct functional analysis on GEMs.

681   CHEA3 is used to conduct the transcription factor analysis [23]. Reactome, KEGG and GO databases

682   are used to conduct more comprehensive analysis using R package "msigdbr"[61].

Estimation of GEM level in bulk and spatial RNA-seq data

685   To estimate the expression level of a GEM in bulk/pseudo-bulk/spatial RNA-seq data, we used

686   two methods: 1) gene set variation analysis (GSVA) of the top 50 weighted genes in each GEM [62].

687   2) mean expression of the top 50 weighted genes in each GEM.

Fitting non-linear relationship between ligand-receptor pairs and GEMs

The ligand-receptor database was downloaded from Nichenet [20]. To fit the dose-response curve

between the expression of ligand-receptor and GEM value, we model the quantity of LR complex

[LR] as the product of expression of L and R based on the fact that $[LR] \propto [L][R]$ using the

pseudo-bulk RNA-seq data. We first scaled the GEM value to [0, 1] and then created a 2-parameter

log-logistic function where the lower limit fixes at 0 and upper limit fixes at 1 using *drc/v3.0.1* [63].

The coefficients and its associated *p*-values were estimated and adjusted[20]. Sigmoid fitting with a

significant adjusted *p*-value (<0.05) demonstrates a good fit between the expression of LR complex

and the scaled GEM values.

699   Assessing conditional independence of *GEM X-LR-GEM Y* triplets

To find out the ligand-receptor pairs that potentially mediate CCC between cells across cell

categories expressing two correlated GEMs, conditional independence tests of $GEM\ X \perp\!\!\!\perp GEM\ Y |$

LR were performed using the *bnlearn/v4.7.1* package in R. We estimated expression status of each

GEM using the GSVA scores (*GSVA/v1.42.0*) based on its top 50 genes. We calculated all pairwise

correlations between GEMs expressed in different cell categories. If a pair of cross-cell-category

GEMs exhibits significant correlation, we further search for LR pairs that rendered conditional

independence ($GEM\ X \perp\!\!\!\perp GEM\ Y | LR$). *P*-values of conditional independence tests were adjusted

to control the false discovery rate using Benjamini-Hochberg method. We set the threshold of the

conditional correlation between *GEM1* and *GEM2* at an adjusted *p*-value > 0.05 as conditional

independent.

We further performed post-hoc proportional tests to remove false positive findings. For each

conditional independent *GEM X—LR—GEM Y* triplet, we investigated the co-expression of *GEM*

X—L in potential signal-originating cells and the co-expression of *R—GEM Y* in potential signal-

receiving cells. To conduct the proportional tests, we first construct a 2 by 2 contingency table to

measure the frequency of co-expression GEM and the tested gene. GEM and gene were considered

presenting if the corresponding values are beyond 0 and not presenting otherwise. Then the

statistics of proportional tests $z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1 - p_0)}{n}}}$ were estimated to test if the proportion of co-

expressing cells is significantly higher than the proportion of not co-expressing cells, where $\hat{p}$ is

the proportion of cells that co-express GEM and tested gene and p_0 is the proportion of cells that

not co-express GEM and tested gene, n is the total number of cells that express such GEM. p-value

of each post-hoc test was also adjusted using the Benjamini-Hochberg method for false discovery

rate control. Ligand-receptor pair that passes both conditional independence test and post-hoc

proportional test was considered valid for downstream analysis.

To prevent the false-positive rate, we set multiple criteria to filter the GEM pairs, including 1) the

correlation between *GEM X* and *L* is over 0.3 based on pseudo-bulk RNA-seq data 2) the

correlation between *LR* and *GEM Y* is over 0.3 based on pseudo-bulk RNA-seq data 3) more than

10% of cells expressing *GEM X* also express ligand gene *L* and more than 10% of cells expressing

GEM Y also express receptor gene *R*. We incorporated the examination of expression at the single-

cell level to confirm the presence of a specific molecule (ligand gene/receptor gene) in the GEM

expressed cells, thereby reducing false positives.

Identifying programs of GEMs

Hierarchical clustering was performed on a Euclidean distance matrix to identify GEM programs

using the pseudo-bulk expression of GEMs.

Evaluating GEMs as potential biomarkers for anti-PD1 therapies

We performed GSVA analysis to determine the expression levels of GEMs for each bulk tumor

sample. Samples were divided to response group if they showed complete response or partial

response to anti-PD-1 whereas non-response group if progressive disease or stable disease were

seen. We employed one-sided Wilcoxon rank test [64] to identify GEMs that exhibited a significant

difference in expression levels between these two groups.

Availability

745 nHDP model designed for analyzing single-cell transcriptome (scGEM):

746 https://github.com/hansolo-bioinfo/scGEM

747 CRC Atlas: https://github.com/hansolo-bioinfo/CRCAtlas

748 Interactive website: http://44.192.10.166:3838/

Acknowledgments

751 This research was supported in part by The National Library of Medicine; National Cancer

752 Institute at the National Institutes of Health grants R00LM013089, R01HG014023,

753 R01LM012011 and R01CA254274. The Department specifically disclaims responsibility for any

754 analysis, interpretations or conclusions.

Reference

1   Xi, Y. & Xu, P. Global colorectal cancer burden in 2020 and projections to 2040. *Transl*
Oncol **14**, 101174, doi:10.1016/j.tranon.2021.101174 (2021).
2   Sahin, I. H. *et al.* Immune checkpoint inhibitors for the treatment of MSI-H/MMR-D
colorectal cancer and a perspective on resistance mechanisms. *Br J Cancer* **121**, 809-818,
doi:10.1038/s41416-019-0599-y (2019).

3   Armingol, E., Officer, A., Harismendy, O. & Lewis, N. E. Deciphering cell-cell
interactions and communication from gene expression. *Nat Rev Genet* **22**, 71-88,
doi:10.1038/s41576-020-00292-x (2021).
4   Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat*
Commun **12**, 1088, doi:10.1038/s41467-021-21246-9 (2021).
5   Giladi, A. *et al.* Dissecting cellular crosstalk by sequencing physically interacting cells.
Nat Biotechnol **38**, 629-637, doi:10.1038/s41587-020-0442-2 (2020).
6   Shao, X., Lu, X., Liao, J., Chen, H. & Fan, X. New avenues for systematically inferring
cell-cell communication: through single-cell transcriptomics data. *Protein Cell* **11**, 866-
880, doi:10.1007/s13238-020-00727-5 (2020).
7   Marx, V. Method of the Year: spatially resolved transcriptomics. *Nat Methods* **18**, 9-14,
doi:10.1038/s41592-020-01033-y (2021).
8   Moses, L. & Pachter, L. Museum of spatial transcriptomics. *Nat Methods* **19**, 534-546,
doi:10.1038/s41592-022-01409-2 (2022).
9   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. *Nat Biotechnol*
36, 411-420, doi:10.1038/nbt.4096 (2018).
10   Luecken, M. D. & Theis, F. J. Current best practices in single-cell RNA-seq analysis: a
tutorial. *Mol Syst Biol* **15**, e8746, doi:10.15252/msb.20188746 (2019).
11   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression
data analysis. *Genome biology* **19**, 15, doi:10.1186/s13059-017-1382-0 (2018).
12   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-1902
e1821, doi:10.1016/j.cell.2019.05.031 (2019).
13   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587
e3529, doi:10.1016/j.cell.2021.04.048 (2021).
14   Kotliar, D. *et al.* Identifying gene expression programs of cell-type identity and cellular
activity with single-cell RNA-Seq. *Elife* **8**, doi:10.7554/eLife.43803 (2019).
15   Zhu, X., Ching, T., Pan, X., Weissman, S. M. & Garmire, L. Detecting heterogeneity in
single-cell RNA-Seq data by non-negative matrix factorization. *PeerJ* **5**, e2888,
doi:10.7717/peerj.2888 (2017).
16   Liu, L. *et al.* Deconvolution of single-cell multi-omics layers reveals regulatory
heterogeneity. *Nat Commun* **10**, 470, doi:10.1038/s41467-018-08205-7 (2019).
17   Venkatasubramanian, M., Chetal, K., Schnell, D. J., Atluri, G. & Salomonis, N.
Resolving single-cell heterogeneity from hundreds of thousands of cells through
sequential hybrid clustering and NMF. *Bioinformatics* **36**, 3773-3780,
doi:10.1093/bioinformatics/btaa201 (2020).
18   Shao, C. & Hofer, T. Robust classification of single-cell transcriptome data by
nonnegative matrix factorization. *Bioinformatics* **33**, 235-242,
doi:10.1093/bioinformatics/btw607 (2017).
19   Efremova, M., Vento-Tormo, M., Teichmann, S. A. & Vento-Tormo, R. CellPhoneDB:
inferring cell-cell communication from combined expression of multi-subunit ligand-
receptor complexes. *Nat Protoc* **15**, 1484-1506, doi:10.1038/s41596-020-0292-x (2020).
20   Browaeys, R., Saelens, W. & Saeys, Y. NicheNet: modeling intercellular communication
by linking ligands to target genes. *Nat Methods* **17**, 159-162, doi:10.1038/s41592-019-
0667-5 (2020).

21 Chen, X. *et al.* An individualized causal framework for learning intercellular
communication networks that define microenvironments of individual tumors. *PLoS*
computational biology **18**, e1010761, doi:10.1371/journal.pcbi.1010761 (2022).
22 Pelka, K. *et al.* Spatially organized multicellular immune hubs in human colorectal
cancer. *Cell* **184**, 4734-4752 e4720, doi:10.1016/j.cell.2021.08.003 (2021).
23 Keenan, A. B. *et al.* ChEA3: transcription factor enrichment analysis by orthogonal
omics integration. *Nucleic Acids Res* **47**, W212-W224, doi:10.1093/nar/gkz446 (2019).
24 Martinez, G. J. *et al.* The transcription factor NFAT promotes exhaustion of activated
CD8(+) T cells. *Immunity* **42**, 265-278, doi:10.1016/j.immuni.2015.01.006 (2015).
25 Fan, J. *et al.* Zinc finger protein 831 promotes apoptosis and enhances chemosensitivity
in breast cancer by acting as a novel transcriptional repressor targeting the STAT3/Bcl2
signaling pathway. *Genes Dis* **11**, 430-448, doi:10.1016/j.gendis.2022.11.023 (2024).
26 Saeed, A. *et al.* Cabozantinib plus durvalumab in advanced gastroesophageal cancer and
other gastrointestinal malignancies: Phase Ib CAMILLA trial results. *Cell Rep Med* **4**,
100916, doi:10.1016/j.xcrm.2023.100916 (2023).
27 . (!!! INVALID CITATION !!! 23-25).
28 Hoeft, K. *et al.* Platelet-instructed SPP1(+) macrophages drive myofibroblast activation
in fibrosis in a CXCL4-dependent manner. *Cell Rep* **42**, 112131,
doi:10.1016/j.celrep.2023.112131 (2023).
29 Qi, J. *et al.* Single-cell and spatial analysis reveal interaction of FAP(+) fibroblasts and
SPP1(+) macrophages in colorectal cancer. *Nat Commun* **13**, 1742, doi:10.1038/s41467-
022-29366-6 (2022).
30 Mao, X. *et al.* Crosstalk between cancer-associated fibroblasts and immune cells in the
tumor microenvironment: new findings and future perspectives. *Mol Cancer* **20**, 131,
doi:10.1186/s12943-021-01428-1 (2021).
31 Chen, Z. *et al.* Single-cell RNA sequencing highlights the role of inflammatory cancer-
associated fibroblasts in bladder urothelial carcinoma. *Nat Commun* **11**, 5077,
doi:10.1038/s41467-020-18916-5 (2020).
32 . (!!! INVALID CITATION !!! 30).
33 Kim, H. D. *et al.* Regorafenib plus nivolumab in unresectable hepatocellular carcinoma:
the phase 2 RENOBATE trial. *Nat Med* **30**, 699-707, doi:10.1038/s41591-024-02824-y
(2024).
34 Rao, D. A. T Cells That Help B Cells in Chronically Inflamed Tissues. *Front Immunol* **9**,
1924, doi:10.3389/fimmu.2018.01924 (2018).
35 Cui, Q. *et al.* Upregulating MMP-1 in carcinoma-associated fibroblasts reduces the
efficacy of Taxotere on breast cancer synergized by Collagen IV. *Oncol Lett* **16**, 3537-
3544, doi:10.3892/ol.2018.9092 (2018).
36 Sun, R. *et al.* Amphiregulin couples IL1RL1(+) regulatory T cells and cancer-associated
fibroblasts to impede antitumor immunity. *Sci Adv* **9**, eadd7399,
doi:10.1126/sciadv.add7399 (2023).
37 Wilk, A. J., Shalek, A. K., Holmes, S. & Blish, C. A. Comparative analysis of cell-cell
communication at single-cell resolution. *Nat Biotechnol*, doi:10.1038/s41587-023-01782-
z (2023).
38 Nonaka, K. *et al.* Th1 polarization in the tumor microenvironment upregulates the
myeloid-derived suppressor-like function of macrophages. *Cell Immunol* **369**, 104437,
doi:10.1016/j.cellimm.2021.104437 (2021).

39   Chu, J. *et al.* Natural killer cells: a promising immunotherapy for cancer. *J Transl Med*
20, 240, doi:10.1186/s12967-022-03437-0 (2022).
40   van der Sluis, T. C. *et al.* OX40 agonism enhances PD-L1 checkpoint blockade by
shifting the cytotoxic T cell differentiation spectrum. *Cell Rep Med* **4**, 100939,
doi:10.1016/j.xcrm.2023.100939 (2023).
41   Ma, Y. *et al.* Combination of PD-1 Inhibitor and OX40 Agonist Induces Tumor Rejection
and Immune Memory in Mouse Models of Pancreatic Cancer. *Gastroenterology* **159**,
306-319 e312, doi:10.1053/j.gastro.2020.03.018 (2020).
42   Lao, J. *et al.* OX40 enhances T cell immune response to PD-1 blockade therapy in non-
small cell lung cancer. *Int Immunopharmacol* **108**, 108813,
doi:10.1016/j.intimp.2022.108813 (2022).
43   Im, Y. & Kim, Y. A Comprehensive Overview of RNA Deconvolution Methods and
Their Application. *Mol Cells* **46**, 99-105, doi:10.14348/molcells.2023.2178 (2023).
44   Zhang, H., Lu, X., Lu, B. & Chen, L. scGEM: Unveiling the Nested Tree-Structured
Gene Co-Expressing Modules in Single Cell Transcriptome Data. *Cancers (Basel)* **15**,
doi:10.3390/cancers15174277 (2023).
45   Davoli, T., Uno, H., Wooten, E. C. & Elledge, S. J. Tumor aneuploidy correlates with
markers of immune evasion and with reduced response to immunotherapy. *Science* **355**,
doi:10.1126/science.aaf8399 (2017).
46   Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression profiles.
Nat Methods **12**, 453-457, doi:10.1038/nmeth.3337 (2015).
47   Liu, Z. *et al.* A Novel Bayesian Framework Infers Driver Activation States and Reveals
Pathway-Oriented Molecular Subtypes in Head and Neck Cancer. *Cancers (Basel)* **14**,
doi:10.3390/cancers14194825 (2022).
48   Joanito, I. *et al.* Single-cell and bulk transcriptome sequencing identifies two epithelial
tumor cell states and refines the consensus molecular classification of colorectal cancer.
Nat Genet **54**, 963-975, doi:10.1038/s41588-022-01100-4 (2022).
49   Wu, Y. *et al.* Spatiotemporal Immune Landscape of Colorectal Cancer Liver Metastasis
at Single-Cell Level. *Cancer Discov* **12**, 134-153, doi:10.1158/2159-8290.CD-21-0316
(2022).
50   Liu, Y. *et al.* Immune phenotypic linkage between colorectal cancer and liver metastasis.
Cancer Cell **40**, 424-437 e425, doi:10.1016/j.ccell.2022.02.013 (2022).
51   Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with
Harmony. *Nat Methods* **16**, 1289-1296, doi:10.1038/s41592-019-0619-0 (2019).
52   Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-
connected communities. *Sci Rep* **9**, 5233, doi:10.1038/s41598-019-41695-z (2019).
53   Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of
single-cell gene expression data. *Nat Biotechnol* **33**, 495-502, doi:10.1038/nbt.3192
(2015).
54   Goldman, M. J. *et al.* Visualizing and interpreting cancer genomics data via the Xena
platform. *Nat Biotechnol* **38**, 675-678, doi:10.1038/s41587-020-0546-8 (2020).
55   Riaz, N. *et al.* Tumor and Microenvironment Evolution during Immunotherapy with
Nivolumab. *Cell* **171**, 934-949 e916, doi:10.1016/j.cell.2017.09.028 (2017).
56   Gide, T. N. *et al.* Distinct Immune Cell Populations Define Response to Anti-PD-1
Monotherapy and Anti-PD-1/Anti-CTLA-4 Combined Therapy. *Cancer Cell* **35**, 238-255
e236, doi:10.1016/j.ccell.2019.01.003 (2019).

57   Liu, D. *et al.* Integrative molecular and clinical modeling of clinical outcomes to PD1
900   blockade in patients with metastatic melanoma. *Nat Med* **25**, 1916-1927,
901   doi:10.1038/s41591-019-0654-5 (2019).
902   58   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-
903   seq data using regularized negative binomial regression. *Genome Biol* **20**, 296,
904   doi:10.1186/s13059-019-1874-1 (2019).
905   59   Miller, B. F., Huang, F., Atta, L., Sahoo, A. & Fan, J. Reference-free cell type
906   deconvolution of multi-cellular pixel-resolution spatially resolved transcriptomics data.
907   *Nature communications* **13**, 2339 (2022).
908   60   Paisley, J., Wang, C., Blei, D. M. & Jordan, M. I. Nested Hierarchical Dirichlet
909   Processes. *IEEE Trans Pattern Anal Mach Intell* **37**, 256-270,
910   doi:10.1109/TPAMI.2014.2318728 (2015).
911   61   Liberzon, A. *et al.* Molecular signatures database (MSigDB) 3.0. *Bioinformatics* **27**,
912   1739-1740, doi:10.1093/bioinformatics/btr260 (2011).
913   62   Hanzelmann, S., Castelo, R. & Guinney, J. GSVA: gene set variation analysis for
914   microarray and RNA-seq data. *BMC Bioinformatics* **14**, 7, doi:10.1186/1471-2105-14-7
915   (2013).
916   63   Ritz, C., Baty, F., Streibig, J. C. & Gerhard, D. Dose-Response Analysis Using R. *PLoS*
917   *One* **10**, e0146021, doi:10.1371/journal.pone.0146021 (2015).
918   64   Forrester, J. C. & Ury, H. K. The Signed-Rank (Wilcoxon) test in the rapid analysis of
919   biological data. *Lancet* **1**, 239-241, doi:10.1016/s0140-6736(69)91245-8 (1969).
920

Supplementary Figures

Supp Fig. 1: UMAP of B, Myeloid and Stromal cells. A. The cell clusters for B and Plasma cells.

B. The cell clusters for myeloid cells. **C.** The cell clusters for stromal cells.

Supp Fig. 2: Comparison between GEMs learned from nHDP and GEMs learned from NMF.

A. The cells expressing NMF5 are the combination of cells expressing TNK15, TNK42 and

TNK51. **B.** The cells expressing NMF58 are the combination of cells expressing TNK17, TNK29,

and TNK50.

Supp Fig. 3: Cosine similarity between top50 GEM associated genes and highly expressed marker genes in T cell subtypes. A. Heatmap of cosine similarities between differentially expressed genes of T cell subtype and top 50 genes of all 85 T cell GEMs learned from nHDP model. GEMs are labeled with respect to their tree levels (L1, L2, and L3) and major T cell subtypes including CD4[+], CD8[+], ILC, MAIT, NK, Treg and proliferation T cells. **B.** The heatmap of cosine similarity for significant T cell GEMs summarized in Fig. 2E are shown.

Supp Fig. 4: The dot plot comparing the similarity between GEMs and traditional cell clusters. A. The dot plot for B and Plasma cells. **B.** The dot plot for myeloid cells. **C.** The dot plot for stromal cells. **D.** The dot plot for TNK cells. The color and size of the dot reflect the average expression and percentage of overlapped cells, respectively.

Supp Fig. 5: The enriched signaling pathways. A. The enriched signaling pathway for myeloid

cells. **B.** The enriched signaling pathway for B and plasma cells. **C.** The enriched signaling

pathway for stromal cells.

Supp Fig. 6: Example of GEMs significantly differentially expressed in responder group and

non-responder group. A. Example of GEMs significantly differentially expressed in responder

group and non-responder group treated with Cabozantinib plus durvalumab (CAMILLA trial). **B.**

The correlation between TNK26 and TNK50 in CAMILLA cohort. **C.** Example of GEMs

significantly differentially expressed in responder group and non-responder group treated with

solo anti-PD-1 immunotherapy (three independent melanoma datasets).

A Inference of cell-type composition in tumor region

🔴🟢 Fibroblast 🟡🔵 Lymphocyte 🟠🔵🟣 Epithelial cell

B

	B21	Stroma68	Epi23
B21	1	0	0
Stroma68	0	1	0.02
Epi23	0	0.02	1

981 **Supp Fig. 7: Exploring the uniqueness of three GEMs in CCC program 2. A.** The inference

 of cell-type composition in tumor region of **Fig. 5D**. **B.** The cosine similarity between B21,

 Stroma68 and Epi23 using the expression level of the top 50 associated genes.

 Suppl Fig 8: Example of GEMs in the proliferation program (CCC program2), which are

 significantly differentially expressed in responders and non-responders treated with

 regorafenib and nivolumab combination therapy.