Earlier snowmelt reduces the strength of carbon sink in montane meadows

Running title: Weaker carbon sink with earlier snowmelt

OK Vought[1,2,3*], SN Kivlin[4], H Shulman[4], PO Sorensen[5,6], DW Inouye[3,7], I Ibáñez[8], P Falb[1], K
Rand[1,9], AT Classen[1,9]

[1]Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI 48109, USA
[2]Institute for Global Change Biology, School for Environment and Sustainability, University of Michigan,
Michigan, Ann Arbor, USA
[3]The Rocky Mountain Biological Laboratory, PO Box 519, Crested Butte, Colorado 81224, USA
[4]Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, Tennessee 37996, USA
[5]Natural Resources Science, University of Rhode Island, Kingston, Rhode Island 02881, USA
[6]Climate and Ecosystem Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California, 94720,
USA
[7]Department of Biology, University of Maryland, College Park, Maryland 20742, USA
[8]School for Environment and Sustainability, University of Michigan, Ann Arbor, MI 48109, USA
[9]University of Michigan Biological Station, Pellston, Michigan 49769, USA

* Corresponding author: Olivia Vought (ovought@umich.edu, +1 315-414-6294)

Abstract

Warming temperatures are changing winters, leading to earlier snowmelt. This shift can

accelerate or extend the growing season, which in turn may affect various plant-mediated

ecosystem functions. Despite its relevance in the carbon cycle, we still know little about how

earlier snowmelt impacts the carbon balance in ecosystems over the growing season. Most

studies rely on interannual variability in snowmelt timing, making it difficult to isolate snowmelt

effects from other confounding variables, e.g, temperature and moisture anomalies. To address

this uncertainty, we investigated how advancing snowmelt affects the carbon cycling of montane

meadows. We experimentally advanced the snowmelt date by approximately nine days and

collected data every two weeks throughout the growing season, including net ecosystem

exchange (NEE), gross primary productivity (GPP), ecosystem respiration (ER), plant

composition, and shrub, graminoid, and forb biomass. Early in the season, GPP was higher in the

early snowmelt plots, though this effect decreased as the season progressed. Our modeling of

cumulative NEE showed that earlier snowmelt reduced the carbon sink strength by around 70%,

likely due to drier soils. Graminoid biomass was 47% higher in plots with earlier snowmelt, but

there was no change in total biomass. As winters warm and snowmelt occurs earlier, plant

productivity will shift earlier in the growing season. However, reduced soil moisture during the

summer will limit productivity, weakening the overall carbon sink and altering the carbon

balance in montane meadows.

Keywords: net ecosystem carbon exchange, ecosystem respiration, gross primary productivity,

carbon cycle, montane meadow, plant community, climate change, snowmelt

Introduction

In mountain regions, the climate is getting warmer and drier, causing earlier snowmelt (Gottlieb & Mankin, 2024; Lambert et al., 2010; Pepin et al., 2015). Snowmelt timing affects plant phenology and growth in montane meadows (Semenchuk et al., 2016; Vorkauf et al., 2021), as it provides a critical moisture pulse for plant growth in water-limited montane meadows (Ganjurjav et al., 2016; Kawai & Kudo, 2021; Kudo, 2021). Earlier snowmelt can either increase (Zhang et al., 2023) or decrease (Gamon et al., 2013) plant productivity and ecosystem respiration (Li et al., 2016), with a lagged effect on carbon fluxes evident for months (Liu et al., 2023). Earlier snowmelt can also interact with warming temperatures to further dry soils and exacerbate climate change effects on ecosystems (Suzuki, 2014). Changes in plant community composition due to earlier snowmelt can either amplify or mitigate these snowmelt effects on carbon cycling by altering the relative abundance of highly productive species or modifying root production, which influences carbon inputs to the soil through changes in root turnover (D'Imperio et al., 2018; Kelsey et al., 2021; Li et al., 2020; Ma et al., 2024). Despite these documented effects, the impact of earlier snowmelt on the carbon balance of montane meadows remains unclear, as most studies rely on natural variations in snowmelt dates rather than experimental manipulations with control plots, which can lead to confounding results, such as temperature and moisture anomalies. Plant communities are also likely adapted to variable snow conditions in systems with large natural snowmelt variation, so they could be less responsive to changing snowmelt (Metz & Tielbörger, 2016). Additionally, the same system is experiencing earlier and later snowmelt year to year with variable snowmelt, but climate change is advancing snowmelt, which an experiment better captures.

Plants play a crucial role in shaping the response of carbon cycling to changes in snowmelt, by

altering their phenology, shifting their species composition (Li et al., 2020; Ma et al., 2024), and

changing how they allocate resources above and belowground (Darrouzet-Nardi et al., 2019;

Sherwood et al., 2017; Vorkauf et al., 2021). With earlier snowmelt, plants often shift their

growth earlier in the season, leading to increased productivity in the spring (Blume-Werry et al.,

2017; Darrouzet-Nardi et al., 2019; Sherwood et al., 2017; Vorkauf et al., 2021). However, by

peak growing season, earlier snowmelt may not significantly change overall biomass (Möhl et

al., 2023). Grasses, in particular, tend to respond to changes in snowmelt timing, often becoming

more abundant following the pulse of moisture from snowmelt (Ma et al., 2024). However,

whether grasses increase (Li et al., 2020; Ma et al., 2024) or decrease (Liu et al., 2018; Rosbakh

et al., 2017) their biomass with less snow and drier conditions depends on the relative rooting

depth of co-occurring species and their ability to uptake water. Drier conditions typically favor

deeper-rooted species, so species with deeper roots could increase with drought (Liu et al.,

2018). However, grasses have lower carbon uptake than other plant functional groups (Ibanez et

al., 2020), so shifts toward grasses create a positive feedback loop, exacerbating climate change

impacts. Interestingly, changes to snowmelt timing do not typically affect root phenology

(Blume-Werry et al., 2017; Darrouzet-Nardi et al., 2019), though experimentally increased snow

depth in some ecosystems can increase (Li et al., 2020) or decrease (D'Imperio et al., 2018) root

production. These changes in root production can influence carbon uptake, as increased root

growth can boost carbon absorption, but greater standing root biomass could also release carbon

via respiration.

Changes in snowmelt timing can affect community and ecosystem-level carbon flux. Earlier

snowmelt can increase plant carbon sequestration by enhancing plant productivity earlier in the

growing season (Chen et al., 2019; Zhang et al., 2023). However, it may also shorten the

growing season if moisture limitations arise later in the growing season (Zona et al., 2022).

While earlier snowmelt could boost gross primary productivity (GPP), it may also increase

ecosystem respiration (ER), neutralizing any impact on NEE (López-Blanco et al., 2017).

Experimental snow depth studies find mixed effects on carbon uptake, with some studies finding

increased uptake (Li et al., 2020) and others finding no effect (Hu et al., 2013). Interestingly,

delaying snowmelt had a more significant impact on plant productivity than increasing snow

depth, likely because early-season rain diminished the moisture effects of deeper snow (Ma et

al., 2024).

To better understand the effects of earlier snowmelt on carbon cycling, we experimentally

advanced the snowmelt date in a montane meadow by nine days. Specifically, we asked: How

does an earlier snowmelt date impact the carbon dynamics of a montane meadow across the

growing season, considering plant biomass and the overall carbon balance? We hypothesized:

1. Total annual biomass aboveground will not change with snowmelt, as drier conditions

later in the growing season would offset any early season increases. Root production will

increase as plants allocate more resources belowground with drier conditions due to

earlier snowmelt.

2. Earlier snowmelt will shift aboveground spring phenology earlier. Root phenology will

be unchanged, as root growth may begin before snowmelt.

3. Grasses, which respond to moisture pulses from snowmelt due to their fibrous roots, will

be relatively abundant early in the season, especially with earlier snowmelt.

4. NEE will initially decrease (carbon uptake increases) with early snowmelt as early-

season carbon uptake rises. However, the overall carbon budget will remain unchanged,

as early-season productivity gains will be offset by declines in late-season productivity

due to soil drying.

Our results highlight how shifts in snowmelt timing affect carbon dynamics, as we measured the

impacts on plant phenology, biomass accumulation, and carbon fluxes across the growing season

in a controlled experiment.

Materials & Methods

Study sites & experimental establishment

In July 2022, we established a snowmelt manipulation experiment in the East River Meadow

near RMBL (38.962396, -106.993848). In a paired design, we established 10, 10 m × 7 m plots

to account for spatial differences, with snowmelt manipulation plots paired with control plots (n

= 5). The plots all have a similar slope and aspect, with the downslope side facing the East River

(west). A minimum of 1.6 m separated each plot. Within each plot, we established two 1 m^2

subplots in the center zone of the plot inside a 1 m buffer from the edge of the plot and marked

them to allow repeated measurements in the same location.

The winter of 2022-2023 had a deep snowpack and reached a maximum depth of 246 cm. The

total snowfall for the 2022-2023 winter was 921 cm, 12% less than the average from 1974-2022,

but the greatest snowpack since the 2018-2019 winter. To perform the snowmelt manipulation,

we used black shade cloths that covered the entirety of the treatment plots (10 m × 7 m) attached

to corner snow poles (Blume-Werry et al., 2017). We deployed the shade cloths on April 20th,

2023, and removed them on May 16th, 2023, when the treatment plots were snow-free. On

average, the control plots melted out on May 25th, 2023, meaning the treatment accelerated

snowmelt by approximately 9 days. The snowmelt date has advanced by about 3.5 days per

decade at this site since 1975 (Iler et al., 2013). We took biweekly carbon flux and plant

measurements from early June to mid-August 2023. The growing season starts when the snow

melts in late May and ends when plants senesce in September.

NDVI, air temperature, & soil moisture measurements

We measured NDVI using Red and Near-IR reflectance measurements from a multispectral

camera system (Micasense Altum v1) mounted on an uncrewed aircraft system (UAS, DJI m300

RTK). The instrument was flown on parallel flight lines with 28 m spacing at a height of 100 m

above ground level, yielding images with approximately 85% forward overlap and 75% side

overlap. We mosaiced, georeferenced, and radiometrically calibrated images from each flight

using photogrammetry software (Agisoft Metashape version 1.7) and a standard protocol

utilizing ground control points and a reflectance calibration panel. We computed NDVI [(NIR-

Red)/(NIR+RED)] at 5 cm resolution for each mosaic before mean-aggregating values to 1 m

resolution for time-series analysis. NDVI values for the inner 9 m $\times$ 6 m of the plot were then

averaged to calculate one NDVI value per plot per UAS flight. Flights were conducted near

midday (10 am - 3 pm local time) in sunny conditions approximately once per week during our

sampling period.

To measure air and soil temperature (ºC) and soil moisture (% volumetric soil water content,

VWC), we used TMS-4 sensors (TOMST, Praha, Czech Republic). We installed a single TMS-4

probe in the summer of 2022 in the center of each plot. We collected air temperature at 15 cm

above the soil surface and soil temperature and moisture at 6 cm below the soil surface level

every 15 minutes throughout the growing season (Wild et al., 2019).

Plant community & plant biomass measurements above and belowground

We quantified plant community composition and biomass using the pin-drop method (McLaren

& Turkington, 2011). We placed a 0.5 m^2 frame on each subplot and dropped a pin at 25 evenly

spaced points, recording the type of plant and the number of times the pin hit to measure plant

diversity and to calculate plant biomass. To calculate plant biomass, we built allometric

equations that correspond pin-drop hits to aboveground plant biomass. To collect biomass data,

we set up 15-20 0.5 m^2 plots haphazardly spread outside the experimental plots. Within each

biomass removal plot, we performed a pin-drop measurement as described above. Then, we

clipped all the aboveground plant biomass at ground level, separated it to the species level, dried

it, and weighed it. We collected plant biomass data each sampling week to account for the

seasonal changes in plant biomass. We made one equation to estimate forb biomass, one for

graminoid biomass, and another for shrub biomass (plant classifications in Supplementary Table

S1). To make each equation, we built a linear regression between the number of hits and biomass

for all functional groups separately. In the equation, we included all the different time points to

account for how biomass would change over the growing season and included time as a random

effect in the model. The forb equation was y $= 0.93x + 0.38$, and the relationship had an r^2 value

of 0.66. The shrub equation was y $= 2.26x + 4.52$, and the relationship had an r^2 value of 0.63.

The graminoid equation was y $= 0.58x + 0.387$, and the relationship had an r^2 value of 0.58. The

x in the equations represents the number of pin-drop hits. We estimated the relative abundance of

each functional group pre-treatment and found no difference in percent cover of the functional

groups between control and earlier snowmelt plots (Figure S1).

To estimate root biomass, we randomly collected a bulk soil core (0-20 cm) using a hammer

corer (6 cm in diameter) from each plot (n $= 5$). We sieved the soil to 2 mm to sort the roots from

the soil core. We then spread the sieved rocks, roots, etc., onto a clean surface and picked out the

roots for 15 minutes per sample. We dried the roots at 60° C for 48 hours and weighed them to

determine root biomass. We sampled nine times throughout the growing season.

Carbon flux measurements

To measure how earlier snowmelt impacted carbon fluxes, we measured net ecosystem exchange

(NEE) and ecosystem respiration (ER) at each time point throughout the growing season and

calculated gross primary productivity (GPP) as the sum of those measurements. We measured

NEE in the subplots using an infrared gas analyzer (Li-Cor, Lincoln, Nebraska, USA) inside an

airtight, translucent chamber with a volume of $0.6m^3$ and an area of $1m^2$ (Prager et al., 2021).

The chamber was sealed to the ground with chains to ensure it was airtight. We took the carbon

measurements for 90 seconds at four different light regimes to calculate NEE inside the chamber.

We used full sun where the photosynthetic photon flux density (PPFD) was greater than 1000

μmol m^{-2} s^{-1}, full darkness where the PPFD was zero μmol m^{-2} s^{-1}, and two in-between light

levels. Full darkness is a measurement of ER. To block the light, we used one and two layers of

plastic shade cloths, each reducing the PPFD by ~50% inside the chamber (i.e., to about 25% of

ambient light with two layers). To measure PPFD, we used an MQ-100 Apogee PAR meter

(Apogee Instruments, Logan, UT, USA), which was mounted in the center of the chamber. We

then made a response curve between time and carbon dioxide concentration and removed any

non-linear parts. To calculate NEE (μmol$\cdot$m$^{-2}\cdot$s^{-1}) from the flux measurements, we used the

below equation (Prager et al., 2021) after we converted the carbon dioxide concentrations to dry

molar fractions:

193
$$NEE = \frac{\rho \times V \times (dC/dT)}{A}$$

In this equation, ρ is the air density (mol air/m^3) calculated from *P/RT*, where *P* is the

atmospheric pressure (Pa), *R* is the universal gas constant (8.314 J·air K^{-1}·mol^{-1}), and *T* is the air

temperature (K). *V* is the volume of the chamber (m^3), dC/dT is the rate of change of carbon

dioxide taken from the chamber (µmol·CO_2 mol^{-1}·air s^{-1}), and *A* is the ground area of the

chamber (m^2). We estimated NEE at PPFD = 800 μmol m^{-2} s^{-1} (NEE$_{800}$) to standardize our

measurements over time and space. We estimated NEE$_{800}$ by creating a light response curve with

NEE against PPFD and fitting a line between our four light measurements. We then solved the

equation for PPFD = 800 μmol m^{-2} s^{-1} (Sundqvist et al., 2020). GPP$_{800}$ is then calculated using

the equation:

203                                $$NEE_{800} = GPP_{800} - ER$$

with a negative NEE value indicating net CO_2 uptake and a positive NEE indicating net CO_2

release to the atmosphere.

Carbon analysis

We used a Bayesian modeling approach to estimate how NEE changed every other hour

throughout the growing season in the control and earlier snowmelt plots. We analyzed NEE at

any particular time point as a function of the average soil moisture (*Soilm$_i$*) and air temperature

(*Airtemp$_i$*) in the plot three days before the NEE measurement was taken and included an

intercept. We estimated distinct soil moisture and air temperature parameters for the control and

earlier snowmelt plots. For each time during the season, NEE at plot *i* was estimated using the

following normal likelihood:

214                                $$NEE_i \sim Normal(\lambda_i, \sigma^2)$$

Process model:

$$\lambda_i = \alpha_{(treatment(i))} + \beta 1_{(treatment(i))} \times Soilm_i + \beta 2_{(treatment(i))} \times Airtemp_i$$

We estimated the parameters from uninformed prior distributions (with a precision parameter):

$\alpha_* \sim Normal(0, 0.0001)$

$\beta^*_* \sim Normal(0, 0.0001)$

$1/\sigma^2 \sim Gamma(0.0001, 0.0001)$

We used the fitted parameters estimates (α, $\beta 1$, and $\beta 2$) and their associated variance to estimate

NEE in all plots. Soil moisture and air temperature averages, calculated every other hour for each

plot, were used as input data to predict NEE values throughout the growing season. We used

JAGS software (Plummer, 2003) and the *rjags* package in R (Plummer, 2023) to run the analysis.

We ran the model for 10,000 iterations following an initial burn-in period of 10,000 iterations.

We used three MCMC chains and monitored their mixing to determine convergence. Each model

parameter had well-mixed chains and a smooth and even posterior distribution after 10,000

iterations. The goodness of fit was calculated as the r^2 between predicted and observed NEE

values to check model fit. All parameter values (means, SDs, and 95% CIs) are in Supplementary

Table S2. Using similar methods and the model structure described above, we also ran predictive

models for GPP and ER.

Data analysis

To determine the effect of earlier snowmelt on total aboveground biomass, forb biomass, shrub

biomass, graminoid biomass, root biomass, NEE, GPP, and ER throughout the growing season,

we ran linear mixed effect models using the package *nlme* (Pinheiro et al., 2023; Pinheiro &

Bates, 2000) in R, version 4.15 (R Core Team, 2023). We ran an identical linear mixed-effects

model for all the analyses. We included the interaction between the treatment and date (as day of

the year, a continuous variable) as fixed effects. The plot was included as a random effect, and

we included a correlation structure to account for temporal autocorrelation. To determine the

effect of earlier snowmelt on total aboveground biomass, forb biomass, shrub biomass,

graminoid biomass, root biomass, NEE, GPP, and ER at each measurement point throughout the

growing season, we calculated the standardized mean difference (SMD) using Hedge's *d* metric

of effect size using the package *effsize* (Torchiano, 2020) in R, version 4.15 (R Core Team,

2023). If SMD > 0, earlier snowmelt increased the variable (a positive effect), if SMD < 0 (a

negative effect), earlier snowmelt caused a decrease in the variable. To examine the relationship

between carbon fluxes and plant community composition, we used distance-based redundancy

analysis (dbRDA). We built six separate models for each time point we sampled throughout the

summer, with GPP, ER, NEE, and treatment in the model (plant composition ~ GPP + ER +

Treatment + Condition(Plot), strata= Plot). The models were run on Bray-Curtis distance

similarity matrices of plant composition, with the plot as a random effect. Each dbRDA model

was tested with 99,999 permutations using the package *Vegan* (Oksanen et al., 2024). We also

ran correlation tests between the plant biomass of each plant functional type (graminoid, shrub,

and forb) and each carbon flux (NEE, GPP, and ER) to see how they were correlated with each

other across the season.

Results

Earlier snowmelt had a minimal effect on soil moisture early in the season but resulted in an 11%

decrease in soil moisture from mid-June to mid-August (treatment × time, Figure S2, Table 1).

Soil moisture changed significantly over time (Figure S2, Table 1). While earlier snowmelt did

not significantly affect air temperature, air temperature fluctuated significantly throughout the

growing season, peaking in late July (Figure S2, Table 1).

Total aboveground biomass increased during the growing season, ranging from an average

minimum of 110 g/m^2 in early June to a maximum of 183 g/m^2 in mid-July. However, earlier

snowmelt did not significantly affect the total aboveground biomass (Figure 1, Table 1). Root

biomass varied within each sampling point, between treatments, and throughout the growing

season, with earlier snowmelt plots showing an advancement in peak root biomass (Figure 1,

Table 1). NDVI was higher in the earlier snowmelt plots early in the season but became more

similar to the control plots as the season progressed (treatment × time, Figure S3, Table 1).

NDVI showed a significant treatment effect, with earlier snowmelt plots being greener (Figure

S3, Table 1).

At the first sampling point in early June, graminoid biomass was significantly higher in the

earlier snowmelt plots (Figure 2b), and on average, treatment plots had 47% more graminoid

biomass throughout the growing season (Figure 2a). Graminoid biomass changed significantly

over the growing season (Table 1). Shrub biomass was 22% lower in the earlier snowmelt plots

over the growing season (Figure 2a), with the effect being more pronounced at the end of the

season than at the beginning of the season (Figure 2b). However, these trends were not

statistically significant. Shrub biomass changed significantly over the growing season (Table 1),

generally increasing throughout the season (Figure 2a). Forb biomass was not affected by earlier

snowmelt but did show significant seasonal changes, peaking in mid-July and declining for the

remainder of the season.

NEE changed significantly over the season (Figure 3a, Table 1). Both the control and snowmelt

plots were carbon sources in the spring but became carbon sinks by early July. At no point was

there a significant difference in NEE between the earlier snowmelt and the control plots (Figure

3b). GPP also varied significantly over the growing season, starting low, peaking in early July,

and then decreasing through the end of the season. The control and earlier snowmelt plots did not

differ significantly, though there was a significant treatment $\times$ time interaction (Table 1). In early

June, the advanced snowmelt plots showed 206% greater GPP than the control plots, but this

effect diminished over the season, and by mid-July, GPP in the snowmelt plots was less than in

the control plots (Figure 3b). ER did not change significantly over the season and was not

significantly impacted by the treatment.

The model fit for predicted NEE over the season (predicted vs. observed r^2) was 0.35. Air

temperature was a significant predictor of NEE in the control plots, such that higher air

temperature correlated with a stronger carbon sink (Figure 4a). Soil moisture was not a

significant parameter, but greater soil moisture was correlated with a weaker carbon sink (Figure

4a). In the earlier snowmelt plots, the relationships between air temperature, soil moisture, and

NEE were weaker, though they had the same trends as the control plots (Figure 4a). Drier soils in

the earlier snowmelt plots and a weaker relationship between soil moisture and air temperature

conditions resulted in smaller daily NEE fluctuations and less overall ER and GPP (Figure 4b).

Over the entire growing season, cumulative NEE was 5051 μmol m^{-2} s^{-1} in the control plots but

only 1468 μmol m^{-2} s^{-1} in the earlier snowmelt plots, a 70% reduction in carbon sink strength

with earlier snowmelt (Figure 4c). Modeled predicted carbon fluxes showed more GPP but also

more ER in the earlier snowmelt plots (Figures S4 & S5).

The dbRDA results from the six models throughout the growing season did not reveal any

significant results (Figure S6). NEE, GPP, ER, and treatment are never significant variables in

the models. Model one is from June 5th ($r^2 = 0.13$), model two is from June 19th ($r^2 = 0.28$),

model three is from July 5th ($r^2 = 0.09$), model four is from July 21st ($r^2 = 0.15$), model 5 is from

August 1st ($r^2 = 0.08$), and model 6 is from August 15th ($r^2 = 0.10$). Although the models were

not significant, *Dasiphora fruticosa* (the most abundant shrub species) and *Festuca thurberi* (the

most abundant grass species) had the largest species scores at each time point. *D. fruticosa* was

generally positioned near GPP in the dbRDA ordination space, while *F. thurberi* was near ER.

Pearson's correlations between plant functional groups and carbon fluxes at each time point

revealed patterns in the data, but these were not statistically significant.

Discussion

Winters are changing, and earlier snowmelt is occurring more frequently with climate change.

However, there is still uncertainty about how earlier snowmelt will impact the carbon balance of

ecosystems. We advanced snowmelt by nine days in a montane and found that earlier snowmelt

reduced the carbon uptake of a montane meadow, likely due to drier soils later in the season.

While earlier snowmelt increased GPP at the start of the growing season, it resulted in lower

cumulative carbon accumulation (less negative NEE) across the entire season. The timing of

aboveground green-up and root growth shifted earlier in the year due to earlier snowmelt. Shrubs

and graminoids showed the strongest responses, with higher graminoid biomass in early

snowmelt plots and greater shrub biomass in control plots. These shifts in plant community

biomass likely influenced carbon flux patterns, as graminoids tended to have a positive

relationship with ER. Relatively small shifts in snowmelt date will have large implications for

carbon storage in montane meadows.

Total aboveground biomass did not change with an earlier snowmelt date, supporting our

hypothesis and previous work suggesting that drought might have a larger effect on total

aboveground biomass than snowmelt timing (Möhl et al., 2023). Contrary to our hypothesis, total

root biomass remained unchanged with earlier snowmelt, despite contradictory findings in other

studies. Some studies show that increased snow leads to more root biomass (Li et al., 2020),

while others show the opposite (D'Imperio et al., 2018). Root biomass typically responds to soil

moisture, increasing in drier ecosystems to optimize water acquisition (Liu et al., 2018).

However, roots can maintain similar biomass while changing branching patterns or their

diameters in response to water stress (Yetgin, 2024). In our experiment, earlier snowmelt may

not have significantly altered soil moisture enough to impact overall root biomass, especially

given how similar soil moisture was between the treatments in the early season.

Although earlier snowmelt did not affect total above or below-ground biomass, it did shift root

phenology and the timing of aboveground green-up. Peak root growth occurred earlier in the

year, which contrasts with previous studies showing that root phenology is relatively

unresponsive to snowmelt timing (Blume-Werry et al., 2017). A recent meta-analysis of 88

studies found that herbaceous root phenology was unaffected by warming, while aboveground

herbaceous plants shifted their growing season earlier (Liu et al., 2022). However, warmer

temperatures throughout the growing season and changes in the snowmelt timing could affect

root phenology differently. Despite peak root biomass shifting earlier with advanced snowmelt,

we did not observe a shift in the timing of aboveground peak biomass. However, earlier

snowmelt did lead to earlier green-up (higher NDVI; Figure S3) and greater productivity (higher

GPP; Figure 3a). The disparity between the treatment effects on the timing of biomass

production and green-up suggests that though the biomass of perennial woody structures

remained stable, green-up timing and productivity were more responsive to earlier snowmelt.

Traditionally, phenology focuses on life history events such as bud burst and flowering (Gezon

et al., 2016; Inouye, 2008; Sherwood et al., 2017; Slatyer et al., 2022). However, focusing on the

timing of ecosystem functions, such as productivity (Chen et al., 2019; Galvagno et al., 2013),

can clarify how changing snowmelt timing affects the carbon cycle on the ecosystem scale.

In plots with earlier snowmelt, graminoid biomass was higher, especially early in the season, and

shrub biomass was lower throughout the season. Graminoids are particularly responsive to

changes in snowmelt timing and winter snowpack, likely due to shifts in soil moisture (Ma et al.,

2024). However, the response of graminoids to drought conditions can vary depending on the

plant community context and the relative soil moisture of the ecosystem (Gherardi & Sala, 2015;

Li et al., 2020; Liu et al., 2018; Rosbakh et al., 2017). Since grasses typically have relatively

shallow fibrous roots, the early-season moisture pulse from snowmelt may have stimulated their

growth, sustaining high biomass throughout the growing season. Previous work supports that

early-season abiotic conditions have a large impact on total biomass throughout the growing

season (Guo et al., 2017; Shaw et al., 2022). Additionally, snowmelt is often coupled with a

pulse of nitrogen (Miller et al., 2009), and graminoids are more responsive to short-term nutrient

changes than shrubs (Carbonell-Silletta et al., 2024). While shrubs tend to increase their biomass

with warming temperatures (Amagai et al., 2018; Broadbent et al., 2022) and have strong

phenological responses to changes in snowmelt timing (Wipf & Rixen, 2010), we found lower

shrub biomass with earlier snowmelt. This could indicate that shrub biomass responses to

changes in snow differ from warming effects, as shrubs tend to be slow-growing and may

respond more to long-term warming experiments than short-term changes in snowmelt timing.

Our study found that early snowmelt had a greater effect on GPP than ER. This is similar to

findings from 119 site-years of eddy flux tower data in the Arctic, where productivity was

impacted by snowmelt timing, but there was little impact on ecosystem respiration (Zona et al.,

2022). Similarly, a meta-analysis of 49 studies looking at the effect of snow depth on carbon

dynamics found no substantial impact of snow depth on respiration (Li et al., 2016), consistent

with our results. One study found that earlier snowmelt influences annual NEE by affecting

respiration; however, this effect is attributed to reductions in winter respiration caused by earlier

snowmelt rather than changes in growing season respiration (Galvagno et al., 2013). We

observed a significant interaction between treatment and time for productivity but not

respiration. This finding is consistent with other studies showing that earlier snowmelt boosts

early-season productivity (Chen et al., 2019; Sherwood et al., 2017; Vorkauf et al., 2021; Wipf &

Rixen, 2010; Zona et al., 2022), although many studies focus more on phenology than on

productivity (Wipf & Rixen, 2010). A review of snow manipulation experiments in Arctic and

alpine ecosystems found that flower phenology was particularly responsive to the snowmelt

timing and tracked earlier snowmelt (Wipf & Rixen, 2010), likely boosting early plant

production. In our study, the increase in GPP with early snowmelt was significant only early in

the season, and this diminished as the season progressed.

When modeling the cumulative effect of earlier snowmelt on the carbon budget, we found that

montane meadows with earlier snowmelt dates act as weaker carbon sinks. This is because

correlations with abiotic factors become weaker, and conditions become drier. The extended

growing season may weaken the relationship between NEE, temperature, and moisture,

potentially making NEE more correlated with light availability. Additionally, drier soils later in

the season disrupted the alignment of peak soil moisture and air temperature, further weakening

the relationship between snowmelt timing and carbon fluxes. Our study is one of the first to

measure NEE in response to experimentally manipulated snowmelt timing. Most studies

examining carbon fluxes in relation to snow changes use natural variation in snowmelt days

across years (Chen et al., 2019; Galvagno et al., 2013; López-Blanco et al., 2017; Zona et al.,

2022), with varying results. Some studies report that earlier snowmelt shifts the growing season

forward, increasing productivity in the spring (Chen et al., 2019; Zona et al., 2022), while others

find no effect on NEE because changes in GPP and ER cancel each other out (López-Blanco et

al., 2017). In some cases, earlier snowmelt decreased NEE due to lower respiration rates

(Galvagno et al., 2013).

Changes in plant community composition can also affect the carbon balance, as graminoids and

shrubs process carbon differently. Although not statistically significant, the dbRDA analysis

suggested that the most abundant graminoid, *F. thurberi*, was most correlated with ER, and the

most abundant shrub, *D. fruticosa*, was more correlated with GPP. This likely reflects

differences in carbon allocations between grasses and shrubs. Shrubs allocate more resources

aboveground, while grasses invest in root production, causing grasses to respire around 51%

more carbon than shrubs (Carbone & Trumbore, 2007). Graminoids also tend to have higher

respiration rates than shrubs (Jin et al., 2010; Quin et al., 2015) and produce more litter, which

can increase respiration during decomposition. Compared to shrubs, grasses require more energy

for photosynthesis (Quin et al., 2015). Our study observed a shift in plant community

composition with earlier snowmelt, with higher grass abundance and lower shrub abundance,

which may explain the lower overall carbon uptake in those plots. Correlations between

functional group biomass and carbon fluxes also showed that higher graminoid biomass resulted

in higher respiration (Figure S7).

Environmental gradients alone cannot fully capture climate change effects as they do not have

any controls, making experiments valuable for new insights as they can pinpoint mechanisms

while accounting for confounding factors (Metz & Tielbörger, 2016; Rysavy et al., 2014).

Integrating natural observations and experiments is key to understanding climate change impacts

(Prager et al., 2022). Approximately 55% of snow studies rely on observations and 45% on

experiments, but most focus on plant phenology or community diversity responses (Slatyer et al.,

2022). Natural gradients usually exhibit larger variations in snowmelt timing than experiments,

potentially leading to greater ecosystem responses (Rixen et al., 2022). However, natural

gradients tend to incorporate areas with frequent early snowmelt due to their slope or aspect,

where vegetation is likely adapted to variable conditions (Rixen et al., 2022). Ecosystem

manipulation experiments offer a way to assess the impacts of earlier snowmelt on ecosystems

where snowmelt variation is historically minimal, which is important as climate change alters

ecosystems with historically less variation. In our experimental meadows, snowmelt has

advanced by 3.5 days per decade (Iler et al., 2013), and we advanced snowmelt by about nine

days. Our results, therefore, estimate the carbon balance approximately three decades into the

future, around 2055.

In conclusion, earlier snowmelt weakens the carbon sink strength in montane meadows due to

abiotic and biotic factors. Abiotic factors include drier soils later in the season—exacerbated by

weak monsoon conditions in summer 2023—while biotic factors include higher graminoid

biomass and lower shrub biomass, with graminoids respiring more. As summer seasons become

drier due to climate change, snowmelt-induced water stress may be exacerbated. Phenological

shifts in plants both above- and belowground will also occur with earlier snowmelt. Our study

underscores how even small environmental changes, such as a less-than-two-week advance in

snowmelt, can significantly affect ecosystems, potentially reducing carbon storage in montane

meadows. Understanding how ecosystems respond to changes in snowmelt timing is crucial for

predicting the impacts of climate variability.

Acknowledgments

This study was supported by the U.S. Department of Energy, Office of Science, Office of

Biological and Environmental Research, Terrestrial Ecosystem Sciences program under award

number DE-FOA-0002392. PS was partially supported by the Watershed Function Scientific

Focus Area at Lawrence Berkeley National Laboratory, funded by the U.S. Department of

Energy, Office of Science, Office of Biological and Environmental Research under Award

Number DE-AC02-05CH11231. NDVI data were provided by Ian Breckheimer and Amanda

Henderson at Rocky Mountain Biological Laboratory. This project was funded by a Rocky

Mountain Biological Laboratory Graduate Student Fellowship, an Institute of Global Change

Biology Fellowship, and the Dr. Nancy Williams Walls Award for Field Research to OKV.

Figures

453
454 **Figure 1.** a) Aboveground biomass and root biomass over time in control and earlier snowmelt
455 plots. b) The effect sizes of the treatment on aboveground biomass and root biomass. The effect
456 sizes are calculated using Hedge's d, comparing the control versus the snowmelt plots. Above
457 the zero line indicates that the biomass value in the snowmelt plots was greater than the control
458 plots.

459
460
461 **Figure 2.** a) Graminoid, shrub, and forb biomass in control and earlier snowmelt plots over time.
462 b) The effect sizes of the treatment on aboveground biomass and root biomass. The effect sizes
463 are calculated using Hedge's d, comparing the control versus the earlier snowmelt plots. Above
464 the zero line indicates that the biomass value in the earlier snowmelt plots was greater than the
465 control plots.

466
Figure 3. a) NEE, GPP, and ER across the growing season in earlier snowmelt and control plots.
Carbon fluxes are on the y-axis, and the date is on the x-axis. b) The effect sizes of the treatment
on NEE, GPP, and ER. The effect sizes are calculated using Hedge's d, comparing the control
versus the snowmelt plots. Values above the zero line indicate that the carbon value in the
snowmelt plots was greater than the control plots. For NEE, any effect size values above the zero
line indicate that earlier snowmelt plots are a greater carbon source.

473
Figure 4. a) Parameter values for the Bayesian model, median plus 95% confidence interval.
Values that do not cross zero are significant. Any number below zero indicates that the abiotic
factor causes a larger carbon sink, and any number above zero indicates that the abiotic factor
causes a greater carbon source. b) NEE from late May to late August in earlier snowmelt and
control plots. Data were modeled using a Bayesian model, and NEE was predicted based on soil
moisture and air temperature values taken from TMS-4 probes. Blue and orange are the 50%
quantile measurements, and the light blue and light orange around is the 95% confidence
interval. NEE values below the zero line indicate a carbon sink, while NEE values above the zero
line indicate a carbon source. c) Cumulative NEE, summed from late May to late August in
control and earlier snowmelt plots.

Tables

Table 1. Linear mixed effect model results for aboveground biomass, belowground biomass,
GPP, ER, and NEE, with a bolded p-value indicating a significant effect.

Response	Treatment			Time			Treatment × Time		
	df	F	p	df	F	p	df	F	p
Total biomass	1,18	0.0335	0.8568	1,98	36.7935	**<0.0001**	1,98	2.1718	0.01438
Shrub biomass	1,9	1.0411	0.3342	1,49	22.4280	**<0.0001**	1,49	2.3947	0.1282
Forb biomass	1,18	0.0398	0.8441	1,97	14.5089	**0.0002**	1,97	0.9353	0.3359
Graminoid biomass	1,18	1.6902	0.2100	1,94	18.5403	**<0.0001**	1,94	0.3538	0.5534
Root biomass	1,8	0.2260	0.6472	1,75	0.2037	0.6531	1,75	0.5689	0.4531
NEE	1,18	0.2036	0.6572	1,98	20.9097	**<0.0001**	1,98	1.9490	0.1658
GPP	1,18	0.0185	0.8934	1,98	17.1742	**0.0001**	1,98	4.9033	**0.0291**
ER	1,18	0.0869	0.7715	1,98	0.1040	0.7478	1,98	2.2621	0.1358
Soil moisture	1,8	0.8500	0.3846	1,20148	149883	**<0.0001**	1,20148	652.37	**<0.0001**
Air temperature	1,8	0.1510	0.7079	1,20148	395.02	**<0.0001**	1,20148	0.0930	0.7601
NDVI	1,8	9.1886	**0.0163**	1,128	73.002	**<0.0001**	1,183	12.229	**0.0006**

References

Amagai, Y., Kudo, G., & Sato, K. (2018). Changes in alpine plant communities under climate change: Dynamics of snow-meadow vegetation in northern Japan over the last 40 years. *Applied Vegetation Science*, *21*(4), 561–571. https://doi.org/10.1111/AVSC.12387

Blume-Werry, G., Jansson, R., & Milbau, A. (2017). Root phenology unresponsive to earlier snowmelt despite advanced above-ground phenology in two subarctic plant communities. *Functional Ecology*, *31*(7), 1493–1502. https://doi.org/10.1111/1365-2435.12853

Broadbent, A. A. D., Bahn, M., Pritchard, W. J., Newbold, L. K., Goodall, T., Guinta, A., Snell, H. S. K., Cordero, I., Michas, A., Grant, H. K., Soto, D. X., Kaufmann, R., Schloter, M., Griffiths, R. I., & Bardgett, R. D. (2022). Shrub expansion modulates belowground impacts of changing snow conditions in alpine grasslands. *Ecology Letters*, *25*(1), 52–64. https://doi.org/10.1111/ELE.13903

Carbone, M. S., & Trumbore, S. E. (2007). Contribution of new photosynthetic assimilates to respiration by perennial grasses and shrubs: Residence times and allocation patterns. *New Phytologist*, *176*(1), 124–135. https://doi.org/10.1111/j.1469-8137.2007.02153.x

Carbonell-Silletta, L., Scholz, F. G., Burek, A., Villa, V. D., Cavallaro, A., Askenazi, J. O., Arias, N. S., Hao, G.-Y., Goldstein, G., & Bucci, S. J. (2024). Nitrogen rather than water

availability limits aboveground primary productivity in an arid ecosystem: Substantial differences between grasses and shrubs. *Ecohydrology*, *17*(3), e2636. https://doi.org/10.1002/eco.2636

Chen, S., Huang, Y., & Wang, G. (2019). Response of vegetation carbon uptake to snow-induced phenological and physiological changes across temperate China. *Science of The Total Environment*, *692*, 188–200. https://doi.org/10.1016/j.scitotenv.2019.07.222

Darrouzet-Nardi, A., Steltzer, H., Sullivan, P. F., Segal, A., Koltz, A. M., Livensperger, C., Schimel, J. P., & Weintraub, M. N. (2019). Limited effects of early snowmelt on plants, decomposers, and soil nutrients in Arctic tundra soils. *Ecology and Evolution*, *9*(4), 1820–1844. https://doi.org/10.1002/ECE3.4870

D'Imperio, L., Arndal, M. F., Nielsen, C. S., Elberling, B., & Schmidt, I. K. (2018). Fast responses of root dynamics to increased snow deposition and summer air temperature in an arctic wetland. *Frontiers in Plant Science*, *9*, 1258. https://doi.org/10.3389/FPLS.2018.01258/BIBTEX

Galvagno, M., Wohlfahrt, G., Cremonese, E., Rossini, M., Colombo, R., Filippa, G., Julitta, T., Manca, G., Siniscalco, C., Morra Di Cella, U., & Migliavacca, M. (2013). Phenology and carbon dioxide source/sink strength of a subalpine grassland in response to an exceptionally short snow season. *Environmental Research Letters*, *8*(2), 025008. https://doi.org/10.1088/1748-9326/8/2/025008

Gamon, J. A., Huemmrich, K. F., Stone, R. S., & Tweedie, C. E. (2013). Spatial and temporal variation in primary productivity (NDVI) of coastal Alaskan tundra: Decreased vegetation growth following earlier snowmelt. *Remote Sensing of Environment*, *129*, 144–153. https://doi.org/10.1016/j.rse.2012.10.030

Ganjurjav, H., Gao, Q., Schwartz, M. W., Zhu, W., Liang, Y., Li, Y., Wan, Y., Cao, X., Williamson, M. A., Jiangcun, W., Guo, H., & Lin, E. (2016). Complex responses of spring vegetation growth to climate in a moisture-limited alpine meadow. *Scientific Reports 2016 6:1*, *6*(1), 1–10. https://doi.org/10.1038/srep23356

Gezon, Z. J., Inouye, D. W., & Irwin, R. E. (2016). Phenological change in a spring ephemeral: Implications for pollination and plant reproduction. *Global Change Biology*, *22*(5), 1779–1793. https://doi.org/10.1111/GCB.13209

Gherardi, L. A., & Sala, O. E. (2015). Enhanced precipitation variability decreases grass- and increases shrub-productivity. *Proceedings of the National Academy of Sciences*, *112*(41), 12735–12740. https://doi.org/10.1073/pnas.1506433112

Gottlieb, A. R., & Mankin, J. S. (2024). Evidence of human influence on Northern Hemisphere snow loss. *Nature*, *625*(7994), 293–300. https://doi.org/10.1038/s41586-023-06794-y

Guo, L., Cheng, J., Luedeling, E., Koerner, S. E., He, J. S., Xu, J., Gang, C., Li, W., Luo, R., & Peng, C. (2017). Critical climate periods for grassland productivity on China's Loess Plateau. *Agricultural and Forest Meteorology*, *233*, 101–109. https://doi.org/10.1016/J.AGRFORMET.2016.11.006

Hu, J., Hopping, K. A., Bump, J. K., Kang, S., & Klein, J. A. (2013). Climate Change and Water Use Partitioning by Different Plant Functional Groups in a Grassland on the Tibetan Plateau. *PLOS ONE*, *8*(9), e75503. https://doi.org/10.1371/JOURNAL.PONE.0075503

Ibanez, M., Altimir, N., Ribas, A., Eugster, W., & Sebastia, M. T. (2020). Phenology and plant functional type dominance drive CO2 exchange in seminatural grasslands in the Pyrenees. *The Journal of Agricultural Science*, *158*(1–2), 3–14. https://doi.org/10.1017/S0021859620000179

Iler, A. M., Høye, T. T., Inouye, D. W., & Schmidt, N. M. (2013). Nonlinear flowering responses to climate: Are species approaching their limits of phenological change? *Philosophical Transactions of the Royal Society B: Biological Sciences*, *368*(1624), 20120489. https://doi.org/10.1098/rstb.2012.0489

Inouye, D. W. (2008). Effects of climate change on phenology, frost damage, and floral abundance of montane wildflowers. *Ecology*, *89*(2), 353–362.

Jin, Z., Dong, Y. S., Qi, Y. C., & An, Z. S. (2010). Soil respiration and net primary productivity in perennial grass and desert shrub ecosystems at the Ordos Plateau of Inner Mongolia, China. *Journal of Arid Environments*, *74*(10), 1248–1256. https://doi.org/10.1016/j.jaridenv.2010.05.018

Kawai, Y., & Kudo, G. (2021). Climate change shifts population structure and demographics of an alpine herb, Anemone narcissiflora ssp. Sachalinensis (Ranunculaceae), along a snowmelt gradient. *Population Ecology*, *63*(3), 260–271. https://doi.org/10.1002/1438-390X.12089

Kelsey, K. C., Pedersen, S. H., Leffler, A. J., Sexton, J. O., Feng, M., & Welker, J. M. (2021). Winter snow and spring temperature have differential effects on vegetation phenology and productivity across Arctic plant communities. *Global Change Biology*, *27*(8), 1572–1586. https://doi.org/10.1111/GCB.15505

Kudo, G. (2021). Habitat-specific effects of flowering advance on fruit-set success of alpine plants: A long-term record of flowering phenology and fruit-set success of Rhododendron aureum. *Alpine Botany*, *131*(1), 53–62. https://doi.org/10.1007/S00035-021-00248-9/FIGURES/6

Lambert, A. M., Miller-Rushing, A. J., & Inouye, D. W. (2010). Changes in snowmelt date and summer precipitation affect the flowering phenology of Erythronium grandiflorum (glacier lily; Liliaceae). *American Journal of Botany*, *97*(9), 1431–1437. https://doi.org/10.3732/AJB.1000095

Li, P., Sayer, E. J., Jia, Z., Liu, W., Wu, Y., Yang, S., Wang, C., Yang, L., Chen, D., Bai, Y., & Liu, L. (2020). Deepened winter snow cover enhances net ecosystem exchange and stabilizes plant community composition and productivity in a temperate grassland. *Global Change Biology*, *26*(5), 3015–3027. https://doi.org/10.1111/GCB.15051

Li, W., Wu, J., Bai, E., Jin, C., Wang, A., Yuan, F., & Guan, D. (2016). Response of terrestrial carbon dynamics to snow cover change: A meta-analysis of experimental manipulation (II). *Soil Biology and Biochemistry*, *103*, 388–393. https://doi.org/10.1016/j.soilbio.2016.09.017

Liu, H., Mi, Z., Lin, L., Wang, Y., Zhang, Z., Zhang, F., Wang, H., Liu, L., Zhu, B., Cao, G., Zhao, X., Sanders, N. J., Classen, A. T., Reich, P. B., & He, J.-S. (2018). Shifting plant species composition in response to climate change stabilizes grassland primary production. *Proceedings of the National Academy of Sciences*, *115*(16), 4051–4056. https://doi.org/10.1073/pnas.1700299114

Liu, H., Wang, H., Li, N., Shao, J., Zhou, X., van Groenigen, K. J., & Thakur, M. P. (2022). Phenological mismatches between above- and belowground plant responses to climate warming. *Nature Climate Change*, *12*(1), 97–102. https://doi.org/10.1038/s41558-021-01244-x

Liu, H., Xiao, P., Zhang, X., Chen, S., Wang, Y., & Wang, W. (2023). Winter snow cover influences growing-season vegetation productivity non-uniformly in the Northern Hemisphere. *Communications Earth & Environment*, *4*(1), 1–10.

https://doi.org/10.1038/s43247-023-01167-9

López-Blanco, E., Lund, M., Williams, M., Tamstorf, M. P., Westergaard-Nielsen, A., Exbrayat,
J. F., Hansen, B. U., & Christensen, T. R. (2017). Exchange of CO2 in Arctic tundra:
Impacts of meteorological variations and biological disturbance. *Biogeosciences*, *14*(19),
4467–4483. https://doi.org/10.5194/bg-14-4467-2017

Ma, W., Hu, J., Zhang, B., Guo, J., Zhang, X., & Wang, Z. (2024). Later-melting rather than
thickening of snowpack enhance the productivity and alter the community composition of
temperate grassland. *Science of The Total Environment*, *923*, 171440.
https://doi.org/10.1016/j.scitotenv.2024.171440

McLaren, J. R., & Turkington, R. (2011). Biomass compensation and plant responses to 7 years
of plant functional group removals. *Journal of Vegetation Science*, *22*(3), 503–515.
https://doi.org/10.1111/j.1654-1103.2011.01263.x

Metz, J., & Tielbörger, K. (2016). Spatial and temporal aridity gradients provide poor proxies for
plant–plant interactions under climate change: A large-scale experiment. *Functional*
Ecology, *30*(1), 20–29. https://doi.org/10.1111/1365-2435.12599

Miller, A. E., Schimel, J. P., Sickman, J. O., Skeen, K., Meixner, T., & Melack, J. M. (2009).
Seasonal variation in nitrogen uptake and turnover in two high-elevation soils:
Mineralization responses are site-dependent. *Biogeochemistry*, *93*(3), 253–270.
https://doi.org/10.1007/S10533-009-9301-4/TABLES/4

Möhl, P., Vorkauf, M., Kahmen, A., & Hiltbrunner, E. (2023). Recurrent summer drought affects
biomass production and community composition independently of snowmelt
manipulation in alpine grassland. *Journal of Ecology*, *111*(11), 2357–2375.
https://doi.org/10.1111/1365-2745.14180

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R.
B., Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M.,
Bolker, B., Borcard, D., Carvalho, G., Chirico, M., Caceres, M. D., Durand, S., …
Weedon, J. (2024). *vegan: Community Ecology Package*. https://CRAN.R-
project.org/package=vegan

Pepin, N., Bradley, R. S., Diaz, H. F., Baraer, M., Caceres, E. B., Forsythe, N., Fowler, H.,
Greenwood, G., Hashmi, M. Z., Liu, X. D., Miller, J. R., Ning, L., Ohmura, A., Palazzi,
E., Rangwala, I., Schöner, W., Severskiy, I., Shahgedanova, M., Wang, M. B., … Yang,
D. Q. (2015). Elevation-dependent warming in mountain regions of the world. *Nature*
Climate Change 2015 5:5, *5*(5), 424–430. https://doi.org/10.1038/NCLIMATE2563

Pinheiro, J. C., & Bates, D. M. (2000). *Mixed-Effects Models in S and S-PLUS*. Springer.
https://doi.org/10.1007/b98882

Pinheiro, J. C., Bates, D., & R Core Team. (2023). *nlme: Linear and Nonlinear Mixed Effects*
Models (R package version 3.1-163). https://CRAN.R-project.org/package=nlme

Plummer, M. (2003). JAGS: A Program for Analysis of Bayesian Graphical Models Using Gibbs
Sampling. *Proceedings of the 3rd International Workshop on Distributed Statistical*
Computing (DSC 2003), 1–10.

Plummer, M. (2023). *rjags: Bayesian Graphical Models using MCMC* (R package version 4-15).
https://CRAN.R-project.org/package=rjags

Prager, C. M., Classen, A. T., Sundqvist, M. K., Barrios-Garcia, M. N., Cameron, E. K., Chen,
L., Chisholm, C., Crowther, T. W., Deslippe, J. R., Grigulis, K., He, J.-S., Henning, J. A.,
Hovenden, M., Høye, T. T. T., Jing, X., Lavorel, S., McLaren, J. R., Metcalfe, D. B.,
Newman, G. S., … Sanders, N. J. (2022). Integrating natural gradients, experiments, and

statistical modeling in a distributed network experiment: An example from the WaRM
Network. *Ecology and Evolution*, *12*(10), e9396. https://doi.org/10.1002/ECE3.9396
Prager, C. M., Jing, X., Henning, J. A., Read, Q. D., Meidl, P., Lavorel, S., Sanders, N. J.,
Sundqvist, M., Wardle, D. A., Classen, A. T., Prager, C. :, Jing, X., Henning, J. A., Read,
Q. D., Meidl, P., Lavorel, S., Sanders, N. J., Sundqvist, M., Wardle, D. A., & Classen, A.
T. (2021). Climate and multiple dimensions of plant diversity regulate ecosystem carbon
exchange along an elevational gradient. *Ecosphere*, *12*(4).
https://doi.org/10.1002/ecs2.3472
Quin, S. L. O., Artz, R. R. E., Coupar, A. M., & Woodin, S. J. (2015). *Calluna vulgaris*-
dominated upland heathland sequesters more CO2 annually than grass-dominated upland
heathland. *Science of The Total Environment*, *505*, 740–747.
https://doi.org/10.1016/j.scitotenv.2014.10.037
R Core Team. (2023). *R: A Language and Environment for Statistical Computing*. R Foundation
for Statistical Computing. https://www.R-project.org/
Rixen, C., Thomas Høye, T., Macek, P., Aerts, R., Alatalo, J. M., Anderson, J. T., Arnold, P. A.,
Barrio, I. C., Bjerke, J. W., Björkman, M. P., Blok, D., Blume-Werry, G., Boike, J.,
Bokhorst, S., Carbognani, M., Christiansen, C. T., Convey, P., Cooper, E. J., Hans
Cornelissen, J. C., … Blok, D. (2022). Winters are changing: Snow effects on Arctic and
alpine tundra ecosystems. *Https://Doi.Org/10.1139/as-2020-0058*, 1–37.
https://doi.org/10.1139/AS-2020-0058
Rosbakh, S., Leingärtner, A., Hoiss, B., Krauss, J., Steffan-Dewenter, I., & Poschlod, P. (2017).
Contrasting Effects of Extreme Drought and Snowmelt Patterns on Mountain Plants
along an Elevation Gradient. *Frontiers in Plant Science*, *8*.
https://doi.org/10.3389/fpls.2017.01478
Rysavy, A., Seifan, M., Sternberg, M., & Tielbörger, K. (2014). Shrub seedling survival under
climate change – Comparing natural and experimental rainfall gradients. *Journal of Arid
Environments*, *111*, 14–21. https://doi.org/10.1016/j.jaridenv.2014.07.004
Semenchuk, P. R., Gillespie, M. A. K., Rumpf, S. B., Baggesen, N., Elberling, B., & Cooper, E.
J. (2016). High Arctic plant phenology is determined by snowmelt patterns but duration
of phenological periods is fixed: An example of periodicity. *Environmental Research
Letters*, *11*(12), 125006. https://doi.org/10.1088/1748-9326/11/12/125006
Shaw, E. A., White, C. T., Silver, W. L., Suding, K. N., & Hallett, L. M. (2022). Intra-annual
precipitation effects on annual grassland productivity and phenology are moderated by
community responses. *Journal of Ecology*, *110*(1), 162–172.
https://doi.org/10.1111/1365-2745.13792
Sherwood, J. A., Debinski, D. M., Caragea, P. C., & Germino, M. J. (2017). Effects of
experimentally reduced snowpack and passive warming on montane meadow plant
phenology and floral resources. *Ecosphere*, *8*(3), e01745.
https://doi.org/10.1002/ECS2.1745
Slatyer, R. A., Umbers, K. D. L., & Arnold, P. A. (2022). Ecological responses to variation in
seasonal snow cover. *Conservation Biology*, *36*(1), e13727.
https://doi.org/10.1111/cobi.13727
Sundqvist, M. K., Sanders, N. J., Dorrepaal, E., Lindén, E., Metcalfe, D. B., Newman, G. S.,
Olofsson, J., Wardle, D. A., & Classen, A. T. (2020). Responses of tundra plant
community carbon flux to experimental warming, dominant species removal and
elevation. *Functional Ecology*, *34*(7), 1497–1506. https://doi.org/10.1111/1365-

690   2435.13567

691 Suzuki, R. O. (2014). Combined effects of warming, snowmelt timing, and soil disturbance on
692   vegetative development in a grassland community. *Plant Ecology*, *215*(12), 1399–1408.
693   https://doi.org/10.1007/s11258-014-0396-x

694 Torchiano, M. (2020). *effsize: Efficient Effect Size Computation* (R package version 0.8.1).
695   https://CRAN.R-project.org/package=effsize

696 Vorkauf, M., Kahmen, A., Körner, C., & Hiltbrunner, E. (2021). Flowering phenology in alpine
697   grassland strongly responds to shifts in snowmelt but weakly to summer drought. *Alpine*
698   *Botany 2021 131:1*, *131*(1), 73–88. https://doi.org/10.1007/S00035-021-00252-Z

699 Wild, J., Kopecký, M., Macek, M., Šanda, M., Jankovec, J., & Haase, T. (2019). Climate at
700   ecologically relevant scales: A new temperature and soil moisture logger for long-term
701   microclimate measurement. *Agricultural and Forest Meteorology*, *268*, 40–47.
702   https://doi.org/10.1016/J.AGRFORMET.2018.12.018

703 Wipf, S., & Rixen, C. (2010). A review of snow manipulation experiments in Arctic and alpine
704   tundra ecosystems. *Polar Research*, *29*(1), Article 1.
705   https://doi.org/10.3402/polar.v29i1.6054

706 Yetgin, A. (2024). Exploring the dynamic nature of root plasticity and morphology in the face of
707   changing environments. *Ecological Frontiers*, *44*(1), 112–119.
708   https://doi.org/10.1016/j.chnaes.2023.07.008

709 Zhang, Q., Zhang, X., Lara, M. J., Li, Z., Xiao, J., Zhao, K., & Hu, T. (2023). Impacts of abiotic
710   and biotic factors on tundra productivity near Utqiaġvik, Alaska. *Environmental*
711   *Research Letters*, *18*(9), 094070. https://doi.org/10.1088/1748-9326/acf7d6

712 Zona, D., Lafleur, P. M., Hufkens, K., Bailey, B., Gioli, B., Burba, G., Goodrich, J. P., Liljedahl,
713   A. K., Euskirchen, E. S., Watts, J. D., Farina, M., Kimball, J. S., Heimann, M., Göckede,
714   M., Pallandt, M., Christensen, T. R., Mastepanov, M., López-Blanco, E., Jackowicz-
715   Korczynski, M., … Oechel, W. C. (2022). Earlier snowmelt may lead to late season
716   declines in plant productivity and carbon sequestration in Arctic tundra ecosystems.
717   *Scientific Reports 2022 12:1*, *12*(1), 1–10. https://doi.org/10.1038/s41598-022-07561-1