Title: CXCL16 mediates nociception and inflammation in murine post-traumatic osteoarthritis

Authors: Lindsey Lammlin[1], Stephen Redding[1], Alexander J. Knights[1], Aanya Mohan[1], Michael D. Newton[1], Rachel F. Bergman[1], Isabelle J. Smith[1], Scarlet C. Howser[1], Sofia Gonzalez-Nolde[1], Phillip M. Rzeczycki[1], Natalie S. Adamczyk[2], Rachel E. Miller[2], *Tristan Maerz[1,3,4]

Affiliations:

[1]Department of Orthopaedic Surgery, University of Michigan; Ann Arbor, Michigan, United States.

[2]Division of Rheumatology, Rush University Medical Center; Chicago, Illinois, United States.

[3]Department of Biomedical Engineering, University of Michigan; Ann Arbor, Michigan, United States.

[4]Division of Rheumatology, Department of Internal Medicine, University of Michigan; Ann Arbor, Michigan, United States.

Corresponding Author Info

*Tristan Maerz, PhD

Department of Orthopaedic Surgery, University of Michigan; Ann Arbor, Michigan, United States.

tmaerz@umich.edu

109 Zina Pitcher Place, Ann Arbor, MI 48109

One Sentence Summary:

CXCL16 regulates synovial inflammation and mediates joint nociception via CXCR6,

highlighting its potential as a therapeutic target for post-traumatic osteoarthritis.

Abstract:

This study investigates the role of the chemokine CXCL16 and its receptor, CXCR6, in post-traumatic osteoarthritis (PTOA) and joint nociception, highlighting the potential of targeting the CXCL16-CXCR6 axis for therapeutically managing joint inflammation and pain. Following joint injury in mice, the CXCL16-CXCR6 signaling axis is activated in synovium, driven by synovial fibroblasts and macrophages. Human OA synovium also exhibited increased CXCL16 and CXCR6 gene expression. CXCL16 stimulated a pro-inflammatory response in fibroblasts and macrophages, contrasting with an anti-inflammatory response observed in mesenchymal progenitor cells. In mice, repeated intra-articular CXCL16 injections induced histological synovitis and sex-dependent activation of inflammatory and fibrotic transcriptional programs in synovium. Repeated CXCL16 joint injections also induced knee hyperalgesia, which was mitigated by co-administration of the CXCR6 antagonist, ML339. A single intra-articular injection of CXCL16 induced acute knee hyperalgesia as early as 30 minutes post-injection, which was completely abrogated by ML339 co-treatment, suggesting direct CXCL16 binding to nociceptor-expressed CXCR6. In a murine PTOA model, systemic CXCR6 antagonism with ML339 alleviated knee hyperalgesia and altered circulating immune cell profiles. Direct stimulation of mouse dorsal root ganglion-derived nociceptive neurons with CXCL16 induced rapid calcium signaling, which was abolished by co-treatment with ML339. These findings establish CXCL16 as a regulator of joint inflammation and identifies the CXCL16-CXCR6 binding mechanism as key in mediating pain-related behaviors and nociceptor activation, offering a therapeutic target for PTOA-related inflammation and pain management.

Main text:

INTRODUCTION

The role of synovitis in promoting multiple pathological manifestations of osteoarthritis (OA), including pain and catabolic articular destruction, is now well-recognized (*1, 2*). In idiopathic OA, synovitis manifests as chronic low-grade tissue inflammation with intermittent flare-ups, and recent literature has demonstrated the existence of different inflammatory endotypes in both late-stage OA and rheumatoid arthritis (RA) (*3-5*). In the setting of post-traumatic osteoarthritis (PTOA), joint injury results in a high degree of acute synovial inflammation, which transitions to a chronic, low-grade nature akin to that observed in idiopathic OA.

A highly coordinated and tightly regulated set of factors orchestrate the local and systemic responses to tissue trauma. However, the immunological mechanisms underpinning joint injury-induced synovitis and its downstream consequences remain poorly described. Among the factors strongly induced following injury are the families of chemokines, secreted by tissue-resident stromal, endothelial, and immune cells. While some chemokines such as CCL2/MCP-1 (*6-10*) have been shown to be involved in OA/PTOA inflammation, the individual functions of most chemokines remain unclear in the context of joint injury and PTOA development. For instance, while CCL5/RANTES was shown to activate IL6 production in human synovial fibroblasts (*11*), indicating a potential deleterious pro-inflammatory effect, its genetic ablation did not alter disease severity in a surgical mouse OA model (*12*). Here, we identified and functionally characterized the chemokine CXCL16 as a synovial-derived factor selectively upregulated and secreted following joint injury. CXCL16 is expressed as a transmembrane protein, and it is cleaved from the cell surface by a disintegrin and metalloproteinase 10 (ADAM10) to act as a soluble ligand (*13*). As a transmembrane protein, CXCL16 is thought to act as an adhesion molecule for leukocytes (*14*) and can also act as a receptor for the soluble ligand form of CXCL16 (*15, 16*). CXCL16 also binds CXCR6, a G-protein coupled receptor and the only classical receptor described to interact with CXCL16 (*17, 18*). The CXCL16-CXCR6 axis has been associated with *in vivo* chemotaxis of macrophages, MSCs, and T cells in lung, liver, cardiac, renal, skin, and rheumatoid arthritic joint tissues (*19-24*). In the context of RA, CXCL16 has been shown to recruit mononuclear cells to synovial

tissue (*25*) and promote FLS proliferation (*26*). CXCL16 has also been shown to be involved in

the recruitment of endothelial progenitor cells, contributing to pathological angiogenesis

observed in RA (*27*). Ablation of CXCR6 ameliorated K/BxN serum-induced arthritis by

mitigating inflammatory cell infiltration, and fewer T cells and macrophages were found in joints

of K/BxN-challenged CXCR6[-/-] mice (*27*). CXCL16 monoclonal antibody treatment was also

shown to be effective in mitigating collagen induced arthritis in mice (*28*). Despite this evidence

from inflammatory and autoimmune-mediated arthritis, no prior study has described the role of

the CXCL16-CXCR6 axis following joint injury and during PTOA progression.

Chemokines are also highly relevant in the context of pain, and pain mechanisms in

OA/PTOA remain poorly described. Chemokines can either directly affect nociception as

agonists of nerve fiber-bound receptors, serving to sensitize nociceptive transmission

underpinning pain, or indirectly by inducing the upregulation of cytokines and neurotrophic

mediators that interact with neurons (*29, 30*). Very few chemokines have been characterized in

their involvement in joint nociception and OA/PTOA-related pain specifically. In a mouse model

of gouty arthritis, the CXCL5-CXCR2 axis was shown to have a role in pain and neurogenic

inflammation as mice with neuronal specific CXCR2 depletion exhibited less severe pain

related-behaviors and reduced production of calcitonin gene-related peptide and substance P

(*31*). Prior work has demonstrated that CCL2 is a key mediator of joint nociception by

interacting with nociceptor-bound CCR2 (*6*). Furthermore, CCL2[-/-] and CCR2[-/-] mice exhibit

delayed onset of pain-related behavior in a surgical PTOA model (*7, 8*). The CXCL16-CXCR6

axis, however, has yet to be described in the context of joint nociception.

The objective of this study was to assess the role of CXCL16 in joint injury-induced,

PTOA-associated inflammation and nociception. We demonstrate that the CXCL16-CXCR6 axis

is activated following traumatic joint injury, that CXCL16 induces pro-inflammatory signaling in

synovial fibroblasts and macrophages, and that CXCL16 is sufficient by itself to induce joint

pain and synovitis in mice *in vivo*. Using *in vitro* Ca^{2+} measurements in neuronal cells derived

from nociceptor-specific GCaMP6s calcium sensor mice and using *in vivo* acute hyperalgesia

experiments, we show that CXCL16 directly agonizes nociceptors and induces knee

hyperalgesia. Finally, employing a clinically-relevant pharmacological CXCR6 antagonist, we

demonstrate that CXCR6 inhibition mitigated the pro-algesic effects of CXCL16, abrogated

CXCL16-induced Ca^{2+} signaling by nociceptors, and ameliorated joint injury-induced knee

hyperalgesia.

RESULTS

Traumatic joint injury activates the CXCL16-CXCR6 axis in synovium

To understand the temporal expression patterns of synovial-derived chemokines and their

receptors following joint injury, we analyzed our bulk RNA-sequencing (RNA-seq) data of

whole mouse synovium (inclusive of Hoffa's fat pad) (*32*). We observed marked injury-induced

perturbation in members of the CCL and CXCL families (**Fig. 1A**), with corresponding changes

in the expression pattern of their respective receptors (**Fig. S1**). We next screened this list for

chemokines with zero or near-zero expression in Sham joints, further focusing on chemokines

with no prior PTOA-related studies. We identified CXCL16, which was upregulated 4.22- and

2.97-fold at 7d and 28d post-ACLR, respectively, with markedly lower P_{adj} values compared to

all other chemokines (**Fig. 1A**). qPCR corroborated the upregulation of *Cxcl16* at additional

timepoints post-ACLR in the synovium and meniscus (**Fig. S2A-B**). CXCL16 is expressed as a

transmembrane protein, in which state it can function as a scavenger receptor for lipoproteins

and bacteria (*33*) in addition to acting as a receptor for the soluble form of CXCL16 (*15, 16*).

Upon cleavage by ADAM10, CXCL16 is shed from the membrane and acts as a soluble

chemokine, shown to mediate leukocyte chemotaxis (*14*) as well as activate the

phosphatidylinositol-4,5-bisphosphate 3-kinase (PI3K)-Akt/PKB and extracellular signal-

regulated kinase mitogen-activated protein kinase (ERK/MAPK) pathways, and mobilize Ca^{2+}

via the G protein-coupled receptor CXCR6 (*34*). Mining of our previously published bulk RNA-

seq data of whole synovium also showed that *Cxcr6* was upregulated 3.30-fold at 7d and 2.87-

fold at 28d post-ACLR (**Fig. S1A**). By qPCR we also observed upregulation of *Adam10* in the

synovium, but not the meniscus following joint injury (**Fig. S2A-B**), demonstrating upregulation

of the entire CXCL16-CXCR6-ADAM10 axis following joint injury.

In a meta-analysis of transcriptomic data of human OA synovium, we also observed

marked alterations in the expression patterns of CCL and CXCL chemokines and their receptors,

relative to healthy synovium (**Fig. S1B-C**). *CXCL16*, *CXCR6*, and *ADAM10* were significantly

upregulated in OA synovium compared to healthy synovium (**Fig. 1B**). Notably, CXCL16 had

the lowest, i.e. most significant composite P_{adj} value compared to all other CCL and CXCL

chemokines (**Fig. S1B**), demonstrating that CXCL16 upregulation is also observed in human OA

and not confined to the acute post-injury period. Uninjured mouse joints and healthy human OA

synovium exhibited very low CXCL16 expression. Concordantly, while CXCL16 protein was

undetectable in the synovial fluid of uninjured joints, the synovial fluid of injured ACLR joints

was enriched in CXCL16 protein (**Fig. 1C**).

To identify the synovial cells expressing CXCL16 (intracellular) and CXCR6 (surface)

proteins, we performed flow cytometry of ACLR synovium compared to healthy contralateral

synovium at both early (3d) and intermediate (14d) timepoints following injury. We observed

large but statistically non-significant increases in the total number of CXCL16+ and CXCR6+

cells in ACLR synovia compared to uninjured synovia (**Fig. 1D**), but only CXCR6+ cells

increased in their proportion to all live cells at 3d post-ACLR (**Fig. 1E**). This is likely due to the

widely reported dramatic increase in total synovial cells after joint injury, corroborated by our

results showing all major synovial cell types elevated at 3d and 14d post-ACLR compared to

contralateral tissue (**Fig. S2C-G**). Intracellular CXCL16 was found to be expressed most

strongly by CD31(-) CD45(-) stromal cells, i.e. synovial fibroblasts, as well as by CD31(+)

CD45(-) endothelial cells, and by CD45(+) immune cells (**Fig. 1F**). Within CD45(+) immune

cells, we observed CXCL16 expression in CD11b(+) CD64(+) macrophages, CD11b(-) CD3(+)

T cells, and other immune subsets (**Fig. 1G**). CXCR6 was predominantly expressed by immune

cells with lower levels of expression in stromal and endothelial cells (**Fig. 1F**). The CXCR6+

immune cells were comprised of ~80-90% macrophages (**Fig. 1G**). Mining of our previously

published murine synovial single-cell RNA sequencing data (*35*) shows that *Cxcl16* transcript is

expressed by fibroblasts and most highly in macrophage and dendritic cells, whereas *Cxcr6*

transcripts are confined to T-cells, and *Adam10* is expressed ubiquitously across synovial cell

populations (**Fig. 1H**).

Taken together, these findings demonstrate that CXCL16 is a disease-specific secreted

factor, expressed by synovial fibroblasts and macrophages, and enriched in the synovium of both

human OA patients and injured mouse joints.

IL1β and TNFα induce CXCL16 expression in joint relevant cell types

Given the observation of injury-induced upregulation of CXCL16 by multiple synovial

cell types, we sought to identify upstream activators of CXCL16 expression (**Fig. 2A**). We found

that IL-1β and TNFα, two key cytokines strongly upregulated in synovium following ACLR (*32,*

36), activated *Cxcl16* and *Cxcr6* transcript upregulation in both primary murine FLS and bone

marrow-derived macrophages (**Fig. 2B-C**). IL-1β and TNFα also caused upregulation of *Adam10*

in FLS (**Fig. 2B**), but downregulation of *Adam10* in BMDM (**Fig. 2C**). We confirmed the

increase of secreted CXCL16 protein in the conditioned medium of IL-1β- and TNFα-treated

FLS, compared to vehicle-treated cells (**Fig. 2B**). We also found IL-1β and TNFα to cause

upregulation of *Cxcr6* in an immortalized chondrogenic cell line, ATDC5s - a culture model of

articular chondrocytes, whereas these cytokines induced downregulation of *Adam10* while

Cxcl16 remained unchanged (**Fig. 2D**). These results demonstrate that the CXCL16-CXCR6 axis

is activated by IL1β and TNFα.

CXCL16 induces cell type-dependent pro- or anti-inflammatory gene expression

We next wanted to assess the potential functional role of CXCL16 in joint-relevant cell

types. We treated primary murine synovial fibroblasts, bone marrow-derived macrophages, and

bone marrow-derived mesenchymal progenitor cells (BMPCs) with increasing doses of

recombinant CXCL16 protein. We found CXCL16 to induce a multi-factorial pro-inflammatory

response in FLS, indicated by both an increase in pro-inflammatory gene expression (*Il1b, Il6*)

and a decrease in anti-inflammatory gene expression (*Il10*) (**Fig. 3A**). BMDMs treated with

CXCL16 similarly had a similar double-edge pro-inflammatory response as summarized by the

increase in macrophage polarization score upon treatment with CXCL16 (**Fig. 3B**). This

macrophage score is an aggregate assessment of macrophage phenotype which is calculated by

the ratio of pro-inflammatory (*Il1b*, *Il6*, *Mmp1b*) to anti-inflammatory (*Cd206*, *Il4*, *Il10*) gene

expression (**Fig. S3A-B**). Interestingly, CXCL16 elicited a cell-type specific response as BMPCs

treated with CXCL16 upregulated their expression of anti-inflammatory transcripts *Il10* and

Il1ra, and the pro-inflammatory transcript *Il1b*, but *Il6* was unchanged (**Fig. 3C**).

In addition to promoting cell-type specific inflammatory responses, CXCL16 contributed

to its own feed forward signaling activation in FLS, BMDM, and BMPCs. CXCL16-treated FLS

upregulated *Cxcl16*, *Cxcr6*, and *Adam10* expression (**Fig. 3A**). BMDMs exhibited similar

responses, with increases in *Cxcl16*, *Cxcr6*, and *Adam10* expression after treatment with

recombinant CXCL16 (**Fig. 3B**). BMPCs exhibited strong upregulation of *Cxcr6* with a less

dramatic induction of *Cxcl16* and *Adam10* (**Fig. 3C**). In an *in vitro* migration assay, BMDMs and

BMPCs also exhibited a dose-dependent chemotactic response to CXCL16 (**Fig. 3D**),

corroborating prior studies on the chemotactic effects of CXCL16 (*19-24*). Taken together, these

in vitro findings demonstrate that CXCL16 induces pro-inflammatory effects in primary FLS and

BMDMs but anti-inflammatory effects in BMPCs.

Repeated intra-articular injections of CXCL16 induces knee hyperalgesia via CXCR6 and

synovitis in healthy murine joints

On the basis of CXCL16 activating pro-inflammatory gene expression in synovial

fibroblasts and BMDMs, we next wanted to assess whether CXCL16 is sufficient by itself to

induce synovitis and joint disease *in vivo*. We injected either 5 µL of vehicle or 0.25 ng (low

dose) or 25 ng (high dose) of recombinant murine CXCL16 into healthy, adult C57BL/6 knee

joints for five consecutive days and performed knee hyperalgesia and histological assessments.

Low and high dose CXCL16-injected limbs, but not vehicle-injected limbs, exhibited reduced

knee withdrawal thresholds at 7d following the first injection, indicative of hyperalgesia (**Fig.**

4A). High dose CXCL16-injected limbs had a sustained reduction in knee withdrawal threshold

up to 14d following the first injection, but this effect was not observed in low dose or vehicle-

injected limbs (**Fig. 4A**). Histological evaluation demonstrated no change in the total post-

traumatic osteoarthritis score, however there were significantly increased total synovitis scores in

CXCL16-injected joints, driven by increased synovial lining hyperplasia and synovial fibrosis

(**Fig. 4B-C**).

To assess the molecular programs that underpin these pathological manifestations, we

performed bulk RNA-seq of whole synovium from a separate cohort of mice treated with the

identical vehicle or high-dose CXCL16 injection regimen as above. In this cohort, CXCL16-

injected joints again exhibited reduced knee withdrawal thresholds 7d post-first injection (**Fig.**

4D). By RNA-seq at the same timepoint, we observed a highly sex-dependent perturbation of the

synovial transcriptome due to CXCL16 administration (**Fig 4E**). Whereas female synovium

exhibited near-absent CXCL16-induced differentially expressed genes (DEGs), male synovia

upregulated cytokines, chemokines, and cytokine/chemokine receptors (e.g. *Ccl2, Ccl7, Cxcl5,*

Il1rl1, Ccr5), genes of macrophage-relevant transcription factors and genes related to

macrophage polarization (e.g. *Junb, Fos, Mt1, Arg1, Fgr*), and genes related to fibroblast

activation and ECM turnover (*Pdpn, Mmp3, Itga5, Lum, Cspg4, Hmcn2*) (**Fig. 4E, Data file S2**).

GSEA pathway analysis demonstrated that synovia from male CXCL16-treated joints had

enriched pathways related to collagen production, immune/inflammatory response, and

neuroinflammation (**Fig. 4F**). Taken together, these findings demonstrate that CXCL16 is

sufficient by itself to induce arthritis-relevant disease manifestations in healthy joints *in vivo*,

characterized by increased hyperalgesia, histological synovitis, and the induction of a synovial

transcriptional program indicative of inflammation, lymphocyte and macrophage activation,

ECM production, and fibroblast proliferation.

To interrogate the receptor mechanism by which CXCL16 induced pain, we co-

administered a CXCR6-specific antagonist, ML339, in the same repeated injection study design.

Male mice were administered five consecutive days of intra-articular vehicle, CXCL16 (200 ng),

or CXCL16 (200 ng) + ML339 (1 mM) injections. CXCL16-injected joints again had reduced

knee withdrawal thresholds and knee withdrawal threshold ratios as compared to vehicle injected

limbs. This hyperalgesia-inducing effect was partially ablated by co-administration of ML339, as

joints injected with combinatorial CXCL16 + ML339 had significantly higher knee withdrawal

threshold and knee withdrawal ratio compared to joints injected with CXCL16 only (**Fig. 4G**).

CXCL16 induces acute knee hyperalgesia via CXCR6

Joint pain is a complex physiological process orchestrated by multiple cell types, ligands,

and receptors. A given mediator may agonize pain pathways via direct signaling, i.e. binding to

joint nociceptor-bound receptors to directly alter nociception, or via indirect means, i.e. inducing

the expression of other factors that directly bind nociceptors. Having demonstrated that CXCL16

is sufficient to induce knee hyperalgesia in a repeated-injection model, we sought to assess

whether this at least partially occurs via direct interactions with joint-innervating nociceptors.

We performed an acute hyperalgesia experiment in which recombinant CXCL16 (25 ng), or

vehicle (sterile PBS) was injected into joints followed by knee withdrawal threshold

measurements at acute timepoints post-injection (30 mins – 4 hrs) (**Fig. 5A**). Vehicle injection

caused minor decreases in knee withdrawal threshold compared to baseline, however, CXCL16

significantly reduced withdrawal thresholds compared to vehicle by 30 mins post-injection in

male mice, with a sustained effect until 7 days (**Fig. 5B-C, Fig. S4A-B**). Strikingly, despite

female mice also exhibiting increased hyperalgesia after repeated injections in the prior

experiment (**Fig. 4A, D**), female mice had absent hyperalgesia responses relative to vehicle in

the acute experiment (**Fig. 5B-C**, **Fig. S4A-B**). The rapid responses observed in this experiment

provide evidence of a direct nociceptive effect, i.e. direct interactions between CXCL16 and

nociceptor-bound receptors (*37, 38*).

We next wanted to ascertain whether the induction of acute hyperalgesia by CXCL16 is

mediated by CXCR6. We conducted another acute hyperalgesia experiment in only male mice

with a higher dose of CXCL16 (250 ng) and intra-articular co-administration of either vehicle or

the CXCR6 antagonist ML339 (1mM) (**Fig. 5D**). Consistent with the prior experiment, CXCL16

induced acute hyperalgesia that persisted until 24 hrs post-injection, and co-administration of

ML339 completely ablated this response (**Fig. 5E-F, Fig. S4C-D**). These findings demonstrate

that CXCL16-induced hyperalgesia is mediated via CXCR6 binding.

Systemic CXCR6 antagonism blocks joint injury-induced hyperalgesia

As joint injury induces the upregulation of multiple cytokines, chemokines, and

neurotrophic mediators (**Fig. 1A**) (*32*) that orchestrate synovitis and hyperalgesia, we next

wanted to assess whether CXCR6 antagonism mitigates injury-induced hyperalgesia. Male mice

underwent ACLR and received 5x intra-peritoneal injections of ML339 (5 mg/kg) or vehicle

followed by assessments of knee hyperalgesia and circulating immune cell populations (**Fig.**

5G). At the study endpoint, 7d post-ACLR, we also conducted an acute hyperalgesia experiment

to measure any potential acute analgesic effects due to systemic injection. Whereas pre-injection

withdrawal thresholds were comparable between treatment groups, ML339-treated mice had

significantly higher withdrawal threshold ratio and absolute withdrawal threshold 2 hrs post-

injection. (**Fig. 5H-I, Fig. S4E-F**). Complete blood counts further revealed that ML339-treated

mice had higher circulating total white blood cell and neutrophil counts (**Fig. 5J**). These findings

indicate that systemically administered ML339 acutely mitigates ACLR-induced knee

hyperalgesia.

CXCL16 induces rapid Ca^{2+} influx in dorsal root ganglia-derived nociceptive neurons via

CXCR6

Given our observation that CXCR6 antagonism mitigated CXCL16-induced acute

hyperalgesia as well as joint injury-induced hyperalgesia, we last sought to determine whether

CXCL16 directly agonizes nociceptive neurons and whether this requires CXCR6. We isolated

and cultured primary neurons derived from the dorsal root ganglia (DRG) of $Na_V1.8$-Cre;

GCaMP6s$^{fl/+}$ mice, which express the fluorescent calcium indicator, GCaMP6s, in nociceptive

neurons, marked by expression of the voltage-gated sodium channel 1.8 ($Na_V1.8$) (*39*). In a time

series experiment, cultured neurons were first treated with a negative control (vehicle only),

followed by a single dose of 50 ng/ml CXCL16, with KCl used as a positive control (**Fig. 6A**).

This was conducted in the presence or absence of 180 µM ML339 to antagonize CXCR6, and

fluorescent signal was monitored and quantified as the change over baseline. CXCL16, but not

vehicle, induced fluorescent responses, indicative of rapid neuronal Ca^{2+} signaling (**Fig. 6B-D**).

Co-treatment with the CXCR6 antagonist ML339 completely ablated CXCL16-induced

fluorescent responses ($P<0.0001$), with no effect on positive control responses to KCl (**Fig. 6B-**

D). Taken together, these findings confirm that CXCL16 directly agonizes nociceptive neurons

by inducing Ca^{2+} signaling, and this effect is abolished by pharmacological inhibition of the

CXCR6 receptor.

DISCUSSION:

This study elucidated the role of the CXCL16-CXCR6 axis in the context of post-

traumatic osteoarthritis (PTOA), with a focus on its contributions to inflammation and

nociception. Our findings demonstrate that joint injury rapidly activates gene and protein

expression of CXCL16 and CXCR6, and that CXCL16 orchestrates synovial inflammation, pain-

related behaviors and nociceptive signaling during PTOA progression. Using a translationally-

relevant small-molecule antagonist against CXCR6, we confirmed that CXCR6 mediates

CXCL16-induced joint hyperalgesia *in vivo* as well as CXCL16-induced Ca^{2+} signaling in DRG-

derived nociceptive neurons *in vitro*.

To our knowledge, this is the first study to demonstrate the role of CXCL16 and its only

known classical receptor, CXCR6, in peripheral nociception. Recently, Wang, et. al.,

demonstrated that CXCL16 expression and CXCL16 mRNA N^6-methyladenosine modification

was increased in rat sympathetic ganglia in a model of spared nerve injury (SNI)-induced

neuropathic pain (*40*). They further showed that intra-DRG administration of CXCL16 induced

mechanical allodynia in naïve rats while ablation of CXCL16 in sympathetic ganglia mitigated

SNI-induced mechanical allodynia, establishing a role for CXCL16 in mediating neuropathic

pain (*40*). Wang, et. al., also observed CXCR6 protein expression in rat DRG neurons (*40*),

which suggests that CXCL16 can directly bind to DRG neurons via CXCR6. These findings

support our premise that CXCL16 can induce knee hyperalgesia through direct binding to

CXCR6+ nociceptive neurons. Of note, Wang, et. al did not empirically test whether CXCL16

operates via CXCR6, and our study did not measure CXCR6 expression by joint-resident

nociceptors, which is an important future direction. However, using a small-molecule CXCR6

antagonist, we confirmed the receptor mechanism of CXCL16-induced effects in cultured DRG

neurons *in vitro* and in CXCL16-induced joint hyperalgesia *in vivo,* providing the first

mechanistic evidence for a CXCL16-CXCR6 binding interaction that promotes peripheral

nociceptive signaling and pain-related behaviors.

CXCL16 has been shown to induce calcium mobilization in a CXCR6-expressing murine

pre-B lymphoma cell line (*17*) and previous studies in rat and mouse DRGs have established the

direct activation of Ca^{2+} mobilization by other chemokines, including CCL2/MCP-1,

CCL5/RANTES, CXCL1/fractalkine (*30*), and CXCL5 (*31*). However, to date, CXCL16 has not

been tested in this context. Our studies advance the understanding of the role of CXCL16-

CXCR6 in calcium mobilization, as we demonstrated that CXCL16 can directly stimulate

calcium signaling in mouse DRG neurons via CXCR6. Although a direct connection between

CXCL16-CXCR6 and pain-related behaviors has not previously been established, CXCL16 has

been connected to the nervous system through roles in neurotransmitter release and neuronal-

glial cellular crosstalk (*41-43*). CXCL16 and CXCR6 have been shown to be expressed by

astrocytes, microglia, and neurons in hippocampal cultures, indicating a potential for CXCL16-

CXCR6 signaling (*42*). Di Castro, et. al, observed that CXCL16 promotes spontaneous GABA

release in pyramidal neurons from the hippocampal CA1 area (*41*). Wang, et al. also showed that

CXCL16 worked synergistically with norepinephrine to increase DRG hyperexcitability and

exacerbate neuropathic pain (*40*). Our finding that CXCL16 induces acute knee hyperalgesia, as

early as 30 minutes post-injection, suggests that CXCL16 is interacting directly with intra-

articular nociceptors, but further studies are required to determine whether this hypothesized

activation is due to modulation of Ca^{2+} influx, neurotransmitter release, or some other

mechanism.

CXCL16 and CXCR6 have been established as inflammatory markers across multiple

diseases including high-grade prostate cancer, which is considered an inflammation associated

cancer (*44*). Lehrke, et. al., also suggest a pro-inflammatory role for CXCL16 in inflammatory

bowel disease as CXCL16 was highly enriched in patients with Crohn's disease or ulcerative

colitis and the elevated CXCL16 levels correlated with level of C-reactive protein (*45*). Our

studies also demonstrate a pro-inflammatory role for CXCL16, showing that CXCL16 elicits a

pro-inflammatory response in synovium *in vivo* and in fibroblasts and macrophages *in vitro*.

However, these responses were not tested alongside the CXCR6 antagonist ML339, and thus we

cannot conclude whether pro-inflammatory effects were mediated by CXCR6. Further, we

demonstrate mitigated CXCL16-induced pain following co-administration with ML339 in a

repeated injection experimental design, but we did not evaluate whether co-administration with

ML339 also mitigated the pro-inflammatory activation of the synovium that was observed in

males. Future studies aim to address these limitations. In a cardiotoxin injection model of muscle

injury, CXCL16 knockout mice had decreased macrophage infiltration but increased neutrophil

infiltration to injured muscle as compared to controls (*46*). CXCL16 knockout mice also had

decreased infiltration of $Ly6C^{hi}$ monocytes and macrophages in the infarcted heart following

myocardial infarction (*47*). These studies and our data support the function of CXCL16 as a

macrophage chemoattractant across multiple tissue types, and our findings extend this

knowledge by demonstrating its pro-inflammatory effects in synovial fibroblasts.

To the author's knowledge, while the CXCR6 antagonist ML339 has been validated *in*

vitro (*48*), *in vivo* use has been limited to a few cancer studies (*49, 50*) and no study to date has

used ML339 *in vivo* under any musculoskeletal health context. While systemic administration of

the CXCR6 antagonist acutely ablated joint injury-induced hyperalgesia, it did not mitigate

overall hyperalgesia development via disease modification at 7d post-ACLR. These findings

indicate that the development of joint hyperalgesia following injury is not solely dependent upon

CXCR6, but that CXCL16-CXCR6 at least partially mediate acute nociception. As CXCL16 is

the only described ligand for CXCR6, we conclude that the injury-induced upregulation of

CXCL16 in the joint, which we confirmed at the protein level in synovial fluid, is partially

responsible for injury-induced hyperalgesia, which can be transiently blocked by targeting the

CXCR6 receptor.

Our findings should be interpreted in light of several limitations. The sex-specific

differences in the synovial transcriptome and acute hyperalgesia observed in response to intra-

articular CXCL16 injection highlight an area needing further investigation. Such studies should

address whether this is related to a dose dependency. This will allow future refinement of

treatment approaches that account for sex as a biological variable. Furthermore, the efficiency of

in vivo CXCR6 antagonism by ML339 is unclear, and residual CXCR6-mediated signaling by

incomplete receptor inhibition may compensate certain responses, which should be addressed by

studies employing genetic knockouts of CXCL16 and CXCR6. While our study provides a

comprehensive examination of the role of CXCL16 within murine models, there is a need to

validate these findings in larger animal models, and human synovial cells and DRGs. Moreover,

our study focused mainly on the acute phases of PTOA, with limited assessments of the long-

term effects of CXCL16 activation or CXCR6 inhibition. Chronic pain and progressive joint

degeneration are characteristic of PTOA, and thus longer-term studies are needed to understand

whether the analgesic effects of CXCR6 inhibition are sustained throughout PTOA progression.

In summary, our study unveils the CXCL16-CXCR6 axis as a novel mediator of joint

inflammation and nociception in PTOA while also establishing pharmacological CXCR6

antagonism as a potential therapeutic strategy to treat joint trauma-related inflammation and

pain.

MATERIALS AND METHODS:

Study design

The objectives of this study were to: 1) Evaluate the CXCL16 signaling axis in the

context of PTOA; 2) establish key mediators and downstream effectors of CXCL16; 3) identify

the sufficiency of CXCL16 in driving synovial inflammation and joint nociception, and finally,

4) establish a receptor mechanism through which CXCL16 exerts its effects. To explore how

CXCL16 expression is altered following murine joint injury and in human OA synovium, we

mined bulk and single cell RNAseq datasets and RNA microarrays from mouse and human

synovium. Flow cytometry was used to identify CXCL16+ and CXCR6+ cells in healthy and

injured mouse synovium, and ELISA was used to measure CXCL16 protein in synovial fluid. *In*

vitro studies examined the effects of pro-inflammatory cytokines and CXCL16 treatment in

murine joint-relevant cell types. The sufficiency of CXCL16 to induce synovial inflammation

and joint nociception *in vivo* was evaluated by repeated joint injections with recombinant

CXCL16 in naïve mice, followed by knee hyperalgesia testing, histological evaluation, and bulk

RNA sequencing of synovium. The receptor mechanism by which repeated intra-articular dosing

of CXCL16 induces knee hyperalgesia was assessed by co-administration of CXCL16 and a

CXCR6 antagonist, ML339. To demonstrate whether CXCL16 was directly agonizing

nociceptors to induce pain, mice were administered a single intra-articular injection of CXCL16

followed by hyperalgesia measurements as early as 30 mins post-injection. This was repeated

with co-administration of ML339 to establish CXCR6 as the receptor mediating this effect. The

role of CXCL16 in driving joint injury-induced pain was tested by administering multiple intra-

peritoneal doses of ML339 following noninvasive joint injury in mice and performing knee

hyperalgesia assessments longitudinally. Finally, to assess whether CXCL16 can directly activate

calcium signaling in nociceptive neurons, we performed *in vitro* calcium imaging of primary

DRG-derived neurons from $Na_V1.8$-Cre; $GCaMP6s^{fl/+}$ mice following treatment with CXCL16,

with or without CXCR6 antagonist co-treatment. Mice were randomly assigned to groups within

an experiment. A blinded operator performed all knee hyperalgesia testing and histopathologic

scoring. Sample sizes are detailed in the figure caption and statistical analyses are described in

the *Data analysis and statistics* methods section.

Animals and noninvasive ACL rupture

All animal studies were approved by the institutional animal care and use committee.

Male and female C57BL/6 mice aged 12-15 weeks (Jackson Laboratories) were used throughout.

To induce joint injury and PTOA, we employed a noninvasive mechanical loading-based anterior

cruciate ligament rupture (ACLR) model (*32, 36*) (**Suppl. Materials and Methods**). Limbs from

either Sham mice (no loading, with anesthesia and analgesia only) or unloaded contralateral

limbs from injured mice served as controls. Naïve male and female $Na_V1.8$-Cre; $GCaMP6s^{fl/+}$

mice (*39*) older than 12 weeks were used to obtain primary DRG cultures for *in vitro* Ca^{2+}

imaging experiments.

Bulk RNA sequencing, single-cell RNA-sequencing, RNA microarray published data mining

Existing, published bulk RNAseq data generated by our group (*32*) was re-analyzed to

assess injury-induced gene expression of all CCL and CXCL family chemokines and their

receptors in Sham, ACLR 7d, and ACLR 28d male and female synovia (**Suppl. Materials and**

Methods).

To assess expression patterns of Cxcl16, Cxcr6, and Adam10 in synovium, existing

published scRNAseq data from mouse and human OA synovium was analyzed using R. Mouse

scRNAseq previously published in our lab (*35*) was mined to examine synovial cell types

expressing *Cxcl16*, *Cxcr6*, and *Adam10*. Feature plots showing the expression of the genes of

interest across the all cells object, which contains integrated Sham, 7d ACLR, and 28d ACLR

synovial cells were generated using the *Shiny* R package.

A meta-analysis of human OA synovium bulk RNAseq and RNA microarray datasets

was performed using weighted Limma-Voom. Data from these is given as $\log_2$(fold change) and

$-\log 10(P_{adj})$. Unadjusted *P*-values from individual studies were combined across datasets into a

composite *P*-value using the weighted *Z*-test, with a BH (e.g. FDR) correction for multiple

comparisons applied following combination. This test is directional, so a positive change in one

study and negative change in another cancel out rather than combining. Composite data is

presented only as a signed $-\log_{10}(P_{adj})$, since $\log_2$(fold change) cannot be validly combined

between RNAseq and RNA microarray datasets. For genes which were missing from some

microarrays, only datasets which contained these genes were included in the composite *P*-value

calculation.

Synovial fluid collection and protein quantification

Synovial fluid was collected from joints as previously described (*36, 51*). In brief,

following euthanasia, a small, consistently-sized alginate sponge was inserted into the joint space

through a small incision made in the suprapatellar capsule. Synovial fluid-soaked sponges were

flash frozen and stored at -80C. Following thawing, sponges were digested via alginate lyase,

and digestates were assayed for CXCL16 content by ELISA (Invitrogen: EMCXCL16). Total

protein was quantified in the same samples via Pierce™ Bradford Plus Protein Assay Kit

(ThermoFisher Scientific, Cat. #23236). CXCL16 concentrations were normalized to total

protein.

Flow cytometry

Whole synovium, inclusive of Hoffa's fat pad, was dissected and enzymatically digested

to yield single-cell suspensions using Liberase TM, Collagenase IV, and DNaseI (*35, 36*). Each

sample consisted of one male and one female synovium pooled at the time of dissection (n=3

ACLR, n=3 contralateral). After digestion, single cell suspensions were stained (**Suppl.**

Materials and Methods). Flow cytometry was performed on a BD LSRFortessa and analysis

was performed in FlowJo v10 software (BD/TreeStar).

Tissue gene expression via RT-qPCR

Dissected joint tissue were homogenized in TRIzol, RNA was isolated, and qPCR was

performed to evaluate the expression of target genes (**Suppl. Materials and Methods**). Data was

normalized to housekeeping genes and reported further normalized to experimental controls

using the $2^{-\Delta\Delta CT}$ method.

Primary cell isolation and in vitro treatments

Primary fibroblast-like synoviocytes (FLS) from knee synovia, bone marrow-derived

macrophages (BMDMs), and bone marrow-derived mesenchymal progenitor cells (BMPCs)

were isolated from naïve adult male and female C57BL/6 mice, as previously described (*35*)

(**Suppl. Materials and Methods).**

To ascertain upstream activators of *Cxcl16* and related gene expression, FLS, BMDMs,

and BMPCs were treated with PBS (control), TNF-α (10 ng/mL), or IL-1β (10 ng/mL) for 48

hours (FLS, BMPCs) or 8 hours (BMDMs) and then harvested and processed for RT-qPCR.

Conditioned media from FLS was also collected for CXCL16 protein quantification via ELISA.

To assess the potential functional role of CXCL16, FLS, BMDMs, and BMPCs were treated with

PBS (control) or CXCL16 (10ng/mL or 50 ng/mL) for 48 hours (FLS, BMPCs) or 8 hours

(BMDMs) and then harvested and processed for RT-qPCR. All *in vitro* experiments were

conducted in n=4-8 biological replicates, i.e. cell isolations from unique mice (**Suppl. Materials**

and Methods).

ATDC5 cell culture

Following differentiation (**Suppl. Materials and Methods**), chondrogenic ATDC5s were

treated with PBS (control), TNF-α (10 ng/mL), or IL-1β (10 ng/mL) for 48 hours, followed by

gene expression studies. Briefly, ATDC5s were lysed with Trizol and frozen at -80°C until RNA

isolation and gene expression studies by RT-qPCR. All experiments were performed in n=4

replicates.

 In vitro chemotaxis of BMDMs and BMPCs

494   BMDMs and BMPCs were separately seeded into transwell insert membranes and

495   combined with a transwell chamber containing either media with vehicle, a positive control (LPS

496   for BMDMs, 5% FBS for BMPCs), or recombinant mouse CXCL16 (10 ng/mL or 50 ng/mL).

497   Cells were allowed to migrate for three days prior to quantification of migrated cell density

498   (**Suppl. Materials and Methods**).

Intraarticular and systemic treatments

Joint injections were administered to the knees of naïve C57Bl/6 mice. All injections

were performed using 33G needles and microsyringes (Hamilton Company). To assess the

disease-promoting effects of CXCL16 in healthy joints, CXCL16 (R&D Systems, Cat. #503-CX)

(0.25 ng or 25 ng in 4 µL of sterile PBS) was injected for five consecutive days into one knee

joint, and vehicle was injected in the contralateral joint (n=3 male, n=3 female per dose). Knee

hyperalgesia testing occurred at 7d and 14d post-first injection. At 14d post-first injection

hindlimbs were harvested for histological evaluation of PTOA and synovitis severity. A separate

group of mice (n=3 male, n= 3 female) underwent a similar regiment of 5 consecutive intra-

articular injections of CXCL16 into one knee joint (25 ng in 4 µL of sterile PBS with .1% BSA)

and vehicle (sterile PBS with .1% BSA) into the contralateral. A baseline measurement of knee

hyperalgesia was taken pre-injection then knee hyperalgesia and synovial harvests for bulk RNA

sequencing occurred 7d post-first injection. In another cohort, mice received 5 consecutive days

of bilateral intra-articular injections of either vehicle (sterile PBS with 0.69% DMSO), CXCL16

(200 ng in 5 µL sterile PBS with 0.69% DMSO), or CXCL16 co-administered with CXCR6

antagonist, ML339 (MedChemExpress, Cat. #HY-122197) (200 ng CXCL16 in 5 µL sterile PBS

with 1mM ML339 and 0.69% DMSO). Knee hyperalgesia was performed at baseline and 7d

post-first injection (n=6 males/group).

For the first acute hyperalgesia experiment in naïve mice, mice received either a single

CXCL16 (25 ng in 4 µL of sterile PBS with 0.1% BSA) or vehicle (sterile PBS with 0.1% BSA)

injection into a single joint (n=3 male, n=3 female). Knee hyperalgesia occurred at baseline and

30 min, 2 hr, 4 hr, 24 hr, and 7d post-injection (**Fig. 5A**). Mice were divided into two groups for

the second acute hyperalgesia experiment in naïve mice (n=9 males/group). One group received

an intra-articular injection of CXCL16 (250 ng in 6 µL sterile PBS containing 0.08% BSA and

0.69% DMSO) in one knee and vehicle (sterile PBS containing 0.08% BSA, and 0.69% DMSO)

in the contralateral. The other group received an intra-articular injection of CXCL16 co-

administered with CXCR6 antagonist, ML339, (250 ng CXCL16 in 6 µL sterile PBS containing

1mM ML339, 0.08% BSA, and 0.69% DMSO) in one knee and vehicle (sterile PBS containing

0.08% BSA, and 0.69% DMSO) in the contralateral. Knee hyperalgesia occurred at baseline and

30 min, 2 hr, 4 hr, and 24 hr (**Fig. 5D**). To assess whether ML339 mitigates ACLR-induced

effects, injured mice (n=9 males/group) received repeated intraperitoneal injections of either

vehicle (corn oil containing 1% DMSO) or ML339 (5 mg/kg of 1.45 mM ML339 in corn oil

containing 1% DMSO). Knee hyperalgesia occurred at baseline, 7d post-ACLR prior to

injection, and 7d post-ACLR at 2 hrs post-injection (**Fig. 5G**). All mice were randomly assigned

to treatment groups within each experiment.

Knee hyperalgesia testing

Knee hyperalgesia testing was performed by a blinded operator (LL) using a Randall-

Selitto device as described previously (*32, 36*). Briefly, pressure was applied to the medial aspect

of the knee and the knee withdrawal threshold was recorded when the mouse gave a response – a

struggle or vocalization. Three measurements were recorded for each limb and averaged together

to obtain the knee withdrawal threshold per limb.

Histological analysis

Whole hindlimbs were processed, stained with Safranin O/Fast Green (SafO), imaged

(**Suppl. Materials and Methods**), and qualitatively graded for OA and synovitis severity using

updated and extended versions (**Table S3-S4**) of an established grading scheme (*52, 53*). All

grading was performed in a blinded fashion (LL). Scores were averaged across all slides to

calculate a limb average, and limbs were averaged to calculate group scores.

Bulk RNA-sequencing and bioinformatic analysis

RNA was isolated from Trizol homogenates of whole synovium, inclusive of Hoffa's fat

pad, then sequenced and quality control was performed (**Data file S1**). All Bulk RNA-

sequencing data analysis including quality control, gene expression analysis, and pathway

analysis was performed in RStudio (R v4.4.0, RStudio v2024.04.1). Differential gene expression

analysis was performed using DESeq2 (*54*). Pathway analysis was performed using the

clusterProfiler (55) implementation of fast gene set enrichment analysis (GSEA), using the GO:

Biological Processes gene ontology terms list (**Suppl. Materials and Methods**).

Complete blood counts

Complete blood count (CBC) analysis was performed using 50-100 µL of blood, obtained

via cardiac puncture at the time of harvest.

Primary DRG neuron culture and Ca^{2+} imaging

Primary DRG cultures were generated from Na$_V$1.8-Cre; GCaMP6s$^{fl/+}$ mice as previously

described (*56*). These mice express the fluorescent calcium indicator, GCaMP6s, in nociceptive

neurons, marked by expression of the voltage-gated sodium channel 1.8 (Na$_V$1.8). L3-L5 DRGs

were collected and pooled from 12 wk-old mice, and male and female cultures were generated

separately. Four DRGs were used for every coverslip plated. Cultures were maintained at 37□,

treated with CXCL16 with or without pre-treatment with a CXCR6 antagonist, ML339 (180

µM), and fluorescent imaging was performed with Zeiss Zen software to measure Ca^{2+} signaling

(n=2-5 well replicates per sex/treatment) (**Suppl. Materials and Methods**). KCl was used as a

positive control.

Fluorescent imaging of the calcium sensor was performed at each step of the stimulations

and a Fiji macro was used to create videos for calculating (Ft-Fo) / Fo, termed "ΔF/Fo". Regions

of interest (ROIs) were circled around individual neurons in Fiji based on responses to KCl and

ΔF/Fo was recorded for each ROI. Graphs for each ROI were created in R or in Python. Each

graph and video file were manually checked to determine responses for each neuron. Neurons

that responded to vehicle were excluded from further analysis.

Statistical analysis

SPSS (v27, IBM, Armonk, NY) and Prism 10.0 (Graphpad, San Diego, CA) were used

for statistical analyses. Paired t-tests compared contralateral and ACLR groups for synovial fluid

ELISA and synovial tissue qPCR and flow cytometry analyses. Ordinary one-way ANOVA with

Tukey's multiple comparison evaluated treatment differences for *in vitro* cytokine and CXCL16

treatments. Repeated measures one-way ANOVA was used to analyze *in vitro* chemotaxis

experiments. Paired t-tests compared knee hyperalgesia threshold and knee withdrawal threshold

ratio between vehicle and CXCL16 injected limbs. Paired t-tests within each dose and across

both doses of CXCL16 compared histological parameters between vehicle injected and CXCL16

injected limbs. Two-way ANOVA was used to compare knee withdrawal threshold and knee

withdrawal threshold ratio between vehicle, CXCL16, and CXCL16 + ML339 injected limbs at

baseline and 7d post-injection (factors: timepoint, group). Linear mixed effects models were used

to compare knee withdrawal threshold in acute hyperalgesia experiments (**Suppl. Materials and**

Methods). Fisher's exact tests within each sex were used to determine associations between the

proportion of cells responsive to CXCL16 or to positive control only, in the presence or absence

of ML339. *P* values below 0.05 were considered significant throughout.

List of Supplementary Materials

Materials and Methods

Fig. S1. Synovial gene expression of chemokine ligands and chemokine receptors is perturbed in
mouse and human OA.

Fig. S2. qPCR corroboration of CXCL16 signaling axis expression and flow cytometry
supplement

Fig. S3. CXCL16 treatment *in vitro* – individual expression of macrophage score genes.

Fig. S4. Acute knee hyperalgesia supplement.

Fig. S5. Bulk RNAseq PCA plot.

Table S1. Antibodies.

Table S2. Gene primers for qPCR.

Table S3. PTOA severity scoring.

Table S4. Synovitis severity scoring.

Data file S1. Synovial bulk RNA-seq – quality control.

Data file S2. Synovial bulk RNA-seq – DEG lists.

References

1. C. R. Scanzello, S. R. Goldring, The role of synovitis in osteoarthritis pathogenesis. *Bone* **51**, 249-257 (2012).
2. E. Sanchez-Lopez, R. Coras, A. Torres, N. E. Lane, M. Guma, Synovial inflammation in osteoarthritis progression. *Nature Reviews Rheumatology* **18**, 258-275 (2022).
3. F. Angelini, P. Widera, A. Mobasheri, J. Blair, A. Struglics, M. Uebelhoer, Y. Henrotin, A. C. Marijnissen, M. Kloppenburg, F. J. Blanco, I. K. Haugen, F. Berenbaum, C. Ladel, J. Larkin, A. C. Bay-Jensen, J. Bacardit, Osteoarthritis endotype discovery via clustering of biochemical marker data. *Annals of the rheumatic diseases* **81**, 666-675 (2022).
4. M. F. Rai, K. H. Collins, A. Lang, T. Maerz, J. Geurts, C. Ruiz-Romero, R. K. June, Y. Ramos, S. J. Rice, S. A. Ali, C. Pastrello, I. Jurisica, C. T. Appleton, J. S. Rockel, M. Kapoor, Three decades of advancements in osteoarthritis research: insights from transcriptomic, proteomic, and metabolomic studies. *Osteoarthritis and cartilage* **32**, 385-397 (2024).
5. J. Blair, C. Bager, A. Platt, M. Karsdal, A.-C. Bay-Jensen, Identification of pathological RA endotypes using blood-based biomarkers reflecting tissue metabolism. A retrospective and explorative analysis of two phase III RA studies. *PLoS One* **14**, e0219980 (2019).
6. S. Ishihara, A. M. Obeidat, D. L. Wokosin, D. Ren, R. J. Miller, A.-M. Malfait, R. E. Miller, The role of intra-articular neuronal CCR2 receptors in knee joint pain associated with experimental osteoarthritis in mice. *Arthritis research & therapy* **23**, 1-12 (2021).
7. J. Miotla Zarebska, A. Chanalaris, C. Driscoll, A. Burleigh, R. E. Miller, A. M. Malfait, B. Stott, T. L. Vincent, CCL2 and CCR2 regulate pain-related behaviour and early gene expression in post-traumatic murine osteoarthritis but contribute little to chondropathy. *Osteoarthritis and Cartilage* **25**, 406-412 (2017).
8. R. E. Miller, P. B. Tran, R. Das, N. Ghoreishi-Haack, D. Ren, R. J. Miller, A.-M. Malfait, CCR2 chemokine receptor signaling mediates pain in experimental osteoarthritis. *Proceedings of the National Academy of Sciences* **109**, 20602-20607 (2012).
9. C. T. G. Appleton, S. E. Usmani, M. A. Pest, V. Pitelka, J. S. Mort, F. Beier, Reduction in disease progression by inhibition of transforming growth factor α–CCL2 signaling in experimental posttraumatic osteoarthritis. *Arthritis & rheumatology* **67**, 2691-2701 (2015).
10. L. Longobardi, J. D. Temple, L. Tagliafierro, H. Willcockson, A. Esposito, N. D'Onofrio, E. Stein, T. Li, T. J. Myers, H. Ozkan, M. L. Balestrieri, V. Ulici, R. F. Loeser, A. Spagnoli, Role of the CC chemokine receptor-2 in a murine model of injury-induced osteoarthritis. *Osteoarthritis and cartilage* **25**, 914-925 (2017).
11. C. H. Tang, C. J. Hsu, Y. C. Fong, The CCL5/CCR5 axis promotes interleukin⎕6 production in human synovial fibroblasts. *Arthritis & Rheumatism* **62**, 3615-3624 (2010).
12. H. Raghu, C. M. Lepus, Q. Wang, H. H. Wong, N. Lingampalli, F. Oliviero, L. Punzi, N. J. Giori, S. B. Goodman, C. R. Chu, J. B. Sokolove, W. H. Robinson, CCL2/CCR2, but not CCL5/CCR5, mediates monocyte recruitment, inflammation and cartilage destruction in osteoarthritis. *Annals of the rheumatic diseases* **76**, 914-922 (2017).
13. S. Abel, C. Hundhausen, R. Mentlein, A. Schulte, T. A. Berkhout, N. Broadway, D. Hartmann, R. Sedlacek, S. Dietrich, B. Muetze, B. Schuster, K. J. Kallen, P. Saftig, S. Rose-John, A. Ludwig, The transmembrane CXC-chemokine ligand 16 is induced by

IFN-gamma and TNF-alpha and shed by the activity of the disintegrin-like
metalloproteinase ADAM10. *Journal of immunology (Baltimore, Md. : 1950)* **172**, 6362-
6372 (2004).
14.   T. Nakayama, K. Hieshima, D. Izawa, Y. Tatsumi, A. Kanamaru, O. Yoshie, Cutting
edge: profile of chemokine receptor expression on human plasma cells accounts for their
efficient recruitment to target tissues. *The Journal of Immunology* **170**, 1136-1140
(2003).
15.   K. Hattermann, H. Gebhardt, S. Krossa, A. Ludwig, R. Lucius, J. Held-Feindt, R.
Mentlein, Transmembrane chemokines act as receptors in a novel mechanism termed
inverse signaling. *eLife* **5**, e10820 (2016).
16.   K. Hattermann, K. Bartsch, H. H. Gebhardt, H. M. Mehdorn, M. Synowitz, A. D.
Schmitt, R. Mentlein, J. Held-Feindt, "Inverse signaling" of the transmembrane
chemokine CXCL16 contributes to proliferative and anti-apoptotic effects in cultured
human meningioma cells. *Cell Communication and Signaling* **14**, 1-11 (2016).
17.   A. Wilbanks, S. C. Zondlo, K. Murphy, S. Mak, D. Soler, P. Langdon, D. P. Andrew, L.
Wu, M. Briskin, Expression cloning of the STRL33/BONZO/TYMSTR ligand reveals
elements of CC, CXC, and CX3C chemokines. *The Journal of Immunology* **166**, 5145-
5154 (2001).
18.   M. Matloubian, A. David, S. Engel, J. E. Ryan, J. G. Cyster, A transmembrane CXC
chemokine is a ligand for HIV-coreceptor Bonzo. *Nature immunology* **1**, 298-304 (2000).
19.   A. N. Wein, S. R. McMaster, S. Takamura, P. R. Dunbar, E. K. Cartwright, S. L.
Hayward, D. T. McManus, T. Shimaoka, S. Ueha, T. Tsukui, T. Masumoto, M. Kurachi,
K. Matsushima, J. E. Kohlmeier, CXCR6 regulates localization of tissue-resident
memory CD8 T cells to the airways. *The Journal of experimental medicine* **216**, 2748-
2762 (2019).
20.   K. Heesch, F. Raczkowski, V. Schumacher, S. Hunemorder, U. Panzer, H. W.
Mittrucker, The function of the chemokine receptor CXCR6 in the T cell response of
mice against Listeria monocytogenes. *PLoS One* **9**, e97701 (2014).
21.   M. J. Butcher, C. I. Wu, T. Waseem, E. V. Galkina, CXCR6 regulates the recruitment of
pro-inflammatory IL-17A-producing T cells into atherosclerotic aortas. *International
immunology* **28**, 255-261 (2016).
22.   K. M. Krawczyk, H. Nilsson, R. Allaoui, D. Lindgren, M. Arvidsson, K. Leandersson, M.
E. Johansson, Papillary renal cell carcinoma-derived chemerin, IL-8, and CXCL16
promote monocyte recruitment and differentiation into foam-cell macrophages.
Laboratory investigation; a journal of technical methods and pathology **97**, 1296-1305
(2017).
23.   R. van der Voort, A. W. van Lieshout, L. W. Toonen, A. W. Sloetjes, W. B. van den
Berg, C. G. Figdor, T. R. Radstake, G. J. Adema, Elevated CXCL16 expression by
synovial macrophages recruits memory T cells into rheumatoid joints. *Arthritis Rheum*
52, 1381-1391 (2005).
24.   N. R. Dhoke, K. Kaushik, A. Das, Cxcr6-Based Mesenchymal Stem Cell Gene Therapy
Potentiates Skin Regeneration in Murine Diabetic Wounds. *Molecular therapy : the
journal of the American Society of Gene Therapy*, (2020).
25.   J. H. Ruth, C. S. Haas, C. C. Park, M. A. Amin, R. J. Martinez, G. K. Haines, 3rd, S.
Shahrara, P. L. Campbell, A. E. Koch, CXCL16-mediated cell recruitment to rheumatoid

arthritis synovial tissue and murine lymph nodes is dependent upon the MAPK pathway. *Arthritis Rheum* **54**, 765-778 (2006).

26.   X. Zhang, J. X. Zhao, L. Sun, X. Y. Liu, [Expression of CXCL16/CXCR6 in fibroblast-like synoviocytes in rheumatoid arthritis and its role in synoviocyte proliferation]. *Beijing da xue xue bao. Yi xue ban = Journal of Peking University. Health sciences* **49**, 663-668 (2017).

27.   T. Isozaki, A. S. Arbab, C. S. Haas, M. A. Amin, M. D. Arendt, A. E. Koch, J. H. Ruth, Evidence that CXCL16 is a potent mediator of angiogenesis and is involved in endothelial progenitor cell chemotaxis : studies in mice with K/BxN serum-induced arthritis. *Arthritis Rheum* **65**, 1736-1746 (2013).

28.   T. Nanki, T. Shimaoka, K. Hayashida, K. Taniguchi, S. Yonehara, N. Miyasaka, Pathogenic role of the CXCL16–CXCR6 pathway in rheumatoid arthritis. *Arthritis & Rheumatism: Official Journal of the American College of Rheumatology* **52**, 3004-3014 (2005).

29.   R. E. Miller, R. J. Miller, A.-M. Malfait, Osteoarthritis joint pain: the cytokine connection. *Cytokine* **70**, 185-193 (2014).

30.   S. B. Oh, P. B. Tran, S. E. Gillard, R. W. Hurley, D. L. Hammond, R. J. Miller, Chemokines and glycoprotein120 produce pain hypersensitivity by directly exciting primary nociceptive neurons. *Journal of Neuroscience* **21**, 5027-5035 (2001).

31.   C. Yin, B. Liu, Z. Dong, S. Shi, C. Peng, Y. Pan, X. Bi, H. Nie, Y. Zhang, Y. Tai, Q. Hu, X. Wang, X. Shao, H. An, J. Fang, C. Wang, B. Liu, CXCL5 activates CXCR2 in nociceptive sensory neurons to drive joint pain and inflammation in experimental gouty arthritis. *Nature Communications* **15**, 3263 (2024).

32.   R. F. Bergman, L. Lammlin, L. Junginger, E. Farrell, S. Goldman, R. Darcy, C. Rasner, A. M. Obeidat, A.-M. Malfait, R. E. Miller, T. Maerz, Sexual dimorphism of the synovial transcriptome underpins greater PTOA disease severity in male mice following joint injury. *Osteoarthritis and Cartilage* **32**, 1060-1073 (2024).

33.   P. J. Gough, K. J. Garton, P. T. Wille, M. Rychlewski, P. J. Dempsey, E. W. Raines, A disintegrin and metalloproteinase 10-mediated cleavage and shedding regulates the cell surface expression of CXC chemokine ligand 16. *The Journal of Immunology* **172**, 3678-3685 (2004).

34.   N. Bao, B. Fu, X. Zhong, S. Jia, Z. Ren, H. Wang, W. Wang, H. Shi, J. Li, F. Ge, Q. Chang, Y. Gong, W. Liu, F. Qiu, S. Xu, T. Li, Role of the CXCR6/CXCL16 axis in autoimmune diseases. *International Immunopharmacology* **121**, 110530 (2023).

35.   A. J. Knights, E. C. Farrell, O. M. Ellis, L. Lammlin, L. M. Junginger, P. M. Rzeczycki, R. F. Bergman, R. Pervez, M. Cruz, E. Knight, D. Farmer, A. A. Samani, C.-L. Wu, K. D. Hankenson, T. Maerz, Synovial fibroblasts assume distinct functional identities and secrete R-spondin 2 in osteoarthritis. *Annals of the rheumatic diseases* **82**, 272-282 (2023).

36.   P. Rzeczycki, C. Rasner, L. Lammlin, L. Junginger, S. Goldman, R. Bergman, S. Redding, A. J. Knights, M. Elliott, T. Maerz, Cannabinoid receptor type 2 is upregulated in synovium following joint injury and mediates anti-inflammatory effects in synovial fibroblasts and macrophages. *Osteoarthritis and cartilage* **29**, 1720-1731 (2021).

37.   R. J. Miller, H. Jung, S. K. Bhangoo, F. A. White, Cytokine and chemokine regulation of sensory neuron function. *Sensory nerves*, 417-449 (2009).

38. H.-G. Schaible, Nociceptive neurons detect cytokines in arthritis. *Arthritis research &*
therapy **16**, 1-9 (2014).
39. A. M. Obeidat, M. J. Wood, N. S. Adamczyk, S. Ishihara, J. Li, L. Wang, D. Ren, D. A.
Bennett, R. J. Miller, A.-M. Malfait, R. E. Miller, Piezo2 expressing nociceptors mediate
mechanical sensitization in experimental osteoarthritis. *Nature communications* **14**, 2479
(2023).
40. C. Wang, A. Di, Y. Wu, M. Liu, M. Wei, Z. Liang, F. Liu, H. Fan, B. Dong, C. Li,
Sprouting sympathetic fibres release CXCL16 and norepinephrine to synergistically
mediate sensory neuronal hyperexcitability in a rodent model of neuropathic pain. *British*
Journal of Anaesthesia, (2025).
41. M. A. Di Castro, F. Trettel, G. Milior, L. Maggi, D. Ragozzino, C. Limatola, The
chemokine CXCL16 modulates neurotransmitter release in hippocampal CA1 area.
Scientific reports **6**, 34633 (2016).
42. M. Rosito, C. Deflorio, C. Limatola, F. Trettel, CXCL16 orchestrates adenosine A3
receptor and MCP-1/CCL2 activity to protect neurons from excitotoxic cell death in the
CNS. *Journal of Neuroscience* **32**, 3154-3163 (2012).
43. M. Rosito, C. Lauro, G. Chece, A. Porzia, L. Monaco, F. Mainiero, M. Catalano, C.
Limatola, F. Trettel, Trasmembrane chemokines CX3CL1 and CXCL16 drive interplay
between neurons, microglia and astrocytes to counteract pMCAO and excitotoxic
neuronal death. *Frontiers in cellular neuroscience* **8**, 193 (2014).
44. M. Darash-Yahana, J. W. Gillespie, S. M. Hewitt, Y.-Y. K. Chen, S. Maeda, I. Stein, S.
P. Singh, R. B. Bedolla, A. Peled, D. A. Troyer, E. Pikarsky, M. Karin, J. M. Farber, The
chemokine CXCL16 and its receptor, CXCR6, as markers and promoters of
inflammation-associated cancers. *PloS one* **4**, e6695 (2009).
45. M. Lehrke, A. Konrad, M. Lehrke, A. Konrad, V. Schachinger, C. Tillack, F. Seibold, R.
Stark, K. G. Parhofer, U. C. Broedl, CXCL16 is a surrogate marker of inflammatory
bowel disease. *Scandinavian journal of gastroenterology* **43**, 283-288 (2008).
46. L. Zhang, L. Ran, G. E. Garcia, X. H. Wang, S. Han, J. Du, W. E. Mitch, Chemokine
CXCL16 regulates neutrophil and macrophage infiltration into injured muscle, promoting
muscle regeneration. *The American journal of pathology* **175**, 2518-2527 (2009).
47. J. Zhang, W. Hao, J. Zhang, T. Li, Y. Ma, Y. Wang, X. Li, W. Cui, J. Du, CXCL16
promotes Ly6Chigh monocyte infiltration and impairs heart function after acute
myocardial infarction. *The Journal of Immunology* **210**, 820-831 (2023).
48. P. M. Hershberger, S. Peddibhotla, E. Sugarman, P. Maloney, D. Key, E. Suyama, K.
Nguyen, S. Vasile, M. Kraft, D. Stonich, Probing the CXCR6/CXCL16 axis: targeting
prevention of prostate cancer metastasis. *Probe Reports from the NIH Molecular*
Libraries Program [Internet], (2014).
49. S. Peddibhotla, P. M. Hershberger, R. J. Kirby, E. Sugarman, P. R. Maloney, E. H.
Sessions, D. Divlianska, C. J. Morfa, D. Terry, A. B. Pinkerton, L. H. Smith, S. Malany,
Discovery of small molecule antagonists of chemokine receptor CXCR6 that arrest tumor
growth in SK-HEP-1 mouse xenografts as a model of hepatocellular carcinoma.
Bioorganic & medicinal chemistry letters **30**, 126899 (2020).
50. T. Chen, C. Qiao, E. Yinwang, S. Wang, X. Wen, Y. Feng, X. Jin, S. Li, Y. Xue, H.
Zhou, W. Zhang, X. Zeng, Z. Wang, H. Sun, L. Jiang, H. Li, B. Li, Z. Cai, Z. Ye, N. Lin,
Natural lung-tropic TH9 cells: a sharp weapon for established lung metastases. *Journal*
for Immunotherapy of Cancer **12**, e009629 (2024).

51. D. R. Seifer, B. D. Furman, F. Guilak, S. A. Olson, S. C. Brooks III, V. B. Kraus, Novel
synovial fluid recovery method allows for quantification of a marker of arthritis in mice.
Osteoarthritis and cartilage **16**, 1532-1538 (2008).
52. C. B. Little, A. Barai, D. Burkhardt, S. M. Smith, A. J. Fosang, Z. Werb, M. Shah, E. W.
Thompson, Matrix metalloproteinase 13-deficient mice are resistant to osteoarthritic
cartilage erosion but not chondrocyte hypertrophy or osteophyte development. *Arthritis*
Rheum **60**, 3723-3733 (2009).
53. M. T. Jackson, B. Moradi, S. Zaki, M. M. Smith, S. McCracken, S. M. Smith, C. J.
Jackson, C. B. Little, Depletion of protease-activated receptor 2 but not protease-
activated receptor 1 may confer protection against osteoarthritis in mice through
extracartilaginous mechanisms. *Arthritis & rheumatology (Hoboken, N.J.)* **66**, 3337-3348
(2014).
54. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome biology* **15**, 1-21 (2014).
55. G. Yu, L.-G. Wang, Y. Han, Q.-Y. He, clusterProfiler: an R package for comparing
biological themes among gene clusters. *Omics: a journal of integrative biology* **16**, 284-
287 (2012).
56. R. E. Miller, S. Ishihara, P. B. Tran, S. B. Golub, K. Last, R. J. Miller, A. J. Fosang, A.-
M. Malfait, An aggrecan fragment drives osteoarthritis pain through Toll-like receptor 2.
JCI insight **3**, e95704 (2018).
805

Acknowledgements:

The authors would like to acknowledge the Michigan Integrative Musculoskeletal Health

Core Center (MiMHC) which is supported by P30AR069620 from NIH/NIAMS for processing

samples for histology and for use of their microscopy equipment. We also wish to acknowledge

the University of Michigan Biomedical Research Core Facilities, specifically the flow cytometry

core, which is a part of the Medical School Office of Research, for use of their flow cytometer.

We would like to thank the Chicago Center on Musculoskeletal Pain (C-COMP) funded by

P30AR079206 for technical assistance with calcium imaging.

 Figures and tables:

816 **Fig. 1. Joint injury activates the CXCL16-CXCR6 axis in synovium.** (**A**) Volcano plots of

817 mouse synovial bulk RNAseq data (*32*) comparing chemokine ligand expression in Sham vs 7d

818 ACLR and Sham vs 28d ACLR. Genes with red datapoints were differentially expressed at P_{adj} <

819 0.05 and $|\log_2 FC| > 0.585$. (**B**) Bubble plot from a meta-analysis of human OA synovial

820 transcriptomics datasets showing expression of *CXCL16*, *CXCR6*, and *ADAM10* relative to

healthy synovium. (**C**) CXCL16 ELISA of synovial fluid in contralateral and injured joints 3d

post-ACLR (n=6 per group). Total number of (**D**) and percent positive (**E**) CXCL16+ and

CXCR6+ cells in contralateral and injured synovium at 3d and 14d post-ACLR (n=3 per group. 1

sample = 1 male + 1 female synovia pooled). (**F**) Proportions plots showing the cell type

breakdown of CXCL16+ and CXCR6+ cells. (**G**) Proportions plots further breaking down the

CXCL16+ and CXCR6+ immune cell types. (**H**) Feature plots of mouse synovial single cell

RNAseq data (*35*) showing *Cxcl16, Cxcr6,* and *Adam10* gene expression distribution across all

cell types. All bars show mean ± SEM.

Fig. 2. Upstream mediators of the CXCL16 signaling axis *in vitro*. (**A**) Fibroblast like

synoviocytes (FLS), bone marrow-derived macrophages (BMDM), and immortalized

chondrogenic cells (ATDC5s) were treated with pro-inflammatory cytokines and perturbation of

the CXCL16 signaling axis was evaluated via qPCR. (**B**) Gene expression of *Cxcl16*, *Cxcr6*, and

Adam10 by qPCR in FLS (n=6 per group) and a CXCL16 ELISA of FLS conditioned media

(n=6-7 per group). (**C**) Gene expression of *Cxcl16*, *Cxcr6*, and *Adam10* by qPCR in BMDMs

(n=6 per group). (**D**) Gene expression of *Cxcl16*, *Cxcr6*, and *Adam10* by qPCR in ATDC5s. (n=4

per group). All bars show mean ± SEM.

839
Fig. 3. Downstream effects of CXCL16 *in vitro*. (**A**) In FLS, gene expression of the CXCL16

signaling axis, pro-inflammatory markers, and anti-inflammatory markers was measured by

qPCR following stimulation with recombinant CXCL16 (n=6 per group). (**B**) In BMDMs, gene

expression of the CXCL16 signaling axis and a macrophage score was measured by qPCR

following stimulation with recombinant CXCL16 (n=6 per group). (**C**) In BMPCs, gene

expression of the CXCL16 signaling axis, pro-inflammatory markers, and anti-inflammatory

markers was measured by qPCR following stimulation with recombinant CXCL16 (n=6 per

group). (**D**) Chemotaxis of BMDMs and BMPCs in response to CXCL16 was measured by

migrated cell density (n=3 per group). All bars show mean ± SEM.

849
Fig. 4. CXCL16-induced hyperalgesia and modulation of the synovial transcriptome. (**A**)
Knee withdrawal threshold following repeated intra-articular injection of CXCL16 (either low
dose or high dose) or vehicle in the contralateral side. Testing occurred at 7d and 14d post-first
injection (n=6 per dose). (**B**) Histopathological scoring of limbs given repeated intra-articular

injections of CXCL16 (either low or high dose) or vehicle in the contralateral side. Total PTOA

score and total synovitis score are shown along with the synovitis subscores of anterior, inferior

synovial lining hyperplasia and synovial fibrosis (n=5-6 mice per dose, n=3-12 images averaged

together per limb). (**C**) Representative low dose CXCL16-injected limbs and vehicle-injected

contralateral limbs. Yellow arrows – synovial lining hyperplasia. Green arrows – synovial

fibrosis. (**D**) Knee withdrawal threshold and knee withdrawal threshold ratio (CXCL16/vehicle)

following repeated intra-articular injection of CXCL16 (high dose only) or vehicle in the

contralateral side (n=3 per sex). (**E**) Venn diagram of synovial bulk RNAseq showing

differentially expressed genes (DEGs) between vehicle and CXCL16 injected joints in males

compared to females (n=3 per group per sex, $|\log_2 FC| > .585$, $P_{adj} < .05$). Heatmap of male

DEGs. Red indicates upregulated and blue indicates downregulated in CXCL16 injected joints

relative to vehicle injected joints. (**F**) Gene set enrichment analysis of male synovia comparing

vehicle vs CXCL16 injected joints. (**G**) Knee withdrawal threshold following repeated bilateral

intra-articular injections of vehicle, CXCL16, or CXCL16 co-administered with ML339 (n=6

mice, i.e. n=12 limbs, per group). All bars show mean ± SEM.

870
Fig. 5. CXCL16-induced acute hyperalgesia via CXCR6. (A) Experimental design for intra-

articular CXCL16 injection followed by acute knee hyperalgesia. Knee withdrawal threshold (**B**)

and knee withdrawal threshold ratio (**C**) at acute timepoints following intra-articular injection of

CXCL16 or vehicle in the contralateral joint (n=3 per sex). (**D**) Experimental design for intra-

articular CXCL16 +/- ML339 injection followed by acute knee hyperalgesia. Knee withdrawal

threshold (**E**) and knee withdrawal threshold ratio (**F**) at acute timepoints following intra-

articular injection of CXCL16 or CXCL16 + ML339, and vehicle in the contralateral joint (n=9

per group). (**G**) Experimental design for intra-peritoneal ML339 injections and acute knee

hyperalgesia post-ACLR. Knee withdrawal threshold (**H**) and knee withdrawal threshold ratio (**I**)

pre- and post-intraperitoneal injection of vehicle or ML339 at 7d post-ACLR (n=8-9 per group).

(**J**) Complete blood count analysis showing total number of circulating white blood cells and

neutrophils from naïve mice and mice treated with vehicle or ML339 7d post-ACLR (n=10-13

per group). All bars show mean ± SEM.

Fig. 6. CXCL16-induced Ca^{2+} in Na$_V$1.8 nociceptive neurons *in vitro*. (**A**) Experimental

design: Murine DRG cultures were pre-treated with either vehicle or ML339, treated with

CXCL16 or CXCL16 + ML339, and then followed by a positive control stimulus, KCl. (**B**)

Quantification of percent responsive cells to CXCL16 stimulation in the presence and absence of

CXCR6 antagonist, ML339. Total number of responsive cells is detailed for each sex. (**C**)

Representative images of each stage of the experiment – pretreatment, treatment, positive

control. (**D**) Representative line plots showing ΔF/Fo of a single responsive cell (responds to

KCl) for each condition.

Funding:

National Science Foundation Graduate Research Fellowship grant DGE 2241144 (LL)

National Institutes of Health K99 grant K99AR081894 (AJK)

Pioneer Fellowship from the University of Michigan (AJK)

Orthopaedic Research Society Scientific Network Award (TM, REM)

F31AR083277 (NSA)

R01AR077019 (REM)

R01AR077019-03S1 (REM)

R01AR080035 (TM)

Department of Orthopaedic Surgery, University of Michigan (TM)

Author contributions:

Conceptualization: LL, SR, REM, TM
Methodology: LL, SR, AJK, MDN, PMR, REM, TM
Investigation: LL, SR, AJK, AM, MDN, RFB, IS, SCH, SGN, PMR, NSA
Visualization: LL, AM, MDN, IS
Funding acquisition: LL, NSA, REM, TM
Writing – original draft: LL, TM
Writing – review & editing: LL, AJK, MDN, NSA, REM, TM

Competing interests:

All authors declare that they have no competing interests.

Data and materials availability:

Bulk RNA-seq data will be publicly available on GEO and all other data are available in the
main text or the supplementary materials.