Transcription factor reveals interconnected regulation of carbon source utilization and

carbon catabolite repression in an oleaginous yeast

Brandon Reyes-Chavez[1,2,*], Joshua D. Kerkaert[1,*], and Lori B. Huberman[1,#]

[1]Plant Pathology and Plant-Microbe Biology Section, School of Integrative Plant Science,

Cornell University, Ithaca, NY 14853

[2]Microbiology Department, Cornell University, Ithaca, NY 14853

*These authors contributed equally to this work.

[#]To whom correspondence may be addressed: huberman@cornell.edu

ABSTRACT

Cells must sense and respond to available nutrients to utilize resources and establish fungal colonies. Saprophytic microbes harvest carbon from plant biomass, which contains insoluble, complex carbohydrates that must be degraded extracellularly prior to import into the cell. Filamentous fungi can degrade these complex carbohydrates. However, while many single cellular microbes can utilize the building blocks of these insoluble polysaccharides, they frequently are unable to degrade insoluble carbohydrates. Cellobiose is a disaccharide breakdown product of cellulose, the most abundant component of plant biomass. We investigated the genetic regulation of cellobiose utilization in the oleaginous, basidiomycete yeast *Rhodotorula (Rhodosporidium) toruloides* using a combination of phylogenetic, genetic, and genomic approaches. We tested the closest *R. toruloides* homolog of the cellulose degradation regulator CLR-2/ClrB for a role in cellobiose utilization. Deletion of this transcription factor, which we named Cbr1, eliminated the ability of cells to grow on media containing cellobiose as the sole carbon source. Although the role of CLR-2/ClrB is limited to regulating the expression of genes involved in the utilization of cellulose and hemicellulose, Cbr1 plays a role not only in cellobiose utilization, but also utilization of tricarboxylic acid (TCA) cycle intermediates. Transcriptional profiling revealed that the genes activated by Cbr1 included genes encoding β-glucosidases and transporters. These β-glucosidases are secreted and are necessary for cellobiose utilization, while one of the transporters is required for utilization of the TCA cycle intermediate, citrate. Additionally, Cbr1 inhibits carbon catabolite repression specifically during utilization of disaccharides, which may have evolved to limit the repression of genes encoding secreted proteins that cleave disaccharides into glucose extracellularly. The co-regulation of cellobiose and TCA cycle utilization with carbon catabolite repression may shed light on alternative methods of the regulation of carbon catabolite repression in fungi.

INTRODUCTION

Cells must sense and respond to nutrients to successfully grow and divide using the resources available. For fungi, nutrient sensing is critical to establish fungal colonies, compete with other microbes, and establish pathogenic and symbiotic relationships with plants and animals. Saprophytic fungi acquire nutrients from dead plant matter. The carbon sources found in plant biomass range from simple sugars, like sucrose and fructose, to complex carbohydrates, such as cellulose, found in plant cell walls.

Utilization of complex polysaccharides requires extracellular degradation to break down the insoluble molecules into soluble building blocks that can be imported into fungal cells. Many yeast species can utilize the soluble building blocks of polysaccharides found in plant biomass (1, 2). These yeast species are excellent choices for growth and engineering in industrial settings because of their uninucleate, single cellular lifestyle and wide carbon utilization repertoire. The basidiomycete, oleaginous yeast *Rhodotorula toruloides* (previously known as *Rhodosporidium toruloides*) can eat the vast majority of soluble breakdown products in lignocellulose (3, 4). *R. toruloides* can also accumulate up to 70% of its biomass as fatty acids in conditions in which carbon is abundant but other nutrients are limiting and produces useful secondary metabolites, such as carotenoids (5–11). This makes *R. toruloides* an excellent choice as a potential industrial fungus for green biotechnology in which products that might otherwise be sourced from petroleum or other environmentally unsustainable sources are produced in microbes fed by agricultural waste or plant matter grown on marginal land (12).

Most of the work to characterize the genetic pathways regulating utilization of carbon sources has been done in Ascomycete fungi (1, 13). In these fungi, a network of activating transcription factors activates the genes necessary for utilization of the carbohydrates found in plant biomass (14). Plant biomass contains a wide array of carbon sources. These varied carbon sources have differing levels of preference for the fungal cell. When a more preferred carbon source is available a process known as carbon catabolite repression downregulates the

expression of genes required for utilization of any less preferred carbon sources present in the

environment (1, 15). In Ascomycete fungi, one of the major regulators of carbon catabolite

repression is the transcription factor CRE-1/CreA/Mig1, which broadly represses the expression

of genes necessary for utilization of any nonpreferred carbon source in the presence of any

more preferred carbon source (15–21). However, the genetic mechanisms through which

Basidiomycete fungi activate genes necessary to utilize nonpreferred carbon sources and

repress these genes in the presence of more preferred carbon sources is significantly less well

studied.

We investigated the genetic mechanisms regulating carbon utilization in the

basidiomycete yeast *R. toruloides*. Cellulose is the most abundant component of plant biomass

and is composed of long chains of glucose linked by β-1,4-glycosidic bonds (22). *R. toruloides*

can utilize the disaccharide building block of cellulose, cellobiose (23). We took a phylogenetic

approach to identify the regulator of cellobiose utilization in *R. toruloides* and identified a

transcription factor homologous to the cellulose degradation regulator CLR-2/ClrB in

Ascomycete fungi (24) that is required for utilization of cellobiose in *R. toruloides*, which we

named Cbr1.

Characterization of Cbr1 revealed that the role of this transcription factor is significantly

expanded in *R. toruloides*, where it is also required for utilization of tricarboxylic acid (TCA)

cycle intermediates. We used transcriptomic analysis to identify genes regulated by Cbr1. These

genes included two genes encoding secreted β-glucosidases and two genes encoding

transporters. These β-glucosidase genes are necessary for cellobiose utilization, while one of

the transporters is required for utilization of the TCA cycle intermediate citrate. Cellobiose and

TCA cycle intermediates are not utilized through similar pathways, suggesting that *R. toruloides*

may inhabit an ecological niche in which these carbon sources are frequently found together.

Further investigation into the role of Cbr1 demonstrated that this transcription factor inhibits

carbon catabolite repression specifically when a disaccharide is present as a nonpreferred

carbon source. This role for Cbr1 contrasts with previously identified genetic mechanisms of

carbon catabolite repression regulation, which are involved in the regulation of genes

responsible for the utilization of any nonpreferred carbon source. Taken together, the varied

roles of Cbr1 in *R. toruloides* metabolic regulation presents a novel mechanism of carbon

catabolite repression in fungi and may provide insights into the ecological niche of this

oleaginous yeast.

RESULTS

Cbr1 is required for wild type utilization of cellobiose and TCA cycle intermediates

Cellulose is the most abundant component of lignocellulose. Although *R. toruloides*

cannot degrade cellulose, it is capable of utilizing the simple sugar breakdown product of

cellulose, cellobiose (23). Several transcription factors are known to regulate cellulose and

cellobiose utilization in filamentous fungi, including CLR-1/ClrA, CLR-2/ClrB, and XlnR (1, 24–

27). To investigate the genetic mechanisms regulating *R. toruloides* cellobiose utilization, we

searched for transcription factors in the *R. toruloides* genome with homology to these regulators

of cellulose and cellobiose utilization. No clear homolog of CLR-1/ClrA or XlnR existed in the *R.*

toruloides genome (Fig S1). However, we did identify a single homolog of CLR-2/ClrB,

RTO4_13588 (Fig 1A and Fig S1).

We hypothesized that RTO4_13588 may play a role in regulating cellobiose utilization in

R. toruloides. To test this hypothesis, we deleted RTO4_13588 and grew the resulting

RTO4_13588Δ strain with cellobiose as the sole carbon source. Growth of cells lacking

RTO4_13588 with glucose as the sole carbon source was indistinguishable from that of wild

type cells (Fig 1B). However, deletion of RTO4_13588 resulted in a complete lack of growth

when cellobiose was provided as the sole carbon source (Fig 1B). Expression of RTO4_13588

under the native promoter at an exogenous location (the *YKU70* locus) to complement the

RTO4_13588 deletion, restored growth of this strain to wild type levels (Fig 1B). The closest

homolog to RTO4_13588 in the model yeast *Saccharomyces cerevisiae* is the transcription

factor Gal4, which regulates galactose utilization in *S. cerevisiae* (Fig 1A) (28). To test whether

RTO4_13588 also plays a role in regulating galactose utilization in *R. toruloides*, we grew wild

type cells and the RTO4_13588Δ strain with galactose as the sole carbon source. There was no

detectable difference in growth between wild type cells and cells lacking RTO4_13588 (Fig

S2A). Thus, we named RTO4_13588, *CBR1* for cellobiose regulator 1.

R. toruloides is capable of growth on a wide variety of soluble carbon sources (3, 23).

We asked whether *CBR1* is also required for growth on any additional carbon sources. The

growth of *cbr1*Δ cells was indistinguishable from that of wild type on all the other sugars we

tested (arabinose, fructose, galacturonic acid, maltose, sucrose, and xylose) (Fig S2A). *R.*

toruloides can use amino acids as a carbon source. However, *CBR1* was not required for

utilization of glutamate, glycine, or L-alanine (Fig S2B). Cells lacking *CBR1* were also capable of

growing as well as wild type cells on media containing acetate and the lipid Tween-80 as the

sole carbon source (Fig S2C). Additionally, both wild type and *cbr1*Δ cells could grow on the

phenolic breakdown products of lignin, coumaric acid and ferulic acid (Fig S2D).

Cells lacking *CBR1* are capable of respiration, as evidenced by their ability to grow with

acetate, lipids, and amino acids as the sole carbon source. Thus, we hypothesized that *cbr1*Δ

cells would be able to grow with TCA cycle intermediates provided as the sole carbon source. To

our surprise, when we grew wild type and *cbr1*Δ cells on media containing citrate, α-

ketoglutarate, succinate, fumarate, and malate as the sole carbon source, the *cbr1*Δ cells had

significant growth defects relative to wild type (Fig 1B). Complementation of the *cbr1* deletion

with *CBR1* expressed under the native promoter at the *YKU70* locus restored growth of the

*cbr1*Δ cells to a level indistinguishable from wild type.

The role for Cbr1 in TCA cycle intermediate utilization could either be an expanded role

for Cbr1 in *R. toruloides* relative to CLR-2/ClrB or a previously uncharacterized role for CLR-

2/ClrB. To distinguish between these two possibilities, we tested whether *clr-2* is required for

TCA cycle intermediate utilization in the model filamentous fungus *Neurospora crassa.* We

directly inoculated *N. crassa* wild type and Δ*clr-2* cells into media containing succinate, α-

ketoglutarate, or malate as the sole carbon source. Wild type *N. crassa* was only capable of

limited growth with TCA cycle intermediates as the sole carbon source, and deletion of *clr-2*

caused no detectable difference in *N. crassa* growth on these carbon sources (Fig S3). This

data suggests that the role for *CBR1* in regulating TCA cycle intermediate utilization is an

expanded role for this transcription factor, relative to the role of Cbr1 orthologs in Ascomycete

filamentous fungi.

Cbr1 activates five genes in response to cellobiose and TCA cycle intermediates

We had three possible hypotheses for the genetic mechanism through which Cbr1 could

regulate utilization of both cellobiose and TCA cycle intermediates: (1) Cbr1 regulates genes

necessary to utilize both cellobiose and TCA cycle intermediates in response to all of these

carbon sources, (2) Cbr1 regulates genes necessary to utilize cellobiose in response to

cellobiose and genes necessary to utilize TCA cycle intermediates in response to TCA cycle

intermediates, and (3) *R. toruloides* utilization of cellobiose and/or TCA cycle intermediates

proceeds through atypical metabolic pathways such that they require the same catabolic

enzymes. To distinguish between these three hypotheses, we used transcriptional profiling. We

grew wild type cells and *cbr1*Δ cells in yeast extract peptone dextrose (YPD) medium overnight

so that the cells were in exponential growth phase, washed the cells three times in yeast

nitrogen base (YNB) medium without a carbon source, and then shifted the cells to YNB

containing 1% cellobiose as the sole carbon source, 1% succinate as the sole carbon source, or

lacking a carbon source for 8 h prior to harvesting the cells for RNA sequencing.

The expression of 776 genes was at least 4-fold differentially expressed between wild

type cells exposed to carbon starvation and cellobiose as the sole carbon source, with 139

genes upregulated and 637 genes downregulated in cells exposed to cellobiose as compared to

carbon starvation. Genes upregulated during exposure to cellobiose were enriched for genes

associated with carbohydrate metabolism, nucleosome assembly, and sulfur metabolism via

selenoamino acid metabolism. Genes downregulated during exposure to cellobiose compared

to carbon starvation were enriched for general metabolic processes, including regulation of

nitrogen utilization, amino acid metabolism, and transport associated genes (Fig 2B-C).

Far fewer genes (373) were at least 4-fold differentially expressed between wild type

cells exposed to carbon starvation and succinate, with 116 upregulated and 257 downregulated

in cells exposed to succinate as compared to carbon starvation. While only selenoamino acid

metabolism was enriched for genes upregulated on succinate, genes associated with nitrogen

utilization, electron transport, and carbohydrate transport were downregulated in succinate

compared to carbon starvation (Fig 2B-C). The majority of genes (289 out of 373) differentially

expressed by at least 4-fold during exposure to succinate as compared to carbon starvation

were also differentially expressed by at least 4-fold during exposure to cellobiose as compared

to carbon starvation. Fifty-seven genes were upregulated in both cellobiose and succinate as

compared to carbon starvation by at least 4-fold. Of the 257 genes downregulated by at least 4-

fold in response to succinate, 228 were also downregulated by at least 4-fold in response to

cellobiose. This supports the hypothesis that similar sets of genes are regulated in response to

both cellobiose and succinate.

Although the expression of many fungal transcription factor genes is upregulated in the

condition in which the transcription factor plays a role, *CBR1* expression was not activated in

response to either cellobiose or succinate as compared to carbon starvation (Fig S4A). To

directly regulate gene expression, Cbr1 must be present in the nucleus. We tagged Cbr1 with

green fluorescent protein (mEGFP) and the histone H2A subunit *HTA1* (RTO4_12090) with

mRuby2 and exposed these cells to glucose, cellobiose, or succinate as the sole carbon source

to identify the localization pattern of Cbr1. Cbr1 colocalized with the histone H2A subunit under

all conditions, suggesting that Cbr1 is expressed and present in the nucleus during exposure to

glucose, cellobiose, and succinate (Fig S4B).

To test our hypotheses for potential connections in the regulation of cellobiose and

succinate utilization, we examined the genes differentially expressed in wild type as compared

to *cbr1Δ* cells. The expression of 635 genes was at least 4-fold differentially expressed in *cbr1Δ*

cells relative to wild type cells during exposure to cellobiose, with 270 genes downregulated and

365 genes upregulated in *cbr1Δ* cells as compared to wild type. Similar to WT cells on

cellobiose compared to carbon starvation, *cbr1Δ* cells on cellobiose had higher levels of

transcripts associated with transport than WT cells. Additionally, genes associated with protein

folding, amino acid and protein metabolism, and sulfur metabolism were downregulated in *cbr1Δ*

cells as compared to wild type cells when exposed to cellobiose (Fig 2B-C). Nearly half of these

genes (281) were also at least 4-fold differentially expressed in wild type cells exposed to

cellobiose as compared to carbon starvation.

Only 22 genes were at least 4-fold differentially expressed between cells lacking *cbr1*

and wild type cells exposed to succinate. The 10 genes downregulated in *cbr1Δ* cells as

compared to wild type cells during exposure to succinate were enriched for several KEGG

categories, including starch and sucrose metabolism, that upon closer inspection were largely

driven by a decrease in expression of two putative β-glucosidases. Similarly, the KEGG

categories enriched in genes downregulated in *cbr1Δ* cells relative to wild type cells in carbon

starvation were largely driven by expression of the two putative β-glucosidase genes (Fig 2B-C).

Thirteen genes were at least 4-fold differentially expressed in *cbr1*Δ cells relative to wild

type cells during exposure to both cellobiose and succinate. Of these 13 genes, 5 were

downregulated and 8 were upregulated in *cbr1*Δ cells relative to wild type during exposure to

both cellobiose and succinate. To distinguish between our possible hypotheses for Cbr1 co-

regulation of cellobiose and succinate, we focused on genes downregulated in cells lacking *cbr1*

as compared to wild type. These genes included two genes encoding predicted β-glucosidases

(RTO4_16716 and RTO4_16717), two genes encoding predicted major facilitator superfamily

(MFS) transporters (RTO4_13825 and RTO4_10339), and a gene encoding a predicted

phosphoglycerate mutase (RTO4_11229). The downregulation of genes encoding β-

glucosidases, which may cleave cellobiose into its constituent glucose molecules, in *cbr1*Δ cells

relative to wild type cells during exposure to both cellobiose and succinate may support our

hypothesis that Cbr1 regulates genes necessary to utilize both cellobiose and TCA cycle

intermediates in response to all of these carbon sources. The eight genes upregulated in *cbr1*Δ

cells relative to wild type during exposure to both cellobiose and succinate, included a gene

encoding a predicted metallo-β-lactamase superfamily protein (RTO4_11764), a gene encoding

a predicted peptidoglycan-N-acetylmuramic acid deacetylase (RTO4_9785), a gene encoding a

predicted cyclin (RTO4_15596), a gene encoding a predicted methyltransferase (RTO4_14040),

and four hypothetical proteins (RTO4_8662, RTO4_9846, RTO4_10117, and RTO4_13502).

β-glucosidase genes expressed in a CBR1-dependent fashion in response to cellobiose and

TCA cycle intermediates are necessary for cellobiose utilization

The expression of predicted β-glucosidase genes in a *CBR1*-dependent fashion in

response to both cellobiose and succinate, led us to hypothesize that Cbr1 regulates genes

necessary to utilize both cellobiose and TCA cycle intermediates in response to all of these

carbon sources. To test this hypothesis, we deleted RTO4_16716 and RTO4_16717, which both

encode predicted β-glucosidases and are adjacent on chromosome 9 of the *R. toruloides*

genome. Deletion of RTO4_16716 and RTO4_16717 had no effect on growth on glucose but

eliminated the ability of *R. toruloides* to grow when cellobiose was provided as the sole carbon

source (Fig 3A). Canonical TCA cycle intermediate utilization pathways do not require β-

glucosidases. Thus, we hypothesized that even though RTO4_16716 and RTO4_16717 were

expressed in a *CBR1*-dependent fashion during succinate exposure, they would not actually be

required for TCA cycle intermediate utilization. Indeed, growth of wild type cells and

RTO4_16716Δ RTO4_16717Δ cells was indistinguishable when succinate or citrate were

provided as the sole carbon source (Fig 3A). This data suggests that RTO4_16716 and

RTO4_16717 are necessary for growth on cellobiose but not the TCA cycle intermediates

succinate or citrate.

Both RTO4_16716 and RTO4_16717 had predicted secretion signals (Fig S5). We

asked whether it was possible to detect β-glucosidase activity in the supernatant of *R. toruloides*

cultures. We grew wild type cells, *cbr1*Δ cells, and RTO4_16716Δ RTO4_16717Δ cells in YNB

with glucose as the sole carbon source overnight to exponential growth phase, back diluted the

cultures into YNB containing glucose as the sole carbon source for 8 doublings, washed them

twice in YNB lacking a carbon source, shifted them to YNB with either glucose, cellobiose, or

succinate as the sole carbon source, and then harvested culture supernatants after 20 hours of

incubation to measure β-glucosidase activity. Wild type cells had no detectable β-glucosidase

activity during growth on glucose (Fig 3B). When wild type cells were grown on cellobiose, we

could detect substantial β-glucosidase activity in the supernatant. However, deletion of *CBR1* or

RTO4_16716 and RTO4_16717 resulted in a complete abolition of β-glucosidase activity (Fig

3B). β-glucosidases are not required for TCA cycle intermediate utilization. However, β-

glucosidase activity was present in wild type succinate cultures at a level that was 30% of wild

type cellobiose supernatants (Fig 3B). Thus, we named RTO4_16716 and RTO4_16717, *BGL1*

and *BGL2*, respectively, for β-glucosidase 1 and 2.

If *BGL1* and *BGL2* are sufficient for growth on cellobiose, we hypothesized that it would

be possible to grow cells lacking *CBR1* or *BGL1* and *BGL2* with cellobiose as the sole carbon

source if the media was supplemented with spent supernatant from wild type cells grown on

cellobiose or succinate. To test this hypothesis, we grew wild type cells in media containing

either glucose, cellobiose, or succinate as the sole carbon source. We harvested supernatant

from these cultures and then supplemented media containing cellobiose as the sole carbon

source with either 6%, 12%, or 25% of the filtered spent supernatant. As expected, supernatants

from wild type cells grown in cellobiose and succinate exhibited β-glucosidase activity, while

supernatant from wild type cells grown in glucose did not (Fig S6A). Wild type cells, *cbr1Δ* cells,

and *bgl1Δ bgl2Δ* cells were then inoculated into this cellobiose media supplemented with spent

supernatant. As a control for growth on any of the nutrients present in the spent supernatant, we

also supplemented media lacking a carbon source with either 6%, 12%, or 25% of the filtered

spent supernatant and subtracted any growth seen in this condition from the growth in the spent

supernatant supplemented cellobiose media (Fig. 3C-E and Fig S6B-G).

To detect growth of the *cbr1Δ* and *bgl1Δ bgl2Δ* cells, we allowed the strains to grow for

250 hours. Minimal growth of cells lacking *CBR1* in media containing cellobiose as the sole

carbon source was detected after approximately one week (Fig 3C-E). The *bgl1Δ bgl2Δ* cells

were never able to grow when cellobiose was provided as the sole carbon source (Fig 3C-E).

When spent supernatant from wild type cells grown on cellobiose was added at 6%, 12%, or

25% the growth of wild type cells was unchanged. However, the *cbr1Δ* cells on cellobiose were

able to grow exponentially when cellobiose spent supernatant was added, while growth of the

bgl1Δ bgl2Δ cells began after approximately 5 days (Fig 3C and Fig S6B-C). Spent supernatant

from wildtype grown on succinate did not impact wildtype growth on cellobiose. However, the

growth of both the *cbr1Δ* and *bgl1Δ bgl2Δ* cells improved significantly upon addition of spent

succinate supernatant (Fig 3D and Fig S6D-E). The addition of 6%, 12%, or 25% spent

supernatant from wild type cells grown on glucose did not improve the growth of *cbr1Δ* with 6%

or 12% supernatant or *bgl1Δ bgl2Δ* cells with any percent supernatant on cellobiose media (Fig

3E and Fig S6F-G). Taken together, our data show that genes necessary for growth on

cellobiose but not TCA cycle intermediates are expressed in a *CBR1*-dependent fashion on both

cellobiose and TCA cycle intermediates.

Transporter genes expressed in a CBR1-dependent fashion in response to cellobiose and TCA

cycle intermediates are necessary for citrate utilization

Two genes encoding MFS transporters were expressed in a *CBR1*-dependent fashion in

response to both cellobiose and succinate: RTO4_13825 and RTO4_10339. We hypothesized

that these transporters may be responsible for growth on cellobiose and/or TCA cycle

intermediates. To test this hypothesis, we deleted RTO4_13825 and RTO4_10339 individually

and grew these strains on media containing glucose, cellobiose, citrate, succinate, α-

ketoglutarate, malate, and fumarate. The growth of cells lacking RTO4_13825 or RTO4_10339

was indistinguishable from that of wild type cells when glucose or cellobiose was provided as

the sole carbon source (Fig 4A). These data suggest that RTO4_13825 and RTO4_10339 are

not necessary for growth when cellobiose is provided as the sole carbon source.

Cells lacking RTO4_13825 were unable to grow when citrate was provided as the sole

carbon source (Fig 4B). However, RTO4_13825Δ cells were able to grow as well as wild type

cells when any of the dicarboxylic acid TCA cycle intermediates, succinate, α-ketoglutarate,

malate, or fumarate, were provided as the sole carbon source (Fig 4B). Thus, we named

RTO4_13825 *TCT1* for tricarboxylic acid transporter 1. In contrast, RTO4_10339 had no

significant growth phenotype on any of the carbon sources we tested.

A gene encoding a predicted phosphoglycerate mutase (RTO4_11229) was expressed in

a *CBR1*-dependent fashion in response to cellobiose and succinate. We hypothesized that it

may play a role in utilization of cellobiose or TCA cycle intermediates. However, the growth of

cells lacking RTO4_11229 in media containing glucose, cellobiose, citrate, succinate, α-

ketoglutarate, malate, and fumarate was indistinguishable from wild type cells (Fig S7).

Cbr1 inhibits carbon catabolite repression during disaccharide utilization

Our data suggest that *R. toruloides* β-glucosidases are secreted into the media where

they cleave the cellobiose β-1,4-glycosidic bond, releasing two glucose molecules (Fig 3). If this

is the case, cellobiose is likely imported into the cell as glucose molecules rather than cellobiose

molecules. Glucose is a preferred carbon source for most fungi and represses genes necessary

for utilization of nonpreferred carbon sources (1, 14, 15). We asked whether glucose was

preferred over cellobiose in *R. toruloides* by growing wild type cells in media with cellobiose as

the sole carbon source and adding the nonhydrolyzable glucose analog 2-deoxyglucose to the

media. Because 2-deoxyglucose cannot be hydrolyzed, it cannot act as a carbon source, but it

is still sensed as glucose. It was possible to completely repress the growth of wild type cells on

cellobiose by adding 0.001% 2-deoxyglucose (Fig 5A).

The extracellular release of glucose by the extracellular β-glucosidase activity of Bgl1

and Bgl2 begs the question of how *R. toruloides* avoids carbon catabolite repression during

cellobiose utilization. We hypothesized that *CBR1* may play a role in inhibiting carbon catabolite

repression. To test this hypothesis, we grew wild type cells and *cbr1Δ* cells in media with a

nonpreferred carbon source and 2-deoxyglucose. We first investigated the role of *CBR1* in

responding to the presence of 2-deoxyglucose in combination with the disaccharide maltose.

When 2-deoxyglucose was added at 0.002%, wild type cells were still able to grow but the

growth of cells lacking *CBR1* were completely inhibited (Fig 5B). A similar phenomenon was

seen when 2-deoxyglucose was added to media containing the disaccharide trehalose. When

0.01% 2-deoxyglucose was added to 1% trehalose the growth of *cbr1Δ* cells was reduced by

40% relative to wild type cells and when 0.02% 2-deoxyglucose was added, *cbr1Δ* cells were

unable to grow, while the growth of wild type cells was only moderately inhibited (Fig 5B).

Complementation of the *cbr1* deletion with *CBR1* placed under its native promoter at the *YKU70*

locus, restored the sensitivity to 2-deoxyglucose to levels indistinguishable from that of wild type

cells (Fig 5B).

In Ascomycete fungi, known regulators of carbon catabolite repression, such as CRE-

1/CreA/Mig1 regulate expression of genes responsible for utilization of all nonpreferred carbon

sources in response to preferred carbon sources (15–21). We hypothesized that carbon

catabolite repression would function similarly in Basidiomycete cells, and thus, *CBR1* would

inhibit carbon catabolite repression for all nonpreferred carbon sources. To test this hypothesis,

we tested the sensitivity of wild type and *cbr1Δ* cells in media containing other classes of

carbohydrates. To our surprise, wild type and *cbr1Δ* cells were equally sensitive to 2-

deoxyglucose when exposed to the monosaccharides fructose and xylose, the amino acid

proline, and the lipid Tween-80 (Fig 5D-G).

To test the role of *CBR1* in regulating carbon catabolite repression in response to

alcohols, we grew wild type and *cbr1Δ* cells in media containing glucose and allyl alcohol, which

is oxidized to the toxic compound acrolein by alcohol dehydrogenases. If carbon catabolite

repression is functioning normally, the presence of glucose should inhibit expression of alcohol

dehydrogenases in the presence of glucose (29). Wild type and *cbr1Δ* cells were equally

sensitive to allyl alcohol when glucose was present (Fig 5H). These data suggest that *CBR1*

inhibits carbon catabolite repression specifically when the available nonpreferred carbon source

is a disaccharide. We asked whether the *N. crassa* Cbr1 ortholog, CLR-2, also played a role in

inhibiting carbon catabolite repression by exposing wild type cells and *Δclr-2* cells to trehalose

or xylose with the addition of 2-deoxyglucose. The response to 2-deoxyglucose in wild type and

Δ*clr-2* cells was indistinguishable (Fig S8), indicating that the role of Cbr1 in inhibiting carbon

catabolite repression during exposure to disaccharides is an expanded role for Cbr1 relative to

the filamentous fungal ortholog CLR-2.

DISCUSSION

The regulation of gene expression in response to nutrient availability requires cells to

integrate information from activating and repressing environmental nutrient signals. There have

been limited investigations of the genetic regulators of cellulose and cellobiose utilization in

Basidiomycetes. A recent study identified Roc1, a regulator of cellulase-encoding genes in the

basidiomycete wood-decaying fungus *Schizophyllum commune* (30). However, homologs of

Roc1 do not exist outside of the Agaricomycetes (30). In Ascomycete filamentous fungi, CLR-

2/ClrB regulates cellulose degradation and utilization through the direct activation of genes

encoding cellulases (24, 31). We identified the CLR-2/ClrB ortholog Cbr1 in the oleaginous,

basidiomycete yeast *R. toruloides* (Fig 1A). Although Ascomycete and Basidiomycete fungi are

separated by approximately 400 million years of evolution (32), our study showed that Cbr1 has

a role in utilization of the cellulose breakdown product cellobiose (Fig 1). *N. crassa* CLR-2 does

not directly regulate the expression of genes encoding β-glucosidases, which are regulated by

CLR-1 (NCU07705) in *N. crassa* (24, 31). However, a role for Cbr1 in cellobiose utilization

suggests that there is conservation in the role of CLR-2/ClrB/Cbr1 between Ascomycete

filamentous fungi and Basidiomycete yeast.

Our characterization of Cbr1 showed that Cbr1 regulates utilization of TCA cycle

intermediates through the regulation of a gene encoding a transporter and regulates cellobiose

utilization through the regulation of genes encoding β-glucosidases (Fig 6). In other fungal

species, genes necessary for utilization of multiple carbon sources are also sometimes

regulated by a single transcription factor. For example, CLR-2 in *N. crassa* regulates cellulose

utilization, mannan utilization, and a small number of genes involved in hemicellulose utilization

(14, 31, 33). In *Trichoderma reesei*, *Penicillium oxalicum*, and some Aspergilli, the transcription

factor Xyr1/XlnR regulates utilization of both cellulose and hemicellulose (25–27). ARA-1/Ara1

regulates both arabinan and galactose utilization in *N. crassa* and *T. reesei* (14, 34), and FarA

regulates utilization of both fatty acids and the lignin component ferulic acid (35, 36). However,

in these examples linkage of these carbon sources can easily be explained either by the

presence of both carbon sources in the plant cell wall (i.e, cellulose and hemicellulose or

arabinan and galactose) or catabolism through the same metabolic pathways (i.e., fatty acids

and ferulic acid).

Our data does not support the hypothesis that cellobiose and TCA cycle intermediates

are catabolized through the same metabolic pathway in *R. toruloides*. The Cbr1-regulated β-

glucosidase genes are critical for cellobiose utilization but not TCA cycle intermediate utilization

(Fig 3), while one of the Cbr1-regulated transporters is critical for citrate utilization but not

cellobiose utilization (Fig 4). Aside from *CBR1* itself, we did not identify any genes that were

necessary for utilization of both cellobiose and TCA cycle intermediates. Thus, we speculate

that the coregulation of cellobiose and TCA cycle intermediates may suggest that *R. toruloides*

frequently encounters cellobiose and TCA cycle intermediates in close proximity. One possible

explanation for this is that *R. toruloides* scavenges cellobiose during cellulose degradation by

filamentous fungi, which also secrete organic acids, such as citrate, to acidify their environment

(37).

Carbon catabolite repression is the genetic mechanism through which fungal cells

prioritize utilization of carbon sources that require less energy to metabolize over carbon

sources that require more energy to metabolize (1). Known genetic mechanisms of regulating

carbon catabolite repression regulate the expression of genes required for utilization of all less

preferred carbon sources when a more preferred carbon source is present (15–21). Cbr1

demonstrates a mechanism through which cells inhibit carbon catabolite repression specifically

during utilization of disaccharides (Fig 5). We speculate that this may allow cells to continue

expressing genes necessary for hydrolyzing disaccharide sugar bonds even when this

hydrolysis produces preferred monosaccharides like glucose (Fig 6). Genetic linkage of carbon

catabolite repression inhibition with the activation of genes necessary to utilize a nonpreferred

carbon source has not previously been described. Additionally, regulation of genes involved in

utilization of a specific class of nonpreferred carbon sources in response to a preferred carbon

source represents a novel form of carbon catabolite repression regulation.

MATERIALS and METHODS

Strains and culturing

R. toruloides strains used in this study are listed in Table S1. All strains were derived

from the wild type reference strain IFO 0880 (also called NBRC 0880, obtained from Biological

Resource Center, NITE (NBRC), Japan). The starting strain for genetic manipulations was in a

non-homologous end-joining deficient *yku70Δ* background (38). *R. toruloides* was transformed

using *Agrobacterium tumefaciens*-mediated transformation using *A. tumefaciens* EHA 105 and

plasmids derived from pGI2 (39) that were previously used for *R. toruloides A. tumefaciens*-

mediated transformation (11). *A. tumefaciens*-mediated transformation was performed as

previously described (11). Gene deletions, complementations, and tags were performed using

homologous recombination of a nourseothricin or G418 resistance cassette with flanking arms

of ~1000 bp. All strains were confirmed by PCR. Complement and tagged strains were further

confirmed by DNA sequencing. We fluorescently tagged the histone H2A subunit gene *HTA1*

(RTO4_12090) with the mRuby2 fluorescent protein at the native locus. The Hta1-mRuby2

tagged strains were generated using a combination of G418 resistance and a 5-

fluorodeoxyuridine (FUDR) sensitive thymidine kinase counterselection marker. The Hta1-

mRuby2 construct was transformed into the *yku70Δ CBR1-GFP* backgrounds using G418 to

select for transformants. Transformed colonies were passaged one time on liquid YPD and

plated onto YNB + 1% glucose + FUDR plates to select for loop-outs of the G418 and thymidine

kinase markers. Colonies were screened by microscopy, PCR, and locus sequencing for correct

integration of the mRuby2 tag and successful removal of the G418 and thymidine kinase

markers. As RTO4_16716 and RTO4_16717 are adjacent in the genome, the *bgl1Δ bgl2Δ* strain

was generated via single transformation of the nourseothricin resistance cassette to remove

both open reading frames from the genome. *R. toruloides* strains were grown from freezer

stocks on YPD (VWR) plates at 28^{o}C for 2 d to 3 d prior to inoculation into liquid YPD to grow

strains at 28^{o}C with constant shaking to exponential phase growth prior to starting experiments.

N. crassa strains used in this study are listed in Table S2. All strains were derived from

the wild type reference strain FGSC 2489 using standard genetic techniques and confirmed

using PCR (40, 41). *N. crassa* cells were grown from freezer stocks on Vogel's minimal medium

(42) + 2% sucrose + 1.5% agar (Bacto) slants for 2 d at 28^{o}C in the dark and 4 d at 28^{o}C in

constant light. *N. crassa* conidia were harvested from slants with sterile double distilled H_2O for

inoculation.

All *R. toruloides* experiments were done using yeast nitrogen base (YNB) with NH_4SO_4

without amino acids (MP Biomedicals) + 12 mM K_2HPO_4 (Fisher) + 38 mM KH_2PO_4 (Fisher) +

100 nM Fe(II)SO_4 pH 6.2 with the indicated carbon source at 1% w/v with the exception of

coumaric acid and ferulic acid, which were added at a concentration of 0.2%, and citrate which

was added at a concentration of 8.5 mM and a pH of 5.5. All *N. crassa* experiments were

performed using Vogel's minimal medium with 50mM NH_4Cl instead of 25mM NH_4NO_3 (VMM

NH_4Cl) and the indicated carbon source at 1% w/v. Carbon sources used are listed in Table S3.

R. toruloides growth experiments

Colonies of *R. toruloides* grown on solid YPD were inoculated into liquid YPD and grown

at 28^{o}C with constant shaking until exponential phase. Cells were pelleted at 4000 x g and

washed three times with YNB lacking a carbon source. Washed cells were used to inoculate

750 µL of media in 48-well plates at an OD 600 of 0.1 for initial carbon screening experiments

(Figure S2) and 0.01 for all subsequent growth experiments. Growth assays were incubated at

28ºC with constant shaking (3 mm orbit, 600 rpm for individual time points and 3 mm orbit 425

rpm for kinetic growth experiments). Kinetic growth experiments were performed using an

Agilent BioTek Epoch2 plate reader with OD 600 measurements taken every 20 or 30 minutes

depending on the experiment.

N. crassa growth experiments

For trehalose and xylose 2-deoxyglucose (TCI) flask growth experiments wild type *mat A*

(FGSC 2489) and Δ*clr2 mat A* conidia were inoculated at 10^6 conidia/mL into flasks with 100 mL

of filter sterilized VMM NH_4Cl with either 1% xylose or 1% trehalose. Filter sterilized 20% 2-

deoxyglucose dissolved in water was added to the 100mL media to achieve xylose + 2-

deoxyglucose concentrations of either 0%, 0.00195%, 0.00391%, or 0.0078125% and trehalose

+ 2-deoxyglucose concentrations of either 0%, 0.04%, 0.0625%, or 0.125%. Cultures were

grown at 28ºC with constant light and shaking at 200rpm. After 24 hours biomass was

autoclaved, harvested, dried for 24 h at 60ºC in a drying oven, and weighed.

For growth of *N. crassa* on TCA cycle intermediates and glucose, wild type mat A (FGSC

2489) and Δ*clr2 mat A* conidia inoculated at 10^6 conidia/mL in 24 deep-well plates with 3 mL of

filter sterilized VMM NH_4Cl with 1% succinate, 1% alpha-ketoglutarate, 1% malate, or 1%

glucose. Cultures were incubated at 28ºC with constant light and shaking (200 rpm). Biomass

from glucose cultures was harvested at 18 h post inoculation and biomass from TCA cycle

intermediate cultures was harvested 7 d post inoculation. Biomass was dried for 24 h in a drying

oven at 60ºC and weighed.

Phylogenetic tree generation

Protein sequences for CLR-1, CLR-2, and XLR-1, from *N. crassa* (OR74A) and ClrA,

ClrB, and XlnR from *Aspergillus nidulans* (A4) were obtained from Mycocosm (43). The protein

sequences were used to BLAST search the proteomes of *N. crassa*, *A. nidulans*,

Saccharomyces cerevisiae (S288C)*,* and *R. toruloides* (IFO 0880). Identified homologs with an

e-value < 1E-10 were aligned via MUSCLE (44) in MEGA11 (45) and the phylogenetic tree was

generated using FastTree (46) using the maximum likelihood method based on the Jones-

Taylor-Thornton matrix-based model and visualized in MEGA11. The reliability of each split in

the tree was computed using the Shimodaira-Hasegawa test on three alternate topologies

around that split.

Microscopy

Fluorescent microscopy was performed using a Nikon Eclipse Ti2 inverted microscope

equipped with a prime BSI express CMOS camera using a CFI Plan Apochromat 100X oil

objective. *HTA1-mRuby2 CBR1-GFP* strains were grown to exponential phase in liquid YPD.

Cells were pelleted at 4000 x g and washed three times with YNB without a carbon source.

Cells were inoculated into a 48-well plate with YNB + 1% w/v of the indicated carbon source at

an OD 600 of 0.2 and incubated at 28ºC with shaking (3mm orbit, 600rpm) for 4 hours. Aliquots

of the cultures were placed onto agarose coated slides and imaged.

RNA sequencing and transcript abundance

Cells were grown to exponential phase in liquid YPD and washed three times in YNB

without a carbon source. Flasks containing 100 mL of YNB + 1% cellobiose, 1% succinate, or

no carbon source were inoculated at an OD 600 of 0.1. Cultures were incubated for 8 hours at

28ºC with shaking (200 rpm). Cells were harvested using centrifugation in 50 mL conical tubes

at 4000 x g. The supernatants were removed, and the cell pellets were flash frozen in liquid

nitrogen. Pellets were stored at -80ºC prior to RNA extraction.

RNA was extracted using the Quick-RNA Fungal/Bacterial Miniprep Kit according to the

manufacturer's instructions. RNA quality was checked via agarose gel electrophoresis and

BioAnalyzer analysis performed at the Transcriptional Regulation & Expression Facility at

Cornell University. RNA was submitted to the California Institute for Quantitative Biosciences at

UC Berkeley (QB3-Berkeley) for library preparation and sequencing. Libraries were prepared

using standard Illumina protocols and RNA was sequenced using an Illumina NextSeq 2000

150PE at a depth of approximately 20 million reads per sample.

The transcript abundance (transcripts per million, TPM) was quantified using Salmon v.

1.10.0 mapping to the *R. toruloides* IFO 0880 genome (v4) (11) using the –validateMappings

setting (47). Differential expression was determined using DESeq2 v. 1.44.0 (48). Genes were

denoted as differentially expressed if there was at least a 4-fold change in expression and the

average TPM across all samples was at least 5. RNAseq data used in this study were deposited

in the Gene Expression Omnibus (GEO) at the National Center for Biotechnology Information

(NCBI) and are accessible through GEO series accession number GSE293943. Expression

data was visualized using the ComplexHeatmap R package (49).

β-glucosidase enzyme activity assays

R. toruloides wild type, *cbr1∆*, and *bgl1∆ bgl2∆* strains were struck out from freezer

stocks on 2% agar yeast extract peptone dextrose (YPD) plates from freezer stocks and

incubated for five days at 28ºC with constant light. Three independent colonies of each strain

were inoculated into flasks with 100 mL of YNB with 1% glucose and grown for 23 h at 28ºC in

constant light with shaking at 200 rpm. Cultures were transferred to 50mL conical tubes, and the

cells were pelleted at 4000 x g. Supernatants were discarded and pellets were resuspended in 5

mL YNB 1% glucose media. Each resuspended culture was then transferred into 3 independent

flasks with 100 mL YNB 1% glucose media at 0.004 OD/mL and cultures were grown for 8

generations at 28ºC, constant light, and shaking at 200 rpm. Cultures were transferred to 50mL

conical tubes, and the cells were pelleted at 4000 x g. Supernatants were discarded and pellets

were washed twice in 5mL YNB media without carbon. Cultures were transferred into YNB 1%

glucose, succinate, or cellobiose at 0.8 OD/mL and grown at 28ºC in constant light with shaking

at 200rpm for 20 h prior to measuring optical density. An aliquot from each culture was

centrifuged at 10,000 x g for 1 m and the culture supernatant was filtere sterilized with a 0.22μm

syringe filter and stored at -80ºC. Filtered supernatant aliquots were thawed on ice and β-

glucosidase activity was assessed by adding 50 uL of 100 mM sodium acetate (pH 5) and 25 uL

of 20 mM p-nitrophenyl-β-D-glucopyranoside (EMD Millipore Corp.) in a 96-well plate. The plate

was incubated on ice for five minutes and 25 uL of each supernatant aliquot was added, leaving

a set without supernatant added for a blank reading, and then incubated at 37ºC for 30 m.

Reactions were quenched by adding 100 uL of 200 mM Na_2CO_3 and 25 uL of each supernatant

aliquot was added to quenched blank wells. Optical densities were measured at 400 nm, and

blank readings were subtracted from their respective assay readings.

Spent media growth assays

The *R. toruloides* wild type strain was struck out from freezer stocks on a 2% agar YPD

plate and incubated at 28ºC for two days with constant light. Single colonies were inoculated

into 15 14 mL culture tubes with 5 mL YPD. The cultures were grown overnight at 28ºC with

constant light and shaking at 250rpm. The following day, five 5 mL cultures were combined into

sterile 50 mL conical tubes for each of three replicates. 25 mL cultures were spun down at 4000

x g and washed three times with YNB no carbon. Cells were inoculated at a density of 0.01

OD/mL into flasks with 100 mL of either YNB 1% glucose, 1% cellobiose, or 1% succinate.

Cultures were grown to an OD 600 of 1, 1.1, or 0.9, for glucose, cellobiose, and succinate,

respectively, prior to centrifugation in sterile 50 mL conicals at 4000 x g for 10 minutes.

The supernatants were sterilized with tube top vacuum 0.22 μm filters. Filter sterilized

supernatants were used the same day to grow wild type, *cbr1Δ*, and *bgl1Δ bgl2Δ* strains at 0.01

OD/mL in YNB with 1% cellobiose and YNB lacking a carbon source with spent, filtered

supernatant added to make up 6%, 12%, or 25% of the total 750 μL culture in 48-well plates.

OD 600 measurements of the 48-well plates were taken as indicated. β-glucosidase activity of

each supernatant was assessed according to β-glucosidase assay protocol above.

Statistical significance tests

All statistical analyses were performed in Rstudio (v2024.12.1; R v4.4.3). All experiments

were performed with a minimum of three biological replicates.

Functional enrichment analysis

Functional annotations for the *R. toruloides* IFO 0880 genome (v4) were downloaded

from Mycocosm

(https://mycocosm.jgi.doe.gov/Rhoto_IFO0880_4/Rhoto_IFO0880_4.home.html). Functional

category enrichment analyses were performed using the compareCluster function in the

clusterProfiler R package with function set to "enricher" (50). Differentially expressed genes

were compared for enrichment against all genes expressed at an average TPM of 5 or greater

across the experiment. The Benjamini-Hochberg procedure was used to correct for multiple

hypothesis testing. Data were visualized using the clusterProfiler package's dotplot function

(50).

ACKNOWLEDGEMENTS

This work was supported by a grant from the National Institute of General Medical

Sciences of the National Institutes of Health (NIH) under Award Number R35GM150926 to

L.B.H. J.D.K. was supported by NIH Ruth L. Kirschstein National Research Service Award

Institutional Research Training Grant 5T32AI145821-03.

REFERENCES

1. J. D. Kerkaert, L. B. Huberman, Regulation of nutrient utilization in filamentous fungi. *Appl.*

Microbiol. Biotechnol. **107**, 5873–5898 (2023).

2. L. B. Huberman, J. Liu, L. Qin, N. L. Glass, Regulation of the lignocellulolytic response in

filamentous fungi. *Fungal Biol. Rev.* **30**, 101–111 (2016).

3. J. Kim, *et al.*, Multi-Omics Driven Metabolic Network Reconstruction and Analysis of

Lignocellulosic Carbon Utilization in Rhodosporidium toruloides. *Front. Bioeng. Biotechnol.*

8, 612832 (2021).

4. J. Yaegashi, *et al.*, Rhodosporidium toruloides: a new platform organism for conversion of

lignocellulose into terpene biofuels and bioproducts. *Biotechnol. Biofuels* **10**, 241 (2017).

5. M. G. Wiebe, K. Koivuranta, M. Penttilä, L. Ruohonen, Lipid production in batch and fed-

batch cultures of Rhodosporidium toruloidesfrom 5 and 6 carbon carbohydrates. *BMC*

Biotechnol. **12**, 26 (2012).

6. S. Wu, C. Hu, G. Jin, X. Zhao, Z. K. Zhao, Phosphate-limitation mediated lipid production by

Rhodosporidium toruloides. *Bioresour. Technol.* **101**, 6124–6129 (2010).

7. S. Wu, X. Zhao, H. Shen, Q. Wang, Z. K. Zhao, Microbial lipid production by

Rhodosporidium toruloides under sulfate-limited conditions. *Bioresour. Technol.* **102**, 1803–

1807 (2011).

8.   Z. Zhu, *et al.*, A multi-omic map of the lipid-producing yeast Rhodosporidium toruloides. *Nat.*

Commun. **3**, 1112 (2012).

9.   J. J. L. Lee, L. Chen, J. Shi, A. Trzcinski, W.-N. Chen, Metabolomic Profiling of

Rhodosporidium toruloides Grown on Glycerol for Carotenoid Production during Different

Growth Phases. *J. Agric. Food Chem.* **62**, 10203–10209 (2014).

10. L. C. Mata-Gómez, J. C. Montañez, A. Méndez-Zavala, C. N. Aguilar, Biotechnological

production of carotenoids by yeasts: an overview. *Microb. Cell Factories* **13**, 12 (2014).

11. S. T. Coradetti, *et al.*, Functional genomics of lipid metabolism in the oleaginous yeast

Rhodosporidium toruloides. *eLife* **7**, e32110 (2018).

12. Y.-K. Park, J.-M. Nicaud, R. Ledesma-Amaro, The Engineering Potential of Rhodosporidium

toruloides as a Workhorse for Biotechnological Applications. *Trends Biotechnol.* **36**, 304–

317 (2018).

13. T. Benocci, M. V. Aguilar-Pontes, M. Zhou, B. Seiboth, R. P. De Vries, Regulators of plant

biomass degradation in ascomycetous fungi. *Biotechnol. Biofuels* **10**, 152 (2017).

14. V. W. Wu, *et al.*, The regulatory and transcriptional landscape associated with carbon

utilization in a filamentous fungus. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 6003–6013 (2020).

15. M. Adnan, *et al.*, Carbon Catabolite Repression in Filamentous Fungi. *Int. J. Mol. Sci.* **19**, 48

(2017).

16. J. O. Nehlin, H. Ronne, Yeast MIG1 repressor is related to the mammalian early growth

response and Wilms' tumour finger proteins. *EMBO J.* **9**, 2891–2898 (1990).

17. H. N. Arst, D. J. Cove, Nitrogen metabolite repression in Aspergillus nidulans. *Mol. Gen.*

Genet. MGG **126**, 111–141 (1973).

18. C. E. A. Dowzer, J. M. Kelly, Analysis of the *creA* Gene, a Regulator of Carbon Catabolite

Repression in *Aspergillus nidulans*. *Mol. Cell. Biol.* **11**, 5701–5709 (1991).

19. Y. Hong, *et al.*, Carbon catabolite repressor MoCreA is required for the asexual development

and pathogenicity of the rice blast fungus. *Fungal Genet. Biol.* **146**, 103496 (2021).

20. J. Strauss, *et al.*, Crel, the carbon catabolite repressor protein from *Trichoderma reesei*.

FEBS Lett. **376**, 103–107 (1995).

21. J. Sun, N. L. Glass, Identification of the CRE-1 Cellulolytic Regulon in Neurospora crassa.

PLoS ONE **6**, e25654 (2011).

22. S. Rongpipi, D. Ye, E. D. Gomez, E. W. Gomez, Progress and Opportunities in the

Characterization of Cellulose – An Important Regulator of Cell Wall Growth and Mechanics.

Front. Plant Sci. **9**, 1894 (2019).

23. H. V. Dinh, *et al.*, A comprehensive genome-scale model for Rhodosporidium toruloides

IFO0880 accounting for functional genomics and phenotypic data. *Metab. Eng. Commun.* **9**,

e00101 (2019).

24. S. T. Coradetti, *et al.*, Conserved and essential transcription factors for cellulase gene

expression in ascomycete fungi. *Proc. Natl. Acad. Sci.* **109**, 7397–7402 (2012).

25. A. R. Stricker, K. Grosstessner-Hain, E. Würleitner, R. L. Mach, Xyr1 (Xylanase Regulator 1)

Regulates both the Hydrolytic Enzyme System and D -Xylose Metabolism in *Hypocrea*

jecorina. *Eukaryot. Cell* **5**, 2128–2137 (2006).

26. Z. Li, *et al.*, Synergistic and Dose-Controlled Regulation of Cellulase Gene Expression in

Penicillium oxalicum. *PLOS Genet.* **11**, e1005509 (2015).

27. N. N. M. E. Van Peij, M. M. C. Gielkens, R. P. De Vries, J. Visser, L. H. De Graaff, The

Transcriptional Activator XlnR Regulates Both Xylanolytic and Endoglucanase Gene

Expression in *Aspergillus niger*. *Appl. Environ. Microbiol.* **64**, 3615–3619 (1998).

28. P. J. Bhat, T. V. S. Murthy, Transcriptional control of the *GAL/MEL* regulon of yeast

Saccharomyces cerevisiae : mechanism of galactose-mediated signal transduction. *Mol.*

Microbiol. **40**, 1059–1066 (2001).

29. R. Megnet, Mutants partially deficient in alcohol dehydrogenase in Schizosaccharomyces

pombe. *Arch. Biochem. Biophys.* **121**, 194–201 (1967).

30. I. M. Marian, *et al.*, The Transcription Factor Roc1 Is a Key Regulator of Cellulose

Degradation in the Wood-Decaying Mushroom *Schizophyllum commune*. *mBio* **13**, e00628-

22 (2022).

31. J. P. Craig, S. T. Coradetti, T. L. Starr, N. L. Glass, Direct Target Network of the Neurospora

crassa Plant Cell Wall Deconstruction Regulators CLR-1, CLR-2, and XLR-1. *mBio* **6**,

e01452-15 (2015).

32. J. W. Taylor, M. L. Berbee, Dating divergences in the Fungal Tree of Life: review and new

analyses. *Mycologia* **98**, 838–849 (2006).

33. A. Samal, *et al.*, Network reconstruction and systems analysis of plant cell wall

deconstruction by Neurospora crassa. *Biotechnol. Biofuels* **10**, 225 (2017).

34. T. Benocci, *et al.*, ARA 1 regulates not only L -arabinose but also D -galactose catabolism in

Trichoderma reesei. *FEBS Lett.* **592**, 60–70 (2018).

35. M. J. Hynes, S. L. Murray, A. Duncan, G. S. Khew, M. A. Davis, Regulatory Genes

Controlling Fatty Acid Catabolism and Peroxisomal Functions in the Filamentous Fungus

Aspergillus nidulans. *Eukaryot. Cell* **5**, 794–805 (2006).

36. M. Arentshorst, *et al.*, Utilization of ferulic acid in Aspergillus niger requires the transcription

factor FarA and a newly identified Far-like protein (FarD) that lacks the canonical

Zn(II)2Cys6 domain. *Front. Fungal Biol.* **3**, 978845 (2022).

37. P. L. Show, *et al.*, Overview of citric acid production from *Aspergillus niger*. *Front. Life Sci.* **8**,

271–283 (2015).

38. S. Zhang, M. Ito, J. M. Skerker, A. P. Arkin, C. V. Rao, Metabolic engineering of the

oleaginous yeast Rhodosporidium toruloides IFO0880 for lipid overproduction during high-

density fermentation. *Appl. Microbiol. Biotechnol.* **100**, 9393–9405 (2016).

39. E. P. Abbott, G. Ianiri, R. Castoria, A. Idnurm, Overcoming recalcitrant transformation and

gene manipulation in Pucciniomycotina yeasts. *Appl. Microbiol. Biotechnol.* **97**, 283–295

(2013).

40. H. V. Colot, *et al.*, A high-throughput gene knockout procedure for *Neurospora* reveals

functions for multiple transcription factors. *Proc. Natl. Acad. Sci.* **103**, 10352–10357 (2006).

41. K. McCluskey, A. Wiest, M. Plamann, The Fungal Genetics Stock Center: a repository for 50

years of fungal genetics research. *J. Biosci.* **35**, 119–126 (2010).

42. H. J. Vogel, A convenient growth medium for Neurospora (medium N). *Microb. Genet Bull*

13, 42–43 (1956).

43. I. V. Grigoriev, *et al.*, MycoCosm portal: gearing up for 1000 fungal genomes. *Nucleic Acids*

Res. **42**, D699–D704 (2014).

44. R. C. Edgar, MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).

45. K. Tamura, G. Stecher, S. Kumar, MEGA11: Molecular Evolutionary Genetics Analysis

Version 11. *Mol. Biol. Evol.* **38**, 3022–3027 (2021).

46. M. N. Price, P. S. Dehal, A. P. Arkin, FastTree 2 – Approximately Maximum-Likelihood Trees

for Large Alignments. *PLoS ONE* **5**, e9490 (2010).

47. R. Patro, G. Duggal, M. I. Love, R. A. Irizarry, C. Kingsford, Salmon provides fast and bias-

aware quantification of transcript expression. *Nat. Methods* **14**, 417–419 (2017).

48. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

49. Z. Gu, Complex heatmap visualization. *iMeta* **1**, e43 (2022).

50. S. Xu, *et al.*, Using clusterProfiler to characterize multiomics data. *Nat. Protoc.* **19**, 3292–

3320 (2024).

 FIGURES

Fig 1. *CBR1* **regulates utilization of cellobiose and TCA cycle intermediates. (A)** Protein

sequences of *N. crassa* CLR-2 and *Aspergillus nidulans* ClrB homologs were used to build a

phylogenetic tree using the maximum likelihood method based on the Jones-Taylor-Thornton

matrix-based model using FastTree (46). The reliability of each split in the tree was computed

using the Shimodaira-Hasegawa test on three alternate topologies around that split. *R.*

toruloides proteins are indicated with maroon diamonds. *N. crassa* proteins are indicated with

orange squares. *A. nidulans* proteins are indicated with green circles. *Saccharomyces*

cerevisiae proteins are indicated with blue triangles. **(B-C)** Growth curves (OD 600) of wild type,

cbr1Δ, and *cbr1Δ+CBR1* complement strains grown in YNB with the indicated carbon source at

8.5 mM for citrate or 1% w/v for all other carbon sources. Dots are the average and colored bars

indicate standard deviation of three biological replicates. *** p < 1E-10 *cbr1Δ* vs WT and *cbr1Δ*

vs *cbr1Δ+CBR1* as determined by two-way ANOVA with a TukeyHSD post-hoc test.

Fig 2. Cbr1 activates five genes in response to cellobiose and TCA cycle intermediates.

(A) Heatmap of the expression of genes at least 4-fold differentially expressed between at least

one of the following comparisons: wild type (WT) cells exposed to cellobiose as compared to

carbon starvation (no carbon), wild type cells exposed to succinate as compared to carbon

starvation, *cbr1Δ* cells as compared to wild type exposed to cellobiose, *cbr1Δ* cells as compared

to wild type exposed to succinate, or *cbr1Δ* cells as compared to wild type exposed to carbon

starvation. Genes were normalized by their average expression across all conditions, log

transformed, and clustered using the Euclidean distance metric. Pink bars indicate genes that

are at least 4-fold differentially expressed in wild type cells exposed to cellobiose as compared

to carbon starvation (NC). Orange bars indicate genes that are at least 4-fold differentially

expressed in wild type cells exposed to succinate as compared to carbon starvation (NC). Blue

bars indicate genes that are at least 4-fold differentially expressed in *cbr1Δ* as compared to wild

type cells exposed to carbon starvation (NC). Black bars indicate genes that are at least 4-fold

differentially expressed in *cbr1Δ* as compared to wild type cells exposed to succinate. Green

bars indicate genes that are at least 4-fold differentially expressed in *cbr1Δ* as compared to wild

type cells exposed to cellobiose. **(B-C)** Genes significantly increased or decreased in

expression for each comparison were tested for gene ontology (GO) **(B)** and KEGG **(C)**

functional category enrichment. Comparisons were only displayed if they had at least one

statistically significantly enriched category for each analysis. Dot size corresponds to the

number of genes differentially expressed for a category divided by total genes in that category.

Color indicates p-value as determined by clusterProfiler (50) with a Benjamini-Hochberg multiple

comparisons correction. **(D)** Heatmap of the expression of genes differentially expressed by at

least 4-fold in *cbr1Δ* as compared to wild type cells exposed to cellobiose and exposed to

succinate. Genes were normalized by their average expression across all conditions, log

transformed, and clustered using the Euclidean distance metric. Gene IDs and annotations are

indicated to the right of the heatmap. The expression of *CBR1* was left out of the heatmaps,

GO, and KEGG enrichment analysis, since the change in expression of this gene was due to its

deletion in the *cbr1Δ* strain.

Fig 3. Cbr1-dependent expression of secreted β-glucosidase genes is necessary for

cellobiose utilization. (A) Growth curves of wild type and *bgl1Δ bgl2Δ* cells grown in YNB with

the indicated carbon source at 8.5 mM for citrate or 1% w/v for all other carbon sources. Dots

are the average and colored bars indicate standard deviation of three biological replicates. *** p

< 1E-10 *bgl1Δ bgl2Δ* vs wild type as determined by two-way ANOVA with a TukeyHSD post-hoc

test. **(B)** β-glucosidase activity secreted into the supernatant of cultures of the indicated strain

grown in YNB with the indicated carbon source at 1% w/v for 20 h relative to the β-glucosidase

activity of wild type cells grown in cellobiose for 20 h. Bars are the averages of three biological

replicates (dots). *** p < 0.001 as determined by two-way ANOVA with a TukeyHSD post-hoc

test. **(C-E)** Growth (OD 600) of the indicated strains in YNB with 1% cellobiose supplemented

with the indicated percentage of spent supernatant from wild type cells grown in YNB with 1%

cellobiose **(C)**, 1% succinate **(D)**, or 1% glucose **(E)** for 6-7 generations. To normalize for the

771   growth of the indicated strain on the spent supernatant alone, the OD 600 of the indicated strain

772   grown on the indicated amount of spent supernatant alone was subtracted. Growth curves

773   without growth on spent supernatant alone subtracted, the growth curves on spent supernatant

774   alone, and the β-glucosidase activity of the spent supernatant are in Fig S6. Lines are the

775   averages of at least 3 biological replicates (dots).

Fig 4. Cbr1-dependent expression of transporter genes is necessary for TCA cycle

intermediate utilization. (A) Growth of the indicated strains in YNB with 1% glucose or 1%

cellobiose. Lines are the averages of three biological replicates (dots). **(B)** Growth of the

indicated strains in YNB with 8.5 mM citrate or 1% w/v of the indicated carbon source. For

growth on 8.5 mM citrate, dots are the average and colored bars are the standard deviation of

three biological replicates. For growth on all other carbon sources, lines are the average of three

biological replicates (dots). *** p < 1E-10 *tct1Δ* vs WT as determined by two-way ANOVA with a

TukeyHSD post-hoc test.

Fig 5. Cbr1 inhibits carbon catabolite repression during disaccharide utilization. (A)

Growth (OD 600) of wild type in YNB + 1% cellobiose supplemented with the indicated

concentration of 2-deoxyglucose. **(B)** Growth after 4 days of the indicated strain in YNB + 1%

maltose supplemented with the indicated concentration of 2-deoxyglucose relative to growth

after 4 days of the indicated strain in YNB + 1% maltose. **(C)** Growth after 6 days of the

indicated strain in YNB + 1% trehalose supplemented with the indicated concentration of 2-

deoxyglucose relative to growth after 6 days of the indicated strain in YNB + 1% trehalose. **(D)**

Growth after 4 days of the indicated strain in YNB + 1% fructose supplemented with the

indicated concentration of 2-deoxyglucose relative to growth after 4 days of the indicated strain

in YNB + 1% fructose. **(E)** Growth after 6 days of the indicated strain in YNB + 1% xylose

supplemented with the indicated concentration of 2-deoxyglucose relative to growth after 6 days

of the indicated strain in YNB + 1% xylose. **(F)** Growth after 5 days of the indicated strain in

YNB + 1% proline supplemented with the indicated concentration of 2-deoxyglucose relative to

growth after 5 days of the indicated strain in YNB + 1% proline. **(G)** Growth after 7 days of the

indicated strain in YNB + 1% Tween-80 supplemented with the indicated concentration of 2-

deoxyglucose relative to growth after 7 days of the indicated strain in YNB + 1% Tween-80. **(H)**

Growth after 3 days of the indicated strain in YNB + 1% glucose supplemented with the

indicated concentration of allyl alcohol relative to growth after 3 days of the indicated strain in

YNB + 1% glucose. *** $p < 1E-9$ *cbr1*Δ vs WT and *cbr1*Δ vs *cbr1*Δ+*CBR1* as determined by two-

way ANOVA with a TukeyHSD post-hoc test.

Fig 6. Model for Cbr1-mediated regulation of cellobiose utilization, TCA cycle

intermediate utilization, and carbon catabolite repression. Cbr1 is activated by the

presence of cellobiose and TCA cycle intermediates. Activated Cbr1 activates the expression of

genes encoding β-glucosidases, transporters, and a phosphoglycerate mutase and represses

the expression of genes encoding a chitin deacetylase, methyltransferase, cyclin, metallo-β-

lactamase, and several hypothetical proteins. The Cbr1-regulated transporter *TCT1* is required

for citrate utilization. The two β-glucosidase genes *BGL1* and *BGL2* encode secreted β-

glucosidases that cleave cellobiose into two glucose molecules. These glucose molecules

activate carbon catabolite repression, which likely represses the expression of genes necessary

to utilize cellobiose and TCA cycle intermediates. Cbr1 plays a role in inhibiting carbon

catabolite repression, specifically when a disaccharide is present as the nonpreferred carbon

source. Solid lines indicate a genetic interaction for which direct genetic or biochemical data

exists. Dotted lines indicate a genetic interaction for which indirect genetic data exists.

822
823