Carbohydrate utilization regulator reveals a noncanonical mechanism of nutrient

differentiation

Brandon Reyes-Chavez[1,2,#], Joshua D. Kerkaert[1,#], and Lori B. Huberman[1,*]

[1]Plant Pathology and Plant-Microbe Biology Section, School of Integrative Plant Science,

Cornell University, Ithaca, NY, USA 14853

[2]Microbiology Department, Cornell University, Ithaca, NY, USA 14853

[#]These authors contributed equally to this work.

*To whom correspondence may be addressed: huberman@cornell.edu

Key words: Carbon catabolite repression, fungi, oleaginous yeast, nutrient sensing, metabolic

regulation

ABSTRACT

Cells must sense and respond to nutrients to survive. To efficiently grow in mixed carbon

environments, microbes repress genes necessary to utilize carbon sources that require

substantial resources to catabolize when a simpler carbon source, such as glucose, is present.

This process is known as carbon catabolite repression. Canonically, in fungi, nutrient sensing

transcriptional networks are composed of carbon source-specific transcription factors that

activate carbon source utilization genes and carbon catabolite repression regulators, which

broadly repress all nonpreferred carbon source utilization genes when a preferred carbohydrate

is present. In contrast to this model, we identified a transcription factor (Cbr1) in the

basidomycete yeast *Rhodotorula* (*Rhodosporidium*) *toruloides* that specifically inhibits glucose-

mediated repression of glucose-glucose disaccharide utilization, presenting a mechanism of

tailored carbon catabolite repression regulation that combats a negative feedback loop formed

when glucose is released during disaccharide utilization. Cbr1 is also required for cellobiose,

carboxylic acid, and fucose utilization. Using transcriptomic and molecular analyses, we

demonstrated that catabolism of these carbon sources is not metabolically linked, but genes

necessary for their utilization are coactivated by Cbr1 in response to each of the carbon

sources. This coactivation suggests *R. toruloides* may encounter these carbon sources

together, potentially during complex interactions among microbes in nature. Coregulation of

nutrient-specific gene activation and carbon catabolite repression by a transcription factor

establishes a previously uncharacterized mechanism for building nutrient sensing transcriptional

networks in fungi. Characterizing diverse nutrient sensing regulatory mechanisms is critical for

understanding resource acquisition during fungal pathogenesis, where carbon catabolite

repression is important for virulence and drug tolerance, and metabolically engineering fungi for

green biotechnology.

INTRODUCTION

Cells must sense and respond to nutrients to successfully grow and divide using the resources available. For fungi, nutrient sensing is critical to establish fungal colonies, compete with other microbes, and establish pathogenic and symbiotic relationships with plants and animals. Most work characterizing genetic pathways regulating carbon source utilization in fungi has been performed in the Ascomycete phylum [1, 2]. In these fungi, a network of carbon source-specific transcription factors activates genes necessary for utilization of available carbon [3]. However, in a mixed carbon environment, it is not necessarily energetically optimal to simultaneously consume all available carbon sources. Thus, when a more preferred carbon source, such as glucose, is available, a process known as carbon catabolite repression broadly downregulates expression of genes required for utilization of all less preferred carbon sources [1, 3-10].

Many devastating human, animal, and plant pathogens fall outside of the Ascomycete phylum [11]. Genetic mechanisms through which these fungi activate genes necessary to utilize nonpreferred carbon sources and repress these genes in the presence of more preferred carbon sources are significantly less well studied. Basidiomycetes diverged from Ascomycetes approximately 400 million years ago and represent over 30% of described fungi [12, 13]. Recent studies in Basidiomycete pathogens have demonstrated that understanding the regulation of carbon utilization in this phylum is critical, since accurately regulating carbon metabolism is not only important for virulence but also changes the efficacy of antifungal drug treatments [14-16].

To address this knowledge gap, we investigated genetic mechanisms regulating carbon utilization in the basidiomycete yeast *Rhodotorula toruloides* (previously *Rhodosporidium toruloides*). Recent studies of Ascomycete yeast species have established these organisms have a range of carbon utilization repertoires, from specialists capable of consuming only a small range of carbohydrates to generalists that have a much broader carbon utilization repertoire [17]. *R. toruloides* is a generalist that can consume most soluble breakdown products

of plant biomass [18, 19]. *R. toruloides* can also accumulate up to 70% of its biomass as fatty

acids in conditions in which carbon is abundant but other nutrients are limiting and produces

useful secondary metabolites, such as carotenoids [20-26]. This makes *R. toruloides* an

excellent choice for metabolic engineering [27], where understanding the regulation of carbon

metabolism is critical [28, 29].

Cellulose is the most abundant plant biomass component and is composed of glucose

chains linked by β-1,4-glycosidic bonds [30]. *R. toruloides* can utilize the disaccharide cellulose

building block, cellobiose [31]. We identified a transcription factor, Cbr1, required for cellobiose

utilization. Cbr1 is homologous to the cellulose degradation regulator CLR-2/ClrB in

Ascomycetes [32], but the role of Cbr1 is significantly expanded in *R. toruloides*, elucidating

alternative mechanisms of building carbon utilization regulatory networks in fungi. Unlike CLR-

2/ClrB, Cbr1 is also required for tricarboxylic acid (TCA) cycle intermediate and fucose

utilization. Cellobiose, fucose, and TCA cycle intermediates are not catabolized through similar

pathways, suggesting *R. toruloides* may inhabit an ecological niche in which these carbon

sources are frequently found together, potentially through cooperation with or exploitation of

saprophytic filamentous fungi.

Further investigation of Cbr1 demonstrated this transcription factor inhibits carbon

catabolite repression specifically of glucose-glucose disaccharide utilization genes. This role for

Cbr1 contrasts with previously identified regulatory mechanisms of carbon catabolite repression,

which regulate utilization of all nonpreferred carbon sources. Taken together, the diverse roles of

Cbr1 in *R. toruloides* metabolic regulation present a previously uncharacterized mechanism of

carbon catabolite repression and nutrient sensing transcriptional network composition in fungi.

RESULTS

Cbr1 is required for wild-type utilization of cellobiose, TCA cycle intermediates, and fucose

R. toruloides can utilize cellobiose, the disaccharide breakdown product of the most

abundant plant cell wall polysaccharide, cellulose [31]. Several transcription factors regulate

cellulose and cellobiose utilization in Ascomycete filamentous fungi, including CLR-1/ClrA, CLR-

2/ClrB, and XlnR [1, 32-35]. To investigate genetic mechanisms regulating *R. toruloides*

cellobiose utilization, we searched for genes in the *R. toruloides* genome with homology to

these transcription factors. No clear homolog of CLR-1/ClrA or XlnR existed in *R. toruloides* (Fig

S1). However, we identified a single homolog of CLR-2/ClrB, RTO4_13588 (Fig 1A and Fig S1).

Fig 1. *CBR1* (RTO4_13588) is involved in utilization of cellobiose, TCA cycle

intermediates, and fucose. (**A**) Protein sequences of *N. crassa* CLR-2 and *Aspergillus*

nidulans ClrB homologs were used to build a phylogenetic tree using the maximum likelihood

method based on the Jones-Taylor-Thornton matrix-based model using FastTree [36]. Numbers

at nodes indicate the reliability of each split in the tree as computed using the Shimodaira-

Hasegawa test on three alternate topologies around that split with 1,000 resamples. *R.*

toruloides proteins are indicated with maroon diamonds. *N. crassa* proteins are indicated with

orange squares. *A. nidulans* proteins are indicated with green circles. *S. cerevisiae* proteins are

indicated with blue triangles. **(B)** Growth curves (OD600) of the indicated strains grown in yeast

nitrogen base (YNB) with the indicated carbon source at 1% except for citrate which was at 8.5

mM. Lines are the average and bars or colored bands are the standard deviation of three

biological replicates. $*p_{adj} < 10^{-4}$ and $**p_{adj} < 10^{-10}$ for $cbr1\Delta$ vs wild type (WT) and $cbr1\Delta$ vs

$cbr1\Delta$ + *CBR1* as determined by pairwise comparisons of estimated marginal means with a

holm multiple comparison correction.

We hypothesized RTO4_13588 may play a role in regulating cellobiose utilization in *R.*

toruloides. To test this hypothesis, we deleted RTO4_13588 and grew the resulting

RTO4_13588Δ strain on cellobiose. Growth of cells lacking RTO4_13588 on glucose was

indistinguishable from wild type cells (Fig 1B). However, deletion of RTO4_13588 resulted in a

complete lack of growth on cellobiose, which could be complemented by exogenous expression

of RTO4_13588 (Fig 1B). The *N. crassa* homolog of Cbr1, CLR-2, also plays a role in mannan

utilization leading us to hypothesize that *CBR1* may be required for utilizing mannose, a major

mannan building block [37], but $cbr1\Delta$ cells grew as well as wild type on mannose (Fig S2A).

The closest homolog to RTO4_13588 in the model yeast *Saccharomyces cerevisiae* is the

transcription factor Gal4, which regulates galactose utilization (Fig 1A) [38]. To test whether

RTO4_13588 also played a role in regulating galactose utilization in *R. toruloides*, we grew wild

type and RTO4_13588Δ cells with galactose as the sole carbon source. There was no

detectable difference in growth between wild type cells and cells lacking RTO4_13588 on

galactose (Fig S2A). Thus, we named RTO4_13588, *CBR1* for <u>c</u>ello<u>b</u>iose <u>r</u>egulator 1.

R. toruloides can grow on a wide variety of soluble carbon sources [18, 31]. We asked

whether *CBR1* was required for growth on any additional carbon sources. Cells lacking *CBR1*

were unable to grow as well as wild type cells on the deoxy hexose sugar fucose, which could

be complemented by exogenous expression of *CBR1* (Fig 1B). The growth of $cbr1\Delta$ cells was

indistinguishable from wild type cells on all other sugars tested (fructose, xylose, arabinose,

galacturonic acid, sucrose, maltose, trehalose, and raffinose) and on the lignin phenolic building

blocks, coumaric and ferulic acid (Fig S2A-S2B). *CBR1* was not required for utilization of the

lipid mimic Tween-80, the fermentation product acetate, or the amino acids L-alanine, proline,

glutamate, or glycine, suggesting *cbr1Δ* cells were capable of respiration (Fig S2C-S2D). Thus,

we hypothesized *cbr1Δ* cells would be able to grow on TCA cycle intermediates. To our surprise,

cbr1Δ cells had significant growth defects on citrate, α-ketoglutarate, succinate, fumarate, or

malate relative to wild type and *cbr1Δ + CBR1* cells (Fig 1B).

The role for Cbr1 in TCA cycle intermediate and fucose utilization could either be an

expanded role for Cbr1 in *R. toruloides* relative to CLR-2/ClrB or a previously uncharacterized

role for CLR-2/ClrB. To distinguish between these two possibilities, we assessed growth of the

model filamentous fungus *Neurospora crassa* wild type and *Δclr-2* cells in media containing

succinate, α-ketoglutarate, malate, or fucose as the sole carbon source. Wild type *N. crassa*

was only capable of limited growth on TCA cycle intermediates or fucose, and deletion of *clr-2*

caused no detectable growth defect (Fig S3). These data suggest the *CBR1* role in regulating

TCA cycle intermediate and fucose utilization is an expanded role for this transcription factor,

relative to Cbr1 orthologs in Ascomycete filamentous fungi.

Cbr1 activates a core regulon of five genes in response to cellobiose, TCA cycle intermediates,

and fucose

We had three possible hypotheses for genetic mechanisms through which Cbr1 could

regulate cellobiose, TCA cycle intermediate, and fucose utilization: (1) Cbr1 regulates genes

necessary to utilize all three classes of carbon sources in response to any of these carbon

sources, (2) Cbr1 regulates genes necessary to utilize each carbon source only in response to

that specific carbon source, and (3) *R. toruloides* utilization of cellobiose, fucose, and/or TCA

cycle intermediates proceeds through atypical metabolic pathways that require the same

transporters and/or catabolic enzymes. To distinguish between these three hypotheses, we

used transcriptional profiling. We exposed wild type and *cbr1Δ* cells to yeast nitrogen base

(YNB) lacking a carbon source (no carbon) or supplemented with 1% cellobiose, fucose, or

succinate for 8 h prior to harvesting the cells for RNA sequencing (RNAseq). The cellobiose,

succinate, and no carbon samples were submitted for whole transcript RNAseq, and the fucose

samples were submitted for 3' RNAseq.

Given the limited transcriptional characterization of *R. toruloides* on cellobiose or

succinate, we first examined the transcriptional profile of wild type cells in these conditions

relative to carbon starvation. The expression of 1168 genes was at least four-fold differentially

expressed in wild type cells exposed to cellobiose compared to carbon starvation, with 155

genes upregulated and 1013 genes downregulated. In wild type cells exposed to succinate, 525

genes were at least four-fold differentially expressed relative to carbon starved cells, with 142

genes upregulated and 383 genes downregulated. Intriguingly, over 75% of genes differentially

expressed on succinate (405/525) were also differentially expressed on cellobiose, supporting

the hypothesis that similar sets of genes are regulated in response to cellobiose and succinate

(Fig S4A-C and Dataset S1-S2).

Cells lacking *CBR1* also showed substantial transcriptional responses to cellobiose and

succinate compared to carbon starvation. Five hundred seventy-nine genes were at least four-

fold differentially expressed in cellobiose compared to carbon starvation. Eighty-one of the 179

genes (45%) upregulated on cellobiose in *cbr1Δ* cells were also upregulated in wild type cells

exposed to cellobiose, and 319 of the 400 genes (80%) downregulated on cellobiose in *cbr1Δ*

cells had the same expression pattern in wild type cells (Fig S4A, S4D, S4F, and Dataset S1-

S2). Similarly, 742 genes were differentially expressed on succinate compared to carbon

starvation in *cbr1Δ* cells. Eighty-six of the 296 genes (29%) upregulated on succinate in *cbr1Δ*

cells were also upregulated in wild type cells exposed to succinate, and 256 of 446 genes (57%)

downregulated on succinate in *cbr1Δ* cells were also downregulated in wild type cells exposed

to succinate (Fig S4B, S4E, S4G, and Dataset S1-S2). The significant transcriptional response

to cellobiose and succinate combined with the high degree of transcriptional similarity exhibited

by *cbr1Δ* and wild type cells suggests despite not being able to utilize cellobiose or succinate

like wild type cells, *cbr1Δ* cells still respond to these carbon sources to some extent.

Although many fungal transcription factor genes are upregulated in the condition in

which the transcription factor plays a role, *CBR1* expression was not induced in response to

cellobiose or succinate compared to carbon starvation (Fig S5A and Dataset S1-S2). To directly

regulate gene expression, Cbr1 must be present in the nucleus. We hypothesized Cbr1 might

be transported into the nucleus under conditions where it plays a role. To test this hypothesis,

we tagged Cbr1 with green fluorescent protein (mEGFP) and the histone H2A subunit *HTA1*

(RTO4_12090) with mRuby2 and exposed these cells to carbon starvation, glucose, cellobiose,

or succinate to identify the localization of Cbr1. Cbr1 colocalized with the histone H2A subunit

under all four conditions, indicating Cbr1 is expressed and in the nucleus during exposure to

glucose, cellobiose, succinate, and carbon starvation (Fig S5B).

To gain insight into the mechanism behind the growth defects of cells lacking *CBR1*, we

examined genes decreased in expression in *cbr1Δ* relative to wild type cells. Using a four-fold

differential expression cutoff, 284, 18, 25, and 25 genes were downregulated in *cbr1Δ* compared

to wild type cells on cellobiose, succinate, carbon starvation, and fucose, respectively (Fig 2A-D

and Dataset S1-S2). In concordance with our hypothesis that cellobiose, carboxylic acid, and

fucose utilization are transcriptionally linked via *CBR1*, genes annotated as part of the Kyoto

Encyclopedia of Genes and Genomes (KEGG) [39] 2-oxocarboxylic acid metabolism category

were enriched in the *CBR1* regulon on cellobiose (Fig 2E). Carbohydrate metabolism KEGG

[39] and gene ontology (GO) [40, 41] related categories were enriched on succinate, no carbon,

and fucose. Further analysis revealed the carbohydrate-related enrichments primarily consisted

of genes predicted to encode β-glucosidases, which cleave cellobiose into two glucose

molecules (Fig 2E). Only five genes were differentially expressed by at least four-fold in at least

three of the four conditions: two encoding putative β-glucosidases (RTO4_16716 and

RTO4_16717) that are adjacent in the *R. toruloides* genome, two encoding putative major

facilitator superfamily transporters (RTO4_10339 and RTO4_13825), and one encoding a

putative phosphoglycerate mutase (RTO4_11229) (Fig 2A-D and Dataset S1-S2).

Fig 2. Cbr1 regulates a core regulon of five genes. (**A-D**) Scatterplots of the $\log_2$(average

expression) of all genes in wild type (WT) cells versus the $\log_2$(fold change) in gene expression

in *cbr1Δ* relative to wild type cells when cells were exposed to (**A**) cellobiose, (**B**) succinate, (**C**)

media lacking a carbon source, or (**D**) fucose. Genes with $p_{adj} < 0.05$ are indicated in red or blue

if they had higher or lower expression in *cbr1Δ* cells than wild type cells, respectively. Genes

differentially expressed by a least four-fold in at least three of the four conditions (*CBR1* core

regulon) are circled in orange and labeled with their protein IDs. Genes in grey were not

significantly differentially expressed in *cbr1Δ* relative to wild type cells. Black dotted lines

indicate four-fold differential expression. Standard RNAseq (**A-C**) was used to determine gene

expression in transcripts per million (TPM) and fold change during exposure to cellobiose,

succinate, and carbon starvation. 3'RNAseq (**D**) was used to determine gene expression in

counts per million (CPM) and fold change during exposure to fucose. (**E**) Genes expressed at

least four-fold lower in *cbr1Δ* cells than wild type cells for each carbon source were tested for

Kyoto Encyclopedia of Genes and Genomes (KEGG) [39] (left) and gene ontology (GO) [40, 41]

(right) functional category enrichment. Comparisons were only displayed if they had at least one

statistically significantly enriched category for each analysis. Dot color represents the number of

genes differentially expressed belonging to a given category. Dot size represents the percent of

genes belonging to a given category that were differentially expressed relative to all genes in

that category. The text of GO biological process and molecular function categories are colored

black and red, respectively. The expression of *CBR1* was left out of the scatterplots, KEGG, and

GO enrichment analyses, since the change in expression of this gene was due to its deletion in

the *cbr1Δ* strain. All RNAseq experiments had three biological replicates.

Cbr1-regulated β-glucosidases are necessary for cellobiose utilization

The *CBR1*-dependent expression of two predicted β-glucosidase genes led us to

hypothesize that one or both β-glucosidase genes is essential for cellobiose utilization. An

amino acid alignment using LALIGN [42] revealed 72% identity and 90% similarity between the

two proteins, suggesting a potential for redundancy, so to test this hypothesis we deleted

RTO4_16716 and RTO4_16717, together and individually. RTO4_16716Δ RTO4_16717Δ cells

grew like wild type cells on glucose, citrate, succinate, and fucose, but were unable to grow on

cellobiose (Fig 3A and Fig S6A). Analysis of the individual deletion mutants revealed that

deletion of RTO4_16717 alone eliminated growth on cellobiose and complementation with

RTO4_16717 restored the ability to grow on cellobiose (Fig 3B). Cells lacking RTO4_16716

 alone grew as well as wild type cells on cellobiose (Fig 3B). These data indicate RTO4_16717

250 encodes the primary β-glucosidase for cellobiose utilization. Thus, we named RTO4_16717 and

251 RTO4_16716, *BGL1* and *BGL2*, respectively, for β-glucosidase 1 and 2.

Fig 3. Cbr1-dependent expression of secreted β-glucosidase genes *BGL1* (RTO4_16717)

and *BGL2* (RTO4_16716) is necessary for cellobiose utilization. (**A-B**) Growth (OD600) of

the indicated strains in the indicated carbon sources. (**C**) β-glucosidase activity secreted by the

indicated strains into culture supernatants after 20 h growth on the indicated carbon sources. β-

glucosidase activity is normalized to the β-glucosidase activity of wild type (WT) cells on

cellobiose. Growth (OD600) at time of supernatant harvest is in Fig S6B. (**D**) *BGL1* expression

of WT cells exposed to the indicated carbon source for 4 h relative to *BGL1* expression during

exposure to cellobiose for 4 h. (**E-F**) Normalized growth (OD600) of the indicated strains in 1%

cellobiose supplemented with the indicated percentage of spent supernatant from wild type cells

grown in (**E**) 1% cellobiose or (**F**) 1% succinate. To account for growth on nutrients remaining in

the spent supernatant, growth was normalized by subtracting the growth of each replicate on

media lacking a carbon source supplemented with the same concentration of spent supernatant.

Growth curves without normalization, growth curves in media without a carbon source

supplemented with spent supernatants, and β-glucosidase activity present in spent

supernatants are in Fig S7. (**A**, **B**, **E**, and **F**) Lines are the average and bars or colored bands

are the standard deviation of three biological replicates. (**C** and **D**) Bars are the average and

dots are the individual data points of (**C**) five (*cbr1Δ*) or six (all other strains) biological replicates

or (**D**) three biological replicates. (**A** and **B**) Colored stars indicate a significant difference in

growth of the respectively colored strain compared to wild type cells and, when applicable, the

respective complement strain. (**E** and **F**) Colored stars indicate a significant difference in growth

of the respectively colored strain compared to growth of that strain without supernatant

supplementation. *$p_{adj} < 0.05$ and **$p_{adj} < 10^{-7}$ as determined by either (**A**, **B**, **E**, and **F**) pairwise

comparisons of estimated marginal means with a holm multiple comparison correction, (**C**) a

two-way ANOVA with a TukeyHSD post-hoc test, or (**D**) a one-way ANOVA with a TukeyHSD

post-hoc test.

Both *BGL1* and *BGL2* have predicted secretion signals, leading us to hypothesize

cellobiose cleavage by β-glucosidases occurs extracellularly. To address this hypothesis, we

harvested supernatants of wild type, *cbr1Δ*, and *cbr1Δ* + *CBR1* cell cultures after 20 h of growth

in glucose, cellobiose, or succinate to measure secreted β-glucosidase activity. β-glucosidase

activity was detected in the culture supernatant of wild type cells exposed to cellobiose

indicating that β-glucosidases are indeed secreted by *R. toruloides* (Fig 3C). Consistent with our

transcriptomics data, and despite the lack of a role for *BGL1* and *BGL2* in carboxylic acid

utilization (Fig S6A), β-glucosidase activity was also detectable in the culture supernatant of wild

type cells on succinate. Deletion of *CBR1* eliminated β-glucosidase activity in the supernatant of

cellobiose and succinate cultures. This activity was restored in the supernatant of cultures of

cbr1Δ + CBR1 cells, indicating the secreted β-glucosidase activity was dependent on *CBR1* (Fig

3C).

Glucose cultures had no detectable β-glucosidase activity, which could either be due to

reduced expression of β-glucosidase genes or reduced secretion of β-glucosidase proteins in

glucose relative to cellobiose and succinate media (Fig 3C). To distinguish between these

possibilities, we performed reverse transcription quantitative PCR (RT-qPCR) on *BGL1* in wild

type cells. Corroborating our RNAseq data, *BGL1* expression was highest on cellobiose,

followed by succinate, and then carbon starvation (Fig 3D). During exposure to glucose, *BGL1*

expression was nearly undetectable, indicating the lack of β-glucosidase activity in the glucose

culture supernatant was due to reduced expression of β-glucosidase genes (Fig 3C and 3D).

These data may indicate β-glucosidase expression is regulated by carbon catabolite repression

and that β-glucosidases potentially act as "scout" enzymes during carbon starvation similar to

some polysaccharide degrading enzymes in some filamentous fungi [1, 3].

Given that *bgl1Δ* cells cannot grow when cellobiose is the sole carbon source (Fig 3B),

we hypothesized Bgl1 was the source of the secreted β-glucosidase activity. To test this

hypothesis, we incubated *bgl1Δ* cells in cellobiose, succinate, or glucose for 20 h. While β-

glucosidase activity in the cellobiose *bgl1Δ* culture supernatant was 22-fold lower than that of

wild type culture supernatants, it was detectable, and β-glucosidase activity of succinate *bgl1Δ*

cultures was even more pronounced (Fig 3C and Fig S6C). This reduction in β-glucosidase

secretion was mitigated in *bgl1Δ + BGL1* cells (Fig 3C). The low level of β-glucosidase activity

remaining in *bgl1Δ* cells could either result from secretion of Bgl2 or an unidentified protein. To

distinguish between these possibilities, we exposed *bgl1Δ bgl2Δ* and *bgl1Δ bgl2Δ + BGL1 BGL2*

cells to cellobiose, succinate, or glucose for 20 h. Cells lacking both *BGL1* and *BGL2* had no

detectable β-glucosidase activity, which was rescued by complementation with *BGL1* and *BGL2*

(Fig 3C and Fig S6C). These data suggested Bgl2 contributes to β-glucosidase activity in

cellobiose and succinate cultures even though *BGL2* was dispensable for growth on cellobiose.

Indeed, cells lacking *BGL2* alone had a 2.8-fold reduction of β-glucosidase activity on both

cellobiose and succinate relative to wild type cells, demonstrating that both Bgl1 and Bgl2 are

responsible for the secreted β-glucosidase activity of *R. toruloides* (Fig 3C).

It was possible either that β-glucosidase genes were the only Cbr1-regulated genes

necessary for cellobiose utilization or that other genes were also required. To distinguish

between these two possibilities, we expressed *BGL1* under the constitutive *AAC1*

(RTO4_12704) promoter [43] in a *cbr1Δ* background (*cbr1Δ* + P_{AAC1}-*BGL1*). Constitutive

expression of *BGL1* in *cbr1Δ* cells resulted in wild type levels of growth on cellobiose when cells

were inoculated at very low densities and a growth advantage on cellobiose relative to wild type

cells at higher starting densities (Fig 3B and Fig S6D and S6E). Correspondingly, *cbr1Δ* + P_{AAC1}-

BGL1 cellobiose culture supernatants had 3.2-fold higher β-glucosidase activity than wild type

culture supernatants (Fig S6F).

If β-glucosidase activity is the only Cbr1-regulated requirement for cellobiose utilization,

we speculated *cbr1Δ* and *bgl1Δ bgl2Δ* cell growth on cellobiose could be rescued by

exogenously supplied β-glucosidases. To test this hypothesis, we inoculated wild type, *cbr1Δ*,

and *bgl1Δ bgl2Δ* cells into cellobiose supplemented with filtered, spent supernatant from wild

type cells grown on cellobiose, succinate, or glucose. This spent supernatant supplementation

enabled significantly improved growth of *cbr1Δ* and *bgl1Δ bgl2Δ* cells, which was not due to

growth on residual carbon in the spent supernatant (Fig 3E, 3F, and Fig S7). Our data

demonstrate genes encoding secreted β-glucosidases, activated in response to cellobiose and

TCA cycle intermediates, are the only Cbr1-regulated requirement for cellobiose utilization.

CBR1-mediated transcriptional response to cellobiose is dependent on extracellular β-

glucosidases

Extracellular degradation of cellobiose by *R. toruloides* provoked the question of whether

cellobiose activates the *CBR1*-dependent transcriptional response to cellobiose or if the

activating signal is a cellobiose catabolic derivative. We differentiated between these

possibilities in two ways.

We first compared the transcriptional profile of *bgl1Δ bgl2Δ* cells exposed to cellobiose,

which have no extracellular β-glucosidase activity but contain wild type copies of *CBR1*, to that

of wild type and *cbr1Δ* cells using 3' RNAseq. The transcriptional profiles of *bgl1Δ bgl2Δ* cells

were more similar to *cbr1Δ* than wild type cells (Fig 4A-C and Fig S8 and Dataset S3-S4).

Thirty-five genes were at least four-fold differentially expressed in *bgl1Δ bgl2Δ* relative to wild

type cells (12 upregulated and 23 downregulated) (Fig 4A and Dataset S3-S4). Twenty-three of

these genes were also at least four-fold differentially expressed in *cbr1Δ* compared to wild type

cells (Fig 4B and Dataset S3-S4). In contrast, only five genes were at least four-fold differentially

expressed in *bgl1Δ bgl2Δ* relative to *cbr1Δ* cells, which included a 38-fold upregulation of the

CBR1-regulated predicted transporter gene RTO4_13825 (Fig 4C and Dataset S3-S4). The

expression pattern of RTO4_13825, raised the question of whether any other genes in the core

Cbr1 regulon were involved in cellobiose sensing or utilization. Cells lacking RTO4_13825,

RTO4_10339, or RTO4_11229 all grew as well as wild type on cellobiose (Fig S9). These data

indicate the signal for the *CBR1*-dependent cellobiose response requires secreted β-

glucosidases, suggesting cellobiose is likely not the signaling molecule for Cbr1 activation.

Fig 4. *CBR1*-mediated transcriptional response to cellobiose is dependent on secreted β-

glucosidases. (**A-C**) Scatterplots of the $\log_2$(average expression) of all genes in (**A-B**) wild type

(WT) or (**C**) *cbr1*Δ cells versus the $\log_2$(fold change) in gene expression for the indicated

comparison when cells were exposed to cellobiose as measured by 3'RNAseq. Genes with p_{adj}

< 0.05 are indicated in red or blue if they had higher or lower expression, respectively, in (**A** and

C) *bgl1*Δ *bgl2*Δ cells or (**B**) *cbr1*Δ cells. The genes in the *CBR1* core regulon are circled in

orange and labeled with their respective protein IDs. The expression of *CBR1*, *BGL1,* and *BGL2*

were left out of the scatterplots involving their respective deletion strains. RNAseq experiments

had three biological replicates. (**D**) *BGL1* expression in wild type cells exposed to carbon

starvation for 1 h followed by exposure to continued carbon starvation or the indicated

concentrations of cellobiose or 2-deoxyglucose (2-DG) as the sole carbon source for 1 h or 4 h.

Expression values were normalized to the average expression of *BGL1* on no carbon for each

timepoint and $\log_2$ transformed. Bars are the average and dots are the values of three biological

replicates. *p_{adj} < 0.05 relative to the no carbon condition for the respective time point, as

determined by two-way ANOVA with a TukeyHSD post-hoc test.

β-glucosidases cleave cellobiose into two glucose molecules. Thus, a second way we

answered the question of whether cellobiose or a cellobiose catabolic derivative is the activation

signal was to determine whether low glucose concentrations activated Cbr1-regulated gene

expression. To mitigate changes in glucose concentration due to metabolism, we used the non-

metabolizable glucose analog 2-deoxyglucose. We exposed wild type cells to 10-fold dilutions of

2-deoxyglucose (100 μM to 0.1 μM), 1% 2-deoxyglucose, 1% cellobiose, or carbon starvation

for 1 h and 4 h and measured *BGL1* transcription via RT-qPCR. As expected, cells exposed to

cellobiose expressed *BGL1* at high levels, and *BGL1* expression was repressed during

exposure to 1% 2-deoxyglucose (Fig 4D). Exposure to 1 μM 2-deoxyglucose resulted in a two to

three-fold increase in *BGL1* expression relative to carbon starvation at both timepoints (Fig 4D).

This *BGL1* activation did not rise to the same magnitude as exposure to cellobiose itself,

perhaps due to differences in glucose concentration, the rate glucose enters the cell, or a lack of

resources due to starvation on 2-deoxyglucose. However, increased *BGL1* expression on 1 μM

2-deoxyglucose suggests a low concentration of glucose, a catabolic cellobiose derivative,

activates β-glucosidase expression.

A Cbr1-regulated transporter gene is necessary for citrate utilization

Two genes encoding predicted major facilitator superfamily transporters (RTO4_13825

and RTO4_10339) and a gene encoding a predicted phosphoglycerate mutase (RTO4_11229)

were also members of the *CBR1* core regulon. We hypothesized these genes may be

necessary for growth on TCA cycle intermediates and/or fucose. To test this hypothesis, we

grew RTO4_13825Δ, RTO4_10339Δ, and RTO4_11229Δ cells on media containing glucose,

fucose, citrate, succinate, α-ketoglutarate, malate, or fumarate. The growth of RTO4_10339Δ

and RTO4_11229Δ cells was indistinguishable from wild type cells under all conditions tested

(Fig S9). Cells lacking RTO4_13825 were unable to grow on citrate but grew as well as wild

type cells on all other carbon sources tested (Fig 5A and Fig S9A). Thus, we named

RTO4_13825 *TCT1* for tricarboxylic acid transporter.

Fig 5. *TCT1* (RTO4_13825) is required for citrate utilization. (**A**) Growth (OD600) of the

indicated strains in YNB 1% glucose or 8.5 mM citrate. Lines are the average and bars are the

standard deviation of three biological replicates. *$p_{adj} < 10^{-7}$ of *cbr1Δ* vs wild type (WT), *tct1Δ* vs

WT, and *tct1Δ* vs *tct1Δ + TCT1* as determined by pairwise comparisons of estimated marginal

means with a holm multiple comparison correction. (**B**) Heatmap of the $\log_2$(fold change) in the

expression of genes at least four-fold differentially expressed in one or more of the indicated

comparisons when cells were exposed to citrate as the sole carbon source, as measured by

3'RNAseq. Orange bars to the right of the heatmap indicate significance using the criteria of p_{adj}

< 0.05 and at least four-fold change in expression, while grey bars indicate genes that do not

meet those criteria in the indicated comparison (N.S., not significant). The expression of *CBR1*

and *TCT1* were left out of the heatmaps since changes in expression were due to their

respective deletions in the *cbr1Δ* and *tct1Δ* strains. RNAseq experiments had three biological

replicates.

To gain further insight into the role of *TCT1*, we conducted RNAseq on wild type, *cbr1*Δ,

and *tct1*Δ cells exposed to citrate for 24 h. Only 20 genes were at least four-fold differentially

expressed in *cbr1*Δ relative to wild type cells (Fig 5B and Dataset S5-S6). Surprisingly, these 20

genes did not include *TCT1* or genes predicted to play a role in citrate metabolism. Of the other

genes in the *CBR1* core regulon, *BGL1* and *BGL2* were downregulated by 6.7 and 3.2-fold,

respectively, in *cbr1*Δ relative to wild type cells, but RTO4_10339 and RTO4_11229 were not

differentially expressed (Fig 5B and Dataset S5-S6). Only one gene, RTO4_15140, had

decreased expression in *tct1*Δ cells relative to wild type cells. RTO4_15140 was also

downregulated in *cbr1*Δ relative to wild type cells (Fig 5B and Dataset S5-S6). RTO4_15140 is

predicted to encode a P-type ATPase transduction domain A containing protein with a single

transmembrane domain, and any potential role in citrate utilization is unclear. The similarity of

*tct1*Δ and wild type cell transcriptional profiles may suggest *tct1*Δ cells can sense citrate, despite

not being able to grow.

Cbr1 inhibits carbon catabolite repression during glucose-glucose disaccharide utilization

Glucose is a highly preferred carbon source in most fungi. Extracellular β-glucosidases

cleave cellobiose into two glucose molecules, which we hypothesized would repress cellobiose

utilization genes via carbon catabolite repression [1, 3, 4]. *BGL1* expression is almost

undetectable when wild type cells are exposed to 1% glucose or at least 100 µM 2-

deoxyglucose (Fig 3D and 4D). Thus, we hypothesized wild type cell growth on cellobiose would

be inhibited by the non-hydrolyzable glucose analog 2-deoxyglucose. Indeed, 61 µM 2-

deoxyglucose fully inhibited *R. toruloides* growth on 1% cellobiose, indicating glucose is

preferred over cellobiose by *R. toruloides* (Fig 6A).

Fig 6. Cbr1 inhibits carbon catabolite repression during glucose-glucose disaccharide

utilization. (**A**) Growth (OD600) of wild type (WT) cells in 1% cellobiose supplemented with the

indicated concentration of 2-deoxyglucose (2-DG). (**B-F**) Growth of the indicated strains on the

indicated carbon sources with the indicated concentration of 2-DG or allyl alcohol (AA). Lines

are the average and bars are the standard deviation of (**A-E**) three or (**F**) four biological

replicates. $*p_{adj} < 0.01$ and $**p_{adj} < 10^{-4}$ for (**A**) 1% cellobiose supplemented with the indicated 2-

DG concentration vs 1% cellobiose or (**B-F**) $cbr1\Delta$ vs WT and $cbr1\Delta$ vs $cbr1\Delta + CBR1$ as

determined by pairwise comparisons of estimated marginal means with a holm multiple

comparison correction.

This carbon preference represents a biological challenge during utilization of substrates

in which glucose is released during catabolism; a negative feedback loop could repress

substrate utilization genes in response to the released glucose, slowing growth on the carbon

source. We hypothesized *CBR1* may inhibit carbon catabolite repression to bypass this negative

feedback loop. To test this hypothesis, we grew wild type, $cbr1\Delta$, and $cbr1\Delta + CBR1$ cells on a

nonpreferred carbon source and 2-deoxyglucose. Cells lacking *CBR1* were significantly more

sensitive to 2-deoxyglucose than wild type or *cbr1Δ* + *CBR1* cells when the glucose-glucose

disaccharides maltose or trehalose were the carbon source, suggesting Cbr1 is a negative

regulator of carbon catabolite repression (Fig 6B).

Canonically, regulators of carbon catabolite repression regulate a broad spectrum of

nonpreferred carbon source utilization genes [1, 3-10]. We hypothesized *CBR1* would play a

similar role in *R. toruloides*. However, the growth of *cbr1Δ* cells exposed to 2-deoxyglucose in

combination with the glucose-fructose disaccharide sucrose, the hexose monosaccharide

fructose, the pentose monosaccharide xylose, the amino acid proline, or the lipid mimic Tween-

80 was indistinguishable from wild type cells (Fig 6C-6E). To measure the role of *CBR1* in

regulating genes involved in alcohol utilization, we used allyl alcohol, which is converted to the

toxic molecule acrolein by alcohol dehydrogenases. Alcohol dehydrogenase expression is

repressed by glucose [44]. Wild type and *cbr1Δ* cells were equally sensitive to allyl alcohol,

indicating glucose-mediated repression of alcohol dehydrogenases is not dependent on Cbr1

(Fig 6F). To determine whether any *CBR1* core regulon genes play a role in carbon catabolite

repression, we tested 2-deoxyglucose sensitivity of *bgl1Δ bgl2Δ*, *tct1Δ*, RTO4_10339Δ, and

RTO4_11229Δ cells. The sensitivity of these strains to 2-deoxyglucose during exposure to

trehalose, maltose, sucrose, proline, xylose, fructose, and tween-80 was indistinguishable from

wild type cells (Fig S10). These data indicate Cbr1 inhibits carbon catabolite repression of

glucose-glucose disaccharides either by directly regulating gene expression or through genes

outside of the *CBR1* core regulon.

A role for Cbr1 Ascomycete homologs in carbon catabolite repression regulation has not

been identified, so we asked whether the *N. crassa* Cbr1 homolog CLR-2 plays a similar role.

The sensitivity of wild type and Δ*clr-2* cells to 2-deoxyglucose was indistinguishable during

growth on the glucose-glucose disaccharide trehalose or the monosaccharide xylose (Fig S11).

These data indicate the role for Cbr1 in *R. toruloides* in inhibiting carbon catabolite repression of

glucose-glucose disaccharides is an expanded role for Cbr1 relative to the model Ascomycete

ortholog CLR-2.

DISCUSSION

Gene regulation in response to nutrient availability requires cells to integrate information

from activating and repressing nutrient signals. Previously described fungal transcriptional

networks regulating carbon utilization are made up of carbon source-specific transcription

factors that activate genes necessary to utilize a particular carbon source when it is present and

broad-acting transcription factors that regulate carbon catabolite repression [1, 2]. However,

studies of these carbon source utilization transcriptional networks have mainly focused on

Ascomycete fungi. We investigated nutrient sensing regulatory mechanisms in a Basidiomycete

yeast and identified a transcription factor that coregulates carbon source-specific gene

activation and carbon catabolite repression specifically of glucose-glucose disaccharide

utilization genes (Fig 7).

Fig 7. Model for Cbr1-mediated regulation of cellobiose, TCA cycle intermediate, and fucose utilization and carbon catabolite repression. Cbr1 is activated by the presence of a low glucose concentration, fucose, and TCA cycle intermediates or their metabolic derivatives. Activated Cbr1 induces the expression of genes encoding β-glucosidases, predicted transporters, and a predicted phosphoglycerate mutase. The Cbr1-regulated predicted transporter gene *TCT1* is required for citrate utilization. Tct1 is placed at the plasma membrane in this model, but the localization of Tct1 is currently unknown and should be the subject of future studies. Transporters for other TCA cycle intermediates and fucose are unknown. The β-glucosidase genes *BGL1* and *BGL2* encode secreted β-glucosidases that cleave cellobiose into two glucose molecules. *BGL1* is required for growth on cellobiose. A high glucose concentration activates carbon catabolite repression, which represses β-glucosidase gene expression and likely represses the expression of genes necessary to utilize fucose and TCA cycle intermediates. Cbr1 plays a role in inhibiting carbon catabolite repression, specifically when a glucose-glucose disaccharide is present as the nonpreferred carbon source.

Cbr1 regulation of cellobiose, TCA cycle intermediate, and fucose utilization may shed light on complex microbial interactions

There have been limited investigations of genetic regulators of cellulose and cellobiose utilization in Basidiomycetes. A recent study identified Roc1 as a regulator of cellulase-encoding genes in the basidiomycete wood-decaying fungus *Schizophyllum commune* [45]. However, homologs of Roc1 do not exist outside of the Agaricomycetes [45]. In *Ganoderma lucidum*, the R2R3-type MYB transcription factor *Gl*Myb activates cellulase gene expression [46]. In Ascomycete filamentous fungi, CLR-2/ClrB regulates cellulose degradation and utilization through direct activation of cellulase genes [32, 47]. We identified the CLR-2/ClrB ortholog Cbr1 in the oleaginous, basidiomycete yeast *R. toruloides* (Fig 1). Although Ascomycete and Basidiomycete fungi are separated by approximately 400 million years of evolution [12], Cbr1 is

required to utilize the cellulose breakdown product cellobiose (Fig 1). *N. crassa* CLR-2 does not

directly regulate the expression of genes encoding β-glucosidases, which are regulated by CLR-

1 [32, 47]. However, although the role of Cbr1 is significantly expanded beyond that of CLR-

2/ClrB to include inhibition of carbon catabolite repression and utilization of TCA cycle

intermediates and fucose, Cbr1 regulation of cellobiose utilization indicates some functional

conservation of CLR-2/ClrB/Cbr1 exists between Ascomycete filamentous fungi and

Basidiomycete yeast.

Cbr1 regulates citrate utilization through the predicted transporter gene *TCT1* and

cellobiose utilization through the β-glucosidase genes *BGL1* and *BGL2* (Fig 7). Genetic

mechanisms through which Cbr1 regulates utilization of fucose and dicarboxylic acid TCA cycle

intermediates remains unclear. It is possible the two uncharacterized genes in the *CBR1* core

regulon, the predicted transporter RTO4_10339 and the predicted phosphoglycerate mutase

RTO4_11229, play roles in fucose and/or dicarboxylic acid utilization that are redundant with

other *CBR1*-regulated genes that did not meet our strict *CBR1* core regulon criteria (Fig 2 and

Fig S4). It will be the role of future studies to characterize roles for RTO4_10339, RTO4_11229,

and other genes that may be responsible for fucose and dicarboxylic acid catabolism in the

CBR1 pathway.

In other fungal species, utilization of multiple carbon sources is occasionally regulated by

a single transcription factor. CLR-2 in *N. crassa* regulates cellulose utilization, mannan

utilization, and a small number of hemicellulose utilization genes [3, 32, 37, 47]. In *Trichoderma*

reesei, *Penicillium oxalicum*, and some Aspergilli, the transcription factor Xyr1/XlnR regulates

cellulose and hemicellulose utilization [33-35]. ARA-1/Ara1 regulates arabinan and galactose

utilization in *N. crassa* and *T. reesei* [3, 48]. FarA regulates utilization of fatty acids and the lignin

component ferulic acid in Aspergilli [49, 50]. In *Candida albicans*, Adr1 is required for growth on

citrate and substrates that feed into the TCA cycle, like glutamate and malate [51]. However, in

these examples linkage of these carbon sources can be explained either by the presence of

both carbon sources in the plant cell wall (i.e., cellulose and hemicellulose or arabinan and

galactose) or shared catabolic pathways (i.e., fatty acids and ferulic acid or citrate, glutamate,

and malate).

Cellobiose, fucose, and TCA cycle intermediates are not linked by known catabolic

pathways, and our data do not support catabolism through the same noncanonical metabolic

pathway in *R. toruloides*. Cbr1-regulated β-glucosidase genes are critical for cellobiose but not

fucose or TCA cycle intermediate utilization (Fig 3 and Fig S6A), while Tct1 is necessary for

citrate but not cellobiose or fucose utilization (Fig 5A and Fig S9A). Aside from *CBR1*, we did not

identify genes necessary for utilization of all three of cellobiose, fucose, and TCA cycle

intermediates.

We speculate coregulation of cellobiose, fucose, and TCA cycle intermediates may

suggest *R. toruloides* encounters cellobiose, fucose, and TCA cycle intermediates together in

nature. One possible explanation is *R. toruloides* scavenges cellobiose and fucose during plant

biomass degradation by filamentous fungi, which secrete organic acids like citrate or succinate

to acidify their environment and can contain fucose in their cell walls [52-54]. β-glucosidase

gene activation that is primed by organic acids secreted by fungi or fucose in fungal cell walls

before encountering cellobiose released from cellulose could confer a growth advantage to *R.*

toruloides over microbes that only activate β-glucosidase genes in response to cellobiose itself.

Indeed, high densities of cells constitutively expressing *BGL1* that do not require time to activate

β-glucosidase genes had a growth advantage over wild type cells that activated β-glucosidase

genes in response to cellobiose (Fig S6D).

Cbr1 regulation of carbon catabolite repression specifically of glucose-glucose disaccharides

disrupts a negative feedback loop for disaccharide utilization

Carbon catabolite repression enables prioritization of carbon sources that require less

energy to metabolize over carbon sources that require more energy to metabolize [1]. Known

carbon catabolite repression genetic mechanisms regulate genes required for utilization of all

less preferred carbon sources when a more preferred carbon source is present [3-10]. In

contrast, Cbr1 inhibits carbon catabolite repression specifically of glucose-glucose disaccharide

utilization (Fig 6). We speculate this mechanism allows continued expression of genes that

hydrolyze disaccharide bonds even when this hydrolysis produces preferred monosaccharides,

combating a negative feedback loop (Fig 7). To our knowledge, genetic linkage via a single

transcription factor of carbon catabolite repression inhibition with activation of genes necessary

for nonpreferred carbon source utilization has not been described in fungi. Additionally,

regulation of genes involved in utilization of a specific class of nonpreferred carbon sources in

response to a preferred carbon source represents a previously uncharacterized form of carbon

catabolite repression regulation.

Fungal pathogens of plants and animals that do not regulate carbon catabolite

repression like wild type cells frequently have reduced virulence [8, 55-57]. Carbon utilization

and carbon catabolite repression regulation by fungi also affects how they respond to antifungal

drugs [14, 15]. Thus, understanding diverse carbon catabolite repression mechanisms present

throughout the fungal kingdom is critical in controlling fungal disease. Additionally, a more

complete understanding of mechanisms of carbon catabolite repression and the diversity of

nutrient sensing transcriptional networks is necessary for harnessing the primary and secondary

metabolic capabilities of the fungal kingdom for biotechnology. Mapping these networks will be

critical in metabolically engineering fungi for green industrial fermentation.

MATERIALS and METHODS

Strains and culturing

R. toruloides strains used in this study are listed in Table S1. All strains were derived

from the wild type reference strain IFO 0880 (also called NBRC 0880, obtained from Biological

Resource Center, NITE (NBRC), Japan). The starting strain for genetic manipulations was the

non-homologous end-joining deficient *yku70Δ* (RTO4_11920Δ) strain [58]. *R. toruloides* was

transformed using *Agrobacterium tumefaciens*-mediated transformation as previously described

[26], using *A. tumefaciens* EHA 105 and plasmids derived from pGI2 [59]. All strains were

confirmed by PCR and/or DNA sequencing of transformed loci. For details see Text S1. *R.*

toruloides strains were grown from freezer stocks on yeast extract peptone dextrose (YPD)

(VWR) + 2% agar (BD Difco™ Bacto™) plates at 28°C for 2-3 d prior to inoculation into liquid

YPD to grow strains at 28°C with constant shaking at 250 rpm to exponential phase growth prior

to starting experiments.

N. crassa strains used in this study are listed in Table S2. These strains were derived

from the wild type reference strain FGSC 2489 and obtained from the Fungal Genetics Stock

Center [60, 61]. *N. crassa* cells were grown from freezer stocks on Vogel's minimal medium [62]

+ 2% sucrose + 1.5% agar (BD Difco™ Bacto™) slants for 2 d at 28°C in the dark and 4 d at

28°C in constant light. *N. crassa* conidia were harvested from slants with sterile double distilled

H_2O for inoculation.

All *R. toruloides* experiments were performed with yeast nitrogen base (YNB) with

NH_4SO_4 without amino acids (MP Biomedicals) + 12 mM K_2HPO_4 (Fisher) + 38 mM KH_2PO_4

(Fisher) + 100 nM Fe(II)SO_4 pH 6.2 with the indicated carbon source at 1% w/v with the

exception of coumaric acid and ferulic acid, which were added at a concentration of 0.2% w/v,

and citrate which was added at a concentration of 8.5 mM. All *N. crassa* experiments were

performed using Vogel's minimal medium [62] with 50 mM NH_4Cl instead of 25 mM NH_4NO_3

(VMM NH_4Cl) and the indicated carbon source at 1% w/v. Carbon sources used are listed in

Table S3.

R. toruloides and N. crassa growth experiments

R. toruloides colonies were inoculated into liquid YPD and grown at 28^{o}C with constant

shaking at 250 rpm to exponential phase. Cells were then washed and inoculated into 750 μL of

the indicated media in 48-well plates at 0.01 OD600/mL or 0.1 OD600/mL, except for the growth

curve in Fig S6D, which was inoculated at 0.5 OD600/mL as measured by an Eppendorf

BioPhotometer with a 10 mm path length. Plates were incubated at 28^{o}C with constant shaking

(3 mm orbit, 600 rpm for individual time points and 3 mm orbit, 425 rpm for kinetic growth

experiments). Growth was measured as OD600 on an Agilent BioTek Synergy HTX for

individual time points or an Agilent BioTek Epoch2 for kinetic growth experiments. For details

see Text S1.

N. crassa conidia were inoculated into either 100 mL of liquid VMM NH$_4$Cl with the

indicated carbon source in 250 mL flasks or 3 mL of liquid VMM NH$_4$Cl with the indicated carbon

source in round-bottomed, deep-well 24-well plates. After 24 h (flasks) or 7-10 d (24-well plates)

biomass was harvested, dried for 24 h in a 60^{o}C drying oven, and weighed. For details see Text

S1.

Phylogenetic tree generation

The Basic Local Alignment Search Tool for proteins (BLASTP) was used to identify

homologs of CLR-1, CLR-2, and XLR-1 from *N. crassa* (OR74A) and ClrA, ClrB, and XlnR from

Aspergillus nidulans (A4) in the proteomes of *N. crassa*, *A. nidulans*, *S. cerevisiae* (S288C)*,* and

R. toruloides (IFO 0880). CLR-1, CLR-2, and XLR-1 from *N. crassa* (OR74A) and ClrA, ClrB,

and XlnR from *A. nidulans* (A4) protein sequences were obtained from Mycocosm [63].

Homologs with E < 10^{-10} were aligned via MUSCLE [64] in MEGA11 [65] and the phylogenetic

tree was generated using FastTree [36] using the maximum likelihood method based on the

Jones-Taylor-Thornton matrix-based model and visualized in MEGA11. The reliability of each

split in the tree was computed using the Shimodaira-Hasegawa test on three alternate

topologies around that split. Trees were visualized in MEGA11 [65] and rooted on the midpoint.

Microscopy

Fluorescent microscopy was performed using a Nikon Eclipse Ti2-E inverted microscope

equipped with a prime BSI express CMOS camera using a CFI Plan Apochromat 100X oil

objective. For details see Text S1.

RNA sequencing and transcript abundance

Flasks containing 100 mL of YNB without a carbon source or YNB with the indicated

carbon source were inoculated with the indicated exponentially growing cells at 0.1 OD600/mL

(measured by an Eppendorf BioPhotometer). Culture conditions and sequencing methods for

each dataset are summarized in Table S4. Cells were harvested and flash frozen in liquid N_2.

RNA was extracted using the Quick-RNA Fungal/Bacterial Miniprep Kit (Zymo), followed by

either treatment with the TURBO DNA-free kit (Invitrogen) for standard RNAseq or DNase I

treatment (NEB) and purification with the Monarch® Spin RNA Cleanup Kit (NEB) for 3'RNAseq.

Standard RNAseq was performed at the California Institute for Quantitative Biosciences at the

University of California Berkeley (QB3-Berkeley) on an Illumina NextSeq 2000 with 150 bp

paired end reads. 3'RNAseq was performed at the Biotechnology Resource Center at Cornell

University on an Illumina NovaSeqX with 150 bp paired end reads.

Reads were aligned to the *R. toruloides* IFO 0880 genome (v4) [26]. For standard

RNAseq, transcript abundance (transcripts per million, TPM) was quantified using Salmon v.

1.10.0 [66]. For 3'RNAseq, reads were aligned using STAR v. 2.7.11b [67], counted with HTSeq

v. 2.0.9 [68], and counts per million (CPM) values were calculated by dividing counts for a gene

by total counts for the sample and multiplying by 10^6. The GFF file used to analyze 3'RNAseq

data is available in Dataset S7. Differential expression was determined using DESeq2 v. 1.50.1

[69]. RNAseq data were deposited in the Gene Expression Omnibus (GEO) at the National

Center for Biotechnology Information (NCBI) and are accessible through GEO series accession

numbers GSE293943 (standard RNAseq) and GSE313844 (3'RNAseq). For details see Text

S1.

Functional enrichment analysis

KEGG and GO enrichment analyses were performed using the enricher function of the

clusterProfiler R package v. 4.18.1 with Benjamini-Hochberg multiple hypothesis testing

correction [70]. For details see Text S1.

Reverse transcription quantitative PCR

To measure *BGL1* expression, wild type cells were exposed to the indicated carbon

source in 100 mL of the indicated media in 250 mL flasks and incubated at 28°C with shaking at

200 rpm for the indicated length of time. Cells were harvested, flash frozen in liquid N_2, and RNA

was purified as described above. RT-qPCR was performed using the Luna$^{®}$ Universal One-Step

RT-qPCR Kit (NEB) on a CFX96^{TM} Real-Time System (Bio-Rad). For details see Text S1.

β-glucosidase enzyme activity assays

β-glucosidase activity was measured using p-nitrophenyl-β-D-glucopyranoside (EMD

Millipore Corp.). For details see Text S1.

Spent media growth assays

Filtered supernatants of wild type cells growing in YNB 1% glucose, YNB 1% cellobiose,

or YNB 1% succinate were added to YNB lacking a carbon source or YNB 1% cellobiose at the

indicated concentration. The indicated strains were inoculated into this media at 0.01 OD600/mL

(measured by an Eppendorf BioPhotometer) in 750 μL in 48-well plates and incubated at 28ºC

with shaking at 600 rpm (3 mm orbit). For details see Text S1.

Statistical significance tests

All statistical analyses were performed in Rstudio (v2025.09.2+418; R v4.5.2) using

either a one-way or two-way ANOVA with a Tukey post-hoc test (bar graphs), a two-tailed

Welch's two-sample *t*-test (bar graphs), or a general linear model with two-tailed post-hoc

analysis using the emmeans function of the emmeans R package [71] with a holm multiple

comparison correction (line graphs). Statistical significance for differential expression of

RNAseq data was determined using DESeq2 [69]. All experiments were performed with a

minimum of three biological replicates. The exact number of replicates for each experiment is

noted in the relevant figure legend.

Data availability

Raw RNAseq data is available in GEO at the NCBI through accession numbers

GSE293943 (standard RNAseq) and GSE313844 (3'RNAseq). Processed RNAseq data is

available in Dataset S1-S6. The numerical values used to generate all other graphs are in

Dataset S9. Strains constructed in this study are available upon request.

ACKNOWLEDGEMENTS

We thank members of the Huberman lab for helpful comments on the manuscript. This

work was supported by a grant from the National Institute of General Medical Sciences of the

National Institutes of Health (NIH) under Award Number R35GM150926 to L.B.H. J.D.K. was

supported by NIH Ruth L. Kirschstein National Research Service Award Institutional Research

Training Grant 5T32AI145821-03. This work utilized the Cornell Institute of Biotechnology

Biotechnology Resource Center Genomics Facility (RRID:SCR_021727) and BioHPC

(RRID:SCR_021757) and the QB3 Genomics core at the University of California Berkeley

(RRID:SCR_022170).

REFERENCES

1. Kerkaert JD, Huberman LB. Regulation of nutrient utilization in filamentous fungi. Appl

Microbiol Biotechnol. 2023;107(19):5873-98. Epub 2023/08/04. doi: 10.1007/s00253-023-

12680-4. PubMed PMID: 37540250; PubMed Central PMCID: PMCPMC10983054.

2. Benocci T, Aguilar-Pontes MV, Zhou M, Seiboth B, de Vries RP. Regulators of plant

biomass degradation in ascomycetous fungi. Biotechnol Biofuels. 2017;10:152. Epub

2017/06/16. doi: 10.1186/s13068-017-0841-x. PubMed PMID: 28616076; PubMed Central

PMCID: PMCPMC5468973.

3. Wu VW, Thieme N, Huberman LB, Dietschmann A, Kowbel DJ, Lee J, et al. The

regulatory and transcriptional landscape associated with carbon utilization in a filamentous

fungus. Proc Natl Acad Sci U S A. 2020;117(11):6003-13. Epub 2020/03/01. doi:

10.1073/pnas.1915611117. PubMed PMID: 32111691; PubMed Central PMCID:

PMCPMC7084071.

4. Adnan M, Zheng W, Islam W, Arif M, Abubakar YS, Wang Z, et al. Carbon catabolite

repression in filamentous fungi. Int J Mol Sci. 2017;19(1). Epub 2018/01/04. doi:

10.3390/ijms19010048. PubMed PMID: 29295552; PubMed Central PMCID:

PMCPMC5795998.

5. Nehlin JO, Ronne H. Yeast *MIG1* repressor is related to the mammalian early growth

response and Wilms' tumour finger proteins. EMBO J. 1990;9(9):2891-8. Epub 1990/09/01.

PubMed PMID: 2167835; PubMed Central PMCID: PMC552004.

6. Arst HN, Jr., Cove DJ. Nitrogen metabolite repression in *Aspergillus nidulans*. Mol Gen

Genet. 1973;126(2):111-41. Epub 1973/11/02. doi: 10.1007/bf00330988. PubMed PMID:

4591376.

7. Dowzer CE, Kelly JM. Analysis of the *creA* gene, a regulator of carbon catabolite

repression in *Aspergillus nidulans*. Mol Cell Biol. 1991;11(11):5701-9. Epub 1991/11/01.

PubMed PMID: 1922072; PubMed Central PMCID: PMC361941.

8.   Hong Y, Cai R, Guo J, Zhong Z, Bao J, Wang Z, et al. Carbon catabolite repressor

MoCreA is required for the asexual development and pathogenicity of the rice blast fungus.

Fungal Genet Biol. 2021;146:103496. Epub 2020/12/09. doi: 10.1016/j.fgb.2020.103496.

PubMed PMID: 33290821.

9.   Strauss J, Mach RL, Zeilinger S, Hartler G, Stöffler G, Wolschek M, et al. Cre1, the

carbon catabolite repressor protein from *Trichoderma reesei*. FEBS Lett. 1995;376(1-2):103-7.

Epub 1995/11/27. doi: 10.1016/0014-5793(95)01255-5. PubMed PMID: 8521952.

10.   Sun J, Glass NL. Identification of the CRE-1 cellulolytic regulon in *Neurospora crassa*.

PloS One. 2011;6(9):e25654.

11.   Li Y, Steenwyk JL, Chang Y, Wang Y, James TY, Stajich JE, et al. A genome-scale

phylogeny of the kingdom Fungi. Curr Biol. 2021;31(8):1653-65.e5. Epub 20210218. doi:

10.1016/j.cub.2021.01.074. PubMed PMID: 33607033; PubMed Central PMCID:

PMCPMC8347878.

12.   Taylor JW, Berbee ML. Dating divergences in the Fungal Tree of Life: review and new

analyses. Mycologia. 2006;98(6):838-49. doi: 10.3852/mycologia.98.6.838. PubMed PMID:

17486961.

13.   Ainsworth GC. Ainsworth & Bisby's dictionary of the fungi: Cabi; 2008.

14.   Chen L, Tian X, Zhang L, Wang W, Hu P, Ma Z, et al. Brain glucose induces tolerance of

Cryptococcus neoformans to amphotericin B during meningitis. Nat Microbiol. 2024;9(2):346-58.

Epub 20240115. doi: 10.1038/s41564-023-01561-1. PubMed PMID: 38225460.

15.   Caza M, Hu G, Price M, Perfect JR, Kronstad JW. The zinc finger protein Mig1 regulates

mitochondrial function and azole drug susceptibility in the pathogenic fungus *Cryptococcus*

neoformans. mSphere. 2016;1(1). Epub 2016/06/16. doi: 10.1128/mSphere.00080-15. PubMed

PMID: 27303693; PubMed Central PMCID: PMCPMC4863601.

16.   Price MS, Betancourt-Quiroz M, Price JL, Toffaletti DL, Vora H, Hu G, et al.

Cryptococcus neoformans requires a functional glycolytic pathway for disease but not

persistence in the host. mBio. 2011;2(3):e00103-11. Epub 2011/06/10. doi:

10.1128/mBio.00103-11. PubMed PMID: 21652778; PubMed Central PMCID:

PMCPMC3110414.

17.   Opulente DA, LaBella AL, Harrison MC, Wolters JF, Liu C, Li Y, et al. Genomic factors

shape carbon and nitrogen metabolic niche breadth across Saccharomycotina yeasts. Science.

2024;384(6694):eadj4503. Epub 20240426. doi: 10.1126/science.adj4503. PubMed PMID:

38662846; PubMed Central PMCID: PMCPMC11298794.

18.   Kim J, Coradetti ST, Kim YM, Gao Y, Yaegashi J, Zucker JD, et al. Multi-omics driven

metabolic network reconstruction and analysis of lignocellulosic carbon utilization in

Rhodosporidium toruloides. Front Bioeng Biotechnol. 2020;8:612832. Epub 2021/02/16. doi:

10.3389/fbioe.2020.612832. PubMed PMID: 33585414; PubMed Central PMCID:

PMCPMC7873862.

19.   Yaegashi J, Kirby J, Ito M, Sun J, Dutta T, Mirsiaghi M, et al. *Rhodosporidium toruloides*:

a new platform organism for conversion of lignocellulose into terpene biofuels and bioproducts.

Biotechnol Biofuels. 2017;10:241. Epub 2017/10/28. doi: 10.1186/s13068-017-0927-5. PubMed

PMID: 29075325; PubMed Central PMCID: PMCPMC5651578.

20.   Wiebe MG, Koivuranta K, Penttilä M, Ruohonen L. Lipid production in batch and fed-

batch cultures of *Rhodosporidium toruloides* from 5 and 6 carbon carbohydrates. BMC

Biotechnol. 2012;12:26. Epub 2012/06/01. doi: 10.1186/1472-6750-12-26. PubMed PMID:

22646156; PubMed Central PMCID: PMCPMC3499381.

21.   Wu S, Hu C, Jin G, Zhao X, Zhao ZK. Phosphate-limitation mediated lipid production by

Rhodosporidium toruloides. Bioresour Technol. 2010;101(15):6124-9. Epub 2010/03/24. doi:

10.1016/j.biortech.2010.02.111. PubMed PMID: 20307977.

22.   Wu S, Zhao X, Shen H, Wang Q, Zhao ZK. Microbial lipid production by *Rhodosporidium

toruloides* under sulfate-limited conditions. Bioresour Technol. 2011;102(2):1803-7. Epub

2010/10/12. doi: 10.1016/j.biortech.2010.09.033. PubMed PMID: 20934330.

23.   Zhu Z, Zhang S, Liu H, Shen H, Lin X, Yang F, et al. A multi-omic map of the lipid-

producing yeast *Rhodosporidium toruloides*. Nat Commun. 2012;3:1112. Epub 2012/10/11. doi:

10.1038/ncomms2112. PubMed PMID: 23047670; PubMed Central PMCID: PMCPMC3493640.

24.   Lee JJ, Chen L, Shi J, Trzcinski A, Chen WN. Metabolomic profiling of *Rhodosporidium*

toruloides grown on glycerol for carotenoid production during different growth phases. J Agric

Food Chem. 2014;62(41):10203-9. Epub 2014/09/26. doi: 10.1021/jf502987q. PubMed PMID:

25254328.

25.   Mata-Gómez LC, Montañez JC, Méndez-Zavala A, Aguilar CN. Biotechnological

production of carotenoids by yeasts: an overview. Microb Cell Fact. 2014;13:12. Epub

2014/01/22. doi: 10.1186/1475-2859-13-12. PubMed PMID: 24443802; PubMed Central

PMCID: PMCPMC3922794.

26.   Coradetti ST, Pinel D, Geiselman GM, Ito M, Mondo SJ, Reilly MC, et al. Functional

genomics of lipid metabolism in the oleaginous yeast *Rhodosporidium toruloides*. Elife.

2018;7:e32110. Epub 2018/03/10. doi: 10.7554/eLife.32110. PubMed PMID: 29521624;

PubMed Central PMCID: PMCPMC5922974.

27.   Park YK, Nicaud JM, Ledesma-Amaro R. The engineering potential of *Rhodosporidium*

toruloides as a workhorse for biotechnological applications. Trends Biotechnol. 2018;36(3):304-

17. Epub 2017/11/15. doi: 10.1016/j.tibtech.2017.10.013. PubMed PMID: 29132754.

28.   Qin N, Li L, Ji X, Pereira R, Chen Y, Yin S, et al. Flux regulation through glycolysis and

respiration is balanced by inositol pyrophosphates in yeast. Cell. 2023;186(4):748-63.e15. Epub

20230208. doi: 10.1016/j.cell.2023.01.014. PubMed PMID: 36758548.

29.   Yu T, Liu Q, Wang X, Liu X, Chen Y, Nielsen J. Metabolic reconfiguration enables

synthetic reductive metabolism in yeast. Nat Metab. 2022;4(11):1551-9. Epub 20221027. doi:

10.1038/s42255-022-00654-1. PubMed PMID: 36302903; PubMed Central PMCID:

PMCPMC9684072.

30.   Rongpipi S, Ye D, Gomez ED, Gomez EW. Progress and opportunities in the

characterization of cellulose - an important regulator of cell wall growth and mechanics. Front

Plant Sci. 2018;9:1894. Epub 2019/03/19. doi: 10.3389/fpls.2018.01894. PubMed PMID:

30881371; PubMed Central PMCID: PMCPMC6405478.

31.   Dinh HV, Suthers PF, Chan SHJ, Shen Y, Xiao T, Deewan A, et al. A comprehensive

genome-scale model for *Rhodosporidium toruloides* IFO0880 accounting for functional

genomics and phenotypic data. Metab Eng Commun. 2019;9:e00101. Epub 20190828. doi:

10.1016/j.mec.2019.e00101. PubMed PMID: 31720216; PubMed Central PMCID:

PMCPMC6838544.

32.   Coradetti ST, Craig JP, Xiong Y, Shock T, Tian C, Glass NL. Conserved and essential

transcription factors for cellulase gene expression in ascomycete fungi. Proc Natl Acad Sci U S

A. 2012;109(19):7397-402.

33.   Stricker AR, Grosstessner-Hain K, Würleitner E, Mach RL. Xyr1 (xylanase regulator 1)

regulates both the hydrolytic enzyme system and D-xylose metabolism in *Hypocrea jecorina*.

Eukaryot Cell. 2006;5(12):2128-37. Epub 2006/10/24. doi: 10.1128/ec.00211-06. PubMed

PMID: 17056741; PubMed Central PMCID: PMCPMC1694815.

34.   Li Z, Yao G, Wu R, Gao L, Kan Q, Liu M, et al. Synergistic and dose-controlled

regulation of cellulase gene expression in *Penicillium oxalicum*. PLoS Genet.

2015;11(9):e1005509. Epub 2015/09/12. doi: 10.1371/journal.pgen.1005509. PubMed PMID:

26360497; PubMed Central PMCID: PMC4567317.

35.   van Peij NN, Gielkens MM, de Vries RP, Visser J, de Graaff LH. The transcriptional

activator XlnR regulates both xylanolytic and endoglucanase gene expression in *Aspergillus*

niger. Appl Environ Microbiol. 1998;64(10):3615-9. Epub 1998/10/06. PubMed PMID: 9758775;

PubMed Central PMCID: PMC106473.

36.   Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees for

large alignments. PLoS One. 2010;5(3):e9490. Epub 2010/03/13. doi:

10.1371/journal.pone.0009490. PubMed PMID: 20224823; PubMed Central PMCID:

PMCPMC2835736.

37.   Samal A, Craig JP, Coradetti ST, Benz JP, Eddy JA, Price ND, et al. Network

reconstruction and systems analysis of plant cell wall deconstruction by *Neurospora crassa*.

Biotechnol Biofuels. 2017;10:225. Epub 2017/09/28. doi: 10.1186/s13068-017-0901-2. PubMed

PMID: 28947916; PubMed Central PMCID: PMCPMC5609067.

38.   Bhat PJ, Murthy TV. Transcriptional control of the GAL/MEL regulon of yeast

Saccharomyces cerevisiae: mechanism of galactose-mediated signal transduction. Mol

Microbiol. 2001;40(5):1059-66. doi: 10.1046/j.1365-2958.2001.02421.x. PubMed PMID:

11401712.

39.   Kanehisa M, Goto S. KEGG: Kyoto encyclopedia of genes and genomes. Nucleic Acids

Res. 2000;28(1):27-30. Epub 1999/12/11. doi: 10.1093/nar/28.1.27. PubMed PMID: 10592173;

PubMed Central PMCID: PMCPMC102409.

40.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology:

tool for the unification of biology. The Gene Ontology Consortium. Nat Genet. 2000;25(1):25-9.

doi: 10.1038/75556. PubMed PMID: 10802651; PubMed Central PMCID: PMCPMC3037419.

41.   Gene Ontology Consortium. The Gene Ontology knowledgebase in 2026. Nucleic Acids

Res. 2025. Epub 20251218. doi: 10.1093/nar/gkaf1292. PubMed PMID: 41413728.

42.   Madeira F, Madhusoodanan N, Lee J, Eusebi A, Niewielska A, Tivey ARN, et al. The

EMBL-EBI Job Dispatcher sequence analysis tools framework in 2024. Nucleic Acids Res.

2024;52(W1):W521-W5. doi: 10.1093/nar/gkae241. PubMed PMID: 38597606; PubMed Central

PMCID: PMCPMC11223882.

43.   Nora LC, Wehrs M, Kim J, Cheng JF, Tarver A, Simmons BA, et al. A toolset of

constitutive promoters for metabolic engineering of *Rhodosporidium toruloides*. Microb Cell

Fact. 2019;18(1):117. Epub 2019/07/01. doi: 10.1186/s12934-019-1167-0. PubMed PMID:

31255171; PubMed Central PMCID: PMCPMC6599526.

44. Megnet R. Mutants partially deficient in alcohol dehydrogenase in *Schizosaccharomyces*

pombe. Arch Biochem Biophys. 1967;121(1):194-201. Epub 1967/07/01. doi: 10.1016/0003-

9861(67)90024-0. PubMed PMID: 4291962.

45. Marian IM, Vonk PJ, Valdes ID, Barry K, Bostock B, Carver A, et al. The transcription

factor Roc1 is a key regulator of cellulose degradation in the wood-decaying mushroom

Schizophyllum commune. mBio. 2022;13(3):e0062822. Epub 20220523. doi:

10.1128/mbio.00628-22. PubMed PMID: 35604096; PubMed Central PMCID:

PMCPMC9239231.

46. Wang Z, Li Y, Qiu H, Li Z, Ji T, Ren A, et al. GlSlt2 positively regulates GlMyb-mediated

cellulose utilization in *Ganoderma lucidum*. mBio. 2025;16(10):e0181225. Epub 20250908. doi:

10.1128/mbio.01812-25. PubMed PMID: 40919811; PubMed Central PMCID:

PMCPMC12506057.

47. Craig JP, Coradetti ST, Starr TL, Glass NL. Direct target network of the *Neurospora*

crassa plant cell wall deconstruction regulators CLR-1, CLR-2, and XLR-1. mBio.

2015;6(5):e01452-15. Epub 2015/10/16. doi: 10.1128/mBio.01452-15. PubMed PMID:

26463163; PubMed Central PMCID: PMC4620465.

48. Benocci T, Aguilar-Pontes MV, Kun RS, Seiboth B, de Vries RP, Daly P. ARA1 regulates

not only L-arabinose but also D-galactose catabolism in *Trichoderma reesei*. FEBS Lett.

2018;592(1):60-70. Epub 2017/12/08. doi: 10.1002/1873-3468.12932. PubMed PMID:

29215697.

49. Hynes MJ, Murray SL, Duncan A, Khew GS, Davis MA. Regulatory genes controlling

fatty acid catabolism and peroxisomal functions in the filamentous fungus *Aspergillus nidulans*.

Eukaryot Cell. 2006;5(5):794-805. Epub 2006/05/10. doi: 10.1128/ec.5.5.794-805.2006.

PubMed PMID: 16682457; PubMed Central PMCID: PMCPMC1459687.

50. Arentshorst M, Reijngoud J, Van Tol DJ, Reid ID, Arendsen Y, Pel HJ, et al. Utilization of

ferulic acid in *Aspergillus niger* requires the transcription factor FarA and a newly identified Far-

like protein (FarD) that lacks the canonical Zn (II) 2Cys6 domain. Frontiers in Fungal Biology.

2022;3:978845.

51. White AM, Mitchell AP. Control of citrate utilization by *Candida albicans* Adr1. mSphere.

2025;10(7):e0031125. Epub 20250611. doi: 10.1128/msphere.00311-25. PubMed PMID:

40497710; PubMed Central PMCID: PMCPMC12306173.

52. Show PL, Oladele KO, Siew QY, Aziz Zakry FA, Lan JC-W, Ling TC. Overview of citric

acid production from *Aspergillus niger*. Frontiers in life science. 2015;8(3):271-83.

53. Gallmetzer M, Meraner J, Burgstaller W. Succinate synthesis and excretion by

Penicillium simplicissimum under aerobic and anaerobic conditions. FEMS Microbiol Lett.

2002;210(2):221-5. doi: 10.1111/j.1574-6968.2002.tb11184.x. PubMed PMID: 12044678.

54. Yugueros SI, Peláez J, Stajich JE, Fuertes-Rabanal M, Sánchez-Vallet A, Largo-Gosens

A, et al. Study of fungal cell wall evolution through its monosaccharide composition: An insight

into fungal species interacting with plants. Cell Surf. 2024;11:100127. Epub 20240525. doi:

10.1016/j.tcsw.2024.100127. PubMed PMID: 38873189; PubMed Central PMCID:

PMCPMC11170279.

55. Beattie SR, Mark KMK, Thammahong A, Ries LNA, Dhingra S, Caffrey-Carr AK, et al.

Filamentous fungal carbon catabolite repression supports metabolic plasticity and stress

responses essential for disease progression. PLoS Pathog. 2017;13(4):e1006340. Epub

2017/04/20. doi: 10.1371/journal.ppat.1006340. PubMed PMID: 28423062; PubMed Central

PMCID: PMC5411099.

56. Ries LNA, Beattie S, Cramer RA, Goldman GH. Overview of carbon and nitrogen

catabolite metabolism in the virulence of human pathogenic fungi. Mol Microbiol.

2018;107(3):277-97. Epub 2017/12/03. doi: 10.1111/mmi.13887. PubMed PMID: 29197127;

PubMed Central PMCID: PMCPMC5777862.

57.   Ospina-Giraldo MD, Mullins E, Kang S. Loss of function of the *Fusarium oxysporum*

SNF1 gene reduces virulence on cabbage and *Arabidopsis*. Curr Genet. 2003;44(1):49-57.

Epub 2003/07/08. doi: 10.1007/s00294-003-0419-y. PubMed PMID: 12845476.

58.   Zhang S, Ito M, Skerker JM, Arkin AP, Rao CV. Metabolic engineering of the oleaginous

yeast *Rhodosporidium toruloides* IFO0880 for lipid overproduction during high-density

fermentation. Appl Microbiol Biotechnol. 2016;100(21):9393-405. Epub 2016/09/30. doi:

10.1007/s00253-016-7815-y. PubMed PMID: 27678117.

59.   Abbott EP, Ianiri G, Castoria R, Idnurm A. Overcoming recalcitrant transformation and

gene manipulation in Pucciniomycotina yeasts. Appl Microbiol Biotechnol. 2013;97(1):283-95.

Epub 20121114. doi: 10.1007/s00253-012-4561-7. PubMed PMID: 23149757.

60.   Colot HV, Park G, Turner GE, Ringelberg C, Crew CM, Litvinkova L, et al. A high-

throughput gene knockout procedure for *Neurospora* reveals functions for multiple transcription

factors. Proc Natl Acad Sci U S A. 2006;103(27):10352-7. Epub 2006/06/28. doi:

10.1073/pnas.0601456103. PubMed PMID: 16801547; PubMed Central PMCID: PMC1482798.

61.   McCluskey K, Wiest A, Plamann M. The Fungal Genetics Stock Center: a repository for

50 years of fungal genetics research. J Biosci. 2010;35(1):119-26. Epub 2010/04/24. PubMed

PMID: 20413916.

62.   Vogel H. A convenient growth medium for *Neurospora* (medium N). Microbial Genetics

Bulletin. 1956;13(4):42-3.

63.   Grigoriev IV, Nikitin R, Haridas S, Kuo A, Ohm R, Otillar R, et al. MycoCosm portal:

gearing up for 1000 fungal genomes. Nucleic Acids Res. 2014;42(Database issue):D699-704.

Epub 2013/12/04. doi: 10.1093/nar/gkt1183. PubMed PMID: 24297253; PubMed Central

PMCID: PMCPMC3965089.

64.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res. 2004;32(5):1792-7. Epub 20040319. doi: 10.1093/nar/gkh340.

PubMed PMID: 15034147; PubMed Central PMCID: PMCPMC390337.

65.   Tamura K, Stecher G, Kumar S. MEGA11: Molecular Evolutionary Genetics Analysis

Version 11. Mol Biol Evol. 2021;38(7):3022-7. doi: 10.1093/molbev/msab120. PubMed PMID:

33892491; PubMed Central PMCID: PMCPMC8233496.

66.   Patro R, Duggal G, Love MI, Irizarry RA, Kingsford C. Salmon provides fast and bias-

aware quantification of transcript expression. Nat Methods. 2017;14(4):417-9. Epub 2017/03/07.

doi: 10.1038/nmeth.4197. PubMed PMID: 28263959; PubMed Central PMCID:

PMCPMC5600148.

67.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21. Epub 20121025. doi:

10.1093/bioinformatics/bts635. PubMed PMID: 23104886; PubMed Central PMCID:

PMCPMC3530905.

68.   Putri GH, Anders S, Pyl PT, Pimanda JE, Zanini F. Analysing high-throughput

sequencing data in Python with HTSeq 2.0. Bioinformatics. 2022;38(10):2943-5. doi:

10.1093/bioinformatics/btac166. PubMed PMID: 35561197; PubMed Central PMCID:

PMCPMC9113351.

69.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550. Epub 2014/12/18. doi:

10.1186/s13059-014-0550-8. PubMed PMID: 25516281; PubMed Central PMCID:

PMCPMC4302049.

70.   Xu S, Hu E, Cai Y, Xie Z, Luo X, Zhan L, et al. Using clusterProfiler to characterize

multiomics data. Nat Protoc. 2024;19(11):3292-320. Epub 20240717. doi: 10.1038/s41596-024-

01020-z. PubMed PMID: 39019974.

71.   Lenth RVP, J. emmeans: Estimated Marginal Means, aka Least-Squares Means. R

package version 2.0.1 ed2025.