Computational Evaluation of Phytochemicals as Potential Anti-HIV Drugs Targeting CCR5 and CXCR4 Receptors

Sadman Sakib Nebir[1], Tawsif Al Arian[2], Bishajit Sarkar[3], Ripa Moni[3*], Salina Malek[3], Umme Salma Zohora[3], Mohammad Shahedur Rahman[3]

[1]Department of Pharmacology, Faculty of Basic Sciences, Bangladesh University of Health Sciences, Dhaka-1216, Bangladesh

[2]Department of Pharmacy, Jahangirnagar University, Dhaka-1342, Bangladesh

[3]Department of Biotechnology and Genetic Engineering, Faculty of Biological Sciences, Jahangirnagar University, Dhaka-1342, Bangladesh

Corresponding author: Ripa Moni

Email of the corresponding author: ripamoni@bgeju.edu.bd

Abstract

HIV is still a major worldwide health concern; hence new therapeutic approaches are needed to fight viral resistance and enhance treatment results. HIV entrance into host cells depends on the C-C chemokine receptor type-5 (CCR5) and C-X-C chemokine receptor type-4 (CXCR4), which makes them potential targets for antiviral medication development. The objective of this study is to computationally evaluate 53 phytochemicals that target the C-C chemokine receptor type-5 (CCR5) and C-X-C chemokine receptor type-4 (CXCR4) as potential anti-HIV medications. Effective anti-HIV medications were projected to be phytochemicals that may inhibit these receptors and so interfere with the HIV life cycle. AutoDock Vina was used to perform the molecular docking investigation from which six phytochemicals capable of inhibiting CCR5 and CXCR4 were identified based on the lowest docking score. The SWISSADME server was utilized to decide the druglikeness properties of these phytochemicals, the ADMETlab server to predict different pharmacokinetic and pharmacodynamic properties, the PASS-Way2Drug server to evaluate their activity spectra, and the RS-WebPredictor server to figure out the metabolism in the body. The six phytochemicals that were capable of suppressing CCR5 and CXCR4 were Withaferin A, Oleanolic Acid, Ursolic Acid, Theaflavine, Camptothecin, and Hypericin. The acceptable docking scores for these phytochemicals were less than -8.8 kcal/mol. They also adhered to Lipinski's rule of five and had promising ADME/toxicity study results, indicating their drug-likeness properties. Also, these compounds produced strong likelihood ratings (around 1.0) in activity spectra analysis and site of metabolism prediction, demonstrating their potency as powerful bioactive agents. Additionally, the molecules were exposed to sophisticated molecular dynamics simulation to assess their binding properties under a simulated physiological environment. Their performance in this investigation was efficient compared to the positive controls. According to the techniques and resources used in this study, the six phytochemicals might be used as alternative HIV therapeutic solutions.

Keywords: Antiviral agents, CCR5, CXCR4, Drug design, HIV, Phytochemicals, Molecular docking, Molecular Dynamics Simulation

1. Introduction

Human immunodeficiency virus (HIV) raids the body's immune system and weakens a person's defenses against many infections and some cancers that people with healthy immune systems can fight more easily. Infected individuals eventually lose their immunological protection and become immune-deficient as the virus kills and damages immune cells. CD4 cell count is commonly used to assess immune function (1). Approximately 40.1 million [33.6-48.6 million] people have died from HIV since the epidemic's inception, while 84.2 million [64.0-113.0 million] individuals have contracted the infection with HIV. At the end of 2021, there were 38.4 million [33.9-43.8 million] HIV-positive individuals worldwide. Moreover, 650000 people died of HIV-related illnesses worldwide in 2021 (2). Within the family Retroviridae, subfamily Orthoretrovirinae, the genus Lentivirus contains the human immunodeficiency virus (HIV) (3). Two identical single-stranded RNA molecules enclosed inside the virus particle's core comprise the HIV genome (3). HIV is a retrovirus. The genetic material is single-stranded RNA, not the double-stranded DNA found in human cells. Retroviruses also have reverse transcriptase, which can convert RNA into DNA and infect humans or host cells with that DNA. When HIV infects a cell, it first binds to the host cell and unites with it. The virus then uses the host cell mechanism to convert viral RNA into DNA and multiply. The new HIV copies then leave the host cell and spread to other cells. Binding and invasion into host cells are the first step in the human immune deficiency virus (HIV) replication cycle, which determines the tropism of the virus and the potential for HIV to impair the human immune system. HIV evades the host immune response by delivering its DNA into the host cell's cytoplasm through a multi-step process (4). In the first phase of the viral replication cycle, HIV entry starts with the virus adhering to the host cell and ends with the cell and viral membranes fusing and the viral core being delivered into the cytoplasm (4). The second step in viral entry and essential for infection is that the envelope (Env) protein binds to the host cell protein CD4, the primary receptor. HIV envelope (Env) protein is comprised of gp41 and gp120 subunits (5, 6), and Env is a highly glycosylated trimmer of the gp41 and gp120 heterodimers. Receptor binding is carried out by the gp120 subunit, which has five relatively conserved domains (C1–C5) and five variable loops (V1–V5), so termed for their relative genetic heterogeneity (4). CD4, an immunoglobulin superfamily member, typically enhances T-cell receptor (TCR) mediated signaling (7). We will focus on the third step of virus entry: co-receptor CCR5 (C-C chemokine receptor type 5) and CXCR4 (C-X-C chemokine receptor type four)

binding. Various HIV strain uses different co-receptors. R5 HIV uses the CCR5 chemokine receptor, X4 HIV uses CXCR4, and R5X4 HIV uses both co-receptors (8). There is no convincing evidence that co-receptors other than CCR5 and CXCR4 are important in supporting HIV-1 infection in vivo. With rare exceptions, only the R5 and R5X4 viruses are typically spread from person to person (9). Chemokine receptors are crucial during the initial stages of CD4+ cells' HIV-1 infection. While CCR5 permits entry of M-tropic HIV-1 strains, CXCR4 is the fusogenic receptor that favors entry of T-tropic HIV-1 strains (10). The fourth step in the virus entry process is moving the viral particle to where productive membrane fusion occurs. Recent research has demonstrated that several viruses usurp cellular transport systems to reach specific locations that are either required for infection or efficient entrance, and HIV may also do the same to reach locations where membrane fusion can take place (11-13). The chemokine receptor CCR5 is expressed on various cell types, including epithelial, endothelial, vascular smooth muscle, and fibroblasts; dendritic cells, macrophages, and memory T cells of the immune system; neurons, microglia, and astrocytes of the central nervous system (14). It was discovered in 1996 that CCR5 is required as a co-receptor for entry of the macrophage tropic HIV strains (15, 16). CCR5 ligands impede viral entry and envelope-mediated fusion (15). The virus prefers CCR5 during initial infection, whereas the alternative co-receptor 'C-X-C chemokine receptor type four' (CXCR4) is utilized considerably later in HIV infection, as the infected person approaches AIDS (17). Neutralizing antibodies, genetic modification, and preventing an integrated latent provirus are some of the fundamentally novel methods being developed for treating HIV-1. Nowadays, most medications target one of the HIV-1 enzymes, such as reverse transcriptase, protease, or integrase. NRTIs (Nucleoside and nucleotide reverse transcriptase inhibitors) and NNRTIs (non-nucleoside reverse transcriptase inhibitors) are the two categories into which reverse transcriptase inhibitors are typically divided (18). Protease inhibitors are a second significant class of clinically utilized inhibitors. In addition to inhibitors of the HIV-1 enzyme, inhibitors that affect other steps in the viral life cycle are being developed. Viral cell entry inhibitors used in HIV infection can be divided into inhibitors of viral-cell membrane fusion and inhibitors of viral envelope proteins' ability to attach to receptors (18). The G-protein coupled chemokine receptors CCR5 and CXCR4 determine the cellular tropism of HIV-1. The primary cell infection is specifically caused by viruses that preferentially employ CCR5 (R5 strains), as this co-receptor is mostly found in macrophages. On the other hand, viruses' ability to replicate themselves is due to their preference for interacting with CXCR4 (X4 strains).

This second co-receptor is mostly expressed in T-lymphocytes (19). These findings imply that CCR5 is a good target for HIV prevention or treatment and that even novel strategies, like anti-CCR5 vaccination, can offer vital scientific understanding and, more importantly, effective tools for the battle against HIV and other immune-based diseases. It has been demonstrated that several types of phytochemicals prevent HIV replication, and various phytochemicals that target the HIV infection cycle are investigated here. Natural products originating from plants continue to be a source for developing novel medicines, such as anti-HIV agents. Despite significant advances in HIV chemotherapy, there is still a need for novel anti-HIV drug development, and medicinal plants can be beneficial in this effort. Several plant species, particularly *Artemisia annua*, *Garcinia edulis*, *Justicia gendarussa*, *Phyllanthus pulcher*, *Rhus chinensis*, *Smilax corbularia*, *Terminalia paniculata*, and *Tuberaria lignosa*, have demonstrated notable anti-HIV activity (20). As a result, the use of plant-based alternatives holds enormous promise for the treatment of HIV patients. Now, the focus is shifting to using natural compounds, such as plant-based alternatives, to produce anti-HIV medications. Although HIV/AIDS is not a new disease, clinical trials are currently being conducted on research based on plant-derived products. Therefore, this computational study aimed to analyze antiviral phytochemicals as potential treatment options for HIV.

Table 01. List of the plant-derived anti-HIV agents that work via CCR5 and CXCR4

Serial	Name of the agents	PubChem CID	Reference
1	Ellagic acid	5281855	(21)
2	Gallic acid	370	(21)
3	Epigallocatechin gallate	65064	(22)
4	Quercetin	5280343	(23)
5	Quercetrin	5280459	(23)
6	Indirubin	10177	(24)
7	Fisetin	5281614	(25)
8	Naringenin	439246	(26)
9	Luteolin	5280445	(27)
10	Apigenin	5280443	(28)

11	Camptothecin	24360	(29)
12	Daidzein	5281708	(30)
13	Kaempferol	5280863	(31)
14	6-Gingerol	442793	(32)
15	Curcumin	969516	(33)
16	Palmatine	19009	(34)
17	Berberine	2353	(35)
18	Magnoflorine	73337	(36)
19	Lanuginosine	97622	(37)
20	Liriodenine	10144	(38)
21	Anonaine	160597	(39)
22	Ursolic acid	64945	(40)
23	Eugenol	3314	(41)
24	Cordycepin	6303	(42)
25	Harmaline	3564	(43)
26	Harmine	5280953	(44)
27	Lycorine	72378	(45)
28	Oliverine	24198103	(46)
29	Hesperetin	72281	(47)
30	Oleanolic acid	10494	(48)
31	Myricetin	5281672	(49)
32	Imperatorin	10212	(50)
33	Bergapten	2355	(51)
34	Silybin	31553	(52)
35	Withaferin A	265237	(53)
36	Cordioside	101915817	(54)

37	Neoandrographolide	9848024	(55)
38	Rotenone	6758	(56)
39	Calanolide A	64972	(57)
40	Pericalline	6436240	(58)
41	Ferruginol	442027	(59)
42	Allicin	65036	(60)
43	Genistein	5280961	(61)
44	Scutellarein	5281697	(62)
45	Capsaicin	1548943	(63)
46	Theaflavin	135403798	(64)
47	Hypericin	3663	(65)
48	Methyl rosmarinate	6479915	(66)
49	Licoleafol	11111496	(67)
50	Nimocinol	178770	(68)
51	Isomeldenin	76316558	(69)
52	Tembetarine	167718	(70)
53	Tinosponone	15215479	(71)

Figure 01. The 3D structures of the target proteins: 1. CCR5 Chemokine Receptor (PDB ID: 4MBS) (72); 2. CXCR4 chemokine receptor in complex (PDB ID: 3ODU) (73). The 3D structures were visualized by Protein Data Bank 3D viewer.

1.1 Role of CCR5 and CXCR4 in terms of HIV infection

Immune cells, such as CD4+ T-cells, essential in the immunological response against viruses, bacteria, and other pathogens, have the protein CCR5 on their surface. CCR5, CCR5, also known as CCCKR-5 or CKR5, is engaged mainly in immune surveillance, inflammatory response, tumor development and metastasis (20)(72)(73), etiology of inflammatory disorders (74)(75)(76), asthma (77)(78), and cancer (72, 73). It is vital in the recruitment of immune cells to inflammatory sites by guiding immune cell movement (chemotaxis) along the chemokine gradient (79) (80). CCR5 modulates the trafficking and effector activities of memory/effector T lymphocytes, macrophages, and immature dendritic cells (81). Aside from its direct role in immune process mediation, it also suppresses learning, memory, and synaptic connections in the brain (82). The human immunodeficiency virus (HIV) uses CCR5 as a co-receptor in order to enter and infect CD4+ T-cells. CCR5 is a seven-transmembrane G protein-coupled receptor (GPCR) that belongs to the class A GCPR family. CCR5 is a GPCR with seven transmembrane a-helices, three extracellular loops, three intracellular loops, an amino-terminal domain, and a carboxyl-terminal domain (83).

HIV is an RNA virus that attaches to CD4 on the surface of target cells through the surface glycoprotein gp120. HIV cannot enter the cell through this contact; it needs a co-receptor, CCR5 or CXCR4, to finish the entrance process. The majority of HIV strains, sometimes referred to as R5-tropic HIV, employ CCR5 as their co-receptor. Natural CCR5 ligands contain chemokines (small chemoattractant cytokines) involved in innate immunity, which are natural suppressors of HIV-1 infection (84-87): macrophage inflammatory proteins CCL3 (MIP-1 a) and CCL4 (MIP-1 b), CCL5 (RANTES - regulated on activation, normal T-cell expressed and secreted), and CCL3L1, the most potent among CCR5 (88). CCL7 (MCP-3) is the main antagonist ligand of the CCR5 receptor (89). The CCR5 receptor is activated by its agonist ligands, which increase cell migration and mediate inflammatory reactions. In summary, CCR5 is a co-receptor essential for HIV entrance into target cells (Figure 3).

Figure 2. Extracellular virions penetrate their target cell through a three-step process: binding to the CD4 receptor, binding to the CCR5 or CXCR4 coreceptors, or both, and membrane fusion. The HIV reverse transcriptase enzyme catalyzes HIV RNA transcription into double-stranded HIV DNA, a phase blocked by nucleoside analogs and non-nucleoside reverse transcriptase inhibitors (NNRTIs). HIV integrase is an enzyme that aids in integrating HIV DNA into host chromosomes. Immature virions are created and bud from the cell surface following transcription and translation of the HIV genome. Since the HIV protease enzyme cleaves polypeptide chains, the virus can mature.

CXCR4, commonly known as "fusin," is one of the most well-studied chemokine receptors due to its previously discovered role as an HIV entrance coreceptor (87). Although CXCR4 is known to bind only CXCL12, another chemokine receptor, CXC receptor 7 (CXCR7, ACKR3, RDC1, CMKOR1, or GPR159), was identified in 2005 as a CXCL12 receptor (90). Further research will

be necessary to determine the precise involvement of the CXCR4-CXCR7-CXCL12 axis in cell migration (87). Certain chemokines, including MIP-1a, MIP-1b, and RANTES, were identified to suppress HIV-1 infection, indicating that the chemokine receptors may be a coreceptor for HIV-1 entrance (91). It has been demonstrated to have an essential role in the pathogenesis of HIV infection. Soon after CCR5, the a-chemokine receptor CXCR4 was identified as the predominant coreceptor for T-tropic HIV-1 isolates that can replicate successfully in T-cell lines and primary CD4+ T cells (88).

CXCR4 and CCR5 chemokine receptors are coreceptors for HIV-1 entrance into CD4+ cells. At the beginning stages of HIV infection, viral segregates frequently utilize CCR5, while later disconnects typically use CXCR4. The pattern of expression and regulation of these chemokine receptors on T cell subsets has crucial implications for AIDS pathogenesis and lymphocyte recirculation.

2. Materials and Methods

2.1 Target Selection and Preparation

In this work, the CCR5 Chemokine Receptor (4MBS) and the CXCR4 Chemokine Receptor (3ODU) were chosen as possible targets. PDB structures of targets with bound ligands were obtained (Protein Data Bank). Structures lacking the physiologically required cofactor and heteroatom were rejected, as were those lacking side chains. By preserving the optimum physiological pH, hydrogen atoms were introduced to the target. Apart for those essential to the aim hypothesis, all water molecules were eliminated. The Kollman Charges were used to determine the partial charges. Nonpolar hydrogens were merged since it is a mechanism for representing unified atoms. AutoGrid was used to generate grid maps, which display a 3D grid of interaction energies with the target.

2.2 Ligand Selection and Preparation

Potential ligands were identified and obtained from PubChem using 3D SDF format structures. As AutoDock Vina cannot read the SDF format, the PyMOL tool was used to convert it to PDB format. All of the PDB files were then converted to PDBQT files and loaded into AutoDock in order to dock with the required targets. Positive control for 4MBS and 3ODU, Maraviroc (Positive control, PubChem CID 3002977), and Plerixafor (Positive control, PubChem CID 65015), were prepared using the same process.

2.3 Receptor Grid Generation

Grid usually confines the active site to shorten specific areas of the receptor protein so that the ligand can dock specifically. In Glide, a grid was generated using the default Van der Waals radius scaling factor of 1.0 and charge cutoff of 0.25. The spacing and npts were set to a default of 0.375 and 40. The energy range and exhaustiveness were 4 and 8, the default value. A cubic box is available (reference ligand active site). Then, the grid box volume was adjusted to $168.307\times 116.536\times 39.126$ for the docking test.

2.4 Pass Prediction:

PASS-Way2Drug (http://way2drug.com/passonline/) online server was used to predict the PASS (Prediction of Activity Spectra for Substances) values of the best selected phytochemicals or

ligands (74). This server can predict over 4000 different types of biological, pharmacological, metabolic, pharmacokinetic, and toxic characteristics of chemical compounds by analyzing its database consisting of 30000 biologically active compounds (75). The canonical smiles of these phytochemicals were used in the PASS analysis, and the probability of a compound being active or the Pa threshold was kept greater than 0.7 or 70% to increase the reliability of the predictions. Thereafter, SMARTCyp (https://smartcyp.sund.ku.dk/mol_to_som) online server was used to carry out the P450 site of metabolism or SOM prediction (76). The server utilizes the following standard equation for conducting the predictions:

Score, $S = E - 8*A - 0.04*SASA$

Where S represents the prediction score, E represents the activation energy score, A is the accessibility measure of the compound in the query determined based on its topological features, and SASA is the Solvent Accessible Surface Area that is also measured from the topological characteristics of the query compound. The lower the prediction score and energy score are, the better the reliability of the prediction is, and vice versa (45). Canonical smiles of the phytochemical were again used in the server to perform the predictions.

2.5 Ligand-based drug-likeness property analysis and ADME/toxicity prediction:

SWISSADME server (http://www.swissadme.ch/) was used to assess the drug-likeness properties of the nine selected ligand molecules (95). In order to predict their various pharmacokinetic and pharmacodynamic properties, the ADMET analysis for each of the ligand molecules was carried out using online-based servers, admetSAR (http://lmmd.ecust.edu.cn/admetsar2/) and ADMETlab (http://admet.scbdd.com/) (96, 97). Both the servers determined the absorption, distribution, metabolism, excretion, and toxicity properties. The numeric and categorical values of the results given by the ADMETlab server were changed into qualitative values according to the explanation and interpretation described in the ADMETlab server (http://admet.scbdd.com/home/interpretation/) for the convenience of interpretation.

2.6 Molecular Dynamic Simulation

Molecular dynamics is a sophisticated automated simulation approach used for evaluating the degree of stability of protein and protein-ligand complex structure at the minuscule stage through demonstrating the behavioral characteristics, interacting arrangement, fluctuation, physical

foundation of function, and structure. The molecular dynamics (MD) simulations were conducted using GROMACS version 2023.1 to investigate the behavior of both Apo protein and Protein-ligand complexes (77). The protein was parameterized for its protein content using the CHARMM General Force Field. The SwissParam server was utilized to perform the ligand topologies (78). The structures underwent 2500 cycles of vacuum minimization using the steepest descent method in order to mitigate any potential steric issues. The solvation of the structure was accomplished through the utilization of the Simple Point Charge (SPC) water model. Subsequently, the system was rendered neutral through the introduction of Na+ and Cl- ions utilizing the gmx genion tool. This measure was implemented in order to maintain the overall electrical neutrality of the system. After minimization, three stages were led in the MD simulation: NVT, NPT, and production. The equilibration of the systems was conducted in two phases. Initially, an NVT equilibration lasting 100 picoseconds was conducted to attain a steady state of the number of particles, volume, and temperature. The purpose of this step was to elevate the system to a temperature of 300 Kelvin. The second step involved conducting a 100 picosecond NPT equilibration, which aimed to maintain the system's pressure and density stability by ensuring an equal number of particles, pressure, and temperature. The simulations involved the imposition of bond constraints on all bonds within the protein, thereby inducing position restraint of the protein group. The constrained conditions of NVT and NPT resulted in the relaxation of water molecules surrounding the protein, leading to a decrease in system entropy. The dynamics were conducted utilizing the Parrinello-Rahman barostat and the v-rescale thermostat (79). The relaxation of the barostat and thermostat persisted for a duration of 100 picoseconds. The application of Linear Constraint Solver algorithm was used to impose constraints on the covalent bonds. The Particle-Mesh Ewald (PME) method was utilized to process the electrostatic interactions. After reaching equilibrium, every framework underwent a production run that lasted for 100 nanoseconds (ns) of simulation time.

3. Results and Discussion

3.1 Molecular Docking

The study conducted molecular docking of identified ligand molecules with their receptor proteins, finding that molecules with the least binding energy were most effective in hindering their receptors due to their expanded binding affinity (Table 2) (89). Among the ligands, Ursolic acid, Oleanolic acid, Withaferin A, Hypericin (inhibit CCR5) and Camptothecin, Ursolic acid, Oleanolic acid, Theaflavin, and Hypericin (inhibit CXCR4), were chosen for their lower free binding energy and comparable binding energy to respective controls Maraviroc and Plerixafor (table 2). An efficient method for computer-aided drug design is molecular docking, which uses particular algorithms to assign affinity scores based on the positions of ligands inside target binding pockets. Since the complex will spend more time in contact, the lowest docking score indicates the highest affinity (90, 91). Strengthening the receptor-ligand interactions requires both hydrophobic and hydrogen interactions (92). Multiple hydrogen and hydrophobic interactions were predicted to be formed by the six best-selected ligands and the other ligands with the target molecules (Figure 9 and Table 2 & 12). Therefore, Ursolic acid, Oleanolic acid, Withaferin A, and Hypericin, Camptothecin, and Theaflavin ligands from each receptor category were chosen and assessed in the subsequent phases of the study.

Table 2. Docking Scores

Receptor	Ligands	Docking Scores / Binding energy (kcal/mol)
CCR5 (PDB: 4MBS)	Ellagic acid	-7.2
	Gallic acid	-5.3
	Epigallocatechin gallate	-7.5
	Quercetin	-7.2
	Quercetrin	-7.3
	Indirubin	-8.4
	Fisetin	-7.4
	Naringenin	-7.4

	Luteolin	-7.4
	Apigenin	-7.5
	Camptothecin	-8.0
	Daidzein	-7.5
	Kaempferol	-7.5
	6-Gingerol	-6.2
	Curcumin	-6.7
	Palmatine	-7.3
	Berberine	-7.7
	Magnoflorine	-7.6
	Lanuginosine	-7.9
	Liriodenine	-8.3
	Anonaine	-8.0
	Ursolic acid	-9.0
	Eugenol	-6.2
	Cordycepin	-6.1
	Harmaline	-6.4
	Harmine	-6.6
	Lycorine	-7.2
	Oliverine	-7.6
	Hesperetin	-7.7
	Oleanolic acid	-9.0
	Myricetin	-7.0
	Imperatorin	-8.1
	Bergapten	-7.2
	Silybin	-8.1
	Withaferin A	-9.2
	Cordioside	-7.3
	Neoandrographolide	-7.9
	Rotenone	-8.7

	Calanolide A	-7.9
	Pericalline	-7.6
	Ferruginol	-8.3
	Allicin	-4.1
	Genistein	-6.8
	Scutellarein	-7.0
	Capsaicin	-7.1
	Theaflavin	-8.7
	Hypericin	-9.0
	Methyl rosmarinate	-6.9
	Licoleafol	-7.9
	Nimocinol	-8.1
	Isomeldenin	-8.1
	Tembetarine	-7.2
	Tinosponone	-7.7
	Maraviroc	-9.0
CXCR4 (PDB: 3ODU)	Ellagic acid	-7.5
	Gallic acid	-6.1
	Epigallocatechin gallate	-8.0
	Quercetin	-7.5
	Quercetrin	-7.8
	Indirubin	-8.0
	Fisetin	-7.5
	Naringenin	-8.1
	Luteolin	-7.9
	Apigenin	-7.5
	Camptothecin	-9.2
	Daidzein	-7.6
	Kaempferol	-7.4
	6-Gingerol	-5.6

	Curcumin	-7.3
	Palmatine	-7.0
	Berberine	-8.1
	Magnoflorine	-8.2
	Lanuginosine	-8.3
	Liriodenine	-8.9
	Anonaine	-8.7
	Ursolic acid	-9.2
	Eugenol	-5.3
	Cordycepin	-6.2
	Harmaline	-7.0
	Harmine	-7.0
	Lycorine	-8.1
	Oliverine	-7.7
	Hesperetin	-7.2
	Oleanolic acid	-9.3
	Myricetin	-7.9
	Imperatorin	-7.6
	Bergapten	-6.9
	Silybin	-8.7
	Withaferin A	-8.5
	Cordioside	-8.5
	Neoandrographolide	-7.4
	Rotenone	-8.0
	Calanolide A	-8.0
	Pericalline	-8.6
	Ferruginol	-7.8
	Allicin	-3.6
	Genistein	-7.6
	Scutellarein	-8.2

	Capsaicin	-6.7
	Theaflavin	-9.0
	Hypericin	-10.1
	Methyl rosmarinate	-6.9
	Licoleafol	-7.9
	Nimocinol	-8.3
	Isomeldenin	-8.5
	Tembetarine	-7.0
	Tinosponone	-8.0
	Plerixafor	-8.1

3.2 Drug-likeness properties

Druglikeness properties of a chemical compound determine the ability of that compound to act as a drug in humans based on its biological and pharmacological properties. Lipinski's rule of five is a set of widely used rules that are used to evaluate a chemical's drug-likeness properties. According to Lipinski's rule of five, an active drug molecule should violate no more than one of these primary criteria: Molecular weight <500 g/mol or Daltons, Num. H-bond donors ≤5, Num. H-bond acceptors ≤10 and partition coefficient log Po/w ≤5. To improve the prediction accuracy, sometimes molar refractivity from 40 to 130 is also considered as a rule of Lipinski's rule of five (93, 94). According to our study, withaferin A and camptothecin followed all the primary criteria of Lipinski's rule of five, reflecting the fact that they might be the best phytochemicals to combat HIV infection among the selected ligands. On the other hand, both hypericin and theaflavin violated more than one primary criterion. Hypericin has a molecular weight of 504.44 g/mol and 06 H-bond donors. Theaflavin has a molecular weight of 564.49 g/mol, 12 H-bond acceptors, and 09 H-bond donors. Other ligands, ursolic acid and oleanolic acid, violated only one primary rule each, and for this reason, they were also found to contain the properties of active drugs. These results are also in agreement with the results generated by the positive controls. Maraviroc followed Lipinski's rule of five with one violation; however, Plerixafor was predicted to be not very active as it violated two primary criteria of Lipinski's rule of five. Considering the molar refractivity, although it is not a primary criterion, only withaferin A and camptothecin were found

to have their values within the acceptable range of 40 to 130. For this reason, among the ligands on which the druglikeness property analysis was conducted, withaferin A and camptothecin can be considered the best ligands or phytochemicals for treating HIV infections. The results of the druglikeness property analysis are listed in Table 3.

Table 3. Drug-likeness properties of selected compounds

Drug-likeness properties	Rules	Ursolic acid	Oleanolic acid	Withaferin A	Hypericin	Camptothecin	Theaflavin	Maraviroc	Plerixafor
Lipinski's rule of five	-	Yes	Yes	Yes	No	Yes	No	Yes	No
Molecular weight (g/mol)	<500	456.70	456.70	470.60	504.44	348.35	564.49	513.67	502.78
Log Po/w	≤5	5.88	6.06	3.45	4.26	2.20	1.31	4.81	0.68
Num. H-bond acceptors	≤10	03	03	06	08	05	12	06	08
Num. H-bond donors	≤5	02	02	02	06	01	09	01	06
Molar refractivity	40 - 130	136.91	136.65	127.49	144.83	95.31	143.98	145.84	182.21

3.3 ADME/T Analysis

ADMET virtual test was conducted in this study to predict the absorption, distribution, metabolism, excretion, and toxicity of the best selected ligands or phytochemicals (Table 4). The Caco-2 permeability represents the permeability and absorption of a chemical compound through the intestinal cell barrier (95). The study found that camptothecin and maraviroc, a positive control, showed predicted Caco-2 permeability, a key efflux membrane transporter in the body, hindering therapeutic agent uptake. So, limiting the activity of p-gp proteins is desirable for the therapeutic

delivery of a drug (90). The study found that withaferin A, camptothecin, theaflavin, and maraviroc inhibited p-gp protein, with camptothecin being the p-gp substrate. Other ligands and positive control faced no issues during cellular uptake. Plasma protein binding (PPB) is a parameter used in deciding the distribution fate of a drug and the lesser bound a drug is, the more effectively it can be distributed to its target organs (91). Camptothecin exhibits undesirable plasma protein binding, while ligands and controls are BBB permeable. Cytochrome P450 enzymes are crucial for drug metabolization, with CYP450 substrates more effectively metabolized. Furthermore, CYP450 inhibition can have beneficial or harmful outcomes depending on the pharmacokinetic properties of the specific drug molecule (92). The study found that withaferin A, a phytochemical, was found to be a substrate for two CYP450 enzymes, indicating that it provided better metabolic activity results than other phytochemicals. The study found that hypericin had the highest excretion half-life of 3.306 hours, followed by theaflavin. Ursolic acid and oleanolic acid were predicted to be non-hERG blockers, hepatotoxic, non-Ames mutagenic, and non-toxic to the liver. Camptothecin and theaflavin were found to be toxic in three parameters. Considering all the parameters of ADMET analysis, withaferin A could be declared as the best-performing phytochemical, followed by ursolic acid and oleanolic acid.

Table 4. ADMET Analysis

ADMET Class	Properties	Ursolic acid	Oleanolic acid	Withaferin A	Hypericin	Camptothecin	Theaflavin	Maraviroc	Plerixafor
Absorption	Caco-2 permeability	Negative	Negative	Negative	Negative	Positive	Negative	Positive	Negative
	Pgp-inhibitor	Negative	Negative	Positive	Negative	Positive	Positive	Positive	Negative
	Pgp-substrate	Negative	Negative	Negative	Negative	Positive	Negative	Negative	Negative
	Human Intestinal Absorption (HIA)	Positive	Positive	Positive	Negative	Negative	Negative	Positive	Positive

Distribution	Plasma Protein Binding (PPB)	Less significant	Less significant	Less significant	Less significant	Significant	Less significant	Less significant	Less significant
	Blood Brain Barrier (BBB)	Positive	Positive	Positive	Positive	Positive	Positive	Positive	Positive
Metabolism	CYP450 1A2 inhibition	Negative	Negative	Negative	Positive	Negative	Negative	Negative	Negative
	CYP450 1A2 substrate	Negative	Negative	Negative	Negative	Negative	Negative	Negative	Positive
	CYP450 3A4 inhibition	Negative	Negative	Negative	Negative	Positive	Negative	Positive	Negative
	CYP450 3A4 substrate	Positive	Positive	Positive	Negative	Positive	Positive	Positive	Negative
	CYP450 2C9 inhibition	Negative	Negative	Negative	Negative	Negative	Negative	Negative	Negative
	CYP450 2C9 substrate	Negative	Negative	Negative	Negative	Negative	Negative	Negative	Negative
	CYP450 2C19 inhibition	Negative	Negative	Negative	Negative	Negative	Negative	Negative	Negative
	CYP450 2C19 substrate	Negative	Negative	Positive	Negative	Negative	Negative	Positive	Positive
	CYP450 2D6 inhibition	Negative	Negative	Negative	Negative	Negative	Negative	Positive	Positive

	CYP450 2D6 substrate	Negative	Negative	Negative	Negative	Negative	Negative	Negative	Positive
Excretion	T1/2 (h)	1.754 (h)	1.735 (h)	1.709 (h)	3.306 (h)	1.664 (h)	2.345 (h)	1.759 (h)	2.038 (h)
Toxicity	hERG blockers	Negative	Negative	Negative	Negative	Positive	Positive	Negative	Positive
	Human Hepatotoxicity (HHT)	Negative	Negative	Positive	Negative	Positive	Negative	Positive	Positive
	Ames mutagenicity	Negative	Negative	Negative	Negative	Negative	Positive	Negative	Negative
	Drug-Induced Liver Injury (DILI)	Negative	Negative	Negative	Positive	Positive	Positive	Positive	Negative

3.4 Pass Prediction

PASS forecast is a technique for investigating the natural exercises of drug-like molecules by considering two likelihood factors, i.e., Dad addressing the likelihood of a compound being dynamic and Pi addressing the possibility of a compound being dormant. Any Pa value more noteworthy than 0.7 portrays a higher probability of a compound showing a specific action or trademark. Hence, the Pa>0.7 edge gives a profoundly solid movement forecast (82). The results of the PASS prediction analysis of this study are listed in Table 5.

This research showed that apart from potential anti-HIV properties (based on the molecular docking study), ursolic acid, oleanolic acid, and withaferin A might have hepatoprotective activity. We can assume that at a certain dosage, these phytochemicals can exert their protective activities on the liver as phytochemicals are proven to have hepatoprotective activities at a certain dosage (96). The phytochemicals were also found to have antitumor and anticancer activities. For

example, ursolic acid, oleanolic acid, and camptothecin were predicted to have the capability to act as Caspase-3 and Caspase-8 stimulants. Caspases mediate programmed cell death or apoptosis and thus depending on the cellular context, they are antitumorigenic in nature. Their activity is used as a marker for measuring the efficacy of cancer therapy (97). Again, hypericin and camptothecin were predicted to inhibit the expression of HIF1A and enhance the expression of TP53. Studies have found that the HIF1A expression increases in a wide variety of cancers in humans (98, 99). TP53 is a tumor suppressor and it is the most commonly mutated gene in many human cancers (100). Camptothecin can be declared as the best phytochemical according to the PASS prediction analysis as it was predicted to exhibit antineoplastic activities against multiple human cancers and found to possess HIF1A expression inhibitory, TP53 expression enhancing, and Caspase 3 stimulating properties.

Table 5. Pass Prediction Analysis

Ursolic acid	Oleanolic acid	Withaferin A	Hypericin	Camptothecin	Theaflavin	Maraviroc	Plerixafor
Insulin promoter (0.970)	Caspase 3 stimulant (0.984)	Antineoplastic (0.916)	CYP2C12 substrate (0.898)	Topoisomerase I inhibitor (0.942)	UGT1A substrate (0.976)	HIV attachment inhibitor (0.906)	Nicotinic alpha2beta2 receptor antagonist (0.857)
Hepatoprotectant (0.961)	Transcription factor NF kappa B stimulant (0.954)	Hepatoprotectant (0.904)	Membrane integrity agonist (0.889)	HIF1A expression inhibitor (0.936)	Membrane integrity agonist (0.972)	CC chemokine receptor antagonist (0.830)	5 Hydroxytryptamine release stimulant (0.858)
Transcription factor NF kappa B stimulant (0.927)	Transcription factor stimulant (0.954)	Macular degeneration treatment (0.887)	Ubiquinol-cytochrome-c reductase inhibitor (0.877)	Antineoplastic (0.929)	HMOX1 expression enhancer (0.966)	CC chemokine 5 receptor antagonist (0.718)	Nicotinic alpha6beta3beta4 alpha5 receptor antagonist (0.828)

Transcription factor stimulant (0.927)	Chemopreventive (0.937)	Antieczematic (0.876)	Alkane 1-monooxygenase inhibitor (0.870)	DNA intercalator (0.895)	CYP1A1 substrate (0.961)	-	Antidepressant (0.771)
Antiprotozoal (Leishmania) (0.915)	Hepatoprotectant (0.930)	Hepatic disorders treatment (0.865)	TP53 expression enhancer (0.846)	Antineoplastic (colorectal cancer) (0.839)	UGT1A6 substrate (0.954)	-	Mood disorders treatment (0.768)
Caspase 3 stimulant (0.912)	Membrane integrity antagonist (0.928)	Immunosuppressant (0.850)	Chlordecone reductase inhibitor (0.844)	Antineoplastic (solid tumors) (0.835)	CYP1A substrate (0.955)	-	Antiischemic, cerebral (0.759)
Apoptosis agonist (0.890)	Caspase 8 stimulant (0.914)	Transcription factor NF kappa B inhibitor (0.777)	HIF1A expression inhibitor (0.839)	Antineoplastic (colon cancer) (0.833)	Antimutagenic (0.940)	-	Mucomembranous protector (0.744)
Membrane integrity antagonist (0.885)	Oxidoreductase inhibitor (0.904)	Antineoplastic (breast cancer) (0.722)	Antineoplastic (0.836)	Antineoplastic (small cell lung cancer) (0.796)	UDP-glucuronosyltransferase substrate (0.927)	-	Phobic disorders treatment (0.757)
Diacylglycerol O-acyltransferase inhibitor (0.882)	Apoptosis agonist (0.901)	Phosphatase inhibitor (0.725)	CYP1A inducer (0.828)	Caspase 3 stimulant (0.772)	Reductant (0.925)	-	Testosterone 17beta-dehydrogenase (NADP+) inhibitor (0.731)
Oxidoreductase inhibitor (0.876)	Phospholipase C inhibitor (0.894)	Oxidoreductase inhibitor (0.707)	Membrane permeability inhibitor (0.831)	TP53 expression enhancer (0.778)	Pectate lyase inhibitor (0.916)	-	-

3.5 P450 SOM Prediction

The P450 site of metabolism (SOM) analysis was conducted to find out the sites of the selected phytochemicals that are more likely to be metabolized by CYP450 3A4, CYP450 2D6, and CYP450 2C9 (Table 6). Emphasis was mainly given on the lowest activation energy and lowest score because lower (activation) energy and lower score represent the higher probability of a site of a molecule being metabolized. Therefore, the sites and the molecules having lower energies and lower scores tend to be better metabolized by the CYP450 enzymes (101, 102). The P450 SOM analysis generated results that are very much in agreement with the metabolism part of the previously done ADMET analysis. According to the SOM analysis, withaferin A showed the best results with its predicted lowest energy of 52.4 and the lowest score of 43.4 CYP450 3A4 section. Therefore, withaferin A should be better metabolized by the CYP450 3A4 enzyme among the selected phytochemicals. In the ADMET analysis, withaferin A also generated the best results among the selected phytochemicals. Furthermore, according to the ADMET analysis, hypericin was the only phytochemical that might not be metabolized well by the CYP450 3A4 enzyme. According to the SOM analysis, hypericin provided the highest energy and highest score, representing its less active nature than the CYP450 3A4 enzyme. Again, according to the ADMET analysis, all the phytochemicals showed poor performance in terms of metabolizing by the enzymes compared to the positive controls. Likewise, the SOM analysis also showed that the energies and the scores generated by the phytochemicals were much higher than those of the positive controls, reflecting their unsatisfactory performance relative to the positive controls. And among the phytochemicals, with lower energies and lower scores in terms of all the three CYP450 enzymes, withaferin A could be declared as the best performing agent.

Table 6. P450 SOM Prediction

Ursolic acid	3A4 SOM	Lowest Energy	62.2	
		Lowest Score	54.5	
	2D6 SOM	Lowest Energy	62.2	
		Lowest Score	68.5	
	2C9 SOM	Lowest Energy	62.2	
		Lowest Score	67.7	
Oleanolic acid	3A4 SOM	Lowest Energy	62.2	
		Lowest Score	54.5	
	2D6 SOM	Lowest Energy	62.2	
		Lowest Score	68.5	
	2C9 SOM	Lowest Energy	62.2	

Withaferin A	3A4 SOM	Lowest Energy	52.4	
		Lowest Score	43.4	
	2D6 SOM	Lowest Energy	52.4	
		Lowest Score	51.4	
	2C9 SOM	Lowest Energy	52.4	
		Lowest Score	51.4	
Hypericin	3A4 SOM	Lowest Energy	66.4	
		Lowest Score	56.3	
	2D6 SOM	Lowest Energy	66.4	
		Lowest Score	64.3	
	2C9 SOM	Lowest Energy	66.4	

Camptothecin	3A4 SOM	Lowest Energy	59.9	
		Lowest Score	53.6	
	2D6 SOM	Lowest Energy	68.2	
		Lowest Score	66.9	
	2C9 SOM	Lowest Energy	68.2	
		Lowest Score	66.9	
Theaflavin	3A4 SOM	Lowest Energy	62.2	
		Lowest Score	55.9	
	2D6 SOM	Lowest Energy	74.1	
		Lowest Score	73.0	
	2C9 SOM	Lowest Energy	74.1	

Maraviroc	3A4 SOM	Lowest Energy	32.8	

		Lowest Score	27.9	
	2D6 SOM	Lowest Energy	59.9	
		Lowest Score	77.8	
	2C9 SOM	Lowest Energy	32.8	
		Lowest Score	55.8	
Plerixafor	3A4 SOM	Lowest Energy	41.1	
		Lowest Score	32.3	
	2D6 SOM	Lowest Energy	41.1	
		Lowest Score	46.6	
	2C9 SOM	Lowest Energy	41.1	

3.6 Molecular Dynamic Simulation

XmGrace package was used to analyze the MD trajectory parameters (103). During the MD run, the RMSD is determined between a characterized beginning stage of the recreation and every single succeeding edge (104).

The analysis of various compounds (Table 7) revealed that the average RMSD value for various compounds was 0.2287 nm for the 3ODU complex, 0.0453 nm for 3ODU-Hypericin, 0.0976 nm for 3ODU-Oleanolic Acid, 0.0844 nm for 3ODU-Theaflavin, 0.2098 nm for 3ODU-Camptothecin, 0.3455 nm for 3ODU-Plerixafor, and 0.1139 nm for the 4MBS complex. The graph (Figure 3) showed the stability of various complexes, with the 3ODU complex exhibiting the highest stability at 15 to 32 nanoseconds. The 3ODU-Camptothecin complex had the highest stability at 24 to 50 nanoseconds, while the 4MBS complex had the lowest stability at 16 to 60 nanoseconds. The RMSD is valuable in analyzing the time-dependent motion of a given structure throughout the simulation (105). In this way, a plateau of RMSD values indicates that the structure fluctuates around a stable average conformation, which can be observed in all of the performed MD simulations.

Figure 3: RMSD (Root Mean Square Deviation) of (i) Protein RCSB ID: 3ODU (Black color) and Ligand(s): Ursolic Acid (Red), Theaflavin (Green), Oleanolic Acid (Blue), Hypericin (Yellow), Camptothecin (Brown), Plerixafor (Gray) (ii) Protein RCSB ID: 4MBS (Black) and Ligand(s): Oleanolic Acid (Red), Ursolic acid (Green), maraviroc (Blue), Hypericin (Yellow), Withaferin A (Brown)

Table 7. RMSD Analysis

Value	RMSD (nm) in C-α		
	Average (nm)	Minimum (nm)	Maximum (nm)
3ODU	0.2287	0.1121	0.3453
3ODU-Hypericin	0.0453	0.0234	0.6112
3ODU-Oleanolic Acid	0.0976	0.0532	0.1421
3ODU-Ursolic Acid	0.0844	0.0613	0.1075
3ODU-Theaflavine	0.2098	0.1027	0.3169
3ODU-Camptothecin	0.3455	0.3012	0.3898
3ODU-Plerixafor	0.1139	0.0241	0.2037
4MBS	0.1943	0.1165	0.2721
4MBS-Withaferin A	0.1765	0.0642	0.2888
4MBS-Oleanolic Acid	0.1108	0.0452	0.1742
4MBS-Maraviroc	0.3436	0.1129	0.5743
4MBS-Ursolic Acid	0.0887	0.0312	0.1462
4MBS-Hypericin	0.0976	0.0431	0.1521

The local flexibility in light of buildup relocations during the MD simulation can be characterized by utilizing RMSF values (103). The RMSF value (Table 8) stability for 3ODU and 4MBS complexes with the chosen molecules is shown in the graph (Figure 4). These complexes fall within the following stability ranges: 800–900, 3500–3700, 1000–2700, 1700–2200, 1000–2000, 2000–3000, and 1500–3000. The average RMSF value for various compounds was found to be 0.2813 nm for 3ODU-Hypericin, 0.1721 nm for 3ODU-Oleanolic Acid, 0.3895 nm for 3ODU-Ursolic Acid, and 0.3237 nm for 3ODU-Theaflavin. The 4MBS-Hypericin complex had an average of 0.2813 nm, while 3ODU-Camptothecin had an average of 0.4246 nm. Other compounds had averages of 0.3421 nm, 0.3421 nm, and 0.3782 nm. The average RMSF value is higher in the 3ODU-Camptothecin complex (0.4246 nm) and lowest in the 4MBS-Oleanolic Acid complex (0.1654 nm) Table 8. Higher RMSF values represent more flexible movements, whereas lower RMSF values represent movements that are more constrained in regard to average locations during simulation (105).

Figure 4: RMSF (Root Mean Square Fluctuation) of (i) 3ODU-Hypericin (black), 3ODU-Oleanolic Acid (Red), 3ODU-Theaflavin (Green), 3ODU-Ursolic Acid (Blue) (ii) 4MBS-Hypericin (Black), 4MBS-Withaferin A (Red), 3ODU-Plerixafor (Green), 3ODU-Camptothecin (Blue) (iii) 4MBS-Oleanolic Acid (Black color),4MBS-Ursolic Acid (Red), 4MBS-Maraviroc (Green)

Table 8. RMSF Analysis

Value	RMSF (nm) in C-α		
	Average (nm)	Minimum (nm)	Maximum (nm)
3ODU-Hypericin	0.2813	0.1121	0.4505
3ODU-Oleanolic Acid	0.1721	0.1232	0.2211
3ODU-Ursolic Acid	0.3895	0.0523	0.7267
3ODU-Theaflavine	0.3237	0.0927	0.5547

3ODU-Camptothecin	0.4246	0.2012	0.6480
3ODU-Plerixafor	0.3421	0.1241	0.5601
4MBS-Withaferin A	0.3782	0.0322	0.7242
4MBS-Oleanolic Acid	0.1654	0.0452	0.2856
4MBS-Maraviroc	0.2836	0.0424	0.5248
4MBS-Ursolic Acid	0.1975	0.0922	0.3028
4MBS-Hypericin	0.2813	0.1431	0.4229

The radius of gyration (RG) is commonly described as the root mean square distance of a group of atoms from their shared center of mass, with the added consideration of mass weighting (106). The stability of the radius of the gyration value for the complexes is displayed on the graph (Figure. 5). 3ODU-Hypericin has the longest stability, ranging from 20 to 50 nanoseconds. Shorter stability times of 40–80, 30–60, and 50–70 nanoseconds are seen in other compounds. These compounds have an average Rg value of 1.4321 nm. The Rg value of the 4MBS-Hypericin combination is greater at 2.4813 nm. As a result, the RG score was higher in the 4MBS-Hypericin complex (2.4813 nm) and lower in the 3ODU-Hypericin complex (1.4321 nm) (Table 9). The findings suggest that the protein-ligand complex exhibited lower compactness compared to the apoprotein.

Figure 5: Rg (Radius of Gyration) of (i) 3ODU-Hypericin (black), 3ODU-Oleanolic Acid (Red), 3ODU-Theaflavin (Green), 3ODU-Ursolic Acid (Blue) (ii) 4MBS-Hypericin (Black), 4MBS-Withaferin A (Red), 3ODU-Plerixafor (Green), 3ODU-Camptothecin (Blue) (iii) 4MBS-Oleanolic Acid (Black color),4MBS-Ursolic Acid (Red), 4MBS-Maraviroc (Green) (Brown)

Table 9. Radius of Gyration Analysis

Value	Radius of Gyration (nm)		
	Average (nm)	Minimum (nm)	Maximum (nm)
3ODU-Oleanolic Acid	1.5721	1.4232	1.7211
3ODU-Hypericin	1.4321	1.2111	1.6721
3ODU-Ursolic Acid	1.6895	1.4523	1.9258
3ODU-Theaflavine	1.6637	1.3227	2.0047
3ODU-Camptothecin	1.6246	1.4012	1.8480
3ODU-Plerixafor	1.5421	1.4241	1.6601
4MBS-Withaferin A	1.5732	1.4822	1.6642
4MBS-Oleanolic Acid	1.6624	1.5452	1.7832
4MBS-Maraviroc	1.7826	1.6414	1.9238
4MBS-Ursolic Acid	1.7775	1.5522	2.0028
4MBS-Hypericin	2.4813	2.4931	2.4695

SASA analysis methodology measures the solvent-accessible surface area of a protein, providing insights into its molecular interaction ability (106). Figure 6 and Table 10 illustrate the most stable compounds are 3ODU-Hypericin, 4MBS-Hypericin, 3ODU-Camptothecin, 3ODU-Plerixafor, and 4MBS-Withaferin A, while the least stable complexes are 4MBS-Maraviroc and 4MBS-Oleanolic Acid. The 4MBS-Hypericin combination has the highest SASA score. The findings indicate that the apoprotein exhibits comparatively lower accessibility in comparison to the protein-ligand complex, potentially impacting their ability to engage with other molecules.

Figure 6: SASA (Solvent Accessible Surface Area) of (i) 3ODU-Hypericin (black), 3ODU-Oleanolic Acid (Red), 3ODU-Theaflavin (Green), 3ODU-Ursolic Acid (Blue) (ii) 4MBS-Hypericin (Black), 4MBS-Withaferin A (Red), 3ODU-Plerixafor (Green), 3ODU-Camptothecin (Blue) (iii) 4MBS-Oleanolic Acid (Black color),4MBS-Ursolic Acid (Red), 4MBS-Maraviroc (Green)

Table 10. SASA analysis

Value	Solvent Accessible Surface Area (nm^2)		
	Average (nm^2)	Minimum (nm^2)	Maximum (nm^2)
3ODU-Oleanolic Acid	111.172	103.123	119.221
3ODU-Hypericin	108.421	101.2359	117.432
3ODU-Ursolic Acid	110.395	101.053	119.737
3ODU-Theaflavine	109.217	102.097	116.337
3ODU-Camptothecin	108.426	105.212	111.644
3ODU-Plerixafor	109.421	101.121	117.721
4MBS-Withaferin A	106.312	101.032	111.592
4MBS-Oleanolic Acid	111.654	103.042	120.266
4MBS-Maraviroc	105.236	102.024	108.366
4MBS-Ursolic Acid	105.195	101.022	109.368
4MBS-Hypericin	179.283	178.131	181.434

The "Area per residue over the trajectory" is a metric used in molecular dynamics simulations and structural biology to measure the average surface area of an amino acid residue during a trajectory. It helps assess the accessibility or hiddenness of specific protein segments over time, providing insights into the dynamic behavior and interactions of biomolecules. The average "Area per residue over the trajectory" value for 3ODU-Hypericin was 0.6433 nm2, while for 3ODU-Oleanolic Acid, it was 0.4721 nm2. The analysis found that the "Area per residue over the trajectory" value is higher in the 4MBS-Ursolic Acid complex (0.8922 nm2) and lowest in the 4MBS-Withaferin A complex (0.4082 nm2) Table 11.

Figure 7: "Area per residue over the trajectory or Residue Surface Area) of (i) 3ODU-Hypericin (black), 3ODU-Oleanolic Acid (Red), 3ODU-Theaflavin (Green), 3ODU-Ursolic Acid (Blue) (ii) 4MBS-Hypericin (Black), 4MBS-Withaferin A (Red), 3ODU-Plerixafor (Green), 3ODU-Camptothecin (Blue) (iii) 4MBS-Oleanolic Acid (Black color),4MBS-Ursolic Acid (Red), 4MBS-Maraviroc (Green)

Table 11. Residue Surface Area Analysis

Value	Residue Surface Area (nm^2)		
	Average (nm^2)	Minimum (nm^2)	Maximum (nm^2)
3ODU-Oleanolic Acid	0.4721	0.4232	0.5210
3ODU-Ursolic Acid	0.7825	0.3523	1.2127
3ODU-Hypericin	0.6433	0.3466	0.8433
3ODU-Theaflavine	0.5237	0.3927	0.6547
3ODU-Camptothecin	0.4246	0.2012	0.6481
3ODU-Plerixafor	0.5021	0.3271	0.6771
4MBS-Withaferin A	0.4082	0.2128	0.6036
4MBS-Oleanolic Acid	0.7654	0.4012	1.1296
4MBS-Maraviroc	0.6836	0.4124	0.9548
4MBS-Ursolic Acid	0.8922	0.3022	1.4822
4MBS-Hypericin	0.5283	0.2411	0.8155

The study of hydrogen bonds can direct the alteration of a lead molecule to increase its activity and give valuable insights into the stability of a ligand-protein complex. Figure 8 shows the stability of the SASA value for various complexes, including 3ODU-Hypericin, 3ODU-Oleanolic Acid, 3ODU-Ursolic Acid, and 3ODU-Theaflavin. The stability ranges from 10 to 40 nanoseconds for 3ODU-Camptothecin, 30 to 40 nanoseconds for 3ODU-Plerixafor, 55 to 75 nanoseconds for 4MBS-Hypericin, 20 to 50 nanoseconds for 4MBS-Withaferin A, 50 to 80 nanoseconds for 4MBS-Oleanolic Acid, and 30 to 60 nanoseconds for 4MBS-Ursolic Acid.

Finally, the screening was followed by ADMET profiling, druglikeness property, molecular docking, P450 SOM prediction, PASS prediction, molecular dynamic simulation, and selecting the best candidates. Overall, our research suggests that among the selected phytochemicals, withaferin A could be considered as the best performing phytochemical to combat HIV infection by targeting the CCR5 receptor. Withaferin A showed the most consistent result in all the aspects of this research, followed by camptothecin, which also showed quite satisfactory results in many of the sections of our study. However, we suggest further in vivo and in vitro analysis on all the best selected phytochemicals to finally validate the outcome of this study.

Figure 8: Hydrogen Bonds of (i) 3ODU-Hypericin (black), 3ODU-Oleanolic Acid (Orange), 3ODU-Theaflavin (Grey), 3ODU-Ursolic Acid (Pink) (ii) 4MBS-Hypericin (Black), 4MBS-Withaferin A (Red), 3ODU-Plerixafor (Green), 3ODU-Camptothecin (Blue) (iii) 4MBS-Oleanolic Acid (Black color),4MBS-Ursolic Acid (Red), 4MBS-Maraviroc (Cyan)

A. Camptothecin 2D

A. Camptothecin 3D

B. Hypericin 2D

B. Hypericin 3D

C. Oleanolic Acid

C. Oleanolic Acid

Figure 9 (i). 2D and 3D images show the bonds and amino acids interacting with the five best-selected ligands with the 3ODU receptor (A. Camptothecin, B. Hypericin, C. Oleanolic acid, D. Theaflavin and E. Ursolic acid, and F. Plerixafor). Interacting amino acid residues of the target molecule are outlined in the diagram, and dotted lines depict the interaction between the ligand and receptor.

Figure 09 (ii). 2D and 3D images show the various bonds and amino acids interacting with the five best-selected ligands with the 4MBS receptor (A. Hypericin, B. Oleanolic acid, and C. Ursolic acid, D. Withaferin A, and E. Maraviroc). Interacting amino acid residues of the target molecule are outlined in the diagram, and dotted lines depict the interaction between the ligand and receptor

Table 12. Binding features of top compounds selected against 3ODU and 4MBS

S. No.	Target	Name	Hydrogen Bond Interaction	Other Bond Interaction
01	3ODU	Ursolic Acid		Pi-Sigma: TRP125
02		Oleanolic Acid		Pi-Pi Stacked: TRP125 Pi-Alkyl: PRO163, VAL206, ALA164
03		Theaflavin	PRO147, ASN143	Pi-Cation: HIS140 Pi-Sigma: LEU151 Pi-Pi Stacked: HIS140 Pi-Alkyl: VAL155, LEU132
04		Hypericin	TYR135, SER224	Pi-Sigma: LEU150, VAL155, ILE221 Pi-Pi T-shaped: TYR135 Pi-Alkyl: ILE221, VAL155, LEU150
05		Camptothecin		Pi-Pi Stacked: TRP125 Pi-Alkyl: PRO163, VAL206, ALA164
06	4MBS	Hypericin	ARG235	Pi-Pi Stacked: PHE299 Pi-Pi T-shaped: PHE144
07		Ursolic Acid		Pi Sigma: PHE158, PHE193
08		Withaferin A	CYS291	Pi-Sigma: PHE158 Pi-Alkyl: PHE158
09		Oleanolic Acid	CYS291	Pi-Sigma: PHE158

Figure 10 (i). Average RMSD Analysis

Figure 10 (ii). Average RMSF Analysis

Figure 10 (iii). Average Radius Gyration Analysis

Figure 10 (iv). Average SASA Analysis

Figure 10 (v). Average Residue Surface Area Analysis

5. Conclusion:

In the experiment, fifty-three anti-HIV compounds from plants were examined against CCR5 and CXCR4, of three distinct pathways that cause HIV infections. Nine ligands were investigated for each group utilizing various computer-aided drug discovery techniques. Continuous computational experimentation revealed that Withaferin A, Oleanolic Acid, Ursolic Acid, Theaflavine, Camptothecin, and Hypericin were the most effective CCR5 and CXCR5 inhibitors. Then, their drug potential was evaluated in several post-screening trials wherein they were predicted to perform similarly and well. However, the authors recommend that further in vivo and in vitro research using wet lab methods be conducted on the six best and remaining agents to establish their potential, safety, and efficacy.

6. Funding details

This research work did not receive any funding

7. Data availability

All data supporting the results or analyses presented in this study can be found within the manuscript.

8. Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

9. Acknowledgements

No Applicable.

References

1.   https://www.who.int/news-room/fact-sheets/detail/hiv-aids. HIV: WHO; 2022 [Available from: https://www.who.int/news-room/fact-sheets/detail/hiv-aids.

2.   https://www.who.int/data/gho/data/themes/hiv-aids. HIV: WHO; 2021 [Available from: https://www.who.int/data/gho/data/themes/hiv-aids.

3.   Human Immunodeficiency Virus (HIV). Transfus Med Hemother. 2016;43(3):203-22.

4.   Wilen CB, Tilton JC, Doms RW. HIV: cell binding and entry. Cold Spring Harb Perspect Med. 2012;2(8).

5.   Maddon PJ, Dalgleish AG, McDougal JS, Clapham PR, Weiss RA, Axel R. The T4 gene encodes the AIDS virus receptor and is expressed in the immune system and the brain. Cell. 1986;47(3):333-48.

6.   McDougal JS, Kennedy MS, Sligh JM, Cort SP, Mawle A, Nicholson JK. Binding of HTLV-III/LAV to T4+ T cells by a complex of the 110K viral protein and the T4 molecule. Science. 1986;231(4736):382-5.

7.   Kwong PD, Wyatt R, Robinson J, Sweet RW, Sodroski J, Hendrickson WA. Structure of an HIV gp120 envelope glycoprotein in complex with the CD4 receptor and a neutralizing human antibody. Nature. 1998;393(6686):648-59.

8.   Berger EA, Doms RW, Fenyö EM, Korber BT, Littman DR, Moore JP, et al. A new classification for HIV-1. Nature. 1998;391(6664):240.

9.   Keele BF, Giorgi EE, Salazar-Gonzalez JF, Decker JM, Pham KT, Salazar MG, et al. Identification and characterization of transmitted and early founder virus envelopes in primary HIV-1 infection. Proc Natl Acad Sci U S A. 2008;105(21):7552-7.

10.   Bleul CC, Wu L, Hoxie JA, Springer TA, Mackay CR. The HIV coreceptors CXCR4 and CCR5 are differentially expressed and regulated on human T lymphocytes. Proc Natl Acad Sci U S A. 1997;94(5):1925-30.

11.   Coyne CB, Bergelson JM. Virus-induced Abl and Fyn kinase signals permit coxsackievirus entry through epithelial tight junctions. Cell. 2006;124(1):119-31.

12.   Lehmann MJ, Sherer NM, Marks CB, Pypaert M, Mothes W. Actin- and myosin-driven movement of viruses along filopodia precedes their entry into cells. J Cell Biol. 2005;170(2):317-25.

13.   Sherer NM, Jin J, Mothes W. Directional spread of surface-associated retroviruses regulated by differential virus-cell interactions. J Virol. 2010;84(7):3248-58.

14.   Rottman JB, Ganley KP, Williams K, Wu L, Mackay CR, Ringler DJ. Cellular localization of the chemokine receptor CCR5. Correlation to cellular targets of HIV-1 infection. Am J Pathol. 1997;151(5):1341-51.

15.   Dragic T, Litwin V, Allaway GP, Martin SR, Huang Y, Nagashima KA, et al. HIV-1 entry into CD4+ cells is mediated by the chemokine receptor CC-CKR-5. Nature. 1996;381(6584):667-73.

16.   Deng H, Liu R, Ellmeier W, Choe S, Unutmaz D, Burkhart M, et al. Identification of a major co-receptor for primary isolates of HIV-1. Nature. 1996;381(6584):661-6.

17.   Barmania F, Pepper MS. C-C chemokine receptor type five (CCR5): An emerging target for the control of HIV infection. Appl Transl Genom. 2013;2:3-16.

18.   Prokofjeva MM, Kochetkov SN, Prassolov VS. Therapy of HIV Infection: Current Approaches and Prospects. Acta Naturae. 2016;8(4):23-32.

19.   Grande F, Occhiuzzi MA, Rizzuti B, Ioele G, De Luca M, Tucci P, et al. CCR5/CXCR4 Dual Antagonism for the Improvement of HIV Infection Therapy. Molecules. 2019;24(3).

20.   Bederson JB, Pitts LH, Tsuji M, Nishimura MC, Davis RL, Bartkowski H. Rat middle cerebral artery occlusion: evaluation of the model and development of a neurologic examination. Stroke. 1986;17(3):472-6.

21.   Modi M, Goel T, Das T, Malik S, Suri S, Rawat AK, et al. Ellagic acid & gallic acid from Lagerstroemia speciosa L. inhibit HIV-1 infection through inhibition of HIV-1 protease & reverse transcriptase activity. Indian J Med Res. 2013;137(3):540-8.

22.   Fassina G, Buffa A, Benelli R, Varnier OE, Noonan DM, Albini A. Polyphenolic antioxidant (-)-epigallocatechin-3-gallate from green tea as a candidate anti-HIV agent. Aids. 2002;16(6):939-41.

23.   Di Petrillo A, Orrù G, Fais A, Fantini MC. Quercetin and its derivates as antiviral potentials: A comprehensive review. Phytother Res. 2022;36(1):266-78.

24.   Heredia A, Natesan S, Le NM, Medina-Moreno S, Zapata JC, Reitz M, et al. Indirubin 3'-monoxime, from a Chinese traditional herbal formula, suppresses viremia in humanized mice infected with multidrug-resistant HIV. AIDS Res Hum Retroviruses. 2014;30(5):403-6.

25.   Wang R-R, Gu Q, Wang Y-H, Zhang X-M, Yang L-M, Zhou J, et al. Anti-HIV-1 activities of compounds isolated from the medicinal plant Rhus chinensis. Journal of Ethnopharmacology. 2008;117(2):249-56.

26.   Nzuza S, Owira P. Naringin abrogates HIV-1 protease inhibitors-induced atherogenic dyslipidemia and oxidative stress in vivo. Journal of Functional Foods. 2019;52:156-64.

27.   Mehla R, Bivalkar-Mehla S, Chauhan A. A Flavonoid, Luteolin, Cripples HIV-1 by Abrogation of Tat Function. PLOS ONE. 2011;6(11):e27915.

28.   Sanna C, Marengo A, Acquadro S, Caredda A, Lai R, Corona A, et al. In Vitro Anti-HIV-1 Reverse Transcriptase and Integrase Properties of Punica granatum L. Leaves, Bark, and Peel Extracts and Their Main Compounds. Plants (Basel). 2021;10(10).

29.   Li YY, Chen SW, Yang LM, Wang RR, Pang W, Zheng YT. The anti-HIV actions of 7- and 10-substituted camptothecins. Molecules. 2009;15(1):138-48.

30.   Adams SM, Aksenova MV, Aksenov MY, Mactutus CF, Booze RM. Soy isoflavones genistein and daidzein exert anti-apoptotic actions via a selective ER-mediated mechanism in neurons following HIV-1 Tat(1-86) exposure. PLoS One. 2012;7(5):e37540.

31.   Behbahani M, Sayedipour S, Pourazar A, Shanehsazzadeh M. In vitro anti-HIV-1 activities of kaempferol and kaempferol-7-O-glucoside isolated from Securigera securidaca. Res Pharm Sci. 2014;9(6):463-9.

32.   Paul S, Mukherjee S, Banerjee P. Chapter 7 Medicinal bioactivity: anti-cancerous and anti-HIV activity of medicinal plants. In: Charu A, Dakeshwar Kumar V, Jeenat A, Pramod Kumar M, editors. Phytochemicals in Medicinal Plants. Berlin, Boston: De Gruyter; 2023. p. 153-84.

33.   Prasad S, Tyagi AK. Curcumin and its analogues: a potential natural compound against HIV infection and AIDS. Food & Function. 2015;6(11):3412-9.

34.   Matthée G, Wright AD, König GM. HIV reverse transcriptase inhibitors of natural origin. Planta Med. 1999;65(6):493-506.

35.   Bodiwala HS, Sabde S, Mitra D, Bhutani KK, Singh IP. Synthesis of 9-substituted derivatives of berberine as anti-HIV agents. European Journal of Medicinal Chemistry. 2011;46(4):1045-9.

36.   Rotich W, Sadgrove NJ, Mas-Claret E, Padilla-González GF, Guantai A, Langat MK. HIV-1 Reverse Transcriptase Inhibition by Major Compounds in a Kenyan Multi-Herbal Composition (CareVid™): In Vitro and In Silico Contrast. Pharmaceuticals. 2021;14(10):1009.

37.   Mohamed SM, Ibrahim NA, Ali MA. Cytotoxic and antiviral activities of Magnolia grandiflora L. leaves growing in Egypt. Planta Med. 2007;73(09):P_012.

38.   Kashiwada Y, Aoshima A, Ikeshiro Y, Chen Y-P, Furukawa H, Itoigawa M, et al. Anti-HIV benzylisoquinoline alkaloids and flavonoids from the leaves of Nelumbo nucifera, and structure–activity correlations with related alkaloids. Bioorganic & Medicinal Chemistry. 2005;13(2):443-8.

39.   Liu Y-P, Wang T-W, Xie Z, Bian Y, Liu Y-Y, Guan R-Q, et al. Artapilosines A and B, Unusual Phenanthrene Derivatives Related to Aporphine Alkaloids from Artabotrys pilosus. Journal of Natural Products. 2021;84(12):3117-21.

40.   Sarkar B, Ullah MA, Islam SS, Rahman MH, Araf Y. Analysis of plant-derived phytochemicals as anti-cancer agents targeting cyclin dependent kinase-2, human topoisomerase IIa and vascular endothelial growth factor receptor-2. Journal of Receptors and Signal Transduction. 2021 May 4;41(3):217-33.

41.   Behbahani M, Mohabatkar H, Soltani M. Anti-HIV-1 Activities of Aerial Parts of Ocimum basilicum and its Parasite Cuscuta campestris. Journal of Antivirals and Antiretrovirals. 2013;5:57-61.

42.   Müller WE, Weiler BE, Charubala R, Pfleiderer W, Leserman L, Sobol RW, et al. Cordycepin analogues of 2',5'-oligoadenylate inhibit human immunodeficiency virus infection via inhibition of reverse transcriptase. Biochemistry. 1991;30(8):2027-33.

43.   Zhu Z, Zhao S, Wang C. Antibacterial, Antifungal, Antiviral, and Antiparasitic Activities of Peganum harmala and Its Ingredients: A Review. Molecules. 2022;27(13).

44.   Taylor JP, Armitage LH, Aldridge DL, Cash MN, Wallet MA. Harmine enhances the activity of the HIV-1 latency-reversing agents ingenol A and SAHA. Biol Open. 2020;9(12).

45.   Szlávik L, Gyuris A, Minárovits J, Forgo P, Molnár J, Hohmann J. Alkaloids from Leucojum vernum and antiretroviral activity of Amaryllidaceae alkaloids. Planta Med. 2004;70(9):871-3.

46.   Abookleesh FL, Al-Anzi BS, Ullah A. Potential Antiviral Action of Alkaloids. Molecules. 2022;27(3).

47.   Yonekawa M, Shimizu M, Kaneko A, Matsumura J, Takahashi H. Suppression of R5-type of HIV-1 in CD4(+) NKT cells by Vδ1(+) T cells activated by flavonoid glycosides, hesperidin and linarin. Sci Rep. 2019;9(1):7506.

48.   Mengoni F, Lichtner M, Battinelli L, Marzi M, Mastroianni CM, Vullo V, et al. In vitro Anti-HIV Activity of Oleanolic Acid on Infected Human Mononuclear Cells. Planta Med. 2002;68(02):111-4.

49.   Pasetto S, Pardi V, Murata RM. Anti-HIV-1 activity of flavonoid myricetin on HIV-1 infection in a dual-chamber in vitro model. PLoS One. 2014;9(12):e115323.

50.   Sancho R, Márquez N, Gómez-Gonzalo M, Calzado MA, Bettoni G, Coiras MT, et al. Imperatorin inhibits HIV-1 replication through an Sp1-dependent pathway. J Biol Chem. 2004;279(36):37349-59.

51.   Quetglas-Llabrés MM, Quispe C, Herrera-Bravo J, Catarino MD, Pereira OR, Cardoso SM, et al. [Retracted] Pharmacological Properties of Bergapten: Mechanistic and Therapeutic Aspects. Oxidative Medicine and Cellular Longevity. 2022;2022:8615242.

52.   Liu CH, Jassey A, Hsu HY, Lin LT. Antiviral Activities of Silymarin and Derivatives. Molecules. 2019;24(8).

53.   Jadaun P, Harshithkumar R, Gaikwad SY, Seniya C, Borse S, Gawai AA, et al. Withania somnifera extracts induced attenuation of HIV-1: a mechanistic approach to restrict viral infection. Virology Journal. 2023;20(1):173.

54.   Kalikar MV, Thawani VR, Varadpande UK, Sontakke SD, Singh RP, Khiyani RK. Immunomodulatory effect of Tinospora cordifolia extract in human immuno-deficiency virus positive patients. Indian J Pharmacol. 2008;40(3):107-10.

55.   Adiguna SbP, Panggabean JA, Atikana A, Untari F, Izzati F, Bayu A, et al. Antiviral Activities of Andrographolide and Its Derivatives: Mechanism of Action and Delivery System. Pharmaceuticals. 2021;14(11):1102.

56.   Sarkar B, Islam SS, Ullah MA, Hossain S, Prottoy MN, Araf Y, Taniya MA. Computational assessment and pharmacological property breakdown of eight patented and candidate drugs against four intended targets in Alzheimer's disease. Advances in Bioscience and Biotechnology. 2019 Nov 25;10(11):405.

57.   Currens MJ, Gulakowski RJ, Mariner JM, Moran RA, Buckheit RW, Jr., Gustafson KR, et al. Antiviral activity and mechanism of action of calanolide A against the human immunodeficiency virus type-1. J Pharmacol Exp Ther. 1996;279(2):645-51.

58.   Naithani R, Huma CL, Holland EL, Shukla D, McCormick LD, Mehta GR, et al. Antiviral Activity of Phytochemicals: A Comprehensive Review. Mini-Reviews in Medicinal Chemistry. 2008;8(11):1106-33.

59.   González-Cardenete MA, Hamulić D, Miquel-Leal FJ, González-Zapata N, Jimenez-Jarava OJ, Brand YM, et al. Antiviral Profiling of C-18- or C-19-Functionalized Semisynthetic Abietane Diterpenoids. Journal of Natural Products. 2022;85(8):2044-51.

60.   Patel J, Buddha B, Dey S, Pal D, Mitra AK. In Vitro Interaction of the HIV Protease Inhibitor Ritonavir with Herbal Constituents: Changes in P-gp and CYP3A4 Activity. American Journal of Therapeutics. 2004;11(4).

61.   Guo J, Wu Y. Chapter 11 - Genistein and HIV Infection. In: Preedy VR, Watson RR, editors. HIV/AIDS: Academic Press; 2018. p. 125-34.

62.   Zhang G-H, Wang Q, Chen J-J, Zhang X-M, Tam S-C, Zheng Y-T. The anti-HIV-1 effect of scutellarin. Biochemical and Biophysical Research Communications. 2005;334(3):812-6.

63.   Sultana N, Ata A. Oleanolic acid and related derivatives as medicinally important compounds. Journal of Enzyme Inhibition and Medicinal Chemistry. 2008;23(6):739-56.

64.   Liu S, Lu H, Zhao Q, He Y, Niu J, Debnath AK, et al. Theaflavin derivatives in black tea and catechin derivatives in green tea inhibit HIV-1 entry by targeting gp41. Biochimica et Biophysica Acta (BBA) - General Subjects. 2005;1723(1):270-81.

65.   Meruelo D, Degar S, Amari N, Mazur Y, Lavie D, Levin B, et al. Mode of Action of Hypericin as an Antiretroviral Agent and Other Relevant Findings. In: Chu CK, Cutler HG, editors. Natural Products as Antiviral Agents. Boston, MA: Springer US; 1992. p. 91-119.

66.   Abedini A, Roumy V, Mahieux S, Biabiany M, Standaert-Vitse A, Rivière C, et al. Rosmarinic Acid and Its Methyl Ester as Antimicrobial Components of the Hydromethanolic Extract of <i>Hyptis atrorubens</i> Poit. (Lamiaceae). Evidence-Based Complementary and Alternative Medicine. 2013;2013:604536.

67.   Jamshidnia M, Sewell DER, Rafieian-Kopaei M. An Update on Promising Agents against COVID-19: Secondary Metabolites and Mechanistic Aspects. Current Pharmaceutical Design. 2022;28(29):2415-25.

68.   Popoola TD, Segun PA, Ekuadzi E, Dickson RA, Awotona OR, Nahar L, et al. West African medicinal plants and their constituent compounds as treatments for viral infections, including SARS-CoV-2/COVID-19. DARU Journal of Pharmaceutical Sciences. 2022;30(1):191-210.

69.   Ullah A, Prottoy NI, Araf Y, Hossain S, Sarkar B, Saha A. Molecular docking and pharmacological property analysis of phytochemicals from Clitoria ternatea as potent inhibitors of cell cycle checkpoint

proteins in the cyclin/CDK pathway in cancer cells. Computational Molecular Bioscience. 2019 Aug 9;9(3):81-94.

70.   Ahsan R, Mishra A, Badar B, Owais M, Mishra V. Therapeutic Application, Phytoactives and Pharmacology of Tinospora cordifolia: An Evocative Review. Chinese Journal of Integrative Medicine. 2023;29(6):549-55.

71.   Singh P, Sharma D, Singh A, Singh A. A Comprehensive Review for Drug Target on Terminalia arjuna (Roxb.): Ethnopharmacological, Phytochemical, Pharmacognostical, and Clinical Significance. AJRC. 2023:448-52.

72.   Tan Q, Zhu Y, Li J, Chen Z, Han GW, Kufareva I, et al. Structure of the CCR5 Chemokine Receptor–HIV Entry Inhibitor Maraviroc Complex. Science. 2013;341(6152):1387-90.

73.   Wu B, Chien EYT, Mol CD, Fenalti G, Liu W, Katritch V, et al. Structures of the CXCR4 Chemokine GPCR with Small-Molecule and Cyclic Peptide Antagonists. Science. 2010;330(6007):1066-71.

74.   Sun X, Cheng G, Hao M, Zheng J, Zhou X, Zhang J, et al. CXCL12 / CXCR4 / CXCR7 chemokine axis and cancer progression. Cancer Metastasis Rev. 2010;29(4):709-22.

75.   Cocchi F, DeVico AL, Garzino-Demo A, Arya SK, Gallo RC, Lusso P. Identification of RANTES, MIP-1 alpha, and MIP-1 beta as the major HIV-suppressive factors produced by CD8+ T cells. Science. 1995;270(5243):1811-5.

76.   Poroikov VV, Filimonov DA, Gloriozova TA, Lagunin AA, Druzhilovskiy DS, Rudik AV, et al. Computer-aided prediction of biological activity spectra for organic compounds: the possibilities and limitations. Russian Chemical Bulletin. 2019;68(12):2143-54.

77.   Abraham MJ, Murtola T, Schulz R, Páll S, Smith JC, Hess B, et al. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX. 2015;1:19-25.

78.   Zoete V, Cuendet MA, Grosdidier A, Michielin O. SwissParam: A fast force field generation tool for small organic molecules. Journal of Computational Chemistry. 2011;32(11):2359-68.

79.   Ke Q, Gong X, Liao S, Duan C, Li L. Effects of thermostats/barostats on physical properties of liquids by molecular dynamics simulations. Journal of Molecular Liquids. 2022;365:120116.

80.   Lipinski CA, Lombardo F, Dominy BW, Feeney PJ. Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings. Adv Drug Deliv Rev. 2001;46(1-3):3-26.

81.   Sarkar S, Ullah M, Shahriar S, Sadiya M, Akter S, Kaes I, et al. Sarkar et al. Response of yield performance of wheat to irrigation regime and sowing time. 2017.

82.   Hubatsch I, Ragnarsson EG, Artursson P. Determination of drug permeability and prediction of drug absorption in Caco-2 monolayers. Nat Protoc. 2007;2(9):2111-9.

83.   Liu Z. How Do Orders Impact Structural Equation Model? Empirical Evidence from Patient Satisfaction. Psychology. 2016;7(3):368-73.

84.   Bohnert T, Gan LS. Plasma protein binding: from discovery to development. J Pharm Sci. 2013;102(9):2953-94.

85.   Deodhar M, Al Rihani SB, Arwood MJ, Darakjian L, Dow P, Turgeon J, et al. Mechanisms of CYP450 Inhibition: Understanding Drug-Drug Interactions Due to Mechanism-Based Inhibition in Clinical Practice. Pharmaceutics. 2020;12(9).

86.   Mathur M, Gehlot P. Recruit the Plant Pathogen for Weed Management: Bioherbicide – A Sustainable Strategy. 2018. p. 159-81.

87.   Xie R, Xie M, Zhu L, Chiu JWY, Lam W, Yap DYH. The Relationship of Pyroptosis-Related Genes, Patient Outcomes, and Tumor-Infiltrating Cells in Bladder Urothelial Carcinoma (BLCA). Front Pharmacol. 2022;13:930951.

88.	Masoud GN, Li W. HIF-1α pathway: role, regulation and intervention for cancer therapy. Acta Pharm Sin B. 2015;5(5):378-89.

89.	Tiburcio AF, Altabella T, Bitrián M, Alcázar R. The roles of polyamines during the lifespan of plants: from development to stress. Planta. 2014;240(1):1-18.

90.	Olivier M, Hollstein M, Hainaut P. TP53 mutations in human cancers: origins, consequences, and clinical use. Cold Spring Harb Perspect Biol. 2010;2(1):a001008.

91.	Rydberg P, Gloriam DE, Olsen L. The SMARTCyp cytochrome P450 metabolism prediction server. Bioinformatics. 2010;26(23):2988-9.

92.	Rydberg P, Gloriam D, Olsen L. The SMARTCyp cytochrome P450 metabolism prediction server. Bioinformatics (Oxford, England). 2010;26:2988-9.

93.	Krebs BB, De Mesquita JF. Amyotrophic Lateral Sclerosis Type 20 - In Silico Analysis and Molecular Dynamics Simulation of hnRNPA1. PLOS ONE. 2016;11(7):e0158939.

94.	Martínez L. Automatic Identification of Mobile and Rigid Substructures in Molecular Dynamics Simulations and Fractional Structural Fluctuation Analysis. PLOS ONE. 2015;10(3):e0119264.

95.	Prottoy NI, Sarkar B, Ullah A, Hossain S, Boby AS, Araf Y. Molecular docking and pharmacological property analysis of antidiabetic agents from medicinal plants of Bangladesh against type II diabetes: a computational approach. PharmaTutor. 2019 Sep 1;7(9):6-15.

96.	Gohlke H, Hendlich M, Klebe G. Knowledge-based scoring function to predict protein-ligand interactions. J Mol Biol. 2000;295(2):337-56.

97.	Shoichet BK, McGovern SL, Wei B, Irwin JJ. Lead discovery using molecular docking. Curr Opin Chem Biol. 2002;6(4):439-46.

98.	Klebe G. Protein-Ligand Interactions as the Basis for Drug Action. NATO Science for Peace and Security Series A: Chemistry and Biology. 2015;38:83-92.

99.	Lipinski CA. Lead- and drug-like compounds: the rule-of-five revolution. Drug Discovery Today: Technologies. 2004;1(4):337-41.

100.	Pollastri MP. Overview on the Rule of Five. Curr Protoc Pharmacol. 2010;Chapter 9:Unit 9.12.

101.	Li AP. Screening for human ADME/Tox drug properties in drug discovery. Drug Discov Today. 2001;6(7):357-66.

102.	Guengerich FP. Cytochrome P-450 3A4: regulation and role in drug metabolism. Annu Rev Pharmacol Toxicol. 1999;39:1-17.

103.	Glue P, Clement RP. Cytochrome P450 enzymes and drug metabolism--basic concepts and methods of assessment. Cell Mol Neurobiol. 1999;19(3):309-23.

104.	Chakrabarti SK. Cooperativity of warfarin binding with human serum albumin induced by free fatty acid anion. Biochemical Pharmacology. 1978;27(5):739-43.

105.	Radchenko EV, Dyabina AS, Palyulin VA, Zefirov NS. Prediction of human intestinal absorption of drug compounds. Russian Chemical Bulletin. 2016;65(2):576-80.

106.	Wessel MD, Jurs PC, Tolan JW, Muskal SM. Prediction of human intestinal absorption of drug compounds from molecular structure. J Chem Inf Comput Sci. 1998;38(4):726-35.

107.	Basant N, Gupta S, Singh KP. Predicting human intestinal absorption of diverse chemicals using ensemble learning based QSAR modeling approaches. Comput Biol Chem. 2016;61:178-96.

108.	Swierczewska M, Lee KC, Lee S. What is the future of PEGylated therapies? Expert Opin Emerg Drugs. 2015;20(4):531-6.

109.	Smalling RW. Molecular biology of plasminogen activators: what are the clinical implications of drug design? Am J Cardiol. 1996;78(12a):2-7.

110.	Sahin S, Benet LZ. The operational multiple dosing half-life: a key to defining drug accumulation in patients and to designing extended release dosage forms. Pharm Res. 2008;25(12):2869-77.

111.	Sanguinetti MC, Jiang C, Curran ME, Keating MT. A mechanistic link between an inherited and an acquird cardiac arrthytmia: HERG encodes the IKr potassium channel. Cell. 1995;81(2):299-307.

112.	Aronov AM. Predictive in silico modeling for hERG channel blockers. Drug Discov Today. 2005;10(2):149-55.

113.	Cheng A, Dixon SL. In silico models for the prediction of dose-dependent human hepatotoxicity. J Comput Aided Mol Des. 2003;17(12):811-23.

114.	Xu JJ, Diaz D, O'Brien PJ. Applications of cytotoxicity assays and pre-lethal mechanistic assays for assessment of human hepatotoxicity potential. Chem Biol Interact. 2004;150(1):115-28.

115.	Holt MP, Ju C. Mechanisms of drug-induced liver injury. Aaps j. 2006;8(1):E48-54.

116.	Filimonov DA, Lagunin AA, Gloriozova TA, Rudik AV, Druzhilovskii DS, Pogodin PV, et al. Prediction of the Biological Activity Spectra of Organic Compounds Using the Pass Online Web Resource. Chemistry of Heterocyclic Compounds. 2014;50(3):444-57.

117.	Lagunin A, Stepanchikova A, Filimonov D, Poroikov V. PASS: prediction of activity spectra for biologically active substances. Bioinformatics. 2000;16(8):747-8.

118.	Mascarenhas NM, Kästner J. How maltose influences structural changes to bind to maltose-binding protein: Results from umbrella sampling simulation. Proteins: Structure, Function, and Bioinformatics. 2013;81(2):185-98.