Apolipoprotein interaction induces shape remodeling and lipid phase separation in giant unilamellar vesicles

Christopher F. Carnahan[1], Wei He[2], Tugba N. Ozturk[2], Yaqing Wang[3], Viviane N. Ngassam[4], Aleksandr Noy[3], Timothy S. Carpenter[2], John C. Voss[5], Matthew A. Coleman[2], Atul N. Parikh[1,4,6]

1. Biophysics Graduate Group, University of California, Davis, CA
2. Biosciences and Biotechnology Division, Lawrence Livermore National Laboratory, Livermore, CA
3. Materials Science Division, Lawrence Livermore National Laboratory, Livermore, CA
4. Department of Biomedical Engineering, University of California, Davis, CA
5. Department of Biochemistry & Molecular Medicine, University of California, Davis, CA
6. Institute for Digital Molecular Analytics & Science, Nanyang Technological University, Singapore

ABSTRACT

Apolipoprotein A-I (ApoA-I) – a 243-residue amphipathic protein containing an N-terminal globular domain and a primarily helical C-terminal lipid binding domain – is a principal protein component of high-density lipoprotein (HDL) or "good" cholesterol, which is an important component of lipid homeostasis in humans. Synthesized in the liver and intestine and excreted in the blood, ApoA-I undergoes complex, cooperative, and dynamic self-assembly with membrane lipids producing unlipidated (or weakly lapidated), nascent discoidal, and mature HDL states. *In vitro* studies establish that the recombination of purified protein and lipids reconstitutes this cooperative self-assembly. However, the kinetic pathways by which these mesoscopic, proteolipidic assemblies form remain incompletely understood. Here, we monitor the dynamics of ApoA-I-membrane interactions through real-time monitoring of morphological changes, which ensue when ApoA-I is incubated with minimal giant unilamellar vesicles (GUVs) composed of single phospholipids or phase-separating phospholipid-cholesterol mixtures. Our fluorescence and atomic force microscopy measurements reveal that the interaction produces discoidal particles while remodeling the parent vesicle in discrete stages involving membrane poration, solute leakage, vesiculation, and lipid-lipid phase separation. We find that the qualitative phenomenology is robust and fully reproducible for different protein mutants and alleles (WT APOA-1, Δ49ApoA-I, ApoE-3, and ApoE-4) and different lipid mixtures (PC, PC+PS), we tested. Our molecular simulations recapitulate the essential shape changes and further reveal the composition dependence of the interactions. Together, these findings depict key steps in the protein-lipid interactions, which guide the assembly of mesoscopic reconstituted lipoproteins and nanodiscs.

INTRODUCTION

Lipids are an amphiphilic molecule that play essential roles in membrane synthesis, energy storage, and cell signaling. The ability to transport them from the sites of absorption (or syntheses) to other cells and tissues

– as a means to distribute metabolic energy – represents a critical step in the evolution of multicellular

organisms (*1*). In animals, this function is enabled by the collective transport of lipids in the form of micelle-

like, colloidal nanoparticles through the aqueous circulatory system.

Collectively termed lipoproteins, these circulating nanoparticles are supramolecular, mesoscopic complexes

consisting of mixtures of several different classes of lipids (e.g., triglycerides, cholesterol, and

phospholipids) and one or more proteins called apolipoproteins (*2*). Morphologically, they are organized into

spheroidal particles consisting of an outer layer of phospholipids, unesterified cholesterol, and proteins,

encapsulating the central core made up of neutral lipids, predominately cholesteryl ester, and triacylglycerols

(*3*). Although most lipoproteins share these structural features, they also exhibit considerable structural,

compositional, and morphological variations. An early basis for the classification and fractionation of the

lipoprotein diversity was the particle density: different lipoproteins floated differentially in repeated

ultracentrifugations with progressively increased densities (*4*). Subsequent efforts revealed additional criteria

for classifying lipoprotein diversity (*5, 6*), including sizes, electrophoretic mobilities, lipid and protein

compositions, and metabolic functions. Currently, the most common classification recognizes six different

classes of human lipoproteins (*2*). These classes include (i) *chylomicrons*, which are >100 nm in diameter

and commonly synthesized in intestine; (ii) *chylomicron remnants,* which form after lipoprotein lipase

hydrolyze the triglycerides of the source chylomicrons; (iii) *very-low-density lipoproteins* (VLDL, d < 1.006

g/mL and diameter, 30-90 nm), produced in the liver; (iv, v) *intermediate-density lipoproteins* (IDL, d =

1.006-1.019 g/ml) and *low-density lipoproteins* (LDL, d = 1.019-1.063 g/mL).), produced by the lipase

activity on VLDLs; and (6) *high-density lipoproteins* (HDL, d = 1.063-1.21 g/mL), which have the smallest

sizes of 8-12 nm in diameter. Within each of these classes, a significant compositional variability exists,

making lipoproteins one of the most compositionally heterogeneous classes of particles in blood.

In addition to being the smallest (~8-12 nm in diameter) and the densest (d = 1.063-1.21 g/mL), HDLs are

also compositionally and functionally distinct (*7*). Metabolically, they play a major role in reverse cholesterol

transport, a mechanism by which circulating HDLs draw cholesterol from the cells of the peripheral tissues

and return it to the liver for excretion (*8, 9*). This role in mediating macrophage cholesterol efflux is thought

to be primarily responsible for the inverse correlation between their concentrations in plasma levels and the

risk for coronary artery disease (*10, 11*), rendering HDLs uniquely atheroprotective. HDLs are also

compositionally distinct from their lipoprotein counterparts. Their lipids are not solubilized by

apolipoprotein B. Instead, they are bounded, most dominantly, by ApoA-I – a single polypeptide of 243

amino acids secreted in the intestine and liver (*12, 13*) – which accounts for ~70% of the total protein mass

of HDL particles (*7*).

In vivo, HDL assembly begins with the activity of ATP-binding cassette transporter ABCA1, a membrane

protein. According to the alternating access mechanism, an activated ABCA1 flips phospholipids from the

cytoplasmic to the outer, exocytoplasmic leaflets of plasma membranes (*14*). The resulting excess lipid

density (and area) in the outer leaflet promotes vesiculation and efflux of phospholipids and cholesterol to

the extracellular space in the form of vesicles. A recent computer simulation (*15*) proposes an alternate view.

It suggests that upon activation, ABCA1 acts as a phospholipid translocase, extracting lipids from the outer

exocytoplasmic leaflet to form a luminal structure, that migrates into the extracellular space as vesicles. In

both cases, lipid-poor ApoA-I activates ABCA1 and interacts with the vesicles produced by the ABCA1

activity in the extracellular space, leading to the formation of nascent HDLs. These newly formed HDLs are

discoidal in morphology. Subsequent esterification of cholesterol by the enzyme lecithin: cholesterol

acyltransferase (LCAT) initiates the formation of the hydrophobic core and induces the shape transformation

from the discoidal to the spheroidal morphology (*16*).

As discrete nanoscale compartments transporting lipids and membrane proteins through aqueous

environments, HDLs and their synthetic counterparts are appealing for many biomolecular technologies (*17*).

Reconstituted *de novo* by incubating pre-determined mixtures of lipids with apolipoprotein (as well as their

truncated variants called membrane scaffold proteins or MSPs) or synthetic polymers (e.g., styrene-maleic

acid, SMA) (*18, 19*), these reconstituted HDLs (rHDLs) – also called nanolipoproteins (NLPs) and nanodiscs

– mimic an intermediate discoidal state of ApoA-I found *in vivo* (*20*). These constructs are proving valuable

for solubilizing membrane proteins (*21-24*), tailoring membrane compositions (*25*), and screening

membrane-protein targeting drugs and therapies (*26, 27*).

For both the biotechnological and biomedical applications of HDLs, understanding molecular mechanisms

by which ApoA-I disrupts topologically closed lipid vesicles is important. Much of our current knowledge

is obtained from computer simulations and *in vitro* reconstitution studies. Two limiting modes of protein-

lipid association in the formation of discoidal rHDLs have been proposed: (i) the picket fence model, where

two ApoA-I molecules wrap around a disc of a single lipid bilayer with adjacent alpha-helices in antiparallel

orientation, parallel to the acyl chain of the phospholipids (*28, 29*) and (ii) the double-belt model, where two

antiparallel ApoA-I are oriented perpendicular to the lipid tails (*30*). The bulk of the experimental evidence

currently favors the belt model, but the hybrid conformations in HDL subpopulations where some of the

ApoA-I helices are perpendicular to the lipid tails have not been fully ruled out.

For either model, the dynamics of the alpha-helical segments of ApoA-I play important roles. ApoA-I is an

amphipathic protein that contains a globular N-terminal domain (residues 1-43) and a lipid-binding C-

terminal domain (residues 44-243). As secreted, lipid-poor ApoA-I features five amphipathic helices

spanning residues 7-44, 54-65, 70-78, 81-115, and 147-178. Together, these helices make up approximately

50% of the protein content (*31*). The free energies of stability for these helices in the lipid-free state are

estimated to be in the range of 3-5 kcal/mol (~5-9 k_bT), sufficient to provide stable helical segments,

however, these values are lower than 5-10 kcal/mol (~9-17 k_bT) typically observed for globular proteins,

suggesting that the protein is highly conformationally dynamic. Together, these features allow ApoA-I to

unfold and refold readily. This structural plasticity is essential to its function, allowing for the stability of

both the discoidal and spheroidal assemblies with varied lipid stoichiometries. Specific protein

conformational changes – specifically the increase in the alpha-helical content of the protein (*32*, *33*) – that

accompany HDL assembly are now well-understood (*34*).

Still, our understanding of how ApoA-I interacts with organized lipid mesophases, such as membranes,

vesicles, and liposomes, remains incomplete. Computer simulations and *in vitro* reconstitution studies

suggest roles of packing density (*35*), packing defects (*36*), and local membrane curvatures in driving the

assembly (*37*). However, the kinetic pathways and structural intermediates – how ApoA-I invades a

topologically closed membrane, disrupts its structure, assembles lipoproteins, and what remains of the parent

membrane – remain incompletely understood. To begin to address these questions here, we probe the real-

time interactions between a truncated isoform of ApoA-I and lipid bilayers organized as giant unilamellar

vesicles. By employing a platform that enables visual characterization, we identify previously unappreciated

previously unrecognized biophysical steps – including poration, vesiculation, and phase separation – which

mark the ApoA-I – membrane interactions *en route* to the formation of HDL-like particles.

RESULTS AND DISCUSSION

We begin by investigating the interaction between ApoA-I and single lipid bilayers. For our study, we used

a N-terminally truncated isoform of ApoA-I in which the first 49 amino acids are removed: Δ49ApoA-I.

This modification has been previously shown to preserve the essential lipoprotein-lipid interactions (*38*).

Our lipid source for reconstituting HDL-like particles is a suspension of protein- and cytosol-free, model

phospholipid membranes of closed topologies, i.e., giant unilamellar vesicles (GUVs, 15-30 µm diameter)

(*39*). We used both single-component and multicomponent GUVs. Our single component GUVs consist of

the mono-unsaturated phospholipid, 1-palmitoyl 2-oleoyl-*sn*-1-glycero-3-phosphocholine (POPC). Our

multi-component GUVs are reconstituted using ternary lipid mixtures composed of POPC, egg-

sphingomyelin (SM), and cholesterol (Chol) in equimolar (1:1:1) and cholesterol poor (2:2:1) proportions.

Depending on composition and temperature, these lipid mixtures are known to organize as a singular,

uniform phase or exhibit phase separation (*40*), forming two co-existing liquid phases: a dense phase

enriched in SM and Chol, designated as the liquid-ordered phase (L_o) and a second, less dense liquid-

disordered phase (L_d) consisting primarily of POPC. Differentiation between the L_o and L_d phases through

fluorescence microscopy is achieved by doping the GUVs with small concentrations (1 mol %) of a phase

sensitive probe, N-lissamine rhodamine dioleoyl phosphatidylethanolamine (Rho-DOPE, Ex/Em, 560/583

nm) (*41*). At the equimolar (1:1:1) composition, the phase diagram predicts the absence of domain formation

at optical length scales (*40*) but does not rule out existence of nanoscale domains (*42*). Consistent with this

prediction, GUVs encapsulating 100 mM sucrose appear optically homogeneous at room temperature in an

osmotically balanced, isotonic medium containing 100 mM glucose. Monitoring the temporal changes using

spinning disc confocal and wide-field fluorescence microscopy allows us to follow the morphological and

topological transformations of these GUVs.

Introducing Δ49ApoA-I to a suspension of single component, POPC GUVs (decorated with 1 mol %

Rhodamine PE and encapsulating 100μM of an NBD-labeled glucose (NBD-G, Ex/Em, 475/550 nm) in the

interior) triggers gross membrane remodeling (**Supplementary Video 1**) characterized by a well-defined

sequence of structural, morphological, and topological transformations (**Fig. 1**).

Δ49ApoA-I-membrane interactions produce microscopic pores

The first major event involves a transient, short-lived appearance of an intensity modulation at the membrane

surface. Here, the uniform fluorescence intensity decorating the equatorial plane in the confocal fluorescence

images of single GUVs is replaced, transiently, by a Janus binary pattern of fluorescence (**Fig. 1a - 17s**)

consisting of (i): a dominant segment with the fluorescence intensity comparable to that prior to the change

and (ii) a smaller arc or a domain, roughly 2 µm in width, with the vanishingly low fluorescence intensity

comparable to the background value ($I_{membrane} / I_{arc} = 1,400$ %). This intensity perturbation lasts less than 2

seconds before homogenizing. Comparing across multiple experiments, we find that the event is

reproducible; it also exhibits considerable variability in the lag time (10-500 s), time after the introduction

of protein that we witness the event, and the lifetime (200 ms – 2 s), the time over which patterned intensity

persists (**Figure S1**). In all cases, these short-lived arcs of low fluorescence are of microscopic (2 - 5 µm in

diameter) dimensions, reminiscent of a transient microscopic pore in the GUV membrane. The variable lag

time we observe is consistent with the stochastic nature of the pore formation.

The appearance of microscopic pores following ApoA-I interaction begs a question: Are the observed pores

related to the known phenomenon of peptide-induced membrane poration? It is now well established that

many proteins and antimicrobial peptides (e.g., defensin, magainin 2, mellitin, almethicin, protegrin, nystatin

and actinoporins to name a few) – in particular, those that adopt amphipathic α-helical conformation upon

interaction with lipid or cellular membranes – disrupts membrane integrity either by forming transmembrane

pores (*43-49*) or by mechanochemically altering membrane permeability through interfacial activity (*50*). In

all cases, AMP-mediated poration involves the formation of spatially localized, nanoscopic pores. typically

1-3 nm (*47, 48, 51, 52*). The transient microscopic pores, we witness, by contrast, are orders of magnitude

larger, suggesting that they are unlikely to be related to peptide-induced membrane poration.

The most likely explanation is that the pores appear because of a cumulated loss of membrane area caused

by ApoA-I-mediated lipid removal and concomitant formation of rHDL or NLP particles (Further discussion

of this scenario is deferred to a later section in the manuscript).

To directly test whether the incubation of GUVs with Δ49ApoA-I produces NLP particles, we investigated

the GUV bathing medium using high-speed atomic force microscopy (HS-AFM). Scanning the deposits of

Figure 1: (**a**) Confocal xy-scan montage of a POPC GUV doped with 1 mol % Rho-B-DOPE (red) and encapsulating 100μM of a soluble, membrane impermeable dye NBD-G (green). The GUV is exposed to 0.88μM Δ49ApoA-I (Supplementary Video 4). At 17 seconds, a pore formation event can be seen (circled in yellow) and persists for ~2 seconds. Leakage of the encapsulated fluorophore depicted in the second row. Intensity characterized by the yellow line. Vesiculation depicted in the third row, occurring rapidly within 1 second. Scale bar 10μm. (**b**) Montage of a DOPC-DOPS-Rho (79:20:1) GUV exposed to 20μM ApoE4. Vesiculation occurs over several minutes and results in clearly spherical daughter vesicles.

the solution on planar substrates (see experimental methods), we find evidence for the appearance of a population of discrete disk-shaped structures even when GUVs are nominally intact (**Fig. 2a**). These structures have an average height of 4.5 ± 1.1 nm and an average diameter of 14.0 ± 4.0 nm (**Fig. 2b & 2c**) – in excellent agreement with the formation of NLPs ([53], [54]). Consistent with earlier reports of lipid exchange between Δ49ApoA-I NLPs and supported bilayers occurring in 15 minutes ([55]), our results support the picture that Δ49ApoA-I stochastically binds the membrane, extracts lipids, and produces NLPs concomitantly rendering the parent GUV tense, yet nominally intact.

These observations of NLP formation can be readily reconciled with those of transient pore formation (see above). The formation of NLPs requires that the lipids be drawn stochastically from the GUV membrane.

With each peptide-promoted lipid loss the residual GUV membrane experiences a reduced lipid density,

which, in turn, raises the membrane tension (*56*). Of course, this view presumes that the vesicular integrity

is not compromised and that the isotonic conditions within the bath ensure the constant volume [Note that

the difference in solute concentration (Δc) between the bath and the encapsulated solvent is initially isotonic.

Thus, the osmotic pressure acts to maintain the encapsulated volume, as prescribed by Van't Hoff's equation

($\Delta\Pi = RT\Delta c$), where R is the gas constant and T the absolute temperature]. Beyond a threshold lateral

tension, the membrane ruptures through the well-known phenomenon of tension-induced membrane

poration (*57-60*). Specifically, it is now known that when membrane tension exceeds ~6 mN/m, energetics

favor pore formation (*59*). This, in turn, sets the stage for additional morphological changes: (*61*) (i) the open

pore permits the redistribution of lipids, allowing the membrane to relieve lateral tension (σ) , and (ii) the

encapsulated liquid (solute and solvent) leaks through the open pore due to the excess of Laplace pressure,

($\Delta P = 2\sigma/R$) decreasing the radius of the vesicle, R. These changes culminate with the healing of the pore

and the resealing of the membrane. With the continued removal of lipids by ApoA-I, the process can be

repeated multiple times.

Solute leakage through a cascade of pore formation events

A key consequence of the formation of transient pores is the accompanying mixing of the encapsulated

solution with the external bath. To track this leakage, we monitor a membrane-impermeable (permeation

constant, ~6-25 x 10^{-9} cm/s) (*62*) fluorescent marker, NBD-glucose (green, 100 µM), encapsulated within a

GUV. Images shown in **Fig. 1a** reveal that the fluorescence intensity of the dye diminishes gradually, until

optical homogeneity is achieved with the surrounding bath. Observing several different GUVs of different

sizes (r = 3-23 µm), we find that the intensity follows a sigmoidal decay (**Fig. 2d**) with variable lag time

(~10-700 s) and persists over a broad window of time (5-94 s) before reaching a homogeneous plateau

(averaging ~20 ± 7 s (**Fig. 2e**)).

The time taken for the encapsulated intensity of NBD-G to homogenize to the background intensity (~20 ±

7 s) (**Fig. 2e**) is significantly longer than the lifetime of any pore formation events we observe (0.2 – 2 s).

Stochastic pore nature and optical limitations of confocal microscopy limit our direct observations of

transient pores to, at most, a single pore per GUV, however, we often observe secondary evidence of

multiple pore formation events through fluctuations in membrane radius (**Supplementary video 2**). It

seems reasonable that the leakage of the dye occurs through multiple pore formation events: multiple

rounds of lipid removal and tension generation, because of the continued Δ49ApoA-I activity (**Fig. S2**). In

a striking parallel, this behavior resembles that of membrane equilibration under hypotonic stress. Here,

repeated cycles of elemental biophysical processes – including osmotically-mediated water influx,

generation of membrane tension, membrane failure through pore formation, solute efflux through open

pore, and pore healing – lead to osmotic equilibration (*58*).

Figure 2: (**a**) AFM images depicting nanodisc appearance. (**b**) Histogram of NLP height and (**c**) NLP width collected using HS-AFM. (**d**) Three examples of NBD-G intensity leakage profiles from POPC GUVs following the addition of ~0.88 µM Δ49ApoA-I (supplementary video 2) (**e**) Relative frequency histogram for full leakage of NBD-G from POPC GUVs (24 GUVs) ($\bar{t}$ = 20.3s).

Vesiculation and lysis of single-component GUVs

The observations of membrane poration, solute, and solvent exchanges above precede the most consequential structural remodeling leading to large-scale morphological and topological transformations because of the interactions of GUVs with Δ49ApoA-I. Beginning at ~98 s (**Fig. 1a**), the epifluorescence images at the equatorial plane of this POPC GUV show a dramatic membrane restructuring. The optically uniform and continuous circle of fluorescence at the equatorial perimeter is replaced by an open arc terminating into a string of closely spaced, smaller circular features with diameters of less than one micrometer (< 1 µm). These features form abruptly over 2 - 3 frames (< 1 s). The fluorescence emission from these clusters is more intense (I_{small}/I_{mother} = 200%) than that of the surrounding mother membrane. **Supplementary video 2** depicts an example where similar topological change occurs in nearly all POPC GUVs during the observed period (17 min).

These observations are reproducible (**Supplementary Videos 1-6**) across multiple independent experiments. Previous studies report slow spontaneous formation of rHDL by WT ApoA-I from POPC LUVs at 37°C (*63*). Additionally, membrane curvature generation in response to ApoA-I has been previously observed (*37*), without significant elaboration. To understand whether our observations were unique to ApoA-I, we carried out additional experiments utilizing wildtype ApoA-I, ApoE-4, and ApoE-3. When these peptides were introduced to GUVs composed of 4:1 DOPC-DOPS, qualitatively similar results observed. As an example, the breakdowns of a 4:1 DOPC-DOPS GUV exposed to 20 µM and 0.44 µM ApoE-4, respectively, are depicted in **Figure 1b** and **Figure S3**. Another example in **Figure S4** presents a montage of a 4:1 DOPC-DOPS GUV vesiculation in response to exposure to 7.4 µM ApoE-3. Together, these observations indicate

that these instances of membrane restructuring are not limited to the WT ApoA-I isoform or its Δ49 variant.

Instead, they reflect a consistent pattern of membrane deformations characterizing the interaction between

apolipoproteins and lipid membranes.

These large-scale membrane deformations are consistent with a known aspect of ApoA-I-membrane

interactions, namely conformational change. It is now well-established that the membrane binding induces

a significant conformational shift in ApoA-I increasing the helical segment from ~50% to ~80% (*32*, *33*).

The increase in the helical segment population then triggers enhanced "interfacial activity (*50*)," which

reflects the increased ability of the protein to (i) bind to the membrane; (ii) partition into the membrane–

water interface, acting as a wedge altering the packing and organization of the lipids within the membrane

(*64*, *65*); and induce an area mismatch (*66*) between the two individual leaflets that make up the phospholipid

bilayer. These mechanical deformations synergistically serve to reduce the energy cost for the creation of

curvature generation (*67*) and thus promote vesiculation, such as we observe.

Coarse-grained molecular dynamics simulations with WT and Δ49ApoA-I

To further characterize the ApoA-I-mediated membrane deformations, we carried out coarse-grained

molecular dynamics simulations. We studied interactions of both the **WT and Δ49ApoA-I** with lipid

bilayers composed of 100% POPC, 4:1 POPC:POPS, 1:1:1 POPC:DPSM:Chol and the PM8 lipid mixture

(See Methods). For all bilayer compositions, we pre-assembled the double-belt and placed it 4 nm away

from the bilayer surface. For each membrane condition, three independent simulations demonstrate that,

without any restraints, the double-belt protein collapses into an 8-shape within the first 35-100 ns of the

simulations and this conformation remains stable for the rest of the simulations regardless of the membrane

composition (**Fig S5. a-d, the first three rows**). With restraints, we ensure that the double-belt conformation

remain as an open ring that is still able to undergo significant conformational changes (**Fig S5. a-d, the last**

three rows). In 100% POPC membranes (**Fig. S5a**), the WT ApoA-I interacts with the surface, either

continuously or intermittently, without significant perturbation in all simulations except for one, in which a

small portion of POPC lipid is captured by the protein and remains deformed throughout the rest of the

simulation. For membranes composed of 4:1 POPC:POPS (**Fig. S5b**), at least some parts of the WT ApoA-

I continuously interact with the membrane. In 4 of 6 independent simulations containing a 4:1 POPC:POPS

membrane, membranes show mild deformations due to interactions with the protein. In 1:1:1

POPC:DPSM:Chol membranes (**Fig. S5c**), the WT ApoA-I intermittently interacts with the membrane,

without significant morphological defects. Finally, in the PM8 membranes (**Fig. S5d**), it takes about 300 ns

to 1 µs for the WT ApoA-I to fully sit on the PM8 membrane surface (**Fig. 3a**) in all independent simulations

and afterwards, stable protein-membrane interactions are maintained. These stable interactions cause

significant deformations on the surface of PM8 membranes in all independent simulations. Particularly in

the simulations where the protein is restrained to remain flexible and open, the protein causes a bud-like

*Figure 3: Coarse-grained molecular dynamics simulations of the WT and Δ49 ApoA-I. The snapshots of (a) the WT ApoA-I and (b)
Δ49 ApoA-I double-belt proteins in four different membranes are taken at every 1 µs throughout a single representative 5 µs-long
simulation trajectory as indicated by the arrow; the first column of each row corresponds to the initial placement of the pre-
assembled double-belt protein 4 nm away from the membrane surface. In the panels a and b, each row corresponds to a system
containing a different membrane composition: from top to bottom, the systems include a 100% POPC, 4:1 POPC:POPS, 1:1:1
CHOL:DPSM:POPC and PM8 (mammalian plasma membrane model) membrane. The representative simulations shown here are
carried out with the restraints ensuring double-belt proteins remain open and flexible. Lipids are color-coded: CHOL in orange,
DIPE in blue, DPSM in gray, PAPC in pink, PAPS in red, PAP6 in dark red, POPC in white, POPE in cyan and POPS in green.
Water and ions are not shown for clarity. Protein backbone beads are colored in dark gray.*

deformation that is the most significant in comparison to the deformations observed in other membrane
compositions.

These simulations were repeated with Δ49ApoA-I double-belt (**Fig. 3b**). Similar interactions with the POPC,
POPC:POPS, POPC:DPSM:Chol and PM8 membranes were observed. The protein induced the most

pronounced bud-like deformations in the PM8 membrane (**Fig. S5d**), and its interactions were subtly

disruptive for 4:1 POPC:POPS membranes (**Fig 3b., row 2**) when compared to interactions by WT ApoA-I

(**Fig 3a., row 2**).

Overall, the computational data show that ApoA-I interactions depend significantly on the lipid composition

of the membrane in a way that the presence of negatively charged lipid species may help mediate embedding

of the protein into the membrane surface. These MD simulations also show that ApoA-I exhibits long term

association with these membrane compositions. Such an association time would be needed for the kinds of

deformation we observe, experimentally. However, the simulations reveal only subtle morphological

deformation in POPC, a variance with our observations of gross membrane remodeling which ensues upon

peptide interaction with POPC GUVs. In contrast, when we substitute the single component membrane for

a more complex, PM8, the full scope of morphological changes becomes apparent.

Δ49ApoA-I binding-induced phase separation in cholesterol-containing GUVs

The lipid specificity of ApoA-I interactions witnessed during the molecular simulations above raises some

general questions: By sequestering and binding specific sub-classes of lipid molecules, does Δ49ApoA-I

sort the remaining pool of membrane lipids? Can the protein association break symmetry and transform

multicomponent, homogeneous membranes into phase-separated states? These questions are not limited to

Δ49ApoA-I but arise in broad classes of protein-membrane interactions where the preferential association

of proteins with a sub-set of molecules locally can shift the phase space and drive global phase separation.

To begin to explore these questions, we incubated a model ternary GUVs composed of an equimolar mixture

(1:1:1) of POPC, SM, and Chol, with Δ49ApoA-I. To enable visualization of membrane phases by

fluorescence, we doped the lipid mixture with small concentrations of two phase-sensitive lipid probes: (i)

1 mol % of Rho-DOPE (red, Ex/Em 560:583), which preferentially associates with the POPC enriched

liquid-disordered phase (L_d), and (ii) 3 mol % NBD-PE (green, Ex/Em 463:536), which selectively partitions

within cholesterol- and SM-rich, liquid-ordered (L_o) phase. Previous work has established that the presence

of these probe lipids at these concentrations does not influence the phase behaviour of the dominant ternary

lipid mixture (*41*).

A selection of images from the time-lapse movies (**Supplementary Video 4 & 5**) shown in **Fig. 4a** and **Fig.**

4b document how Δ49ApoA-I interaction transforms these multi-component GUVs. The GUV is imaged

from a distal plane, capturing a confocal view of the apical hemisphere (away from the substrate), which

initially appears homogeneous (**Fig. 4a 5:27**). After a short delay, following the addition of Δ49ApoA-1

(8:54), the uniform intensity along the perimeter of the membrane is abandoned, replaced by a pattern of

small domains of high intensity of Rho-DOPE (red). In the subsequent several minutes (12:21), membrane

domains elevated in NBD-PE intensity (green) form simultaneously with the complementary domains

enriched in Rho-DOPE. These segments of the membrane continue to grow and increase in intensity while

*Figure 4: Confocal xy-scans of vesiculation in equimolar GUVs composed of POPC, SM and Chol is doped with 1 mol % Rho-B-DOPE (L_d) (**red**) and 3 mol % NBD-PE (L_o) (**green**). (**a**) Montage of **Supplementary Video 4** depicting phase separation and vesiculation following the introduction of ~3.5 µM Δ49ApoA-1. A pore can be seen to appear beginning at 9:01. Scale Bar 20 µm. (**b**) Montage of **Supplementary Video 5** viewing the upper hemisphere of the vesicle. Scale bar 20 µm (**c**) Channel split and merge of collection of vesiculated spheres from Figure 1. Clear maintained structure within the cholesterol-rich domain (green). Scale bar 10 µm.*

traversing to and aggregating within the distal hemisphere of the vesicle, characterizing the incipient phase
separation.

Instances of phase separation precede large-scale morphological remodeling of the membrane. In the minutes

following the appearance of phase-separated domains (12:21), structures of high-intensity NBD-PE (green)

are observed to appear from the membrane. These structures appear circular and folded with an average

diameter of ~2 µm. They also seem to colocalize with the smaller Rho-DOPE domains (**Fig. 4a (13:40) &**

Fig. 4b (17:41)). Over the next several minutes, formation continues, transforming the GUV into a tightly

associating cluster of spherical features (24:23). These observations are fully reproducible and observed in

multiple independent experiments (**Supplementary Videos 4, 5 & 6**). Across these experiments, the size of

these folded structures ranges from 0.3-4.6 µm in diameter, with a density average of ~0.2 µm^{-2}.

These localized substructures, which emerge at the membrane boundary (**Fig 4a.**), form by folding at the

membrane, appear circular, and remain bound at the membrane surface – consistent with localized

vesiculation. This vesiculation of multi-component GUVs reveals a striking separation of membrane lipid

lipids (**Fig. 4c**). This effect is observed most conspicuously for 2:2:1 (POPC:SM:Chol) GUVs (**Fig. 4d**).

Here, the NBD-PE label (green) indicates that the structured portion of the membrane domain is primarily

composed of cholesterol-enriched, sphingomyelin (**Fig. 4c (L_o) & 4d (L_o)**), while the state of cholesterol-

poor, POPC domains reveal no resolvable structure (**Fig. 4c (L_d) & 4d (L_d)**). Other examples of similar,

'colony-like' formations can be seen in **Figure S4**. This segregation of membrane lipids can be attributed to

the differences in the bending moduli (_68, 69_) of the two co-existing phases. For PC:SM:Chol membranes,

the difference in bending rigidity between the liquid-ordered (red, $\kappa_{L_o} = 6.4 \pm 0.3$ (_70_)) and the liquid

disordered (green, $\kappa_{L_d} = 2.6 \pm 0.4$ (_70_)) membrane domains are reflected in the discrepancy between their

structural appearance. The results are the complementary images we observe in **Figures 4c & 4d**.

It is notable that membrane domains form 1:1:1 POPC:SM:Chol GUVs at room temperature, following the

introduction of Δ49ApoA-I. Under these conditions, the phase diagram predicts the absence of phase-

separated domains at optical length scales (_40_). The appearance of phase-separated membrane domains can

be understood as a consequence of the interaction between Δ49ApoA-I and GUV. Because the latter depletes

lipids from the GUV membranes, the residual membranes accumulate lateral tension. It is known that beyond

a critical value, lateral membrane tension can give rise to isothermal phase separation and poration (**Figure**

S1c), occurring transiently during the equilibration of osmotically swollen GUVs (_58_). However, this

mechanism produces pulsatory phase separation as a response to dynamic membrane tension – induced by

transient poration. Therefore, this mechanism cannot explain the steady state phase separation we observe.

Thus, one mechanism likely responsible for inducing membrane phase separation – membrane binding by

ApoA-I. As an amphipathic α-helical protein, ApoA-I embeds partially into the outer leaflet of the lipid

bilayer at the headgroup-water interface. This then creates localized regions of packing and curvature stress

(_71_), which relieve the elastic energy by driving membrane phase separation.

Membrane interaction with amphipathic α-helical adopting peptides have long reported curvature generation

in BAR domains (_72, 73_), GTPases (_74_), annexin B12, α-synuclein (_75_) and clathrin (_76_). Our experiments

suggest that Apolipoprotein interaction with a vesicular membrane leads to curvature generation (**Fig. 1a &**

1b). The physical space of the peptide acts as a wedge within membrane, supported by the presented coarse-

grained MD simulations depicting ApoA-I association with the outer leaflet of varying membrane

compositions (**Fig. 3a & 3b**). Insertion by the helical segments of ApoA-I into a mixed lipid membrane

induces localized region of altered lipid curvature, packing and composition (_71_), reported to create

microdomains capable of separating lipid species into CO-EXISTING liquid phases (_77_). A cooperative

effect of ApoA-I peptides could lead to the instances of micro-scale domains, of which we observe.

Alternatively, we must consider that the biological function of HDL and ApoA-I is the removal of cholesterol

from the cell through reverse cholesterol transport (RCT) (_8_). If, in the process of forming HDL-like

particles, ApoA-I participates in the preferential removal of cholesterol from GUVs comprised of POPC,

egg-SM, and cholesterol in an equimolar ratio, the phase diagram predicts a shift in the membrane state towards a two-domain mixture ([78](#)). Either or both the cooperative effect of the α-helical peptide disruption and the selective removal of cholesterol could be responsible for the sorting of the membrane lipids into the observed membrane domains.

CONCLUSION

In this work, we monitor the interactions between a truncated isoform of ApoA-1 (Δ49ApoA-I) and lipid bilayers organized as giant unilamellar vesicles. We find that morphological transformations of the parent lipid bilayer *en route* to the reconstitution of HDL-like particles *in vitro* result in gross membrane remodeling, which includes the formation of transient, microscopic pores, solute leakage, and vesiculation. The formation of these pores is typical in instances of membrane tension, likely occurring from the physical removal of lipids from the GUVs membrane by Δ49ApoA-I concomitant with nano lipoprotein particle formation. Subsequently, solute leakage through these physical breaks in the membrane is observed to occur over a timeframe beyond the lifetime of individual instances of pore formation, indicating the presence of a cascade of pore formation events. Furthermore, Δ49ApoA-I interaction results in an energetically driven restructuring of the vesicle membrane into smaller, more stable daughter vesicles, concomitant with NLP formation. We propose this observed vesiculation phenomenon is a result of the interfacial alpha-helical peptides embedding into the membrane, forming smaller vesicles upon the lysis of the GUV. Wildtype ApoA-I, ApoE-3, and ApoE-4 additionally show these structural changes when introduced to 4:1 DOPC: POPS membranes, showing this behavior is not limited to the ApoA-I isoform or its Δ49 truncated variant. This lipid selective behavior is additionally observed through a series of coarse-grained molecular dynamic simulations. Introducing Δ49ApoA-I to a collection of equimolar GUVs composed of POPC, egg-sphingomyelin, and cholesterol results in the appearance of micro-phase separated domains, likely occurring through either a cooperative effect by ApoA-I in altering local lipid curvature or by the preferential removal of cholesterol from the vesicular membrane *en route* to HDL-like particle formation.

FUNDING

This work was performed, in part, under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344. Support for WH, TNO, YW, AN TSC and MAC was provided by the LLNL Laboratory Directed Research and Development program (21-ERD-047). LLNL-JRNL-2001736. Contributions of ANP and CFC were supported by a grant from the National Science Foundation through the award: Biomaterials program, division of materials research, National Science Foundation (2342436).

ACKNOWLEDGMENTS

A.N.P & C.F.C acknowledge the preparation of WT ApoA-I, ApoE-3 and ApoE-4 for these experiment by the J.V. lab and graduate student, Maha Al-Habeeb. A.N.P & C.F.C acknowledge the microscopy facilities provided by the UC Davis LMC. T.S.C & T.N.O thank Livermore Computing and the Livermore Computational Grand Challenge for computing time. Figures were assembled, in part, utilizing the online tool Biorender.

MATERIALS AND METHODS

Materials

1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC), 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC), 1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-L-serine (sodium salt) (POPS), egg-sphingomyelin, Cholesterol, and lissamine rhodamine B 1,2-dioleyl-sn-glycero-3-phosphoethanolamine (Rho-B DOPE) were acquired from Avanti Polar Lipids (Alabaster, AL). Glucose was obtained from Sigma-Aldrich (St. Louis, MO), and 2-deoxy2-[(7-nitro-2,1,3-benzoxadiazol-4-yl) amino]-D-glucose (2-NBDG) was obtained from Cayman Chemical (Ann Arbor, MI). Sucrose was obtained from EMD Chemicals (Philadelphia, PA). Chloroform, head-labeled N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)-1,2-dihexadecanoyl-sn-glycero-3phosphoethanolamine, triethylammonium salt (NBD-PE) and head labeled Texas Red™ 1,2-dihexadecanoyl-sn-glycero-3-phosphoethanolamine, Triethylammonium Salt (Texas Red™ DHPE) was obtained from Thermo Fisher Scientific (Waltham, MA). Δ49Apolipoprotein A-1 was provided by the Lawrence Livermore National Laboratory (Livermore, CA). Wildtype Apolipoproteins A-1, E-3 and E-4 were provided by the John Voss lab.

Glass-bottom, 96-well plates were obtained from Cellvis (Mountain View, CA). Indium tin oxide (ITO)-coated glass slides (5-25 Ω) were obtained from Delta Technologies, LTD (Loveland, CO). Manual Teflon-tipped syringes were obtained from Agilent Technologies (Santa Clara, CA).

Preparation of synthetic model membranes

GUVs

GUV preparation via electroformation is a well-established process (*79, 80*). A lipid stock of a desired composition at a concentration of 2 mg/mL is prepared in chloroform. For single component synthetic membranes, POPC (99 mol %) is dissolved with a Rho-B DOPE label (1 mol %). Cholesterol containing membranes are prepared with a combination of POPC, egg-Sphingomyelin, and Cholesterol with the fluorescent labels of Rho-B DOPE (L_d domain) and NBD-PE (L_o domain) at a molar ratio of either 32:32:32:1:3 or 38.4:38.4:19.2:1:3, respectively. DOPC:POPS or DOPC:DOPS lipid mixtures are doped with prepared with Rho-B DOPE label at a molar ratio of 79:20:1. For all lipid stock, 15 µL is deposited onto the conductive side of two ITO-coated slides and spread evenly on half of the slide as it evaporates.

These slides are further dried inside a vacuum desiccator from anywhere from 2 – 24 h. Once dry, a chamber

is formed using a 1-mm-thick rubber O ring (Ace Hardware, Davis, CA) and sealed with high-vacuum grease

(Dow Corning, Midland, MI). After the O ring is adhered to the conductive side of the ITO-slide, the

chamber is hydrated with ~1 mL of a 100 mM Sucrose solution. The chamber is sealed with the second,

inward facing lipid cake, ITO-slide ensuring no trapped air bubbles. The two slides form the chamber in a

way that the halves without dried lipid are facing in opposite directions to avoid the alligator clips from the

function generator from touching both slides, effectively forming a less resistant pathway for any current. A

2.2 V(pp) AC Sine wave is applied across the two slides at a frequency of 10 Hz for 1 h, followed by another

2.2 V(pp) AC Sine wave at a frequency of 2 Hz for 30 minutes. Vesicles containing egg-SM undergo this

step at a temperature of 55 °C. All samples undergo this step covered to protect from light. After GUV

formation, the chamber is disassembled, and the solution is pipetted into a small centrifuge tube and stored

at 4 °C. Vesicles prepared are either used or discarded within a week of preparation.

Preparation of Δ49ApoA-I and WT Apolipoprotein

Cell-free synthesis of Δ49ApoA-I protein:

Cell-free reactions were set up using RTS 500 ProteoMaster *E. coli* HY Kit (Biotechrabbit GmbH, Hannover,

Germany). Reaction components (lysate, reaction mix, feeding mix, amino acid mix, and methionine) were

combined as specified by the manufacturer. Each 1 mL reaction contained 15 µg Δ49ApoA-I plasmid DNA

and 20 µL FluoroTect™ GreenLys tRNA (Promega, Madison, USA). The reactions were incubated at 30

°C, with shaking at 300 rpm for 18 h in a floor shaker. After the reaction, ApoA-I protein is purified using

nickel affinity chromatography. Briefly, 1 mL of 50% slurry complete His-Tag Purification Resin (Roche

Molecular Diagnostics, Basel, Switzerland) was equilibrated with equilibration buffer (50 mM NaH_2PO_4,

300 mM NaCl, pH 8.0) with 10 mM imidazole (Sigma-Aldrich, St Louis, MO) in a 10 mL chromatography

column. The total cell free reaction was mixed with the equilibrated resin and was incubated/nutated at 4 °C

for 1 h. The column was then washed 6 times with 1 mL wash buffer containing (50 mM NaH_2PO_4, 300 mM

NaCl, 20 mM imidazole, pH 8.0). ApoA-I is eluted with 6 x 300 µl elution buffer (50 mM NaH_2PO_4, 300

mM NaCl, 250 mM imidazole, pH 8.0). All elutions were analyzed by SDS-PAGE and peak fractions

containing labeled protein were combined. Pooled fractions were dialyzed in PBS (pH 7.4) and then stored

at 4 °C. Protein levels were quantified using a Qubit protein test according to manufacturer's instruction

(Thermo Fisher Scientific, Carlsbad, CA).

Production and Purification of ApoA-I protein

Human ApoA-I was expressed in *Escherichia coli* strain BL21 Star (DE3)pLysS cells (Life Technologies,

Carlsbad, CA, USA) from the human ApoA-I gene containing a hexa-His affinity tag at the N-terminus. The

ApoA-I gene was cloned into the pEXP-5 plasmid (Novagen, Inc, Madison, WI, USA) and transferred into

the bacteria and cultivated at 37°C in LB medium with 50 µg/mL of ampicillin and 34 µg/mL of

chloramphenicol. Protein expression was induced for 3-4 h following the addition of 0.5 mM isopropyl-beta-

thiogalactopyranoside (Sigma, St Louis, MO, USA). Following cell disruption, ApoA-I was purified from

the soluble fraction of the cells using a His-Trap-Nickel-chelating column (GE Healthcare, Uppsala,

Sweden) and a mobile phase of phosphate-buffered saline (PBS), pH 7.4 with 3 M guanidine. The protein

was washed in PBS (pH 7.4) containing 40 mM imidazole and then eluted with PBS containing 500 mM

imidazole. Imidazole was removed from the protein sample by dialysis (10K MWCO Slide-A-Lyzer,

ThermoFisher) equilibrated with PBS, pH 7.4. Protein purity was analyzed by SDS-PAGE and concentration

was determined using a Nanodrop 2000c spectrophotometer (Thermo scientific, Waltham, MA, USA),

which was based on the extension coefficient (32,490 l mol^{-1}cm^{-1}) and molecular weight (28,279 Da) of

ApoA-I.

Production and purification of ApoE protein

Human ApoE-3 or ApoE-4 protein was expressed in *Escherichia coli* cells and purified as described

previously ([81](#)). Briefly, one liter of cells in LB broth were grown to mid-log phase at 37 °C. These were

then induced with 0.1% arabinose, and incubated for an additional 4h at 37°C. The cells were harvested by

centrifugation, and inclusions bodies isolated and washed as described previously ([82](#)). To purify the protein,

washed inclusion bodies were dissolved in 8 M urea, 200 mM NaCl, and the sample filtered through a 0.2

µm filter and then chromatographed on a SuperDex 200 (Pharmacia) size exclusion column using a mobile

phase of 8M urea, 200 mM NaCl, 10mM TRIS (pH 7.5) and 1mM EDTA. The fraction was then desalted

using a SuperDex 200 column using a mobile phase of 8M urea, 10mM TRIS (pH 7.5) and 1mM EDTA and

then bound to a mono-Q ion exchange column. The column was washed with 8M urea,10 mM TRIS (pH

7.5) and 1mM EDTA, and the protein eluted using a gradient of NaCl. The protein was then re-folded by

dialysis (Slide-A-Lyzer, ThermoFisher) against 0.1M ammonium bicarbonate and then PBS buffer pH 7.4.

The sample was concentrated using centrifugal spin concentrators with a molecular size cutoff of 30 kDa

(Millipore), and the protein concentration determined using the Pierce BCA kit (ThermoFisher, USA).

ApoA-I incubation assays

GUV incubation with Apoprotein

All spinning-disk confocal fluorescence imaging was taken in conjunction with glass-bottom, 96-well plates

(Cellvis). Experiments observed in this way have a final volume of 100 µL per well. 2 µL of GUV solution

prepared in a 100 mM sucrose solution are added to a bath of osmotically balanced, 100 mM Glucose

solution. The difference in density of the GUVs allows them to settle on the bottom of the well. Control

experiments are imaged in these conditions, otherwise 1-5 µM Δ49ApoA-I, WT ApoA-I or ApoE-4 is added

by micropipette slowly (3-5 s) and gently to the center of the well.

GUV leakage assay

GUVs containing soluble dye are prepared as described above; Slides are hydrated with a 100 mM Sucrose & 100 µM of 2-NBDG dye before the electroformation step. Both POPC and cholesterol containing GUVs are prepared in this way. When observing the leakage of dye from the inside of GUVs, incubation with ApoA-I follows the above-described method. When observing the leakage of dye from extracellular space, the bath is instead prepared with 100 mM Glucose & 100 µM 2-NBDG followed by the introduction of GUVs prepared in only 100 mM Sucrose. Spinning-disk confocal fluorescence imaging with 488 nm & 561 nm lasers were used simultaneously to observe any ApoA-I-induced leakage between the intra- and extracellular space. The difference in fluorescence intensities between these spaces is used for quantification of the leakage over time.

Coarse-grained molecular dynamics simulations

The sequence of human ApoA-I was taken from Uniprot database (Entry P02647 ([83])) and used as an input to *Nanodisc Builder* ([84]) in order to generate an atomistic structural model for the ApoA-I double-belt, referred as WT. The mutant sequence where first 49 residues were deleted was also fed into *Nanodisc Builder* to generate a model for the Δ49ApoA-I double-belt, referred as Δ49. Next, the atomistic models were converted to coarse-grained structural models using *martinize2* with Martini 2.2 force field parameters ([85], [86]). The elastic network was added to the structures with a force constant of 900 kJ/mol/nm^2 and a lower (upper) distance cutoff of 0.5 nm (1.2 nm). A 30 nm x 30 nm POPC, 4:1 POPC:POPS, 1:1:1 POPC:DPSM:CHOL, or PM8 bilayer was built with *Insane* ([87]) and solvated with 0.15 M NaCl and 10% antifreeze water (WF) beads; these systems included the coarse-grained WT and Δ49 belt proteins placed 4 nm away from the membrane. PM8 membrane was set as a symmetric bilayer with the lipid composition of the inner leaflet composition of a typical mammalian plasma membrane model used in a previous study ([88]) and included 13.9% POPC, 7.5% PAPC, 5.4% POPE, 16.1% DIPE, 10.8% DPSM 28% CHOL, 16.1% PAPS and 2.2% PAP6.

For each membrane composition, we carried out three independent simulations for 10 µs, 5 µs and 5 µs using the protocol presented by Ozturk et al. ([89]). We carried out these simulations twice: the one set with restraints on the ApoA-I belt dimers and the other set without any restraints. The first set had two distance restraints that were only in effect if the distances became smaller than 8 nm in WT ApoA-I and 6 nm Δ49, respectively, with a force constant of 1000 kJ/mol/nm^2. Two distance variables were defined using the geometric centers of the residues (1) 110-112 and 250-252 and (2) 280-282 & 410-412 for the WT ApoA-I model and (1) 80-82 and 190-192 and (2) 300-302 and 410-412 for the Δ49 ApoA-I model. These restraints did not allow for full closing-up of the belt but still allowed for large conformational changes. The temperature and pressure were kept at 310 K and 1 bar with v-rescale thermostat ([90]) and c-rescale barostat ([91]) during the production runs as indicated in a previous work ([89]) and are listed in the Supplementary Table 1. Note that LINCS order

and iteration were set to 12 and 2 which are crucial when cholesterol is modeled with Martini 2.2 force field

parameters with a time step of 20 fs (*92*). As suggested in a recent study, to avoid unphysical distortions in

the membrane, the outer cutoff and timesteps between neighbor list updates were set to 1.35 nm and 20 (*93*).

All coarse-grained molecular dynamics simulations were carried out with Gromacs 2024.4 (*94*) and the

restraints were applied using the Colvars module (*95*). The simulation snapshots were generated with VMD

(*96*).

Characterization

Fluorescence microscopy

Spinning-disk confocal imaging and measurements were collected on an Intelligent Imaging Innovations

Marianas Digital Microscopy Workstation (3i, Denver, CO) with attached CSU-X1 spinning-disk head

(Yokogawa; Mushanino-shi, Tokyo, Japan), a Quantem512SC EMCCD camera (Photometrics, Tucson, AZ)

and a Flash 4 ORCA-sCMOS camera (Hamamatsu, Hamamatsu-city, Japan). A Zeiss Plan-Fluor 63x (NA

1.4), oil immersion objective (Carl Zeiss, Oberkochen, Germany) was used in the collection of these data.

Rho-B DOPE (Ex/Em 560:583) was exposed with a 50 mW 561 nM laser line, while NBD-PE (Ex/Em

463:536) and 2-NBDG (Ex/Em 465:540) were exposed with a 50 mW 488 nM laser line. Image processing

was done using Fiji a public-domain software; https://imagej.net/software/fiji/), Slidebook digital

microscopy imaging software (3i), and MatLab (The Math Works, Inc. MATLAB Version 2023a).

Size Exclusion Column Purification (SEC)

The NLPs are analyzed and purified via size exclusion chromatography (SEC) with a semi-preparative

Superose 6 10/300 column (GE Healthcare). SEC was run on a Shimadzu LC-20 HPLC system (Shimadzu)

at flow rate 1mL/min with PBS as running buffer. NLP peak was assessed using absorption at 280 nm and

analyzed with instrument software LC Solutions (Shimadzu).

High-Speed AFM characterization of reactant NLPs

A high-speed atomic force microscope (RIBM, Japan) was equipped with a small cantilever (Ultra-Short

Cantilevers (USC, NanoWorld): spring constant, k = 0.15 N/m, resonance frequency, f = 1200 kHz in water)

and was operated in tapping mode at room temperature. The free oscillation amplitude was 0.9 ~ 1.3 nm, and

the typical set-point amplitude was 85% of the free oscillation amplitude. For the sample stage, a freshly

cleaved mica disk with a diameter of 1.5 mm was fixed on a glass rod with a diameter of 1.5 mm and a

height of 2 mm using epoxy glue. The SEC-purified NLPs was diluted 20x in PBS buffer, and 2ul of the

diluted solution was deposited onto the freshly cleaved mica and incubated for 10 minutes. The scanner was

then mounted above the sample chamber with the cantilever immersed in PBS buffer. HS-AFM images were

processed using ImageJ (National Institutes of Health, Bethesda, MD, USA). The height and width

dimensions of each nanodisc were analyzed with Gwyddion software (*97*) and the data was plotted with

Origin.

REFERENCES

1. R. A. Davis, Evolution of processes and regulators of lipoprotein synthesis: From birds to mammals. *Journal*
of Nutrition **127**, S795-S800 (1997).
2. R. W. Mahley, T. L. Innerarity, S. C. Rall, K. H. Weisgraber, PLASMA-LIPOPROTEINS -
APOLIPOPROTEIN STRUCTURE AND FUNCTION. *Journal of Lipid Research* **25**, 1277-1294 (1984).
3. R. Huang *et al.*, Apolipoprotein A-I structural organization in high-density lipoproteins isolated from human
plasma. *Nature Structural & Molecular Biology* **18**, 416-U444 (2011).
4. R. J. Havel, H. A. Eder, J. H. Bragdon, DISTRIBUTION AND CHEMICAL COMPOSITION OF
ULTRACENTRIFUGALLY SEPARATED LIPOPROTEINS IN HUMAN SERUM. *J. Clin. Invest.* **34**,
1345-1353 (1955).
5. C. J. Packard, J. Shepherd, Lipoprotein heterogeneity and apolipoprotein B metabolism. *Arteriosclerosis*
Thrombosis and Vascular Biology **17**, 3542-3556 (1997).
6. R. I. Levy, D. S. Fredrickson, HETEROGENEITY OF PLASMA HIGH DENSITY LIPOPROTEINS. *J.*
Clin. Invest. **44**, 426-+ (1965).
7. A. Kontush *et al.*, in *High Density Lipoproteins: From Biological Understanding to Clinical Exploitation*,
A. VonEckardstein, D. Kardassis, Eds. (2015), vol. 224, pp. 3-51.
8. G. H. Rothblat, M. C. Phillips, High-density lipoprotein heterogeneity and function in reverse cholesterol
transport. *Current Opinion in Lipidology* **21**, 229-238 (2010).
9. C. J. Fielding, P. E. Fielding, MOLECULAR PHYSIOLOGY OF REVERSE CHOLESTEROL
TRANSPORT. *Journal of Lipid Research* **36**, 211-228 (1995).
10. A. R. Tall, Cholesterol efflux pathways and other potential mechanisms involved in the athero-protective
effect of high density lipoproteins. *Journal of Internal Medicine* **263**, 256-273 (2008).
11. A. V. Khera *et al.*, Cholesterol Efflux Capacity, High-Density Lipoprotein Function, and Atherosclerosis. *N*
Engl J Med **364**, 127-135 (2011).
12. S. K. Karathanasis, V. I. Zannis, J. L. Breslow, ISOLATION AND CHARACTERIZATION OF THE
HUMAN APOLIPOPROTEIN-A-I GENE. *Proceedings of the National Academy of Sciences of the United*
States of America-Biological Sciences **80**, 6147-6151 (1983).
13. V. I. Zannis, F. S. Cole, C. L. Jackson, D. M. Kurnit, S. K. Karathanasis, DISTRIBUTION OF
APOLIPOPROTEIN-A-I, APOLIPOPROTEIN-C-II, APOLIPOPROTEIN-C-III, AND
APOLIPOPROTEIN-E MESSENGER-RNA IN FETAL HUMAN-TISSUES - TIME-DEPENDENT
INDUCTION OF APOLIPOPROTEIN-E MESSENGER-RNA BY CULTURES OF HUMAN
MONOCYTE-MACROPHAGES. *Biochemistry* **24**, 4450-4455 (1985).
14. F. Quazi, R. S. Molday, Differential Phospholipid Substrates and Directional Transport by ATP-binding
Cassette Proteins ABCA1, ABCA7, and ABCA4 and Disease-causing Mutants. *Journal of Biological*
Chemistry **288**, 34414-34426 (2013).
15. J. P. Segrest, W. S. Davidson, J. W. Heinecke, Phospholipid transport by ABCA1: the extracellular
translocase or alternating access model? *Current Opinion in Lipidology* **34**, 208-213 (2023).
16. R. S. Rosenson *et al.*, Cholesterol Efflux and Atheroprotection Advancing the Concept of Reverse
Cholesterol Transport. *Circulation* **125**, 1905-1919 (2012).
17. D. A. Bricarello, J. T. Smilowitz, A. M. Zivkovic, J. B. German, A. N. Parikh, Reconstituted Lipoprotein: A
Versatile Class of Biologically-Inspired Nanostructures. *ACS Nano* **5**, 42-57 (2011).
18. T. H. Bayburt, Y. V. Grinkova, S. G. Sligar, Self-assembly of discoidal phospholipid bilayer nanoparticles
with membrane scaffold proteins. *Nano Letters* **2**, 853-856 (2002).
19. I. G. Denisov, Y. V. Grinkova, A. A. Lazarides, S. G. Sligar, Directed self-assembly of monodisperse
phospholipid bilayer nanodiscs with controlled size. *J. Am. Chem. Soc.* **126**, 3477-3487 (2004).
20. R. O. Ryan, Nanobiotechnology applications of reconstituted high density lipoprotein. *Journal of*
Nanobiotechnology **8**, 1-10 (2010).
21. D. A. Bricarello, E. J. Mills, J. Petrlova, J. C. Voss, A. N. Parikh, Ganglioside embedded in reconstituted
lipoprotein binds cholera toxin with elevated affinity. *Journal of Lipid Research* **51**, 2731-2738 (2010).
22. D. A. Bricarello, M. A. Patel, A. N. Parikh, Inhibiting host-pathogen interactions using membrane-based
nanostructures. *Trends in Biotechnology* **30**, 323-330 (2012).
23. L. Rao, R. Tian, X. Y. Chen, Cell-Membrane-Mimicking Nanodecoys against Infectious Diseases. *ACS*
Nano **14**, 2569-2574 (2020).

24. I. G. Denisov, S. G. Sligar, Nanodiscs for structural and functional studies of membrane proteins. *Nature*
Structural & Molecular Biology **23**, 481-486 (2016).
25. I. G. Denisov, S. G. Sligari, Nanodiscs in Membrane Biochemistry and Biophysics. *Chem. Rev.* **117**, 4669-
4713 (2017).
26. K. O. Baskakova, P. K. Kuzmichev, M. S. Karbyshev, Advanced applications of Nanodiscs-based platforms
for antibodies discovery. *Biophysical Chemistry* **313**, (2024).
27. E. K. Makowski, J. S. Schardt, P. M. Tessier, Improving antibody drug development using
bionanotechnology. *Current Opinion in Biotechnology* **74**, (2022).
28. J. H. Wald, E. Goormaghtigh, J. Demeutter, J. M. Ruysschaert, A. Jonas, INVESTIGATION OF THE
LIPID DOMAINS AND APOLIPOPROTEIN ORIENTATION IN RECONSTITUTED HIGH-DENSITY-
LIPOPROTEINS BY FLUORESCENCE AND IR METHODS. *Journal of Biological Chemistry* **265**,
20044-20050 (1990).
29. A. Jonas, K. E. Kezdy, J. H. Wald, DEFINED APOLIPOPROTEIN A-I CONFORMATIONS IN
RECONSTITUTED HIGH-DENSITY LIPOPROTEIN DISKS. *Journal of Biological Chemistry* **264**, 4818-
4824 (1989).
30. J. P. Segrest *et al.*, A detailed molecular belt model for apolipoprotein A-I in discoidal high density
lipoprotein. *Journal of Biological Chemistry* **274**, 31755-31758 (1999).
31. P. S. Chetty *et al.*, Helical structure and stability in human apolipoprotein A-I by hydrogen exchange and
mass spectrometry. *Proceedings of the National Academy of Sciences of the United States of America* **106**,
19005-19010 (2009).
32. H. Saito, S. Lund-Katz, M. C. Phillips, Contributions of domain structure and lipid interaction to the
functionality of exchangeable human apolipoproteins. *Progress in Lipid Research* **43**, 350-380 (2004).
33. H. Saito *et al.*, α-Helix formation is required for high affinity binding of human apolipoprotein A-I to lipids.
Journal of Biological Chemistry **279**, 20974-20981 (2004).
34. C. G. Brouillette *et al.*, Forster resonance energy transfer measurements are consistent with a helical bundle
model for lipid-free apolipoprotein A-I. *Biochemistry* **44**, 16413-16425 (2005).
35. L. Chièze *et al.*, Difference in lipid packing sensitivity of exchangeable apolipoproteins apoA-I and apoA-II:
An important determinant for their distinctive role in lipid metabolism. *Biochimica Et Biophysica Acta-*
Biomembranes **1818**, 2732-2741 (2012).
36. C. P. L. Wan *et al.*, Apolipoprotein-induced conversion of phosphatidylcholine bilayer vesicles into
nanodisks. *Biochimica Et Biophysica Acta-Biomembranes* **1808**, 606-613 (2011).
37. J. Varkey *et al.*, Membrane Curvature Induction and Tubulation Are Common Features of Synucleins and
Apolipoproteins. *Journal of Biological Chemistry* **285**, 32486-32493 (2010).
38. J. A. Cappuccio *et al.*, Cell-free Co-expression of Functional Membrane Proteins and Apolipoprotein,
Forming Soluble Nanolipoprotein Particles. *Molecular & Cellular Proteomics* **7**, 2246-2253 (2008).
39. N. F. Morales-Penningston *et al.*, GUV preparation and imaging: Minimizing artifacts. *Biochimica Et*
Biophysica Acta-Biomembranes **1798**, 1324-1332 (2010).
40. S. L. Veatch, S. L. Keller, Miscibility phase diagrams of giant vesicles containing sphingomyelin. *Physical*
Review Letters **94**, (2005).
41. T. Baumgart, G. Hunt, E. R. Farkas, W. W. Webb, G. W. Feigenson, Fluorescence probe partitioning
between L-o/L-d phases in lipid membranes. *Biochimica Et Biophysica Acta-Biomembranes* **1768**, 2182-
2194 (2007).
42. F. A. Heberle, J. Wu, S. L. Goh, R. S. Petruzielo, G. W. Feigenson, Comparison of Three Ternary Lipid
Bilayer Mixtures: FRET and ESR Reveal Nanodomains. *Biophysical Journal* **99**, 3309-3318 (2010).
43. L. Kristanc, B. Božič, G. Gomišček, The role of sterols in the lipid vesicle response induced by the pore-
forming agent nystatin. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1838**, 2635-2645 (2014).
44. M. Mally, J. Majhenc, S. Svetina, B. Žekš, Mechanisms of Equinatoxin II-Induced Transport through the
Membrane of a Giant Phospholipid Vesicle. *Biophysical Journal* **83**, 944-953 (2002).
45. M. Zasloff, Antimicrobial peptides of multicellular organisms. *Nature* **415**, 389-395 (2002).
46. A. S. Ladokhin, M. E. Selsted, S. H. White, Sizing membrane pores in lipid vesicles by leakage of co-
encapsulated markers: pore formation by melittin. *Biophysical Journal* **72**, 1762-1766 (1997).
47. K. He, S. J. Ludtke, W. T. Heller, H. W. Huang, Mechanism of alamethicin insertion into lipid bilayers.
Biophysical Journal **71**, 2669-2679 (1996).
48. S. J. Ludtke *et al.*, Membrane Pores Induced by Magainin. *Biochemistry* **35**, 13723-13728 (1996).
49. M.-T. Lee, T.-L. Sun, W.-C. Hung, H. W. Huang, Process of inducing pores in membranes by melittin.
Proc. Natl. Acad. Sci. U.S.A. **110**, 14243-14248 (2013).
50. W. C. Wimley, Describing the Mechanism of Antimicrobial Peptide Action with the Interfacial Activity
Model. *Acs Chemical Biology* **5**, 905-917 (2010).
51. K. A. Brogden, Antimicrobial peptides: Pore formers or metabolic inhibitors in bacteria? *Nature Reviews*
Microbiology **3**, 238-250 (2005).

52.   K. Matsuzaki, O. Murase, N. Fujii, K. Miyajima, An Antimicrobial Peptide, Magainin 2, Induced Rapid
Flip-Flop of Phospholipids Coupled with Pore Formation and Peptide Translocation. *Biochemistry* **35**,
11361-11368 (1996).
53.   T. E. Cleveland IV *et al.*, Small-angle X-ray and neutron scattering demonstrates that cell-free expression
produces properly formed disc-shaped nanolipoprotein particles. *Protein science* **27**, 780-789 (2018).
54.   M. A. Coleman *et al.*, Expression and association of the Yersinia pestis translocon proteins, YopB and
YopD, are facilitated by nanolipoprotein particles. *PloS one* **11**, e0150166 (2016).
55.   A. T. Dang, W. He, D. B. Ivey, M. A. Coleman, T. L. Kuhl, Lipid and Protein Transfer between
Nanolipoprotein Particles and Supported Lipid Bilayers. *Langmuir* **35**, 12071-12078 (2019).
56.   A. S. Reddy, D. T. Warshaviak, M. Chachisvilis, Effect of membrane tension on the physical properties of
DOPC lipid bilayer membrane. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1818**, 2271-2281
(2012).
57.   B. Deryagin, Y. V. Gutop, Theory of the breakdown (rupture) of free films. *Kolloidn. Zh* **24**, 374 (1962).
58.   K. Oglęcka, P. Rangamani, B. Liedberg, R. S. Kraut, A. N. Parikh, Oscillatory phase separation in giant
lipid vesicles induced by transmembrane osmotic differentials. *eLife* **3**, e03695 (2014).
59.   S. U. Alam Shibly, C. Ghatak, M. A. Sayem Karal, M. Moniruzzaman, M. Yamazaki, Experimental
Estimation of Membrane Tension Induced by Osmotic Pressure. *Biophys J* **111**, 2190-2201 (2016).
60.   C. Taupin, M. Dvolaitzky, C. Sauterey, Osmotic pressure-induced pores in phospholipid vesicles.
Biochemistry **14**, 4771-4775 (1975).
61.   E. Karatekin *et al.*, Cascades of Transient Pores in Giant Vesicles: Line Tension and Transport. *Biophysical*
Journal **84**, 1734-1749 (2003).
62.   C. Faure *et al.*, Modeling leakage kinetics from multilamellar vesicles for membrane permeability
determination: Application to glucose. *Biophysical Journal* **91**, 4340-4349 (2006).
63.   M. Fukuda *et al.*, Spontaneous reconstitution of discoidal HDL from sphingomyelin-containing model
membranes by apolipoprotein A-I. *Journal of Lipid Research* **48**, 882-889 (2007).
64.   K. Hristova, C. E. Dempsey, S. H. White, Structure, Location, and Lipid Perturbations of Melittin at the
Membrane Interface. *Biophysical Journal* **80**, 801-811 (2001).
65.   J.-H. Lee *et al.*, Thermal Fluctuation and Elasticity of Lipid Vesicles Interacting with Pore-Forming
Peptides. *Phys. Rev. Lett.* **105**, 038101 (2010).
66.   M. P. Sheetz, S. J. Singer, Biological Membranes as Bilayer Couples. A Molecular Mechanism of Drug-
Erythrocyte Interactions. *Proc. Natl. Acad. Sci. U.S.A.* **71**, 4457-4461 (1974).
67.   H. Bouvrais *et al.*, Softening of POPC membranes by magainin. *Biophysical Chemistry* **137**, 7-12 (2008).
68.   P. D. Sambre, J. C. S. Ho, A. N. Parikh, Intravesicular Solute Delivery and Surface Area Regulation in Giant
Unilamellar Vesicles Driven by Cycles of Osmotic Stresses. *J. Am. Chem. Soc.* **146**, 3250-3261 (2024).
69.   Y. Liu, J. Agudo-Canalejo, A. Grafmüller, R. Dimova, R. Lipowsky, Patterns of Flexible Nanotubes Formed
by Liquid-Ordered and Liquid-Disordered Membranes. *ACS Nano* **10**, 463-474 (2016).
70.   R. D. Usery *et al.*, Membrane bending moduli of coexisting liquid phases containing transmembrane
peptide. *Biophysical Journal* **114**, 2152-2164 (2018).
71.   M. A. Zhukovsky, A. Filograna, A. Luini, D. Corda, C. Valente, Protein amphipathic helix insertion: A
mechanism to induce membrane fission. *Frontiers in cell and developmental biology* **7**, 291 (2019).
72.   A. Mahapatra, C. Uysalel, P. Rangamani, The Mechanics and Thermodynamics of Tubule Formation in
Biological Membranes. *J Membrane Biol* **254**, 273-291 (2021).
73.   B. J. Peter *et al.*, BAR Domains as Sensors of Membrane Curvature: The Amphiphysin BAR Structure.
Science **303**, 495-499 (2004).
74.   M. C. S. Lee *et al.*, Sar1p N-Terminal Helix Initiates Membrane Curvature and Completes the Fission of a
COPII Vesicle. *Cell* **122**, 605-617 (2005).
75.   G. Drin, B. Antonny, Amphipathic helices and membrane curvature. *FEBS Letters* **584**, 1840-1847 (2010).
76.   H. T. McMahon, E. Boucrot, Molecular mechanism and physiological functions of clathrin-mediated
endocytosis. *Nat Rev Mol Cell Biol* **12**, 517-533 (2011).
77.   K. V. Pinigin, T. R. Galimzyanov, S. A. Akimov, Amphipathic peptides impede lipid domain fusion in
phase-separated membranes. *Membranes* **11**, 797 (2021).
78.   D. Marsh, Cholesterol-induced fluid membrane domains: A compendium of lipid-raft ternary phase
diagrams. *Biochimica et Biophysica Acta (BBA) - Biomembranes* **1788**, 2114-2123 (2009).
79.   M. I. Angelova, D. S. Dimitrov, Liposome Electroformation.
80.   N. F. Morales-Penningston *et al.*, GUV preparation and imaging: minimizing artifacts. *Biochim Biophys*
Acta **1798**, 1324-1332 (2010).
81.   J. Petrlova *et al.*, A differential association of Apolipoprotein E isoforms with the amyloid-beta oligomer in
solution. *Proteins* **79**, 402-416 (2011).
82.   J. F. Hess, J. C. Voss, P. G. FitzGerald, Real-time observation of coiled-coil domains and subunit assembly
in intermediate filaments. *The Journal of biological chemistry* **277**, 35516-35522 (2002).

83.   T. U. Consortium, UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Research* **47**, D506-D515
807   (2018).
808   84.   L. R. Kjølbye *et al.*, General Protocol for Constructing Molecular Models of Nanodiscs. *Journal of*
809   *Chemical Information and Modeling* **61**, 2869-2883 (2021).
810   85.   P. C. Kroon *et al.*, Martinize2 and Vermouth: Unified Framework for Topology Generation. *arXiv*
811   **2212.01191v3**, (2024).
812   86.   D. H. de Jong *et al.*, Improved Parameters for the Martini Coarse-Grained Protein Force Field. *Journal of*
813   *Chemical Theory and Computation* **9**, 687-697 (2013).
814   87.   T. A. Wassenaar, H. I. Ingólfsson, R. A. Böckmann, D. P. Tieleman, S. J. Marrink, Computational
815   Lipidomics with insane: A Versatile Tool for Generating Custom Membranes for Molecular Simulations.
816   *Journal of Chemical Theory and Computation* **11**, 2144-2155 (2015).
817   88.   H. I. Ingólfsson *et al.*, Capturing Biologically Complex Tissue-Specific Membranes at Different Levels of
818   Compositional Complexity. *The Journal of Physical Chemistry B* **124**, 7819-7829 (2020).
819   89.   T. N. Ozturk *et al.*, in *Methods in Enzymology*. (Academic Press, 2024).
820   90.   G. Bussi, D. Donadio, M. Parrinello, Canonical sampling through velocity rescaling. *The Journal of*
821   *Chemical Physics* **126**, (2007).
822   91.   M. Bernetti, G. Bussi, Pressure control using stochastic cell rescaling. *The Journal of Chemical Physics* **153**,
823   (2020).
824   92.   S. Thallmair, M. Javanainen, B. Fábián, H. Martinez-Seara, S. J. Marrink, Nonconverged Constraints Cause
825   Artificial Temperature Gradients in Lipid Bilayer Simulations. *The Journal of Physical Chemistry B* **125**,
826   9537-9546 (2021).
827   93.   H. Kim, B. Fábián, G. Hummer, Neighbor List Artifacts in Molecular Dynamics Simulations. *Journal of*
828   *Chemical Theory and Computation* **19**, 8919-8929 (2023).
829   94.   D. Van Der Spoel *et al.*, GROMACS: Fast, flexible, and free. *Journal of Computational Chemistry* **26**,
830   1701-1718 (2005).
831   95.   G. Fiorin *et al.*, Expanded Functionality and Portability for the Colvars Library. *The Journal of Physical*
832   *Chemistry B* **128**, 11108-11123 (2024).
833   96.   W. Humphrey, A. Dalke, K. Schulten, VMD: Visual molecular dynamics. *Journal of Molecular Graphics*
834   **14**, 33-38 (1996).
835   97.   D. Nečas, P. Klapetek, Gwyddion: an open-source software for SPM data analysis. *Open Physics* **10**, 181-
836   188 (2012).
837