Title

Seasonal brain regeneration and chromosome instability are linked to selection on DNA repair in *Sorex araneus*

Authors

William R. Thomas[1][*][†], Tanya M. Lama[1,2], Diana Moreno Santillán[3], Marta Farré[4], Cecilia Baldoni[5,6], Linelle Abueg[7], Jennifer Balacco[7], Olivier Fedrigo[7], Giulio Formenti[7], Nivesh Jain[7], Jacquelyn Mountcastle[7], Tatiana Tilley[7], Alan Tracey[7], David Ray[8], Dina K. N. Dechmann[5,6], Dominik von Elverfeldt[9], John Nieland[10], Angelique P. Corthals[11], Erich Jarvis[7,12], Liliana M. Dávalos[1,13][*]

Affiliations

[1]Department of Ecology and Evolution, Stony Brook University; Stony Brook, NY 11794, United States

[2]Department of Biological Sciences, Smith College; Northampton, MA 01063, United States

[3]Department of Integrative Biology, University of California, Berkeley; Berkeley, CA 94720, United States

[4]School of Biosciences, University of Kent; Canterbury, NJ, United Kingdom

[5]Max-Planck Institute for Animal Behavior; Radolfzell, 78315, Germany

[6]University of Konstanz; Konstanz 78464, Germany

[7]The Vertebrate Genomes Lab, Rockefeller University; New York, NY 10065, United States

[8]Department of Biological Sciences, Texas Tech University, Lubbock, TX 79409, USA

[9]Department of Diagnostic and Interventional Radiology, University Medical Center Freiburg, Faculty of Medicine, University of Freiburg; Freiburg 79211, Germany

[10]Health Science and Technology, Aalborg University; Aalborg 9100, Denmark

[11]John Jay College of Criminal Justice; New York, NY 10019, United States

[12]Howard Hughes Medical Institute; Chevy Chase, MD 20815, United States

[13]Consortium for Inter-Disciplinary Environmental Research, Stony Brook University; Stony Brook, NY 11794, United States

† Lead author

*Corresponding authors. Emails: William.thomas@stonybrook.edu, Liliana.davalos@stonybrook.edu

Abstract

Sorex araneus, the Eurasian common shrew, has seasonal brain size plasticity (Dehnel's phenomenon) and abundant intraspecific chromosomal rearrangements, but genomic contributions to these traits remain unknown. We couple a chromosome-scale genome assembly with seasonal brain transcriptomes to discover relationships between molecular changes and both traits. Positively selected genes enriched the Fanconi anemia DNA repair pathway, which prevents the accumulation of chromosomal aberrations, and is likely involved in chromosomal rearrangements (*FANCI*, *FAAP100*). Genes involved in neurogenesis show either signatures of positive selection (*PCDHA6*), seasonal differential expression in the cortex and hippocampus (Notch signaling), or both (*SOX9*), suggesting a role for cellular proliferation in seasonal brain shrinkage and regrowth. Both positive selection and evolutionary upregulation in the shrew hypothalamus of *VEGFA* and *SPHK2* indicate adaptations in hypothalamic metabolic homeostasis have evolved together with Dehnel's phenomenon. These findings reveal genomic changes central to the evolution of both chromosomal instability and cyclical patterns in brain gene expression that characterizes mammalian brain size plasticity.

49 **Teaser**
50   Genomic and expression variations are key to chromosomal instability and seasonal brain
51   plasticity in the common shrew.

52
MAIN TEXT
Introduction
Sorex araneus, the Eurasian common shrew, exhibits two rare and remarkable phenotypes:
extensive intraspecific chromosomal rearrangements (*1*) and Dehnel's phenomenon, or seasonal
size plasticity (*2–5*). While chromosomal rearrangements are common across species boundaries,
populations of *Sor. araneus* have extensive karyotypic diversity within species, forming distinct
chromosomal races (*1*). This shrew also undergoes dramatic, reversible changes in size, including
the brain, in response to seasonal environment pressures (*2–5*); one of the few mammals known to
deploy this wintering strategy (*6–8*). *Sor. araneus* reach an initial size maximum as juveniles in
their first summer, followed by shrinking of most organs in the autumn, especially the brain. Size
reaches a minimum in winter, with rapid regrowth to a second size maximum the following spring
(*2, 3, 5*), simultaneous with pubescence for the single breeding season of their disproportionately
short lifespan (*9*). Research on the evolutionary basis of these traits is, however, very recent and
has been limited to gene expression analyses (*10, 11*).

67
Previous cellular and molecular investigations of Dehnel's phenomenon in *Sor. araneus* have
discovered that seasonal brain size change occurs without a reduction in neuronal number (*12*),
suggesting the evolution of neuroprotective mechanisms. Seasonal gene expression analyses of
the *Sor. araneus* hypothalamus showed enrichment of upregulated genes involved in apoptosis
regulation and cancer pathways during autumn brain shrinkage, highlighting a balance between
cell proliferation and death (*10*). While the anti-apoptotic gene *BCL2L1* was upregulated during
shrinkage, *SPHK2*, which promotes apoptosis (*13, 14*) and increases activity in the brains of
patients with Alzheimer's disease (*15*), was evolutionarily upregulated compared to other
mammals. These findings pointed toward a finely tuned system that enables common shrews to
reversibly regulate brain shrinkage while avoiding the detrimental effects typically associated
with neurodegeneration.

79
Despite those insights, the evolutionary processes involved in the traits that make *Sor. araneus*
unique, extensive chromosomal rearrangements and Dehnel's phenomenon, remain unknown.
While intraspecific karyotypic diversity in mammals is not uncommon (*16, 17*), the more than 75
distinct chromosomal races in *Sor. araneus* indicate unique chromosomal instability (*1*).
Similarly, Dehnel's phenomenon, though most pronounced in the common shrew, is
exceptionally rare among mammals, found only in some red-toothed shrews (Soricinae) (*5*), the
European mole (*Talpa europaea*) (*6*) and two mustelid species (*Mustela erminea, Mustela nivalis*)
(*8*), with a similar phenotype in a domesticated ferret (*Mustela putorius furo*) (*18*) and the pygmy
shrew (*Suncus etruscus*) (*7*). Dehnel's phenomenon is rare, as is the genomic instability of *Sor.*
araneus. While the former must be associated with positive selection to adapt to the
ecophysiological challenges these species face, the latter could result from relaxed negative
selection on genome repair mechanisms.

92
Dehnel's phenomenon is hypothesized to have evolved as an adaptive tradeoff between size and
metabolic demand during winter (*5*). This strategy is critical for the common shrew, which has
one of the highest mammalian basal metabolic rates per unit of body mass (*19*). By reducing the
size of energy expensive tissues, shrews decrease the energy required for their maintenance and
movement (*3, 11, 20–26*). Recent evidence on the metabolic and gene expression changes

concurrent with seasonal size change in the common shrew has lent support to this hypothesis
(*11*). For example, blood metabolomics and liver transcriptomics indicate increased lipid
metabolism and gluconeogenesis during shrinkage (*11*). These may be regulated by adaptive gene
expression of the hypothalamus blood brain barrier (*10*). The evolution of efficient metabolic
regulation systems is therefore expected during seasonal size change.
103
A new chromosome-level genome assembly for the common shrew provides an opportunity to
investigate how selection has shaped both the convergently evolved Dehnel's like phenotypes as
well as the unique chromosomal instability of these shrews. We compared *Sor. araneus*
chromosome rearrangements to a closely related species (*Sun. etruscus*) and applied branch-site
models to identify positively selected genes (PSGs) by comparing the common shrew genome to
39 mammalian genome assemblies. To further relate molecular evolution to Dehnel's
phenomenon, we tested for parallel evolution with three other species exhibiting Dehnel's
phenomenon (*Suncus etruscus*, *Mustela erminea*, *Mustela putorius furo*) and integrated results
with seasonal gene expression data from the common shrew cortex and hippocampus. We
discovered gene enrichment of *Sor. araneus* specific PSGs in the Fanconi anemia pathway
(*FANCI*, *FAAP100*, *PALB2*), associated with DNA repair and longevity, as well as in *VEGFA*, a
candidate hypothalamic regulator of metabolism in the common shrew. Genes associated with
neurogenesis (*PCDHA6, SOX9*) also showed signatures of parallel evolution across species with
Dehnel's phenomenon, complementing the seasonal changes identified in the hippocampus and
cortex linked to cell proliferation (*SOX9*) and Notch signaling. Together, these integrated
genomic and expression analyses implicate changes in genes involved in metabolism and cell
proliferation in the evolution of Dehnel's phenomenon, while also characterizing evolutionary
processes potentially associated with chromosomal rearrangements and longevity.
122
Results
Genome assembly, annotation and synteny
We sequenced, assembled, and annotated a chromosome-scale reference genome assembly for
one male Eurasian common shrew, *Sorex araneus*, following the Vertebrate Genomes Project
(VGP) assembly pipeline (*27*) using a combination of Illumina short-read-, PacBio long-read-,
and Hi-C sequencing. PacBio sequencing data had 35.71X genome coverage with a genome size
of ~2.40Gb. We used Hi-C chromosomal interactions to further manually curated the genome and
were able to anchor >99.8% of the genome to 12 chromosomes (2N=24) (Figure 1A) with a
scaffold N50 of 374MB (Figure 1B) and an estimated a genome size of ~2.23Gb consisting of
73% unique or non-repetitive sequence. By comparison, the previous *Sor. araneus* genome
assembly from the Broad Institute consisted of 12,845 unplaced scaffolds (Table1, Figure 1B).
We were not able to recover the Y chromosome. Annotations using TOGA's machine learning
ortholog classifier identified 42,221 transcripts associated with 17,246 genes (BUSCO
mammalia_odb10 9,005 complete genes; 97.6%), of which 15,980 were one-to-one orthologs to
the human genome (Figure 1C). These numbers are similar to those identified with RNA
sequencing, which found 45,362 transcripts associated with 24,205 coding loci, of which 15,752
are annotated genes.
140
Pairwise alignments between *Sor. araneus* and other species of the order Eulipotyphla with high
quality genome assemblies showed conserved syntenic regions between species. Comparing the
Eurasian common shrew to its closest relative in the data set, the pygmy shrew *Sun. etruscus*, we
found 236 orthologous blocks (Figure 1D). These blocks show that *Sor. araneus* chromosome 10
is orthologous to *Sun. etruscus* chromosome 11, and *Sor. araneus* chromosome 11 is orthologous
to *Sun. etruscus* chromosome 17. Pairwise comparisons also show that large portions of *S.*
araneus chromosome 7 are syntenic with *Sun. etruscus* 3, and *Sor. araneus* 8 with *Sun. etruscus*

14, while *Sor. araneus* chromosome X appears to be fusions of *Sun. etruscus* X, 5, and 12, with
Sor. araneus chromosome 5 being fusions of *Sun. etruscus* 6, 9, and 16. Meanwhile, *Sor. araneus*
chromosomes 2, 3, and 4 show extensive rearrangements compared to *Sun. etruscus*. Lastly, we
inferred the ancestor of Eulipotyphla had 36 chromosomes using DESCHRAMBLER (*28*).

Positive selection in Sorex araneus
We then analyzed the molecular evolution between the common shrew and other mammal species
by testing for positive selection in more than 15,000 single copy orthologs using adaptive branch-
site random effects likelihood (aBSREL) models implemented in the HyPhy suite (v2.5.32) (*29*).
We inferred 676 genes had been influenced by positive selection (PSGs; $p_{adj} < 0.05$) in *Sor.*
araneus (Figure 2A). Gene set enrichment analysis identified eight Kyoto Encyclopedia of Genes
and Genomes (KEGG) pathways enriched by *Sor. araneus* PSGs, consisting of four related to
immunity, including complement and coagulation cascades (4.1-fold enrichment, $p<0.01$),
hematopoietic cell lineage (3.1-fold enrichment, $p<0.05$), primary immunodeficiency (5.1-fold
enrichment, $p<0.05$), and inflammatory bowel disease (3.6-fold enrichment, $p<0.05$).
Additionally, the Fanconi anemia pathway, associated with genomic stability, was enriched (4.3-
fold enrichment, $p<0.05$), including genes *FANCI* (Figure 2), *PALB2, FAAP100,* and *REV3L.*
Exploratory analyses on background branches inferred 2997 genes to be evolving under positive
selection in at least one background species, of which 288 genes showed signals of pervasive
selection throughout the phylogeny (>5 background species). Of the 676 *Sor. araneus* PSGs, 231
genes showed no signal of background selection and were thus specific to this species.

Parallel evolution of Dehnel's phenomenon
We tested for selection in more than 10,000 genes annotated in all the selected mammal species
with Dehnel-like phenotypes (*M. erminea, M. putorius furo, Sun. etruscus*) and inferred positive
selection in over one hundred genes per species, with various degrees of parallel evolution
between species. We inferred 225 genes to be under positive selection ($p_{adj} < 0.05$) in *M. erminea,*
110 genes in *M. putorius furo,* and 904 genes in *Sun. etruscus* (Table 2). Accounting for
background selection, eight genes show signatures of parallel evolution with only episodic (<5
species) background selection (*ASPHD1, PCDHA6, TNK2, B9D1, MEGF8, TTLL10, RNF34,*
LAMTOR4), significantly more genes than expected by chance (100,000 permutations, $p<0.0001$),
suggesting non-independence with lineage-specific traits, i.e., Dehnel's phenomenon (Figure 3).
A single gene, *PCDHA6*, was parallel between *Sor. araneus, M. erminea,* and *M. putorius furo*
with no background selection. MEME analyses were used to detect specific sites evolving under
positive selection on *Sor. araneus* PSGs and found several sites to be convergent across species
with Dehnel's-like phenotypes. Across the eight genes inferred to be under parallel evolution, 10
sites were found to have positive selection in multiple species with Dehnel's-like phenotypes.

Seasonal brain expression
To explore both the regulatory mechanisms of seasonal brain size change and relationships
between molecular evolution and function, we tested for differential expression between the
shrinking (autumn) and regrowth (spring) phases of Dehnel's phenomenon in two brain regions,
the hippocampus and the cortex, and identified any overlapping PSGs. In the hippocampus, we
found 77 differentially expressed genes (DEGs) between autumn and spring, with 52 significantly
upregulated in spring and 25 significantly upregulated in autumn (Figure 4). Using a ranked gene
set enrichment analysis, we discovered an enrichment of 17 pathways (7 upregulated in autumn,
10 upregulated in spring), which included many pathways associated with metabolism; PPAR
signaling ($p_{adj}<0.01$, NES=1.81), fatty acid metabolism ($p_{adj}<0.01$, NES=1.81), and oxidative
phosphorylation ($p_{adj}<0.01$, NES=1.81) (Figure 3B). In the cortex, we identified 122 significant
DEGs, with 46 upregulated in autumn shrinkage and 76 upregulated in spring regrowth. 10

pathways were enriched with genes varying between these seasons, including five that overlapped
in the same direction with those enriched in the hippocampus; ribosome (cortex:$p_{adj}<0.001$,
NES=-2.35; hippocampus: $p_{adj}<0.001$, NES=-2.36), protein export (cortex:$p_{adj}<0.001$, NES=-2.31;
hippocampus: $p_{adj}<0.01$, NES=-2.05), oxidative phosphorylation (cortex:$p_{adj}<0.001$, NES=-2.02;
hippocampus: $p_{adj}<0.01$, NES=-1.78), proteasome (cortex:$p_{adj}<0.01$, NES=-2.05; hippocampus:
$p_{adj}<0.05$, NES=-1.67), and Notch signaling (cortex:$p_{adj}<0.01$, NES=1.98; hippocampus:
$p_{adj}<0.05$, NES=1.70). Of the DEGs, 6 genes in the cortex and 5 genes in the hippocampus
overlapped with PSGs, three of which were found in all three data sets (*SOX9*, *RIN2*, *STMN2*)
(Figure 5).
207
Hypothalamus expression overlap
There were many instances of overlap between PSGs and prior molecular results. Previous
analyses identifying differentially expressed genes in the hypothalamus between seasons of
Dehnel's phenomenon (*10*) only consisted of six overlapping genes with PSGs (*KCNK12*, *COCH*,
RSPH3, *KNDC1*, *PARP4*, *FAN1*). However, comparing PSGs to previous work identifying
branch specific change in hypothalamus expression in *Sor. araneus* (Figure 5), we found overlap
in genes related to hypothalamus development. Eleven PSGs (*KIAA0930*, *LOXL1*, *MLST8*, *PIGR*,
VEGFA, *GALNT12*, *POLDIP3*, *NXPH3*, *SPHK2*, *GRIP2*, *FANCI*) overlapped with those having
higher expression in the *Sor. araneus* hypothalamus compared to other mammalian species.
VEGFA was found in this overlap which we previously hypothesized was important for *Sor.*
araneus hypothalamus fenestration to improve signaling of metabolic demands across the blood
brain barrier. Of the eight overlapping genes, three were *Sor. araneus* specific, *LOXL1*, *GALNT12*
and *FANCI*. Overlap between PSGs and functional data may indicate selection in coding
sequence to alter gene function and regulation.
222
Discussion
Links between chromosomal evolution, genomic stability, and longevity
Robertsonian chromosomal fusions and rearrangements play a central role in karyotype evolution
in the common shrew. This individual *Sor. araneus* had many fewer chromosomes (2N=24) than
the pygmy shrew, *Suncus etruscus* (2N=42), indicating many chromosomal fusions in the
Eurasian common shrew lineage, or fissions in the pygmy shrew lineage have occurred since their
most recent common ancestor (~20mya) (*30*). Ancestral inference supports fusions in *Sor.*
araneus. Pairwise comparisons (Figure 1) between *Sor. araneus* and *Sun. etruscus* show that
while some chromosomes remain conserved (*Sor. araneus* C7 ≅ *Sun. etruscus* C3, *Sor. araneus*
C10 ≅ *Sun. etruscus* C11, *Sor. araneus* C11 ≅ *Sun. etruscus* C17), many are the result of
chromosomal fusions (*Sor. araneus* CX ≅ *Sun. etruscus* CX+ C5+C12, *Sor. araneus* C5 ≅ *Sun.*
etruscus C6+C9+C16, *Sor. araneus* C8 ≅ *Sun. etruscus* C14+C19). Evolutionary relationships for
many chromosomes are less clear, however. Complicating ancestral inference, *Sor. araneus* has
an unusual propensity for chromosomal rearrangements (*31*), with over 75 chromosomal races in
a single species (*31*). Based on chromosome number and geography, the closest land races with
11 autosomes are the Cordon and Pelister groups (*31*), but these populations are ~400 km and
1800 km from the German population, suggesting that this local population may be a distinct
chromosomal group.
241
We found positive selection in genes involved in maintaining genomic stability and their
substitutions may explain the evolution of extreme intraspecific chromosomal variation in this
species (Figure 2). Chromosome size, arrangement, and number can be disrupted by double-
stranded breaks in DNA followed by improper repair (*32*, *33*), but evolved mechanisms either
prevent such damage or ensure accurate repair. The highly conserved FA gene complex (*34*),
comprises 20 genes that identify and repair crosslinked DNA via homologous recombination (*35*).

Several genes associated with this complex were positively selected (six total and four *Sor.*
araneus specific PSGs) in *Sor. araneus*. We found evidence of positive selection in *FANCI*,
whose encoded protein heterodimerizes with FANCD2 to form an ID complex that localizes to
chromatin in response to damage (*35–37*). We also identified several positively selected genes
related to the FANCI-FANCD2 DNA repair complex: *FAAP100*, which is required for core
complex stability with knockouts leading to FA-mediated genomic instability (*38*), and *PALB2*,
whose involvement in the PALB2-BRCA2 complex promotes chromatin stability and
homologous recombination DNA repair pathways (*39, 40*). Substitutions on this pathway in *Sor.*
araneus may impair or alter DNA repair and recombination processes and thus increase the
generation of structural variation. As genetic variation provides the substrate for evolutionary
change, these rearrangements could also further enhance subsequent local adaptation within this
geographically widespread species.
260
In *Sor. araneus*, substitutions in genes in the Fanconi anemia pathway may be associated with
effects of DNA repair on longevity. DNA-damage responses and genomic stability are linked
with lifespan evolution, as DNA must be constantly repaired throughout life to reduce the risk of
cancer and other age-related diseases (*41–44*). Mechanistically, impaired Fanconi anemia DNA
repair pathways are linked to premature aging (*45*) and cancers (*46, 47*). Conversely, strong
positive selection in DNA repair mechanisms, including *FANCI*, has been found in parrots and
other long lived bird species (*48*). In contrast, *Sor. araneus* have a lifespan half that predicted by
their size (*9*), reducing the value of maintaining genomic stability. This would relax negative
selection on DNA repair pathways and, in turn, co-opt them into alternative functions perhaps
related to the propensity for chromosomal rearrangements. But this would also contribute to the
short lifespan of *Sor. araneus*, as such modifications would expose older shrews to age-related
diseases. Genes within this pathway are excellent candidates for testing for involvement in
chromosome evolution of *Sor. araneus* population, as well as longevity, with potential
implications for aging research.
275
Neuronal proliferation, differentiation, death, and Dehnel's phenomenon
Distinct molecular changes likely underlie the evolution of Dehnel's phenomenon, as molecular
parallelism associated with these phenotypes was rare. Tests for parallel molecular evolution
uncovered eight genes overlapping across species with Dehnel's-like traits (Figure 3),
significantly more than expected by chance (p<0.0001). Consistent with an association with
extraordinary brain size plasticity, the gene encoding protocadherin alpha 6, *PCDHA6*, is the only
one to show signatures of parallel positive selection across species with Dehnel's-like phenotypes
without background selection. *PCDHA6* is part of the protocadherin alpha cluster, a group of cell
adhesion molecules central to brain organization, neuronal circuity, axon maintenance, and
synaptic plasticity (*49, 50*). While in mice *PCDHA* knockout leads to miswiring of hippocampal
connectivity despite normal neurogenesis (*51*), in cell lines, *PCDHA6* knockout results in altered
neurite morphology (*52*). Indicating functional conservation, there was high sequence
conservation in species in which this gene was present (15 of 40 assemblies lost or missed the
gene). Substitutions in *S. araneus, M. erminea,* and *M. putorius furo* may contribute to maintain,
rewire, or regenerate synapses during seasonal changes in brain size and morphology, but
PCDHA6 was not detected in brain expression, weakening the case for the genetic changes found
to contribute to brain size plasticity. This dearth of parallel positively selected genes found in
Dehnel's-like phenotypes without background selection makes the common shrew a uniquely
important model for researching brain size plasticity.
295
296 Comparative tests for convergent evolution did not identify strong candidate genes associated
297 with Dehnel's phenomenon across species, so we focused on *Sor. araneus* specific PSGs and

analyzed gene expression, both seasonally variable and evolutionarily upregulated (the latter previously published) (*10*). While to the best of our knowledge there is no prior study of sequence and expression patterns with the massive changes seen in Dehnel's phenomenon, the question of how sequence and expression evolve provides crucial context for our results. First, genes expressed in the brain diverge less in their sequences than those expressed in other tissues (*53*). Second, expression divergence in the brain is also diminished compared to other tissues (*54*). Third, although the rate of nonsynonymous substitutions sometimes correlates with expression divergence (*55*) (e.g., in *Drosophila* (*56*)), this is not always the case especially when analyzed between species (e.g., in birds (*57*)). In short, divergence in either sequence or expression of brain-expressed genes is rare, and the two types of change may or may not be correlated.

Despite the expectation of strongly conserved brain gene expression (*58*), including throughout development (*59*, *60*), we identified seasonal expression plasticity in pathways related to cellular proliferation, differentiation, and death. Previously, expression analyses of the *S. araneus* hypothalamus indicated anti-apoptotic pathways, mediated by *BCL2L1* and *NFKBIA*, are upregulated in autumn (*10*). That suggested the common shrew brain actively regulates pathways to avoid cell loss, consistent with the constant or increasing number of neural cells during brain shrinkage (*12*). But we did not observe upregulation of those pathways in the hippocampus or cortex during autumn. Instead, we identified significant changes in Notch signaling in both regions. In *Sor. araneus*, Notch signaling was under expressed during brain shrinkage (Figure 4). Notch signaling is a highly conserved pathway in animals, well characterized for its role in development (*61*) and a master regulator of plasticity in the adult brain of model organisms (*62*, *63*).

In the *Sor. araneus* hippocampus and cortex, we found autumn changes in gene expression consistent with enhanced neurogenesis. While the precise effects of the Notch signaling decrease are unknown, decreased Notch can promote neural stem cell proliferation and increased adult neurogenesis (*64*), depleting stem cell populations, and hindering neuronal migration (*65*). In spring, *Sor. araneus* revert to elevated expression of *NOTCH1*, which can decrease the rate of neuron establishment between winter and summer in the hippocampus and cortex (*12*). Notch signaling also regulates *SOX9* expression, which is critical for maintaining neural stem cells (*66*, *67*), with *SOX9* knockdown in mice leading to increased neurogenesis (*68*). As with Notch signaling, we observed decreased *SOX9* expression in the cortex and hippocampus of autumn, shrinking *Sor. araneus* (Figures 4 and 5). Consistent with our interpretation of enhanced autumn neurogenesis, winter common shrews increased neuron counts in the hippocampus and cortex compared to summer juveniles (*12*). Despite being highly conserved among background species, as expected for a developmental and brain-expressed gene, *Sor. araneus SOX9* is also under positive selection (Figure 5). Most disease related genetic variation in *SOX9* is associated with developmental disorders, particularly chondrogenesis (cellular bone proliferation) and sex determination (*69–71*), however, coding changes could alter the function of this gene in hitherto unsuspected *Sor. araneus* specific neurogenic processes.

Metabolic homeostasis and Dehnel's phenomenon
PSGs related to both metabolic regulation and cell proliferation (n=11) also overlapped with genes evolutionarily upregulated in the *Sor. araneus* hypothalamus, supporting the hypothesis that Dehnel's phenomenon evolved as metabolic adaptation. *VEGFA* and *SPHK2* were under positive selection in *S. araneus* while also showing an approximate four-fold upregulation in the *Sor. araneus* hypothalamus compared to other mammals (Figure 5) (*10*). Upregulation of *VEGFA* can increase blood brain barrier permeability, improving nutrient sensing in the brain, while upregulation of *SPHK2* may be involved with neuronal cell death, as overexpression of *SPHK2* in

stressed mice cells promotes apoptosis (*13*, *14*). Positively selected substitutions may benefit the common shrew. For example, we propose substitutions in *VEGFA* can alter blood brain barrier vascularization in *S. araneus*, as different isoforms can be pro- or anti-angiogenic (*72*, *73*), while substitutions in *SPHK2* may reduce the apoptotic effect of this gene in the brain, allowing evolutionary overexpression of each in the *Sor. araneus* hypothalamus. Future mechanistic experiments can test both functional hypotheses.

Limitations
We conducted comparative analyses to characterize chromosome evolution and identify positive selection in the protein-coding genes of *Sor. araneus* and other species with Dehnel-like phenotypes, while also examining these results in relation to seasonal and evolutionary changes in brain RNA expression. However, this study has several limitations. To determine if the Radolfzell population is a distinct chromosomal race requires additional cytogenetic verification. While protein coding genes are essential, they are not the sole contributors to phenotypic information within the genome (*74*). A significant portion of the vertebrate genome is non-coding, consisting of Conserved Non-coding Elements (CNEs) and cis-regulatory elements (CREs) that play crucial regulatory roles in gene expression, including in brain development (*75*) and in alternative wintering strategies (*76*). As we explore gene regulation associated with Dehnel's phenomenon across diverse tissues, future studies should investigate the evolution of common shrew CNEs and CREs, feasible as more chromosome-level genome assemblies of Eulipotyphlans are sequenced. Our findings will benefit from mechanistic validation, as gene function in rodent or primate models may differ from those in the shrew, especially in cases involving coding changes that affect protein structure and function. Perturbation experiments, particularly in candidate genes associated with metabolic regulation and cellular proliferation, will provide future insights into the molecular mechanisms of mammalian brain shrinkage and regrowth.

Conclusion
Sorex araneus has evolved a unique set of traits, including a high rate of chromosomal rearrangements and more seasonal brain size plasticity than any other mammal. Yet, the impacts of both selection on protein-coding genes and gene regulation on these traits were unknown. We therefore generated a chromosome-level genome assembly and conducted comparative genomics and seasonal transcriptomic analyses, with three key findings. First, genes in the Fanconi anemia DNA repair pathway (*FANCI*, *FAAP100*, *PALB2*) are under positive selection specific to the *Sor. araneus* lineage. We propose these changes contribute to the Eurasian common shrew's increased chromosomal rearrangements and reduced lifespan. Second, genes and pathways involved in neurogenesis (*SOX9*, Notch signaling, *PCDHA6*) appear critical to the evolution and regulation of Dehnel's phenomenon, suggesting processes related to adult neurogenesis may mitigate the negative effects of decreased brain size. Third, genes previously identified as upregulated in the *Sor. araneus* hypothalamus were also under positive selection in *Sor. araneus* (*VEGFA*, *SPHK2*), highlighting potential synergies between protein-coding and expression adaptations to improve metabolic homeostasis in Dehnel's phenomenon. Positive selection on sequences is concentrated in pathways involved in DNA repair, metabolic homeostasis, and cellular proliferation, processes we propose are integral to the plasticity of the genome and brain of this species.

Materials and Methods
Genome Sequencing and Assembly
A single male European common shrew (*Sorex araneus*) was sampled to generate a highly contiguous reference genome from a population in Radolfzell, Germany (47.9684N, 8.9761 E) under protocols authorized by Regierungspräsidium Freiburg, Baden-Württemberg (35-9185.81/G-19/131). This shrew was euthanized and subsampled into tissues which were

immediately flash frozen using liquid nitrogen, and then stored at -80°C until DNA extraction. Nucleic acid extraction, genome sequencing, and assembly were conducted by the Vertebrate Genomes Laboratory (pipeline v2.0). High-molecular-weight (HMW) DNA was extracted from spleen tissue using the Bionano Prep Animal Tissue DNA Isolation Kit according to manufacturer protocols. HMW DNA was used to prepare libraries for three sequencing types from listed kits: 1) SMRTbell Template Prep Kit for PacBio Sequel II HiFi, 2) Bionano Prep Labelling NLRS for Bionano Genomics DLS optical mapping, and 3) Arima-HiC kit for Arima Hi-C v2 chromatin interactions. Following sequencing, initial sequencing metrics were assessed with GenomeScope (*77*). Contigs were generated by assembling and phasing PacBio reads with Hifiasm (v0.15.4) (*78*). Duplications representing heterozygous contigs were identified and reassigned with the purge duplications workflow (v.1.2.5) (*79*). Contigs were then scaffolded using Bionano optical maps with Bionano Solve (v.3.6.1). The assembly was further assembled into chromosomal-level scaffolds with Hi-C chromatin interaction data using SALSA2 (v2.2) (*80*), followed by manual curation for errors with gEVAL (*81*). Assembly metrics were benchmarked throughout the pipeline using Merqury (v1.1) (*82*), Quast (v5.0.2) (*83*), and BUSCO (v3.0.2) (*84*).

Genome Annotation and Alignment
Gene annotation and orthology inference for *Sor. araneus* and *Mustela nigripes* were predicted with projections from the human genome. First, curated assemblies of these two species were masked for repetitive regions through the creation of a *de novo* repeat library with RepeatModeler (v2.0.2) (*85*) followed by soft masking with RepeatMasker (v4.1.2) (*86*). Masked genomes were then pairwise aligned to the human reference genome (hg38) using lastz (v1.04.15) with the following alignment parameters (BLASTZ_O=400, BLASTZ_E=30, BLASTZ_M=254, chainLinearGap=loose). Sensitivity and specificity of generated pairwise whole-genome alignments were improved using RepeatFiller (v1.0) (*87*) and chainCleaner (*88*). Resultant alignment chains were then used as inputs to predict orthologous gene locations with TOGA (v1.0) (*89*), with annotation completeness for both species assessed with BUSCO v5.2.2 (*84*).

Sor. araneus and *M. nigripes* TOGA annotations were then aligned to 38 mammalian TOGA annotations acquired from https://genome.senckenberg.de//download/TOGA/, including the stoat (*Mustela erminea*), the domesticated ferret (*Mustela putorius furo*), and the pygmy shrew (*Suncus etruscus*), all of which also undergo Dehnel-like growth patterns. Aligned species were chosen to reduce overrepresentation of species in orders or superorders (7 Eulipotyphlans, 4 Chiropterans, 6 Artiodactyls, 1 Perissodactyl, 1 Pholidota, 8 Carnivorans, 6 Glires, 6 Primates, 1 Scandentia), with genome annotations by TOGA requiring >16,000 annotated genes with BUSCOs >80% (Supplemental Data). The longest transcript for each gene was selected and aligned using the MACSE exon-by-exon alignment function of TOGA to produce multiple codon alignments and cleaned with TAPER (v1.0.0) default settings (*90*). Species were removed from each codon alignment prior to selection analyses when multiple copies of the gene were present or the alignment contained a frameshift, with codon alignments including at least 60% of the species (>24) retained for selection analyses.

Selection Models
To infer positive selection for lineages within the phylogeny, we used the adaptive branch-site random effects likelihood (aBSREL v2.2) model (*91*) implemented in HYPHY (Hypothesis Testing using Phylogenies) (v2.5.32) (*29*). First, tree pruning was conducted using ETE Toolkit python wrapper (v3) (*92*), where unrooted trees for each codon alignment were pruned from a larger Bayesian molecular-clock mammalian phylogeny (*93*). Multiple aBSREL tests were run. Branch-site models were used to infer positive selection in genes for *a priori* foreground lineage, including *S. araneus*, and those with Dehnel-like phenotypes (*M. erminea*, *M. p. furo, S.*

etruscus). Then, we conducted an exploratory analysis where all background branches were tested
for potential positive selection. This step identified genes evolving under positive selection
outside of the focal species to test for lineage specificity. To test if genes significantly rejected the
null hypothesis of no positive selection, we conducted a likelihood ratio test (LRT), with p-values
quantified using a chi-squared distribution (χ_1^2) and adjusted for multiple hypothesis tests across
genes and species with a Holm-Bonferroni corrections. Running both a priori and exploratory
analyses allows identifying genes under positive selection in *Sor. araneus* (regardless of
phylogeny-wide selection), genes under positive selection strictly in *Sor. araneus* (no other
branches with positive selection), and genes evolving in parallel related to Dehnel's phenomenon
(under positive selection in *S. araneus*, 2 other foreground lineages, <5 other background
lineages). To detect potential functional parallelism underlying seasonal size plasticity, we ran a
gene set enrichment analysis on our entire candidate gene set using the DAVID functional
annotation tool (*94*) with KEGG Gene Ontologies (GO) and Functional Annotations. Finally, we
used Mixed Effects Model of Evolution (MEME) (v2.1.2) (*95*) analyses to detect which codons
show signals of positive selection per gene using lineages with inferred positive selection. Here,
we used a higher threshold for significance (p<0.1), with a stricter Empirical Bayes Factor
threshold (EBF>100).

Chromosomal Evolution
Structural variant analysis of *Sor. araneus* and ancestral karyotype reconstruction of Eulipotyphla
were conducted with DESCHRAMBLER (v1) (*28*). Using the new chromosome-scale *Sor.*
araneus genome as a reference, we produced pairwise genome alignment chains and nets for five
Eulipotyphla species (*Condylura cristata*, *Galemys pyrenaicus*, *Suncus etruscus*, *Talpa*
occidentalis, Erinaceus europaeus) and two Chiropterans (*Rhinolophus ferrumequinum* and
Phyllostomus discolor; outgroups) with high quality assemblies (>5 million scaffold N50,
BUSCO > 95%). Chains and nets were made with lastz as described above. Conserved segments,
orthology blocks, and predicted ancestral chromosome structure were identified by
DESCHRAMBLER with a 1Mb resolution and visualized using syntenyPlotteR (*96*).

Transcriptomics
Transcriptomic data for the cortex and hippocampus was used to explore the regulatory
mechanisms of brain shrinkage and regrowth through Dehnel's phenomenon and identify further
overlap with PSGs. Shrews were sampled, brain regions dissected, and RNA extracted as
described in previous studies (*10*, *11*). Briefly, common shrews were collected (n=24) across five
seasons (n=4-5/season) of Dehnel's phenomenon (summer juveniles, autumn, winter, spring,
summer adults) from a single German population (47.9684N, 8.9761 E) using a protocol that
minimized trap-related stress and food insufficiency. Shrews were euthanized with vascular
perfusion of PAXgene Tissue Fixative, with brain regions dissected in cold fixative. The cortex
and hippocampus were incubated in PAXgene Tissue Stabilizer for 2-24 hours and snap frozen in
liquid nitrogen (-180°C). RNA was extracted from each region using a modified Qiagen Micro
RNAeasy protocol designed for small amounts of mammalian sensory tissue to reduce heat-
associated RNA degradation. Quality control (nanodrop and RNA ScreenTape), library
preparation (poly-A selection), and sequencing (approx. 15-25 million reads/sample, 150bp PE)
was conducted by Azenta Life Sciences.

For each sample, raw reads were trimmed for adapter sequences and pruned for low quality
sequences using fastp (*97*), and then aligned to the novel NCBI *Sor. araneus* transcriptome
(GCF_027595985.1_mSorAra2.pri_rna.fna) using Kallisto (v0.46.2) (*98*). Counts were
normalized using the DESeq2 (v1.36) (*99*) median of ratios which accounts for library size and
content. We then tested for differential expression in both regions between autumn and spring

individuals. These seasons were chosen as they resemble the shrinking and regrowth phases of Dehnel's phenomenon, and were previously analyzed in the hypothalamus (*100*). Genes were tested for differential expression ($p_{adj}<0.05$) using a Wald test in DESeq2 (*99*), with sex as an additional covariate (~sex + season), followed by Benjamini-Hochberg multiple test correction (*101*). We then ran a KEGG GO enrichment using fsgea (v1.22) (*102*) to describe potential molecular functions or pathways associated with identified differential expression. Lastly, we identified genes that overlapped with genes inferred to be evolving under positive selection.

Acknowledgments

We thank Marion Muturi for support with fieldwork.

Funding:

Human Frontiers of Science Program, award: RGP0013/2019 (DD, JN, LMD)

WRT was supported in part by a Stony Book University Presidential Innovation and Excellence award (LMD)

NSF-DBI 2010853 (TL)

DMS was supported, in part, by NSF-DEB 1838273 and 2032063 (LMD)

APC was supported, in part, by NSF-IOS 2031926 (APC)

Author contributions:

Conceptualization: WRT, LMD

Methodology: WRT, GF, OF, MF, DDMS, TML

Software: MF, TML, MF

Validation: WRT

Formal analysis: WRT

Investigation: WRT

Resources: TML

Data Curation: WRT, LA, JB, NJ, JM, TT, AA, CB

Writing – Original Draft: WRT

Writing – Review and Editing: LMD, DD, TML, DDMS, DE, JN, DR, GF, EJ

Visualization: WRT, GF

Supervision: LMD, GF, EJ

Project administration: LMD

Funding acquisition: LMD, APC, DD

Competing interests: All other authors declare they have no competing interests.

Data and materials availability: All data are available in the main text or the supplementary materials. Cleaned whole-genome alignments, cleaned gene alignments, and selection outputs openly available on Dryad https://doi.org/10.5061/dryad.8gtht770x. Supplementary tables, results, and code deposited and found on Github https://github.com/wrthomas315/Sorex_Genome2. Raw sequencing data located in the NCBI Sequencing Read Archive (BioProject PRJNA941271).

Figures and Tables

Figure 1 – Genome Assembly and Comparisons (A) Using PacBio and Hi-C chromosome interactions and manual curation, the shrew genome assembly was assembled into 12 chromosomes. **(B)** Cumulative percent of each Eulipotyphla genome found on contigs of certain length, showing a highly contiguous shrew genome assembly and longer chromosomes than those of other close relatives. **(C)** Phylogeny of species used in comparative genomic analyses, each with more than 16,000 genes and BUSCO scores greater than 80%. **(D)** Comparisons between *Sor. araneus* and *Sun. etruscus* orthologous blocks. Results highlight orthology between *Sor. araneus* chromosomes 10 and 11 and *Sun. etruscus* chromosomes 11 and 17, as well as fusions of the *Sun. etruscus* X, 5, and 12 and 6, 9, and 16 to form the *Sor. araneus* X chromosome and chromosome 5.

Figure 2 – Gene set enrichment of shrew-specific positively selected genes - (A) Five pathways were significantly enriched (p<0.05) with positively selected genes only under selection in the *Sor. araneus* lineage. The most enriched pathway was the Fanconi anemia complex, consisting of **(B)** *FANCI*, *FAAP100*, and *PALB2*, with >19 positively selected sites per gene detected from MEME. Genetic variability (selected amino acids, insertions, deletions) was found across genes, both within and between protein domains.

Figure 3 – Signatures of parallel evolution in Dehnel's phenomenon - (A) Upset plot of the positively selected genes (PSGs) found in *S. araneus*, species with Dehnel-like phenotypes, and background branches with two degrees of stringency (background selection >0 and background selection >4). Eight genes overlapped across species with Dehnel's-like traits were found. **(B)** The gene encoding protocadherin alpha 6, *PCDHA6*, was the single gene convergently evolving under positive selection in *S. araneus, M. erminea,* and *M. putorius furo* without background selection.

Figure 4 – Seasonal gene expression of hippocampus and cortex - (A) Volcano plot of differentially expressed genes in the cortex and the hippocampus between autumn (red, cortex: 46 upregulated genes, hippocampus: 25 upregulated genes) and spring (blue; cortex: 76 upregulated genes, hippocampus: 52 upregulated genes). **(B)** Gene set enrichment of differentially expressed genes identified 21 total enriched gene sets (p_{adj}<0.05), 5 of which overlap in the same direction between the two brain regions (Notch signaling, proteasome, oxidative phosphorylation, protein export, ribosome).

Figure 5 – Evolutionary and seasonal expression divergence in the brain associated with positive selection in coding sequence – **(A)** Upset plot of the positively selected genes (PSGs) found in *Sor. araneus* compared to significantly varying genes from transcriptomics analyses. **(B)** *SOX9*, involved in neural stem cell maintenance, is the only gene both downregulated in the hippocampus and cortex during brain shrinkage, and has positively selected changes in its coding sequence outside of a known domain. **(C)** Positive selection was also identified in *VEGFA*, involved in blood brain barrier development and function, and *SPHK2*, which regulates neuronal cell death. Both genes are part of a suite of 11 positively selected genes with upregulated expression in *Sor. araneus* hypothalamus compared to other mammals.

586   **Table 1** – Chromosome quality of new *Sorex araneus* genome assembly compared to Broad
587   Institute assembly.

	Broad Institute (sorAra2)	New assembly (mSorAra2.pri)
Total Sequence Length	2,423,158,183	2,405,993,548
Total Ungapped Length	2,192,103,426	2,393,049,584
Number of Scaffolds	12,845	65
Scaffold N50	22,794,405	374,364,821
Scaffold L50	36	3
Number of Contigs	201,420	1,492
Contig N50	22,623	4,024,507
Contig L50	27,135	182
Chromosomes	0	12
Source Location	Piel Island, United Kingdom	Radolfzell, Germany

588
589

590 **Table 2** – Counts of genes tested and found to be under positive selection by species. For all
591 counts statistical significance at p-val$_{adj} < 0.05$.
592

Species	Total Genes	Count including this tip	Species Specific	Convergent (Shrew+2)	Convergent (<5 Background)	Convergent (<1 Background)
Sor. araneus	15,812	676	231	23	8	1
S. etruscus	15,380	904	397	19	6	0
M. erminea	16,675	225	54	21	6	1
M. p. furo	16,638	110	40	7	4	1

593
594

Literature cited

1.   J. B. Searle, J. Zima, P. D. Polly, *Shrews, chromosomes and speciation* (2019).

2.   A. Dehnel, Studies of the genus Sorex L. *Ann. Univ. M. Curie-Sklod.* **4**, 17–104 (1949).

3.   Z. Pucek, Seasonal and age changes in the weight of internal organs of shrews. *Acta Theriol. (Warsz).* **10**,
369–438 (1965).

4.   J. Lázaro, M. Hertel, C. C. Sherwood, M. Muturi, D. K. N. Dechmann, Profound seasonal changes in brain
size and architecture in the common shrew. *Brain Struct. Funct.* **223**, 2823–2840 (2018).

5.   J. Lázaro, D. K. N. Dechmann, Dehnel's phenomenon. *Curr. Biol.* **31**, R463–R465 (2021).

6.   L. Nováková, J. Lázaro, M. Muturi, C. Dullin, D. K. N. Dechmann, Winter conditions , not resource
availability alone , may drive reversible seasonal skull size changes in moles. **9** (2022),
doi:https://doi.org/10.1098/rsos.220652.

7.   S. Ray, M. Li, S. P. Koch, S. Mueller, P. Boehm-Sturm, H. Wang, M. Brecht, R. K. Naumann, Seasonal
plasticity in the adult somatosensory cortex. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 32136–32144 (2020).

8.   S. Lapoint, L. Keicher, M. Wikelski, K. Zub, D. K. N. Dechmann, Growth overshoot and seasonal size
changes in the skulls of two weasel species. *R. Soc. Open Sci.* **4** (2017), doi:10.1098/rsos.160947.

9.   K. Healy, T. Guillerme, S. Finlay, A. Kane, S. B. A. Kelly, D. McClean, D. J. Kelly, I. Donohue, A. L.
Jackson, N. Cooper, Ecology and mode-of-life explain lifespan variation in birds and mammals. *Proc. R. Soc.*
B. **281** (2014), doi:10.1098/rspb.2014.0298.

10.   W. R. Thomas, T. Richter, C. Baldoni, A. P. Corthals, J. Nieland, D. K. N. Dechmann, R. G. Hunter, L. M.
Daávalos, Seasonal and comparative evidence of adaptive gene expression in mammalian brain size
plasticity. *Elife* (2024) (available at https://doi.org/10.7554/eLife.100788.1).

11.   W. R. Thomas, C. Baldoni, Y. Zeng, D. Carlson, J. Holm-Jacobsen, M. Muturi, D. von Elverfeldt, J. Nieland,
D. K. N. Dechmann, A. P. Corthals, L. M. Dávalos, Dynamic metabolic and molecular changes during
seasonal shrinking in Sorex araneus. *bioRxiv*, 1–24 (2024).

12.   B. Smith, thesis, Max Planck Institute for Ornithology (2016).

13.   M. Maceyka, H. Sankala, N. C. Hait, H. Le Stunff, H. Liu, R. Toman, C. Collier, M. Zhang, L. S. Satin, A. H.
Merrill, S. Milstien, S. Spiegel, SphK1 and SphK2, sphingosine kinase isoenzymes with opposing functions
in sphingolipid metabolism. *J. Biol. Chem.* **280**, 37118–37129 (2005).

14.   H. Liu, R. E. Toman, S. K. Goparaju, M. Maceyka, V. E. Nava, H. Sankala, S. G. Payner, M. Bektas, I. Ishii,
J. Chun, S. Milstien, S. Spiegel, Sphingosine kinase type 2 is a putative BH3-only protein that induces
apoptosis. *J. Biol. Chem.* **278**, 40330–40336 (2003).

15.   N. Takasugi, T. Sasaki, K. Suzuki, S. Osawa, H. Isshiki, Y. Hori, N. Shimada, T. Higo, S. Yokoshima, T.
Fukuyama, V. M. Y. Lee, J. Q. Trojanowski, T. Tomita, T. Iwatsubo, BACE1 activity is modulated by cell-
associated sphingosine-1-phosphate. *J. Neurosci.* **31**, 6850–6857 (2011).

16.   T. A. White, M. Bordewich, J. B. Searle, A network approach to study karyotypic evolution: The
chromosomal races of the common shrew (Sorex araneus) and house mouse (Mus musculus) as model
systems. *Syst. Biol.* **59**, 262–276 (2010).

17.   T. C. M. Benathar, C. Y. Nagamachi, L. R. R. Rodrigues, P. C. M. O'Brien, M. A. Ferguson-Smith, F. Yang,
J. C. Pieczarka, Karyotype, evolution and phylogenetic reconstruction in Micronycterinae bats with
implications for the ancestral karyotype of Phyllostomidae. *BMC Evol. Biol.* **19**, 1–13 (2019).

18.   R. Apfelbach, D. Kruska, Postnatal development of the brain of the ferret Mustela putorius F Furo
(Mustelidae, Mammalia). *Int. J. Mamm. Biol.* **44**, 127–131 (1979).

19.   D. Ochocińska, J. R. E. Taylor, Living at the physiological limits: Field and maximum metabolic rates of the
common shrew (Sorex araneus). *Physiol. Biochem. Zool.* **78**, 808–818 (2005).

20.   L. Keicher, M. T. O'Mara, C. C. Voigt, D. K. N. Dechmann, Stable carbon isotopes in breath reveal fast
metabolic incorporation rates and seasonally variable but rapid fat turnover in the common shrew (Sorex
araneus). *J. Exp. Biol.* **220**, 2834–2841 (2017).

21.   Z. Pucek, Seasonal and age change in shrews as an adaptive process. *Symp. zool. Soc. Lond.* **26**, 189–207
(1970).

22.   J. Lázaro, M. Hertel, M. Muturi, D. K. N. Dechmann, Seasonal reversible size changes in the braincase and
mass of common shrews are flexibly modified by environmental conditions. *Sci. Rep.* **9**, 1–10 (2019).

23.   S. Churchfield, L. Rychlik, J. R. E. Taylor, Food resources and foraging habits of the common shrew, Sorex
araneus: Does winter food shortage explain Dehnel's phenomenon? *Oikos.* **121**, 1593–1602 (2012).

24.   J. R. E. Taylor, L. Rychlik, S. Churchfield, Winter reduction in body mass in a very small, nonhibernating
mammal: Consequences for heat loss and metabolic rates. *Physiol. Biochem. Zool.* **86**, 9–18 (2013).

25.   H. Hyvärinen, Wintering strategy of voles and shrews in Findland. *Winter Ecol. Small Mamm.* **10**, 139–148
(1984).

26.   P. J. Schaeffer, M. T. O'Mara, J. Breiholz, L. Keicher, J. Lázaro, M. Muturi, D. K. N. Dechmann, Metabolic
rate in common shrews is unaffected by temperature, leading to lower energetic costs through seasonal size
reduction. *R. Soc. Open Sci.* **7** (2020), doi:10.1098/rsos.191989.

27.   A. Rhie, S. A. McCarthy, O. Fedrigo, J. Damas, G. Formenti, S. Koren, M. Uliano-Silva, W. Chow, A.

Fungtammasan, J. Kim, C. Lee, B. J. Ko, M. Chaisson, G. L. Gedman, L. J. Cantin, F. Thibaud-Nissen, L. Haggerty, I. Bista, M. Smith, B. Haase, J. Mountcastle, S. Winkler, S. Paez, J. Howard, S. C. Vernes, T. M. Lama, F. Grutzner, W. C. Warren, C. N. Balakrishnan, D. Burt, J. M. George, M. T. Biegler, D. Iorns, A. Digby, D. Eason, B. Robertson, T. Edwards, M. Wilkinson, G. Turner, A. Meyer, A. F. Kautt, P. Franchini, H. W. Detrich, H. Svardal, M. Wagner, G. J. P. Naylor, M. Pippel, M. Malinsky, M. Mooney, M. Simbirsky, B. T. Hannigan, T. Pesout, M. Houck, A. Misuraca, S. B. Kingan, R. Hall, Z. Kronenberg, I. Sović, C. Dunn, Z. Ning, A. Hastie, J. Lee, S. Selvaraj, R. E. Green, N. H. Putnam, I. Gut, J. Ghurye, E. Garrison, Y. Sims, J. Collins, S. Pelan, J. Torrance, A. Tracey, J. Wood, R. E. Dagnew, D. Guan, S. E. London, D. F. Clayton, C. V. Mello, S. R. Friedrich, P. V. Lovell, E. Osipova, F. O. Al-Ajli, S. Secomandi, H. Kim, C. Theofanopoulou, M. Hiller, Y. Zhou, R. S. Harris, K. D. Makova, P. Medvedev, J. Hoffman, P. Masterson, K. Clark, F. Martin, K. Howe, P. Flicek, B. P. Walenz, W. Kwak, H. Clawson, M. Diekhans, L. Nassar, B. Paten, R. H. S. Kraus, A. J. Crawford, M. T. P. Gilbert, G. Zhang, B. Venkatesh, R. W. Murphy, K. P. Koepfli, B. Shapiro, W. E. Johnson, F. Di Palma, T. Marques-Bonet, E. C. Teeling, T. Warnow, J. M. Graves, O. A. Ryder, D. Haussler, S. J. O'Brien, J. Korlach, H. A. Lewin, K. Howe, E. W. Myers, R. Durbin, A. M. Phillippy, E. D. Jarvis, Towards complete and error-free genome assemblies of all vertebrate species. *Nature*. **592**, 737–746 (2021).

28. J. Kim, M. Farré, L. Auvil, B. Capitanu, D. M. Larkin, J. Ma, H. A. Lewin, Reconstruction and evolutionary history of eutherian chromosomes. *Proc. Natl. Acad. Sci. U. S. A.* **114**, E5379–E5388 (2017).

29. S. L. Kosakovsky Pond, A. F. Y. Poon, R. Velazquez, S. Weaver, N. L. Hepler, B. Murrell, S. D. Shank, B. R. Magalis, D. Bouvier, A. Nekrutenko, S. Wisotsky, S. J. Spielman, S. D. W. Frost, S. V. Muse, HyPhy 2.5 - A Customizable Platform for Evolutionary Hypothesis Testing Using Phylogenies. *Mol. Biol. Evol.* **37**, 295–299 (2020).

30. S. Dubey, N. Salamin, S. D. Ohdachi, P. Barrière, P. Vogel, Molecular phylogenetics of shrews (Mammalia: Soricidae) reveal timing of transcontinental colonizations. *Mol. Phylogenet. Evol.* **44**, 126–137 (2007).

31. J. B. Searle, J. Zima, P. D. Polly, *Shrews, Chromosomes and Speciation* (Cambridge University Press, Cambridge, 2019).

32. A. Karlsson, D. Deb-Basu, A. Cherry, S. Turner, J. Ford, D. W. Felsher, Defective double-strand DNA break repair and chromosomal translocations by MYC overexpression. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 9974–9979 (2003).

33. I. Mayrose, M. A. Lysak, The Evolution of Chromosome Numbers: Mechanistic Models and Experimental Approaches. *Genome Biol. Evol.* **13**, 1–15 (2021).

34. T. A. Titus, D. R. Selvig, B. Qin, C. Wilson, A. M. Starks, B. A. Roe, J. H. Postlethwait, The Fanconi anemia gene network is conserved from zebrafish to human. *Gene*. **371**, 211–223 (2006).

35. P. Alcón, S. Shakeel, Z. A. Chen, J. Rappsilber, K. J. Patel, L. A. Passmore, FANCD2–FANCI is a clamp stabilized on DNA by monoubiquitination of FANCD2 during DNA repair. *Nat. Struct. Mol. Biol.* **27**, 240–248 (2020).

36. A. Smogorzewska, S. Matsuoka, P. Vinciguerra, E. R. McDonald, K. E. Hurov, J. Luo, B. A. Ballif, S. P. Gygi, K. Hofmann, A. D. D'Andrea, S. J. Elledge, Identification of the FANCI Protein, a Monoubiquitinated FANCD2 Paralog Required for DNA Repair. *Cell*. **129**, 289–301 (2007).

37. P. Alcón, A. P. Kaczmarczyk, K. K. Ray, T. Liolios, G. Guilbaud, T. Sijacki, Y. Shen, S. H. McLaughlin, J. E. Sale, P. Knipscheer, D. S. Rueda, L. A. Passmore, FANCD2–FANCI surveys DNA and recognizes double- to single-stranded junctions. *Nature*. **632**, 1165–1173 (2024).

38. C. Ling, M. Ishiai, A. M. Ali, A. L. Medhurst, K. Neveling, R. Kalb, Z. Yan, Y. Xue, A. B. Oostra, A. D. Auerbach, M. E. Hoatlin, D. Schindler, H. Joenje, J. P. De Winter, M. Takata, A. R. Meetei, W. Wang, FAAP100 is essential for activation of the Fanconi anemia-associated DNA damage response pathway. *EMBO J.* **26**, 2104–2114 (2007).

39. Y. Guo, W. Feng, S. M. H. Sy, M. S. Y. Huen, ATM-dependent phosphorylation of the fanconi anemia protein PALB2 promotes the DNA damage response. *J. Biol. Chem.* **290**, 27545–27556 (2015).

40. B. Xia, Q. Sheng, K. Nakanishi, A. Ohashi, J. Wu, N. Christ, X. Liu, M. Jasin, F. J. Couch, D. M. Livingston, Control of BRCA2 Cellular and Clinical Functions by a Nuclear Partner, PALB2. *Mol. Cell*. **22**, 719–729 (2006).

41. M. Yousefzadeh, C. Henpita, R. Vyas, C. Soto-Palma, P. Robbins, L. Niedernhofer, Dna damage—how and why we age? *Elife*. **10**, 1–17 (2021).

42. S. L. MacRae, M. M. K. Croken, R. B. Calder, A. Aliper, B. Milholland, R. R. White, A. Zhavoronkov, V. N. Gladyshev, A. Seluanov, V. Gorbunova, Z. D. Zhang, J. Vijg, DNA repair in species with extreme lifespan differences. *Aging (Albany. NY)*. **7**, 1171–1184 (2015).

43. B. Debrabant, M. Soerensen, F. Flachsbart, S. Dato, J. Mengel-From, T. Stevnsner, V. A. Bohr, T. A. Kruse, S. Schreiber, A. Nebel, K. Christensen, Q. Tan, L. Christiansen, Human longevity and variation in DNA damage response and repair: Study of the contribution of sub-processes using competitive gene-set analysis. *Eur. J. Hum. Genet.* **22**, 1131–1136 (2014).

44. K. Christensen, T. E. Johnson, J. W. Vaupel, The quest for genetic determinants of human longevity: Challenges and insights. *Nat. Rev. Genet.* **7**, 436–448 (2006).

45. R. M. J. Brosh, M. Bellani, Y. Liu, M. M. Seidman, Fanconi Anemia: a DNA Repair Disorder Characterized
by Accelerated Decline of the Hematopoietic Stem Cell Compartment and Other Features of Aging. *Ageing*
Res Rev. **33**, 67–75 (2017).
46. B. Xia, J. C. Dorsman, N. Ameziane, Y. De Vries, M. A. Rooimans, Q. Sheng, G. Pals, A. Errami, E.
Gluckman, J. Llera, W. Wang, D. M. Livingston, H. Joenje, J. P. De Winter, Fanconi anemia is associated
with a defect in the BRCA2 partner PALB2. *Nat. Genet.* **39**, 159–161 (2007).
47. S. Reid, D. Schindler, H. Hanenberg, K. Barker, S. Hanks, R. Kalb, K. Neveling, P. Kelly, S. Seal, M.
Freund, M. Wurm, S. D. Batish, F. P. Lach, S. Yetgin, H. Neitzel, H. Ariffin, M. Tischkowitz, C. G. Mathew,
A. D. Auerbach, N. Rahman, Biallelic mutations in PALB2 cause Fanconi anemia subtype FA-N and
predispose to childhood cancer. *Nat. Genet.* **39**, 162–164 (2007).
48. M. Wirthlin, N. C. B. Lima, R. L. M. Guedes, A. E. R. Soares, L. G. P. Almeida, N. P. Cavaleiro, G. Loss de
Morais, A. V. Chaves, J. T. Howard, M. de M. Teixeira, P. N. Schneider, F. R. Santos, M. C. Schatz, M. S.
Felipe, C. Y. Miyaki, A. Aleixo, M. P. C. Schneider, E. D. Jarvis, A. T. R. Vasconcelos, F. Prosdocimi, C. V.
Mello, Parrot Genomes and the Evolution of Heightened Longevity and Cognition. *Curr. Biol.* **28**, 4001-
4008.e7 (2018).
49. T. Yagi, M. Takeichi, Cadherin superfamily genes: Functions, genomic organization, and neurologic
diversity. *Genes Dev.* **14**, 1169–1180 (2000).
50. N. Fukuda, T. Fukuda, P. Percipalle, K. Oda, N. Takei, K. Czaplinski, K. Touhara, Y. Yoshihara, T. Sasaoka,
Axonal mRNA binding of hnRNP A/B is crucial for axon targeting and maturation of olfactory sensory
neurons. *Cell Rep.* **42**, 112398 (2023).
51. W. V. Chen, C. L. Nwakeze, C. A. Denny, S. O'Keeffe, M. A. Rieger, G. Mountoufaris, A. Kirner, J. D.
Dougherty, R. Hen, Q. Wu, T. Maniatis, Pcdhαc2 is required for axonal tiling and assembly of serotonergic
circuitries in mice. *Science (80-.).* **356**, 406–411 (2017).
52. K. C. Vadodaria, Y. Ji, M. Skime, A. C. Paquola, T. Nelson, D. Hall-Flavin, K. J. Heard, C. Fredlender, Y.
Deng, J. Elkins, K. Dani, A. T. Le, M. C. Marchetto, R. Weinshilboum, F. H. Gage, Altered serotonergic
circuitry in SSRI-resistant major depressive disorder patient-derived neurons. *Mol. Psychiatry*. **24**, 808–818
(2019).
53. P. Khaitovich, I. Hellmann, W. Enard, K. Nowick, M. Leinweber, H. Franz, G. Weiss, M. Lachmann, S.
Paabo, Parallel Patterns of Evolution in the Genomes and Transcriptomes of Humans and Chimpanzees.
Science (80-.). **309**, 1850–1854 (2005).
54. B. Lemos, C. D. . Meiklejohn, M. Cáceres, D. L. . Hartl, Rates of Divergence in Gene Expression Profiles of
Primates , Mice , and Flies : Stabilizing Selection and Variability among Functional Categories. *Evolution (N.*
Y). **59**, 126–137 (2005).
55. B. Lemos, B. R. Bettencourt, C. D. Meiklejohn, D. L. Hartl, Evolution of proteins and gene expression levels
are coupled in Drosophila and are independently associated with mRNA abundance, protein length, and
number of protein-protein interactions. *Mol. Biol. Evol.* **22**, 1345–1354 (2005).
56. S. V. Nuzhdin, M. L. Wayne, K. L. Harmon, L. M. McIntyre, Common pattern of evolution of gene
expression level and protein sequence in Drosophila. *Mol. Biol. Evol.* **21**, 1308–1317 (2004).
57. R. Dean, P. W. Harrison, A. E. Wright, F. Zimmer, J. E. Mank, Positive selection underlies Faster-Z
evolution of gene expression in birds. *Mol. Biol. Evol.* **32**, 2646–2656 (2015).
58. P. Khaitovich, W. Enard, M. Lachmann, S. Pääbo, Evolution of primate gene expression. *Nat. Rev. Genet.* **7**,
693–702 (2006).
59. M. Sepp, K. Leiss, F. Murat, K. Okonechnikov, P. Joshi, E. Leushkin, L. Spänig, N. Mbengue, C. Schneider,
J. Schmidt, N. Trost, M. Schauer, P. Khaitovich, S. Lisgo, M. Palkovits, P. Giere, L. M. Kutscher, S. Anders,
M. Cardoso-Moreira, I. Sarropoulos, S. M. Pfister, H. Kaessmann, Cellular development and evolution of the
mammalian cerebellum. *Nature*. **625**, 788–796 (2024).
60. M. Cardoso-Moreira, J. Halbert, D. Valloton, B. Velten, C. Chen, Y. Shao, A. Liechti, K. Ascenção, C.
Rummel, S. Ovchinnikova, P. V. Mazin, I. Xenarios, K. Harshman, M. Mort, D. N. Cooper, C. Sandi, M. J.
Soares, P. G. Ferreira, S. Afonso, M. Carneiro, J. M. A. Turner, J. L. VandeBerg, A. Fallahshahroudi, P.
Jensen, R. Behr, S. Lisgo, S. Lindsay, P. Khaitovich, W. Huber, J. Baker, S. Anders, Y. E. Zhang, H.
Kaessmann, Gene expression across mammalian organ development. *Nature*. **571**, 505–509 (2019).
61. S. Artavanis-Tsakonas, M. D. Rand, R. J. Lake, Notch signaling: Cell fate control and signal integration in
development. *Science (80-.).* **284**, 770–776 (1999).
62. K. Yoon, N. Gaiano, Notch signaling in the mammalian central nervous system: Insights from mouse
mutants. *Nat. Neurosci.* **8**, 709–715 (2005).
63. J. L. Ables, J. J. Breunig, A. J. Eisch, P. Rakic, Not(ch) just development: Notch signalling in the adult brain.
Nat. Rev. Neurosci. **12**, 269–283 (2011).
64. P. Chapouton, P. Skupien, B. Hesl, M. Coolen, J. C. Moore, R. Madelaine, E. Kremmer, T. Faus-Kessler, P.
Blader, N. D. Lawson, L. Bally-Cuif, Notch activity levels control the balance between quiescence and
recruitment of adult neural stem cells. *J. Neurosci.* **30**, 7961–7974 (2010).
65. K. Hashimoto-Torii, M. Torii, M. R. Sarkisian, C. M. Bartley, J. Shen, F. Radtke, T. Gridley, N. Šestan, P.

778   Rakic, Interaction between Reelin and Notch Signaling Regulates Neuronal Migration in the Cerebral Cortex.
779   *Neuron*. **60**, 273–284 (2008).

66.   K. I. Vong, C. K. Y. Leung, R. R. Behringer, K. M. Kwan, Sox9 is critical for suppression of neurogenesis
but not initiation of gliogenesis in the cerebellum. *Mol. Brain*. **8**, 0–17 (2015).

67.   C. E. Scott, S. L. Wynn, A. Sesay, C. Cruz, M. Cheung, M. V. G. Gaviro, S. Booth, B. Gao, K. S. E. Cheah,
R. Lovell-Badge, J. Briscoe, SOX9 induces and maintains neural stem cells. *Nat. Neurosci.* **13**, 1181–1189
(2010).

68.   L. C. Cheng, E. Pastrana, M. Tavazoie, F. Doetsch, MiR-124 regulates adult neurogenesis in the
subventricular zone stem cell niche. *Nat. Neurosci.* **12**, 399–408 (2009).

69.   J. Pritchett, V. Athwal, N. Roberts, N. A. Hanley, K. P. Hanley, Understanding the role of SOX9 in acquired
diseases: Lessons from development. *Trends Mol. Med.* **17**, 166–174 (2011).

70.   C. Kwok, P. A. Weller, S. Guioli, J. W. Foster, S. Mansour, O. Zuffardi, H. H. Punnett, M. A. Dominguez-
Steglich, J. David Brook, L. D. Young, P. N. Goodfellow, A. J. Schafer, Mutations in SOX9, the gene
responsible for campomelic dysplasia and autosomal sex reversal. *Am. J. Hum. Genet.* **57**, 1028–1036 (1995).

71.   F. Csukasi, I. Duran, W. Zhang, J. H. Martin, M. Barad, M. Bamshad, M. A. Weis, D. Eyre, D. Krakow, D.
H. Cohn, Dominant-negative SOX9 mutations in campomelic dysplasia. *Hum. Mutat.* **40**, 2344–2352 (2019).

72.   T. Arcondéguy, E. Lacazette, S. Millevoi, H. Prats, C. Touriol, VEGF-A mRNA processing, stability and
translation: A paradigm for intricate regulation of gene expression at the post-transcriptional level. *Nucleic*
Acids Res. **41**, 7997–8010 (2013).

73.   B. Küsters, R. M. W. De Waal, P. Wesseling, K. Verrijp, C. Maass, A. Heerschap, J. O. Barentsz, F. Sweep,
D. J. Ruiter, W. P. J. Leenders, Differential effects of vascular endothelial growth factor A isoforms in a
mouse brain metastasis model of human melanoma. *Cancer Res.* **63**, 5408–5413 (2003).

74.   L. Duret, F. Dorkeld, C. Gautier, Strong conservation of non-coding sequences during vertebrates evolution:
Potential involvement in post-transcriptional regulation of gene expression. *Nucleic Acids Res.* **21**, 2315–
2322 (1993).

75.   X. L. Zhuang, J. J. Zhang, Y. Shao, Y. Ye, C. Y. Chen, L. Zhou, Z. B. Wang, X. Luo, B. Su, Y. G. Yao, D.
N. Cooper, B. X. Hu, L. Wang, X. G. Qi, J. Lin, G. J. Zhang, W. Wang, N. Sheng, D. D. Wu, Integrative
Omics Reveals Rapidly Evolving Regulatory Sequences Driving Primate Brain Evolution. *Mol. Biol. Evol.*
40, 1–15 (2023).

76.   D. Nakayama, T. Makino, Convergent accelerated evolution of mammal-specific conserved non-coding
elements in hibernators. *Sci. Rep.* **14**, 1–9 (2024).

77.   G. W. Vurture, F. J. Sedlazeck, M. Nattestad, C. J. Underwood, H. Fang, J. Gurtowski, M. C. Schatz,
GenomeScope: Fast reference-free genome profiling from short reads. *Bioinformatics*. **33**, 2202–2204 (2017).

78.   H. Cheng, G. T. Concepcion, X. Feng, H. Zhang, H. Li, Haplotype-resolved de novo assembly using phased
assembly graphs with hifiasm. *Nat. Methods*. **18**, 170–175 (2021).

79.   D. Guan, D. Guan, S. A. McCarthy, J. Wood, K. Howe, Y. Wang, R. Durbin, R. Durbin, Identifying and
removing haplotypic duplication in primary genome assemblies. *Bioinformatics*. **36**, 2896–2898 (2020).

80.   J. Ghurye, A. Rhie, B. P. Walenz, A. Schmitt, S. Selvaraj, M. Pop, A. M. Phillippy, S. Koren, Integrating Hi-
C links with assembly graphs for chromosome-scale assembly. *PLoS Comput. Biol.* **15**, 1–19 (2019).

81.   W. Chow, K. Brugger, M. Caccamo, I. Sealy, J. Torrance, K. Howe, GEVAL - A web-based browser for
evaluating genome assemblies. *Bioinformatics*. **32**, 2508–2510 (2016).

82.   A. Rhie, B. P. Walenz, S. Koren, A. M. Phillippy, Merqury: Reference-free quality, completeness, and
phasing assessment for genome assemblies. *Genome Biol.* **21**, 1–27 (2020).

83.   A. Gurevich, V. Saveliev, N. Vyahhi, G. Tesler, QUAST: Quality assessment tool for genome assemblies.
Bioinformatics. **29**, 1072–1075 (2013).

84.   R. M. Waterhouse, M. Seppey, F. A. Simao, M. Manni, P. Ioannidis, G. Klioutchnikov, E. V. Kriventseva, E.
M. Zdobnov, BUSCO applications from quality assessments to gene prediction and phylogenomics. *Mol.*
Biol. Evol. **35**, 543–548 (2018).

85.   A. Smit, R. M. Hubley, thesis.

86.   A. Smit, R. M. Hubley, P. Green, RepeatMasker Open-4.0. *<http://www.repeatmasker.org>*.

87.   E. Osipova, N. Hecker, M. Hiller, RepeatFiller newly identifies megabases of aligning repetitive sequences
and improves annotations of conserved non-exonic elements. *Gigascience*. **8**, 1–10 (2019).

88.   H. G. Suarez, B. E. Langer, P. Ladde, M. Hiller, ChainCleaner improves genome alignment specificity and
sensitivity. *Bioinformatics*. **33**, 1596–1603 (2017).

89.   B. M. Kirilenko, M. Šálek, E. Osipova, D. Jebb, V. Sharma, M. Blumer, A. E. Morales, A.-W. Ahmed, D.-G.
Kontopoulos, L. Hilgers, K. Lindblad-Toh, E. K. Karlsson, M. Hiller, G. Andrews, J. C. Armstrong, M.
Bianchi, B. W. Birren, K. R. Bredemeyer, A. M. Breit, M. J. Christmas, H. Clawson, J. Damas, F. Di Palma,
M. Diekhans, M. X. Dong, E. Eizirik, K. Fan, C. Fanter, N. M. Foley, K. Forsberg-Nilsson, C. J. Garcia, J.
Gatesy, S. Gazal, D. P. Genereux, L. Goodman, J. Grimshaw, M. K. Halsey, A. J. Harris, G. Hickey, M.
Hiller, A. G. Hindle, R. M. Hubley, G. M. Hughes, J. Johnson, D. Juan, I. M. Kaplow, E. K. Karlsson, K. C.
Keough, B. Kirilenko, K.-P. Koepfli, J. M. Korstian, A. Kowalczyk, S. V. Kozyrev, A. J. Lawler, C. Lawless,

T. Lehmann, D. L. Levesque, H. A. Lewin, X. Li, A. Lind, K. Lindblad-Toh, A. Mackay-Smith, V. D.
Marinescu, T. Marques-Bonet, V. C. Mason, J. R. S. Meadows, W. K. Meyer, J. E. Moore, L. R. Moreira, D.
D. Moreno-Santillan, K. M. Morrill, G. Muntané, W. J. Murphy, A. Navarro, M. Nweeia, S. Ortmann, A.
Osmanski, B. Paten, N. S. Paulat, A. R. Pfenning, B. N. Phan, K. S. Pollard, H. E. Pratt, D. A. Ray, S. K.
Reilly, J. R. Rosen, I. Ruf, L. Ryan, O. A. Ryder, P. C. Sabeti, D. E. Schäffer, A. Serres, B. Shapiro, A. F. A.
Smit, M. Springer, C. Srinivasan, C. Steiner, J. M. Storer, K. A. M. Sullivan, P. F. Sullivan, E. Sundström, M.
A. Supple, R. Swofford, J.-E. Talbot, E. Teeling, J. Turner-Maier, A. Valenzuela, F. Wagner, O. Wallerman,
C. Wang, J. Wang, Z. Weng, A. P. Wilder, M. E. Wirthlin, J. R. Xue, X. Zhang, Integrating gene annotation
with orthology inference at scale. *Science (80-.).* **380**, 1–17 (2023).
90.   C. Zhang, Y. Zhao, E. L. Braun, S. Mirarab, TAPER: Pinpointing errors in multiple sequence alignments
despite varying rates of evolution. *Methods Ecol. Evol.* **12**, 2145–2158 (2021).
91.   M. D. Smith, J. O. Wertheim, S. Weaver, B. Murrell, K. Scheffler, S. L. Kosakovsky Pond, Less is more: An
adaptive branch-site random effects model for efficient detection of episodic diversifying selection. *Mol.*
Biol. Evol. **32**, 1342–1353 (2015).
92.   J. Huerta-Cepas, F. Serra, P. Bork, ETE 3: Reconstruction, Analysis, and Visualization of Phylogenomic
Data. *Mol. Biol. Evol.* **33**, 1635–1638 (2016).
93.   S. Álvarez-Carretero, A. U. Tamuri, M. Battini, F. F. Nascimento, E. Carlisle, R. J. Asher, Z. Yang, P. C. J.
Donoghue, M. dos Reis, A species-level timeline of mammal evolution integrating phylogenomic data.
Nature. **602**, 263–267 (2022).
94.   D. W. Huang, B. T. Sherman, R. A. Lempicki, Systematic and integrative analysis of large gene lists using
DAVID bioinformatics resources. *Nat. Protoc.* **4**, 44–57 (2009).
95.   B. Murrell, J. O. Wertheim, S. Moola, T. Weighill, K. Scheffler, S. L. Kosakovsky Pond, Detecting
individual sites subject to episodic diversifying selection. *PLoS Genet.* **8** (2012),
doi:10.1371/journal.pgen.1002764.
96.   S. Quigley, J. Damas, D. M. Larkin, M. Farré, syntenyPlotteR: a user-friendly R package to visualize genome
synteny, ideal for both experienced and novice bioinformaticians. *Bioinforma. Adv.* **3**, 2–4 (2023).
97.   S. Chen, Y. Zhou, Y. Chen, J. Gu, Fastp: An ultra-fast all-in-one FASTQ preprocessor. *Bioinformatics.* **34**,
i884–i890 (2018).
98.   N. L. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal probabilistic RNA-seq quantification. *Nat.*
Biotechnol. **34**, 525–527 (2016).
99.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. *Genome Biol.* **15**, 1–21 (2014).
100.   W. R. Thomas, T. Richter, C. Baldoni, A. P. Corthals, J. Nieland, D. K. N. Dechmann, R. G. Hunter, L. M.
Daávalos, Seasonal and comparative evidence of adaptive gene expression in mammalian brain size
plasticity. *bioRxiv* (2024), doi:https://doi.org/10.1101/2024.06.18.599531.
101.   Y. Benjamini, Y. Hochberg, Controlling the False Discovery Rate : A Practical and Powerful Approach to
Multiple Testing. *J. R. Stat. Soc.* **57**, 289–300 (1995).
102.   G. Korotkevich, V. Sukhov, N. Budin, M. N. Atryomov, A. Sergushichev, Fast gene set enrichment analysis.
bioRxiv. *bioRxiv* (2021).
878
879

A

Interaction Size

427, 225, 204, 186, 87, 51, 15, 10, 8, 4, 4, 3, 3, 2, 2, 2, 1, 1, 1, 1, 1, 1

Legend
- ☀ Summer
- 🍂 Autumn
- ❄ Winter
- ❀ Spring
- 🔴 Selected Sites

Set Size

- Seasonal Hippocampus
- Seasonal Cortex
- Seasonal Hypothalamus
- Comparative Hypothalamus
- *S. araneus* specific PSGs
- *S. araneus* PSGs

600 300 0

B

SOX9 Expression

- ● Cortex
- ■ Hippocampus

SOX9

| HMG-box | No Domain |

AA1   AA200   AA400

C

VEGFA   SPHK2   HYPOTHALAMUS EXPRESSION

- *Rattus novegicus*
- *Mus musculus*
- *Peromyscus sp.*
- *Sciuridae sp.*
- *Pan troglodytes*
- *Homo sapiens*
- *Macaca mulatta*
- *Bos sp.*
- *Ovis orientalis*
- *Sus scrofa*
- *Sorex araneus*

zscore: 3, 2, 1, 0, -1

| Selected Site | Foreground CDS | Background CDS | Deletion/Masked |

0 200 400   0   500   1000

BranchLength 0.00 0.15 0.3

Amino Acid

VEGFA SPHK2 MLST8 GRIP2 FANCI KIAA0930 LOXL1 PIGR GALNT12 POLDIP3 NXPH3