# Enhanced Tactile Coding in Rat Neocortex Under Darkness

Kotaro Yamashiro[1], Shiyori Tanaka[1], Nobuyoshi Matsumoto[1,2†], Yuji Ikegaya[1,2,3†]

[1]Graduate School of Pharmaceutical Sciences, The University of Tokyo, Tokyo, 113-0033, Japan
[2]Institute for AI and Beyond, The University of Tokyo, Tokyo, 113-0033, Japan
[3]Center for Information and Neural Networks, National Institute of Information and Communications Technology, Suita City, Osaka, 565-0871, Japan

**Abbreviated title:** Enhanced Tactile Coding Under Darkness

†To whom correspondence should be addressed

Yuji Ikegaya Ph.D.
Laboratory of Chemical Pharmacology
Graduate School of Pharmaceutical Sciences, The University of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
Tel. : +81-3-5841-4780; Fax: +81-3-5841-4786
E-mail: yuji@ikegaya.jp

Nobuyoshi Matsumoto Ph.D.
Laboratory of Chemical Pharmacology
Graduate School of Pharmaceutical Sciences, The University of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
Tel. : +81-3-5841-4780; Fax: +81-3-5841-4786
E-mail: nobuyoshi@matsumoto.ac

**Conflicts of interest:** The authors declare no competing interests.

**Acknowledgments:** This work was supported by JST ERATO (JPMJER1801), AMED-CREST (24wm0625401h0001; 24wm0625502s0501; 24wm0625207s0101; 24gm1510002s0104), the Institute for AI and Beyond of the University of Tokyo, JSPS Grants-in-Aid for Scientific Research (18H05525, 20K15926, 22K21353), KOSÉ Cosmetology Research Foundation, the Public Foundation of Chubu Science and Technology Center, and Konica Minolta Science and Technology Foundation.

**Abstract**

Sensory systems are known for their adaptability, responding dynamically to changes in environmental conditions. A key example of this adaptability is the enhancement of tactile perception in the absence of visual input. Despite behavioral studies showing visual deprivation can improve tactile discrimination, the underlying neural mechanisms, particularly how tactile neural representations are reorganized during visual deprivation, remain unclear. In this study, we explore how the absence of visual input alters tactile neural encoding in the rat somatosensory cortex (S1). Rats were trained on a custom-designed treadmill with distinct tactile textures (rough and smooth), and local field potentials (LFPs) were recorded from S1 under light and dark conditions. Machine learning techniques, specifically a convolutional neural network, were used to decode the high-dimensional LFP signals. We found that the neural representations of tactile stimuli became more distinct in the dark, indicating a reorganization of sensory processing in S1 when visual input was removed. Notably, conventional amplitude-based analyses failed to capture these changes, highlighting the power of machine learning in uncovering subtle neural patterns. These findings offer new insights into how the brain rapidly adapts tactile processing in response to the loss of visual input, with implications for multisensory integration and potential strategies for sensory rehabilitation.

## Introduction

The primary somatosensory cortex (S1) is integral to the encoding of tactile information, processing sensory inputs from various regions of the body (Delhaye et al., 2018; Serino, 2019; Di Plinio et al., 2020; Piras et al., 2020). This cortical area is crucial for our sense of touch, facilitating the perception and interpretation of sensations such as pressure, vibration, temperature, and pain (Bushnell et al., 1999; Luna et al., 2005; Moulton et al., 2012). These external stimuli are represented as neural activity in S1, enabling animals to differentiate among distinct sensory experiences (Koch and Fuster, 1989; Salinas et al., 2000; Goodwin and Wheat, 2004; Bensmaia et al., 2008). However, the role of S1 extends beyond the mere reception of tactile signals. The neural representations within S1 are characterized by dynamic and flexible properties, shaped not only by feedforward mechanisms but also by factors such as attention, motivation, and cross-modal influences (Driver and Spence, 2000; Eimer and Forster, 2003; Schürmann et al., 2004; Butler et al., 2012; Ziegler et al., 2023). Such plasticity allows for adaptive responses to an ever-changing environment, which is critical for survival (Dijkerman and de Haan, 2007; Abraira and Ginty, 2013). A key manifestation of this adaptability is cross-modal processing, wherein inputs from other sensory modalities, such as visual or auditory information, can modulate tactile perception (Hopkins et al., 2017; Nikbakht et al., 2018; Sugiyama et al., 2019; Bulusu and Lazar, 2024). These findings suggest that S1 is not exclusively dedicated to tactile processing, but instead integrates and adapts sensory information from multiple modalities.

An example of cross-modal processing between S1 and other sensory modalities is the enhancement of tactile abilities in the absence or reduction of visual input. Blind individuals, particularly those who have been blind from an early age, consistently outperform sighted individuals on tasks requiring tactile discrimination, such as two-point discrimination and texture identification (Van Boven et al., 2000; Goldreich and Kanics, 2003; Norman and Bartholomew, 2011; Wong et al., 2011). This phenomenon is caused by repurposing of cortical areas normally devoted to vision or other senses to

enhance processing of touch and other modalities (Rauschecker et al., 1992; Pascual-
Leone and Torres, 1993; Sadato et al., 1996; Burton, 2003; Sathian and Stilla, 2010).
Research indicates that during critical developmental periods, the visual cortex
undergoes functional reorganization, and regions ordinarily involved in visual
processing are repurposed to process stimuli from other sensory modalities, including
tactile input (Sadato et al., 1996; Burton, 2003; Karlen et al., 2006). This reorganization
is not immediate but rather occurs over extended periods (Karlen et al., 2006; Piché et
al., 2007). This capacity for tactile enhancement results from both structural and
functional plasticity, underscoring the brain's remarkable ability to adapt and optimize
sensory processing when one modality is lost or altered.

Interestingly, even sighted individuals can experience improvements in tactile
performance when visual cues are removed or minimized, as evidenced by increased
accuracy in tasks such as Braille reading under low-visibility conditions (Pascual-Leone
and Hamilton, 2001; Kauffman et al., 2002; Facchini and Aglioti, 2003). These findings
suggest that limiting or removing vision may trigger compensatory mechanisms that
enhance the sense of touch, even in those with typical vision (Boroojerdi et al., 2000;
Merabet et al., 2008; Bola et al., 2017). While much of the research on blindness
focuses on the extended plasticity that occurs in response to sensory deprivation—such
as the reorganization of somatosensory maps and the recruitment of visual regions for
tactile processing—these changes typically unfold over long durations or during critical
developmental periods. Thus, although the brain's capacity to reorganize tactile
representation in the absence of visual input is observed in both blind and sighted
individuals, the underlying mechanisms behind long-term and short-term cross-modal
processing are likely distinct. In blind individuals, these processes may involve more
profound, lasting changes in neural circuitry, whereas in sighted individuals, the
adaptations may be more transient and context-dependent. This suggests that different
neural processes drive the change in somatosensory processing seen in response to
visual deprivation, with the brain's reorganization in sighted individuals occurring more

rapidly and reversibly, while in blind individuals, it reflects more sustained and
structural changes over time.

While behavioral evidence clearly enhanced tactile abilities in visually deprived
individuals, the underlying neural mechanisms, particularly those occurring over short
timescales in sighted individuals, remain poorly understood. How the somatosensory
cortex might reorganize to enhance tactile perception within minutes or hours following
visual deprivation remains unknown. fMRI studies have shown changes in cortical
activity in short-term blindfolded individuals (Merabet et al., 2008; Siuda-Krzywicka et
al., 2016; Bola et al., 2017), but how the neural representation of specific tactile stimuli
is altered remains unclear. This gap in understanding arises because tactile
representations in S1 are distributed across high-dimensional neural population codes,
making it challenging for conventional methods to detect the subtle, stimulus-specific
features embedded in these signals (Romo and Salinas, 2001; Pei et al., 2011; Lieber
and Bensmaia, 2019). This highlights a gap in our understanding of whether—and
how—S1's population-level coding of tactile information may shift acutely when visual
input is removed. Thus, more refined techniques and models are needed to investigate
the immediate neural mechanisms underlying this form of cross-modal plasticity.

To address this gap, we focus on adult rats as a model for studying short-term
cross-modal influences in S1. Rodents have well-defined and accessible somatosensory
maps, enabling precise electrode placement for invasive recordings. In this study, we
use local field potentials (LFPs) recorded from S1, which capture network-level
dynamics not always apparent in single-unit spiking data or other imaging modalities.
LFPs provide a rich, high-dimensional signal that can reflect both synchronous and
asynchronous activity across cortical populations. By leveraging a machine learning
framework, we can decode subtle differences in these signals—differences that might
remain undetected through conventional amplitude-based analyses. Furthermore, to
dissociate the effects of attention in the perception of the somatosensory stimulus (Tamè
and Holmes, 2016), we created a custom behavioral paradigm that delivers the

somatosensory stimuli to rats in the most naturalistic manner, while diverting the rat's attention away from the stimuli (Yamashiro et al., 2024). This custom-designed treadmill paradigm allows us to manipulate both tactile (textured floors) and visual (light vs. dark) inputs independently, precisely controlling and isolating the contribution of visual cues to S1's tactile coding.

We hypothesized that visual deprivation acutely alters the neural representation of tactile stimuli in S1, making texture-specific signals more distinct in the absence of visual input. To test this hypothesis, we analyzed high-dimensional LFP data using a convolutional neural network, a type of deep learning model, to examine how S1 encodes different floor textures under light and dark conditions. Our results reveal subtle, yet systematic shifts in neural coding that emerge in the absence of visual input. By demonstrating this rapid reorganization of tactile representations, we provide new insights into the mechanisms of cross-modal processing at short timescales. These findings hold broader implications for sensory rehabilitation and brain–machine interfaces, suggesting that leveraging the dynamic interplay between sensory modalities could lead to improved outcomes for individuals with sensory impairments.

## Materials and Methods

*Animal ethics*

Animal experiments were performed with the approval of the Animal Experiment Ethics Committee at the University of Tokyo (approval numbers: P29–7 and P4–15) and according to the University of Tokyo guidelines for the care and use of laboratory animals. These experimental protocols were carried out following the Fundamental Guidelines for the Proper Conduct of Animal Experiments and Related Activities of the Academic Research Institutions (Ministry of Education, Culture, Sports, Science and Technology, Notice No. 71 of 2006), the Standards for Breeding and Housing of and Pain Alleviation for Experimental Animals (Ministry of the Environment, Notice No. 88 of 2006) and the Guidelines on the Method of Animal Disposal (Prime Minister's Office, Notice No. 40 of 1995). While our experimental protocols have a mandate to humanely euthanize animals if they exhibit any signs of pain, prominent lethargy, and discomfort, such symptoms were not observed in any of the rats tested in this study. All efforts were made to minimize the animals' suffering.

*Behavioral paradigm*

To record LFPs during natural locomotion, we employed a custom-designed, disk-shaped treadmill with a diameter of 90 cm (Figure 1A). The treadmill's running surface was divided into two halves, each featuring a distinct texture: one side was coated with coarse sandpaper (grain #80) and the other with fine sandpaper (grain #1000). In this setup, the rat was placed on the treadmill and secured with a fabric vest (Figure 1B).

Prior to the experiment, rats were water-restricted to ensure motivation. A waterspout was positioned in front of each rat, and its output was synchronized with the treadmill's movement (Figure 1C). Specifically, once the treadmill began to move, 30 μL of water was dispensed from the spout, encouraging the rat to continue walking to

receive its water reward.

Each rat was deprived of water until its body weight reached 85% of its baseline
weight, then it was trained to walk on the treadmill for four days (one hour per day),
with 30 minutes in a light environment (50 lx) followed by 30 minutes in darkness (0
lx). After successful training, the rats were allowed free access to food and water to
regain their original body weight prior to electrode implantation surgery. Following
surgical recovery, the rats were again water-restricted and placed on the treadmill to
assess cross-modal interactions between the visual and tactile systems. In a single
session, LFP recordings were obtained for 10 minutes in the light environment trial,
followed immediately by 10 minutes in the dark environment trial (Figure 1D).
Recording was performed once a day and performed daily for 4-14 days.

During the trials, the trajectory of the forelimb and onsets when the forelimb
came in contact with the floor were automatically identified using a previously
developed deep-learning–based method (Figure 1E) (Yamashiro et al., 2024).

*Animal preparation and surgical procedures*

LFPs were recorded from eleven 9- to 10-week-old Long-Evans rats (Japan SLC,
Shizuoka, Japan) using a custom-designed, 32-channel electrode assembly. This
assembly, fabricated from nichrome wires (761500, A-M Systems, WA, USA), targeted
the right somatosensory cortex (S1) regions corresponding to the forelimb and hindlimb
representations (Figure 1F). Specifically, 18 and 14 electrodes were placed in the
forelimb and the hindlimb subregion respectively (Figure 1G). Each electrode tip was
platinum-coated to reduce impedance to below 200 k$\Omega$ using a nanoZ tester (Plexon,
TX, USA).

At the start of the surgical procedure, each rat was anesthetized with 2–3%
isoflurane gas. A square craniotomy (2–6 mm posterior and 1–5 mm lateral to bregma)

was then created using a dental drill. The electrode assembly was gently lowered
through the cranial window to a depth of approximately 1.5 mm beneath the dura,
targeting layer IV of S1. Additionally, two stainless steel screws were implanted in the
bone above the cerebellum to serve as ground and reference electrodes. The recording
device and electrodes were secured to the skull using stainless steel screws and dental
cement. Following the surgery, each rat was housed individually in a transparent
Plexiglas cage with ad libitum access to food and water for one week to ensure proper
recovery.

LFP recordings from S1

LFPs were referenced to ground, digitized at 30 kHz using the OpenEphys recording
system (http://open-ephys.org) and an RHD 32-channel headstage (C3314, Intan
Technologies, CA, USA), then downsampled to 2 kHz for subsequent analyses
(Yamashiro et al., 2020). Concurrently, video footage was acquired at 60 Hz using a
USB camera module (MCM-303NIR, Gazo, Niigata, Japan), capturing a lateral view of
the rat. Each video frame was synchronized with the neural recordings via strobe
signals.

*Data analysis*

Data was analyzed offline using custom-made scripts in Python3. For box plots, the
centerline shows the median, the box limits show the upper and lower quartiles, and the
whiskers cover the 10−90% percentiles. $P < 0.05$ was considered statistically
significant. All statistical tests were two-sided.

 *LFP analysis*

To see if there were any differences in LFPs, the LFPs were aligned to the foot strike

onsets detected using deep-learning assisted methods. The aligned LFP were then

categorized by the trial (light *vs.* dark) and the floor texture (smooth *vs.* rough). To

analyze the amplitude of the event-related response from the foot strike, a mean trace of

the aligned LFP was calculated for each channel from 32 electrodes for each condition.

*A deep neural network for joint decoding of trial conditions and floor texture*

A custom deep neural network (DNN) model was implemented to predict both the trial

condition (*e.g.*, light *vs.* dark) and floor texture (*e.g.*, smooth *vs*. rough) from time-

aligned, one-dimensional LFP segments, using the PyTorch framework. Our model

architecture was inspired by one-dimensional ResNet-like structures and incorporated

multiheaded outputs for simultaneous prediction of two distinct variables(He et al.,

2015).

The input to the model consisted of one-dimensional LFP signals, arranged as a

tensor with multiple channels (*e.g.*, 32 input channels) over time. The network began by

splitting the input into two parallel convolutional pathways. The first pathway ("left"

branch) applied sequential convolutional and pooling operations with relatively smaller

kernel sizes and strides to incrementally reduce the dimensionality of the signal and

extract fine-grained temporal features. Specifically, the model employed a two-stage

convolutional process that first passed the input through a one-dimentional (1D)

convolution layer with a kernel size of 7, stride of 2, and batch normalization, followed

by a max pooling and an additional convolution layer. Both convolutional layers in the

left branch used ReLU nonlinearities to facilitate stable and efficient feature extraction.

In contrast, the second pathway ("right" branch) processed the input through a

single 1D convolutional layer with a larger kernel size (*e.g.*, 41) and a more aggressive

stride (*e.g.*, stride of 8). This pathway captured broader temporal contexts from the input

signals. Similar to the left branch, the right branch output was batch-normalized and

passed through a ReLU activation function. After these two parallel extractions, the

outputs of the left and right branches were concatenated along the channel dimension,

forming a combined feature representation that integrated both fine- and coarse-grained

temporal information.

The concatenated output was then passed through a max pooling operation,

followed by two residual layers that employed 1D convolutional blocks

(ResidualBlock) to refine feature representations. These residual layers allowed the

network to learn more complex feature hierarchies by facilitating the flow of gradients

during training and improving convergence, while also maintaining temporal resolution

appropriate for downstream decoding.

Subsequently, the processed features were passed through an average pooling

layer to summarize temporal information into a low-dimensional feature vector. This

vector was flattened into a one-dimensional representation and then fed into two

separate fully connected "heads" (fc_head1 and fc_head2). Each head was a simple

linear layer that provided a scalar output value. By concatenating these outputs, the final

layer jointly predicted two target variables from the same underlying features.

In summary, our model combined parallel convolutional branches for initial

feature extraction, residual layers for robust representation learning, and multiheaded

outputs to facilitate joint prediction of trial conditions and floor textures. The model was

implemented in Python using PyTorch, and all parameters were optimized via standard

stochastic gradient–based methods. This architecture allowed efficient and robust

decoding of environmental conditions from LFP signals in both time and frequency

domains.

*Training and evaluating the deep neural network*

Of the 11 recorded rats, data from the initial two rats were utilized to determine the
optimal model architecture. Once the architecture was established, data from the
remaining 9 rats were used for training and evaluation. The raw LFP signals were
downsampled to 10,000 Hz and segmented into 800 ms windows centered on the
footstrike onset (*e.g.*, 400 ms before and 400 ms after). Each LFP segment was assigned
two labels: one for the trial condition (*e.g.*, light *vs*. dark) and one for the floor texture
(*e.g.*, smooth *vs*. rough).

For each rat, the dataset was shuffled, normalized, and then split into training
(80%) and evaluation (20%) subsets. Using the training dataset, model training was
performed for 80 epochs using a batch size of 128 and a learning rate of $10^{-6}$. Binary
cross entropy with logits loss (BCEWithLogitsLoss) served as the objective function.
These parameters were selected to ensure stable convergence and to reduce overfitting.
After the training, the model performance was evaluated using the evaluation dataset.
The confusion matrix was calculated as the number of true positives, false positives,
false negatives, and true negatives, aggregated across all predictions in the evaluation
set.

*Explainability analysis using occlusion and integrated gradients*

To elucidate the internal decision-making processes of the trained deep neural network
(DNN), we employed two established explainability techniques: occlusion and
integrated gradients.

Occlusion: Occlusion analysis involves systematically masking specific input
features to determine their relative contribution to the model's output (Zeiler and
Fergus, 2013). In the present study, one channel was selectively occluded at a time from
the 32-channel LFP input and the sensitivity of each channel was calculated. Sensitivity

was defined as the corresponding change in model performance when the specified
channel was occluded. By conducting these analyses for each of the nine rats, channel-
specific importance scores were obtained and subsequently normalized (z-scored) to
facilitate cross-subject comparisons. Channels whose removal yielded a more
pronounced decrease in model performance were considered more critical for accurate
prediction.

Integrated gradients: Integrated gradients is an attribution method that quantifies
the importance of each input feature by integrating the gradient of the model's output
with respect to the input, transitioning from a baseline input to the actual input
(Sundararajan et al., 2017). This approach produces class activation maps, enabling the
visualization of features most influential for the model's output. Here, integrated
gradients were applied to the LFP segments for each rat, and the resulting class
activation maps were averaged across subjects and trial conditions. These maps allowed
us to identify salient input regions associated with both trial conditions and floor
texture.

**Results**

*Stable locomotion across light and dark conditions*

To investigate how visual input influences tactile processing in S1, we devised an
experimental paradigm where both tactile and visual inputs were independently
manipulated. Rats were placed on a disk-shaped treadmill with two distinct sandpaper
textures, and LFPs were recorded from walking rats. Each rat walked for 10 min in a
light environment (50 lx) and then for 10 minutes in total darkness (0 lx). To make sure
that rat's trajectory was stable across different floor textures and environmental
conditions, gait parameters were extracted from the trajectories using deep-learning–
based analysis (Figure 2A). From the trajectories, swing duration, stance duration, stride

length, and footstrike speed were extracted. All parameters were calculated for each

floor textures and environmental conditions. Comparison of all conditions revealed that

none of these metrics differed significantly between floor textures or environmental

conditions, indicating that overall locomotion remained stable (Figure 2B-E, $P > 0.05$,

one-way analysis of variance (ANOVA) followed by Tukey–Kramer *post hoc* test, $n =$

149, 149, 107 and 107 trials for smooth-light, rough-light, smooth-dark, and rough-dark,

respectively). Thus, any differences in neural activity under these conditions are

unlikely to be driven by altered motor behavior.

*Characteristic negative deflections in S1 LFP upon forelimb contact*

We next analyzed LFPs from S1 using a custom 32-channel electrode array targeting

the forelimb and hindlimb subregions (Figure 1F, G). LFP traces were first aligned to

forelimb contacts with the disk surface. On inspection of a single LFP trace, no apparent

event-related response could be observed (Figure 3A). However, aligning each LFP at

forelimb contact with the floor, the average trace showed a clear response after the onset

(Figure 3B). Across conditions, we observed a clear negative deflection in LFPs that

was most pronounced in the forelimb subregion (Figure 3C). The amplitude of the

response was significantly larger in all rats, consistent with topographic specificity of

S1 (Figure 3D, $P = 1.89 \times 10^{-3}$, $t_{10} = -2.2$, paired *t*-test, $n = 11$ rats).

To see if the event related responses were affected by the change in floor texture

or the environmental condition, the amplitudes were compared. Despite the prominent

negative deflection in LFPs at the forelimb contact with the floor, analyses showed no

substantial differences in signal amplitude (Figure 3E, $P > 0.05$, one-way analysis of

variance (ANOVA) followed by Tukey–Kramer *post hoc* test, $n = 149, 149, 107$ and

107 trials for smooth-light, rough-light, smooth-dark, and rough-dark, respectively)

when comparing rough *vs.* smooth textures (Figure 3F, $P = 0.48$ and $0.84$, $D = 1.03 \times 10^{-}$

$^{1}$ and $7.54 \times 10^{-2}$ for light and dark environments respectively, two-sample Kolmogorov–

Smirnov test, $n$ = 149 and 107 trials from 11 rats for light and dark, respectively) and

light *vs.* dark environments (Figure 5G, $P$ = 0.11, 0.18, $D$ = 0.14 and 0.149 for smooth

and rough textures respectively, two-sample Kolmogorov–Smirnov test, $n$ = 149 and

107 trials from 11 rats for light and dark, respectively). These findings suggest that

simple amplitude metrics may fail to capture more subtle or higher-dimensional

differences in the underlying neuronal population activity.

*Machine learning–based decoding of tactile and visual information*

Given the lack of clear differences in averaged LFP features, we applied a convolutional

neural network (CNN) to decode both the type of floor texture (smooth *vs.* rough) and

the presence or absence of visual cues (light *vs.* dark). The model architecture

incorporated parallel convolutional pathways to capture both macro- and micro-scale

temporal features, followed by residual blocks, and then split into dual output layers to

jointly predict texture and lighting conditions (Figure 4A).

When trained on 80% of the LFP data (segmented around footstrikes), the model

exhibited stable learning curves (Figure 4B) and generalized well to held-out test data

(Figure 4C). Confusion matrices for both texture and lighting classifications were above

chance along the diagonal, demonstrating that the model reliably extracted neural

representations of the floor textures from the LFPs (Figure 4D–F).

*Neural representations become more distinct in darkness*

To understand how absence of the visual cue might refine tactile processing, we

performed explainability analyses on the model's learned representations (Figure 5). We

extracted a 912-dimensional feature vector from the layer preceding the final outputs,

then visualized these high-dimensional embeddings with scatter plots (Figure 5A, B).

Clustering analyses using silhouette scores showed that representations of texture were
more separated in the dark environment, suggesting that reduced visual input enhances
the distinctness of neural codes for tactile stimuli (Figure 5C, $P = 4.83{\times}10^{-2}$, $t_8 = -2.6$,
paired $t$-test, $n = 9$ rats).

To elucidate which electrodes held most information about the floor textures, we
performed occlusion analysis. The results revealed that electrodes in the forelimb
subregion contributed more to successful texture and lighting predictions (Figure 5D, E,
$P = 4.53{\times}10^{-6}$, $t_8 = -5.57$, Student's $t$-test, $n = 9$ rats), which aligns with our observation
that negative deflections in LFPs were largest in forelimb-targeting channels (Figure
2E). Further analysis using feature maps showed spatiotemporal patterns of salient
features unique to each experimental condition (Figure 5F). In particular, the average
activation in the forelimb subregion displayed a temporal shift in the dark condition,
implying an extended processing window for texture information when visual cues are
unavailable (Figure 5G).

Collectively, these results indicate that visual deprivation modifies population-
level activity in S1, yielding more distinctive representations of tactile stimuli. Such
reorganization could provide a neural substrate for enhanced tactile perception under
conditions of reduced or absent visual input.

**Discussion**

This study provides evidence that S1 undergoes rapid and dynamic
reorganization of tactile representations when visual input is removed, even over a short
period of minutes. By combining high-density LFP recordings with advanced machine
learning techniques, we revealed that the neural encoding of tactile stimuli becomes
more distinct under conditions of visual deprivation. Notably, when rats walked in the
dark condition, texture representations in S1 were more distinguishable than when
visual inputs were present. These findings highlight the adaptability of S1 and
underscore the brain's ability to quickly adjust its tactile representations in response to
changing sensory contexts.

Interestingly, our data indicated that forelimb-targeting electrodes played a
crucial role in decoding both tactile textures and lighting conditions. First, the evoked
potential aligned at forelimb contact with the floor was larger at electrodes in forelimb
subregion of the S1. This finding is consistent with the topographic specificity of S1,
where distinct cortical regions are dedicated to processing sensory information from
different parts of the body (Sur et al., 1980; Ewert et al., 2008; Prsa et al., 2019).
Moreover, the occlusion analysis revealed that these forelimb regions were particularly
critical for distinguishing texture and lighting conditions, suggesting that the forelimb
subregion of S1 plays a dominant role in encoding tactile information in this context.
Although this result is intuitive, since LFPs were aligned at forelimb onset, this is the
first to reveal that a distinct cortical region encodes "tactile information" at the specific
part of the body. Visualization of the key channel and temporal features using integrated
gradients revealed results similar to those of the occlusion analysis. Specifically, the
forelimb channels were found to be particularly influential in encoding texture.
Additionally, the temporal window immediately following forelimb contact proved
critical for accurate predictions, with this temporal window being significantly longer in
the dark condition compared to the light condition.

The observed temporal shift in the dark condition, where texture-specific
processing extended over a longer time window, suggests that the absence of visual
input may enhance the retention of tactile information in S1. Previous studies have
shown that neurons in S1 are involved in the short-term retention of information (Zhou
and Fuster, 1996, 1997, 2000), and higher-order sensory areas exhibit prolonged
neuronal firing (Leavitt et al., 2017; Esmaeili and Diamond, 2019). This sustained
activity may reflect enhanced feedback loops within S1 and also from higher cortical
regions. The temporal shift observed in our study could thus indicate sustained neural
representations, possibly as a result of visual deprivation leading to a more prolonged
representation of tactile stimuli. Whether this shift reflects a compensatory mechanism
for the lack of visual input or directly correlates with heightened tactile sensitivity
requires further investigation.

The key point of this study is the use of a custom-designed behavioral paradigm
and CNN model to decode high-dimensional LFP data. Traditional analyses of neural
signals, such as amplitude-based methods or event-related potentials, often overlook
subtle, higher-dimensional features that are embedded in the data  (Yamins and DiCarlo,
2016a, 2016b; Saxena and Cunningham, 2019). By leveraging machine learning, we
were able to extract these fine-grained temporal and spatial patterns, which revealed that
texture-specific neural signals in S1 became more distinct when rats walked in the dark.
These results highlight the power of advanced computational methods in uncovering
previously undetectable shifts in sensory coding, paving the way for similar techniques
to be applied to other forms of high-dimensional neural data, such as multi-electrode
array recordings in freely behaving animals.

Previous studies on tactile enhancement in short-term visually deprived subjects
have shown improvements in tactile acuity and corresponding plastic changes in fMRI
BOLD signals (Pascual-Leone and Hamilton, 2001; Kauffman et al., 2002; Facchini and
Aglioti, 2003). However, due to the low temporal and spatial resolution of fMRI, these
studies could not capture detailed changes in tactile representation within the brain. In

contrast, our study successfully captured population-level activity from S1 using LFP
recordings. Although future research is needed to assess whether rats can more
accurately discriminate between textures in the dark, our findings suggest an underlying
mechanism driving tactile perception enhancement under short-term visual deprivation.

From an evolutionary standpoint, the enhanced tactile sensitivity observed in the
dark could offer a significant advantage. Many species, including rodents, rely on
somatosensation for navigation and foraging in low-visibility environments. The ability
to quickly enhance tactile processing in such conditions could aid in efficient
navigation, resource acquisition, and predator detection. Notably, changes in arousal
levels correlate with tactile sensitivity (Shimaoka et al., 2018; Lee et al., 2020), and
environmental lighting strongly influences arousal (Tamayo et al., 2023). This
mechanism may thus underlie the computational shifts observed in S1, further
emphasizing the dynamic nature of sensory processing. Rather than being a passive
receptor of tactile input, S1 appears to adapt its encoding strategies depending on the
sensory context, optimizing its function based on available visual or other sensory cues.

In addition to advancing our understanding of cross-modal processing, these
findings have potential applications for sensory rehabilitation. Our results suggest that
visual deprivation can rapidly enhance the brain's processing of tactile stimuli, a
principle that could be leveraged in individuals with tactile impairments to improve
performance. This concept could be explored in the context of training protocols such as
those used for Braille reading or haptic feedback devices, where manipulating sensory
input might improve tactile sensitivity (Bolognini et al., 2009). Moreover, the use of
machine learning techniques to decode neural signals holds promise for advancing
brain–machine interfaces and improving assistive technologies for individuals with
sensory deficits.

While this study contributes to our understanding of cross-modal processing,
several limitations must be considered. Our results are correlational, and future research

could incorporate causal manipulations, such as pharmacological interventions or
optogenetic techniques, to better elucidate the underlying mechanisms driving these
changes. Additionally, while rats are a powerful model system for studying sensory
processing, their neural architecture and behaviors may not fully reflect the complexities
of human perception. Further studies that explore the short-term effects of visual
deprivation on sensory encoding, or extend this paradigm to include behavioral readouts
of tactile acuity, would help clarify how changes in neural coding relate to perceptual
performance.

In conclusion, this study provides novel insights into the brain's capacity for
cross-modal adaptation, showing that visual deprivation rapidly reorganizes tactile
processing in S1. The enhanced distinction of tactile stimuli under conditions of visual
deprivation suggests that S1 can flexibly adjust its neural representations in response to
environmental changes. By combining behavioral paradigms, LFP recordings, and
machine learning techniques, this work highlights the value of advanced computational
approaches in understanding the dynamic and adaptive nature of sensory coding. These
findings open the door for future research on sensory rehabilitation and multisensory
interactions, and they underscore the brain's remarkable ability to reorganize itself in
response to changes in sensory input.

**Data availability**

The data that support the findings of this study are available from the corresponding
author upon reasonable request.

**Code availability**

The codes that support the findings of this study are available from the corresponding
author upon reasonable request.

## References

Abraira VE, Ginty DD (2013) The sensory neurons of touch. Neuron 79:618–639.

Bensmaia SJ, Denchev PV, Dammann JF 3rd, Craig JC, Hsiao SS (2008) The representation of
stimulus orientation in the early stages of somatosensory processing. J Neurosci 28:776–
786.

Bola Ł, Siuda-Krzywicka K, Paplińska M, Sumera E, Zimmermann M, Jednoróg K, Marchewka
A, Szwed M (2017) Structural reorganization of the early visual cortex following Braille
training in sighted adults. Sci Rep 7:17448.

Bolognini N, Pascual-Leone A, Fregni F (2009) Using non-invasive brain stimulation to
augment motor training-induced plasticity. J Neuroeng Rehabil 6:8.

Boroojerdi B, Bushara KO, Corwell B, Immisch I, Battaglia F, Muellbacher W, Cohen LG
(2000) Enhanced excitability of the human visual cortex induced by short-term light
deprivation. Cereb Cortex 10:529–534.

Bulusu V, Lazar L (2024) Crossmodal associations between naturally occurring tactile and
sound textures. Perception 53:219–239.

Burton H (2003) Visual cortex activity in early and late blind people. J Neurosci 23:4005–4011.

Bushnell MC, Duncan GH, Hofbauer RK, Ha B, Chen JI, Carrier B (1999) Pain perception: is
there a role for primary somatosensory cortex? Proc Natl Acad Sci U S A 96:7705–7709.

Butler JS, Foxe JJ, Fiebelkorn IC, Mercier MR, Molholm S (2012) Multisensory representation
of frequency across audition and touch: high density electrical mapping reveals early
sensory-perceptual coupling. J Neurosci 32:15338–15344.

Delhaye BP, Long KH, Bensmaia SJ (2018) Neural basis of touch and proprioception in primate
cortex. Compr Physiol 8:1575–1602.

Dijkerman HC, de Haan EHF (2007) Somatosensory processing subserving perception and
action: Dissociations, interactions, and integration. Behav Brain Sci 30:224–230.

Di Plinio S, Perrucci MG, Aleman A, Ebisch SJH (2020) I am Me: Brain systems integrate and
segregate to establish a multidimensional sense of self. Neuroimage 205:116284.

Driver J, Spence C (2000) Multisensory perception: beyond modularity and convergence. Curr
Biol 10:R731–R735.

Eimer M, Forster B (2003) Modulations of early somatosensory ERP components by transient
and sustained spatial attention. Exp Brain Res 151:24–31.

Esmaeili V, Diamond ME (2019) Neuronal correlates of tactile working memory in prefrontal
and vibrissal somatosensory cortex. Cell Rep 27:3167–3181.e5.

Ewert TAS, Vahle-Hinz C, Engel AK (2008) High-frequency whisker vibration is encoded by
phase-locked responses of neurons in the rat's barrel cortex. J Neurosci 28:5359–5368.

Facchini S, Aglioti SM (2003) Short term light deprivation increases tactile spatial acuity in
humans. Neurology 60:1998–1999.

Goldreich D, Kanics IM (2003) Tactile acuity is enhanced in blindness. J Neurosci 23:3439–
3445.

Goodwin AW, Wheat HE (2004) Sensory signals in neural populations underlying tactile
perception and manipulation. Annu Rev Neurosci 27:53–77.

He K, Zhang X, Ren S, Sun J (2015) Deep residual learning for image recognition. arXiv
[csCV] Available at: http://arxiv.org/abs/1512.03385.

Hopkins K, Kass SJ, Blalock LD, Brill JC (2017) Effectiveness of auditory and tactile
crossmodal cues in a dual-task visual and auditory scenario. Ergonomics 60:692–700.

Karlen SJ, Kahn DM, Krubitzer L (2006) Early blindness results in abnormal corticocortical and
thalamocortical connections. Neuroscience 142:843–858.

Kauffman T, Théoret H, Pascual-Leone A (2002) Braille character discrimination in blindfolded
human subjects. Neuroreport 13:571–574.

Koch KW, Fuster JM (1989) Unit activity in monkey parietal cortex related to haptic perception
and temporary memory. Exp Brain Res 76:292–306.

Leavitt ML, Mendoza-Halliday D, Martinez-Trujillo JC (2017) Sustained activity encoding
working memories: Not fully distributed. Trends Neurosci 40:328–346.

Lee CCY, Kheradpezhouh E, Diamond ME, Arabzadeh E (2020) State-dependent changes in
perception and coding in the mouse somatosensory cortex. Cell Rep 32:108197.

Lieber JD, Bensmaia SJ (2019) High-dimensional representation of texture in somatosensory
cortex of primates. Proc Natl Acad Sci U S A 116:3268–3277.

Luna R, Hernández A, Brody CD, Romo R (2005) Neural codes for perceptual discrimination in
primary somatosensory cortex. Nat Neurosci 8:1210–1219.

Merabet LB, Hamilton R, Schlaug G, Swisher JD, Kiriakopoulos ET, Pitskel NB, Kauffman T,
Pascual-Leone A (2008) Rapid and reversible recruitment of early visual cortex for touch.
PLoS One 3:e3046.

Moulton EA, Pendse G, Becerra LR, Borsook D (2012) BOLD responses in somatosensory
cortices better reflect heat sensation than pain. J Neurosci 32:6024–6031.

Nikbakht N, Tafreshiha A, Zoccolan D, Diamond ME (2018) Supralinear and Supramodal
Integration of Visual and Tactile Signals in Rats: Psychophysics and Neuronal
Mechanisms. Neuron 97:626–639.e8.

Norman JF, Bartholomew AN (2011) Blindness enhances tactile acuity and haptic 3-D shape
discrimination. Atten Percept Psychophys 73:2323–2331.

Pascual-Leone A, Hamilton R (2001) The metamodal organization of the brain. Prog Brain Res
134:427–445.

Pascual-Leone A, Torres F (1993) Plasticity of the sensorimotor cortex representation of the
reading finger in Braille readers. Brain 116 ( Pt 1):39–52.

Pei Y-C, Hsiao SS, Craig JC, Bensmaia SJ (2011) Neural mechanisms of tactile motion
integration in somatosensory cortex. Neuron 69:536–547.

Piché M, Chabot N, Bronchti G, Miceli D, Lepore F, Guillemot J-P (2007) Auditory responses
in the visual cortex of neonatally enucleated rats. Neuroscience 145:1144–1156.

Piras F, Vecchio D, Ciullo V, Gili T, Banaj N, Piras F, Spalletta G (2020) Sense of external
agency is sustained by multisensory functional integration in the somatosensory cortex.
Hum Brain Mapp 41:4024–4040.

Prsa M, Morandell K, Cuenu G, Huber D (2019) Feature-selective encoding of substrate
vibrations in the forelimb somatosensory cortex. Nature 567:384–388.

Rauschecker JP, Tian B, Korte M, Egert U (1992) Crossmodal changes in the somatosensory
vibrissa/barrel system of visually deprived animals. Proc Natl Acad Sci U S A 89:5063–
5067.

Romo R, Salinas E (2001) Touch and go: decision-making mechanisms in somatosensation.
Annu Rev Neurosci 24:107–137.

Sadato N, Pascual-Leone A, Grafman J, Ibañez V, Deiber MP, Dold G, Hallett M (1996)
Activation of the primary visual cortex by Braille reading in blind subjects. Nature
380:526–528.

Salinas E, Hernandez A, Zainos A, Romo R (2000) Periodicity and firing rate as candidate
neural codes for the frequency of vibrotactile stimuli. J Neurosci 20:5503–5515.

Sathian K, Stilla R (2010) Cross-modal plasticity of tactile perception in blindness. Restor
Neurol Neurosci 28:271–281.

Saxena S, Cunningham JP (2019) Towards the neural population doctrine. Curr Opin Neurobiol
55:103–111.

Schürmann M, Caetano G, Jousmäki V, Hari R (2004) Hands help hearing: facilitatory
audiotactile interaction at low sound-intensity levels. J Acoust Soc Am 115:830–832.

Serino A (2019) Peripersonal space (PPS) as a multisensory interface between the individual
and the environment, defining the space of the self. Neurosci Biobehav Rev 99:138–159.

Shimaoka D, Harris KD, Carandini M (2018) Effects of arousal on mouse sensory cortex

depend on modality. Cell Rep 22:3160–3167.

Siuda-Krzywicka K, Bola Ł, Paplińska M, Sumera E, Jednoróg K, Marchewka A, Śliwińska
MW, Amedi A, Szwed M (2016) Massive cortical reorganization in sighted Braille readers.
Available at: https://elifesciences.org/articles/10762 [Accessed February 18, 2025].

Sugiyama S, Kinukawa T, Takeuchi N, Nishihara M, Shioiri T, Inui K (2019) Tactile Cross-
Modal Acceleration Effects on Auditory Steady-State Response. Front Integr Neurosci
13:496317.

Sundararajan M, Taly A, Yan Q (2017) Axiomatic attribution for deep networks. arXiv [csLG]
Available at: http://arxiv.org/abs/1703.01365.

Sur M, Merzenich MM, Kaas JH (1980) Magnification, receptive-field area, and "hypercolumn"
size in areas 3b and 1 of somatosensory cortex in owl monkeys. J Neurophysiol 44:295–
311.

Tamayo E, Mouland JW, Lucas RJ, Brown TM (2023) Regulation of mouse exploratory
behaviour by irradiance and cone-opponent signals. BMC Biol 21:178.

Tamè L, Holmes NP (2016) Involvement of human primary somatosensory cortex in
vibrotactile detection depends on task demand. Neuroimage 138:184–196.

Van Boven RW, Hamilton RH, Kauffman T, Keenan JP, Pascual-Leone A (2000) Tactile spatial
resolution in blind braille readers. Neurology 54:2230–2236.

Wong M, Gnanakumaran V, Goldreich D (2011) Tactile spatial acuity enhancement in
blindness: evidence for experience-dependent mechanisms. J Neurosci 31:7028–7037.

Yamashiro K, Aoki M, Matsumoto N, Ikegaya Y (2020) Polyherbal formulation enhancing
cerebral slow waves in sleeping rats. Biol Pharm Bull 43:1356–1360.

Yamashiro K, Ikegaya Y, Matsumoto N (2024) Automatic detection of foot-strike onsets in a
rhythmic forelimb movement. Neurosci Res 206:41–50.

Yamins DL, DiCarlo JJ (2016a) Eight open questions in the computational modeling of higher
sensory cortex. Curr Opin Neurobiol 37:114–120.

Yamins DLK, DiCarlo JJ (2016b) Using goal-driven deep learning models to understand
sensory cortex. Nat Neurosci 19:356–365.

Zeiler MD, Fergus R (2013) Visualizing and understanding convolutional networks. arXiv
[csCV] Available at: http://arxiv.org/abs/1311.2901.

Zhou YD, Fuster JM (1996) Mnemonic neuronal activity in somatosensory cortex. Proc Natl
Acad Sci U S A 93:10533–10537.

Zhou YD, Fuster JM (1997) Neuronal activity of somatosensory cortex in a cross-modal (visuo-
haptic) memory task. Exp Brain Res 116:551–555.

Zhou YD, Fuster JM (2000) Visuo-tactile cross-modal associations in cortical somatosensory
cells. Proc Natl Acad Sci U S A 97:9777–9782.

Ziegler K, Folkard R, Gonzalez AJ, Burghardt J, Antharvedi-Goda S, Martin-Cortecero J,
Isaías-Camacho E, Kaushalya S, Tan LL, Kuner T, Acuna C, Kuner R, Mease RA, Groh A
(2023) Primary somatosensory cortex bidirectionally modulates sensory gain and
nociceptive behavior in a layer-specific manner. Nat Commun 14:2999.

**Figure 1. Behavioral paradigm and limb movement assessment with concurrent LFP recordings.**

(**A**) A diagram of the disk-shaped treadmill used in the experiment. One half of the disk is covered with #80 sandpaper, and the other half with #2000 sandpaper. (**B**) A frame from the video capturing a walking rat from a left-side perspective. (**C**) The motivation scheme. The rats were water-deprived prior to the experiments. A water port was coupled with the movement of the treadmill so that when the rat walked on the treadmill, the water would come out. This way, the rats were always motivated to walk during the whole session. (**D**) The experimental protocol, where each rat walked for 10 minutes in light (50 lx) and then for 10 minutes in darkness (0 lx). (**E**) An example trajectory of the elbow and wrist joints from one session, plotted with the shoulder joint fixed in the coordinate space. (**F**) A schematic illustrating the forelimb and hindlimb subregions of S1. (**G**) A custom 32-channel electrode array used to record LFPs from these subregions.

**Figure 2. Comparison of gait parameters across textures and environmental conditions.**

(**A**) Swing phase *vs*. stance phase, illustrated with video frames (left: swing, right: stance). (**B**) Normalized swing duration measured for each rat under different textures (smooth *vs*. rough)  and environmental condition (light *vs*. dark). There were no significant differences among trial conditions. (**C–E**) Stance duration, stride length, and footstrike speed, respectively, under the same conditions as in B. None of these parameters differed significantly across texture types or lighting conditions.

**Figure 3. LFP recordings in rat S1 during walking.**

(**A**) A single representative LFP trace aligned to a forelimb contact. (**B**) An example of an averaged LFP trace from one session, aligned to forelimb contact. (**C**) The electrode montage and averaged LFP at each electrode. Left: The electrode montage showing all 32 recording sites. Right: Averaged LFP signals aligned to forelimb contacts with the floor, shown for each electrode depicted in the left panel. (**D**) Comparison of amplitudes

between hindlimb and forelimb subregions, aggregated across all 11 rats. $P = 1.89 \times 10^{-3}$,

$t_{10}$ = -2.2, paired t-test, $n = 11$ rats. (**E**) Comparison of averaged amplitudes within the

same trial for different floor textures and environmental conditions. $P > 0.05$, one-way

analysis of variance (ANOVA) followed by Tukey–Kramer post hoc test, $n = 149, 149$,

107 and 107 trials for smooth-light, rough-light, smooth-dark, and rough-dark,

respectively. (**F**) Cumulative probability distributions of mean amplitude from each

session, compared across different textures. $P = 4.84 \times 10^{-1}$, $8.35 \times 10^{-1}$, $D = 1.03 \times 10^{-1}$

and $7.54 \times 10^{-2}$ for light and dark environments respectively, two-sample Kolmogorov–

Smirnov test, $n = 149$ and 107 trials from 11 rats for light and dark, respectively. (**G**)

Cumulative probability distributions of mean amplitude from each session, compared

across light and dark environments. $P = 1.05 \times 10^{-1}$, $1.83 \times 10^{-1}$, $D = 0.14$ and 0.149 for

smooth and rough textures respectively, two-sample Kolmogorov–Smirnov test, $n = 149$

and 107 trials from 11 rats for light and dark, respectively. Abbreviations: ERP, event-

related potential; LFP, local field potential; S1, primary somatosensory cortex.

**Figure 4. Model-based prediction of texture and environmental conditions from**
**LFP.**

(**A**) The deep learning model architecture. The LFP input is processed through two
parallel pathways for macro- and micro-scale feature extraction, followed by residual
blocks that feed into two output heads for floor texture (Smooth *vs.* Rough) and the
environmental condition (Light *vs.* Dark). (**B**) Training performance for a single
representative rat. The left graph shows accuracy curves for texture (blue) and lighting
(yellow), and the right graph shows the loss curves. (**C**) Testing performance for the
same rat. The model exhibits good generalization, as accuracy increases and loss
decreases on held-out data. (**D**) Confusion matrix for texture classification for all rats.
Values above chance on the diagonal indicate successful texture prediction. Note that all
values in the same row add up to 1. (**E**) Same as D, but for environmental conditions.
(**F**) Combined confusion matrix for texture and trial predictions. The model performs

well on both tasks across all rats. Abbreviations: avgpool, average pooling layer; conv,

convolutional layer; maxpool, max pooling layer; LFP, local field potential

**Figure 5. Neural representations become distinct in dark environments than in light.**

(**A**) A 912-dimensional feature vector is extracted from the layer preceding the final output. (**B**) A scatter plot of these features from one rat shows individual LFP segments (aligned to forelimb contact). Light orange and dark orange correspond to the light conditions (smooth, rough), while light blue and dark blue correspond to the dark conditions (smooth, rough). (**C**) Silhouette scores across all nine rats, showing that the dark condition yields higher scores and thus more distinct neural representations. $P = 4.83 \times 10^{-2}$, $t_8 = -2.6$, paired $t$-test, $n = 9$ rats. (**D**) A pseudo-color map based on occlusion analysis, illustrating the contribution of each electrode in the forelimb and hindlimb

subregions. Hotter regions indicate higher importance for the model's predictions. (**E**)

Forelimb channels exhibit higher occlusion sensitivity than hindlimb channels,

highlighting the forelimb's dominant role when the foot contacts the floor. $P = 4.53 \times 10^{-6}$,

$t_{16}$ = -5.57, Student's *t*-test, $n$ = 9 rats. (**F**) Activation maps generated via integrated

gradients highlight key input features responsible for accurate model predictions of

texture and environmental conditions. Activation scores show each feature's impact on

the model's output relative to a reference baseline: high positive scores denote features

that strongly affect the predicted class. The onset of forelimb contact is aligned to time

zero. (**G**) Activation scores averaged over forelimb electrodes for floor texture (left) and

the environmental conditions (right). A temporal lag in the dark condition suggests an

extended processing window for floor texture when visual cues are absent.

**A**

Feature extraction

**B**

Light-Smooth
Light-Rough
Dark-Smooth
Dark-Rough

PC 2

PC 1

**C**

(×10⁻³)

Silhouette score

Light    Dark

**D**

Forelimb

Sensitivity (z-scored)

Hindlimb

**E**

Sensitivity (z-scored)

Hind    Fore

**F**

Smooth

Light

Fore

Hind

Rough

Dark

Fore

Hind

Channel

Attribution score

Time (ms)

**G**

Attribution score

Rough

Smooth

Dark

Light

Time (ms)