Enhanced Tactile Coding in Rat Neocortex Under Darkness

Kotaro Yamashiro[1], Shiyori Tanaka[1], Nobuyoshi Matsumoto[1,2†], Yuji Ikegaya[1,2,3†]

[1]Graduate School of Pharmaceutical Sciences, The University of Tokyo, Tokyo, 113-0033, Japan
[2]Institute for AI and Beyond, The University of Tokyo, Tokyo, 113-0033, Japan
[3]Center for Information and Neural Networks, National Institute of Information and Communications Technology, Suita City, Osaka, 565-0871, Japan

Abbreviated title: Enhanced Tactile Coding Under Darkness

†To whom correspondence should be addressed

Yuji Ikegaya Ph.D.
Laboratory of Chemical Pharmacology
Graduate School of Pharmaceutical Sciences, The University of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
Tel. : +81-3-5841-4780; Fax: +81-3-5841-4786
E-mail: yuji@ikegaya.jp

Nobuyoshi Matsumoto Ph.D.
Laboratory of Chemical Pharmacology
Graduate School of Pharmaceutical Sciences, The University of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
Tel. : +81-3-5841-4780; Fax: +81-3-5841-4786
E-mail: nobuyoshi@matsumoto.ac

Conflicts of interest: The authors declare no competing interests.

Acknowledgments: This work was supported by JST ERATO (JPMJER1801), AMED-CREST (24wm0625401h0001; 24wm0625502s0501; 24wm0625207s0101; 24gm1510002s0104), the Institute for AI and Beyond of the University of Tokyo, JSPS Grants-in-Aid for Scientific Research (18H05525, 20K15926, 22K21353, 22J22097), KOSÉ Cosmetology Research Foundation, the Public Foundation of Chubu Science and Technology Center, and Konica Minolta Science and Technology Foundation.

Abstract

Sensory systems are known for their adaptability, responding dynamically to changes in environmental conditions. A key example of this adaptability is the enhancement of tactile perception in the absence of visual input. Despite behavioral studies showing visual deprivation can improve tactile discrimination, the underlying neural mechanisms, particularly how tactile neural representations are reorganized during visual deprivation, remain unclear. In this study, we explore how the absence of visual input alters tactile neural encoding in the rat somatosensory cortex (S1). Rats were trained on a custom-designed treadmill with distinct tactile textures (rough and smooth), and local field potentials (LFPs) were recorded from S1 under light and dark conditions. Machine learning techniques, specifically a convolutional neural network, were used to decode the high-dimensional LFP signals. We found that the neural representations of tactile stimuli became more distinct in the dark, indicating a reorganization of sensory processing in S1 when visual input was removed. Notably, conventional amplitude-based analyses failed to capture these changes, highlighting the power of deep learning in uncovering subtle neural patterns. These findings offer new insights into how the brain rapidly adapts tactile processing in response to the loss of visual input, with implications for multisensory integration.

Introduction

The primary somatosensory cortex (S1) is integral to the encoding of tactile information, processing sensory inputs from various regions of the body (Delhaye et al., 2018; Di Plinio et al., 2020; Piras et al., 2020; Serino, 2019). This cortical area is crucial for our sense of touch, facilitating the perception and interpretation of sensations such as pressure, vibration, temperature, and pain (Bushnell et al., 1999; Luna et al., 2005; Moulton et al., 2012). These external stimuli are represented by dynamic neural activity in S1, enabling animals to discriminate distinct sensory experiences (Bensmaia et al., 2008; Goodwin and Wheat, 2004; Koch and Fuster, 1989; Salinas et al., 2000). Crucially, neural representations in S1 are highly adaptable, shaped not only by feedforward mechanisms but also by attention, motivation, and inputs from other sensory modalities (Butler et al., 2012; Driver and Spence, 2000; Eimer and Forster, 2003; Schürmann et al., 2004; Ziegler et al., 2023). This adaptability facilitates rapid adjustments to changing environmental contexts, supporting survival and optimal sensory processing (Abraira and Ginty, 2013; Dijkerman and de Haan, 2007).

One prominent example of such adaptability is cross-modal processing, wherein the absence or reduction of visual input enhances tactile abilities. (Bulusu and Lazar, 2024; Hopkins et al., 2017; Nikbakht et al., 2018; Sugiyama et al., 2019). For instance, prolonged visual deprivation, such as blindness from an early age, results in heightened tactile discrimination abilities (Goldreich and Kanics, 2003; Norman and Bartholomew, 2011; Van Boven et al., 2000; Wong et al., 2011), driven by cortical reorganization where visual cortical areas are repurposed to support tactile processing (Burton, 2003; Karlen et al., 2006; Sadato et al., 1996). Notably, however, tactile performance improvements are also observed in sighted individuals during short-term visual deprivation (Facchini and Aglioti, 2003; Kauffman et al., 2002; Pascual-Leone and Hamilton, 2001a). Such individuals demonstrate enhanced tactile discrimination during temporary darkness or blindfolding, indicating that rapid, context-dependent compensatory mechanisms occur even without long-term structural changes (Bola et al.,

2017; Boroojerdi et al., 2000; Merabet et al., 2008). Thus, although tactile enhancement

has been studied extensively in the context of long-term sensory deprivation, the neural

mechanisms underlying short-term, context-dependent modulation remain unclear.

The neural representation of tactile stimuli within S1 involves complex,

distributed activity patterns, posing significant challenges for conventional analyses that

primarily measure signal amplitude. Thus, the subtle, high-dimensional neural dynamics

that may underlie immediate adaptations to temporary visual deprivation are difficult to

detect using traditional methods. Addressing this gap requires refined experimental

paradigms and advanced analytical tools to capture how tactile representations

reorganize on short timescales.

To examine short-term reorganization in tactile processing, we used a behavioral

paradigm using rats trained to walk naturally on a treadmill featuring distinct tactile

textures (Yamashiro et al., 2024). . By manipulating visual input (light *vs*. dark

conditions) while recording local field potentials (LFPs) from S1, we aimed to

characterize rapid shifts in tactile neural representations. LFPs provide a rich, high-

dimensional signal reflecting both synchronous and asynchronous activity across

cortical populations. Concurrently, we applied deep learning techniques to decode

subtle differences in neural activity patterns that traditional amplitude-based analyses

could not detect. We hypothesized that temporary visual deprivation would enhance the

neural distinction between tactile stimuli, leading to more differentiated representations

of textures in the dark condition.

By investigating these rapid, context-dependent shifts in neural coding, this

study provides valuable insights into the brain's capacity for sensory adaptation. Our

findings not only shed light on how visual deprivation influences tactile processing in

the short term, but also emphasize the potential of deep learning techniques in

uncovering nuanced neural reorganization.

Materials and Methods

Animal ethics

Animal experiments were performed with the approval of the Animal Experiment Ethics Committee at the University of Tokyo (approval numbers: P29–7 and P4–15) and according to the University of Tokyo guidelines for the care and use of laboratory animals. These experimental protocols were carried out following the Fundamental Guidelines for the Proper Conduct of Animal Experiments and Related Activities of the Academic Research Institutions (Ministry of Education, Culture, Sports, Science and Technology, Notice No. 71 of 2006), the Standards for Breeding and Housing of and Pain Alleviation for Experimental Animals (Ministry of the Environment, Notice No. 88 of 2006) and the Guidelines on the Method of Animal Disposal (Prime Minister's Office, Notice No. 40 of 1995). While our experimental protocols have a mandate to humanely euthanize animals if they exhibit any signs of pain, prominent lethargy, and discomfort, such symptoms were not observed in any of the rats tested in this study. All efforts were made to minimize the animals' suffering.

Behavioral paradigm

To record LFPs during natural locomotion, we employed a custom-designed, disk-shaped treadmill with a diameter of 90 cm (Figure 1A). The treadmill's running surface was divided into two halves, each featuring a distinct texture: one side was coated with coarse sandpaper (grain #80) and the other with fine sandpaper (grain #1000). In this setup, the rat was placed on the treadmill and secured with a fabric vest (Figure 1B).

Before the experiment, rats were water-restricted to ensure motivation. A waterspout was positioned in front of each rat, and its output was synchronized with the treadmill's movement (Figure 1C). Specifically, once the treadmill began to move, 30 µL of water was dispensed from the spout, encouraging the rat to continue walking to

receive its water reward.

Each rat was deprived of water until its body weight reached 85% of its baseline

weight, then it was trained to walk on the treadmill for four days (one hour per day),

with 30 minutes in a light environment (50 lx) followed by 30 minutes in darkness (0

lx). After successful training, the rats were allowed free access to food and water to

regain their original body weight before electrode implantation surgery. Following

surgical recovery, the rats were water-restricted again and placed on the treadmill to

assess cross-modal interactions between the visual and tactile systems. In a single

session, LFP recordings were obtained in the light environment trial, followed

immediately by the dark environment trial (Figure 1D). For subset of sessions, the

order of the light and dark environment was swapped, dark environment trial preceding

the dark environment trial. Each trial lasted for approximately 10 minutes. Recording

was performed once a day for 4-14 days.

During the trials, the trajectory of the forelimb and onsets when the forelimb

came in contact with the floor were automatically identified using a previously

developed deep-learning–based method (Figure 1E) (Yamashiro et al., 2024).

Animal preparation and surgical procedures

LFPs were recorded from eleven 9- to 10-week-old Long-Evans rats (Japan SLC,

Shizuoka, Japan) using a custom-designed, 32-channel electrode assembly. This

assembly, fabricated from nichrome wires (761500, A-M Systems, WA, USA), targeted

the right somatosensory cortex (S1) regions corresponding to the forelimb and hindlimb

representations (Figure 1F). Specifically, 18 and 14 electrodes were placed in the

forelimb and the hindlimb subregion respectively (Figure 1G). Each electrode tip was

platinum-coated to reduce impedance to below 200 kΩ using a nanoZ tester (Plexon,

TX, USA).

At the start of the surgical procedure, each rat was anesthetized with 2–3%

isoflurane gas. A square craniotomy (2–6 mm posterior and 1–5 mm lateral to bregma)

was then created using a dental drill. The electrode assembly was gently lowered

through the cranial window to a depth of approximately 1.5 mm beneath the dura,

targeting layer IV of S1. Additionally, two stainless steel screws were implanted in the

bone above the cerebellum to serve as ground and reference electrodes. The recording

device and electrodes were secured to the skull using stainless steel screws and dental

cement. Following the surgery, each rat was housed individually in a transparent

Plexiglas cage with ad libitum access to food and water for one week to ensure proper

recovery.

LFP recordings from S1

LFPs were referenced to ground, digitized at 30 kHz using the OpenEphys recording

system (http://open-ephys.org) and an RHD 32-channel headstage (C3314, Intan

Technologies, CA, USA), then resampled to 10 kHz for subsequent analyses

(Yamashiro et al., 2020). In parallel, video was acquired at 60 Hz using a USB camera

module (MCM-303NIR, Gazo, Niigata, Japan), capturing a lateral view of the rat. Each

video frame was synchronized with the neural recordings using strobe signals. For a

subset of recordings (n = 2 rats), speed sensors were attached to the running disk to

monitor locomotor speed, which was simultaneously recorded through the OpenEphys

analog input.

Data analysis

Data was analyzed offline using custom-made scripts in Python3. For box plots, the

centerline shows the median, the box limits show the upper and lower quartiles,

and the whiskers cover the 10−90% percentiles. $P < 0.05$ was considered

statistically significant. All statistical *t*-tests were two-sided, and the bootstrap method

was applied for multiple comparisons.

LFP analysis

To see if there were any differences in LFPs, the LFPs were aligned to the foot strike

onsets detected using deep-learning assisted methods. The aligned LFP were then

categorized by the trial (light *vs.* dark) and the floor texture (smooth *vs.* rough). To

analyze the amplitude of the event-related response from the foot strike, a mean trace of

the aligned LFP was calculated for each channel from 32 electrodes for each condition.

Foot-strike detection and all LFP alignment were intentionally forelimb-locked, because

the behavioral paradigm was designed for forepaw texture contact (Figure 1E) and our

array oversampled S1-forelimb (18 channels) relative to S1-hindlimb (14 channels;

Figure 1G).

A deep neural network for joint decoding of trial conditions and floor texture

A custom deep neural network (DNN) model was implemented to predict both the trial

condition (*e.g.*, light *vs.* dark) and floor texture (*e.g.*, smooth *vs.* rough) from time-

aligned, one-dimensional LFP segments, using the PyTorch framework. Our model

architecture was inspired by one-dimensional ResNet-like structures and incorporated

multiheaded outputs for simultaneous prediction of two distinct variables (He et al.,

2015).

The input to the model consisted of one-dimensional LFP signals, arranged as a

tensor with multiple channels (*e.g.*, 32 input channels) over time. The network began by

splitting the input into two parallel convolutional pathways. The first pathway ("left"

branch) applied sequential convolutional and pooling operations with relatively smaller
kernel sizes and strides to incrementally reduce the dimensionality of the signal and
extract fine-grained temporal features. Specifically, the model employed a two-stage
convolutional process that first passed the input through a one-dimensional (1D)
convolution layer with a kernel size of 7, stride of 2, and batch normalization, followed
by a max pooling and an additional convolution layer. Both convolutional layers in the
left branch used ReLU nonlinearities to facilitate stable and efficient feature extraction.

In contrast, the second pathway ("right" branch) processed the input through a
single 1D convolutional layer with a larger kernel size (*e.g.*, 41) and a more aggressive
stride (*e.g.*, stride of 8). This pathway captured broader temporal contexts from the input
signals. Similar to the left branch, the right branch output was batch-normalized and
passed through a ReLU activation function. After these two parallel extractions, the
outputs of the left and right branches were concatenated along the channel dimension,
forming a combined feature representation that integrated both fine- and coarse-grained
temporal information.

The concatenated output was then passed through a max pooling operation,
followed by two residual layers that employed 1D convolutional blocks
(ResidualBlock) to refine feature representations. These residual layers allowed the
network to learn more complex feature hierarchies by facilitating the flow of gradients
during training and improving convergence, while also maintaining temporal resolution
appropriate for downstream decoding.

Subsequently, the processed features were passed through an average pooling
layer to summarize temporal information into a low-dimensional feature vector. This
vector was flattened into a one-dimensional representation and then fed into two
separate fully connected "heads" (fc_head1 and fc_head2). Each head was a simple
linear layer that provided a scalar output value. By concatenating these outputs, the final
layer jointly predicted two target variables from the same underlying features.

In summary, our model combined parallel convolutional branches for initial
feature extraction, residual layers for robust representation learning, and multiheaded
outputs to facilitate joint prediction of trial conditions and floor textures. The model was
implemented in Python using PyTorch, and all parameters were optimized via standard
stochastic gradient–based methods. This architecture allowed efficient and robust
decoding of environmental conditions from LFP signals in both time and frequency
domains.

Training and evaluating the deep neural network

Of the 11 recorded rats, data from the initial two rats were utilized to determine the
optimal model architecture. Once the architecture was established, data from the
remaining 9 rats were used for training and evaluation. The raw LFP signals were
resampled from 30 kHz to 10 kHz and segmented into 800 ms windows centered on the
footstrike onset (*e.g.*, 400 ms before and 400 ms after). Each LFP segment was assigned
two labels: one for the trial condition (*e.g.*, light *vs*. dark) and one for the floor texture
(*e.g.*, smooth *vs*. rough).

For each rat, the dataset was shuffled, normalized, and then subjected to 5-fold
cross-validation. In this procedure, the data were partitioned into five equal subsets; in
each fold, four subsets (80%) were used for training, and the remaining subset (20%)
was used for evaluation. Model training was performed for 80 epochs using a batch size
of 128 and a learning rate of 10^{-6}. Binary cross-entropy with logits loss
(BCEWithLogitsLoss) served as the objective function. These parameters were selected
to ensure stable convergence and to reduce overfitting. After the training, the model's
performance was evaluated using the evaluation dataset. The confusion matrix was
calculated as the number of true positives, false positives, false negatives, and true
negatives, aggregated across all predictions in the evaluation set.

Evaluation of cluster separability between light and dark conditions

To examine how texture representations differed between light and dark conditions, we
assessed cluster separability of the intermediate features extracted from the deep-
learning model. Specifically, after the model was trained, we extracted the output from
the penultimate layer (a 912-dimensional feature vector for each trial). Cluster
separability was quantified using the silhouette score, which measures the relative
distances between within-cluster and between-cluster samples.

Silhouette scores were calculated for the two texture classes (smooth and rough)
within each lighting condition, using the Euclidean distance metric applied to the
intermediate feature representations. For each trial, the silhouette score s is defined as:

$$s = \frac{b - a}{\max(a, b)}$$

where a is the average distance between a point and all other points in the same cluster
(intra-cluster distance), and b is the minimum average distance between a point and all
points in the nearest different cluster (inter-cluster distance). The silhouette score,
therefore, provides a normalized index of cluster separability, with larger values
indicating more distinct clustering. To obtain an animal-level measure, silhouette scores
were averaged across cross-validation within each rat.

Explainability analysis using occlusion and integrated gradients

To elucidate the internal decision-making processes of the trained deep neural network
(DNN), we employed two established explainability techniques: occlusion and
integrated gradients.

Occlusion: Occlusion analysis involves systematically masking specific input

features to determine their relative contribution to the model's output (Zeiler and Fergus,

2013). In the present study, one channel was selectively occluded at a time from the 32-

channel LFP input and the sensitivity of each channel was calculated. Sensitivity was

defined as the corresponding change in model performance when the specified channel

was occluded. By conducting these analyses for each of the nine rats, channel-specific

importance scores were obtained and subsequently normalized (z-scored) to facilitate

cross-subject comparisons. Channels whose removal yielded a more pronounced

decrease in model performance were considered more critical for accurate prediction.

Integrated gradients: Integrated gradients is an attribution method that quantifies

the importance of each input feature by integrating the gradient of the model's output

with respect to the input, transitioning from a baseline input to the actual input

(Sundararajan et al., 2017). This approach produces class activation maps, enabling the

visualization of features most influential for the model's output. Here, integrated

gradients were applied to the LFP segments for each rat, and the resulting class

activation maps were averaged across subjects and trial conditions. These maps allowed

us to identify salient input regions associated with both trial conditions and floor texture.

Results

Stable locomotion across light and dark conditions

To investigate how visual input influences tactile processing in S1, we devised an

experimental paradigm where both tactile and visual inputs were independently

manipulated. Rats were placed on a disk-shaped treadmill with two distinct sandpaper

textures, and LFPs were recorded from walking rats. Each rat walked for 10 min in a

light environment (50 lx) and then for 10 minutes in total darkness (0 lx).

To make sure that rat's trajectory was stable across different floor textures and

environmental conditions, gait parameters were extracted from the trajectories using

deep-learning–based analysis (Figure 2A). From the trajectories, swing duration, stance

duration, stride length, and footstrike speed were extracted. All parameters were

calculated for each floor textures and environmental conditions. Comparison of all

conditions revealed that none of these metrics differed significantly between floor

textures or environmental conditions, indicating that overall locomotion remained stable

(Figure 2B-E, $P > 0.05$, one-way analysis of variance (ANOVA) followed by

Tukey☐Kramer *post hoc* test, $n = 149, 149, 107$ and 107 trials for smooth-light, rough-

light, smooth-dark, and rough-dark, respectively). Locomotor-speed traces recorded

from subset of rats. The result indicated that walking velocity was stable across trials,

not affected by the lighting conditions (Supplementary Figure 1). Thus, any differences

in neural activity under these conditions are unlikely to be driven by altered motor

behavior.

Characteristic of S1 LFP upon forelimb contact

We next analyzed LFPs from S1 using a custom 32-channel electrode array targeting

the forelimb and hindlimb subregions (Figure 1F, G). LFP traces were first aligned to

forelimb contacts with the disk surface. On inspection of a single LFP trace, no apparent

event-related response could be observed (Figure 3A). However, aligning each LFP at

forelimb contact with the floor, the average trace showed a clear response after the onset

(Figure 3B). Across conditions, we observed a clear negative deflection in LFPs that

was most pronounced in the forelimb subregion (Figure 3C). The amplitude of the

response was significantly larger in all rats, consistent with the topographic specificity

of S1 (Figure 3D, $P = 1.89 \times 10^{-3}$, $t_{10} = -2.2$, paired *t*-test, $n = 11$ rats). This was expected

since the epochs were forelimb-locked, making larger deflections and greater

importance of forelimb channels consistent with somatotopy.

To see if the event-related responses were affected by the change in floor texture

or the environmental condition, the amplitudes were compared. Despite the prominent

negative deflection in LFPs at the forelimb contact with the floor, analyses showed no

substantial differences in signal amplitude (Figure 3E, $P > 0.05$, one-way analysis of

variance (ANOVA) followed by Tukey☐Kramer *post hoc* test, $n = 149, 149, 107$ and

107 trials for smooth-light, rough-light, smooth-dark, and rough-dark, respectively)

when comparing rough *vs.* smooth textures (Figure 3F, $P = 0.48$ and 0.84, $D = 1.03 \times 10^{-1}$

and 7.54×10^{-2} for light and dark environments respectively, two-sample

Kolmogorov☐Smirnov test, $n = 149$ and 107 trials from 11 rats for light and dark,

respectively) and light *vs.* dark environments (Figure 6G, $P = 0.11, 0.18$, $D = 0.14$ and

0.149 for smooth and rough textures respectively, two-sample Kolmogorov☐Smirnov

test, $n = 149$ and 107 trials from 11 rats for light and dark, respectively).

To further investigate, we computed grand-average LFP waveforms across

combinations of floor texture (smooth *vs.* rough) and lighting condition (light *vs.* dark)

(Figure 4A). Visual inspection of the mean waveforms revealed no distinct or prominent

features. We then calculated channel-wise correlations between textures within each

lighting condition (i.e., light-smooth *vs.* light-rough; dark-smooth *vs.* dark-rough). To

assess whether lighting modulated texture-related correlations, we subtracted the

correlation coefficients obtained in the dark from those in the light and averaged the

differences across animals (Figure 4B). The correlation of LFP waveforms between

different textures was stronger in the light condition, suggesting that in the dark

environment, the average LFP waveform is more distinct. Additionally, we examined

channel–channel correlation matrices across all 32 channels for each condition (Figure

4C). While stronger correlations were observed within individual S1 subregions, no

systematic differences emerged between lighting conditions or between textures. To

probe the frequency domain of the LFP, we computed time–frequency (wavelet)

spectrograms across the floor texture and lighting condition combinations. However, no

clear differences were observed across these combinations. Taken together, these results

suggest that simple amplitude-based metrics may not fully capture subtler or higher-

dimensional variations in underlying neuronal activity.

Machine learning–based decoding of tactile and visual information

Given the absence of clear differences in averaged LFP features, we next examined how
representations of textures in the LFPs varied across environments using machine
learning approaches. We first implemented a support vector machine (SVM) classifier
with a radial basis function (RBF) kernel. To reduce dimensionality, principal
component analysis (PCA) was applied to the raw LFP traces (Supplementary Figure
2A). The SVM was then trained using 5-fold cross-validation, where, in each fold, 80%
of the dataset was used to classify two types of labels: texture (smooth *vs*. rough) and
trial (light *vs*. dark). Model performance was evaluated on the remaining 20% of the
data. The overall classification accuracy was at chance level (~50%), indicating that the
PCA–SVM combination failed to extract informative features for texture or lighting
condition from the LFP traces (Supplementary Figure 2B).

We next employed a deep learning model, a convolutional neural network
(CNN), to classify both floor texture (smooth *vs*. rough) and trial (light *vs*. dark). The
CNN architecture incorporated parallel convolutional pathways designed to capture
both macro- and micro-scale temporal dynamics, followed by residual blocks. The
network then branched into dual output layers to jointly predict texture and lighting
conditions (Figure 5A).

Similarly to SVM, the model was trained and evaluated using 5-fold cross-
validation. In training, the model exhibited stable learning curves (Figure 5B) and also
generalized well to held-out test data (Figure 5C). Confusion matrices for both texture
and lighting classifications were above chance along the diagonal, demonstrating that
the model reliably extracted neural representations of the floor textures from the LFPs
(Figure 5D–F).

 Neural representations become more distinct in darkness

To understand how the absence of the visual cue might refine tactile processing, we
performed explainability analyses on the model's learned representations (Figure 6). We
extracted a 912-dimensional feature vector from the layer preceding the final outputs,
then visualized these high-dimensional embeddings with scatter plots (Figure 6A, B).
Clustering analyses using silhouette scores showed that representations of texture were
more separated in the dark environment, suggesting that reduced visual input enhances
the distinctness of neural dynamics that code tactile stimuli (Figure 6C, $P = 2.31 \times 10^{-2}$, t_8
$= -2.8$, paired *t*-test, $n = 9$ rats, Supplementary Table 1). To account for trial order,
silhouette scores were also calculated in sessions where dark trials preceded light trials.
In this case, only an increasing trend was observed, with no significant difference
detected (Supplementary Figure 3A, B $P = 2.9 \times 10^{-1}$, $t_5 = -1.16$, paired *t*-test, $n = 6$ rats,
Supplementary Table 2).

To assess whether the observed differences in cluster separability between light
and dark conditions could arise by chance, we performed a permutation test. Within
each rat, the darkness/light labels were randomly shuffled across trials while preserving
the number of trials per condition. For each permutation, silhouette scores were
recalculated from the layer embeddings of the trained network. This procedure was
repeated 1000 times to generate an empirical null distribution of cluster separability.
The observed silhouette scores for the dark condition consistently exceeded the 95th
percentile of the null distribution, confirming that the enhanced separability in darkness
reflects systematic differences in neural representations rather than random variation
(Supplementary Figure 4).

To elucidate which electrodes held the most information about the floor textures,
we performed occlusion analysis. The results revealed that electrodes in the forelimb
subregion contributed more to successful texture and lighting predictions (Figure 6D, E,
$P = 4.53 \times 10^{-6}$, $t_8 = -5.57$, Student's *t*-test, $n = 9$ rats), which aligns with our observation

that negative deflections in LFPs were largest in forelimb-targeting channels (Figure

2E). Further analysis using class activation maps from integrated gradients showed

spatiotemporal patterns of salient features unique to each experimental condition

(Figure 6F). In particular, the average attribution score in the forelimb subregion

displayed a temporal shift in the dark condition, implying an extended processing

window for texture information when visual cues are unavailable (Figure 6G,

Supplementary Figure 5).

Collectively, these results indicate that visual deprivation modifies population-

level activity in S1, yielding more distinctive representations of tactile stimuli. Such

reorganization could provide a neural substrate for enhanced tactile perception under

conditions of reduced or absent visual input.

Discussion

This study provides evidence that S1 undergoes rapid reorganization of tactile
representations when visual input is removed, even over a short period. Using high-
density LFP recordings and deep learning techniques, we show that the neural encoding
of tactile stimuli becomes more distinct under conditions of visual deprivation.
Specifically, when rats were exposed to darkness, texture representations in S1 were
more clearly distinguishable than under normal visual conditions. This demonstrates the
adaptability of S1 and its capacity to rapidly adjust to changing sensory contexts.

A key finding was the significant increase in silhouette scores when texture
representations were decoded in the dark. Silhouette scores, which quantify the
separability of neural features, were notably higher under dark conditions, suggesting
that visual deprivation sharpens the distinctions between tactile representations. While
motivational factors, such as water restriction, could potentially confound the results,
trial order reversals showed only a slight increase in silhouette scores during the dark
condition. If motivation alone were responsible for the enhanced discriminability,
silhouette scores should have been elevated during the later light trial in the reversed
sequence. This supports the conclusion that the reorganization in S1 is more likely a
direct result of visual deprivation rather than solely motivation or arousal. Although
differences in silhouette scores were not significant in the reversed trials, this
observation may reflect a lingering effect of prior visual deprivation, consistent with
previous studies in humans showing that even brief periods of visual deprivation, such
as blindfolding, can enhance tactile sensitivity. These findings suggest a robust and
lasting effect of visual deprivation on tactile processing.

Our results also highlight the somatotopic specificity of tactile encoding in S1.
The evoked potential in the forelimb subregion of S1 was significantly larger than in the
hindlimb subregion, reflecting the topographic organization of S1, where distinct
cortical areas process sensory inputs from different body parts (Ewert et al., 2008; Prsa

et al., 2019; Sur et al., 1980). Further, occlusion analysis revealed that forelimb

subregions were critical for distinguishing between tactile textures and lighting

conditions. This finding underscores the dominance of the forelimb subregion in

encoding tactile information, especially during the specific task of walking and

contacting different textures. Integrated gradient analysis confirmed the importance of

forelimb channels in encoding texture information and showed that the temporal

window of processing following forelimb contact was extended in the dark compared to

the light condition. Additionally, comparative analysis of the integrated gradient and

time-frequency maps revealed a negative correlation in the 50-60 Hz frequency range,

while a positive correlation was observed in the 80-90 Hz range. These findings suggest

that frequency-specific patterns of LFP activity in different conditions are closely linked

to the texture representations captured by the CNN model.

The extended temporal window in the dark condition suggests that visual

deprivation may enhance the retention of tactile information in S1. Previous studies

have shown that S1 neurons are involved in the short-term retention of tactile

information (Zhou and Fuster, 2000, 1997, 1996), and prolonged neuronal firing in

higher sensory areas may contribute to this sustained activity (Esmaeili and Diamond,

2019; Leavitt et al., 2017). Our findings suggest that the dark condition could lead to

more prolonged neural representations of tactile stimuli. Whether this reflects a

compensatory mechanism for the absence of visual input or a correlation with

heightened tactile sensitivity requires further exploration.

Environmental illumination can also influence the arousal state, which in turn

affects cortical dynamics as reflected in LFPs. Dark environments typically promote

exploratory behavior and increase arousal, modulating cortical activity through

irradiance- and cone-opponent–dependent pathways to arousal circuits like the locus

coeruleus and basal forebrain (Tamayo et al., 2023). Arousal in S1 enhances both

baseline activity and stimulus encoding, with neuromodulatory drive sharpening

temporal precision (Eggermann et al., 2014; Poulet and Petersen, 2008; Shimaoka et al.,

2018). These state-dependent effects may account for the enhanced separability of

tactile representations observed in the dark, despite minimal differences in average

evoked amplitude (Lee et al., 2020; McGinley et al., 2015; Shimaoka et al., 2018; Vinck

et al., 2015). The shift toward a desynchronized, high-arousal state in darkness likely

reduces low-frequency shared variability and boosts fast-timescale signal components,

improving the separability of neural clusters in a high-dimensional LFP space.

Moreover, neuromodulatory engagement during heightened arousal can extend effective

integration windows and enhance gain in task-relevant networks, consistent with the

longer temporal window for accurate predictions seen in forelimb channels.

These findings highlight the limitations of traditional analysis methods, such as

amplitude-based metrics and event-related potentials, which often fail to capture subtle,

higher-dimensional features in neural signals (Saxena and Cunningham, 2019; D. L.

Yamins and DiCarlo, 2016; D. L. K. Yamins and DiCarlo, 2016). In contrast, our deep-

learning approach revealed fine-grained spatiotemporal patterns in LFPs, showcasing

enhanced texture-specific separability under visual deprivation. This demonstrates the

power of advanced computational tools to uncover previously inaccessible shifts in

sensory coding, with broad applicability to other high-dimensional neural datasets, such

as those from multi-electrode arrays in freely behaving animals. However, it's important

to note that each model was trained specifically for each rat, and inter-animal

generalization remains a challenge, due to differences in electrode placement and

individual brain structure. Despite this limitation, the model's ability to extract complex,

high-dimensional patterns from the dataset remains evident, and the results show

significant progress in detecting sensory changes.

While fMRI studies have demonstrated tactile enhancement and corresponding

neural reorganization during short-term visual deprivation (Facchini and Aglioti, 2003;

Kauffman et al., 2002; Pascual-Leone and Hamilton, 2001b), the low temporal and

spatial resolution of fMRI limits its ability to capture detailed changes in neural

representations. Our study, through LFP recordings, provides a more nuanced picture of

the population-level activity underlying tactile perception enhancement in the absence

of visual input. Future studies could extend this work to assess whether rats can

discriminate textures more accurately in the dark and explore the direct relationship

between neural coding and perceptual performance.

From an evolutionary perspective, enhanced tactile sensitivity in low-visibility

environments provides an adaptive advantage. Many species, including rodents, rely on

somatosensation for navigation and foraging when visual information is scarce. The

ability to rapidly enhance tactile processing in such conditions could aid in efficient

resource acquisition and predator detection. Given the influence of arousal on tactile

sensitivity (Lee et al., 2020; Shimaoka et al., 2018), the dynamic nature of sensory

processing becomes evident—S1 does not merely respond passively to tactile input but

actively adapts its encoding strategies based on available sensory cues.

Several limitations warrant further exploration. First, if arousal contributes to the

effects observed in S1 during light–dark transitions, we predict that arousal indices,

such as pupil or whisking activity, should correlate with LFP spectral composition and

decoding accuracy on a trial-by-trial basis (McGinley et al., 2015; Reimer et al., 2014;

Shimaoka et al., 2018). Second, manipulating neuromodulatory tone through

optogenetic activation of arousal circuits should modulate the separability of texture

representations, independent of illumination (Eggermann et al., 2014; Harris and Thiele,

2011). Future experiments integrating real-time arousal measures and LFP analyses will

be crucial in disentangling the contributions of arousal versus light–dark effects on

cross-modal reorganization in S1. Additionally, while rats serve as a powerful model for

sensory processing, their neural architecture may not fully capture the complexities of

human perception. Extending this paradigm to assess perceptual changes in tactile

acuity will further clarify the relationship between neural coding and perceptual

performance.

In conclusion, our study provides new insights into the brain's remarkable

ability to reorganize its sensory processing in response to changes in sensory input. By

demonstrating that visual deprivation rapidly reconfigures tactile processing in S1, we

highlight the brain's flexibility in adapting to environmental changes. This work

underscores the potential of combining behavioral paradigms, LFP recordings, and deep

learning techniques to understand the dynamic and adaptive nature of sensory coding,

paving the way for future research in multisensory interactions.

Data availability

The data that support the findings of this study are available from the corresponding

author upon request.

Code availability

Custom code generated during this study for data analysis are available at

https://github.com/UT-yakusaku/Yamashiro-eLife-2024.

References

Abraira VE, Ginty DD. 2013. The sensory neurons of touch. *Neuron* **79**:618–639.
Bensmaia SJ, Denchev PV, Dammann JF 3rd, Craig JC, Hsiao SS. 2008. The representation of
stimulus orientation in the early stages of somatosensory processing. *J Neurosci* **28**:776–
786.
Bola Ł, Siuda-Krzywicka K, Paplińska M, Sumera E, Zimmermann M, Jednoróg K, Marchewka
A, Szwed M. 2017. Structural reorganization of the early visual cortex following Braille
training in sighted adults. *Sci Rep* **7**:17448.
Boroojerdi B, Bushara KO, Corwell B, Immisch I, Battaglia F, Muellbacher W, Cohen LG.
2000. Enhanced excitability of the human visual cortex induced by short-term light
deprivation. *Cereb Cortex* **10**:529–534.
Bulusu V, Lazar L. 2024. Crossmodal associations between naturally occurring tactile and
sound textures. *Perception* **53**:219–239.
Burton H. 2003. Visual cortex activity in early and late blind people. *J Neurosci* **23**:4005–4011.
Bushnell MC, Duncan GH, Hofbauer RK, Ha B, Chen JI, Carrier B. 1999. Pain perception: is
there a role for primary somatosensory cortex? *Proc Natl Acad Sci U S A* **96**:7705–7709.
Butler JS, Foxe JJ, Fiebelkorn IC, Mercier MR, Molholm S. 2012. Multisensory representation
of frequency across audition and touch: high density electrical mapping reveals early
sensory-perceptual coupling. *J Neurosci* **32**:15338–15344.
Delhaye BP, Long KH, Bensmaia SJ. 2018. Neural basis of touch and proprioception in primate
cortex. *Compr Physiol* **8**:1575–1602.
Dijkerman HC, de Haan EHF. 2007. Somatosensory processing subserving perception and
action: Dissociations, interactions, and integration. *Behav Brain Sci* **30**:224–230.
Di Plinio S, Perrucci MG, Aleman A, Ebisch SJH. 2020. I am Me: Brain systems integrate and
segregate to establish a multidimensional sense of self. *Neuroimage* **205**:116284.
Driver J, Spence C. 2000. Multisensory perception: beyond modularity and convergence. *Curr*
Biol **10**:R731–5.
Eggermann E, Kremer Y, Crochet S, Petersen CCH. 2014. Cholinergic signals in mouse barrel
cortex during active whisker sensing. *Cell Rep* **9**:1654–1660.
Eimer M, Forster B. 2003. Modulations of early somatosensory ERP components by transient
and sustained spatial attention. *Exp Brain Res* **151**:24–31.
Esmaeili V, Diamond ME. 2019. Neuronal correlates of tactile working memory in prefrontal
and vibrissal somatosensory cortex. *Cell Rep* **27**:3167–3181.e5.
Ewert TAS, Vahle-Hinz C, Engel AK. 2008. High-frequency whisker vibration is encoded by
phase-locked responses of neurons in the rat's barrel cortex. *J Neurosci* **28**:5359–5368.
Facchini S, Aglioti SM. 2003. Short term light deprivation increases tactile spatial acuity in
humans. *Neurology* **60**:1998–1999.
Goldreich D, Kanics IM. 2003. Tactile acuity is enhanced in blindness. *J Neurosci* **23**:3439–
3445.
Goodwin AW, Wheat HE. 2004. Sensory signals in neural populations underlying tactile
perception and manipulation. *Annu Rev Neurosci* **27**:53–77.
Harris KD, Thiele A. 2011. Cortical State and Attention. *Nature Reviews Neuroscience* **12**:509.
He K, Zhang X, Ren S, Sun J. 2015. Deep residual learning for image recognition. *arXiv*
[csCV].
Hopkins K, Kass SJ, Blalock LD, Brill JC. 2017. Effectiveness of auditory and tactile
crossmodal cues in a dual-task visual and auditory scenario. *Ergonomics* **60**:692–700.

Karlen SJ, Kahn DM, Krubitzer L. 2006. Early blindness results in abnormal corticocortical and
thalamocortical connections. *Neuroscience* **142**:843–858.
Kauffman T, Théoret H, Pascual-Leone A. 2002. Braille character discrimination in blindfolded
human subjects. *Neuroreport* **13**:571–574.
Koch KW, Fuster JM. 1989. Unit activity in monkey parietal cortex related to haptic perception
and temporary memory. *Exp Brain Res* **76**:292–306.
Leavitt ML, Mendoza-Halliday D, Martinez-Trujillo JC. 2017. Sustained activity encoding
working memories: Not fully distributed. *Trends Neurosci* **40**:328–346.
Lee CCY, Kheradpezhouh E, Diamond ME, Arabzadeh E. 2020. State-dependent changes in
perception and coding in the mouse somatosensory cortex. *Cell Rep* **32**:108197.
Luna R, Hernández A, Brody CD, Romo R. 2005. Neural codes for perceptual discrimination in
primary somatosensory cortex. *Nat Neurosci* **8**:1210–1219.
McGinley MJ, David SV, McCormick DA. 2015. Cortical Membrane Potential Signature of
Optimal States for Sensory Signal Detection. *Neuron* **87**:179–192.
Merabet LB, Hamilton R, Schlaug G, Swisher JD, Kiriakopoulos ET, Pitskel NB, Kauffman T,
Pascual-Leone A. 2008. Rapid and reversible recruitment of early visual cortex for touch.
PLoS One **3**:e3046.
Moulton EA, Pendse G, Becerra LR, Borsook D. 2012. BOLD responses in somatosensory
cortices better reflect heat sensation than pain. *J Neurosci* **32**:6024–6031.
Nikbakht N, Tafreshiha A, Zoccolan D, Diamond ME. 2018. Supralinear and Supramodal
Integration of Visual and Tactile Signals in Rats: Psychophysics and Neuronal
Mechanisms. *Neuron* **97**:626–639.e8.
Norman JF, Bartholomew AN. 2011. Blindness enhances tactile acuity and haptic 3-D shape
discrimination. *Atten Percept Psychophys* **73**:2323–2331.
Pascual-Leone A, Hamilton R. 2001a. The metamodal organization of the brain. *Prog Brain Res*
134:427–445.
Pascual-Leone A, Hamilton R. 2001b. The metamodal organization of the brain. *Prog Brain Res*
134:427–445.
Piras F, Vecchio D, Ciullo V, Gili T, Banaj N, Piras F, Spalletta G. 2020. Sense of external
agency is sustained by multisensory functional integration in the somatosensory cortex.
Hum Brain Mapp **41**:4024–4040.
Poulet JFA, Petersen CCH. 2008. Internal brain state regulates membrane potential synchrony
in barrel cortex of behaving mice. *Nature* **454**:881–885.
Prsa M, Morandell K, Cuenu G, Huber D. 2019. Feature-selective encoding of substrate
vibrations in the forelimb somatosensory cortex. *Nature* **567**:384–388.
Reimer J, Froudarakis E, Cadwell CR, Yatsenko D, Denfield GH, Tolias AS. 2014. Pupil
fluctuations track fast switching of cortical states during quiet wakefulness. *Neuron*
84:355–362.
Sadato N, Pascual-Leone A, Grafman J, Ibañez V, Deiber MP, Dold G, Hallett M. 1996.
Activation of the primary visual cortex by Braille reading in blind subjects. *Nature*
380:526–528.
Salinas E, Hernandez A, Zainos A, Romo R. 2000. Periodicity and firing rate as candidate
neural codes for the frequency of vibrotactile stimuli. *J Neurosci* **20**:5503–5515.
Saxena S, Cunningham JP. 2019. Towards the neural population doctrine. *Curr Opin Neurobiol*
55:103–111.
Schürmann M, Caetano G, Jousmäki V, Hari R. 2004. Hands help hearing: facilitatory
audiotactile interaction at low sound-intensity levels. *J Acoust Soc Am* **115**:830–832.

Serino A. 2019. Peripersonal space (PPS) as a multisensory interface between the individual and
the environment, defining the space of the self. *Neurosci Biobehav Rev* **99**:138–159.
Shimaoka D, Harris KD, Carandini M. 2018. Effects of arousal on mouse sensory cortex depend
on modality. *Cell Rep* **22**:3160–3167.
Sugiyama S, Kinukawa T, Takeuchi N, Nishihara M, Shioiri T, Inui K. 2019. Tactile Cross-
Modal Acceleration Effects on Auditory Steady-State Response. *Front Integr Neurosci*
13:496317.
Sundararajan M, Taly A, Yan Q. 2017. Axiomatic attribution for deep networks. *arXiv [csLG]*.
Sur M, Merzenich MM, Kaas JH. 1980. Magnification, receptive-field area, and "hypercolumn"
size in areas 3b and 1 of somatosensory cortex in owl monkeys. *J Neurophysiol* **44**:295–
311.
Tamayo E, Mouland JW, Lucas RJ, Brown TM. 2023. Regulation of mouse exploratory
behaviour by irradiance and cone-opponent signals. *BMC Biol* **21**:178.
Van Boven RW, Hamilton RH, Kauffman T, Keenan JP, Pascual-Leone A. 2000. Tactile spatial
resolution in blind braille readers. *Neurology* **54**:2230–2236.
Vinck M, Batista-Brito R, Knoblich U, Cardin JA. 2015. Arousal and locomotion make distinct
contributions to cortical activity patterns and visual encoding. *Neuron* **86**:740–754.
Wong M, Gnanakumaran V, Goldreich D. 2011. Tactile spatial acuity enhancement in
blindness: evidence for experience-dependent mechanisms. *J Neurosci* **31**:7028–7037.
Yamashiro K, Aoki M, Matsumoto N, Ikegaya Y. 2020. Polyherbal formulation enhancing
cerebral slow waves in sleeping rats. *Biol Pharm Bull* **43**:1356–1360.
Yamashiro K, Ikegaya Y, Matsumoto N. 2024. Automatic detection of foot-strike onsets in a
rhythmic forelimb movement. *Neurosci Res* **206**:41–50.
Yamins DL, DiCarlo JJ. 2016. Eight open questions in the computational modeling of higher
sensory cortex. *Curr Opin Neurobiol* **37**:114–120.
Yamins DLK, DiCarlo JJ. 2016. Using goal-driven deep learning models to understand sensory
cortex. *Nat Neurosci* **19**:356–365.
Zeiler MD, Fergus R. 2013. Visualizing and understanding convolutional networks. *arXiv*
[csCV].
Zhou YD, Fuster JM. 2000. Visuo-tactile cross-modal associations in cortical somatosensory
cells. *Proc Natl Acad Sci U S A* **97**:9777–9782.
Zhou YD, Fuster JM. 1997. Neuronal activity of somatosensory cortex in a cross-modal (visuo-
haptic) memory task. *Exp Brain Res* **116**:551–555.
Zhou YD, Fuster JM. 1996. Mnemonic neuronal activity in somatosensory cortex. *Proc Natl*
Acad Sci U S A **93**:10533–10537.
Ziegler K, Folkard R, Gonzalez AJ, Burghardt J, Antharvedi-Goda S, Martin-Cortecero J,
Isaías-Camacho E, Kaushalya S, Tan LL, Kuner T, Acuna C, Kuner R, Mease RA, Groh A.
2023. Primary somatosensory cortex bidirectionally modulates sensory gain and
nociceptive behavior in a layer-specific manner. *Nat Commun* **14**:2999.

Figures

Figure 1. Behavioral paradigm and limb movement assessment with concurrent
LFP recordings.

(**A**) A diagram of the disk-shaped treadmill used in the experiment. One half of the disk
is covered with #80 sandpaper, and the other half with #2000 sandpaper. (**B**) A frame
from the video capturing a walking rat from a left-side perspective. (**C**) The motivation
scheme. The rats were water-deprived prior to the experiments. A water port was
coupled with the movement of the treadmill so that when the rat walked on the treadmill,
the water would come out. This way, the rats were always motivated to walk during the
whole session. (**D**) The experimental protocol, where each rat walked for 10 minutes in
light (50 lx) and then for 10 minutes in darkness (0 lx). (**E**) An example trajectory of the
elbow and wrist joints from one session, plotted with the shoulder joint fixed in the
coordinate space (shoulder: *yellow*, elbow: *green*, wrist: *cyan*). (**F**) A schematic

illustrating the forelimb and hindlimb subregions of S1. (**G**) A custom 32-channel

electrode array used to record LFPs from these subregions.

Figure 2. Comparison of gait parameters across textures and environmental conditions.

(**A**) Swing phase *vs.* stance phase, illustrated with video frames (*left*: swing, *right*: stance). (**B**) Normalized swing duration measured for each rat under different textures (smooth *vs.* rough) and environmental condition (light *vs.* dark). Light orange and dark orange correspond to the light conditions (smooth, rough), while light blue and dark blue correspond to the dark conditions (smooth, rough). There were no significant differences among trial conditions. (**C–E**) Stance duration, stride length, and footstrike speed, respectively, under the same conditions as in B. None of these parameters differed significantly across texture types or lighting conditions.

Figure 3. LFP recordings in rat S1 during walking.

(**A**) A single representative LFP trace aligned to a forelimb contact. (**B**) An example of an averaged LFP trace from one session, aligned to forelimb contact. (**C**) The electrode montage and averaged LFP at each electrode. *Left*: The electrode montage showing all 32 recording sites. *Right*: Averaged LFP signals aligned to forelimb contacts with the floor, shown for each electrode depicted in the left panel. (**D**) Comparison of amplitudes

between hindlimb and forelimb subregions, aggregated across all 11 rats. $P = 1.89 \times 10^{-3}$,

$t_{10} = -2.2$, paired t-test, $n = 11$ rats. (**E**) Comparison of averaged amplitudes within the

same trial for different floor textures and environmental conditions. $P > 0.05$, one-way

analysis of variance (ANOVA) followed by Tukey☐Kramer post hoc test, $n = 149, 149,$

107 and 107 trials for smooth-light (light-orange), rough-light (dark-orange), smooth-

dark (light-blue), and rough-dark (dark-blue), respectively. (**F**) Cumulative probability

distributions of mean amplitude from each session, compared across different textures.

$P = 4.84 \times 10^{-1}, 8.35 \times 10^{-1}, D = 1.03 \times 10^{-1}$ and 7.54×10^{-2} for light and dark environments

respectively, two-sample Kolmogorov☐Smirnov test, $n = 149$ and 107 trials from 11

rats for light and dark, respectively. (**G**) Cumulative probability distributions of mean

amplitude from each session, compared across light and dark environments. $P =$

$1.05 \times 10^{-1}, 1.83 \times 10^{-1}, D = 0.14$ and 0.149 for smooth and rough textures respectively,

two-sample Kolmogorov☐Smirnov test, $n = 149$ and 107 trials from 11 rats for light

and dark, respectively. Abbreviations: ERP, event-related potential; LFP, local field

potential; S1, primary somatosensory cortex.

Figure 4. LFP characteristics across textures and lighting

(**A**) Example averaged LFP traces from one session, aligned to forelimb contact: light–

smooth, light–rough, dark–smooth, dark–rough (left to right). (**B**) Channel correlation

difference map (light − dark) computed from average LFP waveforms (Pearson's *r*).

Higher values indicate stronger channel-wise correlations in light relative to dark. (**C**)

Correlation matrices shown separately for each texture and lighting condition. Matrices

exhibit similar within–S1-subregion structure, with no clear texture- or light-dependent

differences. (**D**) Example time–frequency (wavelet) spectrogram of LFP power from

one session, aligned to forelimb contact. Dotted line indicates the cone of influence.

Color scales represent wavelet coefficient magnitude (C) and corresponding power (D).

Figure 5. Model-based prediction of texture and environmental conditions from LFP.

(**A**) The deep learning model architecture. The LFP input is processed through two parallel pathways for macro- and micro-scale feature extraction, followed by residual blocks that feed into two output heads for floor texture (Smooth *vs.* Rough) and the environmental condition (Light *vs.* Dark). (**B**) Training performance for a single representative rat. The left graph shows accuracy curves for texture (blue) and lighting (yellow), and the right graph shows the loss curves. (**C**) Testing performance for the same rat. The model exhibits good generalization, as accuracy increases and loss decreases on held-out data. (**D**) Confusion matrix for texture classification for all rats. Values above chance on the diagonal indicate successful texture prediction. Note that all values in the same row add up to 1. (**E**) Same as D, but for environmental conditions. (**F**) Combined confusion matrix for texture and trial predictions. The model performs

769   well on both tasks across all rats. Abbreviations: avgpool, average pooling layer; conv,

770   convolutional layer; maxpool, max pooling layer; LFP, local field potential.

Figure 6. Neural representations are more distinct in dark environments than in light environments

(**A**) A 912-dimensional feature vector is extracted from the layer preceding the final output. (**B**) A scatter plot of these features from one rat shows individual LFP segments (aligned to forelimb contact). Light orange and dark orange correspond to the light conditions (smooth, rough), while light blue and dark blue correspond to the dark conditions (smooth, rough). (**C**) Silhouette scores across all nine rats, showing that the dark condition yields higher scores and thus more distinct neural representations. $P = 2.31 \times 10^{-2}$, $t_8 = -2.8$, paired t-test, $n = 9$ rats. (**D**) A pseudo-color map based on occlusion analysis, illustrating the contribution of each electrode in the forelimb and hindlimb subregions. Hotter regions indicate higher importance for the model's predictions. (**E**)

Forelimb channels exhibit higher occlusion sensitivity than hindlimb channels, highlighting the forelimb's dominant role when the foot contacts the floor. $P = 4.53 \times 10^{-6}$, $t_{16} = -5.57$, Student's *t*-test, $n = 9$ rats. (**F**) Class activation maps generated via integrated gradients highlight key input features responsible for accurate model predictions of texture and environmental conditions. Attribution scores show each feature's impact on the model's output relative to a reference baseline: high positive scores denote features that strongly affect the predicted class. The onset of forelimb contact is aligned to time zero. (**G**) Attribution scores averaged over forelimb electrodes for floor texture (*left*) and the environmental conditions (*right*). A temporal lag in the dark condition suggests an extended processing window for floor texture when visual cues are absent.