Glycosylated IgG antibodies contribute to the recovery of haemorrhagic fever with renal syndrome patients

Chuansong Quan[1,2,3†], Lu Wang[3,4†], Jiming Gao[5†], Yaoni Li[6†], Xiaoyu Xu[7†], Houqiang Li[3], Zixuan Gao[7], Wenxu Ruan[7], Hongzhi Liu[3], Qian Li[3], Weijia Xing[3], Liqiong Zhao[6], Michael J. Carr[8,9], Weifeng Shi[10,11*], Haifeng Hou[3*]

[1]Key Laboratory of Emerging Infectious Diseases in Universities of Shandong, Shandong First Medical University & Shandong Academy of Medical Sciences, Ji'nan, 250117, China

[2]The Second Affiliated Hospital of Shandong First Medical University, Taian, 271000, China

[3]School of Public Health, Shandong First Medical University & Shandong Academy of Medical Sciences, Ji'nan, 250117, China

[4]Medical Records and Statistics Management Office, Tengzhou Central People's Hospital, Zaozhuang, 277500, China

[5]Department of Neurobiology and Physiology, School of Clinical and Basic Medicine, Shandong First Medical University& Shandong Academy of Medical Sciences, Ji'nan, 250117, China

[6]Baoji Central Hospital, Baoji, 721008, China

[7]School of Life Sciences, Shandong First Medical University & Shandong Academy of Medical Sciences, Taian, 271016, China.

[8]National Virus Reference Laboratory, School of Medicine, University College Dublin, D04 E1W1, Ireland.

[9]International Collaboration Unit, International Institute for Zoonosis Control, Hokkaido University, N20 W10 Kita-ku, Sapporo 001-0020, Japan.

[10]Ruijin Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai,

200025, China.

[11]Shanghai Institute of Virology, Shanghai Jiao Tong University School of Medicine,

Shanghai, 200025, China

*Correspondence:

Haifeng Hou: hfhou@sdfmu.edu.cn

Weifeng Shi: shiwf@ioz.ac.cn

†These authors contributed equally to this work

Abstract

Haemorrhagic fever with renal syndrome (HFRS) is a fatal disease caused by Hantaan virus (HTNV) infection. Humoral immunity is essential for effective viral clearance, however, the glycosylation characteristics of immunoglobulin G (IgG) in HFRS patients is not well known. Peripheral blood mononuclear cells from HFRS patients were obtained for B subset analysis using scRNA-seq and flow cytometry. HTNV specific IgG antibody titers were detected by enzyme-linked immunosorbent assay, and IgG glycosylation was analyzed by ultra-performance liquid chromatography. The proportions of the antibody-secreting memory (ASM) B cells and plasmablasts (PB) were significantly expanded among acute HFRS patients. We discovered significantly increased fucosylated IgG and decreased bisecting N-acetylglucosamine during the convalescent phase of HTNV infection. Meanwhile, positive correlations were observed between ASM subsets and galactosylation/sialylation in the IgG Fc region, and between PB subsets and sialylation. Notably, the glycosylation-related genes, such as *RPN1 and RPN2*, were primarily expressed differentially in the ASM and PB subclusters which were enriched in the N-glycosylation modifications of proteins through asparagine. Our findings indicated that IgG N-glycosylation may play a crucial role in combating HTNV infection and contributing to clinical recovery, which provided new insights for optimizing glycoengineered therapeutic antibodies.

Keywords: Haemorrhagic fever with renal syndrome (HFRS); Hantaan virus (HTNV); B cell; Immunoglobulin G (IgG); Glycosylation; Inflammation

Introduction

The causative agent of haemorrhagic fever with renal syndrome (HFRS) is Hantaan

virus, which belongs to the family *Hantaviridae* and is primarily transmitted by rodents

(Zheng et al., 2019). China is the most heavily affected country by HFRS, accounting

for >90% of global cases, and a total of 32,462 cases of HFRS were reported in China

during 2019–2022, with a fatality rate ranging from 0.1 to 15% (R. Li et al., 2023)

(https://www.chinacdc.cn).

Hantaviruses are tri-segmented, single-stranded, negative-sense RNA viruses, whose

genomes consist of three regions: large (L), medium (M), and small (S). The

glycoproteins Gn and Gc, encoded by the M segment, can infect target cells - primarily

vascular endothelial cells - via β3 integrin receptors (Pizarro et al., 2019).

Simultaneously, they could also infect other cell types, such as mononuclear

macrophages and dendritic cells, leading to systemic viral infection. Although

hantavirus replication is thought to occur primarily in the vascular endothelium without

direct cytopathic effects, a plethora of innate immune cells mediate host antiviral

defenses. These include natural killer cells, neutrophils, monocytes, and macrophages,

together with pattern recognition receptors (PRRs), interferons (IFNs), antiviral

proteins, and complement activation, e.g., via the pentraxin 3 (PTX3) pathway, which

can exacerbate HFRS disease progression leading to immunopathological damage

through cytokine/chemokine production, cytoskeletal rearrangements in endothelial

cells, ultimately amplifying vascular dysfunction (Tariq & Kim, 2022). Rapid and

effective humoral immune responses, however, such as neutralizing antibody responses

targeting the glycoproteins Gn/Gc, contribute to rapid recovery from HFRS and are

critical for protection from severe disease (Engdahl & Crowe, 2020; Li et al., 2020).

Immunoglobulin G (IgG) N-linked glycosylation mediates critical functions

modulating antiviral immunity during viral infection. Changes in the conserved N-

linked glycan Asn297 in the Fc region of IgG typically by fucosylation, galactosylation,

or sialylation can alter antibody effector function. A reduction in core fucosylation

decreases IgG binding to NK cell FcγRIIIa promotes antibody-dependent cellular

cytotoxicity (ADCC) necessary for clearance of viruses, including SARS-CoV-2,

dengue and HIV-1 whereas sialylation can attenuate immune responses resulting in

immune evasion (Ash et al., 2022; Haslund-Gourley et al., 2024; Hou et al., 2021; Wang

et al., 2017). Changes in IgG and other protein N-linked glycosylation profiles therefore

shape virus-host interactions and disease progression.

Importantly, there have not been prior studies specifically examining plasma IgG N-

glycome profiles derived from chromatographic peak data in HFRS patients,

particularly in relation to seroconversion status. This gap in our knowledge motivated

our systematic investigation of both total and virus-specific IgG glycosylation

dynamics during acute infection. In the present study, we combined scRNA-seq and

flow cytometry to reveal the phenotypes of B cell responses during Hantaan virus

(HTNV) infection, and explored the transcriptomic features of reactive B cell subsets.

A total of 166 HFRS patients, including 65 paired HFRS samples, were employed to

profile the IgG-Fc glycosylation pattern and to investigate the potential regulatory role

of IgG-Fc glycosylation in HFRS pathogenesis.

Results

B cell compositional characteristics in HFRS patients

To characterize the humoral immune profiles in HFRS patients, we enrolled 166

suspected HTNV-infected patients who were admitted to Baoji Central Hospital in

Shaanxi Province, China, between October 2019 and January 2022. Among them, 65

met the inclusion criteria and were included in the study (Figure 1). We identified a

total of eight cell subpopulations, including subsets of CD4 T cells, CD8 T cells, CD14

monocytes, NK cells, B cells, platelets, endothelial progenitor cells, and red blood cells

(Figure 2—figure supplement 1A and B). The proportions of these cell types were

comparable between acute HFRS patients and healthy controls, with no significant

batch effects observed (Figure 2—figure supplement 1C). CD4 T cells (31.4%) were

the predominant subset prior to HTNV infection, whereas CD8 T cells (44%) became

the dominant population post-infection. Notably, the proportions of NK cells (15.9% to

5.6%, $p<0.001$) and platelets (3.1% to 0.6%, $p<0.01$) significantly decreased, whereas

the level of CD14 monocytes (15.9% to 32.3%, $p<0.05$) nearly doubled. The overall

proportions of B cells, however, did not exhibit significant changes (Figure 2—figure

supplement 1D and E).

To precisely elucidate the role of B cell subsets in anti-HTNV infection, we classified

B cells into eight subsets, including antibody-secreting memory B cells (ASM), double-

negative B cells (DN), intermediate memory B cells (IM), marginal zone-like B cells

(MZB), naive B cells (naive B), plasmablasts (PB), quiescent resting memory B cells

(RM), and exhausted tissue-like memory B cells (TLM) (Figure 2A and B, and Figure

2—figure supplement 1F). We discovered that significant expansion of ASM, PB, and

RM cell populations in acute HFRS patients compared to healthy controls,

accompanied by contraction of DN, IM, and naive B cell compartments post-infection

(Figure 2C). In healthy controls, naive B cells (42.87%), DN B cells (26.65%), and

TLM B cells (16.13%) constituted the majority of the B cell population. However, ASM

(30.96%), PB cells (11.12%) and RM B cells (2.88%), became the dominant subsets

after HTNV infection (Figure 2D and Supplementary file 1).

To validate the accuracy of our single-cell subpopulation analysis, we employed flow

cytometry to quantify the B cell subsets in acute and convalescent HFRS patients

(Figure 2E). The results demonstrated that ASM (39.48% vs. 24.57%, p<0.001) and PB

(23.97% vs. 2.61%, p <0.001) subsets were significantly amplified in the acute phase

compared to the convalescent phase (Figure 2F). Conversely, the number of naive B

cells (32.78% vs. 48.92%, p=0.003) and CSM cells (18.65% vs. 27.15%, p=0.008)

recovered rapidly in the convalescent stage (Figure 2F), consistent with the

observations at the transcriptome level. No significant differences were observed for

other B cell subtypes between the acute and convalescent stages (Figure 2—figure

supplement 2). These results demonstrated that several B cell subpopulations were

effectively activated following HTNV infection.

Dynamic characteristics of the IgG N-glycome in HFRS patients

HFRS patients were further stratified into three age groups (≤44, 45-59, and ≥60

years). IgG glycosylation was analyzed by ultra-performance liquid chromatography.

As shown in Supplementary file 2, no significant differences in IgG glycosylation were

observed between age groups, except for galactosylation (p=0.001). Similarly, no sex-

based differences in IgG-Fc glycosylation were noted (Supplementary file 3). However,

the levels of IgG with bisecting GlcNAc (15.72% vs. 14.51%, p<0.001), galactosylated

IgG (74.26% vs. 71.15%, p<0.001), and sialylated IgG (22.44% vs. 21.62%, p<0.001)

were significantly higher in the acute phase compared to the convalescent phase.

Conversely, fucosylated IgG levels (94.29% vs. 94.89%, p<0.001) were lower in the

acute phase (Supplementary file 4), indicating that IgG N-glycosylation plays a critical

role in the anti-infection process. Multivariate linear regression was employed to

mitigate potential confounding by genetic and environmental factors in the glycomics

analysis. While no significant associations were observed for most glycan models

(fucosylation, p=0.526; bisecting GlcNAc, p=0.069; and sialylation, p=0.058), we

discovered sex showed a potentially influential effect on galactosylation (p=0.001)

(Supplementary files 5-8). These results suggest that while most glycan features appear

unaffected by the examined covariates, galactosylation may be subject to sex-specific

biological regulation.

Among the 130 paired blood samples from 65 patients, 38 (29.2%) were negative for

HTNV nucleocapsid protein-specific IgG antibodies, and 92 (70.8%) were seropositive.

For the 24 patients with both seronegative and seropositive samples, levels of bisecting

GlcNAc (16.16% vs. 14.54%, p<0.001), galactosylation (74.12% vs. 70.94%, p=0.006),

and sialylation (22.92% vs. 21.80%, p=0.008) were higher in the seronegative period,

whereas that of fucosylation was lower (94.36% vs. 94.74%, p= 0.024) in this period

(Figure 3A). Notably, bisecting GlcNAc and sialylated IgG levels decreased as HTNV

nucleocapsid protein (NP)-specific antibody titers increased (Figure 3B), highlighting

a potential link between antibody levels and IgG-Fc glycosylation.

The IgG N-glycome in HFRS patients with seroconversion

A four-fold or greater increase in HTNV NP-specific antibody titers usually indicates

a protective humoral immune response during the acute phase (Onwuchekwa et al.,

2023). We observed a significant increase in the Fc fucosylation (94.18% vs. 94.48%,

p=0.044) and a decrease in bisecting GlcNAc (16.41% vs. 14.71%, p=0.001) post the

four-fold antibody titer increase compared to the baseline. However, no significant

differences were observed in galactosylation and sialylation (Figure 3C), suggesting

that fucosylation and bisecting GlcNAc may play key roles in anti-HTNV infection and

recovery.

The potential source of the IgG N-glycome during HTNV infection

Positive correlations were observed between the ASM subsets and both

galactosylation (p=0.017, r_s=0.418) and sialylation (p=0.008, r_s=0.458) in the antibody

Fc region, as well as between the PB subsets and sialylation (p=0.036, r_s=0.372) (Figure

4A-C). Therefore, we speculated that galactosylated antibodies may primarily originate from the ASM subsets, while sialylated antibodies may derive from the ASM and PB subsets.

To investigate the process of glycosylated antibody production secreted by the ASM and PB subsets, we identified differentially expressed genes (DEGs) among the eight B cell subpopulations. Notably, the most DEGs were observed in the ASM (470), naive (312), and TLM (333) B subpopulations, respectively (Figure 4—figure supplement 1A and Supplementary file 9). Specifically, among all the upregulated genes, the expressions of glycosylation-related, mitochondrial respiratory-related, and peptide chain elongation-related genes among the ASM, PB, and RM populations were elevated after HTNV infection, including *RPN2*, *MT-ND6*, and *RPS26* (Figure 4—figure supplement 1B and Supplementary file 10). On the contrary, the upregulated genes in the naive, DN, and IM populations were related to anti-inflammatory and immunosuppressive effects, inhibition of the antioxidative function, response to environmental stress, and B-cell growth and development, such as *TSC22D3*, *TXNIP*, *DUSP1*, and *KLF6*, whereas the downregulated genes played an important role in the inflammatory response, cell motility, and antigen presentation, including *LTB*, *ACTG1*, and *HLA-DRB5* (Figure 4—figure supplement 1C and Supplementary file 11).

Unexpectedly, we observed significant changes in the expression of glycosylation-related genes (GRGs) in the ASM, PB, and RM subclusters during HTNV infection compared to healthy controls. *RPN1* and *RPN2* were prominently upregulated, which encoded subunits of the oligosaccharyltransferase complex (Figure 4D). This implied that the process of glycosylation during HTNV infection was rapidly activated among the ASM and PB subclusters. In the ASM subcluster, the fold change of GRGs from all four glycosylation modification-related genes, aside from *RPN1* and *RPN2*, was similar,

suggesting that ASM may undergo complex and diverse glycosylation modification

processes.

To enhance our understanding of the transcriptomic characteristics of GRG-high-

expressing PB and ASM subclusters, we conducted a comprehensive pathway

enrichment analysis and identified significant enrichment of pathways related to

glycosylation modifications, including the endoplasmic reticulum protein-containing

complex, rough endoplasmic reticulum, and protein N-linked glycosylation via

asparagine (Figure 4E). Additionally, gene sets associated with glycosylation

modifications showed positive regulation of these pathways (p=0.002, Figure 4F).

These results indicated that glycosylation modifications in the ASM, PB, and RM

subpopulations were dynamically regulated during the acute phase of HFRS, likely to

satisfy the amplified anti-HTNV antibody requirement and accelerate viral clearance.

Discussion

IgG glycosylation is a key regulator of inflammatory responses for infectious

diseases, acting as a molecular switch between anti-inflammatory and pro-

inflammatory effects upon antigenic challenge. One of the most significant findings of

the present study is the elevation in IgG fucosylation and reduction in bisecting GlcNAc

levels at the time of a four-fold increases in the HTNV NP-specific antibody titers.

Afucosylation is known to enhance antibody-dependent cellular cytotoxicity (ADCC)

by increasing the affinity of IgG-Fc for the FcγRIIIa receptor on NK cells, leading to

heightened production of inflammatory cytokines such as IL-1β, IL-6, IFN-γ, and TNF-

α (Larsen et al., 2021; Lu, Suscovich, Fortune, & Alter, 2018; Wieczorek, Braicu,

Oliveira-Ferrer, Sehouli, & Blanchard, 2020). Increase in bisecting GlcNAc linked to

Fc induces enhanced affinity for the FcγIIIa receptor and enhanced ADCC function of

the antibodies (Giron et al., 2020; Hou et al., 2021; Irvine & Alter, 2020). However,

rather than directly influencing the Fc receptor binding, this modification enhances antibody functionality through afucosylation, which subsequently increases the binding affinity to the FcgRIIIa receptor (Ackerman et al., 2018; Jennewein & Alter, 2017). Our findings provide evidence that the dynamic alterations in glycosylation patterns, characterized by increased fucosylation and decreased bisecting GlcNAc levels during viral infection, are involved in disease pathogenesis and clinical recovery, suggesting their dual utility as both diagnostic biomarkers and promising therapeutic targets.

We also observed elevated galactosylation levels in the acute phase compared to the convalescent phase, a phenomenon also reported in SARS-CoV-2 infections but not in chronic diseases and cancers (Hou et al., 2021; Liu et al., 2018). Increased galactosylation may expand the distance between CH2 domains, exposing key amino acids for FcγRIIIa binding and enhancing ADCC functions (Houde, Peng, Berkowitz, & Engen, 2010). Simultaneously, the increase in the ASM subset, which positively correlated with galactosylation, may reflect polyclonal activation of memory B cells driven by pattern recognition receptors and cytokines, leading to the production of galactosylated antibodies. Similarly, sialylation levels were higher in the acute patients, potentially linked to the expansion of the ASM and PB subsets. In this study, the PB subset was significantly elevated in the acute phase and positively correlated with sialylation, suggesting that rapidly proliferating plasmablasts may contribute to the secretion of sialylated antibodies; however, the mechanisms behind this require further investigation.

Antibody-related glycogenes are significantly activated following Hantaan virus infection. We noted that ribophorin I and II (RPN1 and RPN2) were significantly upregulated in the ASM/IM/PB/RM subsets after Hantaan virus infection, which linked the high mannose oligosaccharides with asparagine residues found in the Asn-X-

Ser/Thr consensus motif (Hwang et al., 2025). We speculate that they continuously attach the synthesized glycan chains to the constant region of antibodies during antibody synthesis. Similarly, fucosyltransferase 8 (FUT8) in the ASM subset, catalyzing the alpha1-2, alpha1-3, and alpha1-4 fucose addition (Wang & Ravetch, 2019; Yang et al., 2015), was downregulated in the mRNA translation, and the levels of fucosylated antibodies were naturally lower in the acute HFRS patients. Meanwhile, the beta-1,4-galactosyltransferase (beta4GalT) gene expression was significantly elevated in the ASM subpopulation during the acute phase, which also correlated with increased levels of galactosylated antibodies in serum (Wang & Ravetch, 2019). However, we did not observe significant upward changes in sialyltransferase mRNA expression in the acute HFRS patients, similar with the finding from severe COVID-19 cohorts (Haslund-Gourley et al., 2024). The neuraminidase 1 (NEU1) gene is strikingly upregulated and may potentially explain the decreased sialylation on the secreted HTNV-specific IgG antibodies during convalescence. Overall, the glycosylation of immunoglobulin G is regulated by a large network of B-cell glycogenes during HTNV infection.

Despite establishing a well-characterized patient cohort and performing systematic IgG glycosylation profiling based on HTNV NP antibody status, this study has several noteworthy limitations. Most notably, while preliminary comparisons suggested similar patterns between virus-specific and total IgG N-glycome, our total plasma IgG analysis may have introduced confounding factors in the observed associations. This methodological constraint could potentially affect the interpretation of certain disease-specific glycosylation signatures.

Our findings demonstrate that the IgG-Fc glycosylation plays a pivotal role in HTNV infection and rapid recovery. Fucosylated oligosaccharides and the absence of bisecting

GlcNAc are associated with favorable clinical outcomes. Importantly, specific B cell subsets, particularly ASMs and PBs, upregulated the secretion of galactosylated and sialylated antibodies, respectively. These findings would deepen our understanding of antibody glycosylation in antivirus immunity and provide a foundation for future studies aimed at optimizing glycoengineered therapeutic antibodies.

Materials and Methods

Study participants

Clinical specimens were collected from HFRS patients who were hospitalized in Baoji Central Hospital between October 2019 and January 2022. Patients were categorized into four clinical subtypes (mild, moderate, severe, and critical) based on the diagnostic criteria for HFRS issued by the Ministry of Health (Ma et al., 2015). This study was approved by the ethics committee of the Shandong First Medical University & Shandong Academy of Medical Sciences (R201937). Written informed consent was obtained from each participant or their guardians.

The clinical course of HFRS is grouped into acute (febrile, hypotensive, and oliguric stages) and convalescent (diuretic and convalescent stages) phases. The acute phase was defined as within 12 days of illness onset, and the convalescent phase was defined as a period of illness lasting 13 days or longer (Tang et al., 2019; Zhang et al., 2022). The earliest sample was selected if there were multiple blood samples available in the acute phase and the last available sample before discharge was selected if there were multiple blood samples in the convalescent phase.

Single-Cell RNA Sequencing and public database

Flow cytometry antibody staining was conducted as previously described (Chakraborty et al., 2022). Dead cell and red blood cell were removed using the BD FACSAria™ III cell sorter, and peripheral blood mononuclear cells (PBMCs) were

collected for single-cell sequencing. The library construction was conducted as

previously described (Jin et al., 2024).

We obtained single-cell transcriptome sequencing data from the Gene Expression

Omnibus (GEO) database (GSE161354), containing six HFRS patients and two healthy

volunteers, as well as healthy control samples from the Genome Sequence Archive

(GSA) database (HRA000203), matched with patient demographic information.

Single-Cell Data Analysis

The FastQC software was used to evaluate the data obtained to ensure the quality of

the raw sequencing data. The raw data were mapped to the human reference genome

(GRCh38, https://cf.10xgenomics.com/supp/cell-exp/refdata-gex-GRCh38-2020-A.tar.

gz) using Cell Ranger. The Seurat R package (Version 4.4.0) was utilized for the

merging and clustering of single-cell data. The Harmony R package and the anchor

module of Seurat were used to remove batch effects between samples and groups for

cell clustering. t-distributed Stochastic Neighbor Embedding (t-SNE) and Uniform

Manifold Approximation and Projection (UMAP) were employed for dimensionality

reduction and visualization of individual cells.

Cell Type Annotation

UMAP and t-SNE were used to reduce the dimension of all cells and to cluster them

in a two-dimensional space based on shared features. The Single R (Version 2.2.0)

package was used to infer independently the cell origin of each single cell based on a

reference transcriptome dataset of pure cell types, for unbiased cell type identification

from single-cell RNA sequencing data. Subsequently, specific highly-expressed genes

were combined with those derived from review of the literature for manual annotation

to ultimately determine cell types. Classic biomarkers for specific cell types were used

to identify cells in different clusters.

Differential Gene Identification and Functional Analysis

The *FindMarkers()* function in the Seurat package was used to identify DEGs between different cell groups, with criteria of $|log2FC| > 0.25$ and p-value<0.05, pct.1>0.25. The ClusterProfiler R package was utilized for GO, KEGG, and GSEA enrichment analyses, and the ggplot2 and enrichplot R packages were used for visualization.

Measurement of HTNV specific antibodies

Levels of serum IgM and IgG antibodies to HTNV NP were assessed through the enzyme-linked immunosorbent assay (ELISA) kit (Wantai BioPharm, Beijing, China) in accordance with the manufacturer's instructions.

Analysis of IgG glycans

The process of isolating, labelling, purifying, and analyzing plasma IgG was performed using previously established methods (Hou et al., 2019; Pengcheng Liu, 2023). Briefly, the diluted plasma samples were transferred onto a 96-well protein G monolithic plate (BIA Separations, Slovenia) for the isolation of IgG. The isolated IgG was eluted with 1 mL of 0.1 M formic acid and was immediately neutralized with 170 µL of 1M ammonium bicarbonate.

The released N-glycans were labelled with 2-aminobenzamide (2-AB) and were then purified from a mixture of 100% acetonitrile and ultrapure water in a 1:1 ratio (v/v). This was then analyzed by hydrophilic interaction liquid chromatography using ultra-performance liquid chromatography (HILIC-UPLC; Walters Corporation, Milford, MA) (Hou et al., 2019). As previously reported, the chromatograms were separated into 24 IgG glycan peaks (GPs) (Menni et al., 2018).

Our study incorporated both biological and technical replicates to ensure a robust glycomic profiling analysis. Specifically, we analyzed paired acute/convalescent-phase

samples from 65 confirmed HFRS patients to assess inter-individual biological variability, while technical reproducibility was validated through comparison with standard chromatographic peak plots (Vučković et al., 2016). This dual-replicate strategy enabled a comprehensive evaluation of both biological heterogeneity and assay precision, and the coefficient of variation for samples were below 16%.

Virus-specific B-cell detection

Virus-specific B-cell against HTNV were performed as previously described (Liechti & Roederer, 2019; Song et al., 2018). All antibodies used for flow cytometry were purchased from BioLegend or BD Biosciences. Antibody combination was as follows: CD3-FITC (cat, 300406), CD19-PerCP-Cy5.5 (cat, 561295), CD21-APC (cat, 559867), CD27-DAPI (cat, 562513), CD38-PE (cat, 555460), IgD-Amcyan (cat, 563034), IgG-PE-CY7 (cat, 561298), and IgM-BV605 (cat, 562977).

Statistical analysis

Categorical variables were compared using chi-square tests. Normality of numeric variables was assessed with the Kolmogorov-Smirnov test; non-normally distributed continuous variables were analyzed using non-parametric tests. The Benjamini - Hochberg (BH) method was used to adjust the raw p-values from DEG analysis, controlling the false discovery rate (FDR). Statistical analyses were performed with R software version 4.1.1 (R Core Team, New Zealand) and GraphPad Prism 8 (GraphPad Software, San Diego, CA), with a significance level of $\alpha = 0.05$ (two-tailed).

Funding

Funder	Grant reference number	Author
the Natural Science Foundation of Shandong Province	Grant numbers ZR2020QH133	Chuansong Quan
the Natural Science Foundation of Shandong Province	Grant numbers ZR2022MH082	Haifeng Hou

the Natural Science Foundation of Shandong Province	Grant numbers ZR2024MH235	Weijia Xing
the Natural Science Foundation of Shandong Province	Grant numbers ZR201911090028	Yaoni Li
Shaanxi Provincial Natural Science Basic Research Program of Shaanxi Province	Grant number 2023-JC-YB-741	Yaoni Li

Data availability

The raw RNAseq data used for this study is available at the BioProject database (the accession number: PRJNA1238574) under the digital object identifier: https://www.ncbi.nlm.nih.gov/bioproject/PRJNA1238574.

The following datasets were reanalysed:

Zhang H, Wang Y (2022) Increased CD4+CD8+ double positive T cells during Hantaan virus infection ID GSE161354. In the public domain at GEO https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE161354.

Zheng Y, Liu X (2020) A human circulating immune cell landscape in aging and COVID-19 ID HRA000203. In the public domain at GSA https://ngdc.cncb.ac.cn/gsa-human/browse/HRA000203.

Disclosure statement

No potential conflict of interest was reported by the author(s).

References

Zheng, Z., Wang, P., Wang, Z., Zhang, D., Wang, X., Zuo, S., & Li, X. (2019). The characteristics of current natural foci of hemorrhagic fever with renal syndrome in Shandong Province, China, 2012-2015. PLoS Negl Trop Dis, 13(5), e0007148. doi:10.1371/journal.pntd.0007148

Li, R., Sun, J., Chen, Y., Fan, X., Wang, X., Zhang, X., . . . Liu, Z. (2023). Clinical and laboratory features and factors predicting disease severity in pediatric patients with hemorrhagic fever with renal syndrome caused by Hantaan virus. J Med Virol, 95(1), e28339. doi:10.1002/jmv.28339

Pizarro, E., Navarrete, M., Mendez, C., Zaror, L., Mansilla, C., Tapia, M., . . . Rodríguez, E. M. (2019). Immunocytochemical and ultrastructural evidence supporting that Andes Hantavirus (ANDV) is transmitted person-to-person through the respiratory and/or salivary pathways. Front Microbiol, 10, 2992. doi:10.3389/fmicb.2019.02992

Tariq, M., & Kim, D. M. (2022). Hemorrhagic fever with renal syndrome: literature review, epidemiology, clinical picture and pathogenesis. Infect Chemother, 54(1), 1–19. doi:10.3947/ic.2021.0148

Engdahl, T. B., & Crowe, J. E., Jr. (2020). Humoral immunity to Hantavirus infection. mSphere, 5(4). doi:10.1128/mSphere.00482-20

Li, Y., Quan, C., Xing, W., Wang, P., Gao, J., Zhang, Z., . . . Shi, W. (2020). Rapid humoral immune responses are required for recovery from haemorrhagic fever with renal syndrome patients. Emerg Microbes Infect, 9(1), 2303–2314. doi:10.1080/22221751.2020.1830717

Ash, M. K., Bhimalli, P. P., Cho, B. K., Mattamana, B. B., Gambut, S., Tarhoni, I., . . . Schneider, J. R. (2022). Bulk IgG glycosylation predicts COVID-19 severity and vaccine antibody response. Cell Rep, 41(11), 111799.

doi:10.1016/j.celrep.2022.111799

Haslund-Gourley, B. S., Woloszczuk, K., Hou, J., Connors, J., Cusimano, G., Bell, M., . . . Comunale, M. A. (2024). IgM N-glycosylation correlates with COVID-19 severity and rate of complement deposition. Nat Commun, 15(1), 404. doi:10.1038/s41467-023-44211-0

Hou, H., Yang, H., Liu, P., Huang, C., Wang, M., Li, Y., . . . Wang, W. (2021). Profile of immunoglobulin G N-glycome in COVID-19 patients: a case-control study. Front Immunol, 12, 748566. doi:10.3389/fimmu.2021.748566

Wang, T. T., Sewatanon, J., Memoli, M. J., Wrammert, J., Bournazos, S., Bhaumik, S. K., . . . Ravetch, J. V. (2017). IgG antibodies to dengue enhanced for FcγRIIIA binding determine disease severity. Science, 355(6323), 395–398. doi:10.1126/science.aai8128

Onwuchekwa, C., Moreo, L. M., Menon, S., Machado, B., Curcio, D., Kalina, W., . . . Begier, E. (2023). Underascertainment of respiratory syncytial virus infection in adults due to diagnostic testing limitations: a systematic literature review and meta-analysis. J Infect Dis, 228(2), 173–184. doi:10.1093/infdis/jiad012

Larsen, M. D., de Graaf, E. L., Sonneveld, M. E., Plomp, H. R., Nouta, J., Hoepel, W., . . . Vidarsson, G. (2021). Afucosylated IgG characterizes enveloped viral responses and correlates with COVID-19 severity. Science, 371(6532). doi:10.1126/science.abc8378

Lu, L. L., Suscovich, T. J., Fortune, S. M., & Alter, G. (2018). Beyond binding: antibody effector functions in infectious diseases. Nat Rev Immunol, 18(1), 46–61. doi:10.1038/nri.2017.106

Wieczorek, M., Braicu, E. I., Oliveira-Ferrer, L., Sehouli, J., & Blanchard, V. (2020). Immunoglobulin G subclass-specific glycosylation changes in primary epithelial

ovarian cancer. Front Immunol, 11, 654. doi:10.3389/fimmu.2020.00654

Giron, L. B., Colomb, F., Papasavvas, E., Azzoni, L., Yin, X., Fair, M., . . . Abdel-Mohsen, M. (2020). Interferon-α alters host glycosylation machinery during treated HIV infection. EBioMedicine, 59, 102945. doi:10.1016/j.ebiom.2020.102945

Irvine, E. B., & Alter, G. (2020). Understanding the role of antibody glycosylation through the lens of severe viral and bacterial diseases. Glycobiology, 30(4), 241–253. doi:10.1093/glycob/cwaa018

Ackerman, M. E., Das, J., Pittala, S., Broge, T., Linde, C., Suscovich, T. J., . . . Alter, G. (2018). Route of immunization defines multiple mechanisms of vaccine-mediated protection against SIV. Nat Med, 24(10), 1590–1598. doi:10.1038/s41591-018-0161-0

Jennewein, M. F., & Alter, G. (2017). The immunoregulatory roles of antibody glycosylation. Trends Immunol, 38(5), 358–372. doi:10.1016/j.it.2017.02.004

Liu, D., Chu, X., Wang, H., Dong, J., Ge, S. Q., Zhao, Z. Y., . . . Wang, W. (2018). The changes of immunoglobulin G N-glycosylation in blood lipids and dyslipidaemia. J Transl Med, 16(1), 235. doi:10.1186/s12967-018-1616-2

Houde, D., Peng, Y., Berkowitz, S. A., & Engen, J. R. (2010). Post-translational modifications differentially affect IgG1 conformation and receptor binding. Mol Cell Proteomics, 9(8), 1716–1728. doi:10.1074/mcp.M900540-MCP200

Hwang, H. J., Kang, D., Shin, J., Jung, J., Ko, S., Jung, K. H., . . . Lee, J. S. (2025). Therapy-induced senescent cancer cells contribute to cancer progression by promoting ribophorin 1-dependent PD-L1 upregulation. Nat Commun, 16(1), 353. doi:10.1038/s41467-024-54132-1

Wang, T. T., & Ravetch, J. V. (2019). Functional diversification of IgGs through Fc glycosylation. J Clin Invest, 129(9), 3492–3498. doi:10.1172/jci130029

Yang, Z., Wang, S., Halim, A., Schulz, M. A., Frodin, M., Rahman, S. H., . . . Clausen,

H. (2015). Engineered CHO cells for production of diverse, homogeneous

glycoproteins. Nat Biotechnol, 33(8), 842–844. doi:10.1038/nbt.3280

Ma, Y., Yuan, B., Zhuang, R., Zhang, Y., Liu, B., Zhang, C., . . . Jin, B. (2015).

Hantaan virus infection induces both Th1 and ThGranzyme B+ cell immune responses

that associated with viral control and clinical outcome in humans. PLoS Pathog, 11(4),

e1004788. doi:10.1371/journal.ppat.1004788

Tang, K., Zhang, C., Zhang, Y., Zhang, Y., Du, H., Jin, B., & Ma, Y. (2019). Elevated

plasma interleukin 34 levels correlate with disease severity-reflecting parameters of

patients with haemorrhagic fever with renal syndrome. Infect Dis (Lond), 51(11-12),

847–853. doi:10.1080/23744235.2019.1672887

Zhang, H., Wang, Y., Ma, Y., Tang, K., Zhang, C., Wang, M., . . . Zhang, Y. (2022).

Increased CD4(+)CD8(+) double positive T cells during Hantaan virus infection.

Viruses, 14(10). doi:10.3390/v14102243

Chakraborty, S., Gonzalez, J. C., Sievers, B. L., Mallajosyula, V., Chakraborty, S.,

Dubey, M., . . . Wang, T. T. (2022). Early non-neutralizing, afucosylated antibody

responses are associated with COVID-19 severity. Sci Transl Med, 14(635), eabm7853.

doi:10.1126/scitranslmed.abm7853

Jin, C., Chen, Y., Wang, Y., Li, J., Liang, J., Zheng, S., . . . Yang, H. (2024). Single-

cell RNA sequencing reveals special basal cells and fibroblasts in idiopathic pulmonary

fibrosis. Sci Rep, 14(1), 15778. doi:10.1038/s41598-024-66947-5

Hou, H., Xu, X., Sun, F., Zhang, X., Dong, H., Wang, L., . . . Wang, Y. (2019).

Hyperuricemia is associated with immunoglobulin G N-glycosylation: a community-

based study of glycan biomarkers. Omics, 23(12), 660–667.

doi:10.1089/omi.2019.0004

Pengcheng Liu, X. W., Aishe Dun, Yutong Li, Houqiang Li, Lu Wang, Yichun Zhang,

Cancan Li, Jinxia Zhang, Xiaoyu Zhang, Lixing Ma f, Haifeng Hou. (2023). High-

throughput profiling of serological immunoglobulin G N-glycome as a noninvasive

biomarker of gastrointestinal cancers. Engineering, 26(7), 44–53.

doi:10.1016/j.eng.2023.02.008

Menni, C., Gudelj, I., Macdonald-Dunlop, E., Mangino, M., Zierer, J., Bešić, E., . . .

Valdes, A. M. (2018). Glycosylation profile of immunoglobulin G is cross-sectionally

associated with cardiovascular disease risk score and subclinical atherosclerosis in two

independent cohorts. Circ Res, 122(11), 1555–1564.

doi:10.1161/circresaha.117.312174

Vučković, F., Theodoratou, E., Thaçi, K., Timofeeva, M., Vojta, A., Štambuk, J., . . .

Lauc, G. (2016). IgG glycome in colorectal cancer. Clin Cancer Res, 22(12), 3078–

3086. doi:10.1158/1078-0432.Ccr-15-1867

Liechti, T., & Roederer, M. (2019). OMIP-051 - 28-color flow cytometry panel to

characterize B cells and myeloid cells. Cytometry A, 95(2), 150–155.

doi:10.1002/cyto.a.23689

Song, P., Zheng, N., Liu, Y., Tian, C., Wu, X., Ma, X., . . . Wu, Z. (2018). Deficient

humoral responses and disrupted B-cell immunity are associated with fatal SFTSV

infection. Nat Commun, 9(1), 3328. doi:10.1038/s41467-018-05746-9

    Figure legends

Figure 1. Flow chart of HFRS patient recruitment in Baoji Central Hospital in

Shaanxi province from November 2019 to January 2022.

Figure 2. Immunophenotypic remodeling in the B-cell subsets during acute HFRS.

(A) t-distributed stochastic neighbor embedding (t-SNE) plot showing antibody

secreting memory B cells (ASM), double negative (DN) B cells, intermediate memory

cells (IM), marginal zone-like cells (MZB), naive B cells, plasmablasts (PB), quiescent

resting memory cells (RM), and exhausted tissue-like memory B cells (TLM) of

PBMCs identified using an integrated and classification analysis. (B) t-SNE projection

of canonical markers, including CD19, CD27, CD38, IgD, and IgM. (C) Proportions of

the eight B cell subsets, colored by the healthy group (green) and HTNV groups (red).

Boxplot features: minimum box, 25th percentile; center, median; maximum box, 75th

percentile. (D) Frequency of the eight B cell subsets in between the healthy group (n =

8) and HTNV groups (n = 15). (E) FACS gating strategy for the measurement of the B-

cell subsets: activated memory B cells (CD21$^-$ CD27$^+$, AM), RM B cells (CD21$^+$

CD27$^+$), IM B cells (CD21$^+$ CD27$^-$), TLM B cells (CD21$^-$ CD27$^-$), naive B cells (CD27$^-$

IgD$^+$), MZB B cells (CD27$^+$ IgD$^+$), ASM B cells (CD27$^+$ IgD$^-$), DN B cells (CD27$^-$ IgD$^-$),

PB (CD38$^+$ CD27$^+$), and class-switched memory B cells (CD38$^-$ CD27$^+$, CSM),

respectively. (F) The proportion of ASM, CSM, naive B, and PB cells in CD19$^+$ B cells

in the acute and convalescent HFRS patients. Data are presented as mean ± SEM in

panels C and F. *p<0.05 (Student's t-test).

Figure 3. Glycosylation modification of antibodies associated with HTNV infection.

(A) Changes of different glycosylation types in both HTNV NP-specific IgG negative

and positive plasma from 24 HFRS patients. (B) Differential IgG glycosylation patterns

across antibody titer levels quantified by ELISA. (C) Comparison of different

glycosylation levels before and after the fourfold increase in the IgG antibody titers.

*$p<0.05$, **$p<0.01$, ***$p<0.001$. For paired samples, Wilcoxon Signed Rank Test was

used to assess the difference.

Figure 4. Glycosylation modifications of antibodies primarily deprived from

antibody-secreting and plasmablast subpopulations.

(A) The correlation between the proportion of PB cells and the sialylation level. (B)

The correlation between the proportion of ASM cells and the sialylation level. (C) The

correlation between the proportion of ASM cells and the galactosylation level. (D) Dot

plot shows the expression levels of glycosylation-related genes in the eight B cell

groups. The pink color represents mannose related genes, the green color represents N-

glycosylation-related genes, the orange color represents galactosylation-related genes,

the purple represents sialylation-related genes, and the red color represents fucosylation

related genes. (E) Enriched pathways in the plasmablast and ASM subsets by GO

enrichment analysis of the DEGs. (F) The GSEA map presents the enrichment score of

GRGs in N-Glycan biosynthesis.

Supplementary materials

Supplementary file 1. The proportion of different B cell subpopulations between healthy and acute HFRS groups.

Supplementary file 2. Difference in main IgG glycome features of HFRS patients between age groups.

Supplementary file 3. Difference in main IgG glycome features of HFRS patients between sex groups.

Supplementary file 4. Relative abundance (%) of the main IgG-Fc glycome features in HFRS patients.

Supplementary file 5. Results of multiple linear regression analysis for fucosylation.

APTT, activated partial thromboplastin time; MCHC, mean corpuscular hemoglobin concentration; NEUT, neutrophil; PLT, platelet; WBC, white blood cell; CRE, creatinine; Cys-C, cystatin C.

Supplementary file 6. Results of multiple linear regression analysis for bisecting GlcNAc.

APTT, activated partial thromboplastin time; MCHC, mean corpuscular hemoglobin concentration; NEUT, neutrophil; PLT, platelet; WBC, white blood cell; CRE, creatinine; Cys-C, cystatin C.

Supplementary file 7. Results of multiple linear regression analysis for galactosylation.

APTT, activated partial thromboplastin time; MCHC, mean corpuscular hemoglobin concentration; NEUT, neutrophil; PLT, platelet; WBC, white blood cell; CRE, creatinine; Cys-C, cystatin C.

Supplementary file 8. Results of multiple linear regression analysis for sialylation.

APTT, activated partial thromboplastin time; MCHC, mean corpuscular hemoglobin

concentration; NEUT, neutrophil; PLT, platelet; WBC, white blood cell; CRE,

creatinine; Cys-C, cystatin C.

Supplementary file 9. Differentially expressed genes among different B cell

subgroups in healthy and acute HFRS groups.

Supplementary file 10. Differential expression genes in the ASM, PB and RM B

cell subpopulations.

Supplementary file 11. Differential expression genes in the DN, IM and naive B

cell subpopulations.

Figure 2—figure supplement 1. Single-cell transcriptomes of PBMCs from

patients with HFRS.

(A) Uniform manifold approximation and projection (UMAP) presentation of the major

peripheral immune cell types in the PBMCs from 15 HFRS patients and 8 healthy

controls. (B) Dot plot shows the expression levels of canonical marker-related genes in

the different subpopulations. (C) UMAP presentation of the major peripheral immune

cell types among HFRS and healthy control groups. (D) Proportions of the eight cell

subsets, colored by the healthy group (green) and HTNV groups (red). Boxplot features:

minimum box, 25th percentile; center, median; maximum box, 75th percentile. (E)

Frequency of the eight cell subsets in between the healthy group (n = 8) and the HTNV

group (n = 15). (F) The UMAP plot shows the differentiation trajectories of different

cell types calculated by pseudotime analysis.

Figure 2—figure supplement 2. Dynamic analysis of the B-cell subsets in HFRS

patients.

(A) Expression of MZB in CD19$^+$ B cells. (B) Expression of DN in CD19$^+$ B cells. (C)

Expression of TLM in CD19$^+$ B cells. (D) Expression of AM in CD19$^+$ B cells. (E)

Expression of RM in CD19$^+$ B cells. (F) Expression of IM in CD19$^+$ B cells.

Figure 4—figure supplement 1. The differentially expressed genes and their

functional changes in the B cell subsets post HTNV infection.

(A) The volcano plot shows the differentially expressed genes of different B cell subsets

in acute HFRS patients. The red font represents the upregulation of the B cell

subpopulations, and the blue font represents the downregulation of the B cell

subpopulations. (B and C) The volcano plot displays the differentially expressed genes

in upregulation B cell subpopulations including antibody secreting memory B cells,

plasmablasts, and quiescent resting memory B cells (B), and downregulation B cell

subpopulations including naive B cells, double negative B cells, and intermediate

memory B cells (C).