Mosquito sex under lock and key

Leah Houri-Zeevi[1,2]*, Madison M. Walker[1,2], Jacopo Razzauti[1,3], Anurag Sharma[4], H. Amalia Pasolli[4], Leslie B. Vosshall[1,2,5]*

[1]Laboratory of Neurogenetics and Behavior, The Rockefeller University, New York, NY 10065 USA
[2]Howard Hughes Medical Institute, New York, NY 10065 USA
[3]Price Family Center for the Social Brain, The Rockefeller University, New York, NY 10065 USA
[4]Electron Microscopy Resource Center, The Rockefeller University, New York, NY, 10065 USA
[5]Kavli Neural Systems Institute, The Rockefeller University, New York, NY, USA 10065 USA
*Correspondence: Leah.Houri@gmail.com (L.H-Z.), leslie@rockefeller.edu (L.B.V.) (Lead Contact)

Summary

Female mosquitoes typically mate once in a lifetime, making this singular mating decision critically important for the female. Yet, mosquito mating has been historically viewed as male-guided, with the female exerting little control. To understand this contradiction, we investigated initiation of successful mating in mosquitoes. Here we show that females of two invasive mosquito species, the yellow fever mosquito (*Aedes aegypti*) and the Asian tiger mosquito (*Aedes albopictus*), actively control mating through a previously undescribed behavior of female genital tip elongation. This female genital response is triggered by a rapidly-evolving secondary male genital structure that together act as a "lock-and-key" mechanism. We identified important differences in the genital interactions of the two species and show that *Aedes albopictus* males, which have larger secondary male genital structures, can bypass the female genital response of *Aedes aegypti* females but not of females of their own species. Coupled with previous observations that *Aedes aegypti* females are effectively sterilized by such cross-species mating, the ability of male *Aedes albopictus* to "pick the lock" of heterospecific females provides a potential mechanism for observed local extinction of *Aedes aegypti* when they live in shared territory with *Aedes albopictus*. Our results redefine mosquito mating as a female-controlled process, with implications for the evolution of reproductive barriers and the population dynamics of these two globally invasive mosquito species.

Introduction

Female evaluation and eventual choice of sexual partners shape evolutionary trajectories at multiple levels. These powerful selective forces can drive rapid changes to genomes and genetic programs, nervous systems, morphologies, behaviors, and more[1]. The act of mating is the ultimate product of these processes, with various factors and mechanisms operating together to ensure correct mate choice and successful transfer of genetic material. Females who mate with an unfit male or a male of the wrong species endanger their lineage.

In the highly invasive yellow fever mosquito (*Aedes aegypti*) and Asian tiger mosquito (*Aedes albopictus*), female choice is a high-stakes decision because the female typically mates only once during her lifetime[2–5]. Following this single mating, the female actively stores and nourishes the sperm in internal sperm reservoirs known as *spermathecae*. Females draw on the spermathecae for sperm to fertilize eggs over multiple cycles of blood-feeding and egg production for the rest of their lives[6–8]. With a single and crucial mating event in a lifetime, one might expect elaborate and intricate courtship and mating rituals that would ensure the female correctly assesses and actively chooses her mate. Such female choice would allow her to block mating attempts from unsuitable males, particularly those of another species[9]. Instead, mating in *Aedes aegypti* mosquitoes typically lasts for 14 seconds and seemingly involves minimal courtship[10], with the female described to be mostly passive and exerting little if any control over the process[11]. In previous studies, females were observed to kick males as a form of resistance, but this behavior was shown by both virgin and previously mated

females[10]. It has been suggested that male and female mosquitoes harmonize their wingbeat as a form of courtship[12,13], but more recent work has challenged the existence of such a mechanism[14]. Together, these observations create a conceptual tension: the female's mating decision is crucial, but she is thought to have no control over the process.

Aedes aegypti and *Aedes albopictus* are globally invasive mosquito species that transmit dangerous arboviruses to hundreds of millions of people every year[15,16]. While *Aedes aegypti* was first introduced into the New World ~500 years ago likely through the Atlantic Slave Trade[17], *Aedes albopictus* only arrived in the last 50 years, and the two species now share broad territories in the Americas[18]. These two *Aedes* species are separated by ~35 million years of evolution[19], were geographically separated throughout most of the evolutionary history, and cross-species mating produces no viable progeny[20]. Despite their genetic incompatibility, cross-species mating is observed in wild populations, with an unexplained bias toward *Aedes albopictus* males mating with *Aedes aegypti* females rather than the reciprocal cross-species mating[21]. *Aedes albopictus* males that mate with *Aedes aegypti* females induce a remating block in those females that prevents them from mating with males of their own species, rendering them sterile[21,22]. This mode of reproductive interference by cross-species mating is termed *satyrization*[23–25]. Satyrization is hypothesized to contribute to the ability of *Aedes albopictus* to outcompete *Aedes aegypti* mosquitoes in the southeastern United States, where they share the same territory, leading to observed local extinction of *Aedes aegypti* [24,26]. The basis for the observed species asymmetry in cross-species mating frequency and satyrization is not known.

In this study, we investigated the behavioral sequence leading to mating in mosquitoes and the mechanisms controlling cross-species mating. We report the first observation of a genital behavior of the female, *female genital tip elongation*, that controls mating in mosquitoes. Only virgin females elongate their genital tips in response to male genital contact. In *Aedes aegypti*, the female genital response is triggered by rapidly evolving secondary male genital structures called *gonostyli*. Fine surgical manipulations of gonostyli dramatically reduced insemination success. Female genital tip elongation also controls mating in *Aedes albopictus* and we identified key differences in genital behaviors and interactions in the two species. Strikingly, we found that *Aedes albopictus* males can bypass female genital tip elongation of both virgin and previously mated *Aedes aegypti* females and initiate mating without the female genital response. These findings correct the long-standing view that the female mosquito plays a passive role in mating and suggest specific mechanisms for reproductive interference by an invasive species that "picks the lock" of the female of the other species.

Results

The female remating block is established rapidly
Aedes aegypti males pester females and attempt to copulate with them regardless of whether females are virgins or previously mated (**Figure S1A,B**). Because mosquito copulation is rapid[27], it is not possible to know if these high intensity pestering events and copulation attempts result in insemination, and whether a previously mated female will remate under laboratory conditions. To accurately assess the extent of remating, we generated two transgenic mosquito strains in which sperm express either green fluorescent protein (mNeonGreen) (**Figure S1C**) or red fluorescent protein (DsRed2) (**Figure S1D**) under control of the β2Tubulin promoter, which confers expression of the fluorescent proteins in mature sperm[28]. Males from these strains were indistinguishable from wild-type males in their ability to inseminate females (**Figure S1E,F**), and females mated to either transgenic strain produced the same number of eggs and resulted in equivalent larval hatch rates compared to females mated with wild-type males (**Figure S1G-H**).

Aedes aegypti female mosquitoes have three spermathecae, only two of which are typically filled with
sperm[7,8]. When we mated transgenic males to wild-type females, we observed rare instances of both
red fluorescent and green fluorescent sperm in the spermathecae, which we define as evidence of
remating (**Figure S1I**). 91% of females continually co-housed with a mixed group of males for 7 days
had a single sperm color in their spermathecae (**Figure S2A,B**), confirming prior observations that
most female *Aedes aegypti* females mate only once in a lifetime[3]. A limitation of this experiment is the
inability to distinguish single inseminations from multiple inseminations with sperm of the same color,
potentially leading to underestimation of the actual remating rates. To address this, we performed
experiments in which females were sequentially exposed to males with different sperm colors. We
removed the first males and replaced with second males of a different sperm color at different post-
mating intervals (**Figure S2C**). We found that 86% of females have an established remating block
two hours after mating, and this increased to 95.8% at 7 days post mating (**Figure S2D,E**). The
remating block was similar in younger and older females (**Figure S2D,E**). Our results show that most
female *Aedes aegypti* block remating rapidly after a single mating event, with a small number of
females capable of mating more than once.
115
Female genital behavior controls mating in *Aedes aegypti*
What is the behavioral basis of the remating block? Video recordings of 3-dimensional free flight
interactions (**Figure S3A-I** and **Video 1**[29]) and contact interaction of a string-tethered female and a
free flying male (**Figure S4A-K** and **Videos 2,3**[29]) revealed that previously mated females spent less
time in contact with males when in free flight (**Figure S3B-D**), and that previously mated females kick
the male more (**Figure S4F**). Although the male contacted the genitalia of the female numerous times
regardless of her mating status (**Figure S4E**), in most of the cases these genital contact events
remained superficial. A tighter genital connection was only observed in virgins[4], which we term *genital
interlocking*. Genital interlocking was observed only once per virgin female and was never observed
in previously mated females (**Figure S4G,H**). These observations point to genital interlocking as the
critical difference between receptive and unreceptive females, prompting us to investigate how this
process is controlled.
128
129 The female vaginal lips are hidden between the ventral side of the 8th sternite and the post-genital
130 plate that sits above the cercal plates (**Figure 1A** and **Figure S5A,B**). The penis-equivalent and
131 sperm transfer organ of the male, the *aedeagus*, is located superior to the secondary genital
132 structures and is sheathed by the *paraproct*[11,30] (**Figure 1B** and **Figure S5C,D**). For intromission, the
133 aedeagus emerges and fits into the female vaginal tract[11].
134
Prior studies of mosquito copulation described a multi-step process for the full formation of a genital
connection and intromission that was presumed to be entirely guided by the male[11,31,32]. Similar to our
results, prior work showed that mating attempts between males and previously mated females are
limited to a superficial genital connection[4]. To better understand the process that leads to successful
mating and genital interlocking, we video recorded male-female interactions in high resolution,
specifically focusing on the genital behavior of the male and the female. This was accomplished by
tethering the female to a metal pin to limit her movement and provide a consistent focal point for our
video recordings (**Figure 1C** and **Video 4**[29]). We recorded videos of virgin and previously mated
females interacting with free-flying males and scrutinized these video recordings frame-by-frame for
male-female genital interactions. With this approach, we discovered three major steps that lead to
successful mating in *Aedes aegypti* mosquitoes. First, the male genitalia contact the female genital tip
(**Figure 1D**). Second, in response to genital contact by the male, virgin females elongated their
genital tip (**Figure 1E**), a previously undescribed female mosquito genital behavior. Third, this act of
female genital response was followed by successful genital interlocking. Importantly, previously
mated females never elongated their genital tips in response to male genital contact and never
engaged in genital interlocking (**Figure 1F**).

151
Figure 1: **Female genital tip elongation controls mating in _Aedes aegypti_ mosquitoes**
(**A,B**). Anatomy of the (**A**) female and (**B**) male genital tips represented by schematics (left) and scanning electron
microscopy images (right). Magenta structures in schematics indicate (**A**) the female vaginal tract and bursa (sealed by the
vaginal lips at the bottom) and (**B**) the male sperm transfer organ, the aedeagus. White asterisks indicate the location of (**A**)
the female vaginal lips and (**B**) the male aedeagus, both of which are covered by other genital structures.
(**C**). Experimental setup for recording genital interactions of a pin-tethered female and free-flying males.
(**D-F**). Representative video still images of the tips of the male and female during the three major steps towards successful
mating in _Aedes aegypti_ mosquitoes and number of events per 5 minutes of video recordings of (**D**) male genital contact,
(**E**) female genital tip elongation, and (**F**) genital interlocking in virgin and previously mated female video recordings. n=12
virgin and n=9 previously mated video recordings with 1 female and 5 males. Time stamp represents time since male genital
contact. Median (thick dotted black line) and 25/75 quartiles (thin dotted black line) are indicated. ns=not significant (Mann-
Whitney test)
(**G**). Schematic showing quantification strategy for measuring genital behavior.
(**H**). Quantification of indicated genital behaviors in virgin and previously mated female video recordings. n=6 tracked video
recordings per condition. Average traces are shown with 95% confidence intervals.
See also **Figures S1-S5**.

To quantify these genital movements, we developed a manual tracking strategy for measuring genital
behavior in the four seconds after male genital contact was initiated. We extracted measurements of
the female dorsal genital tip length as a readout of female genital tip elongation length, and the
shared surface between male and female genitalia as a readout of genital interlocking (**Figure 1G-H**).
We found that the female elongated her genital tip to approximately twice its resting length, reaching
an average maximum length of 280 µm. Female genital tip elongation always preceded the genital
interlocking-associated increase in the connection surface between the male and the female,
reaching an average of 390 µm of connection during genital interlocking (**Figure 1H** and **Figure**
S5E,F). These dynamics were only observed in genital interactions of virgin females with males.
Previously mated females showed no change in either female dorsal genital tip length or male-female
shared surface (**Figure 1H** and **Video 5**[29]). Video recordings of sequential interactions of tethered
females with male groups with different fluorescent sperm colors revealed that two females that
elongated their genital tip twice were inseminated by sperm of both colors (**Figure S5G-K**). Together,
these observations strongly suggest that female mosquitoes actively control mating through
elongation of their genital tip.

Secondary male genital structures trigger the female genital response
Having shown that female genital tip elongation can follow male genital contact, we asked what about
this male contact triggers the female to elongate her genital tip. To answer this, we analyzed male-
female genital interactions prior to the female genital response (**Figure 2A**). We observed that male
genital contact is characterized by the male inserting hook-like genital structures known as the
gonostyli (singular: gonostylus) into the female genital tip (**Figure 2A,B**). The gonostyli are secondary
male genital structures that are not directly involved in intromission and insemination and rest on the
sides of the female cercal plates during copulation[11,32] (**Figure 2C**). Frame-by-frame examination of
videos of pin-tethered females and free-flying males showed that when the male was about to initiate
male genital contact, the gonostyli extended, slid into the ventral side of the female genital tip, and
vibrated (**Figure 2D** and **Video 4**[29]). In virgin females, this sequence was followed by female genital
tip elongation. This response was rapid, with females beginning their tip elongation an average of 784
milliseconds after initiation of male genital contact. In previously mated females, males vibrated their
gonostyli against the female genital tip for the duration of male genital contact, but the females did not
respond (see **Video 5**[29]).

To determine the role of gonostyli insertion prior to female genital tip elongation we tested the effect
of surgical gonostyli manipulation on male-female genital interactions with a pin-tethered female
(**Figure 2E** and **Figure S6A**). In all 14 video recordings, virgin females paired with mock-treated
males elongated their genital tips in response to male genital contact and genital interlocking was
achieved in all cases (**Figure 2F**). In contrast, in 11 of 14 video recordings of virgin females paired
with males with gonostyli removed, the females did not elongate their genital tips in response to male
genital contact and genital interlocking was never achieved (**Figure 2G**). In the three cases that
females elongated their genital tips in response to male genital contact with gonostyli-less males, the
male attempted to genitally interlock but failed. Of 14 video recordings of virgin females paired with
males with dislocated gonostyli, none elongated their genital tip in response to male genital contact
and genital interlocking was never attempted or achieved (**Figure 2H**).

Gonostyli are bilaterally symmetric curved cuticular structures with two arms and one small claw at
the tip of each arm. Using a free-flight setup, we tested how finer gonostyli manipulations affect
insemination success (**Figure 2I,J** and **Figure S6B-C**). Across replicates, we observed that mock-
treated males successfully inseminated 100% of the females they were paired with. Bilateral removal
or truncation of both sides of the gonostyli nearly eliminated insemination success, whereas removal
of only the two claws had a modest impact on insemination success (**Figure 2K**). Unilateral removal
of a gonostylus, truncation, or removal of a claw reduced insemination success 3-19% relative to

mock-treated males, indicating that for most females, one fully intact gonostylus is sufficient to induce

female genital tip elongation (**Figure 2K**).

Figure 2: **Gonostyli trigger female genital tip elongation and are required for successful insemination**
(**A**). Schematics and photographs of dorsal and lateral views of male genital contact.
(**B**). Scanning electron microscopy image of the male genitalia, with gonostyli pseudo-colored in green (top) and a
schematic of the gonostyli (bottom).
(**C**). Cryo-scanning electron microscopy images of mosquito genitalia during copulation. Gonostyli are pseudo-colored
green.
(**D**). The position and behavior of the gonostyli prior to and during male genital contact.
(**E**). A schematic of the experimental setup for testing the effect of gonostyli manipulation on genital behavior (top) and the
indications of the different genital interaction steps and measurements (bottom).
(**F-H**). Genital behavior and representative tracking dynamics in mating interactions with (**F**) mock-treated males or male
with (**G**) gonostyli removed or (**H**) dislocated (motionless). N=14 5-minute video recordings per condition.
(**I**). Schematic of gonostyli manipulations examined for insemination success.
(**J**). A schematic of the experimental setup for testing insemination success of mock treated and manipulated gonostyli
males.
(**K**). Insemination rates of females co-housed with mock treated and manipulated gonostyli males. Overall averaged percent
insemination success across trials and dot plots showing insemination success rates per trial are displayed. N=4 trials per
condition.
(**L**). A schematic of the experimental setup for testing insemination success in competition.
(**M**). Insemination success rates of intact (mock-treated) males or males with manipulated gonostyli when challenged by
competition with mock-treated males. Median (thick dotted black line) and 25/75 quartiles (thin dotted black line) are
indicated. Yellow triangle indicates theoretical insemination success rates calculated from a no-competition scenario (panel
K). N=3 trials with each condition tested twice per trial, switching between green and red sperm males as the manipulated
group.
See also **Figures S6-S7**.

To ask if gonostyli manipulation impacts insemination success in the context of competition with
males with intact gonostyli, we paired females with mock-treated males with one sperm color and
surgically manipulated males with a second sperm color (**Figure 2L**). Unsurprisingly, males with
bilateral gonostyli removal or truncation always lost to mock-treated males. Remarkably, males with
unilateral gonostylus removal who were ~80% successful in inseminating females without competition
saw their insemination success rate drop to an average of 11-15% in competition compared to the
theoretical insemination success rate of 40% calculated from a no-competition scenario (**Figure 2M**).
Even more dramatically, males lacking only the claws at the tip of their gonostyli who were 93%
successful in inseminating females in a non-competitive environment were outcompeted by mock-
treated males and had an average insemination success rate of 15% compared to the theoretical no-
competition success rate of 46.5% (**Figure 2M**). Together, these results suggest a critical role for the
gonostyli in triggering female receptivity via the female genital tip elongation response.
259
Initiation of successful mating in *Aedes albopictus* mosquitoes
The system of a male genital structure that promotes female receptivity and successful insemination
suggests a "lock and key" mechanism between male and female genitalia. In this framework, rapid
evolution of animal genitalia can safeguard the species barrier through structural compatibility and
incompatibility[33–35]. Indeed, male gonostyli are rapidly evolving structures that assume different
shapes across the mosquito tree of life[36] (**Figure S7**). This variation may be particularly important to
prevent cross-species mating in species that share the same habitat.
267
268 We asked how genital interactions during initiation of successful mating vary in another *Aedes*
269 species, *Aedes albopictus*, which has expanded globally over the past 50 years and now frequently
270 shares territories with *Aedes aegypti*[37]. While the structure of the female genital tip is similar between
271 the two species (**Figure S8A**), male *Aedes albopictus* gonostyli are significantly larger in both length
272 and thickness compared to male *Aedes aegypti* gonostyli (**Figure 3A** and **Figure S8B**).
273
We examined male-female genital interactions in *Aedes albopictus* using our high-resolution video
recording setup of a pin-tethered female and free-flying males (**Figure 3B**). We found that *Aedes*
albopictus mating follows the same behavioral sequence of male genital contact, female genital tip
elongation, and genital interlocking that we observed in *Aedes aegypti* (**Figure 3C-E** and **Video 6**[29]).

In *Aedes albopictus*, female genital tip elongation is required for genital interlocking and is absent in
previously mated females, exactly as we observed in *Aedes aegypti* (**Figure 3D,F**, **Figure S8C,D** and
Videos 6,7[29]). This demonstrates that female control over mating, and the behavioral pathway
leading to successful mating, is shared in mosquitoes with ~35 million years of evolutionary distance.
282

283
284
Figure 3: Comparison of genital interactions and behaviors in *Aedes albopictus* and *Aedes aegypti*
(**A**). Scanning electron microscopy images of the ventral side of *Aedes aegypti* and *Aedes albopictus* male genitalia.
Gonostyli are pseudo-colored orange (*Aedes aegypti*) and turquoise (*Aedes albopictus*)
(**B**). Experimental setup for recording of genital behavior of a pin-tethered female and free-flying males.
(**C-E**). Representative video still image of the tips of the male and female during the three major steps towards successful
mating in *Aedes albopictus* mosquitoes and number of events per 5 minutes of video recordings of (**C**) male genital contact,
(**D**) female genital tip elongation, and (**E**) genital interlocking – in virgin and previously mated female video recordings. N=12
video recordings with 1 female and 5 males per condition. Time stamp represents time since the depicted video still image
of male genital contact. Median (thick dotted black line) and 25/75 quartiles (thin dotted black line) are indicated. Data were
analyzed using the Mann-Whitney test. ns=not significant.
(**F**). Average quantification and 95% confidence interval of female dorsal genital tip length and male-female shared surface
length in genital interactions between males and virgin or previously mated female video recordings. n=6 video tracked per
condition.
(**G,H**). Unsolicited genital interlocking attempts by males in same-species mating in (**G**) *Aedes aegypti* and (**H**) *Aedes*
albopictus mosquitoes. Each bar represents the number of events per examined recording. Data are pooled from video
recordings of virgin and previously mated females. Data were analyzed using the Mann-Whitney test. ns=not significant.

Despite these similarities, we identified three fundamental differences in genital behaviors of the two species. First, during male genital contact, the large gonostyli of *Aedes albopictus* males were positioned dorsally and externally (**Figure 3C**) while the smaller *Aedes aegypti* gonostyli were positioned ventrally and internally to the female genital tip (**Figure 1D** and **2D**). Second, *Aedes albopictus* males tried to genitally interlock with females even in the absence of female genital tip elongation, a behavior that we observed only once in *Aedes aegypti* (**Figure 3G,H** and **Video 7**[29]). Third, while elongation of the female genital tip reached similar lengths in both species during the female genital response, only *Aedes albopictus* females displayed dramatic retraction of their genital tip at all steps preceding the genital tip elongation response that signals their receptivity (**Figure 3I**).

Lock and key functionality in cross-species genital interactions

Having shown that genital interactions of male and female mosquitoes differ between *Aedes aegypti* and *Aedes albopictus* during initiation of successful mating, we asked how these differences translate to genital interactions between the two species. Because these species mate at low frequency under laboratory conditions, we examined genital interactions using pin-tethered pairs to manually bring the male into close contact with the female (**Figure 4A**).

Same-species genital interactions of females and intact males followed the same dynamics that we observed before. In both species, the animals followed the behavioral sequence of male genital contact, female genital tip elongation, and genital interlocking (**Figure 4B,C** and **Videos 8,9**[29]). While the gonostyli trigger female genital tip elongation in *Aedes aegypti* (**Figure 2E-G** and **Figure 4D**), *Aedes albopictus* males without gonostyli were able to successfully induce female genital tip elongation in 10 of 12 pairs but were unable to proceed to genital interlocking (**Figure 4E**). This suggests that *Aedes albopictus* males use other unknown mechanisms to trigger female genital tip elongation in females of their own species. As observed before, previously mated females of both species never elongated their genital tip, and the pair did not proceed to genital interlocking (**Figure 4F,G**).

In cross species mating, when intact *Aedes aegypti* males with their small gonostyli were paired with virgin *Aedes albopictus* females with their retracted genital tip, females rarely elongated their genital tip (4 of 30 pairs) and genital interlocking was achieved in only one of these 4 pairs (**Figure 4H**). When *Aedes albopictus* males with their big gonostyli were paired with *Aedes aegypti* females with their extended genital tip, female genital tip elongation was a similarly rare event (4 of 30 pairs) although successful genital interlocking occurred in 3 of these pairs (**Figure 4I**). Strikingly, in 11 of 30 pairs, *Aedes albopictus* males bypassed *Aedes aegypti* female genital tip elongation and moved directly to genital interlocking (**Figure 4I**). This unique ability of male *Aedes albopictus* required intact gonostyli (**Figure 4J,K**).

Given that *Aedes albopictus* are able to bypass the female genital response, we reasoned that these males should be able to overcome the remating block of previously mated *Aedes aegypti* females. We therefore carried out cross-species mating experiments with previously mated females. We first found that *Aedes aegypti* males were unable to trigger female genital tip elongation or proceed to genital interlocking with previously mated *Aedes albopictus* females. We conclude that *Aedes aegypti* males are unable to remate with previously mated females from their own species or the other species (**Figure 4F,L**). In contrast, we found that *Aedes albopictus* males were able proceed to genital interlocking with previously mated *Aedes aegypti* females in half (11 of 22) of the interacting pairs (**Figure 4M,N** and **Video 10**[29]). In all cases, *Aedes albopictus* males bypassed female *Aedes*

353 *aegypti* genital tip elongation and proceeded immediately to cross-species genital interlocking with
354 these previously mated females. Male *Aedes albopictus* were never observed to bypass the female
355 genital response when attempting to mate with females of their own species (**Figure 3C-F**, **Figure**
356 **4G**, and **Figure S8C,D**).

357
358 Together, these results reveal a notable asymmetry in cross-species mating: *Aedes albopictus* males
359 can bypass the female response and break the remating block across species, potentially through
360 mechanical advantage conferred by their larger gonostyli and the extended genital tip of female
361 *Aedes aegypti*.
362

Figure 4: **Same-species and cross-species genital interactions in *Aedes aegypti* and *Aedes albopictus***
(**A**)**.** A schematic of the experimental setup and legend for pin-tethered male and female genital interaction video recordings.
(**B-G**)**.** Same-species genital interactions of (**B**,**C**) virgin females and intact males together with video still images of the
genital interactions, of (**D**,**E**) virgin females and males with their gonostyli removed, and of (**F**,**G**) previously mated females
and intact males.
(**H-M**)**.** Cross-species genital interactions of (**H**,**I**) virgin females and intact males together with video still images of the
genital interactions, of (**J**,**K**) virgin females and males with their gonostyli removed, and of (**L**,**M**) previously mated females
and intact males.
(**N**). Video still images of genital interactions between previously mated *Aedes aegypti* female and intact *Aedes albopictus*
male. Female genital tip elongation response is bypassed directly to genital interlocking.
The number of video recordings per condition is indicated on each panel.
375
376

Discussion

In this study we established that a female genital behavior controls mating in *Aedes aegypti* and
Aedes albopictus mosquitoes. In *Aedes aegypti*, the female genital response is triggered by the
gonostyli, rapidly evolving secondary male genital structures. We identified key differences in genital
interactions of *Aedes aegypti* and *Aedes albopictus* and found that *Aedes albopictus* males, with their
larger gonostyli, can bypass the genital response of *Aedes aegypti* females but not females of their
own species.
384
It is not known how the interaction of the male gonostyli with the female genital tip unlocks the female
genital response. Our observations support several possible mechanisms that are not mutually
exclusive. First, the impressive variability and rapid evolution of the gonostyli structure across
species[36,38] suggest that information might be relayed by the shape of the gonostylus itself. Second,
physical interactions of the gonostyli with female genitalia could provide an additional component to
the code. Prior to female genital tip elongation, the gonostyli are inserted into the female genital tip
and vibrate rapidly. It is possible that the vibration rhythm of the gonostyli is recognized by the female
via a yet unknown mechanosensory pathway and creates a unique code across species. Third, it is
also possible that gonostyli transmit contact chemosensory cues that together with its vibration
pattern could trigger the female genital response.
395
It will be interesting to understand how the female senses male gonostyli stimulation and the
regulation of the motor response governing genital tip elongation. The cercal and the post-genital
plates at the tip of the female genitalia form a triangular structure that is densely covered with
sensilla. These sensilla are described to be mostly mechanosensory[39]. The post-genital plate is
typically pushed by the gonostyli during male genital contact and is in contact with them during the
vibration phase. It is possible that genital plate sensilla have a role in recognizing gonostyli insertion
and vibration, or they may recognize additional cues including chemical substances. Alternatively,
and perhaps synergistically, the tips of the gonostyli contact a membranal-like tissue at the base of
the post-genital plate and near the vaginal lips. This tissue could be internally innervated to sense
and recognize gonostyli stimulation. Understanding the sensory components of the circuitry that
control female genital tip elongation would provide better understanding of the interactions between
male gonostyli and the female genital tip, as well as access to neuronal regulation of mating in female
mosquitoes, and how this regulation changes rapidly after a single mating event.
409
Our findings align with the "lock and key" theory for genital evolution. This framework, first formulated
in 1844[33,34], states that the rapid evolution of animal genitalia is a mechanism for safeguarding the
species barrier through structural compatibility and incompatibility. In our system, the gonostyli
function as the key to unlock the female genital response in *Aedes aegypti* to initiate successful
mating. While the male shows morphological variation of the gonostyli across the two species studied
here, the female side of the mechanism appears to be mostly behavioral. This system better matches
updated versions of the "lock and key" mechanism that focus on female sensory reception and

behavior over structural differences[35]. Understanding the interactions or structures that serve to stimulate the female genital responses in other systems, and how the rapid evolution of the male genitalia corresponds to the female sensory capacity and behavioral response across species, would provide fascinating insights into the evolution of mating systems, male-female co-evolution, and cross-species competition.

Over the past 50 years, *Aedes albopictus* mosquitoes have successfully invaded Europe and the Americas from their original geographic range in Asia. They readily adapt to new environments and outcompete other mosquito species in shared territories[18,40]. Our finding that male *Aedes albopictus* can bypass the female genital response in *Aedes aegypti* may provide a mechanism for the competitive and invasive capabilities of *Aedes albopictus.* Cross-species mating between *Aedes aegypti* and *Aedes albopictus* does not yield viable progeny and prevents *Aedes aegypti* females from mating again[23,24]. One interesting question for future investigation is whether *Aedes albopictus* males who mate with previously mated *Aedes aegypti* females impact the viability of same-species sperm stored in the spermathecae of the female from the prior mating with an *Aedes aegypti* male, potentially leading to another form of reproductive interference Finally, in some regions of the world *Aedes albopictus* are weak "satyrs"[41]. It would interesting whether variation in gonostyli shape and behavior might underlie this difference.

Aedes aegypti and *Aedes albopictus* mosquitoes transmit dangerous arboviruses such as dengue, Zika, and chikungunya to humans[42,43]. Reducing mosquito population sizes is a major public health priority to lower the burden of disease. Much of the effort to control mosquito population relies on mating as a point of intervention. The Sterile Insect Technique and Incompatible Insect Technique involve releasing millions of sterile or Wolbachia-infected males to targeted areas[44]. When these males successfully mate with wild females, the females produce no viable progeny and are effectively sterilized[45], reducing mosquito population size and viral transmission. In this study we describe the process leading to successful mating in *Aedes aegypti* and *Aedes albopictus* mosquitoes and the role of the female in gating it. Our findings provide a framework for enhancing mosquito control methods by accounting for the newly discovered role of the female in mating and the male's ability to induce this receptivity.

Resource availability
Lead contact
Correspondence and material requests should be addressed to L.H.Z. (Leah.Houri@gmail.com)

Materials availability
Plasmids generated in this study have been deposited to Addgene (#236085, #236086). All mosquito strains generated in this study are freely available on request.

Data and code availability
Data are publicly available at the paper's GitHub depository[46] and Zenodo record #15133603[29].

Acknowledgements
We thank members of the Vosshall Lab and Philipp Brand, Tomas Kay, Maximilian M.L. Knott, Orli Snir, and Itai A. Toker for discussion and comments on the manuscript; Gloria Gordon, Libby Mejia, and Melissa Dallesandro for strain maintenance, and Mona Liu for early contributions to the project. We thank Peter Atkinson, Rob Harrell, and Laura Harrington for providing the PB{3xP3-EGFP; β2Tubulin-dsRed2} plasmid. We thank the F.M Kirby Foundation for support in acquiring the JEOL JSM-IT500HR scanning electron microscope. This work was supported by a Junior Fellowship of the Simons Society of Fellows and the European Molecular Biology Organization long-term fellowship (EMBO ALTF 1103-2019) (L.H-Z.), and a graduate fellowship from the Price Family Center for the

468 Social Brain and from the Boehringer Ingelheim Fonds (J.R.). L.B.V. is an Investigator of the Howard
469 Hughes Medical Institute.
470

Author contributions

L.H-Z. carried out or developed and supervised all the experiments and data analyses in the paper
with additional contributions from co-authors. M.M.W. carried out the free flight and pin-tethered video
recordings and gonostyli manipulation experiments. J.R. developed the 3D tracking analysis for
mosquitoes in free flight and pose estimation analysis for string-tethered video recordings (**Extended
Data Fig. 3, 4i-k**). A.S. and H.A.P. processed the mosquito samples for scanning electron microscopy
and obtained the images. L.H-Z. and L.B.V. together conceived the study, designed the figures, and
wrote the paper with input from all authors.
479

Declaration of Interests

481 The authors declare no competing interests.
482

Open access statement

Supplemental Figures

Figure S1: **Transgenic fluorescent sperm strains to score insemination. Related to Figure 1.**
A. B. Mating attempts observed for 7 minutes in cages of males and virgin (**A**) or previously mated (**B**) females.
N=3 trials, 15 male and 15 female mosquitoes each. Black line indicates the median point.
C. D. Schematic of transgenic fluorescent sperm lines expressing mNeonGreen (**C**) or dsRed2 (**D**).
E. Animals used in F-H.

F. Schematic of experimental setup (left) and results (right) for testing insemination rates of transgenic males.
N=5 trials with 50 females and 50 males each. Median (thick dotted line) and 25/75 quartiles (thin dotted line)
are indicated. Data were analyzed using the Mann-Whitney test. ns=not significant.
G. Schematic of experimental setup for testing egg laying and hatch rate.
H. Results for egg laying and hatch rate, plotted separately (median – thick dotted line – and 25/27 quartiles –
thin dotted line – indicated). n=36-40 eggs/larvae obtained from single females were tested per condition. Data
were analyzed using the Mann-Whitney test. ns=not significant.
I. Example images of female spermathecae containing green-fluorescent sperm (top), red-fluorescent sperm
(middle), or both (bottom). Note green autofluorescence of spermathecal ducts in all samples.

508
509
Figure S2: **Quantifying remating rates in female *Aedes aegypti*. Related to Figure 1.**
A. Schematic of experimental setup for testing remating rates following 7-days of co-housing with males with
two different fluorescent sperm colors.
B. Scoring of single or double sperm color of spermathecae from 7-day co-housed females. Each circle is one
female. N=3 trials with 60 females each.
C. Schematic of experimental setup for testing remating rates following introduction of males with two different
fluorescent sperm colors at various post-mating intervals.
D. **E**. Scoring of single or double sperm color in spermathecae from 6-day-old (**D**) and 12-day old (**E**) females at
different post-mating intervals. Data are presented as total scored females across the 3 trials of the experiment.
Each circle is one female. N=3 trials with 40 females each.

521
Figure S3: **Examination of distance in free flight interactions of a mosquito pair. Related to Figure 1.**
A. Schematic of experimental setup for 3-dimensional recording of free flying male and female mosquitoes (1
each) (top) and example of the male and female trajectories over the course of the 7-minute recording (bottom).
B,C. Raster of distance between pairs of mosquitoes in (**B**) virgin female and (**C**) previously mated female
recordings. Virgin females #2 and #4 were not inseminated after 7 minutes. N=7 recordings per condition.
D. Summary of time spent in the different distance categories: contact (<1 cm), mid-range (1-5 cm), and far (5-
20 cm). Median (thick dotted line) and 25/75 quartiles (thin dotted line) are indicated. Data were analyzed using
the Mann-Whitney test. ns=not significant.
E, F. Distance analysis of pairs of (**E**) virgin females or (**F**) males. N=5 (**E**) and N=4 recordings (**F**).
G. Summary of time spent in the different distance categories: contact (<1 cm), mid-range (1-5 cm), and far (5-
20 cm). Median (thick dotted line) and 25/75 quartiles (thin dotted line) are indicated. Data were analyzed using
the Mann-Whitney test. ns=not significant.
H,I Summary of total distance flown by (**h**) females and (**i**) males across all recordings. Median (thick dotted
black line) and 25/75 quartiles (thin dotted black line) are indicated on violin plots. N=7 animals plotted for virgin
and previously mated recordings. N=10 (**h**) and N=8 (**i**). Data were analyzed using the Mann-Whitney test.
ns=not significant.

538
539
Figure S4: **Examination of male-female behavioral interactions in string-tethered female and free-flying**
male recordings. Related to Figure 1.
A. Experimental setup and a video still image of a string-tethered female recording, paired with a free flying
male. N=25 recordings per condition.
B. Behavioral categories scored in this assay. Ethogram represents data from virgin recording #8. Note that
these data are included in the ethograms in panels i-n below.
C-G. Heatmaps and violin plots of each scored behavior and a summary of events per minute, average event
duration, and proportion of total time per behavior in recordings of virgin and previously mated female. Median

(thick dotted line) and 25/75 quartiles (thin dotted line) are indicated. Heatmap is sorted by time until first genital
contact event. Each row in the heatmap represents an individual recording, repeated per behavior: (**C**) female
flight, (**D**) male-female contact, (**E**) genital contact, (**F**) female kicks male, and (**G**) genital interlocking. Median
(thick dotted line) and 25/75 quartiles (thin dotted line) are indicated. Data were analyzed using the Mann-
Whitney test. ns=not significant.
H. Virgin female recordings aligned to the point of genital interlocking and showing ensuing genital contact
events until the end of each recording (yellow hexagon). Recording of virgin female #18, that did not interlock
during the recording, is excluded from the plot.
I. Analysis setup of pose estimation using SLEAP[55] during genital contact events in string-tethered female
recordings.
J,K. Dynamics of the first 3 seconds of genital contact events, showing the shared angle (**J**) between the male
and the female, and the sinuosity (**K**) of the male (tan) and female (turquoise) plotted separately for the
indicated event types.

561
562
Figure S5: **Female and male genital anatomy and female genital tip elongation as the gating mechanism**
for mating and remating. Related to Figure 1.
A. Reproductions of anatomical drawings female *Aedes aegypti* genital structures (after Jobling, 1987) [30].
Female tip is presented in a lateral view (left) and caudal view with the vaginal lips spread open (right). Vaginal
lips are colored pink.
B. Scanning electron microscopy of female tip.
C. Reproductions of anatomical drawings male *Aedes aegypti* genital structures (after Jobling, 1987) [30]. Male tip
is presented in a lateral view (left) and ventral view (right).
D. Scanning electron microscopy of male tip.
E,F. Individual tracks of female dorsal genital tip length and the male-female shared surface length during
genital interactions of (**e**) virgin and (**f**) previously mated females.

G. Schematic of experimental setup for recording genital behavior and scoring sperm color during sequential
mating.
H-K. Genital behaviors and post-recording scoring of sperm color in spermathecae. Each row corresponds to a
recording of a single female. N=13 recordings with 1 female and two groups of 5 males each.
578

579
580
Figure S6: Gonostyli manipulation and mating attempts of intact and gonostyli-manipulated males.
Related to Figure 2.
A. The step-by-step procedure for removing the gonostyli. **B**. Representative snapshots of intact gonostyli and
the different gonostyli manipulations. Gonostyli are pseudo-colored orange. **C**. Mating attempts observed for 5
minutes in cages of virgin females and intact or gonostyli-manipulated males. N=3 trials per male condition, with
15 males and 15 virgin females each. Cages were gently tapped every minute to induce flight activity, which
induces a corresponding increase in mating attempts. Black line (intact) and dotted lines (males with the
indicated gonostylus manipulation, color coded in legend at bottom) connect the mean mating attempts per
indicated timepoint. Error bars represent standard deviation.

Limatus durhamii   Haemagogus equinus   Culex abominator   Culiseta inornata   Deinocerites epitedeus

Wyeomyia smithii   Psorophora virescens   Aedes auroides   Uranotaenia lowii   Sabethes cyaneus

591
592   **Figure S7: Male genitalia in different mosquito species. Related to Figure 2.**
593   Scientific drawings of male genitalia of the indicated species across *Culicine* genera with putative gonostyli
594   structures colored in green. (reproduced from Howard et al. 1912)[36]

595
596
Figure S8: ***Aedes albopictus* genital anatomy and genital interaction dynamics. Related to Figure 3.**
598 **A,B.** Scanning electron microscopy images of (**A**) female and (**B**) male genitalia of *Aedes albopictus*.
599 **C,D**. Individual tracks of female dorsal genital tip and male-female shared surface dynamics during genital
600 interactions in *Aedes albopictus* from recordings of (**C**) virgin females or (**D**) previously mated females.
601
602

Methods

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Animals used in this study

The following mosquito species and strains were used in this study:

Aedes aegypti: Liverpool wild-type laboratory strain (LVP), PB{3xP3-EGFP; β2Tubulin-dsRed2} (red fluorescent sperm strain, LVP background), and PB{3xP3-dsRed2; β2Tubulin-mNeonGreen} (green-fluorescent sperm strain, LVP background). *Aedes albopictus*: Foshan inbred strain Pavia A (FPA) wild-type laboratory strain. Unless otherwise noted, animals used in this study were between 3 days old and 12 days old, defined as days after adult eclosion. Both males and females were sugar fed and unless otherwise specified females were not blood fed. We define "previously mated" females as those that were at least 24 hours post mating. The LVP strain (*Aedes aegypti* wild type) was initially collected in west Africa and has been maintained at the Liverpool School of Tropical Medicine since 1936[47]. The FPA strain (*Aedes albopictus* wild type) was initially isolated in southeast Asia in 1981 and has been cultivated in the lab since then[48]

METHOD DETAILS

Mosquito rearing

Mosquito strains were reared in a constant temperature and humidity environmental room at 26°C with 70 to 80% relative humidity and daily 14-hour light and 10-hour dark cycle (lights on at 7 a.m.). All behavioral experiments, except pin-tethered male and female mating (**Figure 4**), were performed at 26°C with 70 to 80% relative humidity. Pin-tethered male and female mating experiments were performed at ambient temperature and humidity. Experiments were performed between 9 a.m. and 5 p.m. Embryos were hatched in hatching broth, which was made by crushing one pellet of fish food (TetraMin Tropical Tablets, Pet Mountain, YT16110M) using a mortar and pestle, mixing the powder into 850 ml of deionized water and autoclaving it. The following day, 2500 ml of deionized water were added to the hatched larvae, and they were fed two fish food tablets per day until pupation. Pupae were sexed based on size and shape and male and female pupae placed into separate small cups (VRW, Cat. No. 89009-664) with deionized water. Since even a single male could "contaminate" the virgin female cage, the female pupal cups were covered with a net until eclosion. Upon eclosion, animals were aspirated out in small groups and verified to be female before releasing into designated virgin female cage (30cm^3 BugDorm-1 Insect Rearing Cage, MegaView Science). After eclosion, adult mosquitoes had access to 10% sucrose (w/v in distilled water), which was delivered in a Boston clear round 60 ml glass bottle (Fisher Scientific, FB02911944) with sugar dispensed through a saturated cotton dental wick (Richmond Dental, 201205). For egg collection, female mosquitoes were blood-fed on mice or sheep's blood (HemoStat Laboratories, DSB100) mixture with ATP (20 mM ATP in 25 mM aqueous sodium bicarbonate added to the blood to a 2 mM final concentration) and metal blood pucks as previously described[49]. Eggs were dried at 26°C and 80% humidity for three days, then stored at ambient temperature and humidity for up to 4 months.

Transgenic strain generation

Red and green fluorescent sperm lines were generated using PiggyBac transposition-based transgenesis. The PB{3xP3-EGFP; β2Tubulin-dsRed2} plasmid was kindly provided by Robert Harrell at the Insect Transformation Facility, with the permission of Dr. Peter Atkinson[50] and Dr. Laura Harrington. Transgenic males with this plasmid produce red fluorescent sperm under the control of the β2Tubulin promoter, which confers marker expression in mature sperm. The 3XP3 promoter is used as a marker of transgene insertion and confers marker expression in the eyes. To generate a new strain that produces green fluorescent sperm, the PB{3xP3-EGFP; β2Tubulin-dsRed2} plasmid was modified using the NEBuilder HiFi DNA assembly kit (NEB E5520S) to replace the fluorescent proteins with dsRed2 under the 3xP3 promoter (eye marker), and mNeonGreen. Full plasmid details and sequences are available at AddGene (plasmids #236086 and #236085). Each plasmid was injected at 200 ng/µL together with PiggyBac transposase mRNA (200 ng/µL) to induce transposition-

based genomic integration. Injections were carried out in house using the Zeiss SteREO Discovery
V.8 microscope (Zeiss, 495007-9880-010) and Sutter Instrument injection system (Sutter Instrument
MPS-200 controller, FG-MPC385; XenoWorks Digital Microinjector Sutter Instrument, FG-BRE) into
mosquito embryos collected 1 hour after being laid using a standard injection protocol[51]. Injected G0
male and female animals were reared until adulthood and crossed to LVP animals. G1 animals were
screened for the fluorescent eye marker and two founder animals per injected plasmid were isolated.
Lines were then backcrossed to LVP for 3 generations before being inbred. After validating that all
lines show wild-type levels of mating success and male fertility, one line per injected plasmid was
randomly selected for further study. Transgenic animals were sorted as pupae under a fluorescent
microscope (Nikon SMZ18, NIS-Elements AR software) at 475 nm (green fluorescence) or 549 nm
(red fluorescence) to confirm 3XP3-derived fluorescence from the transgenic insertion in eyes (both
sexes) and testes (males) and separated based on sex as described above.
Rearing and testing transgenic animals (Figure S1E-H)
Reproductive behavior of transgenic males was assessed as follows:
Insemination success: per tested group, 50 wild-type females were separately introduced to 50 males
of the three different genotypes (wild type, red fluorescent sperm, green-fluorescent sperm) for one
hour. Females were cold anesthetized for five minutes and dissected to score sperm presence in their
spermathecae. Experiments were carried out in 5 independent biological replicates.
<u>*Egg laying and hatch rate*</u>: per tested group, 60 wild-type females were introduced to 60 males of the
different genotypes (wild type, red-fluorescent sperm, green-fluorescent sperm) for a period of two
hours. Following the mating period, 10 females of each group were dissected to ensure successful
mating. 2 days post mating, the females were blood fed using Sheep's blood (HemoStat
Laboratories, DSB100) mixture with ATP and metal blood pucks as described before[49]. The few
females that were not fully engorged were discarded. Fully engorged females were placed in
individual vials for egg laying at 5 days after blood feeding and were allowed to lay eggs for 24 hours.
Egg-laying vials were made using plastic Drosophila vials (VWR, 25 mm diameter, 95 mm length,
75813–164) with 2–3 ml of deionized water, and a 55 mm diameter Whatman filter paper (GE
Healthcare, WHA1001055) folded into a cone at the bottom of the vial to serve as a moist egg-laying
substrate as previously described[52,53]. The females were then dissected to ensure insemination, and
the eggs were dried for >24h. The number of eggs per female was counted under the dissection
scope using a hand counter. Egg papers were hatched in individual cups, and larvae were counted 4
days later using images captured by a Logitech camera (Logitech, C922x Pro Stream Webcam) and
the Cell Counter plugin on FIJI/ImageJ (NIH) as previously described[49].
686
Insemination and remating assays
Spermathecae dissection and scoring
Spermathecae dissection: Females were aspirated from their cage and into a paper cup (Solo,
KHB16A-J8000) that was covered with a square of 0.8 mm Polyester Mosquito Netting (American
Home Habitat, F03A-PONO-MOSQ-M008-WT), secured by two rubber bands (Universal Rubber
Bands, Size 12, 0.04" Gauge, Beige, 1 lb Box, 2,500/Pack, UNV00112, Universal) to prevent
mosquitoes from escaping. The cup was placed on wet ice for 5 minutes until the females were cold-
anesthetized. The top of the cup was tapped intermittently to move the females to the bottom of the
cup. 10 ml of 1X PBS (10X PBS buffer, AM9625, Invitrogen; diluted in Millipore water) was pipetted
onto a microscope slide (Fisherbrand Superfrost Plus, 12-550-15, Fisher Scientific). Once cold-
anesthetized, a single female was placed on the slide next to the 1xPBS drop under a dissection
scope (Nikon, Model C-PS), ventral side up. Using Dumont #5 forceps (Roboz Surgical Store, RS-
4978), the seventh sternite was gently pressed down, allowing the abdominal tip of the female to
extend from the abdomen. With this segment extended, the abdominal tip was easily pulled and
removed from the body using forceps and dragged into the 1X PBS drop, and the carcass was
discarded from the slide. The spermathecae were then isolated from the remaining tissue, cleaned,

and placed on the edge of the drop of 1X PBS so that they could be clearly observed for scoring insemination.

Scoring insemination: For insemination scoring, the microscope slide was placed under a compound microscope (SW-350B, SWIFT) with the spermathecae centered in field of view under an air 10x lens with bright field illumination. The female was scored as virgin if the spermathecal sacs were empty and was scored as mated if the spermathecae were full, indicated by sperm presence and darkened color of the sacs. Sperm starts to rotate in the female spermathecae ~24h after insemination[54]. Using the same methods as for scoring insemination, sperm rotation was examined as an additional verification method for ensuring that previously mated females were indeed mated at the beginning of recording in our behavioral experiments.

Scoring fluorescent sperm: Following spermathecae dissection, the slide was placed under a fluorescent microscope (Nikon SMZ18) with the spermathecae centered in the field of view and focus was manually adjusted. The spermathecae were examined under a 10X air lens. Samples were imaged at 475 nm light to score green fluorescent sperm and 549 nm light to score red fluorescent sperm. Fluorescence in both channels was scored as a remated female and no fluorescent sperm was scored as a virgin female. Such females were confirmed to be virgin using the insemination scoring procedure described above.

Remating rates after 7 days of co-housing (Figure S2A,B)
60 virgin females (4 days old, wild type) were aspirated into a cage with 30 green-fluorescent sperm males and 30 red-fluorescent sperm males. The animals were left undisturbed for 7 days, after which the females were aspirated from the cage, spermathecae dissected, and sperm color scored.

Remating rates at different intervals post-mating (Figure S2C-E)
Per tested interval, 40 virgin females (ages differ per experiment type, see below) were placed in a cage after eclosion. The first male group (red or green-fluorescent sperm) was aspirated into the cage and had access to the females for 30 minutes. At the end of the 30 minutes, the males were aspirated out and discarded. After the specified post mating intervals, the second male group (green or red fluorescent sperm, the other color from the first mating group) were introduced to the cage and had access to the females for 120 minutes. At the end of 120 minutes, the second males were aspirated out of the cage and discarded. The first mating was staggered at intervals prior to a second mating challenge that was held on the same day. The females were dissected the following day and their spermathecae were examined under a fluorescent microscope to score sperm presence and color. Females that did not have the first male sperm color or that only had the second male color, reflecting a failure to mate with the first males, were discarded from the analysis. In each experiment, we also had a group of females that were not introduced to the first males, and were only introduced to the second male group, to verify that successful mating could be obtained under conditions of the second mating (see GitHub[46] for full results). To avoid any strain-specific biases in mating, we used the green and red sperm males reciprocally as the first male or second male group and we did not detect any differences that arose from the order of introduction (see GitHub[46] for variation across trials). Two separate experiments were performed to test remating rates at different intervals post mating:

6 day-old females: the females were challenged with remating (2nd male group) at six days post eclosion. First matings were staggered prior to the remating challenge and included the examination of the following intervals post initial mating: 2 hours, 6 hours, 24 hours, 48 hours, 72 hours, and no first mating (control group). n=40 females per post-mating interval tested, N=3 trials.

12 day-old females: the females were challenged with remating (2nd male group) at twelve days post eclosion. First matings were staggered prior to the remating challenge and included the examination of the following intervals post initial mating: 2 hours, 48 hours, 96 hours, 120 hours, 144 hours, 168 hours, and no first mating (control group). n=40 females per post-mating interval tested, N=3 trials. Per experiment and trial, all intervals were tested in parallel.

Behavioral assays and analysis
Observing mating attempts (Figure S1A,B)
Comparison of virgin and previously mated females: 15 virgin females and 15 previously mated females (7 days old, wild type) were allowed to acclimate in 30 cm^3 cages for >6 hours. The cages were then deidentified by another lab member and each cage was examined separately for mating attempts in the following way: 15 males (7 days old, wild type) were aspirated into the cage, followed by a blow of exhaled air to induce flight activity in the cage. The cage was gently tapped every minute to further encourage flight. Mating attempts were scored every 15 seconds by visual observation and indicating the number of mating attempts per timepoint on a spreadsheet. Observation and scoring were carried out for 7 minutes per cage. n=15 pairs per condition tested (virgin or previously mated females), N=3 trials.

Comparison of intact and manipulated gonostyli males: 20 5 day-old males were dissected or mock-treated per group. The following day (>24 hours), any dead males were removed from the cage, together with extra males left – to reach a total male amount of 15 males. The cages were deidentified by another lab member and each cage was examined separately for mating attempts in the following way: 15 virgin females (6 days old, wild type) were aspirated into the cage, followed by a blow of exhaled air to induce flight activity in the cage. The cage was gently tapped every minute to further encourage flight. Mating attempts were scored every 15 seconds by visual observation and indicating the number of mating attempts per timepoint on a spreadsheet. Observation and scoring were carried out for 5 minutes per cage. n=15 pairs per condition tested (intact or gonostyli-manipulated males), N=3 trials.

3-dimensional (3D) recordings of mosquitoes in free flight (Figure S3)
Setup: A 14 cm^3 custom clear acrylic cage (see GitHub[46] for design file) was positioned together with a mirror (10-inch standing cosmetic mirror, DEAYOU, B09XR343HB) angled at 45° to the cage. The camera (sensor: Basler ace acA2040-120um, 35-924; lens: Arducam C-mount Lens for Pi HQ Camera 16mm Focal length, LN045, Arducam) was mounted on a 6-inch stainless steel optical post (Ø1/2 inch Optical Post SS 8-32 Setscrew ¼ inch-20 Tap L = 6 inch 5 Pack, ThorLabs, TR6-P5) for recordings.
Mosquito handling: Animals were 4-8 days old at the time of recording. The first animal (female, if recordings were of a male-female pairs) was added to the cage and recording started. After a minute, the second animal (male, if recordings were of a male-female pair) was aspirated into the cage, and the setup was recorded for a total of seven minutes from that point. Exhaled air was blown into the cage through a mouth aspirator (Oral Aspirator with HEPA Filter Model 612, John W. Hock Company) every 30 seconds to encourage flight. Upon completion of the recording, the female was aspirated out of the cage and her spermathecae dissected to score insemination.
Analysis: Recorded videos were processed to isolate both the real and reflected views of the behavioral chamber. A custom Python script was used to crop each original video, generating two separate videos for each recording: one for the real view and another for the reflected view (mirror). The trajectories of the mosquitoes in each view were reconstructed separately using two distinct top-down multi-animal classifiers trained in SLEAP (Social LEAP Estimates Animal Poses[55]), one for each view, to detect the position of each mosquito over time. Mosquito identities were verified *a posteriori* by visual inspection, ensuring consistent tracking across frames and between the paired cropped videos derived from the same original recording. After obtaining the tracks for each video, a data integration process was conducted to reconstruct the 3D trajectories. For each video, a scaling factor was determined by the ratio of the vertical position in the real view to the corresponding vertical position in the reflected view for each mosquito. For appropriate 3D reconstruction, we corrected optical distortions by establishing a unified linear relationship between the scaling factor and z-position (i.e. the horizontal position in the reflected view) for each video. This approach accounts for

possible aberrations resulting from perspective effects. The resulting 3D trajectories were then
interpolated and smoothed to ensure continuity. From the reconstructed 3D trajectories, two
parameters were estimated: The Euclidean distance between mosquitoes, and the total distance
covered by each mosquito throughout the recording.

String-tethered female and free flying male interactions (Figure S4)

Setup: A 17.5 cm^3 polyester and plastic cage (BugDorm Insect Rearing Cage #4S1515)
was used for recording, with the clear side facing the camera. A piece of nylon fabric previously worn
on arm by a human volunteer was positioned on top of the cage to provide human odor, which
enhances flight activity (as described in[10]). Nylon sleeves were prepared by using scissors to remove
the fabric from the tip of the stocking foot of knee highs (L'eggs brand Everyday, Amazon), so that the
modified stocking was open on both ends[56]. A camera (sensor: Basler ace acA2040-90um, 89-989;
lens: Edmund Optics 16mm/ F1.8, 333304) was mounted on a 6-inch stainless steel optical post
(Ø1/2-inch Optical Post SS 8-32 Setscrew ¼ inch-20 Tap L = 6 inch 5 Pack, ThorLabs, TR6-P5) for
recordings.

Mosquito handling: Animals were 4-8 days old at the time of recording. Female mating status was
deidentified by a lab member. Females were cold-anesthetized and gently placed on a microscope
slide (Fisherbrand Superfrost Plus, 12-550-15, Fisher Scientific) with their dorsal side up. 10 μL of
Bondic UV glue (Bondic UV Liquid Plastic Welder, SK8024 BONDIC) was applied to the tip of a 7 cm
long clear nylon thread (Clear nylon invisible thread, 017531051647, Blulu) and then attached to the
dorsal side of the thorax of the female. UV light (Bondic UV Liquid Plastic Welder, SK8024, BONDIC)
was used to cure the glue for 30 seconds. The free end of the string was then attached to the top part
of the cage using a reusable glue pad (Uhu Tac Removable and Reusable Glue Pads, SAU99683,
UHU) At the start of the recording, the female was prompted to fly by blowing exhaled air into the
cage. Once in flight, a male was aspirated into the cage and recording started, and interactions were
recorded over an 8-minute period. Throughout the recording, the cage was occasionally tapped
gently to stimulate the female's flight. Females were dissected ~5 minutes after the completion of
recording and their spermathecae examined for the presence of sperm.

Manual scoring and analysis of behavioral categories: Recordings were deidentified during the
scoring process. Recorded videos were analyzed using BORIS[57] v. 7.10.7 software. Videos were
uploaded to the manual behavioral scoring interface on BORIS and the following behaviors were
scored: female flight (state), contact (state), genital contact (state), kicking (state), genital interlocking
(state). To accurately define the beginning and end of recordings, "Start" and "End" (events) were
scored per video as indicated by the aspiration of the male into and out of the cage, respectively.
Binary score of behaviors per second was extracted using the "behaviors binary table" option and
plotted as heatmaps on GraphPad Prism (10.4.0). Number of events, average duration, and
proportion of total time spent per behavior were extracted using the "synthetic time budget" analysis
option and plotted as violins on PRISM.

Pose estimation analysis: genital contact and genital interlocking events were initially extracted from
the original videos using a custom Python script to process BORIS annotation files. All resulting
videos started 2 seconds before the annotation timestamp and lasted until the end of the annotated
behavioral event. These were encoded using the open-source FFmpeg library to ensure readability.
Next, we trained and implemented a top-down multi-animal pose estimation model in SLEAP[55] to
track five anatomical landmarks per mosquito: the center of the thorax and four sequential points
along the abdomen. These points were arbitrarily named, from proximal to distal, abdominal point 1,
abdominal point 2, abdominal point 3 and abdominal point 4, with abdominal point 4 representing the
abdominal tip. When visible, the lateral characteristic white markings on *Aedes aegypti* abdomens
served as reference points for consistent labeling of abdominal segments. After applying the trained
model to all processed videos, 181 out of 274 videos met our quality standards for further analysis.
This analysis was conducted to determine whether gross differences in the angle formed between the
male and female mosquitoes during contact could be informative of their mating status. To
characterize the spatial relationship between coupled mosquitoes, we calculated inter-segmental

angles between consecutive body parts using the abdominal tips (Abd4) as pivot points. Angle
dynamics were analyzed across the entire interaction window to identify potential differences
between mating conditions. The following analysis pipeline was developed in Python to process the
tracking data: genital contact initiation was defined using temporal and spatial thresholds based on
the Euclidean distance between the abdominal tips (Abd4) of paired mosquitoes. We focused our
analysis on sustained interactions, defined as continuous genital contact lasting at least 3 seconds
(90 frames at 30 frames per second), yielding 174 files out of 181 for final analysis. Videos were
categorized into four conditions based on female mating status and interaction type: genital contact in
virgin females, genital interlocking, post-genital interlocking genital contact, and genital contact in
previously mated females. The results show a consistent pattern across mating conditions with a
rapid initial increase from 75° to 90° in the first second followed by a gradual stabilization. No clear
statistical differences were observed for the angle formed at the abdominal tips between conditions.
Next, we focused on the contribution of male and females to these observed angle dynamics.
Because the abdomen of mosquitoes is a supple, we decided to analyze the abdominal curvature of
both males and females. To quantify abdominal curvature during mating events, we approximated the
distal abdominal segment (Abd2-Abd4) using Bezier curves. A Bézier curve is a smooth mathematical
curve defined by control points, where the curve passes through the endpoints and is shaped by
intermediate control points that influence its curvature. This method allows for flexible fitting of curved
trajectories to biological data while maintaining mathematical smoothness and avoiding abrupt
directional changes. Sinuosity was calculated as the ratio between the Bezier curve length and the
straight-line distance between Abd2 and Abd4. Females maintained lower, more stable sinuosity
compared to males, with overlapping confidence intervals across all conditions. Thus, Bezier curve
analysis demonstrated minimal variation in abdominal curvature across conditions, indicating that
posture alone may not be a reliable predictor of successful mating events. Altogether, these
measurements suggest that the basic biomechanical configuration during male-female coupling
remains relatively constant regardless of female mating status or interaction outcome.
Recordings of pin-tethered female and free flying males interactions in high-resolution
Setup: A 14 cm^3 custom clear acrylic cage was used for recordings (see GitHub[46] for design file). A
camera (sensor: Basler ace acA2040-120um, 35-924; lens: Computar V5024-MPZ C-Mount 50 mm
Machine Vision Fixed Focal Lens, BH COV5024MPZ) was mounted on a 6-inch stainless steel optical
post (Ø1/2 inch Optical Post SS 8-32 Setscrew ¼ inch-20 Tap L = 6 inch 5 Pack, ThorLabs, TR6-P5),
and the focal point was carefully adjusted to focus on the female's abdominal tip.
Mosquito handling: Animals were 4-8 days old at the time of recording. Females were cold-
anesthetized and gently placed on a microscope slide (Fisherbrand Superfrost Plus, 12-550-15,
Fisher Scientific) with their dorsal side up. 10 ml of Bondic UV glue (Bondic UV Liquid Plastic Welder,
SK8024, BONDIC) was applied to the head of a ball head pin (Brass head ball pin, Waydress,
B09FGFJKQH) and then attached to the thorax of the female. UV light (Bondic UV Liquid Plastic
Welder, SK8024, BONDIC) was used to cure the glue for 30 seconds. The free end of the string was
then attached to the top part of the cage using reusable glue pad (Uhu Tac Removable and Reusable
Glue Pads, SAU99683, UHU). At the start of the recording, the female was prompted to fly by blowing
exhaled air into the cage. Once in flight, 5 males were aspirated into the cage and recording started,
and interactions were recorded over a 5-minute period. Throughout the recording, the cage was
occasionally tapped gently to stimulate the female's flight. Females were dissected ~5 minutes after
the completion of recording and their spermathecae examined for the presence of sperm.
Manual tracking of female dorsal genital tip length and male-female shared surface: Video recordings
of genital interactions of a pin-tethered females and free-flying males were trimmed few seconds
before the initiation of genital contact with a male. One genital contact event was tracked per video,
selected based on whether genital interlocking took place (virgin females), or the first clearly visible
interaction event (previously mated females). Videos were then imported to ImageJ[58] using the
"FFmpeg" import option. The videos were then manually tracked using the "Plugins->Tracking-
>Manual Tracking" option. Per video, tracking started at the first frame in which the male and female

genitalia made contact and tracked for at least 150 frames (on a 37 frames-per-second recording rate), corresponding to 4 seconds. Each track was focusing on a specific point: Track 1= upper end of the female dorsal genital tip. Track 2 = lower end of the female dorsal genital tip. Track 3 = leftmost point of shared contact surface between the male and the female. Track 4 = rightmost point of shared contact surface. Three length measurements were taken of the width of the stem of the metal pin, averaged, and used as a length reference indicating a 0.5 μm real world length. Tracks per sample were then imported to an Excel file and x-y coordinates were used to calculate length of the dorsal genital tip of the female and the shared surface per frame using the following distance formula:

$d = \sqrt{((x_2-x_1)^2 + (y_1-y_2)^2)}$

For the average plots, measurements were averaged independently per frame (using the initiation of genital contact as the first aligned frame) and 95% confidence interval was calculated using t-distribution on R (see GitHub[46]).

Scoring genital interlocking attempts: Video recordings of pin-tethered females – virgin and previously mated – and free-flying males were examined for instances of genital interlocking attempts by males in the absence of the female genital tip elongation response. Genital interlocking attempts were scored when the male genitalia pushed the female genital tip towards the center of the male genitalia, and additional male genital structures (the paraproct and aedeagus) emerged.

Female retraction: To calculate female retraction of dorsal genital tip per behavioral stage, the following measurements were obtained and calculated:

Tracking pre and post leg contact: female dorsal genital tip length and male-female shared surface were tracked as described above ("manual tracking"). The first leg contact event was trimmed per video, and each video was then analyzed frame by frame to locate the point in which a leg of the female (any leg) contacted the male. Using this frame as reference, we tracked the female dorsal genital tip and shared surface 75 frames prior to leg contact and 75 frames following leg contact, to track a total of 150 frames in a 37 frames-per-second recording. Per behavioral stage, measurements of 10 frames were extracted and averaged using the following criteria:

No contact: the first 10 frames that were measured in the pre-contact tracking. *Leg contact*: the 10 frames surrounding the minimum length of the female dorsal genital tip during leg contact. *Male genital contact*: the first 10 frames that were measured in the male genital contact tracking. *Female acceptance*: the 10 frames surrounding the maximum length of the female dorsal genital tip.

Remating Recordings (Figure S5G-K)

Setup: A 14 cm^3 custom clear acrylic cage was used for recordings. A camera (sensor: Basler ace acA2040-120um, 35-924; lens: Computar V5024-MPZ C-Mount 50mm Machine Vision Fixed Focal Lens, BH COV5024MPZ) was mounted on a 6-inch stainless steel optical post (Ø1/2 inch Optical Post SS 8-32 Setscrew ¼ inch-20 Tap L = 6 inch 5 Pack, ThorLabs, TR6-P5) and the focal point was carefully adjusted to focus on the female's abdominal tip.

Mosquito handling: Animals were 4-8 days old at the time of recording. Females were cold-anesthetized and gently placed on a microscope slide (Fisherbrand Superfrost Plus, 12-550-15, Fisher Scientific) with their dorsal side up using forceps (Dumont Tweezers, RS-4978). 10 ml of Bondic UV glue (Bondic UV Liquid Plastic Welder, SK8024, BONDIC) was applied to either string (Clear nylon invisible thread, 017531051647, Blulu) or the head of a ball head pin (Brass head ball pin, Waydress, B09FGFJKQH) and then attached to the thorax of the female. UV light (Bondic UV Liquid Plastic Welder, SK8024, BONDIC) was used to cure the glue for 30 seconds. The free end of the string was then attached to the top part of the cage using reusable glue pad (Uhu Tac Removable and Reusable Glue Pads, SAU99683, UHU). At the start of the recording, the female was prompted to fly by blowing exhaled air into the cage. Once in flight, 5 green-sperm males were aspirated into the cage and recording started. The recording was monitored in real-time to detect female acceptance and genital interlocking. Once genital interlocking and mating occurred and the male detached from the female, the green-sperm males were quickly aspirated out of the cage and 5 red-sperm males were immediately aspirated into the cage and interactions were recorded over an additional 5-minute period. If mating did not take place, the green-sperm males were removed from

the cage after 5 minutes of recording and replaced with the red-sperm males for an additional 5
minutes of recording. Throughout the recording, the female was stimulated to fly by occasional puffs
of exhaled air. Females were dissected ~10 minutes after the completion of recording and their
spermathecae examined under a fluorescent microscope to score presence and color of sperm.
Pin-tethered male and female genital interaction recordings of same-species and cross-
species genital interactions (Figure 4)
Setup: A dissection microscope (Nikon SMZ645), with the eyepieces removed and replaced with the
iDuOptics iPhone mount (LabCam for iPhone, iDu Optics, B0799QTHVR) and an iPhone X attached.
The built-in recording software was used, recording at 30 frames per second.
Mosquito handling: Females and males were collected separately in net-covered paper cups (Solo,
KHB16A-J8000) and cold-anesthetized. The female was gently placed on a microscope slide
(Fisherbrand Superfrost Plus, 12-550-15, Fisher Scientific) with her dorsal side up using forceps
(Dumont Tweezers, RS-4978). 10 ml of Bondic UV glue (Bondic UV Liquid Plastic Welder, SK8024,
BONDIC) was applied to the head of a ball head pin (Brass head ball pin, Waydress, B09FGFJKQH)
and then attached to the thorax of the female. UV light (Bondic UV Liquid Plastic Welder, SK8024,
BONDIC) was used to cure the glue for 30 seconds. The female was then tethered perpendicular and
elevated 5 cm above the dissection scope surface using a reusable glue pad (Uhu Tac Removable
and Reusable Glue Pads, SAU99683, UHU). The orientation of the female was confirmed through the
iPhone X so that her lateral left side was being observed, with her ventral side facing up in the
recording. The male was similarly treated with the UV glue and a ball head pin but was not tethered
so that his position could be controlled and adjusted by the experimenter. The tethered female was
brought into focus and her terminalia was centered in the field of view of the iPhone X. The video
recording was started and the experimenter, by holding the edge of the male's metal pin by hand,
gently positioned the male so that its abdominal tip touches the female tip and the pair is in genital
contact. The process did not involve forceful pushing and instead the abdominal tips of the pair were
gently "brushed" against each other. Throughout the recording, we ensured that the female's
terminalia was consistently in focus by manually adjusting the focus. Recordings lasted until genital
interlocking was complete or until the 8-minute mark was reached. The female was allowed to
recover for 15 minutes while still tethered to the metal pin before her spermathecae were dissected
and scored for the presence of sperm.
Scoring behaviors in pin-tethered male and female genital interaction recordings: Videos were
deidentified and each video was individually scored for the three characterized behaviors: 1) male
genital contact, 2) female genital tip elongation, and 3) genital interlocking. Since the position of the
male was controlled in this experimental setup, male genital contact was scored as a postural change
of the male tip aimed towards a tighter connection with the female tip, and the rapid vibration of the
gonostyli when present. Female genital tip elongation was scored as extension of the female genital
tip from its resting or retracted position. Genital interlocking was scored when the male and female
formed a tight connection of their genitalia, and was further verified by gently pulling the male away
from the female to check that the animals do not separate and remain interlocked.
Gonostylus manipulation assays
Surgical manipulation of the gonostylus (**Figure 2G-M** and **Figure S6A-B**)
General procedure: 3-day old males were cold-anesthetized on ice and were gently placed on a
microscope slide (Fisherbrand Superfrost Plus, 12-550-15, Fisher Scientific) ventral-side up under a
dissection scope (Nikon SMZ645). Focusing in on the genital tip of the male, forceps (Dumont
Tweezers, RS-4978) were used to gently extend the gonostyli from their resting position and make
the following manipulations:
Mock treatment: Males were cold-anesthetized for 15 minutes and placed in a paper cup (Solo,
KHB16A-J8000) for recovery.
Removed gonostyli: Forceps were used to pinch at the base of the gonostylus and remove the entire
structure, with minimal invasiveness of the hairs.

Truncated gonostyli: Forceps were used to pinch at the half-length mark of the gonostyli and remove
the distal end of the structure.
Claws removed: Forceps were used to pinch at the tip of the gonostyli to remove the claws.
Dislocation (immobilize the structures): Forceps were used to gently press alongside the base of the
gonostyli resulting in lack of muscle control to the gonostyli and their immobilization
Following the surgical manipulations, males were placed in a paper cup (Solo, KHB16A-J8000) and
allowed to recover for 24 hours in a $30 cm^3$ BugDorm-1 Insect Rearing Cage (MegaView Science). 24
hours after the manipulation the males were checked for their ability to fly by blowing exhaled air into
the cage, and the paper cup was checked to see if any males did not survive the procedure.
Insemination success of males with manipulated gonostyli
Per manipulation condition, 30 virgin females were placed into the cage of 30 males – all with the
same gonostyli manipulation – which were added 24 hours prior for their post-manipulation recovery.
The animals were left undisturbed for 48 hours to allow ample time for mating. At the end of the 48-
hour period, females were aspirated out of the cage and their spermathecae were scored for sperm
presence. N=4 trials.
Competition success: using the two transgenic strains, green-fluorescent and red fluorescent sperm
males, we conducted mating competition assays between males with manipulated and intact (mock
treated) gonostyli. 20 virgin females (4 days old, wild type) were placed in a $30 cm^3$ cage, to which 15
manipulated and 15 intact males were added. Per replicate and tested manipulation, a reciprocal use
of the two sperm color strains was carried out to prevent strain biases in competitive mating. For each
replicate one tested group had the red sperm males manipulated and the greens intact, and another
group had the green sperm males manipulated (same manipulation) and the red intact. The males
and females were co-housed for 48 hours, after which the females were aspirated out and their
spermathecae were dissected and examined under a fluorescent microscope to score sperm
presence and color. Females which had two sperm colors present (remating) were omitted from the
analysis. Full results are available on GitHub[46].
1032
Scanning Electron Microscopy
Conventional scanning electron microscopy
7-day old male and virgin female mosquitoes were fixed in 2% glutaraldehyde, 4% paraformaldehyde,
in 0.1M sodium cacodylate buffer pH 7.2 (0.4M buffer - Electron Microscopy Sciences, 11655-1G -
diluted with Millipore water), for 3 hours at room temperature. After the initial fixation, the terminus of
the female abdomen was severed with a razor blade between the 4th-5th abdominal segments, and
the anterior portions were discarded. Fixation of the abdominal tip continued for 2 hours at room
temperature and then overnight at 4ºC. After rinsing three times with cacodylate buffer, samples were
dehydrated in a graded series of ethanol (30%, 50%, 70%, 90%) for 10 minutes each and three times
in 100% ethanol with molecular sieves for 15 minutes each and then critical point dried in a Tousimis
Autosamdri 931. After mounting, samples were sputter coated with 10 nm of iridium with a Leica
ACE600 coater. Samples were imaged on a JEOL JSM-IT500HR scanning electron microscope at
5.0 kV. Adobe Illustrator was used to generate falsely colored SEM images.
1046
Cryo-scanning electron microscopy
30 7-day old virgin females were placed in a $30 cm^3$ cage. 30 males were then aspirated to the cage
and the mosquitoes were carefully observed by eye to detect genital interlocking events. Upon
detection of a pair in genital interlocking, the experimenter quickly aspirated the couple using a mouth
aspirator (Oral Aspirator with HEPA Filter Model 612, John W. Hock Company) directly into liquid
nitrogen. The frozen couple was mounted on a cryo-stub (JEOL, Cat. JU2010535) precooled at liquid
nitrogen temperature. Samples were imaged within 30 minutes on a JEOL JSM-IT500HR scanning
electron microscope under low vacuum mode (30 Pa of pressure) at 20 KV.
1055

QUANTIFICATION AND STATISTICAL ANALYSIS
Recording software

BASLER video recording software was used on a Dell Latitude 14 Rugged laptop (Model type: P46G001) to capture the recordings for the 3D, string-tethered, pin-tethered, and remating behavioral assays.

Code and analysis plots

Plots created in this study were generated using Python (3.11), GraphPad Prism (10.4.0), and R (3.6.1). Code development was assisted by Claude (Anthropic, 2024-2025), with all code being manually reviewed and validated by the authors. Details and code are available at GitHub (https://github.com/VosshallLab/HouriZeevi_Vosshall-2025).

ADDITIONAL RESOURCES

Data, Videos 1-10 and analysis code used in this publication are available at
Zenodo (https://zenodo.org/records/15133603)
and GitHub (https://github.com/VosshallLab/HouriZeevi_Vosshall-2025).

Supplemental video legends

Available on Zenodo (https://zenodo.org/records/15133603)[29]

Video 1: Free-flying male and female 3D video recording and tracking {*Aedes aegypti*}

Video 2: String-tethered female and free flying male video recording annotated for behaviors {*Aedes aegypti*}

Video 3: Male genital contact with string-tethered previously mated female annotated for pose estimate analysis {*Aedes aegypti*}

Video 4: Genital interactions of a pin-tethered virgin female and a free-flying male {*Aedes aegypti*}

Video 5: Genital interactions of a pin-tethered previously mated female and a free-flying male {*Aedes aegypti*}

Video 6: Genital interactions of a pin-tethered virgin female and a free-flying male {*Aedes albopictus*}

Video 7: Genital interactions of a pin-tethered previously mated female and a free-flying male {*Aedes albopictus*}

Video 8: Pin-tethered genital interactions {virgin *Aedes aegypti* female X *Aedes aegypti* male}

Video 9: Pin-tethered genital interactions {virgin *Aedes albopictus* female X *Aedes albopictus* male}

Video 10: Pin-tethered genital interactions {previously mated *Aedes aegypti* female X *Aedes albopictus* male}

References

1. Futuyma, D., and Kirkpatrick, M. (2022). Evolution 5th edition. (Oxford University Press).

2. Helinski, M.E.H., Deewatthanawong, P., Sirot, L.K., Wolfner, M.F., and Harrington, L.C. (2012). Duration and dose-dependency of female sexual receptivity responses to seminal fluid proteins in Aedes albopictus and Ae. aegypti mosquitoes. J. Insect Physiol. *58*, 1307–1313. https://doi.org/10.1016/j.jinsphys.2012.07.003.

3. Degner, E.C., and Harrington, L.C. (2016). Polyandry depends on postmating time interval in the dengue vector Aedes aegypti. Am. J. Trop. Med. Hyg. *94*, 780–785. https://doi.org/10.4269/ajtmh.15-0893.

4. Gwadz, R.W., Craig, G.B., and Hickey, W.A. (1971). Female sexual behavior as the mechanism rendering Aedes aegypti refractory to insemination. Biol. Bull. *140*, 201–214. https://doi.org/10.2307/1540069.

5. Craig, G.B. (1967). Mosquitoes: Female monogamy induced by male accessory gland substance. Science *156*, 1499–1501. https://doi.org/10.1126/science.156.3781.1499.

6. Jones, J.C., and Wheeler, R.E. (1965). Studies on spermathecal filling in aedes aegypti (Linnaeus). II. Experimental. Biol. Bull. https://doi.org/10.2307/1539731.

7. Degner, E.C., and Harrington, L.C. (2016). A mosquito sperm's journey from male ejaculate to egg: Mechanisms, molecules, and methods for exploration. Mol. Reprod. Dev. *83*, 897–911. https://doi.org/10.1002/mrd.22653.

8. Clements, A.N. (1999). Genitalia and associated organs. In The Biology of Mosquitoes, Volume 2: Sensory Reception and Behaviour CABI Books., pp. 333–359. https://doi.org/10.1079/9780851993133.0034.

9. Rosenthal, G.G. (2017). Mate Choice: The Evolution of Sexual Decision Making from Microbes to Humans (Princeton University Press).

10. Aldersley, A., and Cator, L.J. (2019). Female resistance and harmonic convergence influence male mating success in Aedes aegypti. Sci. Rep. *9*, 2145. https://doi.org/10.1038/s41598-019-38599-3.

11. Spielman, A. (1964). The mechanics of copulation in Aedes Aegypti. Biol. Bull. *127*, 324–344. https://doi.org/10.2307/1539230.

12. Cator, L.J., Arthur, B.J., Harrington, L.C., and Hoy, R.R. (2009). Harmonic convergence in the love songs of the dengue vector mosquito. Science *323*, 1077–1079. https://doi.org/10.1126/science.1166541.

13. Cator, L.J., and Harrington, L.C. (2011). The harmonic convergence of fathers predicts the mating success of sons in *Aedes aegypti*. Anim. Behav. *82*, 627–633. https://doi.org/10.1016/j.anbehav.2011.07.013.

14. Somers, J., Georgiades, M., Su, M.P., Bagi, J., Andrés, M., Alampounti, A., Mills, G., Ntabaliba, W., Moore, S.J., Spaccapelo, R., et al. (2022). Hitting the right note at the right time: Circadian control of audibility in Anopheles mosquito mating swarms is mediated by flight tones. Sci. Adv. *8*, eabl4844. https://doi.org/10.1126/sciadv.abl4844.

15. Leta, S., Beyene, T.J., Clercq, E.M.D., Amenu, K., Kraemer, M.U.G., and Revie, C.W.
(2018). Global risk mapping for major diseases transmitted by Aedes aegypti and Aedes
albopictus. Int. J. Infect. Dis. *67*, 25–35. https://doi.org/10.1016/j.ijid.2017.11.026.

16. Kraemer, M.U., Sinka, M.E., Duda, K.A., Mylne, A.Q., Shearer, F.M., Barker, C.M., Moore,
C.G., Carvalho, R.G., Coelho, G.E., Van Bortel, W., et al. (2015). The global distribution
of the arbovirus vectors Aedes aegypti and Ae. albopictus. eLife *4*, e08347.
https://doi.org/10.7554/eLife.08347.

17. Powell, J.R., Gloria-Soria, A., and Kotsakiozi, P. (2018). Recent history of Aedes aegypti:
vector genomics and epidemiology records. Bioscience *68*, 854–860.
https://doi.org/10.1093/biosci/biy119.

18. Juliano, S.A. (1998). Species introduction and replacement among mosquitoes:
interspecific resource competition or apparent competition? Ecology *79*, 255–268.
https://doi.org/10.1890/0012-9658(1998)079[0255:SIARAM]2.0.CO;2.

19. Soghigian, J., Sither, C., Justi, S.A., Morinaga, G., Cassel, B.K., Vitek, C.J., Livdahl, T.,
Xia, S., Gloria-Soria, A., Powell, J.R., et al. (2023). Phylogenomics reveals the history of
host use in mosquitoes. Nat. Commun. *14*, 6252. https://doi.org/10.1038/s41467-023-
41764-y.

20. Leahy, M.G., and Craig Jr., G.B. (1967). Barriers to hybridization between Aedes Aegypti
and Aedes Albopictus (diptera: Culicidae),. Evolution *21*, 41–58.
https://doi.org/10.1111/j.1558-5646.1967.tb00129.x.

21. Zhou, J., Liu, S., Liu, H., Xie, Z., Liu, L., Lin, L., Jiang, J., Yang, M., Zhou, G., Gu, J., et al.
(2022). Interspecific mating bias may drive Aedes albopictus displacement of Aedes
aegypti during its range expansion. PNAS Nexus *1*, pgac041.
https://doi.org/10.1093/pnasnexus/pgac041.

22. Tripet, F., Lounibos, L.P., Robbins, D., Moran, J., Nishimura, N., and Blosser, E.M. (2011).
Competitive reduction by satyrization? Evidence for interspecific mating in nature and
asymmetric reproductive competition between invasive mosquito vectors. Am. J. Trop.
Med. Hyg. *85*, 265–270. https://doi.org/10.4269/ajtmh.2011.10-0677.

23. Carrasquilla, M.C., and Lounibos, L.P. (2015). Satyrization without evidence of successful
insemination from interspecific mating between invasive mosquitoes. Biol. Lett. *11*,
20150527. https://doi.org/10.1098/rsbl.2015.0527.

24. Bargielowski, I.E., and Lounibos, L.P. (2016). Satyrization and satyrization-resistance in
competitive displacements of invasive mosquito species. Insect Sci. *23*, 162–174.
https://doi.org/10.1111/1744-7917.12291.

25. Ribeiro, J.M.C., and Spielman, A. (1986). The satyr effect: a model predicting parapatry
and species extinction. Am. Nat. *128*, 513–528. https://doi.org/10.1086/284584.

26. O'Meara, G.F., Evans, L.F., Gettman, A.D., and Cuda, J.P. (1995). Spread of Aedes
albopictus and decline of Ae. aegypti (Diptera: Culicidae) in Florida. J. Med. Entomol. *32*,
554–562. https://doi.org/10.1093/jmedent/32.4.554.

27. Roth, L.M. (1948). A study of mosquito behavior. An experimental laboratory study of the
sexual behavior of Aedes aegypti (Linnaeus). Am. Midl. Nat. *40*, 265–352.
https://doi.org/10.2307/2421604.

28. Smith, R.C., Walter, M.F., Hice, R.H., O'Brochta, D.A., and Atkinson, P.W. (2007). Testis-
specific expression of the β2 tubulin promoter of Aedes aegypti and its application as a
genetic sex-separation marker. Insect Mol. Biol. *16*, 61–71. https://doi.org/10.1111/j.1365-
2583.2006.00701.x.

29. Houri-Zeevi, L., Walker, M.M., Razzauti, J., Sharma, A., Pasolli, H.A., and Vosshall, L.B.
Zenodo 15133603: Mosquito sex under lock and key.
https://doi.org/10.5281/zenodo.15133603 https://doi.org/10.5281/zenodo.15133603.

30. Jobling, B. (1987). Anatomical Drawings of Biting Flies (British Museum (Natural
History)).

31. Clements, A.N. (1999). Mating. In The Biology of Mosquitoes, Volume 2: Sensory
Reception and Behaviour CABI Books., pp. 333–359.
https://doi.org/10.1079/9780851993133.0034.

32. Jones, J.C., and Wheeler, R.E. (1965). An analytical study of coitus in Aedes aegypti
(Linnaeus). J. Morphol. *117*, 401–423. https://doi.org/10.1002/jmor.1051170306.

33. Masly, J.P. (2012). 170 Years of "lock-and-key": genital morphology and reproductive
isolation. Int. J. Evol. Biol. *2012*, 247352. https://doi.org/10.1155/2012/247352.

34. Dufour, L. (1844). Anatomie générale des Diptères. Ann. Sci. Nat. *1*, 244.

35. Eberhard, W.G. (2013). Sexual Selection and Animal Genitalia (Harvard University Press)
https://doi.org/10.4159/harvard.9780674330702.

36. Howard, L.O., Howard, L.O., Dyar, H.G., and Knab, F. (1912). The mosquitoes of North
and Central America and the West Indies (Carnegie Institution of Washington)
https://doi.org/10.5962/bhl.title.54495.

37. Kraemer, M.U.G., Reiner, R.C., Brady, O.J., Messina, J.P., Gilbert, M., Pigott, D.M., Yi, D.,
Johnson, K., Earl, L., Marczak, L.B., et al. (2019). Past and future spread of the arbovirus
vectors Aedes aegypti and Aedes albopictus. Nat. Microbiol. *4*, 854–863.
https://doi.org/10.1038/s41564-019-0376-y.

38. Hartberg, W.K., and Craig Jr., G.B. (1973). Gene-controlled morphological differences in
male genitalia of Aedes aegypti and Aedes mascarensis. Mosq. News *33*, 206–214.

39. Rossignol, P.A., and McIver, S.B. (1977). Fine structure and role in behavior of sensilla
on the terminalia of Aedes aegypti (L.) (Diptera: Culicidae). J. Morphol. *151*, 419–437.
https://doi.org/10.1002/jmor.1051510307.

40. Livdahl, T.P., and Willey, M.S. (1991). Prospects for an invasion: competition between
Aedes albopictus and native Aedes triseriatus. Science *253*, 189–191.
https://doi.org/10.1126/science.1853204.

41. Honório, N.A., Carrasquilla, M.C., Bargielowski, I.E., Nishimura, N., Swan, T., and
Lounibos, L.P. (2018). Male origin determines satyrization potential of Aedes aegypti by
invasive Aedes albopictus. Biol. Invasions *20*, 653–664. https://doi.org/10.1007/s10530-
017-1565-3.

42. Aedes aegypti - Factsheet for experts (2017). https://www.ecdc.europa.eu/en/disease-
vectors/facts/mosquito-factsheets/aedes-aegypti.

43. Aedes albopictus - Factsheet for experts (2017). https://www.ecdc.europa.eu/en/disease-
vectors/facts/mosquito-factsheets/aedes-albopictus.

44. Zheng, X., Zhang, D., Li, Y., Yang, C., Wu, Y., Liang, X., Liang, Y., Pan, X., Hu, L., Sun,
Q., et al. (2019). Incompatible and sterile insect techniques combined eliminate
mosquitoes. Nature *572*, 56–61. https://doi.org/10.1038/s41586-019-1407-9.

45. Dyck, V.A., Hendrichs, J., and Robinson, A.S. eds. (2021). Sterile Insect Technique:
Principles And Practice In Area-Wide Integrated Pest Management 2nd ed. (CRC Press)
https://doi.org/10.1201/9781003035572.

46. Houri-Zeevi, L., Walker, M.M., Razzauti, J., Sharma, A., Pasolli, H.A., and Vosshall, L.B.
(2025). GitHub: Mosquito sex under lock and key.
https://github.com/VosshallLab/HouriZeevi_Vosshall-2025

47. Gloria-Soria, A., Soghigian, J., Kellner, D., and Powell, J.R. (2019). Genetic diversity of
laboratory strains and implications for research: The case of Aedes aegypti. PLoS Negl.
Trop. Dis. *13*, e0007930. https://doi.org/10.1371/journal.pntd.0007930.

48. Battaglia, V., Gabrieli, P., Brandini, S., Capodiferro, M.R., Javier, P.A., Chen, X.-G., Achilli,
A., Semino, O., Gomulski, L.M., Malacrida, A.R., et al. (2016). The Worldwide spread of
the tiger mosquito as revealed by mitogenome haplogroup diversity. Front. Genet. *7*.
https://doi.org/10.3389/fgene.2016.00208.

49. Venkataraman, K., Shai, N., Lakhiani, P., Zylka, S., Zhao, J., Herre, M., Zeng, J., Neal,
L.A., Molina, H., Zhao, L., et al. (2023). Two novel, tightly linked, and rapidly evolving
genes underlie Aedes aegypti mosquito reproductive resilience during drought. eLife *12*,
e80489. https://doi.org/10.7554/eLife.80489.

50. Smith, R.C., Walter, M.F., Hice, R.H., O'Brochta, D.A., and Atkinson, P.W. (2007). Testis-
specific expression of the beta2 tubulin promoter of Aedes aegypti and its application as
a genetic sex-separation marker. Insect Mol. Biol. *16*, 61–71.
https://doi.org/10.1111/j.1365-2583.2006.00701.x.

51. Jasinskiene, N., Juhn, J., and James, A.A. (2007). Microinjection of A. aegypti embryos to
obtain transgenic mosquitoes. J. Vis. Exp. JoVE, 219. https://doi.org/10.3791/219.

52. Matthews, B.J., Younger, M.A., and Vosshall, L.B. (2019). The ion channel ppk301
controls freshwater egg-laying in the mosquito Aedes aegypti. eLife *8*, e43963.
https://doi.org/10.7554/eLife.43963.

53. Duvall, L.B., Ramos-Espiritu, L., Barsoum, K.E., Glickman, J.F., and Vosshall, L.B.
(2019). Small-molecule agonists of Ae. aegypti Neuropeptide Y Receptor block mosquito
biting. Cell *176*, 687-701.e5. https://doi.org/10.1016/j.cell.2018.12.004.

54. Noble, J.M., Degner, E.C., Harrington, L.C., and Kourkoutis, L.F. (2019). Cryo-Electron
Microscopy reveals that sperm modification coincides with female fertility in the mosquito
Aedes aegypti. Sci. Rep. *9*, 18537. https://doi.org/10.1038/s41598-019-54920-6.

55. Pereira, T.D., Tabris, N., Matsliah, A., Turner, D.M., Li, J., Ravindranath, S.,
Papadoyannis, E.S., Normand, E., Deutsch, D.S., Wang, Z.Y., et al. (2022). SLEAP: A
deep learning system for multi-animal pose tracking. Nat. Methods *19*, 486–495.
https://doi.org/10.1038/s41592-022-01426-1.

56. De Obaldia, M.E., Morita, T., Dedmon, L.C., Boehmler, D.J., Jiang, C.S., Zeledon, E.V.,
Cross, J.R., and Vosshall, L.B. (2022). Differential mosquito attraction to humans is
associated with skin-derived carboxylic acid levels. Cell *185*, 4099-4116.e13.
https://doi.org/10.1016/j.cell.2022.09.034.

57. Friard, O., and Gamba, M. (2016). BORIS: a free, versatile open-source event-logging
software for video/audio coding and live observations. Methods Ecol. Evol. *7*, 1325–1330.
https://doi.org/10.1111/2041-210X.12584.

58. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source
platform for biological-image analysis. Nat. Methods *9*, 676–682.
https://doi.org/10.1038/nmeth.2019.

1262