A noncanonical Pol III-dependent, Microprocessor-independent biogenesis pathway generates a germline enriched miRNA family

Rima M. Sakhawala[1,2], Karl-Frédéric Vieux[1,3], Dustin Haskell[4], Guoyun Yu[1], Anna Zinovyeva[4], Katherine McJunkin[1]

[1]Laboratory of Cellular and Developmental Biology, National Institute of Diabetes and Digestive and Kidney Diseases Intramural Research Program, National Institutes of Health, Bethesda, Maryland, USA.

[2]Department of Biology, Johns Hopkins University, Baltimore, Maryland, USA.

[3]Current Address: Department of Biology and Biotechnology, Worcester Polytechnic Institute, Worcester, MA

[4]Division of Biology, Kansas State University, Manhattan, Kansas, USA

Abstract

MicroRNAs (miRNAs) are short RNAs that post-transcriptionally regulate gene expression and play critical roles in development and differentiation. In canonical miRNA biogenesis, primary miRNAs are transcribed from intergenic loci or intronic regions by RNA polymerase II, sequentially cleaved by the Microprocessor complex and Dicer, and resulting mature miRNAs are loaded into Argonaute to repress target mRNAs. A minority of miRNAs are generated via noncanonical biogenesis pathways that bypass the Microprocessor and/or Dicer. Here, we describe a new Pol III-dependent, Microprocessor-independent, and Dicer-dependent biogenesis pathway exemplified by the *mir-1829* family in *C. elegans*. The *mir-1829* family is germline enriched and resides in unusually long introns of three host genes that have no apparent overlapping functions. Using multiple approaches, we determined that the *mir-1829* family is derived from independent Pol III transcripts with consistent transcription start sites. Unlike other Pol III-dependent miRNAs, the *mir-1829* family small RNAs are the dominant species derived from their loci, rather than fragments of a larger functional noncoding RNA. We extend these findings, identifying additional Pol III-transcribed and noncanonical miRNAs in *C. elegans* and human datasets. These young, noncanonical miRNAs may represent an early snapshot in the evolution of de novo miRNA genes.

Keywords: RNA Polymerase III, Drosha, DGCR8, Pasha, *pash-1*, ALG-5, *mir-8196, mir-4937, mir-4521, mir-4638*

Introduction

MicroRNAs (miRNAs) are small, ~22-nucleotide regulatory RNAs that selectively repress gene expression during development and differentiation. In canonical miRNA biogenesis, primary miRNAs are transcribed from intergenic loci or intronic regions by RNA polymerase II (Ha and Kim 2014). These transcripts are then processed by the Microprocessor, a protein complex that consists of the catalytic RNase III enzyme Drosha and a homodimer of the RNA-binding protein DGCR8 (known as DRSH-1 and PASH-1 in *C. elegans*, respectively) (Lee et al. 2003; Denli et al. 2004; Gregory et al. 2004; Han et al. 2004). The Microprocessor recognizes and cleaves the stem of hairpin structures of primary miRNAs (pri-miRNAs) to produce miRNA precursors, which are further processed into small RNA duplexes by the RNase III enzyme Dicer (DCR-1 in *C. elegans*) (Hutvágner et al. 2001; Knight and Bass 2001; Grishok et al. 2001). One strand of the resulting duplex is preferentially loaded into an Argonaute protein, forming the miRNA-Induced Silencing Complex (miRISC), which can target mRNAs via base pairing to their 3′ UTR to repress translation and/or mediate the recruitment of factors that cause mRNA decay (Gebert and MacRae 2019).

Beyond canonical miRNA biogenesis, multiple alternative biogenesis pathways bypass the Microprocessor and/or Dicer. One of the best-studied examples is mirtron biogenesis, in which a short intron takes on a hairpin-like structure after debranching; this hairpin is then processed as a miRNA precursor by Dicer, thus bypassing the requirement for Microprocessor (Berezikov et al. 2007; Okamura et al. 2007; Ruby et al. 2007). Some mirtron-like molecules, termed "Agotrons"

bypass both Microprocessor and Dicer and associate with Argonautes as full-length introns (Hansen et al. 2016). Short, 5′ 7-methylguanine (m^{7}G) capped Pol II transcripts can also bypass the Microprocessor, being directly processed by Dicer before Argonaute loading (Sheng et al. 2018; Babiarz et al. 2008; Xie et al. 2013; Zamudio et al. 2014). Dicer-independent biogenesis is exemplified by vertebrate miR-451, which is cleaved by Microprocessor, but then relies on the catalytic activity of Argonaute to further its maturation (Cheloufi et al., 2010; Cifuentes et al., 2010; S. Yang et al., 2010).

In addition to these Pol II transcripts that undergo noncanonical processing, Pol III-derived transcripts can also be fragmented – generally in a Microprocessor-independent, Dicer-dependent manner – giving rise to Argonaute-loaded small RNAs. Regions of tRNAs that form hairpins can be further processed by Dicer to produce miRNA-like small RNAs (Babiarz et al. 2008; Kuscu et al. 2018; Maute et al. 2013; Martinez et al. 2017). Similarly, specific snoRNAs and the small NF90 associated RNA A (snaR-A) serve as sources for distinct Dicer-dependent small RNA species (Ender et al. 2008; Lemus-Diaz et al. 2020; Stribling et al. 2021). Dicer can also generate small RNAs from hairpins formed by transcribed SINE elements (Babiarz et al. 2008). Overall, diverse Pol III transcripts give rise to less abundant small RNA products that can function as miRNAs.

In this study, we present a new noncanonical miRNA biogenesis pathway exemplified by the *mir-1829* family in *C. elegans*. Although they reside within long introns of Pol II-derived host genes, the *mir-1829* family members are derived from independent transcriptional units transcribed by Pol III. The *mir-1829* family is Microprocessor-independent but Dicer-dependent. The *mir-1829* family is enriched in the germline and, in particular, enriched in ALG-5, a germline specific Argonaute protein. Unlike previously-identified Pol III-dependent miRNAs, the *mir-1829* family small RNAs appear to be the primary product of these loci, not derivatives of a more

abundant functional species. We further expanded our search for additional noncanonical and Pol III-derived miRNAs by cross-referencing datasets in which Microprocessor, Dicer, and Pol III are depleted, identifying multiple novel candidates in *C. elegans* and human samples. Both the *mir-1829* family and additional candidate miRNAs are evolutionarily young; thus, these miRNAs may represent a snapshot of miRNA loci at an early stage of their emergence.

Results

The *mir-1829* family is Microprocessor-independent

While investigating miRNA decay in *C. elegans* adults using a temperature-sensitive (ts) allele of the DGCR8 ortholog *pash-1* (Vieux et al. 2021; Lehrbach et al. 2012), we noticed a small portion of miRNAs that appeared to be insensitive to *pash-1* inactivation (Figure 1A, x-axis). These *pash-1*-insensitive miRNAs included both known Microprocessor-independent mirtrons (Chung et al. 2011; Jan et al. 2011; Ruby et al. 2007) and uncharacterized miRNAs such as the *mir-1829* family. To determine whether the *mir-1829* family is also Microprocessor-independent, we compared the *pash-1(ts)* data to another dataset in which RNAi and the auxin induced degron system were used together (RNAiD) to stringently deplete the Microprocessor in embryos (Dexheimer et al. 2020) (Figure 1A, y-axis). In comparing the two datasets, we observe that most miRNAs are sensitive to Microprocessor depletion in both datasets as shown by their decreased abundance (Figure 1A). The imperfect correlation of the amplitudes of miRNA depletion between the two datasets likely stems from the different miRNA complements present in the developmental stages assayed (adults versus embryos). Like in *pash-1(ts)*, Microprocessor-independent mirtrons and uncharacterized miRNAs were also insensitive to Microprocessor depletion by RNAiD. Among the novel Microprocessor-insensitive miRNAs, the vast majority are members of the *mir-1829* family. The *mir-1829* family is comprised of four miRNA loci, *mir-1829a, mir-1829b, mir-*

1829c, and *mir-4812*, with highly similar sequences, including a shared seed sequence in the 3p-derived strand. Both 5p and 3p miRNAs of these four loci are Microprocessor-independent. (A fifth locus, *F39B1.3*, bears high similarity to those of the *mir-1829* family, but does not appear to be expressed.) Thus, we hypothesized that the *mir-1829* family are representatives of a new noncanonical Microprocessor-independent class of miRNAs.

Canonical miRNAs are characterized by having a ~95-nt (~35bp) stem loop hairpin which serves as an optimal substrate for Microprocessor cleavage (Auyeung et al. 2013; Fang and Bartel 2015; Nguyen et al. 2023; Kang et al. 2021; Kim et al. 2021; Partin et al. 2020; Jin et al. 2020) (Figure 1B). Mirtrons, which are derived from splicing, bypass the Microprocessor and serve as substrates for Dicer cleavage. Mirtrons generally have shorter hairpins than canonical miRNAs, lacking a basal stem that is characteristic of Microprocessor substrates (Westholm and Lai 2011) (Figure 1B). To determine whether newly-identified Microprocessor-independent miRNA candidates also lack a basal stem, we examined the distribution of miRNAs derived from primary transcripts containing only a "short hairpin" (<32bp) among the *pash-1(ts)* and Microprocessor RNAiD data. While "short hairpins" were present among Microprocessor-sensitive miRNAs, the Microprocessor-insensitive miRNAs were highly enriched for "short hairpins", highlighting both known mirtrons and novel Microprocessor-independent candidates like the *mir-1829* family (Figure 1A-B, D).

The structure of the *mir-1829* family – along with their Microprocessor-insensitivity – suggests that they are poor Microprocessor substrates (Figure 1B, D). Consistent with this, the *mir-1829* family were among the least-efficiently cleaved pri-miRNAs in *in vitro* processing assays performed using purified *C. elegans* Drosha (Figure 1C, x-axis) or the whole Microprocessor complex (Figure 1C, y-axis) (Nguyen et al. 2023).

We further confirmed these published sequencing datasets by examining miRNA levels via northern blotting and RT-qPCR. We used strains in which PASH-1 and/or DRSH-1 are tagged with the auxin inducible degron (AID) to deplete components of the Microprocessor for 24 h (Dexheimer et al. 2020). Using stringent northern blot conditions, we observed depletion of a canonical miRNA (*mir-1*) in samples depleted of PASH-1 and/or DRSH-1, demonstrating efficient inactivation of Microprocessor (Figure 1E). In contrast, mature *mir-1829b-5p* and *mir-1829c-5p* were not diminished in samples depleted of the Microprocessor; in fact, they were modestly up-regulated (Figure 1E). (Among the *mir-1829* family, only the 5p guide strands of *mir-1829b* and *mir-1829c* were detectable by northern. We similarly focused on *mir-1829b-5p* and *mir-1829c-5p* for all RT-qPCR experiments because they were also the most robustly detectable *mir-1829* family members in these assays.) Similar to northern blots, RT-qPCR showed slightly higher levels of *mir-1829b-5p* and *mir-1829c-5p* in the same samples (Figure 1F). The increased accumulation of noncanonical miRNAs upon Microprocessor inactivation has been noted previously (Kim et al. 2016; Sheng et al. 2018; Babiarz et al. 2008) and likely reflects reduced competition from more abundant canonical miRNAs for Argonaute loading and/or Dicer-mediated processing. Overall, these data confirm deep sequencing data indicating that Microprocessor inactivation does not impair *mir-1829* family biogenesis. The caveat that the lack of sensitivity to Microprocessor depletion could be attributed to an unusually long half-life of the *mir-1829* family is addressed below in experiments demonstrating Dicer-dependence (see below).

The *mir-1829* family is germline-enriched

C. elegans has 19 Argonaute proteins, including three that are primarily dedicated to loading miRNAs: Argonaute-like gene 1 (ALG-1), ALG-2, and ALG-5 (Seroussi et al. 2023). ALG-1 and ALG-2 are both expressed ubiquitously in the soma, and ALG-2 is also expressed in

the germline and early embryos (Aalto et al. 2018; Vasquez-Rifo et al. 2012). ALG-5 expression is restricted to the germline (Seroussi et al. 2023; Brown et al. 2017). To examine potential downstream functions of the Microprocessor-independent *mir-1829* family, we examined its Argonaute loading profile and tissue of expression. According to small RNA sequencing of immunoprecipitations of *C. elegans* Argonautes (IP-sRNA-seq), all members of the *mir-1829* family are highly enriched in ALG-5 IPs and modestly enriched in ALG-2 IPs (Brown et al. 2017; Seroussi et al. 2023). Building upon these data, we sought to determine the relative distribution of the *mir-1829* family in each miRNA Argonaute via IP-RT-qPCR from strains in which ALG-1, -2, or -5 is tagged at its endogenous locus with a 3xFLAG epitope (as well as GFP or mKate2) (Aalto et al. 2018; Brown et al. 2017). We observed that *mir-1829b-5p* is evenly distributed between ALG-2 and ALG-5, whereas *mir-1829c-5p* is evenly distributed between all three miRNA Argonautes (Figure S1A).

We reasoned that our IP-RT-qPCR data could be consistent with IP-sRNA-seq data (in which higher amplitude of enrichment was observed in ALG-5 IPs than ALG-1/2 IPs) if low abundance of ALG-5 imparts higher dynamic range to ALG-5 IP enrichment scores than those for ALG-1 and ALG-2. We therefore determined the relative expression of ALG-1, ALG-2, and ALG-5 in whole animal adult samples; because each Argonaute is endogenously tagged with 3xFLAG, the anti-FLAG signal can be compared across strains to assess relative Argonaute abundance (Figure S1B, input lanes). Accordingly, ALG-1 makes up ~60% of the miRNA Argonaute pool, whereas ALG-2 makes up ~37% and ALG-5 a mere ~3% in whole adult hermaphrodite samples (Figure S1C). Therefore, the dynamic range of enrichment from whole animal samples is greater for ALG-5 IPs than those of ALG-1 and ALG-2, explaining the even distribution of *mir-1829b/c* across multiple Argonautes. Although *mir-1829* family members can be detected in some somatic

tissues (Alberti et al. 2018; Brosnan et al. 2021), the exclusive loading of *mir-1829b-5p* in ALG-5 and ALG-2 suggests that its expression is very strongly germline-enriched, limiting association with ALG-1 (which is not expressed in the germline). The even distribution of *mir-1829c-5p* across ALG-1/2/5 suggests that its expression is strongly enriched in the germline (given that ~33% of the miRNA is loaded into the 3% of the Argonaute pool comprised of ALG-5), along with some expression in the soma. Additionally, germline-specific profiling of miRNAs shows that all *mir-1829* family members are highly expressed in the distal germline during the mitosis to meiosis transition (Diag et al. 2018).

Because of its germline-enriched expression, we sought to determine if the *mir-1829* family plays a role in germline function, specifically in fertility. Using CRISPR-Cas9, we generated strains that contain various combinations of *mir-1829* knockouts. Reduced brood size and embryonic lethality were initially observed in a *mir-1829a* deletion [*mir-1829a(del)*] and a full knockout of the *mir-1829* family [*mir-1829 family(del)*] at elevated temperatures. However, these phenotypes were not rescued by a second round of CRISPR that restored the *mir-1829a* sequence [*mir-1829a(rescue)*] and thus are due to off-target effects of CRISPR (Figure S1D). The *mir-1829* family members reside in introns of protein coding genes (see also below), and the host gene of *mir-1829a*, *gas-1*, is required for normal fertility (Morgan and Sedensky 1994). We therefore redesigned the *mir-1829a* deletion to minimize effects on *gas-1* expression by deleting the whole *mir-1829a* host intron and fusing the flanking exons together [*mir-1829a(intron del)*]. Neither this strain nor a quadruple mutant strain containing this *mir-1829a(intron del)* design along with knockout of all other family members [*mir-1829 family(null)*] showed abnormal brood or embryonic viability phenotypes. Therefore, the *mir-1829* family is dispensable for normal fitness in laboratory conditions.

ALG-5 has a demonstrated role in promoting the mitosis to meiosis transition in the *C. elegans* germline, evidenced by strong enhancement of a the *glp-1(gf)* tumorous phenotype by *alg-5(null)* (Brenner et al. 2022) (Figure S1E). Because of the expression of the *mir-1829* family in the distal germline and its loading in ALG-5, we examined its role in this context. We observed that the quadruple knockout of *mir-1829* does not modify the *glp-1(gf)* phenotype (Figure S1E); thus, it is not the primary miRNA guiding ALG-5's function in this context. Overall, the biological function of the *mir-1829* family remains to be determined.

The mir-1829 **family is differentially transcribed from its host genes**

The four *mir-1829* family loci reside in the introns of three protein coding genes of apparently unrelated function (*gas-1*, *piki-1*, and F20D1.3). As mentioned above, data from the SPACEGERM (Spatial *Caenorhabditis elegans* germline expression of mRNA & miRNA) atlas indicate that the *mir-1829* family peak in expression in the distal germline (Diag et al. 2018) (Figure 2A). Curiously, the host genes peak in expression in the proximal gonad, suggesting differential expression of the miRNAs and their host genes (Figure 2A). Published ChIP-seq data of Pol III subunit RPC-1 further supports the idea that the *mir-1829* family is transcribed independently from its host genes and by Poll III (Araya et al. 2014); the *mir-1829* loci (*mir-1829* precursors shown in pink) are marked by prominent Pol III ChIP peaks, and these peaks are just as prominent as those at tRNA loci (purple boxes) (Figure 2B-C).

To investigate whether the differential expression between host gene and miRNA is due to differential transcription, we performed 5′ Rapid Amplification of cDNA Ends (RACE) on RNA isolated from embryo, L4, and adult tissues using primers that bind in the apical loop of the miRNA hairpins. For each of the *mir-1829* family loci, multiple 5′ RACE reads mapped within the host intron of the miRNA, suggesting alternative proximal transcription start sites of transcripts

independent of the protein-coding host genes (Figure 2B-C, Table S3). For three of the four members of the *mir-1829* family (*mir-1829a-c*), multiple 5′ RACE reads mapped consistently 101-103bp upstream of the precursor, which further suggests the *mir-1829* family members share conserved regulatory elements that drive their independent transcription.

The mir-1829 family is transcribed by Pol III

Because results of 5′ RACE together with published Pol III ChIP suggest that the *mir-1829* family may arise from independent Pol III transcriptional units, we sought additional evidence of their Pol III dependence. Type II Pol III transcripts (e.g. tRNAs) are defined by having Box A and Box B motif promoter elements located downstream of the transcription start sites (TSS) (Dieci et al. 2007) (Figure 3A). For each *mir-1829* family member, we searched upstream of the precursor for the Box A and Box B consensus sequence motifs previously defined for *C. elegans* (Ikegami and Lieb 2013; Stutzman et al. 2020). We determined that all four members have a putative Box A and Box B motif ~79-87bp and ~35-42bp upstream of the precursor, respectively. The positions of the motifs are downstream of the prominent -102bp TSS identified by 5′ RACE for *mir-1829a*, *mir-1829b*, and *mir-1829c* (Figure 3A). Additionally, all the *mir-1829* family loci have 5Ts immediately downstream of the precursor (3-4bp downstream of the 3p arm), and another stretch of 4-5Ts further downstream. These features further support our hypothesis that these are Pol III transcripts since typical Type II Pol III genes have terminator sequences of 5Ts immediately downstream of the gene body (Ikegami and Lieb 2013).

To determine whether the Box A/B motifs are required for *mir-1829* biogenesis, we performed promoter bashing experiments for *mir-1829b*. As a control, we used the *mir-1829 family(null)* mutant in which all the family members are deleted; in this background, a 326bp deletion that removes *mir-1829b* abolishes expression of the miRNA without disrupting the host

gene transcript, F20D1.3 (Figure 3B). The *mir-1829b* locus contains a putative Box A motif, a second degenerate Box A motif, and a putative Box B motif. We first deleted all the promoter motifs, an 82bp deletion, keeping the TSS intact (Box A/B del). This mutant resulted in abolished expression of the miRNA without disrupting the host gene transcript. To ensure that any changes in expression observed were not due to disruption of the secondary structure, we designed a new mutant (Box A/B scramble) in which we scrambled the sequence within a 62bp window, removing the Box A and Box B sequence motifs while maintaining secondary structure of the transcript as predicted by RNAfold (Lorenz et al. 2011; Gruber et al. 2008). Similar to the Box A/B del mutant, we observed abrogated miRNA expression, without disruption of the host gene transcript (Figure 3B).

We next mutated each motif individually. To maintain structure for the Box A scramble mutant, we introduced additional mutations into the degenerate Box A motif, since they form a hairpin according to the RNAfold prediction. In this Box A scramble mutant, we observed abrogated *mir-1829b-5p* expression (Figure 3B). We separately mutated only the degenerate Box A motif in a manner that is not predicted to maintain secondary structure (2nd Box A scramble); this resulted in no change in expression of *mir-1829b-5p* (Figure 3B). Lastly, in the Box B scramble mutant, we maintained predicted structure while disrupting the sequence of the Box B motif and observed loss of *mir-1829b-5p* expression. Together, these results indicate that the Box A and Box B motifs, individually, are each required for *mir-1829b* expression, whereas the degenerate Box A motif is not.

Building on the promoter bashing experiments, we attempted to express the predicted *mir-1829b* locus from an ectopic location. We re-integrated a 510bp minimal transcriptional unit – spanning 125bp upstream of the TSS to 222bp downstream of the precursor – into the *mir-1829*

family(null) background at a known safe harbor locus (the position of MosSCI locus *ttTi5605* in Chromosome II). Not only did we observe expression of *mir-1829b-5p* in this rescue strain, but expression was increased nearly eight-fold compared to wild type (Figure 3C). The observed expression of the minimal transcriptional unit demonstrates that the 510bp fragment contains all the required regulatory elements necessary for *mir-1829b* expression.

To further test the model that the *mir-1829* family is a Pol III transcript, we depleted Pol III using the AID system. To do this, we tagged the Pol III catalytic subunit RPC-1 with a 3xFLAG::AID tag using CRISPR in a strain that expresses TIR1 ubiquitously, resulting in efficient RPC-1 depletion upon auxin treatment (Figure S2). We assessed expression of *mir-1829b-5p*, *mir-1829c-5p*, and their respective host gene transcripts after 24 h of auxin treatment using RT-qPCR. Depletion of Pol III by auxin treatment of the AID-tagged RPC-1 strain did not impact levels of the Pol II-transcribed host gene transcripts but strongly abrogated expression of *mir-1829* family members (Figure 3D). In contrast, *mir-1829b/c* levels were unaffected in RPC-1 degron strains lacking auxin treatment, signifying that tagging RPC-1 alone does not disrupt its function (Figure 3D). Furthermore, the significant depletion of *mir-1829b/c* was not due to off target effects of auxin treatment or TIR1 expression, as observed by the lack of miRNA depletion in strains that do not contain tagged RPC-1 (Figure 3D – "TIR1 +/- auxin"). Overall, depletion of RPC-1 along with the mutation of BoxA/B motifs support the model that the *mir-1829* family is transcribed by Pol III.

To determine if Pol II also plays a role in the transcription of the *mir-1829* family, we deleted the promoter regions of the host genes with the rationale that this would remove Pol II occupancy from these loci. We deleted 1.1kb and 1.7kb upstream of the start codon for F20D1.3 and *piki-1*, respectively (Figure 3E). We observed strong reduction of host gene expression levels

in their respective mutants via RT-qPCR, suggesting that we effectively removed Pol II occupancy (Figure 3F). In the *piki-1* promoter deletion mutant, there was no change in *mir-1829c-5p* levels. In the F20D1.3 promoter deletion mutant, expression of *mir-1829b-5p* was slightly increased (Figure 3F). We hypothesize that there may be slight competition between Pol II and Pol III for the F20D1.3/*mir-1829b* locus, since deletion of the F20D1.3 promoter region (Figure 3F) or ectopically expressing *mir-1829b* from another locus (Figure 3C) results in increased expression of *mir-1829b-5p*. Together, these results validate the model that the *mir-1829* family is primarily transcribed by Pol III, with no detectable contribution from Pol II-driven host gene transcription.

mir-1829 family biogenesis requires Dicer, but not DRH-1, RDE-3 or NSUN-2

Having demonstrated that the *mir-1829* family is Pol III-transcribed and Microprocessor-independent, we sought to further elucidate its biogenesis. To do so, we employed the AID system to determine the dependence of *mir-1829* biogenesis on Dicer (DCR-1). DCR-1 was tagged with 3xFLAG-AID at its endogenous locus and crossed into a background expressing TIR1 ubiquitously or only in the germline, resulting in dramatic or moderate depletion of overall DCR-1, as expected (Figure S2). We first performed northern blotting to examine the abundance of the mature miRNA species and the accumulation of the precursor. As a control, we probed for the canonical miRNA *mir-1*, which is primarily expressed in the pharynx and muscle (Brosnan et al. 2021; Simon et al. 2008; Xu et al. 2024; Gutiérrez-Pérez et al. 2021). We observed reduction of mature *mir-1* and accumulation of its precursor in the samples where Dicer is depleted ubiquitously but not in the case of germline-only Dicer depletion (Figure 4A). To achieve maximal signal for *mir-1829b/c,* low stringency blotting conditions were used that do not distinguish between *mir-1829b-5p* and *mir-1829c-5p* when using a *mir-1829b-5p* probe, as demonstrated by detection of pure synthetic miRNA species corresponding to both sequences (Figure 4A, right side). We

observe decreased *mir-1829b/c* expression in all the Dicer-depleted samples, including those in which Dicer is depleted only in the germline, consistent with strong germline-enrichment of the *mir-1829* family (Figure 4A). Despite low stringency probing conditions, the *mir-1829* family is still near the limit of detection (Figure 4A). This low signal may contribute to our failure to detect accumulation of the precursor species of *mir-1829b/c* as expected upon Dicer depletion.

Due to the limited detection of the *mir-1829* family by northern blotting, we further assessed miRNA levels using RT-qPCR. We observed significant reduction of both *mir-1829b-5p* and *mir-1829c-5p* in samples in which Dicer is ubiquitously depleted (Figure 4B). Furthermore, the level of reduction in *mir-1829b-5p* in germline-only Dicer depletion samples nearly mirrors the level seen in the ubiquitously depleted samples (Figure 4B), further supporting germline-restricted expression of *mir-1829b-5p*. We observe milder depletion of *mir-1829c-5p* in the germline Dicer depletion samples compared to depletion in the whole worm (Figure 4B). This result further supports that *mir-1829c* is expressed in the both the germline and the soma as noted above (Figure S1A).

To further assess Dicer dependence, we performed sRNA-seq of samples in which Dicer is depleted ubiquitously or in the germline (Tables S6-S7). Overall, almost all miRNAs appear to be Dicer-dependent – many of which are statistically significant after spike-in normalization (PAdj > 0.05) (Figure 4C, Table S6). The *mir-1829* family behaves similarly to canonical miRNAs in that it is sensitive to Dicer depletion (Figure 4C, red circles). Consistent with all results discussed above, we also observed depletion of the *mir-1829* family members when Dicer was depleted in the germline alone, along with depletion of a small set of germline-enriched canonical miRNAs (Figure S3, Table S7).

To process long dsRNA substrates, *C. elegans* Dicer works in concert with two cofactors, double stranded RNA (dsRNA) binding protein RDE-4 and RIG-I-like receptor DRH-1 (Tabara et al. 2002; Duchaine et al. 2006; Lu et al. 2009; Consalvo et al. 2024). Since the *mir-1829* family is noncanonically processed but requires Dicer, we tested the role of RDE-4 and DRH-1 in *mir-1829* biogenesis using knockout strains. From RT-qPCR experiments, we observed no change in *mir-1829b/c* in the *rde-4* and *drh-1* mutant strains compared to wild type (Figure S4A). Thus, Dicer functions in *mir-1829* maturation independent of RDE-4 and DRH-1.

We also investigated the role of *nsun-2* in *mir-1829* biogenesis. Vault RNAs can be further processed by Dicer to produced small RNA fragments (svRNAs) that function in RNAi when loaded into Argonaute (Persson et al. 2009). In cell lines, NSun2, an RNA methyltransferase that modifies cytosine, methylates vault RNA to prime them for dicing (Hussain et al. 2013). To determine if the *C. elegans* homolog *nsun-2* plays a role in *mir-1829* biogenesis, we introduced an 3xFLAG-AID tag to the C-terminus of NSUN-2 by CRISPR. After 24 h auxin treatment in a ubiquitous TIR1 background, although NSUN-2 levels decreased approximately 90% (Figure S2), *mir-1829b/c* levels remained unchanged (Figure S4B). Thus, *nsun-2* does not appear to play a role in *mir-1829* family biogenesis.

The 3′ end of the *mir-1829* family precursor is likely generated in the nucleus

Although they differ in biogenesis of their precursor, mirtrons are similar to the *mir-1829* class of miRNAs in their Microprocessor independence and Dicer dependence. Mirtron biogenesis is counteracted by untemplated 3′ end uridylation of the precursor species, which destabilizes these intermediates (Bortolamiol-Becet et al. 2015; Reimão-Pinto et al. 2015). Activity of terminal nucleotidyl transferases on miRNA precursor substrates can be detected through sequencing of mature miRNA populations since the untemplated nucleotide additions (tails) can persist after

Dicer-mediated maturation. In particular, tailing that is strongly skewed toward 3p-derived miRNA strands is likely a signature of an enzyme that acts on miRNA precursor substrates; such activity would not result in tailing of 5p-derived miRNA strands whose 3′ ends are not available for modification until after Dicer-mediated precursor cleavage. We previously observed that miRNA A-tailing in *C. elegans* is strongly skewed toward 3p-derived miRNAs, suggesting that an adenyltransferase acts on miRNA precursors (Vieux et al. 2021). To determine whether *mir-1829* family biogenesis is regulated by tailing of the precursor, similar to mirtrons, we examined the sequences of the mature strands for evidence of tailing (which may have been deposited on either precursor or mature miRNA species). We observed various tailed species of *mir-1829* family members with a clear enrichment of adenylation on the 3p-derived strands, suggestive of adenylation of the precursors (Figure 5A). Since GLDR-2 is responsible for A-tailing of miRNA precursors in *C. elegans*, we reanalyzed published data to examine if GLDR-2 is also responsible for tailing of the *mir-1829* family (Vieux et al. 2021). We observed loss of A-tailing on 3p arms of *mir-1829* family members in *gldr-2(null)* mutants (Figure 5B) (Vieux et al. 2021). To determine if A-tailing is involved in *mir-1829* biogenesis, we evaluated mature miRNA levels in the presence and absence of GLDR-2. We observed no change in the levels of expression, suggesting that, although *mir-1829* precursors are adenylated by GLDR-2, A-tailing is dispensable for *mir-1829* biogenesis (Figure 5B) (Vieux et al. 2021).

The A-tailing of the 3p arms of the *mir-1829* family suggests that a precursor-like intermediate (in which the 3′ end of the 3p arm is available for tailing) precedes Dicer-mediated maturation of the mature duplex, despite our inability to detect this intermediate on northern blots. We reasoned that localization of GLDR-2 may further inform the order of events of *mir-1829* biogenesis. A GFP tag fused to GLDR-2 at its endogenous locus reveals that GLDR-2 specifically

localizes to the nucleus (Figure 5C). This suggests that the cleavage event(s) that gives rise to the miRNA precursor – or at least maturation of the precursor 3′ end – occurs in the nucleus prior to Dicer-mediated cleavage in the cytoplasm (Figure 5D). Thus, we have delineated a new miRNA biogenesis pathway, as exemplified by *mir-1829* family, that involves the transcription of an atypical primary miRNA transcript by Pol III, initial Microprocessor-independent maturation steps in the nucleus, followed by Dicer-mediated cleavage in the cytoplasm (Figure 5D).

Noncanonical miRNA biogenesis extends beyond the *mir-1829* family

Having defined atypical features of the *mir-1829* family's biogenesis, we sought to extend these findings by identifying additional noncanonical miRNAs. First, we re-examined datasets in which Microprocessor is inactivated (Vieux et al. 2021; Dexheimer et al. 2020). For these analyses, we used uncurated miRBase annotations (Kozomara et al. 2019), rather than MirGeneDB which stringently filters for features of canonical miRNAs and evolutionary conservation (Fromm et al. 2020). Among Microprocessor-insensitive miRNAs, we observed many annotated mirtrons as expected (Figure 6A, Tables S10-11) (Chung et al. 2011; Jan et al. 2011; Ruby et al. 2007). In both Microprocessor inactivation datasets, we also observed that *mir-8196a-3p, mir-8196b-3p*, and *mir-8199* were not depleted (Figure 6A, Tables S10-11) (Vieux et al. 2021; Dexheimer et al. 2020). In contrast, all of these were decreased upon Dicer depletion over the same time scale (24 h), demonstrating that a long half-life for these small RNAs is not responsible for their perdurance in the setting of Microprocessor inactivation (Table S10). Based on its sequence and loading into ERGO-1 (Seroussi et al. 2023), *mir-8199* appears to be a misannotated 26G class endo-siRNA. *mir-8196a/b-3p* are more intriguing because they are bound in part by ALG-1 and ALG-2 (Seroussi et al. 2023), and therefore may function as noncanonical miRNAs. *mir-8196b* lies 65bp downstream of a recently annotated 21U piRNA on the X chromosome (Seroussi et al. 2023)

(Figure S5A). Similarly, *mir-8196a* similarly lies 71bp downstream of an unannotated 21U piRNA, also on the X chromosome; the upstream RNA meets the requirements of a 21U based on its sequence, loading in PRG-1, and depletion in *prg-1^{-/-}* (Figure 6B). This consistent proximity to a 21U locus raises the possibility that this relative position may contribute to *mir-8196a/b* transcription or biogenesis. Consistent with this, *mir-8196a/b* are germline-enriched, as evidenced by their significant reduction in samples in which Dicer is depleted only in the germline (Table S12).

Multiple Dicer-independent small RNAs were also detected (Figure 6C, Tables S8,10-11). These include misannotated 21U piRNAs that persist in mirBase that were recently noted (Seroussi et al. 2023). Thus, these small RNAs act as positive controls for Dicer-independence, similarly to mirtrons for Microprocessor independence. We also find that *mir-5549-3p* is an additional misannotated 21U RNA encoded on chromosome III; it is 21-nt in length with U in the 5′ position, bound by PRG-1, and depleted in *prg-1^{-/-}* (Seroussi et al. 2023) (Figure S5B). Finally, we identify *mir-4937* as a Dicer-independent small RNA (see more below).

We also sought to further define Pol III-dependent miRNAs. To this end, we performed sRNA-seq on samples in which RPC-1 was depleted using the AID system and overlapped these data with published Pol III ChIP-seq data to identify bona fide Pol III-transcribed miRNAs (Figure 6D, Tables S9-11). The strongest candidate is again *mir-4937*, which is RPC-1-dependent and has prominent Pol III ChIP peaks (Figure 6E). Consistent with these regulatory elements and Pol III-dependence, a previous study noted homology between *mir-4937* and a tRNA locus, mostly within putative Box A and Box B motifs (Roberts et al. 2013). Although *mir-4937* did not meet our stringent criteria for defining novel Microprocessor-independent small RNAs, its depletion in *pash-1(ts)* is marginal and not called as significant (Table S10), and its predicted secondary

structure suggests that its primary transcript would make a poor Microprocessor substrate (Figure S5C). Thus, *mir-4937* is likely a Pol III-transcribed Microprocessor- and Dicer-independent small RNA. *mir-4937* was not robustly immunoprecipitated with any of the *C. elegans* Argonautes by Seroussi, *et al*. (all average IP <0.2 reads per million) (Seroussi et al. 2023); thus, its functional class remains unclear. Nonetheless, previous studies identified it as a small RNA that increases during *C. elegans* aging (De Lencastre et al. 2010).

To determine whether miRNAs can derive from independent Pol III transcripts outside of *C. elegans,* we also examined ChIP datasets of Pol III subunits in three human cell lines: Hela, K562, and HEK293 (Oler et al., 2010; Dunham et al., 2012; Gerber et al., 2020, respectively). Manual curation of HEK293 RPC62 ChIP confirmed that *mir-4521* and *mir-4638* had strong Pol III ChIP signals. Whereas *mir-4638* is too lowly expressed to assess its biogenesis requirements, *mir-4521* shows evidence of Drosha and Dicer independence in HEK293 cells (Figure S5D) (Rybak-Wolf et al. 2014). Thus, evolutionarily young small RNAs may also arise from Pol III transcription and act as miRNAs in other lineages.

Discussion

In this study, we identified a new noncanonical miRNA biogenesis pathway that circumvents cleavage by the Microprocessor but relies on Dicer processing (Figures 1, 4). Using a variety of techniques, we have determined that the *mir-1829* family in *C. elegans* is derived from independent transcripts that are transcribed by Pol III (Figures 2, 3). Although these miRNA loci reside in the long introns of three Pol II-transcribed host genes, there is no detectable contribution of Pol II in *mir-1829* transcription (Figure 3F). Many classes of Pol III transcripts – including snaR-A, tRNAs, and SINEs – generate miRNAs via cleavage of a larger functional transcript (Babiarz et al. 2008; Martinez et al. 2017; Maute et al. 2013; Stribling et al. 2021). Unlike these

transcripts, the *mir-1829* family exists exclusively as independent miRNA transcripts. The small RNAs generated by these loci are the only detectable species by northern blot, demonstrating that the *mir-1829* family members are the primary product of these loci, thus differentiating this unique biogenesis pathway from previously identified Pol III-transcribed miRNAs.

Given that the Pol III-transcribed *mir-1829* family resides in the long introns of its Pol II-transcribed host genes, we speculate that there may be competition for Pol II and Pol III for the F20D1.3/*mir-1829b* locus. When the *mir-1829b* locus is expressed from an ectopic location in the *mir-1829 family(null)* background, *mir-1829b-5p* expression dramatically increases in comparison to levels in wild type (Figure 3C). Additionally, inhibiting F20D1.3 transcription by removing its promoter region – and thereby removing Pol II occupancy – results in a modest increase of *mir-1829b-5p* expression (Figure 3F). These results suggest that there is transcription interference of Pol III by Pol II at this locus. Previous studies have shown that Pol II can repress the transcription of tRNA genes (Lukoszek et al. 2013; Gerber et al. 2020). Since the *mir-1829* family is considered under the same class of Pol III transcripts as tRNAs (i.e. Type II), we hypothesize Pol II has an inhibitory effect on *mir-1829b* transcription. However, the relationship between Pol II and Pol III is complex as emerging reports suggest that Pol II can promote Pol III transcription and vice versa (Gerber et al. 2020; Jiang et al. 2022; Listerman et al. 2007; K C et al. 2024).

While we have demonstrated that the *mir-1829* family miRNAs require Pol III and Dicer for their biogenesis, additional required factors remain unidentified. We provide evidence that an initial cleavage step that matures the 3′ end of the precursor occurs in the nucleus: this end is modified by the nuclear nucleotidyltransferase GLDR-2 (Figure 5). If cleavage that generates the 5′ end of the precursor also occurs in the nucleus, then the precursor would be indistinguishable

from a canonical miRNA precursor, enabling nuclear export. Whether these cleavage events are coupled or separate events remains to be seen.

Building on previous work that characterized the expression patterns and binding complements of *C. elegans* Argonaute proteins, we show that the *mir-1829* family's expression is strongly germline enriched. Multiple lines of evidence suggest that *mir-1829b-5p* is very highly germline enriched, whereas a larger portion of *mir-1829c-5p* is expressed in the soma. An unusually large portion of the *mir-1829* family is loaded in the newly-characterized germline-specific Argonaute ALG-5. While two roles of ALG-5 in germline biology have been demonstrated, the targets of ALG-5 in these processes have yet to be discovered. Moreover, whether and how ALG-5 represses its targets is not yet known, and it has even been suggested to promote target expression (Liontis et al. 2023). We favor the model that ALG-5 represses its targets, given its strong homology with ALG-1/2 and their shared small RNA cargo, but direct targets must first be identified to test this model.

What is the contribution of the *mir-1829* family to animal physiology? Despite their enrichment in the germline, this miRNA family is not required for viability, germline function, or the robustness of the mitosis to meiosis transition. While these miRNAs are *C. elegans*-specific, they are preserved in the numerous wild isolates that have been collected around the world, suggesting that they may provide a selective advantage (Crombie et al. 2024). An interesting aspect of their future study will be to determine whether the 5p or 3p strands are the biologically important species. Although the 5p strands are more abundant (via qPCR calibrated with synthetic RNA standards, data not shown), the 3p strands have greater sequence conservation, including across the functional seed region.

We expanded our search for additional noncanonical and Pol III-dependent miRNAs by comparing datasets in which Microprocessor, Dicer, or Pol III function are inactivated. In *C. elegans*, we identify 1) misannotated and unannotated 26G and 21U RNAs, 2) a pair of identical Microprocessor-independent small RNAs (*mir-8196a/b*) whose biogenesis may rely on their close proximity to a 21U RNA and 3) a Pol III-dependent small RNA (*mir-4937*) that requires neither Microprocessor nor Dicer for its biogenesis. In human cells, we identify two candidates for Pol III-transcribed small RNAs that do not appear to be derived from a larger functional species, at least one of which is likely Microprocessor- and Dicer-independent. Notably, all of these newly identified noncanonical miRNAs are recently-evolved. Since miRNA loci are predicted to gradually evolve toward perfect biogenesis substrates (Berezikov 2011; Liu et al. 2008), these small RNAs and the *mir-1829* family may be substrates for further evolution to canonical miRNAs.

De novo miRNAs are likely to target many mRNAs, having potentially deleterious effects and preventing their retention in the genome. Chen and Rajewsky suggested a model for miRNA gene evolution in which miRNAs first gain a low level of expression, possibly in a tissue-restricted manner (Chen and Rajewsky 2007). Then, deleterious target sites can be purged, and finally, miRNA level and domain of expression can increase, providing a selective advantage by repressing the remaining target mRNAs. The *mir-1829* family and other identified noncanonical small RNAs like *mir-8196a/b* may be in the early stages of this trajectory, given their low level and tissue-restricted expression. Overall, these miRNAs fit multiple theoretical predictions of the characteristics of young de novo miRNA loci, thus representing a snapshot along the potential evolutionary trajectory to conserved canonical miRNA genes.

Materials and methods

C. elegans growth and maintenance

C. elegans were maintained on NGM plates seeded with OP50 at 20°C except where otherwise noted (Stiernagle 2006). For all experiments, embryos were harvested from gravid adults by hypochlorite treatment and washed thoroughly with M9 before hatching overnight in M9 supplemented with 1mM cholesterol at 20°C. Synchronized L1s were grown at 20°C to the appropriate stage as indicated. Strains and alleles used in this study are listed in Tables S1 and S2. For experiments using the auxin-inducible degron, synchronized L1s were grown for 48 h on NGM plates at 20°C, then washed onto 4mM auxin NGM plates at the L4 stage and grown for an additional 24 h at 20°C before harvesting.

CRISPR/Cas9-mediated genome editing

All strains generated in this study were made by CRISPR-mediated editing as previously described (Yang et al. 2020). Deletions of the *mir-1829* family members were generated by simultaneous CRISPR of two loci, in which one or both repair templates were integrated as indicated in Table S2.

Brood size and embryonic viability assays

Brood size assays were conducted at 25°C as previously described (Kotagama et al. 2024). Worms were moved onto a new plate every 24 h for four days following the L4 stage. After removing the parent hermaphrodite, the plates with embryos were incubated for another 24 h before assessing embryonic viability.

For *glp-1(ar202)* experiments, all strains were maintained at 15°C. A synchronized egg lay was performed for 3 h to isolate age-matched cohorts; these embryos were allowed to hatch at 15°C and then upshifted to 20°C 24 h after the synchronized egg lay. The worms were then kept at 20°C and single worms were followed to begin brood counts as above.

Immunoprecipitation

Synchronized L1s were grown at 20°C for approximately 65 h (to first day of adulthood) prior to pellet collection. Frozen pellets were resuspended with volume of 2x Lysis buffer corresponding to the volume of the pellet. 1x Lysis buffer [30 mM HEPES pH 7.4, 50mM KCl, 0.1% Triton X-100, 2 mM MgCl$_2$, 10% glycerol, 2 mM DTT, 80 units/mL of RNAse inhibitor, 1x cOmplete Mini protease inhibitor cocktail (Roche)] was added to bring volume of all the samples to 1mL. Samples were subjected to 4-6 rounds of sonication (10 cycles of 30s on, 30s off per round) in a Bioruptor Pico, then clarified by centrifugation at >15,000g for 20 min at 4°C. Pierce Reducing Agent Compatible BCA Protein Assay was used to quantify protein prior to IP. IP inputs were 1ml of lysate at 5mg/mL for ALG-5 IP or 2mg/mL for all other samples. Inputs were combined with 50µl of washed Anti-FLAG M2 Magnetic Bead slurry (Millipore Sigma) and incubated with gentle agitation at 4°C overnight. After three washes (500µl of 30mM 30 mM HEPES pH 7.4, 100mM KCl, 0.1% Triton X-100, 2 mM MgCl$_2$, 10% glycerol, 2 mM DTT, 80 units/mL of RNAse inhibitor, 1x cOmplete Mini protease inhibitor), beads were resuspended in 1ml of 1x lysis buffer. Input, supernatant, and IP samples were saved in 50µl aliquots which were either run on 4–20% TGX gels (Bio-Rad) or used for RNA extraction (see below).

RNA isolation

Frozen samples were resuspended with four pellet volumes of Trizol reagent (Fischer Scientific), then subjected to three freeze-thaw cycles on dry ice, followed by vortexing for 15 min at room temperature. RNA isolation was then performed according to the Trizol manufacturer's instructions.

Extra care was taken to recover an equal proportion of RNA from each sample from IP workflows (input, supernatant, and IP). To this end, 82µl dH$_2$O and 2.5µl yeast RNA (10mg/mL - Invitrogen) were added to 50µl aliquots of input, supernatant, or IP. 400µl Trizol LS (Thermo Fisher

Scientific) was added to each sample, followed by 15 min of vortexing at room temperature. 106.25µl chloroform was then added to each sample prior to a 10-min centrifugation at >15,000g. A consistent volume (200µl) of the aqueous phase was transferred to a new tube and mixed with an equal volume phenol: chloroform: isoamyl alcohol (25:24:1, pH=4.5). After 10-min centrifugation at >15,000g, 100µl of the aqueous phase was transferred to a new tube, and RNA was precipitated with one volume (100µl) isopropanol overnight at -20°C. Pellets were washed with 75% ethanol and resuspended in 28µl dH$_2$O.

Rapid amplification of cDNA ends (RACE)

5′ ends of primary *mir-1829* family transcripts were determined using 5′/3′ RACE Kit, 2nd Generation Kit (Sigma-Aldrich). For cDNA synthesis, 1µl of ~1µg/µl RNA sample was used. The amplification steps were carried out using Q5 High-Fidelity PCR Kit (NEB). For primers used, see Table S2. Purified PCR products were cloned into pCR™-Blunt II-TOPO vector using Zero Blunt® TOPO® PCR Cloning Kit (Invitrogen).

Taqman miRNA qPCR

For each sample, 1.66µl of total RNA at 200ng/µl was used per 5µl reverse transcription reaction using the TaqMan MicroRNA Reverse Transcription kit (ThermoFisher). Completed RT reactions were diluted 1:4 with dH$_2$O. Then, 1.33µl of diluted product was used in a 5µl qPCR reaction with Taqman miRNA probes and Taqman Universal Mastermix II with UNG (ThermoFisher) and were run in triplicate on the Applied Biosystems QuantStudio Pro 6.

Reverse transcription-quantitative polymerase chain reaction (RT-qPCR) for mRNAs

Reverse transcription qPCR of mRNA was performed using KAPA SYBR® FAST One-Step qRT-PCR Master Mix (2X) Kit (Fisher Scientific) and gene-specific primers on the Applied Biosystems

QuantStudio Pro 6. The primer sequences are listed in Table S2. For each reaction, 1.25µl of total RNA at 10ng/µl was used in a 5µl qPCR reaction, performing technical triplicate reactions.

Immunoblotting

Protein samples were prepared and quantified as previously described (Kotagama et al. 2024). Samples were run on 4-15% Mini-PROTEAN TGX gels (Bio-Rad) unless otherwise noted. All blots were blocked in TBST containing 5% BSA (AID experiments) or 5% milk (IP experiments) for 1 h at room temperature. Membranes were probed overnight at 4°C with primary antibody, followed by F(ab')2-Goat anti-Mouse IgG (H+L) Secondary Antibody, HRP (ThermoFisher) diluted 1:5000. Primary antibodies used were anti-FLAG M2 antibody (Sigma) diluted 1:1000, anti-Alpha Tubulin antibody DM1A (Abcam, ab7291) diluted 1:5000 (AID experiments) or AA4.3 (Developmental Studies Hybridoma Bank) diluted 1:5000 (IP experiments). ProtoGlow ECL (National Diagnostics) was used for signal detection of FLAG-tagged proteins, whereas Pierce ECL Western Blotting Substrate (Thermo Scientific) was used for signal detection of tubulin. Signal intensity was measured using Amersham Imager 680 (GE) and quantified using Image Studio Lite. For re-probing, membranes were stripped at 50°C for 45 min with stripping buffer (2% SDS, 62.5 mM Tris HCl pH 6.8, 0.8% β-mercaptoethanol).

Northern blot

Total RNA was separated on 15% urea gel (1.5mm) and transferred to Amersham Hybond- NX membranes. Membranes were crosslinked using 0.16M l-ethyl-3-(3-dimethylaminopropyl) carbodiimide (EDC) (Sigma) in 0.13M 1-methylimidazole, pH=8 at 60°C for 30 min as previously described (Pall et al. 2007). DNA oligos complementary to miRNAs of interest and U6 were 5′ end-labelled with ATP-γ-[^{32}P] unless otherwise noted. For Microprocessor depletion experiments, 25µg of RNA was loaded and stringent probing conditions were used: blots were hybridized

overnight in hybridization buffer (7% SDS, 0.2 M NaPO$_4$ pH=7.2 at 42°C and washed four times

for 10 min at 42°C (2x with 2 x SSPE, 0.1% SDS, 1x with 1 x SSPE, 0.1% SDS and 1x with 0.5

x SSPE, 0.1% SDS) before exposing blots. For Dicer depletion experiments, 10μg RNA was

loaded, and low stringency probing conditions were used: blots were hybridized overnight in

ExpressHyb Hybridization Solution (Takara Bio USA) at 32°C and washed once with 2x SSC +

0.1% SDS at 32°C for 10 min and then once more with 1x SSC + 0.1% SDS at 32°C for another

10 min before exposing blots. In between each probe, membranes were stripped by incubating in

a solution of boiling 0.04% SDS at room temperature for 20 min, with shaking, for a total of three

times.

RNA Structure Predictions

Structure of the primary and precursor sequences of *mir-1829* family members were predicted

using the minimum free energy (MFE) structure on RNAfold (Gruber et al. 2008; Lorenz et al.

2011). The Vienna format output for the primary structure was loaded into RNAcanvas (Johnson

and Simon 2023) and the predicted substructure of the precursor was applied when this region

differed in the primary transcript structure prediction.

Small RNA-seq sample preparation and analysis

Library preparation was performed using the NEBNext Small RNA Library Prep Set for Illumina

as previously described (Donnelly et al. 2022). Pooled samples were sequenced on a NextSeq 2000.

Raw sequence data is accessible under GEO accession number GSE282009. Briefly, sequence

analysis was performed on the NIH High Performance Computing Cluster. Libraries were trimmed

using Cutadapt 4.4 (Martin 2011), mapped using bowtie2 2.5.1 with settings '–no-unal –end-to-

end –sensitive' (Langmead and Salzberg 2012), and the resulting bam files were sorted and

indexed using samtools 1.19 (Danecek et al. 2021). Reads were assigned to miRNAs using htseq

2.0.4 with settings '--mode union --nonunique fraction -a 0' (Anders et al. 2015). Analysis of miRNA expression using DESeq2 was performed similarly to previously described (Kotagama et al. 2024), with the modification that size factors were calculated by dividing spike-in reads in each sample by the mean spike-in reads per sample. Human small RNA datasets were analyzed similarly, except that reads were trimmed and filtered for quality using Trimgalore 0.6.7 (DOI 10.5281/zenodo.5127898) with default settings.

GLDR-2 Imaging

Adult worms raised at 15°C were mounted in 15μl of M9 with 5mM levamisole on a 2% agarose pad. Slides were cover slipped then imaged on a Nikon C2 confocal microscope using a 60x water immersion objective.

Pol III ChIP analysis

C. elegans Pol III ChIP peaks were accessed at ENCODE (Accession file ENCFF924VCM) (Araya et al. 2014). For human ChIP data analysis, low quality bases and adaptors were trimmed from raw sequence reads with cutadapt v2.7 using 12 parameters -q 20 -a AGATCGGAAGAGC -minimum-length 25. Trimmed reads were aligned to the human GRCh38.p14 genome assembly using Bowtie2 v2.3.5 (Langmead and Salzberg 2012). Multimapping reads were removed with samtools v1.9 (Li et al. 2009) using the "view" subcommand and the additional argument "-q 20", and duplicate reads were removed with the Picard v2.21.4 (http://broadinstitute.github.io/picard/index.html) Mark Duplicates tool. Peaks were called for each replicate individually using macs v2.2.7.1 (Zhang et al. 2008). Pooled peaks were called by providing multiple BAMs of all the replicates to macs2. miRNAs containing prominent Pol III peaks were called using bedtools intersect with the miRBase annotation file for the respective species (Quinlan and Hall 2010).

Competing Interest Statement

The authors have no competing interests to declare.

Acknowledgments

We thank members of the McJunkin lab and Baltimore Worm Club for helpful discussions and WormBase. Some of the strains used in this study were provided by the Caenorhabditis Genetics Center (CGC), which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440). We thank Eleanor Maine and Tim Schedl for helpful suggestions. This work supported by the NIDDK IRP (ZIADK075147).

Author Contributions

R.S. and K.M. conceived the study. R.S. acquired, analyzed, and interpreted the data. K.F.V. generated reagents and acquired data. D.H. and A.Z. generated reagents. R.S., G.Y., and K.M analyzed primary and published deep sequencing data. R.S. and K.M wrote the manuscript.

References

Aalto AP, Nicastro IA, Broughton JP, Chipman LB, Schreiner WP, Chen JS, Pasquinelli AE. 2018. Opposing roles of microRNA Argonautes during Caenorhabditis elegans aging. *PLoS Genet* **14**.

Alberti C, Manzenreither RA, Sowemimo I, Burkard TR, Wang J, Mahofsky K, Ameres SL, Cochella L. 2018. Cell-type specific sequencing of microRNAs from complex animal tissues. *Nat Methods* **15**: 283–289.

Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**: 166–169.

Araya CL, Kawli T, Kundaje A, Jiang L, Wu B, Vafeados D, Terrell R, Weissdepp P, Gevirtzman L, MacE D, et al. 2014. Regulatory analysis of the C. Elegans genome with spatiotemporal resolution. *Nature* **512**: 400–405.

Auyeung VC, Ulitsky I, McGeary SE, Bartel DP. 2013. Beyond secondary structure: primary-sequence determinants license pri-miRNA hairpins for processing. *Cell* **152**: 844–858.

Babiarz JE, Ruby JG, Wang Y, Bartel DP, Blelloch R. 2008. Mouse ES cells express endogenous shRNAs, siRNAs, and other Microprocessor-independent, Dicer-dependent small RNAs. *Genes Dev* **22**: 2773–2785.

Behlke MA, Dames SA, McDonald WH, Gould KL, Devor EJ, Walder JA. 2000. Use of high specific activity StarFire oligonucleotide probes to visualize low-abundance pre-mRNA splicing intermediates in S. pombe. *Biotechniques* **29**: 892–897.

Berezikov E. 2011. Evolution of microRNA diversity and regulation in animals. *Nat Rev Genet* **12**: 846–860.

Berezikov E, Chung WJ, Willis J, Cuppen E, Lai EC. 2007. Mammalian mirtron genes. *Mol Cell* **28**: 328–336.

Bortolamiol-Becet D, Hu F, Jee D, Wen J, Okamura K, Lin CJ, Ameres SL, Lai EC. 2015. Selective Suppression of the Splicing-Mediated MicroRNA Pathway by the Terminal Uridyltransferase Tailor. *Mol Cell* **59**: 217–228.

Brenner JL, Jyo EM, Mohammad A, Fox P, Jones V, Mardis E, Schedl T, Maine EM. 2022. TRIM-NHL protein, NHL-2, modulates cell fate choices in the C. elegans germ line. *Dev Biol* **491**: 43–55.

Brosnan CA, Palmer AJ, Zuryn S. 2021. Cell-type-specific profiling of loaded miRNAs from Caenorhabditis elegans reveals spatial and temporal flexibility in Argonaute loading. *Nat Commun* **12**: 2194.

Brown KC, Svendsen JM, Tucci RM, Montgomery BE, Montgomery TA. 2017. ALG-5 is a miRNA-associated Argonaute required for proper developmental timing in the Caenorhabditis elegans germline. *Nucleic Acids Res* **45**: 9093–9107.

Cheloufi S, Dos Santos CO, Chong MMW, Hannon GJ. 2010. A dicer-independent miRNA biogenesis pathway that requires Ago catalysis. *Nature* **465**: 584–589.

Chen K, Rajewsky N. 2007. The evolution of gene regulation by transcription factors and microRNAs. *Nat Rev Genet* **8**: 93–103.

Chung WJ, Agius P, Westholm JO, Chen M, Okamura K, Robine N, Leslie CS, Lai EC. 2011. Computational and experimental identification of mirtrons in Drosophila melanogaster and Caenorhabditis elegans. *Genome Res* **21**: 286–300.

Cifuentes D, Xue H, Taylor DW, Patnode H, Mishima Y, Cheloufi S, Ma E, Mane S, Hannon GJ, Lawson ND, et al. 2010. A novel miRNA processing pathway independent of Dicer requires Argonaute2 catalytic activity. *Science* **328**: 1694–1698.

Consalvo CD, Aderounmu AM, Donelick HM, Aruscavage PJ, Eckert DM, Shen PS, Bass BL. 2024. Caenorhabditis elegans Dicer acts with the RIG-I-like helicase DRH-1 and RDE-4 to cleave dsRNA. *Elife* **13**.

Crombie TA, McKeown R, Moya ND, Evans KS, Widmayer SJ, LaGrassa V, Roman N, Tursunova O, Zhang G, Gibson SB, et al. 2024. CaeNDR, the Caenorhabditis Natural Diversity Resource. *Nucleic Acids Res* **52**: D850–D858.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T, McCarthy SA, Davies RM. 2021. Twelve years of SAMtools and BCFtools. *Gigascience* **10**.

De Lencastre A, Pincus Z, Zhou K, Kato M, Lee SS, Slack FJ. 2010. MicroRNAs both promote and antagonize longevity in C. elegans. *Curr Biol* **20**: 2159–2168.

Denli AM, Tops BBJ, Plasterk RHA, Ketting RF, Hannon GJ. 2004. Processing of primary microRNAs by the Microprocessor complex. *Nature* **432**: 231–235.

Dexheimer PJ, Wang J, Cochella L. 2020. Two MicroRNAs Are Sufficient for Embryonic Patterning in C. elegans. *Curr Biol* **30**: 5058-5065.e5.

Diag A, Schilling M, Klironomos F, Ayoub S, Rajewsky N. 2018. Spatiotemporal m(i)RNA Architecture and 3′ UTR Regulation in the C. elegans Germline. *Dev Cell* **47**: 785-800.e8.

Dieci G, Fiorino G, Castelnuovo M, Teichmann M, Pagano A. 2007. The expanding RNA polymerase III transcriptome. *Trends Genet* **23**: 614–622.

Donnelly BF, Yang B, Grimme AL, Vieux K-F, Liu C-Y, Zhou L, McJunkin K. 2022. The developmentally timed decay of an essential microRNA family is seed-sequence dependent. *Cell Rep* **40**: 111154.

Duchaine TF, Wohlschlegel JA, Kennedy S, Bei Y, Conte D, Pang KM, Brownell DR, Harding S, Mitani S, Ruvkun G, et al. 2006. Functional Proteomics Reveals the Biochemical Niche of C. elegans DCR-1 in Multiple Small-RNA-Mediated Pathways. *Cell* **124**: 343–354.

Dunham I, Kundaje A, Aldred SF, Collins PJ, Davis CA, Doyle F, Epstein CB, Frietze S, Harrow J, Kaul R, et al. 2012. An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**: 57–74.

Ender C, Krek A, Friedländer MR, Beitzinger M, Weinmann L, Chen W, Pfeffer S, Rajewsky N, Meister G. 2008. A human snoRNA with microRNA-like functions. *Mol Cell* **32**: 519–528.

Fang W, Bartel DP. 2015. The Menu of Features that Define Primary MicroRNAs and Enable De Novo Design of MicroRNA Genes. *Mol Cell* **60**: 131–145.

Fromm B, Domanska D, Høye E, Ovchinnikov V, Kang W, Aparicio-Puerta E, Johansen M, Flatmark K, Mathelier A, Hovig E, et al. 2020. MirGeneDB 2.0: the metazoan microRNA complement. *Nucleic Acids Res* **48**: D132.

Gebert LFR, MacRae IJ. 2019. Regulation of microRNA function in animals. *Nat Rev Mol Cell Biol* **20**: 21–37.

Gerber A, Ito K, Chu CS, Roeder RG. 2020. Gene-Specific Control of tRNA Expression by RNA Polymerase II. *Mol Cell* **78**: 765–778.

Gregory RI, Yan KP, Amuthan G, Chendrimada T, Doratotaj B, Cooch N, Shiekhattar R. 2004. The Microprocessor complex mediates the genesis of microRNAs. *Nature* **432**: 235–240.

Grishok A, Pasquinelli AE, Conte D, Li N, Parrish S, Ha I, Baillie DL, Fire A, Ruvkun G, Mello CC. 2001. Genes and mechanisms related to RNA interference regulate expression of the small temporal RNAs that control C. elegans developmental timing. *Cell* **106**: 23–34.

Gruber AR, Lorenz R, Bernhart SH, Neuböck R, Hofacker IL. 2008. The Vienna RNA websuite. *Nucleic Acids Res* **36**.

Gutiérrez-Pérez P, Santillán EM, Lendl T, Wang J, Schrempf A, Steinacker TL, Asparuhova M, Brandstetter M, Haselbach D, Cochella L. 2021. miR-1 sustains muscle physiology by controlling V-ATPase complex assembly. *Sci Adv* **7**: eabh1434.

Ha M, Kim VN. 2014. Regulation of microRNA biogenesis. *Nat Rev Mol Cell Biol* **15**: 509–524.

Han J, Lee Y, Yeom KH, Kim YK, Jin H, Kim VN. 2004. The Drosha-DGCR8 complex in primary microRNA processing. *Genes Dev* **18**: 3016–3027.

Hansen TB, Venø MT, Jensen TI, Schaefer A, Damgaard CK, Kjems J. 2016. Argonaute-associated short introns are a novel class of gene regulators. *Nat Commun* **7**: 11538.

Hussain S, Sajini AA, Blanco S, Dietmann S, Lombard P, Sugimoto Y, Paramor M, Gleeson JG, Odom DT, Ule J, et al. 2013. NSun2-mediated cytosine-5 methylation of vault noncoding RNA determines its processing into regulatory small RNAs. *Cell Rep* **4**: 255–261.

Hutvágner G, McLachlan J, Pasquinelli AE, Bálint É, Tuschl T, Zamore PD. 2001. A cellular function for the RNA-interference enzyme Dicer in the maturation of the let-7 small temporal RNA. *Science* **293**: 834–838.

Ikegami K, Lieb JD. 2013. Integral Nuclear Pore Proteins Bind to Pol III-Transcribed Genes and Are Required for Pol III Transcript Processing in C. elegans. *Mol Cell* **51**: 840–849.

Jan CH, Friedman RC, Ruby JG, Bartel DP. 2011. Formation, Regulation and Evolution of Caenorhabditis elegans 3′UTRs. *Nature* **469**: 97.

Jiang Y, Huang J, Tian K, Yi X, Zheng H, Zhu Y, Guo T, Ji X. 2022. Cross-regulome profiling of RNA polymerases highlights the regulatory role of polymerase III on mRNA transcription by maintaining local chromatin architecture. *Genome Biol* **23**.

Jin W, Wang J, Liu C-P, Wang H-W, Xu R-M. 2020. Structural Basis for pri-miRNA Recognition by Drosha. *Mol Cell* **78**: 423-433.e5.

Johnson PZ, Simon AE. 2023. RNAcanvas: interactive drawing and exploration of nucleic acid structures. *Nucleic Acids Res* **51**: W501–W508.

K C R, Cheng R, Zhou S, Lizarazo S, Smith DJ, Van Bortle K. 2024. Evidence of RNA polymerase III recruitment and transcription at protein-coding gene promoters. *Mol Cell* **84**: 4111-4124.e5.

Kang W, Fromm B, Houben AJ, Høye E, Bezdan D, Arnan C, Thrane K, Asp M, Johnson R, Biryukova I, et al. 2021. MapToCleave: High-throughput profiling of microRNA biogenesis in living cells. *Cell Rep* **37**: 110015.

Kim K, Baek SC, Lee Y-Y, Bastiaanssen C, Kim J, Kim H, Kim VN. 2021. A quantitative map of human primary microRNA processing sites. *Mol Cell* **81**: 3422-3439.e11.

Kim YK, Kim B, Kim VN. 2016. Re-evaluation of the roles of DROSHA, Export in 5, and DICER in microRNA biogenesis. *Proc Natl Acad Sci U S A* **113**: E1881–E1889.

Knight SW, Bass BL. 2001. A role for the RNase III enzyme DCR-1 in RNA interference and germ line development in Caenorhabditis elegans. *Science* **293**: 2269–2271.

Kotagama K, Grimme AL, Braviner L, Yang B, Sakhawala RM, Yu G, Benner LK, Joshua-Tor L, McJunkin K. 2024. Catalytic residues of microRNA Argonautes play a modest role in microRNA star strand destabilization in C. elegans. *Nucleic Acids Res* **52**: 4985–5001.

Kozomara A, Birgaoanu M, Griffiths-Jones S. 2019. miRBase: from microRNA sequences to function. *Nucleic Acids Res* **47**: D155–D162.

Kuscu C, Kumar P, Kiran M, Su Z, Malik A, Dutta A. 2018. tRNA fragments (tRFs) guide Ago to regulate gene expression post-transcriptionally in a Dicer-independent manner. *RNA* **24**: 1093–1105.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods* **9**: 357–359.

Lee Y, Ahn C, Han J, Choi H, Kim J, Yim J, Lee J, Provost P, Rådmark O, Kim S, et al. 2003. The nuclear RNase III Drosha initiates microRNA processing. *Nature* **425**: 415–419.

Lehrbach NJ, Castro C, Murfitt KJ, Abreu-Goodger C, Griffin JL, Miska EA. 2012. Post-developmental microRNA expression is required for normal physiology, and regulates aging in parallel to insulin/IGF-1 signaling in C. elegans. *Rna* **18**: 2220–2235.

Lemus-Diaz N, Ferreira RR, Bohnsack KE, Gruber J, Bohnsack MT. 2020. The human box C/D snoRNA U3 is a miRNA source and miR-U3 regulates expression of sortin nexin 27. *Nucleic Acids Res* **48**: 8074–8089.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R. 2009. The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**: 2078–2079.

Liontis T, Verma K, Grishok A. 2023. DOT-1.1 (DOT1L) deficiency in C. elegans leads to small RNA-dependent gene activation. *BBA advances* **3**.

Listerman I, Bledau AS, Grishina I, Neugebauer KM. 2007. Extragenic accumulation of RNA polymerase II enhances transcription by RNA polymerase III. *PLoS Genet* **3**: 2268–2277.

Liu N, Okamura K, Tyler DM, Phillips MD, Chung WJ, Lai EC. 2008. The evolution and functional diversification of animal microRNA genes. *Cell Res* **18**: 985–996.

Lorenz R, Bernhart SH, Höner zu Siederdissen C, Tafer H, Flamm C, Stadler PF, Hofacker IL. 2011. ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**.

Lu R, Yigit E, Li WX, Ding SW. 2009. An RIG-I-Like RNA Helicase Mediates Antiviral RNAi Downstream of Viral siRNA Biogenesis in Caenorhabditis elegans. *PLoS Pathog* **5**: 1000286.

Lukoszek R, Mueller-Roeber B, Ignatova Z. 2013. Interplay between polymerase II- and polymerase III-assisted expression of overlapping genes. *FEBS Lett* **587**: 3692–3695.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* **17**: 10–12.

Martinez G, Choudury SG, Slotkin RK. 2017. tRNA-derived small RNAs target transposable element transcripts. *Nucleic Acids Res* **45**: 5142–5152.

Maute RL, Schneider C, Sumazin P, Holmes A, Califano A, Basso K, Dalla-Favera R. 2013. tRNA-derived microRNA modulates proliferation and the DNA damage response and is down-regulated in B cell lymphoma. *Proc Natl Acad Sci U S A* **110**: 1404–1409.

Morgan PG, Sedensky MM. 1994. Mutations conferring new patterns of sensitivity to volatile anesthetics in Caenorhabditis elegans. *Anesthesiology* **81**: 888–898.

Nguyen TL, Nguyen TD, Ngo MK, Nguyen TA. 2023. Dissection of the Caenorhabditis elegans Microprocessor. *Nucleic Acids Res* **51**: 1512–1527.

Okamura K, Hagen JW, Duan H, Tyler DM, Lai EC. 2007. The mirtron pathway generates microRNA-class regulatory RNAs in Drosophila. *Cell* **130**: 89–100.

Oler AJ, Alla RK, Roberts DN, Wong A, Hollenhorst PC, Chandler KJ, Cassiday PA, Nelson CA, Hagedorn CH, Graves BJ, et al. 2010. Human RNA polymerase III transcriptomes and relationships to Pol II promoter chromatin and enhancer-binding factors. *Nat Struct Mol Biol* **17**: 620–628.

Pall GS, Codony-Servat C, Byrne J, Ritchie L, Hamilton A. 2007. Carbodiimide-mediated cross-linking of RNA to nylon membranes improves the detection of siRNA, miRNA and piRNA by northern blot. *Nucleic Acids Res* **35**: e60.

Partin AC, Zhang K, Jeong B-C, Herrell E, Li S, Chiu W, Nam Y. 2020. Cryo-EM Structures of Human Drosha and DGCR8 in Complex with Primary MicroRNA. *Mol Cell* **78**: 411-422.e4.

Persson H, Kvist A, Vallon-Christersson J, Medstrand P, Borg A, Rovira C. 2009. The non-coding RNA of the multidrug resistance-linked vault particle encodes multiple regulatory small RNAs. *Nat Cell Biol* **11**: 1268–1271.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features. *Bioinformatics* **26**: 841–842.

Reimão-Pinto MM, Ignatova V, Burkard TR, Hung JH, Manzenreither RA, Sowemimo I, Herzog VA, Reichholf B, Fariña-Lopez S, Ameres SL. 2015. Uridylation of RNA Hairpins by Tailor Confines the Emergence of MicroRNAs in Drosophila. *Mol Cell* **59**: 203–216.

Roberts JT, Cooper EA, Favreau CJ, Howell JS, Lane LG, Mills JE, Newman DC, Perry TJ, Russell ME, Wallace BM, et al. 2013. Continuing analysis of microRNA origins: Formation from transposable element insertions and noncoding RNA mutations. *Mob Genet Elements* **3**: e27755.

Ruby JG, Jan CH, Bartel DP. 2007. Intronic microRNA precursors that bypass Drosha processing. *Nature* **448**: 83–86.

Rybak-Wolf A, Jens M, Murakawa Y, Herzog M, Landthaler M, Rajewsky N. 2014. A variety of dicer substrates in human and C. elegans. *Cell* **159**: 1153–1167.

Seroussi U, Lugowski A, Wadi L, Lao RX, Willis AR, Zhao W, Sundby AE, Charlesworth AG, Reinke AW, Claycomb JM. 2023. A comprehensive survey of c. Elegans argonaute proteins reveals organismwide gene regulatory networks and functions. *Elife* **12**.

Sheng P, Fields C, Aadland K, Wei T, Kolaczkowski O, Gu T, Kolaczkowski B, Xie M. 2018. Dicer cleaves 5'-extended microRNA precursors originating from RNA polymerase II transcription start sites. *Nucleic Acids Res* **46**: 5737–5752.

Simon DJ, Madison JM, Conery AL, Thompson-Peer KL, Soskis M, Ruvkun GB, Kaplan JM, Kim JK. 2008. The microRNA miR-1 regulates a MEF-2-dependent retrograde signal at neuromuscular junctions. *Cell* **133**: 903–915.

Stiernagle T. 2006. Maintenance of C. elegans. *WormBook* 1–11.

Stribling D, Lei Y, Guardia CM, Li L, Fields CJ, Nowialis P, Opavsky R, Renne R, Xie M. 2021. A noncanonical microRNA derived from the snaR-A noncoding RNA targets a metastasis inhibitor. *RNA* **27**: 694–709.

Stutzman A V., Liang AS, Beilinson V, Ikegami K. 2020. Transcription-independent TFIIIC-bound sites cluster near heterochromatin boundaries within lamina-associated domains in C. elegans. *Epigenetics Chromatin* **13**: 1.

Tabara H, Yigit E, Siomi H, Mello CC. 2002. The dsRNA Binding Protein RDE-4 Interacts with RDE-1, DCR-1, and a DExH-Box Helicase to Direct RNAi in C. elegans. *Cell* **109**: 861–871.

Vasquez-Rifo A, Jannot G, Armisen J, Labouesse M, Bukhari SIA, Rondeau EL, Miska EA, Simard MJ. 2012. Developmental characterization of the microRNA-specific C. elegans Argonautes alg-1 and alg-2. *PLoS One* **7**.

Vieux KF, Prothro KP, Kelley LH, Palmer C, Maine EM, Veksler-Lublinsky I, McJunkin K. 2021. Screening by deep sequencing reveals mediators of microRNA tailing in C. elegans. *Nucleic Acids Res* **49**: 11167–11180.

Westholm JO, Lai EC. 2011. Mirtrons: microRNA biogenesis via splicing. *Biochimie* **93**: 1897–1904.

Xie M, Li M, Vilborg A, Lee N, Shu M Di, Yartseva V, Šestan N, Steitz JA. 2013. Mammalian 5′-Capped MicroRNA Precursors that Generate a Single MicroRNA. *Cell* **155**: 1568–1580.

Xu W, Liu J, Qi H, Si R, Zhao Z, Tao Z, Bai Y, Hu S, Sun X, Cong Y, et al. 2024. A lineage-resolved cartography of microRNA promoter activity in C. elegans empowers multidimensional developmental analysis. *Nat Commun* **15**: 2783.

Yang B, Schwartz M, McJunkin K. 2020. In vivo CRISPR screening for phenotypic targets of the mir-35-42 family in C. elegans. *Genes Dev* **34**: 1227–1238.

Yang S, Maurin T, Robine N, Rasmussen KD, Jeffrey KL, Chandwani R, Papapetrou EP, Sadelain M, O'Carroll D, Lai EC. 2010. Conserved vertebrate mir-451 provides a platform for Dicer-independent, Ago2-mediated microRNA biogenesis. *Proc Natl Acad Sci U S A* **107**: 15163–15168.

Zamudio JR, Kelly TJ, Sharp PA. 2014. Argonaute-bound small RNAs from promoter-proximal RNA polymerase II. *Cell* **156**: 920–934.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nussbaum C, Myers RM, Brown M, Li W, et al. 2008. Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**.

Figure 1. The *mir-1829* family is Microprocessor-independent. A) Comparison of log₂(fold change) of miRNAs 24h post-upshift to restrictive temperature in *pash-1(ts)* vs. wild type (Vieux et al. 2021) (x-axis) to those in RNAiD depletion of the Microprocessor (Dexheimer et al. 2020) (y-axis). miRNAs are annotated as having a "short hairpin" if predicted structure on MirGeneDB has <32bp (Fromm et al. 2020). A small number of miRNAs are insensitive to Microprocessor depletion in both datasets, including mirtrons and the *mir-1829* family. Only miRNAs with baseMean ≥ 10 in the *pash-1(ts)* data set are plotted. B) Schematics of miRNA structures emphasizing long basal stem of canonical miRNAs and lack thereof in mirtrons and the *mir-1829* family. C) Cleavage efficiency of Drosha (x-axis) or Microprocessor (y-axis) as reported (Nguyen et al. 2023). The *mir-1829* family is among the least favorable substrates. D) Predicted secondary

structures of the *pri-mir-1829b* and *pri-mir-1829c.* E) Northern blot of total RNA from adult animals exposed to auxin for 24h starting at the L4 stage. *mir-1829b-5p*, *mir-1829c-5p*, and U6 probes were 5′ end-labelled, whereas *mir-1* probe was labelled according to Starfire method (Behlke et al. 2000). F) Same samples used for northern blotting were assayed by RT-qPCR. miRNA expression is normalized to a small RNA control (*sn2429*), then further normalized to wild type samples. Mean and standard deviation of two biological replicates shown. Two-way ANOVA showed no significant changes.

Figure 2. The *mir-1829* family shows evidence of differential expression from its host genes in the germline. A) Schematic of relative expression levels of the *mir-1829* family and its host genes in the *C. elegans* germline. Data from SPACEGERM (Diag et al. 2018) indicates that the *mir-1829* family (pink) peaks in expression in the distal germline, whereas its host genes (black) peak in expression in the proximal germline. B) Prominent Pol III ChIP-seq peaks (Araya et al. 2014) reside at the *mir-1829* loci (pink). Arrowheads: 5′ RACE reads cloned from RNA of indicated stage. Putative Pol III promoter sequence motifs upstream of the miRNA precursors are indicated.

Figure 3. *mir-1829* family members are transcribed by Pol III. A) MUSCLE alignment of the *mir-1829* family putative Pol III promoter sequence motifs. B) Schematic of CRISPR alleles generated at the *mir-1829b* locus (left) and qPCR of resulting miRNA and host gene expression (right). C) Schematic of genomic sources of *mir-1829b* (left), including re-integration of a 510bp minimal *mir-1829b* transcriptional unit in a *mir-1829 family(null)* mutant background (bottom left). qPCR of *mir-1829b* expression (right). D) qPCR of miRNAs and host gene transcripts in AID-tagged RPC-1 strain with ubiquitous TIR1 (MCJ666) or TIR1 alone (MLC1040). Auxin treatment was 24h, beginning at the L4 stage. E) Schematics of promoter deletions for two host genes of the *mir-1829* family. F) qPCR in host gene promoter deletion samples. B-D, F) miRNA expression is normalized to a small RNA control (*sn2429*) and then further normalized to wild type levels. Host gene expression was normalized to GAPDH (*gpd-1*) and then to wild type levels. Mean and SD shown. Two-way ANOVA. ****$p<0.0001$, *$p<0.05$.

Figure 4. The *mir-1829* family is Dicer-dependent. (A-C) All samples were exposed to auxin for 24h starting at the L4 stage, then harvested as adults. A) Northern blot using low stringency conditions (see methods). Right panels are lanes loaded with synthetic RNA oligos, demonstrating recognition of both *mir-1829b-5p* and *mir-1829c-5p* with the *mir-1829b-5p* probe under these conditions. B) miRNA qPCR normalized to a small RNA control (*sn2429*) and then further normalized to levels in control that only carries Dicer AID tag (UY212). Mean and SD shown. Two-way ANOVA. ****$p<0.0001$ C) Small RNA-seq. MA plot showing average abundance on x-axis and $\log_2$(fold change) in DCR-1::AID; ubiquitous TIR1 (MCJ387) compared to DCR-1::AID tag alone (UY212) on y-axis. miRNAs showing significant sensitivity to Dicer depletion are shown in blue-green, and the *mir-1829* family is highlighted in red open circles. Three biological replicates of each genotype were analyzed using DESeq2.

Figure 5. The *mir-1829* family primary transcripts are likely cleaved in the nucleus. A) Percent untemplated 3′ nucleotide additions (tailing) of the *mir-1829* family members in wild type. B) Percent A-tailing in wild type and *gldr-2(null)* (top). Abundance of each mature species shown below (bottom). (A-B) All *mir-1829* family member strands with >1RPM average abundance are shown. C) Representative images of 3xFLAG::AID::GFP::GLDR-2. D) Model for new miRNA biogenesis pathway of the *mir-1829* family in contrast to canonical miRNA biogenesis.

Figure 6. Noncanonical miRNA biogenesis pathway extends beyond the *mir-1829* family. A-C) Small RNA-seq results in samples depleted for 24h of functional (A) PASH-1 (C) Dicer (DCR-1) or (D) RPC-1. All miRBase annotations are plotted, and relevant noncanonical small RNAs are highlighted. B) Genome browser tracks showing proximity of *mir-8196a* to an upstream unannotated 21U RNA. 5′-independent small RNA cloning following polyphosphatase treatment also shows 22G RNAs generated from the locus. *mir-8196a* but not the 21U shows Dicer-dependence. E) Genome browser track showing RPC-1 (a Pol III subunit) ChIP at the *mir-4937* locus. F) Genome browser tracks showing RPC62 (Pol III) ChIP in HEK293 cells at the *mir-4638* and *mir-4521* loci.