1

3
4
5
6   **Vascular endothelial-specific loss of TGF-beta signaling as a model for**
7   **choroidal neovascularization and central nervous system vascular inflammation**
8
9
10
11   Yanshu Wang[1,4], Amir Rattner[1,4], Zhongming Li[1,4], Philip M. Smallwood[1,4], and Jeremy Nathans[1,2,3,4]
12
13
14   [1]Department of Molecular Biology and Genetics, Johns Hopkins University School of Medicine,
15   Baltimore, United States
16
17   [2]Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, United States.
18
19   [3]Department of Ophthalmology, Johns Hopkins University School of Medicine, Baltimore, United States.
20
21   [4]Howard Hughes Medical Institute, Johns Hopkins University School of Medicine, Baltimore, United
22   States.
23
24
Number of pages: 26
Number of figures: 9
Number of tables: 0
Supplemental text pages: 4
Number of supplemental figures: 18
Number of supplemental tables: 2
31
32
Address for editorial correspondence:
Dr. Jeremy Nathans
805 PCTB
725 North Wolfe Street
Johns Hopkins University School of Medicine
Baltimore, MD 21205
tel: 410 955 4679
fax: 410 614 0827
email: jnathans@jhmi.edu
42
43
44   **Conflict of interest statement**
45   The authors have declared that no conflict of interest exists.
46
47
48

49
50 **Abstract**
51

In mice, postnatal endothelial cell (EC)-specific knockout of the genes coding for Transforming Growth
Factor-Beta Receptor (TGFBR)1 and/or TGFBR2 eliminates TGF-beta signaling in vascular ECs and leads
to distinctive central nervous system (CNS) vascular phenotypes. Knockout mice exhibit (1) reduced
intra-retinal vascularization, (2) choroidal neovascularization with occasional anastomoses connecting
choroidal and intraretinal vasculatures, (3) infiltration of diverse immune cells into the retina,
including macrophages, T-cells, B-cells, NK cells, and dendritic cells, (4) a close physical association
between immune cells and retinal vasculature, (5) a pro-inflammatory transcriptional state in CNS ECs,
with increased ICAM1 immunoreactivity, and (6) increased Smooth Muscle Actin immunostaining in
CNS pericytes. Comparisons of the retinal phenotype with two other genetic models of retinal
hypovascularization – loss of Norrin/Fzd4 signaling and loss of VEGF signaling – shows that the
immune cell infiltrate is greatest with loss of TGF-beta signaling, more modest with loss of Norrin/Fzd4
signaling, and undetectable with loss of VEGF signaling. The phenotypes caused by loss of TGF-beta
signaling in ECs recapitulate some of the cardinal features of retinal and neurologic diseases associated
with vascular inflammation. These observations suggest that therapies that promote TGF-beta-
dependent anti-inflammatory responses in ECs could represent a promising strategy for disease
modulation.
68
69
70
71 **Introduction**
72
In the central nervous system (CNS), vascular development, structure, and function are tightly regulated
by multiple signaling pathways, including the vascular endothelial growth factor (VEGF), Wnt, Notch, and
TGF-beta pathways (Wälchli et al., 2015, 2023; Rattner et al., 2022). These signals control angogenesis,
vascular remodeling, and vascular permeability. With respect to permeability, the vasculature serving
most of the CNS is distinguished from non-CNS vasculature by greatly reduced permeability, a
specialization refered to as the blood-brain barrier (BBB) or, in the retina, the blood-retina barrier (BRB).
79
Disorders of CNS vascular structure and function are a major cause of morbidity and mortality (Ropper
et al., 2023). These include arteriovenous malformations, vasculitis, atherosclerosis, small vessel
disease, moyamoya, neovascular age-related macular degeneration (AMD), and diabetic retinopathy.
Additionally, many CNS diseases or disorders that are not primarily vascular – including Alzheimer
disease, multiple sclerosis, epilepsy, stroke, infection, and traumatic brain injury – are associated with
altered vascular function, most commonly with reduced BBB integrity (Profaci et al., 2020; Chen et al.,
2024).
87
A common feature of many CNS disorders is neuro-inflammation. Although vascular endothelial cells
(ECs) are not generally considered to be part of the immune system, in some locations and under some
conditions they acquire properties characteristic of immune cells, including secretion of cytokines,
surface display of co-stimulatory or co-inhibitory receptors, and antigen presentation in association with
MHC class II proteins (Pober and Sessa, 2014; Amersfoort et al., 2022). The best characterized role for
ECs in immune system function is as a site for binding and extravasation of circulating immune cells. As
first demonstrated in lymphoid organs, ECs can recruit immune cells by expressing chemokines and cell-
surface adhesion proteins that interact with cognate receptors on immune cells (Sackstein, 2005;
Blanchard and Girard, 2021). In the context of inflammation, increased production of chemokines and

adhesion proteins by ECs in non-lymphoid organs leads to the local recruitment of circulating immune
cells and their subsequent extravasation (Denes et al., 2024). Under non-inflammatory conditions, ECs
are maintained in a quiescent state. In CNS ECs, quiescence is maintained in part by the actions of
astrocyte-derived Sonic Hedgehog, with the result that few immune cells other than resident microglia
are found within the CNS (Alvarez et al., 2011).
102
TGF-beta signaling regulates inflammation in many tissues and in many cell types, as determined by the
phenotypes of ligand, receptor, and down-stream effector (SMAD) knockout (KO) mice (Shull et al.,
1992; Kulkarni et al. 1993; Travis and Shepard, 2014; Massagué and Sheppard, 2023). TGF-beta signaling
also controls vascular development, with loss-of-function mutations in TGF-beta receptor 1
(TGFBR1/ALK5), TGFBR2, the EC accessory receptor Endoglin, or various SMADs producing defects in
endothelial and/or pericyte development and early lethality (Goumans et al., 2009). The principal
ligands for the TGFBR1-TGFBR2 heterodimer are the three highly homologous TGF-beta family members
(Heldin and Moustakas, 2016). TGF-beta ligands are initially secreted as inactive ("latent") complexes
that include inhibitory subunits, and the dimeric ligands are subsequently released by the catalytic
action of integrins (Shi et al., 2011; Dong et al., 2017). Constitutive loss of both TGF-beta1 and TGF-
beta3, loss of integrins expressed by glia, or loss of TGFBR2 in endothelial cells produce similar defects in
CNS angiogenesis and CNS vascular integrity (Cambier et al., 2005; Mu et al, 2008; Aluwihare et al., 2009;
Nguyen et al. 2011; Allinson et al, 2012; Arnold et al., 2012, 2014).
116
The role of TGF-beta signaling in retinal vascular development has been studied with CreER/LoxP
conditional KO mice, both to facilitate survival beyond the early lethality of constitutive KO alleles and to
permit the study of cell-type specific KOs. Postnatal loss of TGFBR2 in all ocular cell types leads to
microaneurysms, leaky capillaries, retinal hemorrhages, reactive microglia, and pericyte abnormalities, a
picture that closely resembles the pathologies associated with severe diabetic retinopathy (Braunger et
al., 2015). Postnatal EC-specific loss of TGFBR2 leads to defective retinal angiogenesis, the absence of an
intraretinal capillary plexus, and choroidal neovascularization (CNV) (Allinson et al., 2011, Schlecht et al.,
2017; Zarkada et al., 2021). The present study was undertaken to more fully define the cellular and
molecular defects associated with endothelial-specific loss of TGF-beta signaling in the CNS, with an
emphasis on the retina. In particular, we have sought to define the relationship between endothelial
TGF-beta signaling and inflamation, as inflammation is a likely driver of diabetic retinopathy and AMD
(Wang et al., 2024).
129
130
131 **Results**
132
133 **Choroidal neovascularization with loss of endothelial TGF-beta signaling**
134
Retinal ECs were visualized with GS-lectin, anti-PECAM1, or anti-CLDN5 in retinas from young adult mice
with early postnatal EC-specific loss of *Tgfbr1* or *Tgfbr2* [*VEcadCreER;Tgfbr1$^{CKO/-}$* or
VEcadCreER;Tgfbr2$^{CKO/-}$ mice treated with 4- hydroxy-tamoxifen (4HT) between postnatal day (P)3 and
P5]. These retinas show moderate disorganization of the three nuclear layers and a near absence of
capillaries in the outermost tier of the retinal vasculature in the outer plexiform layer (OPL) (Figure 1;
Figure 1 – figure supplement 1), consistent with earlier descriptions of EC-specific inactivation of *Tgfbr1*
(Allinson et al., 2011, Schlecht et al., 2017; Zarkada et al., 2021). These retinas also show multiple
regions of choroidal (i.e., subretinal) neovascularization (CNV) as highlighted by the white arrows in
Figure 1A and B (middle row) [see also Figure 1C (right), and Figure 1 – figure supplement 1B].
Subretinal zones with normal morphology or with CNV are also seen in toluidine blue-stained epon-

embedded sections (Figure 1 – figure supplement 2). Quantification of whole eye cross-sections shows
an average of 2-3 zones of CNV per section in *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas, but no detectable CNV in
phenotypically WT control retinas (typically *VEcadCreER;Tgfbr1$^{CKO/+}$* or *VEcadCreER;Tgfbr2$^{CKO/+}$*) or in
retinas with severely reduced intraretinal vascular development secondary to loss of Norrin/Fzd4
signaling (*Ndp^{KO}*; Figure 1 – figure supplement 1C).

Throughout this study, our analyses of EC-specific KO of *Tgfbr1* (*VEcadCreER;TgfbrR1$^{CKO/-}$*), *Tgfbr2*
(*VEcadCreER;TgfbrR2$^{CKO/-}$*) and of both *Tgfbr1* and *Tgfbr2* (*VEcadCreER;TgfbrR1$^{CKO/-}$;Tgfbr2$^{CKO/-}$*) indicate
that the vascular phenotypes in the brain and retina are virtually identical across all three genotypes,
consistent with current models that envision TGFBR1-TGFBR2 heterodimers as the active receptor
complex (e.g., compare Figures 1A and 1B, and compare Figure 1 – figure supplements 1 and 3). In the
figures that follow, all analyses have been conducted with EC-specific KO of *Tgfbr1*. For completeness,
we also include some examples from EC-specific KO of *Tgfbr2* or EC-specific double KO of *Tgfbr1* and
Tgfbr2.

In wild type (WT) mice, all retinal ECs are PECAM1+ (Platelet and Endothelial Cell Adhesion Molecule-1)
and CLDN5+ (Claudin5, a marker of BBB and BRB vasculature) and all of the choroidal ECs are PLVAP+
(Plasmalemma Vesicle-Associated Protein, a marker of permeable vasculature) (Figure 1B and 1C; Figure
1 – figure supplement 3). In mice with EC-specific deletion of *Tgfbr1* or *Tgfbr2,* all of the choroidal ECs
are PLVAP+, all of the retinal ECs remain PLVAP-, and most, but not all, retinal ECs are CLDN5+ (Figure 1B
and 1C; Figure 1 – figure supplement 3). We note that this PLVAP phenotype contrasts with the
phenotype caused by mutations in the Norrin/Fzd4 pathway, which converts all retinal ECs to PLVAP+
and all capillary and vein ECs to CLDN5- (Wang et al., 2018).

Mutant retinas also exhibit occasional anastomoses between the subretinal and retinal vasculatures, a
feature never observed in control retinas (Figure 1B, white arrows in bottom panels). The intraretinal
segments of the communicating vessels are CLDN5+/PLVAP- and the subretinal segments, which
invariably arise in a zone of CNV, are CLDN5-/PLVAP+ (Figure 1B), implying a mixed origin for these
vessels. The attenuated intraretinal vasculature produced by deficient TGF-beta signaling is associated
with retinal hypoxia as determined by localized accumulation of Hypoxia Inducible Factor (HIF)1-alpha in
the nuclei of retinal parenchymal cells (Figure 1 – figure supplement 4).

In mutant retinas, CD45+ immune cells are abundant and many of these cells are closely associated with
the subretinal and inner retinal vasculatures (Figure 1A and 1C; Figure 1 – figure supplement 1A and 1B).
In control retinas, the only CD45+ cells are the relatively sparse microglia. In the several non-CNS tissues
examined – heart, kidney, liver, and lung – the density of CD45+ cells appears to be unaffected in
VEcadCreER;Tgfbr1$^{CKO/-}$ mice (Figure 1 – figure supplement 5). Immunostaining for RPE65, a marker for
the retinal pigment epithelium (RPE), and toluidine blue-staining of epon-embedded retina sections both
show RPE displacement at the sites of CNV (Figure 1C; Figure 1 – figure supplements 1B and 2). In
mutant retinas, there is little change in pericyte NG2 immunostaining (Figure 1 – figure supplement 3).

As the phenotypes associated with EC-specific deletion of *Tgfbr1* or *Tgfbr2* involve both endothelial cells
and immune cells, it is important to ask whether Cre-mediated recombination directed by *VEcadCreER*
is, in fact, EC-specific. With two *LoxP-stop-LoxP* (*LSL*) reporters that express nuclear-localized GFP, *R26-*
LSL-mtdT-2A-nlsGFP (Wang et al., 2018) and *R26-LSL-SUN1-sfGFP-6xmyc* (Mo et al., 2015), *VEcadCreER*
directs recombination exclusively in ECs in the retina, cerebellum, heart, kidney, and lung, as judged by
cellular morphology, EC identification with PECAM1 immunostaining, and co-localization of GFP with the
EC-specific transcription factor ERG (Figure 1 – figure supplement 6). Additionally, CD45 localization in

immune cells is mutually exclusive with *R26-LSL-SUN1-sfGFP-6xmyc* reporter expression in the choroid,
small intestine, and retina (figure 1 – figure supplement 7). In these experiments, the occasional
exceptions to ERG and GFP co-localization consist of ECs in which Cre-mediated recombination failed to
occur (e.g., Figure 1 – figure supplement 6B, white arrows in the kidney and lung panels). As a further
test of *VEcadCreER* specificity, we asked whether the EC-specific recombination of the *R26-LSL-SUN1-*
sfGFP-6xmyc reporter observed on a WT *Tgfbr1* background was retained in a *Tgfbr1$^{CKO/-}$* background.
As seen in Figure 1 – figure supplement 8, the CD45+ immune cells that accumulate in the retina lacking
endothelial *Tgfbr1* do not express the GFP reporter, which retains its pattern of EC specificity. Based on
these experiments, we conclude that all of the phenotypes observed in *VEcadCreER;Tgfbr1$^{CKO/-}$* and
VEcadCreER;Tgfbr2$^{CKO/-}$ mice follow from the loss of TGFBR function exclusively within ECs, and, more
specifically, that all immune cell phenotypes in the retina are secondary to a change in EC properties.
204
VEcadCreER;Tgfbr1$^{CKO/-}$ retina flatmounts show large numbers of vascular tufts (Figure 2A). This
aberrant vascular architecture, which is seen in both *VEcadCreER;Tgfbr1$^{CKO/-}$* and *VEcadCreER;Tgfbr2$^{CKO/-}$*
mice, shows some similarities to the previously reported vascular architecture of *Fzd4$^{-/-}$* and *Ndp^{KO}* mice
(Luhmann et al., 2005; Ye at al., 2009). In particular, retinas with each of these genotypes variably
display disorganized and hypertrophic superficial vessels that give rise to intraretinal vascular tufts
instead of the well-organized trilayered retinal vasculature. The morphologies and locations of the
intraretinal vascular tufts are similar among all of these mutant genotypes, as shown in Figure 2B. At
P14, when endothelial tip cells are building the central tier of intra-retinal capillaries in wild type
(control) retinas (white arrows in the upper left panel in Figure 2B), ECs in *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas
have formed only a few rudimentary capillaries or disconnected vascular tufts (lower left panels in
Figure 2B). In addition to their distinctive morphology, the vascular tufts express low levels of PLVAP
and they show an accumulation of the low molecular weight intravascular tracer sulfo-NHS-biotin, an
indication of increased vascular permeability (Figure 2C).
218
219
220   **Immune cell phenotype in the retina with loss of endothelial TGF-beta signaling**
221
To more rigorously characterize the immune response in *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas and to compare
this response to that observed with loss of Norrin/Fzd4 signaling (*Ndp^{KO}* retinas and *Fzd4$^{-/-}$* retinas), flat
mount retinas from P15-P30 mice were immunostained for CD45/PTPRC (multiple immune cell types),
F4-80 (monocytes and macrophages), PU.1/SPI1 (myeloid cells), IBA1/AIF-1 (microglia and
macrophages), and CD3E (T-cells) (Figure 3A-C). Each of these markers shows a marked increase in
immune cells in *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas compared to age-matched controls. Quantifying the
density of cells staining for CD45, PU.1, F4-80, and CD3E shows that, for each of these markers,
endothelial loss of TGF-beta signaling is associated with the highest immune cell increase and loss of
Norrin/Fzd4 signaling is associated with a more modest immune cell increase (Figure 3D-G).
231
In the quantification of CD45+ cells in Figure 3, cells with resident microglial morphology were not
counted. As seen by immunostaining for ASC (Apoptosis-associated speck-like protein containing a
CARD; nuclei) and CD45 (plasma membrane), microglia are present in control retina flatmounts at a
density of 25-30 cells per 450 um x 450 um area in each of the three retinal layers in which they reside
[retinal ganglion cell layer (RGC), inner plexiform layer (IPL), and outer plexiform layer (OPL)], for a total
density of ~85 cells per 450 um x 450 um area (Figure 3 – figure supplement 1). The density of CD45+
cells in *Fzd4$^{-/-}$* and *Ndp^{KO}* retinas in excess of the density of CD45+ cells in control retinas is ~100 per 450
um x 450 um area, and the excess density of CD45+ cells in *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas is ~200 per
450 um x 450 um area (Figure 3D).

241
In flatmounts of control choroids, CD45+ cells are present at a density of 100-150 cells per 645 um x 645
um area, and in flatmounts of *VEcadCreER;Tgfbr1$^{CKO/-}$* choroid, CD45+ cells are present at approximately
twice that density, with substantial scatter in the *VEcadCreER;Tgfbr1$^{CKO/-}$* data (Figure 3 – figure
supplement 2). A curious feature of CD45+ cells in the *VEcadCreER;Tgfbr1$^{CKO/-}$* retina is the large number
of cells that are positive for cleaved Caspase 3, a marker of apoptosis (Figure 3 – figure supplement 3).
Very few retinal cells of any other type are positive for cleaved Caspase 3. In control retinas, and in *Fzd4^{-}*
$^{/-}$ and *Ndp^{KO}* retinas, very few cells are positive for cleaved Caspase 3, despite the presence of excess
CD45+ cells in *Fzd4$^{-/-}$* and *Ndp^{KO}* retinas (Figure 3 – figure supplement 3A; quantified for *Fzd4$^{-/-}$* in Figure 3
– figure supplement 3B). These data imply that in the *VEcadCreER;Tgfbr1$^{CKO/-}$* retina there is rapid
turnover of CD45+ cells, with new cells replenishing the population as resident cells are eliminated.
252
253
254 **Immune cell phenotyping in the retina by single nucleus RNAseq**
255
To obtain an unbiased assessment of the immune cell types present in the *VEcadCreER;Tgfbr1$^{CKO/-}$* retina,
P14 control and *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas were compared by single nucleus (sn)RNAseq (Figure 4).
For both genotypes, immune cells represent only a minor fraction of retinal cells (Figure 4A). While
microglia are present in both control and *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas, all of the other immune cells in
the combined snRNAseq data set were derived from *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas (Figure 4B). As
determined by the patterns of expression of known immune cell markers (BD Biosciences, 2024), the
immune cell population in *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas encompasses the full spectrum of major cell
classes: B-cells, T-cells, dendritic cells, macrophages, natural killer (NK)-cells, and microglia (Figure 4C-E).
In contrast to the immune cells in the *Aire$^{-/-}$* mouse model of autoimmune uveoretinitis (Heng et al.,
2019), which self-organize into tertiary lymphoid organs, the immune cells in *VEcadCreER;Tgfbr1$^{CKO/-}$*
retinas show little evidence of spatial organization beyond an asssociation with the vasculature (Figure
3A-C, and text below).
268
269
270 **Hypovascularization secondary to loss of retinal VEGF signaling does not lead to immune cell**
271 **infiltration**
272
The presence of immune cells in retinas lacking endothelial TGF-beta signaling or Norrin/Fzd4 signaling
could potentially arise from immune cell infiltration secondary to the hypoxic stress within the inner
retina that results from reduced vasculaturization (Figure 1 – figure supplement 4). Alternately, it could
reflect changes in the intrinsic properties of retinal ECs that promote immune cell recruitment and
extravasation. As a first step in distinguishing between these models, we generated and studied a
distinct retinal hypovascularization model – *Chx10-Cre;Vegfa$^{CKO/CKO}$* – in which deletion of VEGF-A
(hereafter, VEGF) in retinal Muller glia and neurons (Rowan and Cepko, 2004) greatly reduces intra-
retinal vascularization.
281
VEGF production by surface astrocytes guides the first stage of retinal angiogenesis, during which
endothelial cells grow outward from the optic disc along the vitreal face of the retina (Stone et al. 1995;
Rattner et al., 2019). The second stage of retinal angiogenesis involves EC growth into the retina and is
driven by VEGF production from cells within the inner retina (Rattner et al., 2019). In *Chx10-*
Cre;Vegfa$^{CKO/CKO}$ mice, the first stage of retinal angiogenesis is largely unaffected, but the second stage
fails to occur (Figure 5A-C). In retinas lacking endothelial TGF-beta signaling or Norrin/Fzd4 signaling,
high levels of retinal VEGF drive excess proliferation of ECs on the vitreal face of the retina and in intra-

retinal tufts. In contrast, the lack of retina-derived VEGF in *Chx10-Cre;Vegfa*$^{CKO/CKO}$ mice results in little or
no additional proliferation of ECs (Figure 5A-C).
291
To compare immune cell infiltration between control and *Chx10-Cre;Vegfa*$^{CKO/CKO}$ retinas, flatmount
retinas were immunostained for ASC and CD45 (Figure 5D). Both sets of retinas exhibited similar
numbers of immune cells, almost all of which appear to be microglia as judged by their morphology and
distribution (Figure 5E). These data imply that retinal hypo-vascularization and inner retinal hypoxia *per*
se does not lead to immune cell infiltration into the retina. Instead, the simplest explanation for this
data is that the absence of endothelial TGF-beta signaling – and, to a lesser extent, the absence of
Norrin/Fzd4 signaling, but not the absence of VEGF signaling – leads to a pro-inflammatory vascular state
that attracts immune cells. This second hypothesis is explored in the text and figures below.
300
301
302 **Close physical association between retinal vasculature and immune cells with loss of endothelial TGF-**
303 **beta signaling**
304
A close association between blood vessels and CD45+ immune cells was observed in the retina cross-
sections in Figure 1. To visualize these associations in intact retinas, flatmounts were immunstained for
PECAM1, ASC, and CD45 (Figure 6). In control retinas, resident microglia are the only CD45+/ASC+ cells
in the retina and their cell bodies and processes tile the retina in a manner that appears to be
independent of the locations of blood vessels (Figure 6, Figure 3 – figure supplement 1, and Figure 6 –
figure supplements 1-3). By contrast, in retinas with loss of EC TGF-beta signaling, many immune cells
are closely associated with the vasculature (Figure 6 and Figure 6 – figure supplements 1-3). Aberrant
vessels that penetrate the full thickness of the outer retina are associated along their length with CD45+
cells, as seen in serial Z-plane images at both low magnification (Figure 6 – figure supplements 1 and 2)
and high magnification (Figure 6 – figure supplement 3).
315
The subcellular distribution of PECAM1 on retinal ECs is non-uniform in both mutant and control retinas
(Figure 6D).   Close inspection of *VEcadCreER;Tgfbr1*$^{CKO/-}$ retinas shows that in venous ECs there are
numerous ~5 um diameter zones of reduced PECAM1 immunostaining, often accompanied by a rim of
increased staining (white arrows in Figure 6D). These zones coincide with the locations of CD45+
immune cells, implying a direct physical association between immune cells and ECs (Figure 6D). This
association is only rarely observed in control retinas, where immune cells other than microglia are
sparse (Figure 6D). In *VEcadCreER;Tgfbr1*$^{CKO/-}$ retinas, the association of leukocytes with veins is
reminiscent of leukocyte adhesion to high endothelial venules in lymphoid organs (Blanchard and Girard,
2021). These data imply that ECs lacking TGF-beta signaling bind immune cells, likely promoting their
recruitment and extravasation.
326
327
328 **Increase in ICAM1 in retinal endothelial cells and altered pericyte properties with loss of endothelial**
329 **TGF-beta signaling**
330
As noted in the Introduction, under inflammatory conditions, ECs secrete cytokines, express adhesion
and co-stimulatory proteins to recruit and activate immune cells, and display immunogenic peptides
bound to MHC proteins (Pober and Sessions, 2014; Amersfoort et al., 2022). Intercellular adhesion
molecule 1 (ICAM1) is a general marker for vascular inflammation. Endothelial ICAM1 acts as a receptor
for Lymphocyte Function-Associated Antigen 1 (LFA1), an integrin broadly expressed on leukocytes, and
ICAM1-LFA1 binding promotes transmigration of leukocytes from blood to tissue (Ding et al., 1999).

Immunostaining for ICAM1 in *VEcadCreER;Tgfbr1$^{CKO/-}$*, *Fzd4$^{-/-}$*, and *Ndp^{KO}* retinas – each paired with
littermate control retinas processed in parallel – revealed a ~2-fold increase in endothelial ICAM1 in
Fzd4$^{-/-}$ and *Ndp^{KO}* retinas and a ~7-fold increase in *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas (Figure 7A and B, and
Figure 7 – figure supplement 1). Control retina flatmounts show weak ICAM1 immunstaining in veins
and undetectable ICAM1 in arteries and capillaries, whereas *VEcadCreER;Tgfbr1$^{CKO/-}$* retina flatmounts
show accumulation of ICAM1 in veins and in many small diameter vessels. The elevations in EC ICAM1 in
these models of retinal hypovascularization correlate with the levels of retinal immune cells (Figure 3),
suggesting that the level of vascular inflammation, and, more specifically, the level of ICAM1, is one
determinant of the rate of egress of immune cells from blood to retina.
346
Changes in EC phenotype can lead to changes in the characterstics of pericytes, as seen, for example,
with defects in Platelet-Derived Growth Factor signaling (Lindahl and Betsholz, 1998). Although pericyte
coverage appears to be minimally affected by loss of TGF-beta signaling (Figure 1 – figure supplement 3),
we observed a change in the relationship between NG2, a pericyte marker, and smooth muscle actin
(SMA), a smooth muscle marker, in flatmounts of *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas compared to control
retinas. In control retinas, SMA immunostaining is intense on arteries, patchy on veins, and nearly
undetectable on capillaries, with the latter staining strongly for NG2 (Figure 7C). In
VEcadCreER;Tgfbr1$^{CKO/-}$ retinas, artery and vein SMA immunostaining is minimally affected, but
capillaries and vascular tuft ECs immunostain for both NG2 and SMA (Figure 7C and D). These
differences could reflect higher levels of SMA in *VEcadCreER;Tgfbr1$^{CKO/-}$* pericytes or a change in actin
polymerization state. The latter possibility is based on the observations of Alarcon-Martinez et al (2018)
that immunostaining for SMA in mouse retinal pericytes is strongly enhanced by actin polymerization.
359
360
361   **Transient defects in the BBB, altered pericyte properties, and localized immune infiltrates in the brain**
362   **with loss of endothelial TGF-beta signaling**
363
To determine whether the phenotypes associated with EC loss of TGF-beta signaling extend to CNS
territories beyond the retina, we surveyed the brain in *VEcadCreER;Tgfbr1$^{CKO/-}$* mice for
leakage/transport of IgG into the parenchyma, pericyte changes (SMA immunostaining), vascular
inflammation (ICAM1 immunostaining), and immune cell infiltration (Figure 8 and Figure 8 – figure
supplement 1). Interestingly, IgG accumulates in the *VEcadCreER;Tgfbr1$^{CKO/-}$* brain parenchyma at P14
before resolving over the next week, imply a transient pro-inflammatory state within the brain (Figure
8A). As in the retina, in *VEcadCreER;Tgfbr1$^{CKO/-}$* brains there was increased SMA immunoreactivity in
capillary-associated pericytes (Figure 8B). In *VEcadCreER;Tgfbr1$^{CKO/-}$* brains, there was scattered
accumulation of immune cells and variably increased EC ICAM1 levels (Figure 8 – figure supplement 1).
373
After P30, mice with EC-specific loss of TGFBR1, TGFBR2, or both TGFBR1 and TGFBR2 exhibit an increase
in the density of focal brain lesions that are characterized by bleeding and an accumulation of immune
cells (Figure 8C). These brain lesions, which appear to be small cerebrovascular events (strokes), may
explain the observation that otherwise healthy adult mice with EC-specific loss of TGF-beta signaling die
at a rate of ~20% per month. Taken together, the histologic and mortality patterns suggest that mice
with defects in TGF-beta signaling are prone to lethal cerebrovascular events.
380
381
382   **Altered brain endothelial gene expression with loss of endothelial TGF-beta signaling**
383

To obtain an unbiased assessment of gene expression changes in brain ECs lacking TGF-beta signaling,
vascular fragments from P14 control and *VEcadCreER;Tgfbr1$^{CKO/-}$* brains were enriched by centrifugation
through Ficoll and used as input for single nucleus (sn)RNAseq (Figure 9). The paucity of ECs in CNS
tissue (several percent of cells) and the difficulty in freeing ECs from the extra-cellular matrix limit the
yield of CNS ECs when using enzymatic dissociation methods and FACS sorting. By contrast, the Ficoll
centrifugation protocol can rapidly enrich a large number of vascular fragments because it starts with a
large mass of brain tissue (one or more mouse brains). We chose snRNAseq rather than single cell
(sc)RNAseq because snRNAseq has two advantages: (a) input vascular fragments can be frozen prior to
nuclear isolation, and (2) the tissue is not subject to enzymatic dissociation, during which time changes
in gene expression can occur (Lacar et al., 2016; Hrvatin et al., 2018).

snRNAseq on vascular fragments provides a snapshot of the transcriptomes of all of the major CNS cell
classes, with ECs nuclei enriched to ~15% of the total (Figure 9A). It is evident from visual inspection of
the shapes and positions of cell clusters in the UMAP plots that gene expression in ECs undergoes a
substantial change with EC-specific loss of TGF-beta signaling, whereas gene expression in other cell
types undergoes far smaller changes. A volcano plot of transcriptome changes within the EC cluster
shows several dozen transcripts with fold changes >4 and -log$_{10}$ p-values greater than 50 (chosen as a
conservative p-value cut-off for this comparison of 2475 control vs 2944 mutant nuclei) (Figure 9B).
Figure 9C shows UMAP plots for individual transcripts that exhibit increased abundance in mutant ECs
(*Ahr*, *Nr5a2*, *Pcdh17*, and *Tgfbr3*), decreased abundance in mutant ECs (*Stra6*, *AU021092*, and *Htra3*), or
little or no change in mutant ECs (*Icam1* and *Icam2*). In *VEcadCreER;Tgfbr1$^{CKO/-}$* ECs, the elevation in
Tgfbr3 transcripts, which code for Beta-glycan, an accessory (type III) TGFBR receptor, could represent a
homeostatic response to reduced TGF-beta signaling since Beta-glycan enhances the binding of TGF-beta
ligands to the TGFBR1-TGFBR2 complex (Heldin and Moustakas, 2016). The lack of a significant change
in *Icam1* transcript levels implies that the increase in ICAM1 immunostaining is a post-translational
phenomenon, either an increase in protein level or an increase in protein accessibility.

Gene Set Enrichment Analysis (GSEA) confirms that the greatest changes in gene expression among CNS
cell types in *VEcadCreER;Tgfbr1$^{CKO/-}$* mice are in ECs, and it shows that EC transcriptome changes
connected to the cell cycle and inflammation are the dominant categories (Figure 9D). The latter
includes the categories "Interferon (IFN) gamma response", "Interferon (IFN) alpha response",
"IL2/STAT5 signaling", and "Inflammation". An analysis of two categories of inflammation-associated
proteins – components of the NF-kappa-B (NFkB) pathway and the integrins – shows (1) constitutive
enrichment in CNS ECs of transcripts coding for the p50 subunit of NFkB (*Nfkb1*) and, most dramatically,
NFkB inhibitor alpha (*Nfkbia*), and (2) with loss of TGF-beta signaling, upregulation of transcripts coding
for multiple integrins, including alpha1, alpha2, alpha4, alpha6, and beta1 (Figure 9 – figure supplement
1A and 1B). Figure 9 – figure supplement 1D presents a pictorial summary of the trancripts coding for
those chemokines, adhesion proteins, integrins, and TNF receptor superfamily members that are up-
regulated in CNS ECs in *VEcadCreER;Tgfbr1$^{CKO/-}$* mice.

Immunostaining of whole mount retinas for the p65 subunit of NFkB shows enrichment in both control
and *VEcadCreER;Tgfbr1$^{CKO/-}$* vasculature relative to non-vascular cells (Figure 9 – figure supplement 1C).
Immunostaining of whole mount retinas for integrin alpha2 (ITGA2) shows undetectable staining in ECs
in control vasculature and clear staining in *VEcadCreER;Tgfbr1$^{CKO/-}$* vasculature.   The change in ITGA2
immunostaining is in the same direction as the upregulation of *Itga2* transcripts in
VEcadCreER;Tgfbr1$^{CKO/-}$ ECs (Figure 9 – figure supplement 1B).

Immunostaining of whole mount retinas for integrin alpha4 (ITGA4) shows enrichment in both control
and *VEcadCreER;Tgfbr1$^{CKO/-}$* vasculature relative to non-vascular cells (Figure 9 – figure supplement 1C).
Integrin alpha4/beta1 (Very Late Antigen-4; VLA-4) is a heterodimer adhesion protein expressed on ECs
and immune cells that is implicated in immune cell homing. It is also one of several integrins expressed
by ECs that mediate cell-cell and cell-matrix interactions (Guerrero and McCarty, 2018; Aman and
Margadant, 2023). In *VEcadCreER;Tgfbr1$^{CKO/-}$* ECs, there is little change in vascular ITGA4
immunostaining intensity, but an increase in *Itga4* transcript levels (Figure 9 – figure supplement 1B).
These data suggest that there is an increase in both synthesis and degradation of ITGA4 in
VEcadCreER;Tgfbr1$^{CKO/-}$ ECs.
440
The large number of transcriptional changes observed in control vs. *VEcadCreER;Tgfbr1$^{CKO/-}$* CNS ECs
imply substantial changes at the level of transcription and, by implication, the expression of transcription
factors (TFs). Four TFs with the most dramatic increases in *VEcadCreER;Tgfbr1$^{CKO/-}$* ECs are shown in the
UMAP plots in Figure 9C and Figure 9 – figure supplement 2A: the Aryl Hydrocarbon Receptor (*Ahr*), a
basic helix-loop-helix TF that reduces inflammatory responses in a variety of tissues and cell types
(Stockinger et al., 2024); Nuclear Receptor Subfamily 5 Group A Member 2 (*Nr5a2*), also referred to as
Liver Receptor Homologue-1, a TF with multiple roles in development and metabolism (Fayard et al.,
2004); and two members of the Thymocyte Selection-Associated High Mobility Group Box gene family
(*Tox* and *Tox3*), a TF family that is required for the development of innate immune cells and T-cells
(Aliahmad et al., 2012). Immunostaining of retina flatmounts shows a several-fold increase in TOX levels
in *VEcadCreER;Tgfbr1$^{CKO/-}$* ECs compared to control ECs, with localization of the signal to the nucleus
(Figure 9 – figure supplement 2B).
453
454
455 **Discussion**
456
The experiments described here show that postnatal EC-specific loss of TGF-beta signaling in mice leads
to aberrant angiogenesis in the retina and a pro-inflammatory state within the retina and brain. More
specifically, EC-specific loss of TGF-beta signaling leads to: (1) reduced intra-retinal vascularization, (2)
choroidal neovascularization with occasional anastomoses connecting choroidal and intraretinal
vasculatures, (3) infiltration of diverse immune cells into the retina, including macrophages, T-cells, B-
cells, NK cells, and dendritic cells, (4) a close physical association between immune cells and vasculature,
(5) a pro-inflammatory transcriptional state in CNS ECs, with increased ICAM1 immunoreactivity, and (6)
increased SMA immunostaining in pericytes. A striking feature of the phenotypes studied here is their
CNS specificity. Despite Cre-mediated recombination in ECs throughout the body directed by
VEcadCreER, the vascular and inflammatory phenotypes associated with EC-specific loss of TGF-beta
signaling appear to be largely confined to the CNS.
468
Comparisons of the retinal phenotype with two other genetic models of retinal hypovascularization –
loss of Norrin/Fzd4 signaling and loss of VEGF signaling – reveal interesting differences. While loss of
either Norrin/Fzd4 or VEGF signaling leads to a near absence of intra-retinal capillaries – similar to loss of
TGF-beta signaling – the models differ in their immune phenotypes. The immune cell infiltrate is
greatest with loss of TGF-beta signaling, more modest with loss of Norrin/Fzd4 signaling, and
undetectable with loss of VEGF signaling. Interestingly, cleaved Caspase 3 is abundant in immune cells
with loss of TGF-beta signaling but it is extremely rare with loss of Norrin/Fzd4 signaling.
476
477
478 **A model for choroidal neovascularization**

479
Choroidal neovascularization is the hallmark feature of neovascular AMD. The original classification
scheme for neovascularization in AMD posited two categories: occult, or type 1, neovessels were those
located in the sub-RPE space, and classical, or type 2, vessels were those located in the subretinal space
(i.e., between the RPE and retina) (Gass, 1997). More recently, a third type of neovessel has been
recognized that consists of an anastomosis between choroidal and retinal vasculatures (Freund et al.,
2008), and these type 3 vessels are observed in approximately one-third of eyes with newly diagnosed
neaovascular AMD (Jung et al., 2014). Among patients with type 3 vessels in only one eye, the
probability that the other eye will develop type 3 vessels within three years is close to 100% (Gross et al.,
2005).
489
Multiple animal models of ocular neovascularization have been described (Grossniklaus et al., 2010;
Qiang et al., 2021). In the simplest model of CNV, a focal laser-induced injury to the RPE and Bruch's
membrane (the extracellular matrix that separates the RPE and the choroidal vessels) leads to growth of
choroidal vessels through the lesion (Fabian-Jessing et al. 2022). This response reveals the intrinsic
angiogenic potential of the choroidal vasculature, which is normally held in check by Bruch's membrane
and an intact RPE monolayer. Models of CNV that likely have greater relevance to pathogenic
mechanisms in AMD have been developed in mice by altering – either singly or in combination –
lipid/cholesterol metabolism, inflammation, and oxidative damage. These models include (1) KO of
Cyp27a1, a ubiquitously expressed cytochrome P450 (Omarova et al., 2012), (2) *Ccr2*/*Ccl2* double KO
(Takeda et al., 2009), (3) ApoE over-expression combined with a high-fat diet (Malek et al., 2005), (4)
Ccl2/*Cx3cr1* double KO combined with a low omega-3 polyunstaured fatty acid diet (Chan et al., 2008),
and (5) KO of Superoxide dismutase-1 (*Sod1*) and aging for at least one year (Imamura et al., 2006). We
note that CNV was observed in the Ndp^{KO} mice studied by Beck and colleagues (2018), although we did
not observe CNV in our Ndp^{KO} mice.
504
Genetic models for pathologic angiogenesis that may be relevant to type 3 neovascularization include (1)
over-expression of VEGF in rod photoreceptors (Tobe et al., 1998; Ohno-Matsui et al., 2002), (2) RPE-
specific KO of the Von Hippel-Lindau gene (*Vhl*), which leads to activation of the hypoxia response and
excessive production of VEGF (Lange et al., 2012), (3) rod photoreceptor-specific KO of *Vegfr1*, which
codes for FLT-1, a decoy receptor that reduces VEGF signaling (Luo et al., 2013), and (4) KO of the Very
Low-Density Lipoprotein Receptor (*Vldlr*), which presumably perturbs lipid metabolism (Heckenliveley et
al., 2003; Hu et al., 2008).
512
513 Loss of TGF-beta signaling in ECs is mechansitically distinct from all of the above-mentioned models, and
514 it has the added practical advantage that multiple CNV tufts are present in virtually every eye by three
515 months of age without the need for dietary of other interventions (Figure 1 – figure supplement 1).
516
517
518 **Inflammation in retinal and brain vascular disease**
519
Local and/or systemic inflammatory markers, such as cytokines, are elevated in patients with AMD,
diabetic retinopathy, and retinal vein occlusion (Tang et al., 2011; Jung et al., 2014; Kauppinen et al,
2016). One characteristic of the retinal inflammatory state is increased leukocyte adhesion to retinal
ECs, which is observed in patients with diabetic retinopathy and may be a causal factor in vaso-occlusive
events (Chibber et al., 2007). In rats with streptozotocin-induced diabetes, ICAM1 and CD18 are
elevated in retinal ECs, and genetic ablation of *Icam1* or *Cd18* or treatment with an ICAM1-blocking mAb
reduces retinal leukostasis and vascular leakage (Miyamoto et al., 1999; Joussen et al., 2004).

Neuroinflammation in general, and vascular neuroinflammation in particular, is also a feature of multiple
non-retinal CNS diseases, including Alzheimer disease, stroke, and multiple sclerosis (Ritson et al., 2024).
529
Current evidence points to a combination of RPE oxidative damage, sub-RPE inflammatory cells, and
complement activation as pathogenic mechanisms in neovascular AMD. In the early stages of AMD,
inflammatory cells accumulate in the choroid and complement proteins accumulate in subretinal
deposits, and, in the more advanced neovascular stage of AMD, choroidal neovascular tufts are
associated with inflammatory cells (Kauppinen et al, 2016; Armento et al., 2021; Heloterä and
Kaarniranta, 2022). Genetic evidence for a causal role of the complement system comes from the
elevated AMD risk associated with the Y402H and I162V variants in the complement factor H gene, and
less commonly with variants in the genes coding for complement factor 3 (C3), complement factor I
(CFI), and the complement regulator SERPING1 (Montezuma et al., 2007).
539
540 In sum, the evidence presented here and in Schlecht et al. (2017) shows that loss of TGF-beta signaling in
541 CNS ECs recapitulates some of the cardinal features of retinal and neurologic diseases associated with
542 vascular inflammation. These experiments suggest that enhancing TGF-beta-dependent anti-
543 inflammatory responses in ECs could represent a promising strategy for disease modulation (Muniyandi
544 et al., 2023; Hu et al., 2018; Hachana and Larrivée, 2022; Ravichandran and Heneka, 2024).
545
546
547
548 **Materials and methods**
549
Mice
The following mouse lines were used: *VEcadCreER* (Monvoisin et al., 2006); $Tgfbr1^{CKO}$ (Larsson et al.,
2001; JAX 028701); $Tgfbr2^{CKO}$ (Leveen et al., 2002; JAX 012603); $Fzd4^{KO}$ (Wang et al., 2001; JAX 012823);
Ndp^{KO} (Ye et al., 2009; JAX 012287); $VEGF^{CKO}$ (Gerber et al., 1999); *Chx10-Cre* (Rowan and Cepko, 2004);
R26-LSL-SUN1-sfGFP (Mo et al., 2015; JAX 030952), and *R26-LSL-tdTomato-2A-H2B-GFP* (Wang et al.,
2018; JAX 030867). *Ndp* is located on the X-chromosome and therefore we refer to both female $Ndp^{-/-}$
and male $Ndp^{-/Y}$ mice as Ndp^{KO}. All mice were housed and handled according to the approved
Institutional Animal Care and Use Committee protocol of the Johns Hopkins Medical Institutions
(protocol MO19M429).
559
560 **4-hydroxytamoxifen preparation and administration**
561 4HT (Sigma-Aldrich H7904-25MG) was dissolved at 20 mg/ml in ethanol by extensive vortexing.
562 Sunflower seed oil (Sigma-Aldrich S5007) was added to dilute the 4HT to 2 mg/ml, and aliquots were
563 stored at -80^{o}C. Thawed aliquots were mixed well before injections. All injections were performed
564 intraperitoneally.
565
Antibodies and other reagents
The following antibodies were used for tissue immunohistochemistry: rat anti-mouse PLVAP/MECA-32
(BD Biosciences 553849); rat anti-mouse CD31 (BD Biosciences 553370); rat anti-mouse ICAM-1
(Invitrogen 14-0542-82); rat anti-mouse F4/80 (BIO RAD MCA497G); rat anti-mouse CD206 (BioRad
MCA2235); rat anti-mouse PU.1/Spi-1 (R&D Systems MAB7124); mouse mAb anti-alpha SMA, Cy3
conjugate (Sigma-Aldrich C6198); mouse mAb anti-CLDN5, Alexa Fluor 488 conjugate (Thermo Fisher
Scientific 352588); mouse mAb anti-RPE65, Dylight 550 conjugate (Invitrogen MA5-16044); rabbit anti-
Collagen IV (Novus Biologicals NB120-6586); rabbit anti-NG2 Chondroitin Sulfate Proteoglycan (Millipore
AB5320); rabbit mAb anti-ASC/TMS1 (Cell Signaling 67824S); rabbit mAb anti-cleaved Caspase-3 (Cell

Signaling 9664S); rabbit mAb anti-HIF-1alpha (Cell Signaling 36169S); rabbit mAb anti-P-SMAD1/5/9 (Cell
Signaling 13820S); Armenian hamster anti-CD3e (Invitrogen 14-0031-82); goat anti-mouse CD45 (R&D
Systems AF114); goat anti-Iba1 (Novus Biologicals NB100-1028); chicken anti-GFP (Aves Labs GFP-1020);
rabbit mAb anti-NFkappaB NF-κB p65 (D14E12; Cell Signaling Technology 8242S); rabbit mAb anti-
Integrin alpha 2 (ITGA2; clone GEB, BosterBio M01933); rabbit mAb anti-Integrin alpha 4 (ITGA4; D2E1;
Cell Signaling Technology 8440); rabbit mAb anti-TOX/TOX2 (E6G5O; Cell Signaling Technology 36778S).
Alexa Fluor-labeled secondary antibodies and GS Lectin (Isolectin GS-IB4) were from Thermo Fisher
Scientific. Alexa Fluor-labeled secondary goat anti-Armenian hamster IgG antibodies were from
BioLegend. Texas Red Streptavidin was from Vector Laboratories (SA-5006). Sulfo-NHS-biotin was from
Thermo Fisher Scientific (21217).
585
Tissue processing and immunohistochemistry
Tissues were prepared and processed for immunohistochemical analysis as described by Wang et al.
(2012) and Zhou et al. (2014). In brief, mice were deeply anesthetized with ketamine and xylazine and
then perfused via the cardiac route with 1% paraformaldehyde (PFA) in phosphate buffered saline (PBS).
Non-ocular tissues were dissected and dehydrated in 100% cold methanol overnight at 4^{o}C. Tissues
were re-hydrated the following day in 1x PBS at 4^{o}C for at least 3 hours before embedding in 3%
agarose. Tissue sections of 100-200 μm thickness were cut using a Leica vibratome.
593
For whole-mount retinas, intact eyes were immersion fixed in 1% PFA in PBS at room temperature for 1
hour before the retinas were dissected. For eye sections, enucleated eyes were imbedded in Optimal
Cutting Temperature (OCT) compound (Tissue-Tek 4853) and frozen in dry ice. Embedded eyes were cut
into 14 μm sections with a Zeiss cryostat and stored on glass slides at -80^{o}C. For immunostaining,
sections were warmed to room temperature, fixed in 1% PFA at room temperature for 30 minutes, and
washed in PBS before pre-blocking.
600
601 For vascular permeability analysis, mice were injected intraperitoneally with Sulfo-NHS-biotin (100-200
602 μl of 20 mg/ml Sulfo-NHS-biotin in PBS) 30-60 minutes prior to intracardiac perfusion. Covalently bound
603 biotin was visualized in tissue sections or in whole-mount retinas with Texas Red conjugated
604 Streptavidin.
605
Tissue sections or whole-mount retinas were permeabilized in PBSTC (1x PBS + 1% Triton X-100 + 0.1mM
$CaCl_2$) overnight at 4^{o}C, and subsequently incubated overnight at 4^{o}C with primary antibodies, diluted in
1x PBSTC + 7% normal goat or donkey serum. Samples were washed at least 6 times with 1x PBSTC over
the course of 6-8 hours and subsequently incubated overnight at 4^{o}C with secondary antibodies diluted
in 1x PBSTC + 7% normal goat or donkey serum. The next day, samples were washed at least 6 times
with 1x PBSTC over the course of 6 hours and mounted in Fluoromount G (SouthernBiotech 0100-01).
612
613 **Epon embedding and processing**
614 Following cardiac perfusion with 2% PFA and 2% glutaraldehyde in PBS, eyes were immersion fixed in
615 the same fixative overnight at 4^{o}C, treated for 90 minutes in osmium tetroxide on ice, dehydrated in an
616 ethanol series, embedded in Epon, sectioned at 0.5 um thickness, and stained with toluidine blue.
617
618 **Confocal microscopy**
619 Confocal images were captured with a Zeiss LSM700 confocal microscope (20x or 40x objective) using
620 Zen Black 2012 software, and processed with Image J, Adobe Photoshop, and Adobe Illustrator software.
621 For experiments with control and mutant tissues, tissue processing, confocal imaging, and image
622 processing were performed identically across genotypes unless stated otherwise.

623
CNV quantification
For quantification of retinal CNV in frozen sections of whole eyes, 14 um sections for analysis were
spaced >200 um apart so that each retina was sparsely sampled.

Image analysis for cell counts
The density of immune cells was quantified from retina or choroid flatmount images by manually
counting cells using the point-and-click "Cell counter" tool in Fiji-ImageJ (https:// imagej. net/ software/
fiji/). From each quadrant of the retina, a 450 um x 450 um square was selected. For the choroid, the
regions were 645 um x 645 um.

Image analysis for immunostaining intensities
SMA, NG2, and PECAM1 immunostaining intensities were quantified from images of retina flatmounts
that had been processed identically and imaged with identical confocal microscope settings. From each
quadrant of the retina, a 450 um x 450 um square was selected that lacked large arteries or veins. Using
the "Analyze" tool and "Mean grey value" function Fiji-ImageJ (https:// imagej. net/ software/ fiji/), the
mean intensity values were determined for the individual immunostaining channel. The same method
was used for ICAM1 intensities, except the region analyzed consisted of an entire quadrant of retina.
For each experiment, which included experimental and control retinas, the mean pixel intensity in the
ICAM1 channel was scaled for both experimental and control retinas so that the mean of the control
value was set to 1.0.

Purification of brain vascular fragments
Vascular fragments were purified from P14 control and *VEcadCreER;Tgfbr1$^{CKO/-}$* mouse brains as
described previously (Hartz et al., 2018), with some modifications. For each preparation, one mouse
was anesthetized using isoflurane, surface sterilized with 70% ethanol, sacrificed by cervical dislocation,
and its brain dissected and transferred to a 10 cm tissue culture dish. The brain was minced into small
pieces (~1 × 1 mm) with a razor blade (100 strokes in each orthogonal direction) with two drops of
Dulbecco's Phosphate Buffered Saline (DPBS; no calcium, no magnesium) added to maintain moisture
(Gibco, 14190144).

The minced brain tissue was suspended in 10 ml of ice-cold DPBS supplemented with 5 mM EDTA
(Invitrogen, AM9262) and 60 U/ml of RNasin-Plus RNase Inhibitor (Promega, N2615), and gently
homogenized using a smooth pestle in a Thomas Pestle Tissue Grinder (3431E45) on ice with 5 strokes.
The sample was incubated on ice for 10 minutes, and then gently homogenized with 40 more strokes.
To monitor the release of brain capillaries, 5 µl of the homogenate was briefly incubated at room
temperature with DAPI and Isolectin GS-IB4 From *Griffonia simplicifolia* conjugated with Alexa Fluor™
488 (Invitrogen, I21411) and then visualized under a fluorescent microscope.

The brain vascular fragments were pelleted by two rounds of centrifugation through 15% Ficoll PM 400
(Sigma-Aldrich, F4375). For the first round of centrifugation, 10 ml of the homogenate was vigorously
mixed with 10 ml of 30% Ficoll PM 400. The sample was aliquoted into three 50 ml heavy-walled glass
centrifuge tubes with round bottoms (Marienfeld, 3933041) and each tube was loaded with 6.7 ml of
sample and then centrifuged at 5,800 x g for 20 min at 4°C. The pellets, containing enriched vascular
fragments, were saved. For the second round of centrifugation, the supernatant from the first round of
centrifugation, which included a layer of myelin at its top, was transferred to a fresh 50 ml glass tube
and vigorously mixed. The sample was aliquoted into three heavy-walled glass centrifuge tubes and
centrifuged as described for the first round. The pellets from the two rounds of centrifugation were

671 resuspended and pooled in a total volume of 10 ml of DPBS supplemented with 1% BSA and 40 U/ml of
672 RNasin-Plus RNase Inhibitor ("DPBS+BR"). The sample was centrifuged in a heavy-walled glass centrifuge
673 tube with a round bottom at 300 x g for 10 min at 4°C. The pellets were resuspended and pooled in a
674 total volume of 1 ml of DPBS+BR by gently pipetting 10 times with a 1 ml pipette tip.
675
The 1 ml suspension was filtered through a Nylon Mesh Filter (Tisch Scientific; 300 μm mesh, ME17240)
to remove large tissue fragments. The material trapped by the filter was rinsed one time with 1 ml of
DPBS+BR. The resulting 2-ml suspension was then filtered through a PluriStrainer 20 μm filter (Cell
Strainer, 43-50020-03) to capture vascular fragments. The PluriStrainer 20 μm filter was placed on top
of a 50-ml tube and loaded in several steps with the suspension, which was allowed to drain by gravity
flow. The vascular fragments retained on the filter were then washed once with 1 ml of DPBS+BR. The
vascular fragments were recovered from the surface of the filter with several washes with 1 ml of
DPBS+BR. The resulting 3-ml sample was centrifuged in a 12 ml heavy-walled glass centrifuge tube with
a round bottom (Marienfeld, 3933011, 12 ml) at 300 x g for 10 min at 4°C. The pellet was resuspended
in 500 ul of DPBS buffer with 40 U/ml of RNasin Plus RNase Inhibitor using a 1 ml Low Retention Pipette
Tip (1000 uL Filtered Pipette Tips for Rainin LTS Pipette - RNase and DNase Free; 3840/CS, LTS-1000FT-
CS). The suspended vascular fragments were placed in a 1.5-ml Low Retention tube (Thermo Scientific,
3451) and centrifuged at 300 x g for 10 min at 4°C. The supernatant was removed, and the pellet was
frozen on dry ice and stored at -80°C.
690
snRNAseq
Single nucleus (sn)RNA-seq libraries were prepared using the PIPseq T20 3' Single Cell RNA Kit v4.0 PLUS
(Fluent BioSiences; Clark et al., 2023). Frozen tissue was suspended in homogenization buffer (0.25 M
sucrose, 25 mM KCl, 5 mM $MgCl_2$, 20 mM Tricine-KOH,pH 7.8) supplemented with 1 mM DTT, 0.15 mM
spermine, 0.5 mM spermidine, EDTA free protease inhibitor (Roche 11836 170 001), 0.5% IGEPAL-630,
and 40 U/mL Protector RNase Inhibitor (Sigma, 03335402001). Samples were homogenized in a 2 mL
Dounce homogenizer, using 15 strokes with a loose-fitting pestle followed by 30 strokes with a tight-
fitting pestle. The sample was filtered through a 10 μm filter (CellTrix, Sysmex, 04-004-2324), and nuclei
were pelleted in Low Retention 1.5 mL microcentrifuge tubes for 5 min at 500 x g at 4^oC. Nuclei were
washed twice with Nuclei Suspension Buffer (supplied in the Fluent BioSciences kit) supplemented with
1% BSA and 40 U/mL Protector RNase Inhibitor. Nuclei were counted, and ~40,000 nuclei were used for
library production following the Fluent BioSiences protocol. The resulting snRNA-seq libraries were
sequenced on an Illumina NovaSeq X Plus sequencer. Vascular fragments from two mouse brains were
pooled for each snRNAseq library.
705
Analysis of snRNAseq data
Reads were aligned with the PIPseeker program (Fluent BioSiences, version 3.3.0) using the pipseeker-
gex-reference-GRCm39-2023.04 index provided by Fluent BioSiences. The CellBender program was used
to detect empty droplets and remove background (version 0.3.2; Fleming et al., 2023). The SOLO
program was used to remove doublets (scVI-tools 1.2.2-post2; Bernstein et al., 2020). The data were
normalized using a regularized negative binomial regression algorithm implemented with the
SCTransform function (Hafemeister and Satija, 2019). UMAP dimensional reduction was performed
using the R uwot package (https://github.com/jlmelville/uwot) integrated into the Seurat R package
(Melville, 2022). Data for the various scatter plots were extracted using the Seurat AverageExpression
function, and differential gene expression was analyzed using the Seurat FindMarkers function. The
Wilcoxon Rank Sum test was used to calculate p-values. The p-values were adjusted with a Bonferroni
correction using all genes in the dataset. Data exploration, analysis, and plotting were performed using

RStudio (RStudio Team, 2020), the tidyverse collection of R packages (Wickham, 2017), and ggplot2
(Wickham, 2009). Dotplots were generated with the default settings, including default normalization.
720
721 **GSEA analysis**
722 For Gene Set Enrichment Analysis (GSEA; Subramanian et al., 2005), genes were ranked by the fold
723 expression change between control and mutant datasets. The ranked gene list was used to detect
724 enriched gene sets within the Broad Institute Hallmark Gene Sets using the fgsea R package
725 (https://github.com/ctlab/fgsea; Korotkevich et al., 2019).
726
Statistical analysis
All statistical values are presented as mean ± SD. The Wilcoxon rank sum test was used to measure
statistical significance. Statistical tests were carried out using the following web sites:
https://www.socscistatistics.com/tests/signedranks/default2.aspx
and https://www.omnicalculator.com/statistics/wilcoxon-rank-sum-test#how-do-i-calculate-wilcoxon-
rank-sum-test. The statistical significance is represented graphically as n.s., not significant (i.e. $p>0.05$);
*, $p <0.05$; **, $p < 0.01$; ***, $p < 0.001$; ****, $p < 0.0001$.
734
735 **Data availability**
736 The snRNAseq data have been deposited in the Gene Expression Omnibus (GEO) database (GSEXX).
737
738
739

740 **Acknowledgements**
741 Supported by the Howard Hughes Medical Institute. The authors thank David Mohr (Genetic Resources
742 Core Facility, Johns Hopkins School of Medicine) for assistance with NextGen sequencing, and Philip
743 Seegren for helpful comments on the manuscript.

744
745
Author Contributions
Yanshu Wang - Conceptualization, Formal analysis, Investigation, Methodology, Data curation,
Validation, Writing - original draft, Writing - review and editing.
Amir Rattner - Formal analysis, Investigation, Methodology, Data curation, Validation, Writing - original
draft, Writing - review and editing.
Zhongming Li - Investigation, Methodology, Validation.
Philip M. Smallwood - Investigation, Methodology.
Jeremy Nathans - Conceptualization, Formal analysis, Investigation, Methodology, Validation,
Visualization, Writing - original draft, Writing - review and editing, Funding acquisition, Project
administration, Supervision.
756
757
758
759 **References**
760

761 Alarcon-Martinez L, Yilmaz-Ozcan S, Yemisci M, Schallek J, Kılıç K, Can A, Di Polo A, Dalkara T. (2018)
762 Capillary pericytes express α-smooth muscle actin, which requires prevention of filamentous-actin
763 depolymerization for detection. Elife 7:e34861.

765 Aliahmad P, Seksenyan A, Kaye J. (2012) The many roles of TOX in the immune system. Curr Opin
766 Immunol 24:173-177.

767
768 Allinson KR, Lee HS, Fruttiger M, McCarty JH, Arthur HM. (2012) Endothelial expression of TGFbeta type
769 II receptor is required to maintain vascular integrity during postnatal development of the central
770 nervous system. PLoS One 7:e39336.

771
772 Aluwihare P, Mu Z, Zhao Z, Yu D, Weinreb PH, Horan GS, Violette SM, Munger JS. (2009) Mice that lack
773 activity of alphavbeta6- and alphavbeta8-integrins reproduce the abnormalities of Tgfb1- and Tgfb3-null
774 mice. J Cell Sci 122:227-232.

775
776 Alvarez JI, Dodelet-Devillers A, Kebir H, Ifergan I, Fabre PJ, Terouz S, Sabbagh M, Wosik K, Bourbonnière
777 L, Bernard M, van Horssen J, de Vries HE, Charron F, Prat A. (2011) The Hedgehog pathway promotes
778 blood-brain barrier integrity and CNS immune quiescence. Science 334:1727-1731.

779
780 Aman J, Margadant C. (2023) Integrin-Dependent Cell-Matrix Adhesion in Endothelial Health and
781 Disease. Circ Res 132:355-378.

782
783 Amersfoort J, Eelen G, Carmeliet P. (2022) Immunomodulation by endothelial cells - partnering up with
784 the immune system? Nat Rev Immunol 22:576-588

785
786 Armento A, Ueffing M, Clark SJ. (2021) The complement system in age-related macular degeneration.
787 Cell Mol Life Sci 78:4487-4505.

788
789 Arnold TD, Ferrero GM, Qiu H, Phan IT, Akhurst RJ, Huang EJ, Reichardt LF. (2012) Defective retinal
790 vascular endothelial cell development as a consequence of impaired integrin αVβ8-mediated activation
791 of transforming growth factor-β. J Neurosci 32:1197-1206.

792
793 Arnold TD, Niaudet C, Pang MF, Siegenthaler J, Gaengel K, Jung B, Ferrero GM, Mukouyama YS, Fuxe J,
794 Akhurst R, Betsholtz C, Sheppard D, Reichardt LF. (2014) Excessive vascular sprouting underlies cerebral
795 hemorrhage in mice lacking αVβ8-TGFβ signaling in the brain. Development 141:4489-4499.

796
797 BD Biosciences (2024) CD Marker Handbook.
798 https://www.bdbiosciences.com/content/dam/bdb/marketing-documents/cd_marker_handbook.pdf

799
800 Beck SC, Karlstetter M, Garcia Garrido M, Feng Y, Dannhausen K, Mühlfriedel R, Sothilingam V, Seebauer
801 B, Berger W, Hammes HP, Seeliger MW, Langmann T. (2018) Cystoid edema, neovascularization and
802 inflammatory processes in the murine Norrin-deficient retina. Sci Rep 8:5970.

803
804 Bernstein NJ, Fong NL, Lam I, Roy MA, Hendrickson DG, Kelley DR. (2020) Solo: Doublet Identification in
805 Single-Cell RNA-Seq via Semi-Supervised Deep Learning. Cell Syst 11:95-101

806
807 Blanchard L, Girard JP. (2021) High endothelial venules (HEVs) in immunity, inflammation and cancer.
808 Angiogenesis 24:719-753.

809
Braunger BM, Leimbeck SV, Schlecht A, Volz C, Jägle H, Tamm ER. (2015) Deletion of ocular transforming
growth factor β signaling mimics essential characteristics of diabetic retinopathy. Am J Pathol 185:1749-
1768.

Cambier S, Gline S, Mu D, Collins R, Araya J, Dolganov G, Einheber S, Boudreau N, Nishimura SL. (2005)
Integrin alpha(v)beta8-mediated activation of transforming growth factor-beta by perivascular
astrocytes: an angiogenic control switch. Am J Pathol 166:1883-1894.

Chan CC, Ross RJ, Shen D, Ding X, Majumdar Z, Bojanowski CM, Zhou M, Salem N Jr, Bonner R, Tuo J.
(2008) Ccl2/Cx3cr1-deficient mice: an animal model for age-related macular degeneration. Ophthalmic
Res 40:124-128.

Chen T, Dai Y, Hu C, Lin Z, Wang S, Yang J, Zeng L, Li S, Li W. (2024) Cellular and molecular mechanisms of
the blood-brain barrier dysfunction in neurodegenerative diseases. Fluids Barriers CNS 21:60.

Chibber R, Ben-Mahmud BM, Chibber S, Kohner EM. (2007) Leukocytes in diabetic retinopathy. Curr
Diabetes Rev 3:3-14.

Clark IC, Fontanez KM, Meltzer RH, Xue Y, Hayford C, May-Zhang A, D'Amato C, Osman A, Zhang JQ,
Hettige P, Ishibashi JSA, Delley CL, Weisgerber DW, Replogle JM, Jost M, Phong KT, Kennedy VE, Peretz
CAC, Kim EA, Song S, Karlon W, Weissman JS, Smith CC, Gartner ZJ, Abate AR. (2023) Microfluidics-free
single-cell genomics with templated emulsification. Nat Biotechnol 41:1557-1566.

Denes A, Hansen CE, Oezorhan U, Figuerola S, de Vries HE, Sorokin L, Planas AM, Engelhardt B,
Schwaninger M. (2024) Endothelial cells and macrophages as allies in the healthy and diseased brain.
Acta Neuropathol 147:38.

Ding ZM, Babensee JE, Simon SI, Lu H, Perrard JL, Bullard DC, Dai XY, Bromley SK, Dustin ML, Entman ML,
Smith CW, Ballantyne CM. (1999) Relative contribution of LFA-1 and Mac-1 to neutrophil adhesion and
migration. J Immunol 163:5029-5038.

Dong X, Zhao B, Iacob RE, Zhu J, Koksal AC, Lu C, Engen JR, Springer TA. (2017) Force interacts with
macromolecular structure in activation of TGF-beta. Nature 542:55-59.

Fabian-Jessing BK, Jakobsen TS, Jensen EG, Alsing S, Hansen S, Aagaard L, Askou AL, Bek T, Corydon TJ.
(2022) Animal Models of Choroidal Neovascularization: A Systematic Review. Invest Ophthalmol Vis Sci
63:11.

Fayard E, Auwerx J, Schoonjans K. (2004) LRH-1: an orphan nuclear receptor involved in development,
metabolism and steroidogenesis. Trends Cell Biol 14:250-260.

Fleming SJ, Chaffin MD, Arduini A, Akkad AD, Banks E, Marioni JC, Philippakis AA, Ellinor PT, Babadi M.
(2023) Unsupervised removal of systematic background noise from droplet-based single-cell
experiments using CellBender. Nat Methods 20:1323-1335

Freund KB, Ho IV, Barbazetto IA, Koizumi H, Laud K, Ferrara D, Matsumoto Y, Sorenson JA, Yannuzzi L.
(2008) Type 3 neovascularization: the expanded spectrum of retinal angiomatous proliferation.
Retina 28:201-211.

Gass, JD (1997) Stereoscopic Atlas of Macular Diseases. St. Louis: CV Mosby.

Gerber HP, Hillan KJ, Ryan AM, Kowalski J, Keller GA, Rangell L, Wright BD, Radtke F, Aguet M, Ferrara N.
(1999) VEGF is required for growth and survival in neonatal mice. Development 126:1149-1159.
863
864 Goumans MJ, Liu Z, ten Dijke P. (2009) TGF-beta signaling in vascular biology and dysfunction. Cell Res
865 19:116-127.
866
867 Gross NE, Aizman A, Brucker A, Klancnik JM Jr, Yannuzzi LA. (2005) Nature and risk of neovascularization
868 in the fellow eye of patients with unilateral retinal angiomatous proliferation. Retina 25:713-718.
869
870 Grossniklaus HE, Kang SJ, Berglin L. (2010) Animal models of choroidal and retinal neovascularization.
871 Prog Retin Eye Res 29:500-519.
872
873 Guerrero PA, McCarty JH. (2018) Integrins in Vascular Development and Pathology. Adv Pharmacol
874 81:129-153.
875
876 Hachana S, Larrivée B. (2022) TGF-β Superfamily Signaling in the Eye: Implications for Ocular Pathologies.
877 Cells 11:2336.
878
879 Hafemeister C, Satija R. (2019) Normalization and variance stabilization of single-cell RNA-seq data using
880 regularized negative binomial regression. Genome Biol 20:296.
881
882 Hartz AMS, Schulz JA, Sokola BS, Edelmann SE, Shen AN, Rempe RG, Zhong Y, Seblani NE, Bauer B. (2018)
883 Isolation of Cerebral Capillaries from Fresh Human Brain Tissue. J Vis Exp 12:57346.
884
885 Heckenlively JR, Hawes NL, Friedlander M, Nusinowitz S, Hurd R, Davisson M, Chang B. (2003) Mouse
886 model of subretinal neovascularization with choroidal anastomosis. Retina 23:518-522.
887
888 Heldin CH, Moustakas A. (2016) Signaling Receptors for TGF-β Family Members. Cold Spring Harb
889 Perspect Biol 8:a022053.
890
891 Helotera H, Kaarniranta K. (2022) A Linkage between Angiogenesis and Inflammation in Neovascular
892 Age-Related Macular Degeneration. Cells 11:3453.
893
894 Heng JS, Hackett SF, Stein-O'Brien GL, Winer BL, Williams J, Goff LA, Nathans J. (2019) Comprehensive
895 analysis of a mouse model of spontaneous uveoretinitis using single-cell RNA sequencing. Proc Natl Acad
896 Sci U S A 116:26734-26744.
897
898 Hrvatin S, Hochbaum DR, Nagy MA, Cicconet M, Robertson K, Cheadle L, Zilionis R, Ratner A, Borges-
899 Monroy R, Klein AM, Sabatini BL, Greenberg ME. (2018) Single-cell analysis of experience-dependent
900 transcriptomic states in the mouse visual cortex. Nat Neurosci 21:120-129.
901
902 Hu W, Jiang A, Liang J, Meng H, Chang B, Gao H, Qiao X. (2008) Expression of VLDLR in the retina and
903 evolution of subretinal neovascularization in the knockout mouse model's retinal angiomatous
904 proliferation. Invest Ophthalmol Vis Sci 49:407-415.
905
906 Hu X, Leak RK, Thomson AW, Yu F, Xia Y, Wechsler LR, Chen J. (2018) Promises and limitations of immune
907 cell-based therapies in neurological disorders. Nat Rev Neurol 14:559-568.
908

909   Imamura Y, Noda S, Hashizume K, Shinoda K, Yamaguchi M, Uchiyama S, Shimizu T, Mizushima Y,
910   Shirasawa T, Tsubota K. (2006) Drusen, choroidal neovascularization, and retinal pigment epithelium
911   dysfunction in SOD1-deficient mice: a model of age-related macular degeneration. Proc Natl Acad Sci
912   USA 103:11282-11287.
913
914   Joussen AM, Poulaki V, Le ML, Koizumi K, Esser C, Janicki H, Schraermeyer U, Kociok N, Fauser S, Kirchhof
915   B, Kern TS, Adamis AP. (2004) A central role for inflammation in the pathogenesis of diabetic
916   retinopathy. FASEB J 18:1450.
917
918   Jung JJ, Chen CY, Mrejen S, Gallego-Pinazo R, Xu L, Marsiglia M, Boddu S, Freund KB. (2014) The
919   incidence of neovascular subtypes in newly diagnosed neovascular age-related macular degeneration.
920   Am J Ophthalmol 158:769-779.
921
922   Jung SH, Kim KA, Sohn SW, Yang SJ. (2014) Association of aqueous humor cytokines with the
923   development of retinal ischemia and recurrent macular edema in retinal vein occlusion. Invest
924   Ophthalmol Vis Sci 55:2290-2296.
925
926   Kauppinen A, Paterno JJ, Blasiak J, Salminen A, Kaarniranta K. (2016) Inflammation and its role in age-
927   related macular degeneration. Cell Mol Life Sci 73:1765-1786.
928
929   Korotkevich G, Sukhov V, Sergushichev A. 2019. Fast gene set enrichment analysis. bioRxiv. DOI:
930   https://doi. org/10.1101/060012
931
932   Kulkarni AB, Huh CG, Becker D, Geiser A, Lyght M, Flanders KC, Roberts AB, Sporn MB, Ward JM, Karlsson
933   S. (1993) Transforming growth factor beta 1 null mutation in mice causes excessive inflammatory
934   response and early death. Proc Natl Acad Sci U S A 90:770-774.
935
936   Lacar B, Linker SB, Jaeger BN, Krishnaswami SR, Barron JJ, Kelder MJE, Parylak SL, Paquola ACM,
937   Venepally P, Novotny M, O'Connor C, Fitzpatrick C, Erwin JA, Hsu JY, Husband D, McConnell MJ, Lasken
938   R, Gage FH. (2016) Nuclear RNA-seq of single neurons reveals molecular signatures of activation. Nat
939   Commun 7:11022.
940
941   Lange CA, Luhmann UF, Mowat FM, Georgiadis A, West EL, Abrahams S, Sayed H, Powner MB, Fruttiger
942   M, Smith AJ, Sowden JC, Maxwell PH, Ali RR, Bainbridge JW. (2012) Von Hippel-Lindau protein in the RPE
943   is essential for normal ocular growth and vascular development. Development 139:2340-2350.
944
945   Larsson J, Goumans MJ, Sjöstrand LJ, van Rooijen MA, Ward D, Levéen P, Xu X, ten Dijke P, Mummery CL,
946   Karlsson S. (2001) Abnormal angiogenesis but intact hematopoietic potential in TGF-beta type I receptor-
947   deficient mice. EMBO J 20:1663-1673.
948
949   Levéen P, Larsson J, Ehinger M, Cilio CM, Sundler M, Sjöstrand LJ, Holmdahl R, Karlsson S. (2002) Induced
950   disruption of the transforming growth factor beta type II receptor gene in mice causes a lethal
951   inflammatory disorder that is transplantable. Blood 100:560-568.
952
953   Lindahl P, Betsholtz C. (1998) Not all myofibroblasts are alike: revisiting the role of PDGF-A and PDGF-B
954   using PDGF-targeted mice. Curr Opin Nephrol Hypertens 7:21-26.
955
956   Luhmann UF, Lin J, Acar N, Lammel S, Feil S, Grimm C, Seeliger MW, Hammes HP, Berger W.

957   (2005) Role of the Norrie disease pseudoglioma gene in sprouting angiogenesis during development of
958   the retinal vasculature. Invest Ophthalmol Vis Sci 46:3372-3382.
959

960   Luo L, Uehara H, Zhang X, Das SK, Olsen T, Holt D, Simonis JM, Jackman K, Singh N, Miya TR, Huang W,
961   Ahmed F, Bastos-Carvalho A, Le YZ, Mamalis C, Chiodo VA, Hauswirth WW, Baffi J, Lacal PM, Orecchia A,
962   Ferrara N, Gao G, Young-Hee K, Fu Y, Owen L, Albuquerque R, Baehr W, Thomas K, Li DY, Chalam KV,
963   Shibuya M, Grisanti S, Wilson DJ, Ambati J, Ambati BK. (2013) Photoreceptor avascular privilege is
964   shielded by soluble VEGF receptor-1. Elife 2:e00324.
965

966   Malek G, Johnson LV, Mace BE, Saloupis P, Schmechel DE, Rickman DW, Toth CA, Sullivan PM, Bowes
967   Rickman C. (2005) Apolipoprotein E allele-dependent pathogenesis: a model for age-related retinal
968   degeneration. Proc Natl Acad Sci USA 102:11900-11905.
969

970   Massagué J, Sheppard D. (2023) TGF-β signaling in health and disease. Cell 186:4007-4037.
971

972   Melville J. (2022) Uwot: the uniform manifold approximation and projection (UMAP) method for
973   Dimensionality reduction. 0.1.14. R Package. https://cran.r-project.org/web/packages/uwot/index.html
974

975   Miyamoto K, Khosrof S, Bursell SE, Rohan R, Murata T, Clermont AC, Aiello LP, Ogura Y, Adamis AP.
976   (1999) Prevention of leukostasis and vascular leakage in streptozotocin-induced diabetic retinopathy via
977   intercellular adhesion molecule-1 inhibition. Proc Natl Acad Sci USA 96:10836-10841.
978

979   Mo A, Mukamel EA, Davis FP, Luo C, Henry GL, Picard S, Urich MA, Nery JR, Sejnowski TJ, Lister R, Eddy
980   SR, Ecker JR, Nathans J. (2015) Epigenomic Signatures of Neuronal Diversity in the Mammalian Brain.
981   Neuron 86:1369-1384.
982

983   Montezuma SR, Sobrin L, Seddon JM. (2007) Review of genetics in age related macular degeneration.
984   Semin Ophthalmol 22:229-240.
985

986   Monvoisin A, Alva JA, Hofmann JJ, Zovein AC, Lane TF, Iruela-Arispe ML. (2006) VE-cadherin-CreERT2
987   transgenic mouse: a model for inducible recombination in the endothelium. Dev Dyn 235:3413-3422.
988

989   Mu Z, Yang Z, Yu D, Zhao Z, Munger JS. (2008) TGFbeta1 and TGFbeta3 are partially redundant effectors
990   in brain vascular morphogenesis. Mech Dev 125:508-516.
991

992   Muniyandi A, Hartman GD, Song Y, Mijit M, Kelley MR, Corson TW. (2023) Beyond VEGF: Targeting
993   Inflammation and Other Pathways for Treatment of Retinal Disease. J Pharmacol Exp Ther 386:15-25.
994

995   Nguyen HL, Lee YJ, Shin J, Lee E, Park SO, McCarty JH, Oh SP. (2011) TGF-β signaling in endothelial cells,
996   but not neuroepithelial cells, is essential for cerebral vascular development. Lab Invest 91:1554-1563.
997

998   Ohno-Matsui K, Hirose A, Yamamoto S, Saikia J, Okamoto N, Gehlbach P, Duh EJ, Hackett S, Chang M,
999   Bok D, Zack DJ, Campochiaro PA. (2002) Inducible expression of vascular endothelial growth factor in
1000   adult mice causes severe proliferative retinopathy and retinal detachment. Am J Pathol 160:711-719.
1001

1002   Omarova S, Charvet CD, Reem RE, Mast N, Zheng W, Huang S, Peachey NS, Pikuleva IA. (2012) Abnormal
1003   vascularization in mouse retina with dysregulated retinal cholesterol homeostasis. J Clin Invest
1004   122:3012-3023.

1005
1006 Pober JS, Sessa WC. (2014) Inflammation and the blood microvascular system. Cold Spring Harb Perspect
1007 Biol 7:a016345.

1008
1009 Profaci CP, Munji RN, Pulido RS, Daneman R. J (2020) The blood-brain barrier in health and disease:
1010 Important unanswered questions. Exp Med 217:e20190062.

1011
1012 Qiang W, Wei R, Chen Y, Chen D. (2021) Clinical Pathological Features and Current Animal Models of
1013 Type 3 Macular Neovascularization. Front Neurosci 15:734860.

1014
1015 Rattner A, Williams J, Nathans J. (2019) Roles of HIFs and VEGF in angiogenesis in the retina and brain. J
1016 Clin Invest 129:3807-3820.

1017
1018 Rattner A, Wang Y, Nathans J. (2022) Signaling Pathways in Neurovascular Development. Annu Rev
1019 Neurosci 45:87-108.

1020
1021 Ravichandran KA, Heneka MT. (2024) Inflammasomes in neurological disorders - mechanisms and
1022 therapeutic potential. Nat Rev Neurol 20:67-83.

1023
1024 Ritson M, Wheeler-Jones CPD, Stolp HB. (2024) Endothelial dysfunction in neurodegenerative disease: Is
1025 endothelial inflammation an overlooked druggable target? J Neuroimmunol 391:578363.

1026
1027 Ropper AH, Samuels MA, Klein JP, Prasad S. (2023) Adams and Victor's Priciples of Neurology (twelfth
1028 edition). McGraw-Hill.

1029
1030 Rowan S, Cepko CL. (2004) Genetic analysis of the homeodomain transcription factor Chx10 in the retina
1031 using a novel multifunctional BAC transgenic mouse reporter. Dev Biol 271:388-402.

1032
1033 RStudio Team. (2020) RStudio: Integrated Development Environment for R. RStudio. PBC, Boston,
1034 MA. https://rstudio. com

1035
1036 Sackstein R. (2005) The lymphocyte homing receptors: gatekeepers of the multistep paradigm. Curr Opin
1037 Hematol 12:444-450.

1038
1039 Schlecht A, Leimbeck SV, Jägle H, Feuchtinger A, Tamm ER, Braunger BM. (2017) Deletion of Endothelial
1040 Transforming Growth Factor-β Signaling Leads to Choroidal Neovascularization. Am J Pathol 187:2570-
1041 2589.

1042
1043 Shi M, Zhu J, Wang R, Chen X, Mi L, Walz T, Springer TA. (2011) Latent TGF-beta structure and activation.
1044 Nature 474:343-349.

1045
1046 Shull MM, Ormsby I, Kier AB, Pawlowski S, Diebold RJ, Yin M, Allen R, Sidman C, Proetzel G, Calvin D, et
1047 al. (1992) Targeted disruption of the mouse transforming growth factor-beta 1 gene results in multifocal
1048 inflammatory disease. Nature 359:693-699.

1049
1050 Stockinger B, Diaz OE, Wincent E. (2024) The influence of AHR on immune and tissue biology. EMBO Mol
1051 Med 16:2290-2298.
1052

Stone J, Itin A, Alon T, Pe'er J, Gnessin H, Chan-Ling T, Keshet E. (1995) Development of retinal
vasculature is mediated by hypoxia-induced vascular endothelial growth factor (VEGF) expression by
neuroglia. J Neurosci 15:4738-4747.

Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A, Pomeroy SL,
Golub TR, Lander ES, Mesirov JP. (2005) Gene set enrichment analysis: A knowledge-based approach for
interpreting genome-wide expression profiles. Proc Natl Acad Sci USA 102:15545–15550.

Takeda A, Baffi JZ, Kleinman ME, Cho WG, Nozaki M, Yamada K, Kaneko H, Albuquerque RJ, Dridi S, Saito
K, Raisler BJ, Budd SJ, Geisen P, Munitz A, Ambati BK, Green MG, Ishibashi T, Wright JD, Humbles AA,
Gerard CJ, Ogura Y, Pan Y, Smith JR, Grisanti S, Hartnett ME, Rothenberg ME, Ambati J. (2009) CCR3 is a
target for age-related macular degeneration diagnosis and therapy. Nature 460:225-230.

Tang J, Kern TS. (2014) Inflammation in diabetic retinopathy. Prog Retin Eye Res 30:343-358.

Tobe T, Okamoto N, Vinores MA, Derevjanik NL, Vinores SA, Zack DJ, Campochiaro PA. (1998) Evolution
of neovascularization in mice with overexpression of vascular endothelial growth factor in
photoreceptors. Invest Ophthalmol Vis Sci 39:180-188.

Travis MA, Sheppard D. (2014) TGF-β activation and function in immunity. Annu Rev Immunol 32:51-82.

Wälchli T, Wacker A, Frei K, Regli L, Schwab ME, Hoerstrup SP, Gerhardt H, Engelhardt B. (2015) Wiring
the Vascular Network with Neural Cues: A CNS Perspective. Neuron 87:271-296.

Wälchli T, Bisschop J, Carmeliet P, Zadeh G, Monnier PP, De Bock K, Radovanovic I. (2023) Shaping the
brain vasculature in development and disease in the single-cell era. Nat Rev Neurosci 24:271-298.

Wang Y, Huso D, Cahill H, Ryugo D, Nathans J. (2001) Progressive cerebellar, auditory, and esophageal
dysfunction caused by targeted disruption of the frizzled-4 gene. J Neurosci 21:4761-4771.

Wang Y, Rattner A, Zhou Y, Williams J, Smallwood PM, Nathans J. (2012) Norrin/Frizzled4 signaling in
retinal vascular development and blood brain barrier plasticity. Cell 151:1332-1344.

Wang Y, Cho C, Williams J, Smallwood PM, Zhang C, Junge HJ, Nathans J. (2018) Interplay of the Norrin
and Wnt7a/Wnt7b signaling systems in blood-brain barrier and blood-retina barrier development and
maintenance. Proc Natl Acad Sci U S A 115:E11827-E11836.

Wang Y, Gao S, Cao F, Yang H, Lei F, Hou S. (2024) Ocular immune-related diseases: molecular
mechanisms and therapy. MedComm 5:e70021.

Wickham H. (2009) Ggplot2: Elegant Graphics for Data Analysis. New York: Springer-Verlag.

Wickham H. (2017) tidyverse: Easily Install and Load the 'Tidyverse'. 1.2.1. R Package. https://CRAN.R-
project.org/package=tidyverse

Ye X, Wang Y, Cahill H, Yu M, Badea TC, Smallwood PM, Peachey NS, Nathans J. (2009) Norrin, frizzled-4,
and Lrp5 signaling in endothelial cells controls a genetic program for retinal vascularization. Cell
139:285-298.

1101

1102 Zarkada G, Howard JP, Xiao X, Park H, Bizou M, Leclerc S, Künzel SE, Boisseau B, Li J, Cagnone G, Joyal JS,
1103 Andelfinger G, Eichmann A, Dubrac A. (2021) Specialized endothelial tip cells guide neuroretina
1104 vascularization and blood-retina-barrier formation. Dev Cell 56:2237-2251

1105

1106 Zhou Y, Wang Y, Tischfield M, Williams J, Smallwood PM, Rattner A, Taketo MM, Nathans J.
1107 (2014) Canonical WNT signaling components in vascular development and barrier formation.
1108 J Clin Invest 124:3825-3846.

1109

1110

1111 **Figure Legends**

1112

Figure 1. EC-specific loss of TGF-beta signaling leads to attenuated retinal vascular development, CNV,
and anastomoses between retinal and choroidal vasculatures. (A) *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas
showing CNV (white arrows in central panels), and vascular invasion of the outer nuclear layer (white
arrows in lower panels), both with associated CD45+ immune cells. (B) *VEcadCreER;Tgfbr2$^{CKO/-}$* retinas
showing CNV (white arrows in central panels), and an anastomosis between retinal and choroidal
vasculatures (white arrows in lower panels), with intra-retinal EC marker CLDN5, choroidal EC marker
PLVAP, and pan-vascular marker COL4 (collagen4). (C) Upper right panels, in the *VEcadCreER;Tgfbr1$^{CKO/-}$*
retina, CNV (white arrows) is derived from choroidal vasculature, marked by PLVAP. Lower right panels,
CNV (white arrows) is present in the subretinal space, i.e. on the retinal side of the RPE, which is marked
by RPE65. Abbreviations: Ch., choroid; ONL, outer nuclear layer; INL, inner nuclear layer; GCL, ganglion
cell layer; RPE, retinal pigment epithelium. The ages of the mice are indicated in postnatal days (P) for
this and all other figures. Scale bars, 100 um.

1125

Figure 2. Vascular architecture in retinas with EC-specific loss of TGF-beta signaling or global loss of
Norrin/Fzd4 signaling. (A) COL4 immunostaining of control and *VEcadCreER;Tgfbr1$^{CKO/-}$* flatmount retinas
showing arteries, veins, and capillaries in the control retina and a high density of vascular tufts in the
VEcadCreER;Tgfbr1$^{CKO/-}$ retina. (B) False color images from the indicated genotypes and ages showing a
stacked Z-series of flatmount retinas color-coded by the depth of the vasculature. For control retinas
the blue-green-red color scheme corresponds to the inner two-thirds of the retina: blue, vitreal surface;
green, ganglion cell layer and inner plexiform layer; and red, inner nuclear layer and outer plexiform
layer. For mutant retinas, the blue-green-red color scheme corresponds to a shallower depth, as the
most deeply penetrating vascular tufts go only as far as the inner edge of the inner nuclear layer: blue,
vitreal surface; green, ganglion cell layer; and red, inner plexiform layer. Left column of three panels:
P14 control retina (upper image; white arrows point to tip cells in the IPL) and two regions from a P14
VEcadCreER;Tgfbr1$^{CKO/-}$ retina (lower). Center column of three panels: P26 control retina (upper) and
two regions from a P26 *VEcadCreER;Tgfbr1$^{CKO/-}$* retina (lower). Right column of three panels, ~P30 *Ndp^{KO}*
retina (upper) and two regions from a ~P30 *Fzd4$^{-/-}$* retina (lower). All images are at the same
magnfication. (C) Flatmount of a P37 *VEcadCreER;Tgfbr1$^{CKO/-}$* retina showing PLVAP expression and
Sulfo-NHS-biotin accumulation in vascular tufts. The vascular tufts are variably present with loss of
TGFBR1 or TGFBR2. Scale bar in (A), 1 mm. Scale bar in (B), 100 um. Scale bar in (C), 200 um.

1143

Figure 3. Immune cells in retinas with EC-specific loss of TGF-beta signaling or global loss of Norrin/Fzd4
signaling. (A) Control and *VEcadCreER;Tgfbr1$^{CKO/-}$* retina flatmounts (left pair of panels) with enlarged
insets (right pair of panels). (B) Control and *VEcadCreER;Tgfbr1$^{CKO/-}$* retina flatmounts (left panel) and
insets (right three panels). (C) Control and *VEcadCreER;Tgfbr1$^{CKO/-}$;Tgfbr2$^{CKO/-}$* retina flatmounts, and
control and *Fzd4$^{-/-}$* retina flatmounts. (D-G) Quantification of numbers of cells positive for the indicated

markers in 450 um x 450 um zones in the midperiphery of retina flatmounts from mice of the indicated
genotypes. In this and other panels with statistical comparisons, the bars represent mean +/- standard
deviation, and p-values, calculated using the Wilcoxon rank sum test, are shown as * <0.05, ** <0.01,
*** < 0.001, and **** <0.0001. Scale bars in whole retina panels, 1 mm. Scale bars in all other panels,
100 um.
1154
Figure 4. Single nucleus RNA sequencing from control and *VEcadCreER;Tgfbr1$^{CKO/-}$* retinas at P14. (A)
Identification of cell clusters in a Uniform Manifold Approximation and Projection (UMAP) plot, based on
established markers of retinal gene expression. The pooled control and *VEcadCreER;Tgfbr1$^{CKO/-}$* data are
shown. The horizontal arrow points to immune cells, almost entirely derived from the
VEcadCreER;Tgfbr1$^{CKO/-}$ samples, as shown in (B). (B) The contributions of control and
VEcadCreER;Tgfbr1$^{CKO/-}$ retinas to the immune cell cluster. (C) UMAP plot identifying immune cell types
within the immune cell cluster, based on established markers as shown in (D) and (E). (D) Abundances of
select transcripts in different immune cell types, as defined in (B). (E) UMAP plots for select markers
from (D) showing immune cell type-specific expression.
1164
Figure 5. Vascular anatomy and absence of immune cell infiltration in *Chx10-Cre;Vegfa$^{CKO/CKO}$* retinas.
(A) Sections of control and *Chx10-Cre;Vegfa$^{CKO/CKO}$* retinas immunostained with anti-PECAM1 to visualize
the vasculature. (B) False color images of control and *Chx10-Cre;Vegfa$^{CKO/CKO}$* flatmount retinas showing
a stacked Z-series color-coded by the depth of PECAM1 immunstained vasculature. Blue, vitreal surface;
green, inner plexiform layer; red, outer plexiform layer. (C) Control and *Chx10-Cre;Vegfa$^{CKO/CKO}$*
flatmount retinas immunostained with anti-PECAM1. (D) Control and *Chx10-Cre;Vegfa$^{CKO/CKO}$* flatmount
retinas immunostained for ASC and CD45. (E) Quantification of CD45+ cells. Scale bars: (A), (B) and (D),
100 um; (C) 500 um.
1173
Figure 6. Close association between immune cells and retinal vasculature with EC-specific loss of TGF-
beta signaling. (A) Upper six panels show a control retina flatmount. Lower six panels show a
VEcadCreER;Tgfbr1$^{CKO/+}$;Tgfbr2$^{CKO/-}$ retina flatmount. ASC and CD45 label immune cells, including
microglia. The Z-plane is indicated by the numbers at the bottom of each image. The nerve fiber layer
(Z-plane 3) and the inner plexiform layer (Z-planes 10-11) are shown schematically in (B) and (C). (B) and
(C), retinal schematics showing the relationship of the vasculature and the three retinal layers. Confocal
Z-planes are numbered at right. (D) Immune cells and venous ECs in control and *VEcadCreER;Tgfbr1$^{CKO/-}$*
retinas. In the lower image, three of the "impressions" of CD45+ immune in the distribution of PECAM1
on the EC surface are highlighted with white arrows. A, artery; V, vein. NFL, nerve fiber layer; GCL,
ganglion cell layer; IPL, inner plexiform layer; INL, inner nuclear layer; OPL, outer plexiform layer; ONL,
outer nuclear layer; IS/OS, inner segment/outer segment. Scale bars in (A), 100 um. Scale bar in (D), 50
um.
1186
Figure 7. ICAM1 in ECs and SMA in pericytes in retinas with EC-specific loss of TGF-beta signaling. (A)
ICAM1 in the retinal vasculature with EC-specific loss of TGF-beta signaling. Immunostaining conditions
and image capture settings were identical across genotypes. (B) Quantification of the relative intensities
of ICAM1 immunostaining in control vs. *VEcadCreER;Tgfbr1$^{CKO/-}$* , control vs *Fzd4$^{-/-}$* retinas, and control vs
Ndp^{KO} retinas. (C) Left, control retina flatmount showing strong SMA immunstaining in arteries,
weak/patchy SMA immunstaining in veins, and undetectable SMA immunstaining in capillaries. Right,
VEcadCreER;Tgfbr1$^{CKO/-}$ retina flatmount showing SMA immunostaining in all vessels, including vascular
tufts. NG2 immunostaining (a pericyte marker) is shown in the images below. (D) Quantification of the
relative intensities of PECAM1, NG2, and SMA immunostaining in flatmount control and
VEcadCreER;Tgfbr1$^{CKO/-}$ retinas. Bars represent mean +/- standard deviation, and p-values, calculated

using the Wilcoxon rank sum test, are shown as * <0.05, ** <0.01, *** < 0.001, and **** <0.0001. In (A)
and (C), the retinal periphery is at the right. Scale bars in (A) and (C), 500 um.
1199
Figure 8. Transient IgG extravasation, SMA accumulation in pericytes, and immune cell infiltration in the
brains of mice with EC-specific loss of TGF-beta signaling. (A) Endogenous IgG in control and
VEcadCreER;Tgfbr1$^{CKO/-}$ brains at P14 and P24. IgG accumulation is minimal in control brains at P14 and
P24, but is readily detectable in *VEcadCreER;Tgfbr1$^{CKO/-}$* brains at P14 but not at P24. (B) ECs (visualized
with PECAM1) and pericytes (visualized with NG2) in control and *VEcadCreER;Tgfbr1$^{CKO/-}$*brains at P14.
SMA immunostaining visualizes arterioles (continuous staining) and veins (patchy staining) in control and
*VEcadCreER;Tgfbr1$^{CKO/-}$*brains, and a subset of capillary-associated pericytes in *VEcadCreER;Tgfbr1$^{CKO/-}$*
brains. (C) Immune cells (CD45+ and F4-80+) in control and *VEcadCreER;Tgfbr1$^{CKO/-}$* brains at at P35.
Control brains have minimal numbers of immune cells other than resident microglia.
*VEcadCreER;Tgfbr1$^{CKO/-}$*brains show localized regions with concentrated accumulations of immune cells.
White squares marked with letters in the sagittal brain images (upper) are enlarged below. Scale bars, 1
mm for whole brain images and 200 um for enlarged images.
1212
Figure 9. snRNAseq of control and *VEcadCreER;Tgfbr1$^{CKO/-}$* vascular fragments enriched from the brain at
P14. (A) UMAP plots showing cell clusters encampassing the major cell types in the mouse brain,
enriched for ECs, pericytes, and vascular smooth muscle cells (vSMCs). The locations of the EC clusters in
the *VEcadCreER;Tgfbr1$^{CKO/-}$* and control UMAP plots (vertical red arrows) are shifted, indicating
substantial changes in their transcriptomes. Other cell cluster locations are largely unchanged. (B)
Volcano plot showing transcripts from the EC cluster in control vs. *VEcadCreER;Tgfbr1$^{CKO/-}$* snRNAseq.
The labeled transcripts have adjusted -$\log_{10}$ p-values greater than 50. (C) UMAP plots as in (A)
highlighting individual transcripts, with the EC cluster indicated by a vertical red arrow. Left column,
UMAP plots for transcripts that are up-regulated in *VEcadCreER;Tgfbr1$^{CKO/-}$* ECs. Central column, UMAP
plots for transcripts that are down-regulated in *VEcadCreER;Tgfbr1$^{CKO/-}$* ECs. Right column, *Icam1*, *Icam2*,
and *Tgfbr3*. (D) Gene set enrichment analysis (GSEA) for the EC, glial, and astrocyte clusters in (A)
showing the degree of enrichment in *VEcadCreER;Tgfbr1$^{CKO/-}$* brains. The data used to generate the
volcano plot in (B) and the GSEA in (D) are in Supplemental Table 1. NES, normalized enrichment score.
*, adjusted p-value <0.05.
1227
1228

Wang et al., Figure 1

A

Control — P30 | *VEcad-CreER;Tgfbr1^{CKO/-}* — P28

COL4

B

Control — P14, P26 | *Ndp^{KO}* — ~P30

VEcad-CreER;Tgfbr1^{CKO/-} — P14, P26 | *Fzd4^{-/-}* — ~P30

P14, P26, ~P30

PECAM1 (vitreal surface, IPL, OPL)

C

VEcad-CreER;Tgfbr2^{CKO/-}

CLDN5 PLVAP | Sulfo-NHS-biotin — P37 | CLDN5 PLVAP Sulfo-NHS-biotin

Wang et al., Figure 2

Wang et al., Figure 3

Wang et al., Figure 4

Wang et al., Figure 5

Wang et al., Figure 6

Wang et al., Figure 8

Wang et al., Figure 9