Plant host identity drives *Andropogon gerardii* rhizobiome assembly strategies under increasing abiotic stress

Brooke Vogt[1], Anna Kazarina[1], Jack Sytsma[1], Bryttan Adams[1], Leslie Rodela[1], Dasik Clouse[1], Reece Keller[1], Darcy Thompson[1], Helen Winters[1], Maggie Wagner[2], Ari Jumpponen[1], Loretta Johnson[1], and Sonny T.M. Lee[1*]

[1] Division of Biology, Kansas State University, Manhattan, Kansas, USA

[2] Department of Ecology and Evolutionary Biology and the Kansas Biological Survey, University of Kansas, Lawrence, Kansas, USA

[*] corresponding author

Abstract

Predicted changes in precipitation threaten tallgrass prairies by altering the soil

microbial communities that are essential for plant resilience. *Andropogon gerardii*, a

dominant grass in tallgrass prairies, spans the contiguous North American precipitation

gradient. However, it remains unclear to what extent the rhizosphere microbiomes

(rhizobiomes) are influenced by the plant-host environmental interaction. To assess how

environmental and host factors shape the rhizobiome, we surveyed *A. gerardii*

populations across 25 remnant prairie sites (June–August 2023) within its native range

in the United States, characterizing the microbiomes in the rhizosphere and soils using

16S amplicon sequencing. We demonstrated that while geographic location largely

structured both rhizosphere and soil communities, regional precipitation (60-day rainfall)

emerged as a primary driver of the microbial community assembly. We observed

distinct microbial divides across the dry and wet regions of the North American "arid-

humid divide." Importantly, we found the first compelling large-scale evidence that

regional precipitation has a profound influence on rhizobiome assembly. In the most arid

regions, rhizosphere microbial communities exhibited significantly more predicted

stochasticity than those in the local soil and contained taxa related to host-benefiting

functions. Our study suggests that intensified host-driven selection for specific microbial

variants occurs under heightened abiotic stress, highlighting the host's pivotal role in

shaping its rhizobiome composition in challenging environments.

Keywords
Microbiome, Microbial Community Assembly, Host-microbe Interactions, Climate
Change, 100th Meridian, Plant-microbe interactions

Introduction

Changes in the assembly and function of plant host-associated microbiomes can be influenced by a mosaic of factors, such as abiotic stress induced by climate change[1–4]. Over the past 10,000 years, the dominant tallgrass prairie grass, *Andropogon gerardii,* has evolved into genetically distinct ecotypic varieties based on precipitation (dry, mesic, and wet ecotypic varieties) across North America[5–9]. Microevolutionary processes have modified the morphological and physiological traits of these varieties, enabling them to survive and thrive in regions with varying precipitation levels, temperatures, and soil physicochemical properties[6,10–12].

Some studies suggest that plant host-associated rhizobiomes are specific to the plants' locations, indicating that plants of the same host variety and across different host varieties are distinct[13,14]. This view asserts that rhizobiome assembly is more strongly influenced by various local abiotic stresses as well as environmental and edaphic characteristics than by regional similarities in precipitation and climate. Other studies propose that rhizobiomes of the same host variety are similar across various locations, whereas those of different host varieties differ from each other[15,16]. According to this perspective, precipitation-based abiotic stress drives the host plants to select similar microorganisms, resulting in analogous rhizobiome communities within the same host variety[17,18]. These contradicting hypotheses suggest a lack of detailed information on how abiotic, precipitation-based stress affects the rhizobiomes of *A. gerardii* varieties.

Prior reciprocal garden experiments[19,20] examining the rhizobiomes of the dry, mesic, and wet varieties of *A. gerardii* have supported the idea that host-associated rhizobiomes are primarily influenced by regional precipitation. In these studies, *A. gerardii* recruited rhizobiomes similar to those at its native sites, supporting the home-field advantage hypothesis[19,20]. If microbial assembly is driven by drought stress, then different forms of this stress should lead to the host selecting for different specialists.

While the differences among dry, mesic, and wet varieties of *A. gerardii* have evolved over thousands of years due to natural selection, how quickly their host-associated rhizobiome can adapt to long-term changes in precipitation remains to be seen. The consequences of a rhizbiome's ability to adjust to different forms of precipitation-based abiotic stress are crucial. In this century, the 100th meridian ("arid

humid" divide) has shifted eastward, a trend predicted to accelerate due to the
compounding effects of climate change[5,21–24]. Understanding the host rhizobiome
assembly and its vulnerabilities is crucial, mainly because *A. gerardii* constitutes up to
80% of the biomass in tallgrass prairies[25]. In addition, central tallgrass prairies are
critically endangered, with only 4.4% remaining in North America[6,8,25–28]. Rapid shifts in
precipitation patterns may lead to a decline in *A. gerardii* if their microbiomes are closely
linked to local ecological and soil microbial communities, and cannot adapt to long-term
precipitation shifts. If *A. gerardii* microbiomes are solely influenced by local conditions,
then long-term variations in rainfall could pose a significant threat to the host's health,
longevity, and survival.

In this current study, we aimed to evaluate how regional precipitation conditions
influence rhizobiome assembly. We further investigated whether the plant host variety
has a greater influence on the associated rhizobiomes in areas experiencing more
significant precipitation-based abiotic stress. To understand the composition of the
microbial communities associated with the ecotypic varieties of *A. gerardii*, we
conducted a biogeographical survey of 25 locations across the contiguous United
States. We analyzed 16S rRNA amplicon data from local soil (LS) and plant host-
associated rhizobiome (RB) samples. Our study addressed three key questions: I) How
do local and regional characteristics influence microbial community composition? II)
How does plant host selection affect rhizobiome composition and diversity compared to
local soil, and is biotic filtering consistent across abiotic stress levels? III) How do local
soil and rhizobiome communities compare in similarity across different precipitation-
related host variety ecotypes?

Methods

Sample collection and preparation

Sample collection took place between June and August of 2023. We selected 25
sample locations based on the historical distribution of *A. gerardii*, aiming to represent
variations in precipitation[29], soil physicochemical properties, climate, and host variety
across the contiguous United States (Supplementary Table S1). We collected local soil
and rhizosphere samples from six haphazardly chosen *A. gerardii* plants at each

location. From each plant, we sampled one soil core (10 cm deep and 1.25 cm in
diameter) near the base of the plant tillers. This core sample was manually
homogenized in a sterile plastic bag. From the homogenized sample, three aliquots of
roots (approximately 20 cm in length each) were manually collected, gently shaken to
remove any loose soil, and placed into three microcentrifuge tubes. From the remaining
soil, we collected three soil aliquots (~1g each) into three microcentrifuge tubes.

Any remaining soil (without roots) from the six plant replicates that were not used
for the local soil or rhizobiome samples were stored in sterile bags on ice for the soil
physicochemical property analyses (n=25 samples). All samples were stored on ice in a
cooler until they were frozen at -80°C within 96 hours after sampling for further
processing.

Soil physicochemical properties

The soil physicochemical property analyses were conducted at Kansas State
University's Soil Testing Laboratory (https://www.agronomy.k-state.edu/ outreach-and-
services/soil-testing-lab/soil-analysis/researchers.html). The analyses were performed
on the local soil samples (n = 25). Detailed procedures are in the revised version of the
Recommended Chemical Soil Test Procedures for the North Central Region
manual[30,31]. However, a summary of the methods for each test is provided below.

Samples were analyzed for various physicochemical properties, including soil
pH, total nitrogen, carbon content (both organic and inorganic), phosphorus content,
and texture (percentages of sand, silt, and clay). Before analyses, the samples were
dried in a drying oven at 60 °C overnight. The dry soils were passed through a 2 mm
sieve to break up aggregates and ensure homogeneity.

For pH analysis, soil was mixed with an equal amount of deionized water in a 1:1
solution[30,32–34]. The mixture was stirred for 5 seconds or until a uniform soil-water slurry
was achieved. After allowing it to sit for 10 minutes, pH measurements were taken using
a Skalar SP50 Robotic Analyzer (Skalar Inc.; Georgia, USA).

Phosphorus concentrations were measured in parts per million (ppm P) using the
Mehlich III method on 2 grams of soil[30,35,36]. The soil was suspended in a 1:10 solution
with the standard Mehlich III solution, which consists of 0.2 N acetic acid ($CH_{\square}COOH$),

0.25 N ammonium nitrate (NH□NO□), 0.015 N ammonium fluoride (NH□F), 0.013 N

nitric acid (HNO□), and 0.001 N EDTA (ethylenediaminetetraacetic

acid)(HOOCCH$_2$)$_2$NCH$_2$CH$_2$N(CH$_2$COOH)$_2$). The mixture was shaken at 200 excursions

per minute for 5 minutes. After shaking, the slurry was filtered through a Whatman No.

42 filter. From the clear filtered slurry, 2 mL was transferred to a test tube, and 8 mL of

the working solution was added. The working solution was prepared by combining 25

mL of Murphy-Riley Solution A (which contains 0.051 N ammonium molybdate

[(NH□)□Mo□O□□•4H□O] and 0.004 N antimony potassium tartrate

[K□Sb□(C□H□O□)□•3H□O]) with 10 mL of Solution B (0.749 N ascorbic acid

[C□H□O□]) in a total volume of 1 L.

After letting the mixture sit for 10 minutes, the samples were loaded in triplicate

onto a 96-well plate and analyzed using a Lachat QuikChem 8500 Series 2 photometric

colorimeter (Hach Inc.; Colorado, USA) at a wavelength of 882 nm, following the

orthophosphate method (12-115-01-1-A). Phosphorus measurements were determined

by comparing the samples against a 500 ppm potassium dihydrogen phosphate

(KH□PO□) stock solution, diluted to a 50 ppm standard solution before loading into the

96-well plate.

Total nitrogen and total carbon (including organic and inorganic components)

analyses were conducted on 0.35 grams of soil using a LECO TruSpec CN analyzer

(LECO; Michigan, USA), following the procedures outlined in the operation manual[37].

The samples were encapsulated and securely sealed in the loading head. They were

then purged to eliminate any atmospheric gases that could interfere with the

measurements. Once all atmospheric gases were removed, the samples were

combusted at 950°C to ensure rapid combustion while flushed with pure oxygen. A

consistent aliquot of gases was then homogenized and separated. The total percentage

of carbon and nitrogen was calculated based on weight percentages using the formula:

Total % of Carbon or Nitrogen = (Mass of C or N / Total Sample Mass) * 100. The

analysis showed a standard deviation of 2.18% for soil carbon and 3.35% for soil

nitrogen.

Soil texture was determined using the modified hydrometer method[38,39] on a 50-

gram soil sample. First, the soil was placed in a dispersion cup, and then 100 mL of a

5% sodium hexametaphosphate solution ($Na_6(PO_3)_6$) was added. The mixture was

stirred for 30 to 60 seconds until uniform. This slurry was then transferred to a 1000 mL

cylinder filled to the 1000 mL mark with deionized water. The mixture was allowed to sit

overnight to settle. To begin the experiment, a plunger was inserted, and the sample

was mixed for an additional 30 seconds to ensure homogeneity. A Bouyoucos

hydrometer (Fisherbrand™; Soil Analysis ASTM Hydrometer) was then placed in the

slurry. After 40 seconds, the initial hydrometer reading was recorded. A second reading

was taken after 2 hours.

The percentages of sand, silt, and clay were calculated using the following

formulas: - % Clay = (Final Reading * 100 / mass of sample) - % Silt = (Initial Reading *

100 / mass of sample) - % Clay - % Sand = 100% - % Silt - % Clay All readings were

adjusted using a blank sample (100 mL of the 5% dispersion solution plus 880 mL of

deionized water) and applying temperature and density corrections. For every one °F

above 68 °F, 0.2 was added to the reading, while for every one °F below 68 °F, 0.2 was

subtracted from the reading.

DNA extraction, 16S rRNA amplicon sequencing and analyses

We extracted DNA from the samples using the E.Z.N.A.® Soil DNA Kit (Omega

BIO-TEK; Georgia, USA) following the manufacturer's protocol with minimal

modifications[40]. The only adjustment for the local soil samples was the use of glass

beads for cell lysis, performed with a Qiagen TissueLyser II (Qiagen; Hilden, Germany)

at 30 revolutions per second for 10 minutes. For each of the 25 locations, we extracted

three aliquots separately and selected one from each of the six plants based on DNA

concentration and quality. The final elution volume for each local soil sample (n = 150)

was 100 μL.

Similar to the soil samples, we extracted DNA from the rhizosphere samples

using the E.Z.N.A.® Soil DNA Kit (Omega BIO-TEK; Georgia, USA) but with significant

modifications to the manufacturer's protocol to ensure optimal DNA quality and yield.

For these samples, we combined three aliquots into one during the binding phase of the

protocol. Additionally, we used liquid nitrogen to physically macerate the roots into a fine

powder, and then lysed the tissues using the TissueLyser II for 16 minutes at 30

revolutions per second. Depending on the sample location, we adjusted the final elution
volumes for the rhizobiome samples, ranging from 30 µL to 150 µL.

We used the Qubit™ dsDNA Broad-range (BR) DNA quantification kit on a
Qubit™ 4 Fluorometer (Thermo Fisher Scientific, Invitrogen™; Massachusetts, USA) to
determine the sample eDNA concentration. The quality of the DNA was assessed by
measuring the A260/A280 and A260/A230 ratios on a Nanodrop™ One
Spectrophotometer (Thermo Fisher Scientific, Invitrogen™; Massachusetts, USA). After
quantifying the DNA, we normalized each sample to a concentration of 5 ng/µL using
molecular-grade DNA/RNA-free water.

The normalized eDNA was sequenced at the Kansas State University Integrative
Genomics Facility (https://www.k-state.edu/igenomics/). The DNA amplification was
performed using the 515F and 806R primers[41] (16S rRNA v4 amplicon region), and
sequenced with 2 x 300 cycles on the Illumina MiSeq (Illumina; California, USA). We
utilized QIIME 2 v. 2024.5[42] to process the raw reads. We applied the QIIME 2
cutadapt[43] plugin to remove the primer sequences.

Subsequently, we eliminated sequences with an error rate greater than 0.1 and
those with undetermined bases or without a primer. For further quality control, we used
DADA2[44] with consistent parameters across various runs to trim the reads, ensuring
that the 25th percentile of reads had a quality score above 15 (Forward = 270 and
Reverse = 250). To mitigate biases caused by differences in sequence depth across
samples in subsequent diversity analyses, we rarefied the dataset to 15,000 reads per
sample. We used the QIIME 2 pipeline to compute alpha (Shannon's Entropy (H'),
Faith's phylogenetic diversity (Faith PD), and number of observed features (S_{obs})) and
beta diversity metrics. Finally, we employed the pre-trained classifier in QIIME 2 for
bacterial taxonomic classification, referencing the SILVA v. 138 database[45]. After
generating the ASV table, we removed all ASVs assigned to Eukarya, chloroplasts, or
mitochondria, as well as those that remained unresolved or unclassified at the domain
level, or included non-specific reads before statistical analyses.

To methodically tease apart the influence of environment and plant host on the
rhizobiome composition, we analyzed beta dispersion to examine how community
assembly strategies influenced the distance to the centroid between dry and wet local

soil and rhizobiome samples, focusing on whether sample distribution was driven more
by stochastic (random) or deterministic processes. We defined samples where beta
dispersion distances close to 1 indicate increased stochasticity, whereas smaller
distances (0.8) demonstrate more predictability[46].

To assess specific microbial families that contribute to stochasticity, we
performed oligotyping analysis. For this analysis, we employed the QIIME 2 oligotype
function[47] (v. 3.2) and the minimum entropy decomposition (MED) algorithm[48] to identify
oligotypes within our aligned 16S rRNA amplicon sequences. We aligned our
sequences using the MUSCLE function[49]. To ensure an unbiased taxonomic
assessment of oligotypes across various sample types, we set the number of
components to 7 with a minimum substantive abundance parameter of 1 and included
all features. These parameters were determined based on the entropy analysis of
oligotype entropy peaks. We retained (LS: n = 2,248) and (RB: n = 2,876) oligotypes in
this analysis.

In our taxonomic evaluation, we extracted specific features belonging to specific
families from our previous taxonomic assessment (SILVA v. 138[45]). We filtered features
that   belonged   to   the   *Micromonosporaceae,   Nitrosphaeraceae*,
Xiphinematobacteraceae,   Vicinamibacteraceae,   Rhodanobacteraceae,   and
Pedosphaeraceae families because of their significant representation and overall
abundance across all samples. We set the number of components to 2 for the
taxonomic analysis while maintaining a minimum substantive abundance parameter of
1.   Following   the   taxonomic   oligotyping   analysis,   *Micromonosporaceae,*
Nitrosphaeraceae,   *Xiphinematobacteraceae,   Vicinamibacteraceae,*
Rhodanobacteraceae, and *Pedosphaeraceae* families yielded the following total number
of oligotypes: (LS: n = 9 and RB: n = 10; LS: n = 9 and RB: n = 10; LS: n = 14 and RB:
n = 13; LS and RB: n = 16; LS: n = 6 and RB: n = 7; and LS and RB: n = 16).

Statistical analyses and data visualization
We used RStudio version 4.4.2[50,51] for all statistical analyses and figure
generation. We defined significance for all tests as having a p-value or adjusted p-value
(used when appropriate, given the analysis) less than 0.05.

For our studies of soil/climate properties, alpha diversity, and relationships
between these factors, we utilized the car[52], dplyr[53], devtools[54], dunn.test[55],
pairwiseAdonis[56], tidyr[57], tidyverse[58], ggplot2[59], FSA[60], broom[61], rstatix[62], and vegan[63]
packages to evaluate how unique the soil's physicochemical properties were based on
the sample's location, precipitation, and sample cluster category.

We used one-way ANOVAs detect differences among the factors including soil
pH, phosphorus concentration (ppm P), carbon and nitrogen content (% total C and N),
C:N ratio, soil texture (% sand, % silt, and % clay), and precipitation (rainfall) concerning
the sample location and cluster category (dry, ambiguous, and wet). Furthermore, we
analyzed the alpha diversity metrics, sample location, cluster, and pH/precipitation
gradients using a one-way ANOVA to compare the means of the S_{obs}, Faith's PD, and
H' for the local soil and rhizobiome samples. We performed a two-way ANOVA to
analyze the relationship between our alpha diversity metrics, cluster categories, and
pH/precipitation. Additionally, we also used a one-way ANOVA to define statistical
differences among the beta dispersion means for each location and sample type.

While ANOVA identified differences among the factors, we performed Tukey
HSD post hoc tests to determine where those differences were. Additionally, we applied
the Kruskal-Wallis rank sum test to evaluate pairwise comparisons between sample
locations and clusters for both the local soil and rhizobiome samples, analyzing our soil
properties and precipitation (rainfall) data. Using both parametric and non-parametric
approaches allowed us to validate our findings across different statistical assumptions
and ensured the robustness of our results. Finally, we conducted PERMANOVA
analyses to determine whether bacterial community composition varied in relation to
location, precipitation cluster category, soil properties, and precipitation. We then
performed PERMANOVA pairwise comparisons to explore the pairs of groups
individually between the factors described in our PERMANOVA analyses. In our
analysis of how microbial diversity is impacted by climate and soil properties, we
performed both t-test and linear regression analyses to further explore these
relationships.

To assess differences in the variability of community composition within each
cluster category and evaluate category-specific dispersion patterns, we conducted

analyses of multivariate homogeneity of group dispersions (employing the betadisper

function). Using the car[52] and vegan[63] packages, we assessed the differences between

groups based on the mean distance to the centroid for each category. This evaluation

helped us determine the probability of overlap between the cluster categories. Within

our betadisper test, we performed a PERMANOVA, a permutation test, and a pairwise

permutation test to examine sample heterogeneity within each cluster category. Finally,

we used Pearson's Chi-squared tests in the tidyverse[58] and readr[64] packages to find

statistical differences in ASV composition between sample types (LS and RB) and the

cluster categories.

We used beta dispersion distances and the beta-group-significance function in

QIIME 2 to infer stochasticity within a location and between sample types, thereby

surmising the underlying strategies of community assembly. For the oligotyping

analysis, we employed the beta dispersion tests previously described.

We utilized the RStudio packages ggplot2[59], patchwork[65], dplyr[53], multcomp[66],

multcompView[67], agricolae[68], tidyr[57], venn[69], tidyverse[58], vegan[63], cluster[70], sf[71], maps[72],

reshape2[73], pheatmap[74], RColorBrewer[75], car[52], gridExtra[76], phyloseq[77], igraph[78],

ggnewscale[79], readr[64], ggthemes[80], ggtext[81], ggVennDiagram[82], randomcoloR[83],

stringr[84], and VennDiagram[85] to visualize the analyses performed in this study. All raw

sequence data for this study are publicly available in the NCBI SRA under BioProject

accession number PRJNA1249957. All bioinformatic and R codes are deposited in

GitHub   (https://github.com/SonnyTMLee/Plant-host-identity-drives-rhizobiome-

assembly-under-increasing-abiotic-stress.git).

Results and Discussion

We processed a total of 13,087,481 (local soil) and 16,677,859 (rhizobiome) raw

reads. After quality control, we retained 11,759,436 reads (2,462 ASVs) for the local soil

and 14,627,083 reads (3,039 ASVs) for the rhizobiome (Supplementary Table S2). In

the local soil samples, 92% (2,267 ASVs) of the ASVs resolved to the genus level, and

64% (1,567 ASVs) resolved to the species level. We observed similar resolution across

the rhizobiome samples, where 93% (2,815 ASVs) resolved to the genus and 65%

(1,985 ASVs) resolved to the species level. The five most abundant phyla in the local

soil samples were Proteobacteria (22%, 532 ASVs), Actinobacteriota (11%, 278 ASVs), Bacteroidota (9.8%, 242 ASVs), Chloroflexi (8.1%, 200 ASVs), and Acidobacteriota (7.6%, 187 ASVs) (Supplementary Table S3). We observed similar dominant phyla across the rhizobiome samples, with Proteobacteria (23%, 691 ASVs) being the most dominant, followed by Actinobacteriota (10%, 315 ASVs), Bacteroidota (9.9%, 302 ASVs), Planctomycetota (7.5%, 229 ASVs), and Chloroflexi (7.5%, 228 ASVs) (Supplementary Table S3). Of all our samples, 0.12% (3 ASVs) from the local soil and 0.13% (4 ASVs) from the rhizobiome were unclassified.

Regional precipitation shaped both local soil and plant host-associated microbial communities

We used non-metric multidimensional scaling (NMDS) and optimal silhouette scores from k-means clustering, and identified two distinct clusters among both the local soil (silhouette score = 0.21) and rhizobiome (silhouette score = 0.22) samples (Figure 1A, 1B). We observed that regional precipitation significantly influenced the two identified clusters - "wet-categorized" and "dry-categorized". Samples in the transition zone (overlap) between the wet and dry were classified as "ambiguously-categorized." The ten samples organized as dry-categorized local soil (DC_{LS}) were: MT1, SD1, ND1, MN1, CO1, KS2, NM1, NE2, OK1, and TX3. Three samples (KS1, NE1, and IA1) belonged to the ambiguously categorized local soil (AC_{LS}), and 12 samples (WI1, MI1, IN1, MO1, IL1, TX1, TX2, LA1, AL1, NC1, and SC1) were organized as wet-categorized local soil (WC_{LS}). The dry-categorized local soil samples had the same assignments as the dry-categorized rhizobiome samples (DC_{RB}). However, the sample assignments exhibited slight changes from the local soil for the ambiguously- (AC_{RB}) and wet- (WC_{RB}) categorized rhizobiomes, in which IA1 and NE1 were reclassified into the WC_{RB} category. We surmised that the dry-categorized samples had more stable sample assignments than the wet or ambiguous-categorized samples due to long-lasting relationships between the dry variety of *A. gerardii* and its associated rhizobiome[6,19,86]. In contrast, the wet and ambiguously categorized clusters may be less tied to their respective host varieties, possibly due to the shifting of historical precipitation boundaries, bringing aridity to historically moist locations for centuries[24,87,88]. This

change was most striking as a result of the acceleration of eastward shifts in regional

precipitation zones encompassing the 100th meridian, with predictive models indicating

that these transitions are likely to accelerate within this century[24].

To provide deeper insights into the impact of regional precipitation on the local

soil and rhizobiome composition, we used the NOAA[29] (v. 2) database for precipitation

(rainfall) data, selected the maximum precipitation over 60 days, relative to the annual

rainfall at the sample locations, and mapped the distribution of our clustering analysis

across the precipitation gradient (Figure 1C). We observed a clear divide between the

dry and wet categories, separating distinctly along the 100th meridian, often referred to

as the "arid-humid divide," with the ambiguous samples comprising the mesic transition

zone. The clustering of our samples across the 100th meridian suggests that

precipitation influences the microbiome composition across geographically diverse

locations[89]. This indicates that the selective pressures exerted by precipitation may

surpass the effects of local climate, geographic, and soil property differences[90,91].

Our one-way ANOVA showed differences among the categories and their

precipitation depth (LS: $p < 2.2 * 10^{-16}$ and RB: $p < 2.2 * 10^{-16}$; Precipitation: DC_{LS} =

221.5 ± 46.2 mm and $DC_{RB} = 224.3 \pm 45.7$ mm, $AC_{LS} = 312.4 \pm 15.0$ mm and AC_{RB} =

325.1 ± 0 mm, and $WC_{LS} = 360.9 \pm 55.9$ mm, and $WC_{RB} = 350.5 \pm 54.864$ mm).

Kruscal-Wallis post hoc pairwise tests showed that for the local soil, the precipitation

depth between the DC_{LS} and the WC_{LS} was most significant (p-adj = 3.92×10^{-20}),

followed by the DC_{LS} vs. AC_{LS} (p-adj = 3.34×10^{-5}) and WC_{LS} vs. AC_{LS} (p-adj = $1.02 \times$

10^{-1}). This trend was also observed for the rhizobiome samples DC_{RB} vs. WC_{RB} (p-adj =

2.08×10^{-9}), DC_{RB} vs. MC_{RB} (p-adj = 4.08×10^{-2}), and WC_{RB} vs. AC_{RB} (p-adj = 1.00).

We further used PERMANOVA and showed a significant impact of rainfall on

local soil and rhizobiome community structures (p = 0.001). However, though

PERMANOVA pairwise analysis between our categories showed significantly different

bacterial composition between all of the local soil category comparisons (adj p-values =

0.003), comparisons between WC_{RB} and $AC_{RB,}$ as well as DC_{RB} and $AC_{RB,}$ were

insignificant (adj p-values = 0.16 and 0.07, respectively). Our results corroborated with

those of another study[20], showing that regional precipitation influences local soil

communities more than plant host-associated rhizobiome communities. Here, we

demonstrated the influence and regulation of the plant host on its associated rhizobiome

community structure in response to adverse hydrologic conditions[1,92,93]. Next, we

wanted to ask if and to what extent physicochemical soil properties contributed to this

variation.

Soil physicochemical properties and bacterial diversity were primarily influenced

by geographic location rather than regional precipitation

We observed that the soil physicochemical properties (Supplementary Table S4)

were more closely related to individual locations than when we performed analysis

based on grouping the locations by our cluster categories: wet, ambiguous, and dry

(Figure 2A). All soil properties were statistically significant between the locations

(Supplementary Table S5). Nevertheless, only soil pH (LS: $p = 1.05 \times 10^{-12}$), sand

content (LS: $p = 1.02 \times 10^{-2}$), and silt content (LS: $p = 1.90 \times 10^{-4}$) were significantly

different across our cluster categories. We postulate that the lack of significant

differences within our community-defined clusters arose because the extreme variability

of individual local soil properties in specific locations was mitigated, resulting in

comparable soil characteristics over extensive regions[94].

Our results suggested that local soil communities were significantly influenced by

soil pH (ANOVA: $DC_{LS} = 7.00 \pm 0.74$, $AC_{LS} = 7.27 \pm 0.71$, $WC_{LS} = 6.17 \pm 0.64$; Kruskal-

Wallis: DC_{LS} vs. WC_{LS} ($p = 5.40 \times 10^{-8}$), the DC_{LS} vs. AC_{LS} ($p = 0.02$), and the WC_{LS} vs.

AC_{LS} ($p = 1.60 \times 10^{-5}$), whereas in the rhizobiome, root exudates from the host plant

could be acting as buffers to maintain pH homeostasis, thus mitigating these

differences[95]. Although, in the rhizobiome, the only significant difference from our

PERMANOVA pairwise analysis was observed between the DC_{RB} and WC_{RB}, ($p =$

0.003), suggesting that increased rainfall in the wetter regions could have led to the

leaching of basic cations from the soil, resulting in a lower pH; while soils with limited

rainfall retained these cations, making them more alkaline[96].

Our findings further indicated that the ratio (R:pH) between rainfall (R) and soil

pH was significant in the local soil and rhizobiome across the cluster categories ($p =$

0.001). The rainfall and pH ratio were significant for all local soil categories ($p = 0.003$)

for all interactions (Supplementary Table S5). However, for the rhizobiome, only the

interaction between the DC_{RB} and WC_{RB} was significant ($p = 0.003$). The relationships

intertwining soil pH, rainfall, and bacterial diversity are complicated. While pH and

precipitation emerged as primary drivers of change in bacterial diversity[97,98], the extent

to which these two variables directly influenced bacterial community composition

remains unclear. However, common trends regarding pH and precipitation impacts on

specific microbial alpha diversity indices have been well documented[89,90,98,99], leading

us to investigate these relationships.

We observed that, across the 25 locations, alpha diversity metrics

(Supplementary Table S6) were significantly tied to locations (ANOVA: (LS: S_{obs} p =

1.06×10^{-8}, Faith PD p = 4.91×10^{-10}, and H' p = 7.38×10^{-10}) and (RB: S_{obs} p = $6.56 \times$

10^{-6}, Faith p = 4.54×10^{-8}, and H' p = 1.98×10^{-5}); Figure 2B). On the other hand, while

local soil alpha diversity metrics significantly differed across the local soil cluster

categories (ANOVA: (LS: S_{obs} p = 6.02×10^{-5}, Faith PD = 1.55×10^{-3}, and H' p = $8.75 \times$

10^{-6})); there was no significance between the rhizobiome samples possibly due to

shielding from the host (Supplementary Table S5).

We examined how pH and precipitation gradients influenced the organization of

our cluster categories in terms of microbial diversity and abundance (Supplementary

Figure S1). We found that precipitation depth significantly affected all alpha diversity

metrics across (ANOVA: LS and RB (S_{obs} p = 0.05, Faith PD p = $3.82 * 10^{-3}$, H' p = 7.17

$* 10^{-3}$) and between the sample types ((LS vs. RB) ($p < 2 * 10^{-16}$)). Similar results were

observed in our linear regression analysis comparing the alpha diversity metrics of

rhizobiomes, ordered by varying levels of rainfall (Supplementary Figure S2).

Pairwise tests further revealed that the most dramatic differences in diversity

were observed between the following precipitation (as 60 day-maximum rainfall)

categories across sample types: (H': (LS: p-adj = $2.21 * 10^{-3}$ (9 to pH 9.5 vs. 12.5 to pH

13), RB: p-adj = $3.21 * 10^{-3}$ (6.5 to 7 vs. 15 to 15.5)); Sobs: (LS: p-adj = $2.05 * 10^{-3}$ (9 to

9.5 vs. 12.5 to 13), RB: p-adj = $3.99 * 10^{-3}$ (6.5 to 7 vs. 16.5 to 17)); and Faith PD: (LS:

p-adj = $2.21 * 10^{-3}$ (13 to 13.5 vs. 16.5 to 17), RB: p-adj = $6.05 * 10^{-4}$ (14 to 14.5 vs. 16.5

to 17)) (Figure 2C; Supplementary Table S7). While the alpha diversity metrics in our

study were more stable across categories of increased rainfall[100], the local soil revealed

the most substantial differences in alpha diversity, mainly among moderate precipitation

categories. At the same time, the rhizobiome exhibited more variable changes,
particularly between the extreme precipitation groups. This suggests that soil microbial
communities were more responsive to subtle deviations in rainfall within our "ambiguous
buffer zone" separating arid and humid areas[100,101]. However, because of the mitigating
effects of the host, the rhizobiome appeared to be more resistant to intermediate
precipitation fluctuations, only showing differences among precipitation extremes[102].

Linear regression analysis revealed significant changes of relative abundance
across the precipitation gradient for seven phyla: *Crenarchaeota*, *Desulfobacterota*,
MBNT15, *Planctomycetota*, *RCP2-54*, *Verrucomicrobiota*, and *WPS-2* with p-values of p
$= 4.00 * 10^{-2}$, $p = 1.00 * 10^{-2}$, $p = 9.41 * 10^{-3}$, $p = 2.11 * 10^{-2}$, $p = 1.09 * 10^{-2}$, $p = 3.46 *$
10^{-4}, and $p = 9.64 * 10^{-4}$ respectively (Supplementary Figure S3). All except
Crenarchaeota (slope $= -1.22 * 10^{-4}$; $R^2 = 0.306$) increased in relative abundance with
greater precipitation depth. We hypothesize that the decrease in *Crenarchaeota* might
be due to their roles as ammonia-oxidizing archaea (AOA), which might not be in high
demand   or   feasible   under   waterlogged   conditions[103].

In contrast, no significant changes in alpha diversity (ANOVA: (S_{obs} p $= 0.73$,
Faith PD $= 0.59$, H' p $= 0.49$)) or bacterial relative abundance were found across the pH
gradient; with significant changes only observed when comparing sample types (p $< 2 *$
10^{-16}) and categorized cluster regarding Faith PD (p $= 2.08 * 10^{-3}$) (Figure 2D).
However, two-way ANOVA revealed significance regarding the relationship between
Shannon diversity and pH*cluster category (p $= 1.54 * 10^{-2}$). As with rainfall we
conducted a linear regression analysis to compare the alpha diversity metrics of
rhizobiomes and local soils, organized by decreasing pH (Supplementary Figure S4).

Pairwise analysis indicated no significant differences in alpha diversity metrics
among rhizobiome categories and only minimal differences across local soil pH
categories, suggesting that while community composition may be influenced, overall
richness remains relatively stable (Supplementary Table S7). The minimal variation in
microbial diversity across pH subdivisions suggests that legacy effects of the host plant
variety may exert a stronger influence on current plant-microbe associations[10,104,105].
This interpretation is further supported by findings from our previous reciprocal garden

experiment[19], in which host wet varieties exhibited reduced richness (S_{Obs}) in dry soil
compared to ambiguous and dry host varieties grown under the same conditions.

Collaborating with our previous findings that precipitation is the most critical
factor in driving differences in host-associated rhizobiome composition under "abiotic
extremes," led us to ask the following question: How does the interaction between
precipitation, sample type, and sample category assignment influence the distribution of
ASVs among our samples?

Prokaryotic community composition revealed patterns that differentiated our
clustering categories

We observed that the local soil and rhizobiome samples had similar microbial
community composition trends across and within our cluster categories (Figure 3A).
Interestingly, we observed that the dry-categorized samples had community
compositions more similar to each other than the wet-categorized samples, suggesting
that increased abiotic stress might have a filtering effect on the microbial community
structures[106,107].

We found that beta diversity dispersion significantly differed among clustering
categories for both local soil and rhizobiome communities (PERMANOVA, p = 0.001; F-
statistic LS: 28.4, RB: 24.9), with BetaDisper distances to the median exhibiting
inconsistent levels of within-group variability (DC_{LS} = 0.33, DC_{RB} = 0.30; AC_{LS} = 0.21,
AC_{RB} = 0.20; WC_{LS} = 0.33, WC_{RB} = 0.34). Furthermore pairwise comparisons showed
significant differences in betadispersion between all of the categories (DC_{LS} vs. MC_{LS}: p
= 1.57 × 10^{-13}, AC_{LS} vs. WC_{LS}: p = 1.35 × 10^{-9}; DC_{RB} vs. MC_{RB}: p = 1.75 × 10^{-7}, AC_{RB} vs.
WC_{RB}: p = 4.79 × 10^{-6}, and DC_{RB} vs. WC_{RB}: p = 1.00 × 10^{-3}), except for one interaction
(DC_{LS} vs. WC_{LS}: p = 0.91). These results support our previous findings that the microbial
communities within each cluster can vary significantly, with some cluster categories
exhibiting more pronounced heterogeneity (dry and wet) than others ambiguous [19].

Although the dry and the wet cluster categories did not differ significantly in terms
of ASV composition comparing sample types, the following scenarios prompted us to
investigate differences in cluster categorized microbial composition: I) We noticed that
the dry rhizobiome samples are more similar to each other than dry local soil; II)

additionally, the dry rhizobiome composition is very different from the wet rhizobiome

(Figure 3B). While the shared ASVs across all cluster categories (41.2% LS and 34.4%

RB) represented the highest relative abundance, the wet-categorized cluster had the

second-highest number of unique ASVs (20.1% LS and 23.8% RB), followed by the dry-

categorized cluster (15.3% LS and 11.9% RB) (Figure 3C). Across the sample types

(LS and RB), we observed that 26.5% of ASVs were shared among all categories and

samples (Figure 3D, Supplementary Table S8). The wet-categorized cluster had more

unique ASVs than the dry-categorized cluster. We deduce that this difference could be

linked to various microbial legacy effects and adaptation to abiotic stress[108,109]. In this

context, the dry-categorized cluster might prefer specialist microbes over generalists,

reducing the overall number of unique ASVs[110,111]. Correspondingly, the small number

of unique ASVs between clusters and sample types might result from phylogenetic

divergence and variations in niche space, where microbes adapted to specific

precipitation pressures may not suit contrasting environments[112]. We further confirmed

that there were significant differences in unique ASVs across cluster categories (Chi-

squared, $p = 2.2 \times 10^{-16}$), and between wet and dry categories ($p = 0.05$), demonstrating

that differences in unique ASVs were primarily due to cluster category, rather than

sample type ($p = 0.557$). Our results suggest that complex environmental and climate

factors have a more profound impact on shaping the number of unique ASVs than the

localized effects of individual plant hosts.

Stochastic processes influenced community composition more in the local soil

than in the rhizobiome in arid locations

Both local soil and rhizobiome samples showed high beta dispersion distances

across all of our dry and wet locations and cluster categories, common among

grassland microbiomes (Figure 4A, 4B; Supplementary Table S9)[113]. We observed

notable differences between the dry cluster categorized local soil and rhizobiome

communities, in which 8 out of 10 comparisons showed significant differences in beta

dispersion (Dry: TX3: $p = 3.43 \times 10^{-5}$, NE1: $p = 5.41 \times 10^{-6}$, OK1: $p = 1.45 \times 10^{-11}$, KS2:

$p = 3.08 \times 10^{-2}$, ND1: $p = 1.20 \times 10^{-3}$, MT1: $p = 6.02 \times 10^{-3}$, SD1: $p = 5.06 \times 10^{-3}$, NM1:

p = 1.52 × 10^{-3}, and CO1: p = 8.69 × 10^{-4}), whereas only one comparison for wet-

categorized clusters was significant (Wet: MI1: p = 3.02 × 10^{-2}).

Notably, among the two driest locations, CO1 and NM1, the local soil had

significantly higher beta dispersion than their rhizobiome communities, a trend unique to

the dry-categorized clusters. We hypothesize that dispersal limitation and host influence

on microhabitat stability and selection might drive the observed differences in

stochasticity among various sample types in the driest locations. This could result from

dry clusters reducing dispersal effects by fostering specialized microbial communities

shaped by abiotic stress, leading to a more stable rhizobiome community structure than

the local soil[114]. With the dry-associated cluster category rhizobiome demonstrating less

stochasticity, we next asked what microbial populations in the rhizobiome could

contribute to the similarity and stability associated with the dry variety plant hosts.

Oligotyping analysis reveals contrasting variant compositions across cluster

categories and sample types for six prokaryotic families

To analyze fine-scale ecological patterns in local soil and rhizobiome samples

exclusively across wet and dry cluster categories, we conducted an unbiased taxonomic

oligotype assessment for all our ASVs. Our clustering for oligotyping analysis mirrored

the clustering of our ASVs. This suggests that the impact of rare oligotypes within each

taxon from each location was limited, while precipitation and ecological pressures drove

common trends between ASVs and their variants at each location (Supplementary

Figure S5, Supplementary Table S10).

For both sample types, the permutation test for homogeneity revealed significant

differences in betadispersion between the cluster categories for each sample type (LS

and RB: p = 0.001, LS: F = 10.9, RB: F = 14.2). However, we only observed differences

between the dry and wet categorized clusters within the rhizobiome ((LS distance to

centroid: 0.24 and 0.26, p = 0.13) and (RB distance to centroid: 0.21 and 0.26, p = 4.25

x 10^{-6}) for dry and wet, respectively) (Figure 5A). The distinctions between dry and wet

rhizobiomes could originate from more substantial selection pressures and reduced

microbial diversity at our drier locations. Furthermore, the contrast between dry local soil

and rhizobiome might be more pronounced due to host selection[115], microhabitat

stabilization[116], and environmental filtering[110].

Given the differences between the dry and wet categorized clusters and across

the two sample types, we aimed to evaluate the oligotypic patterns within specific ASV

families to examine the abundance of particular community oligotypes among closely

related species of microorganisms (Figure 5B and 5C; Supplementary Table S11).

Taxonomic analysis revealed distinct results across the categorized clusters and

sample types among the *Micromonosporaceae, Nitrosphaeraceae*, and

Xiphinematobacteraceae versus the *Vicinamibacteraceae, Rhodanobacteraceae,* and

Pedosphaeraceae families. For the *Micromonosporaceae, Nitrosphaeraceae*, and

Xiphinematobacteraceae families, the highest relatively abundant oligotypes in the local

soil differed significantly from those in the rhizobiome, with very few "core" variants

between sample types (Figure 5B). The similarities between the dry and wet rhizobiome

oligotypes compared to the local soil suggested that host selection might favor specific

oligotypes over others[117], regardless of the location and variant composition of the local

soil for members of the *Micromonosporaceae* family. This observed host influence might

stem from notably, that many species in the *Micromonosporaceae* family are frequently

associated with plant rhizospheres due to their production of plant growth regulators,

including auxins and gibberellins[118,119]. Additionally, the *Nitrosphaeraceae* family

constitutes ammonia-oxidizing archaea (AOA), which may benefit host nutrient

acquisition[120]. The fact that both of these families had similar oligotypic distribution

patterns supported the "stress-induced selection" hypothesis, where, under abiotic

stress, plants would likely retain the microorganisms within the rhizobiome that benefit

the plant[121,122].

However, in the *Vicinamibacteraceae, Rhodanobacteraceae,* and

Pedosphaeraceae families, there was greater similarity of oligotypes across the

cateorgized clusters and sample types, indicating a more extensive variety of "core"

variants (Figure 5C). While more shared oligotypes existed, each location was more

distinct than observed in the *Micromonosporaceae* family. We speculated that these

differences arose because members of the *Pedosphaeraceae* family are more "free-

living" than those associated with the *Micromonosporaceae* family[123]. Because of this

minimized host selection, its local soil and rhizobiome oligotype composition were more

similar within a location, meaning that the oligotypes found in the rhizobiome most likely

originated from the local soil, with little to no host filtering[116,117].

Despite these apparent distinctions among these families, common trends

emerged. Generally, the wet local soil and rhizobiome contained more oligotypes than

the dry local soil and rhizobiome locations, reflecting the filtering effect from both the

environment and plant host[124,125]. Additionally, the rhizobiome consistently had more

oligotypes than the local soil, suggesting that the plant host could support a higher

diversity of microorganisms within a family due to increased nutrients and reduced

fluctuating stress associated with the environment. Additionally, we theorized that

microhabitat stability and the increased number of features present for the rhizobiome in

both   families   (*Micromonosporaceae*:   LS: n   =   329   and   RB: n   =   587;

Xiphinematobacteraceae: LS: n = 251 and RB: n = 305; *Nitrosphaeraceae:* LS: n = 196

and RB: n = 205; *Vicinamibacteraceae:* LS: n = 758 and RB: n = 932;

Rhodanobacteraceae: LS: n = 136 and RB: n = 318; *Pedosphaeraceae*: LS: n = 755

and RB: n = 1,083) might have contributed to the rhizobiome samples having more

variants regardless of cluster assignment[126]. Considering the oligotypic analysis here,

we suggest that as the abiotic stress of the environment increases, the host influence

on the rhizobiome community exerts a greater pressure.

Conclusion

Precipitation   significantly   shapes   the   compositions   of   microbiome

communities[127]. However, discrepancies between precipitation and the strength of host

interaction can lead to varying impacts on community structure, some of which can be

exacerbated under increased abiotic stress[128,129]. Here, we demonstrated that regional

precipitation profoundly affected and categorized the microbiomes of *A. gerardii* into two

distinct cluster categories (i.e., dry, wet) and one ambiguous cluster category that

comprised the intersection between the dry and the wet clusters. Notably, the

geographic span of our categorized clusters crossed distinguishable areas of the 100[th]

meridian, known as North America's "arid-humid divide." We speculate that our

categorized clusters' composition and geographic span might be linked to the *A. gerardii*
host varieties' diversity and legacy effects[9,12,130].

Additionally, our findings showed that although pH and precipitation depth had a
more pronounced effect on local soil than on rhizobiome communities, alterations in
rhizobiome diversity were more pronounced at precipitation extremes. We surmise that
these differences might arise from host variety (ecotypic) effects through microhabitat
stabilization. By stratifying our samples by type and respective clusters, we further
identified large-scale trends in microbial community composition, particularly in the
differences between the dry and wet categorized clusters. In this study, we observed
significant variations in the relative abundances of specific phyla within sample-type
clusters. This suggests specialized roles of specific phyla responding to precipitation-
related stress and underscores potential precipitation-community structure relationships.

Furthermore, for the first time, we showed notable differences in rhizobiome and
local soil microbial diversity with reduced precipitation, revealing possible links between
abiotic stress and microbial community assembly strategies (stochastic vs.
deterministic). We further demonstrated that regional precipitation was a critical factor in
rhizobiome assembly, with the rhizosphere communities in the aridest locations
exhibiting more predicted forms of randomness than the local soil, implying plant host-
associated biotic filtering under extreme abiotic stress (Figure 6). Our findings highlight
the shift toward reduced stochasticity in the microbial communities under intensified
abiotic stress. This underscores the pivotal role of regional precipitation in shaping
plant-associated rhizobiomes and suggests that plant hosts might rely heavily on biotic
filtering mechanisms to mitigate extreme environmental challenges.

Funding
This study is supported by the United States Department of Agriculture, National Institute of Food and Agriculture (USDA NIFA), project award (Number 2020-67019-3180) and the National Science Foundation CAREER Award (Number 2238633) as well as by the National Science Foundation Award (Number OIA-1656006) with matching support from the State of Kansas through the Kansas Board of Regents.

Acknowledgments
We thank everyone involved in sampling and data acquisition; this research would not have been possible without your contributions. We also thank Tanner Riche and Qinghong Ran for their helpful discussions on data analysis. We appreciate the work conducted by Alina Akhunova, Jose Ruiz Ortiz, and Samantha Elledge at the Integrated Genomic Facility at Kansas State University (https://www.k-state.edu/igenomics/). In addition, we thank the Kansas State University Soil Testing Laboratory for helping with the soil property analyses (https://www.agronomy.k-state.edu/outreach-and-services/soil-testing-lab/).
We acknowledge that this research was conducted on land that is the homeland of many Native Nations, who have long served as stewards of North American tallgrass prairies. We honor their histories, cultures, and traditions and recognize their contributions to the scientific, ecological, and historical knowledge of tallgrass prairie ecosystems.

Conflicts of Interest
The authors stated no conflicts of interest.

Author Contributions
A.K., J.S., D.C., H.W., M.W., S.B., A.J., L.J., and S.T.M.L. collected and processed samples. B.V., B.A., L.R., R.K., D.C., and D.T. extracted, quantified, and normalized genomic DNA. B.V. and A.K. conducted the bioinformatic analysis of 16S rRNA sequencing. B.V. carried out the statistical analyses. B.V. and S.T.M.L. authored the manuscript, developed figures, and prepared all supplementary files. A.J. and L.J. assisted with revisions of the manuscript and figures. S.T.M.L., A.J., L.J., M.W. and S.B. secured funding for this study. All authors reviewed, edited, and endorsed the last version of this manuscript.

Author's Information
Sonny T.M. Lee:
    X: @sonnytmlee
    Bluesky: @sonnytmlee.bsky.social
    LinkedIn: https://www.linkedin.com/in/sonny-tm-lee-a25b9120
    ResearchGate: https://www.researchgate.net/profile/Sonny-Lee-3
Brooke Vogt:
    X: @VogtBrookeM
    Bluesky: @brookevogt.bsky.social
    LinkedIn: www.linkedin.com/in/brooke-vogt-5a351035b

680   ResearchGate: https://www.researchgate.net/profile/Brooke-Vogt
681

Figures

Figure 1: *Precipitation-associated sample clustering and geospatial analysis*

(A) NMDS clustering of local soil samples with k-means clustering and optimized silhouette scores showing distinct groupings with 95% confidence ellipses. Each point represents a sample, colored according to NOAA's 60-day maximum precipitation data at the collection site.

(B) NMDS clustering of rhizobiome samples with the same clustering criteria. Sample points are uniquely marked and colored by precipitation (rainfall), consistent with part A.

(C) Map of biogeographical survey sites, with markers colored by NOAA's 60-day maximum precipitation (circle) values. Stacked bar graphs within the map display the proportion of local soil samples assigned to each NMDS-defined group at each site. The adjacent pie charts summarize group assignments for rhizobiome samples.

Figure 2: *Cluster soil physicochemical properties and alpha diversity across cluster categories and soil property gradients*

(A) Violin plots showing the distribution of soil physicochemical properties across assigned local soil cluster categories. Statistical differences among cluster categories were assessed using ANOVA with Tukey's HSD post hoc tests. Red and blue panels represent dry and wet cluster categories, respectively.

(B) Alpha diversity metrics, including observed features, Shannon entropy, and Faith's PD, are grouped by cluster categories for local soil and rhizobiome samples. Significance was evaluated in the same manner as in part A. Red and blue panels represent dry and wet clusters, respectively.

(C) Comparison of alpha diversity metrics between local soil and rhizobiome samples across environmental precipitation (rainfall) categories (all samples used) (assessed using independent t-tests. Statistical significance is indicated as follows: ns (grey) = $p > 0.05$, * (orange) $p < 0.05$, ** (red) $p < 0.01$, *** (brown) $p < 0.001$.

(D) Comparison of alpha diversity metrics between local soil and rhizobiome samples across pH categories (all samples used), also assessed in the same manner as in part C.

Statistical significance is indicated as follows: ns (grey) = $p > 0.05$, * (orange) $p < 0.05$, ** (red) $p < 0.01$, *** (brown) $p < 0.001$.

Figure 3: *Beta diversity patterns and ASV distribution across cluster categories and sample types*

(A) Bray–Curtis dissimilarity heatmaps illustrating large-scale community composition trends in (A) local soil and (B) rhizobiome samples, clustered by their cluster assignments (dry and wet).

(B) PCoA plots of beta dispersion, showing community variability within cluster category (dry and wet) for each sample type. The accompanying box plots display the distance of each sample to the group centroid.

Statistical significance is defined as follows: ns = p > 0.05, * p < 0.05, ** p < 0.01,
*** p < 0.001.
(C) Venn diagrams depicting the distribution of shared and unique ASVs across
dry and wet cluster categories for local soil and rhizobiome samples.
(D) Combined Venn diagram comparing ASV overlap between local soil and
rhizobiome sample types across dry and wet cluster categories.
733
Figure 4: *Within-site community variability across local soil and rhizobiome
samples*
(A) Beta dispersion analysis of local soil samples, assessing within-site variability
across locations. Sample points are colored by their cluster category, with lines
representing the mean distance to the centroid for each category and shaded
bars indicating standard deviation. Sites are arranged in order of decreasing
precipitation.
(B) Beta dispersion analysis of rhizobiome samples, with sample coloring and
site arrangement as described in part A.
Statistical comparisons between local soil and rhizobiome samples at each site
indicate significant differences in beta dispersion, suggesting variation in
community stochasticity across sample types.
Statistical significance is denoted as follows: ns = p > 0.05, * p < 0.05, ** p <
0.01, *** p < 0.001.
748
Figure 5: *Community variant structure and oligotype composition across cluster
category and sample types*
(A) PCoA plots of beta dispersion based on taxonomically unbiased oligotyping
analysis, illustrating community variability within cluster categories for each
sample type. Accompanying box plots display the distance of individual samples
to their group centroid, describing within-group variability. Statistical significance
is denoted as follows: ns = p > 0.05, * p < 0.05, ** p < 0.01, *** p < 0.001.
(B) The top panels represent host-selected communities. Stacked bar plots
showing the relative abundance (y-axis) of individual oligotypes within selected
prokaryotic families, stratified by sample type and dry versus wet-categorized
locations (x-axis). Each color corresponds to a distinct oligotype variant. Details
of the oligotypes are in Supplementary Tables 10 and 11.
(C) Oligotypes of microbial populations hypothesized to be non-host-selected
communities. Stacked bar plots showing the relative abundance (y-axis) of
individual oligotypes within selected prokaryotic families, stratified by sample type
and dry versus wet-categorized locations (x-axis). Each color corresponds to a
distinct oligotype variant. Details of the oligotypes are in Supplementary Tables
10 and 11.
767
Figure 6: *Conceptual model summarizing the influence of precipitation-driven
abiotic stress on soil and rhizobiome community assembly*
The top panel illustrates a subdivision of the United States precipitation gradient,
highlighting two locations, A and B, to describe two scenarios based on our
findings.

The second panel (Location A) illustrates a high-stress environment with limited
precipitation. In this scenario, plant host selection has a strong influence on
rhizobiome community assembly, leading to divergence from the surrounding soil
community and the enrichment of specialist taxa. This shift reflects a reduction in
environmental influence and an increase in host-mediated filtering under abiotic
stress. Here, the stochasticity of the rhizobiome becomes more uniform, with
more similar plant-associated rhizobiome compositions, as site aridity increases
while the local soil remains highly variable.
In contrast, the bottom panel (Location B) depicts a low-stress environment
characterized by high precipitation. Under these conditions, environmental
influences dominate, while the impact of plant hosts is minimal. Consequently,
local soil and rhizobiome communities exhibit high variability. Generalist taxa
dominate both communities. Here, the stochasticity of the local soil and
rhizobiome remains high, showing a highly variable plant-associated rhizobiome,
and precipitation has a minimal effect.
788
789
790

Supplementary Figures

Supplementary Figure S1: *Microbial community composition across climate and soil property gradients*
Stacked bar plots showing the relative abundance of ASVs classified at the phylum level, stratified by cluster category, environmental pH categories (top), and precipitation (bottom) gradients. This visualization highlights compositional shifts in microbial communities along gradients of climate and soil properties.

Supplementary Figure S2: *Precipitation linear regression analysis on alpha diversity*
Linear regression analysis was conducted to evaluate the relationship between precipitation and alpha diversity metrics—including S_{obs}, H', and Faith PD—in local soil (left panels) and rhizobiome (right panels) communities. Samples were arranged along a gradient of decreasing precipitation and visualized using violin plots. The plot color denotes the location's cluster category (dry and wet). Displayed in each of the plots are the p and R^2 values from the regression analysis.

Supplementary Figure S3: *Linear regression analysis of phylum-level microbial abundance across categories and sample types with increasing precipitation depth*
Linear models assessed phyla whose relative abundances significantly correlated with precipitation across different categories and sample types (local soil and rhizobiome). Only phyla exhibiting significant relationships with precipitation ($p < 0.05$) are shown, highlighting taxa potentially responsive to moisture variability along the precipitation gradient.

Supplementary Figure S4: *pH linear regression analysis on alpha diversity*
Linear regression analysis was conducted to evaluate the relationship between pH and alpha diversity metrics—including S_{obs}, H', and Faith PD—in local soil (left panels) and rhizobiome (right panels) communities. Samples were arranged along a gradient of decreasing pH and visualized using violin plots. The plot color denotes the location's cluster category (dry and wet). Displayed in each of the plots are the p and R^2 values from the regression analysis.

Supplementary Figure S5: *Taxonomically unbiased oligotype clustering*
NMDS taxonomically unbiased oligotype clustering of local soil (top) and rhizobiome (bottom) variants within each sample with k-means clustering and optimized silhouette scores showing distinct groupings with 95% confidence ellipses. Each point represents a sample, colored according to NOAA's 60-day maximum precipitation data at each site.

837
838 **Supplementary Tables**
839
840 Supplementary Table S1: Site Locations with Precipitation Metrics
841
842 Supplementary Table S2: QIIME Local Soil and Rhizobiome Sequences Before and
843 After   Quality   Trimming
844
845 Supplementary Table S3: ASV Assignments for the Local Soil and Rhizobiome Samples
846
847 Supplementary Table S4: Soil Chemistry
848
849 Supplementary Table S5: Soil Properties and Alpha Diversity Statistical Analyses
850
851 Supplementary Table S6: Alpha Diversity for the Local Soil and Rhizobiome Samples
852
853 Supplementary Table S7: Precipitation and pH Gradient t-Test Results
854
855 Supplementary Table S8: ASV Distribution by Sample Type and Cluster Category
856
857 Supplementary Table S9: Beta Dispersion Distance for Local Soil and Rhizobiome
858 Samples within a Location
859
860 Supplementary Table S10: Unbiased Taxonomic Oligotyping Results
861
862 Supplementary Table S11: Oligotyping Results for Individual Families
863

References

1. Bandopadhyay, S., Li, X., Bowsher, A. W., Last, R. L. & Shade, A. Disentangling plant- and environment-mediated drivers of active rhizosphere bacterial community dynamics during short-term drought. *Nat. Commun.* **15**, 6347 (2024).

2. Bei, Q., Moser, G., Wu, X., Müller, C. & Liesack, W. Metatranscriptomics reveals climate change effects on the rhizosphere microbiomes in European grassland. *Soil Biol. Biochem.* **138**, 107604 (2019).

3. Direct and indirect effects of climate change on soil microbial and soil microbial☐plant interactions: What lies ahead? doi:10.1890/ES15-00217.1.

4. Zeglin, L. H. *et al.* Altered precipitation regime affects the function and composition of soil microbial communities on multiple time scales. *Ecology* **94**, 2334–2345 (2013).

5. Axelrod, D. I. Rise of the Grassland Biome, Central North America. *Bot. Rev.* **51**, 163–201 (1985).

6. Maricle, B. R., Caudle, K. L., Lindsey, K. J., Baer, S. G. & Johnson, L. C. Effects of Extreme Drought on Photosynthesis and Water Potential of Andropogon gerardii (Big Bluestem) Ecotypes in Common Gardens Across Kansas. *tkas* **120**, 1–16 (2017).

7. Kramer, D. L. *et al.* Drought tolerance in ecotypes of big bluestem (Andropogon gerardii) relates to above-ground surface area: Results from a common garden experiment. *Flora* **246-247**, 52–60 (2018).

8. Galliart, M. *et al.* Adaptive genetic potential and plasticity of trait variation in the foundation prairie grass Andropogon gerardii across the US Great Plains' climate gradient: Implications for climate change and restoration. *Evol. Appl.* **13**, 2333–2356 (2020).

9. Mendola, M. L., Baer, S. G., Johnson, L. C. & Maricle, B. R. The role of ecotypic variation and the environment on biomass and nitrogen in a dominant prairie grass. *Ecology* **96**, 2433–2445 (2015).

10. Olsen, J. T., Caudle, K. L., Johnson, L. C., Baer, S. G. & Maricle, B. R. Environmental and genetic variation in leaf anatomy among populations of Andropogon gerardii (Poaceae) along a precipitation gradient. *Am. J. Bot.* **100**, 1957–1968 (2013).

11. Barnes, P. W. Adaptation to Water Stress in the Big Bluestem-Sand Bluestem Complex. *Ecology* **66**, 1908–1920 (1985).

12. Johnson, L. C. *et al.* Intraspecific variation of a dominant grass and local adaptation in reciprocal garden communities along a US Great Plains' precipitation gradient: implications for grassland restoration with climate change. *Evol. Appl.* **8**, 705–723 (2015).

13. Wang, X.-B. *et al.* Habitat-specific patterns and drivers of bacterial β-diversity in China's drylands. *ISME J.* **11**, 1345–1358 (2017).

14. Cai, Z., Wang, X., Bhadra, S. & Gao, Q. Distinct factors drive the assembly of quinoa-associated microbiomes along elevation. *Plant Soil* **448**, 55–69 (2020).

15. Alegria Terrazas, R. *et al.* A footprint of plant eco-geographic adaptation on the composition of the barley rhizosphere bacterial microbiota. *Sci. Rep.* **10**, 12916 (2020).

16. Pang, Z., Xu, P. & Yu, D. Environmental adaptation of the root microbiome in two
rice ecotypes. *Microbiol. Res.* **241**, 126588 (2020).
17. Marasco, R. *et al.* Rhizosheath microbial community assembly of sympatric desert
speargrasses is independent of the plant host. *Microbiome* **6**, 215 (2018).
18. Cowan, D. A. *et al.* Biogeographical survey of soil microbiomes across sub-Saharan
Africa: structure, drivers, and predicted climate-driven changes. *Microbiome* **10**, 131
(2022).
19. Kazarina Anna *et al.* Home-field advantage affects the local adaptive interaction
between Andropogon gerardii ecotypes and root-associated bacterial communities.
Microbiology Spectrum **11**, e00208–23 (2023).
20. Sarkar, S. *et al.* Bacterial but Not Fungal Rhizosphere Community Composition
Differ among Perennial Grass Ecotypes under Abiotic Environmental Stress.
Microbiol Spectr **10**, e0239121 (2022).
21. Seager, R. *et al.* Projections of declining surface-water availability for the
southwestern United States. *Nat. Clim. Chang.* **3**, 482–486 (2012).
22. Seager, R. *et al.* Dynamical and Thermodynamical Causes of Large-Scale Changes
in the Hydrological Cycle over North America in Response to Global Warming. *J.
Clim.* **27**, 7921–7948 (2014).
23. Cook, B. I., Smerdon, J. E., Seager, R. & Coats, S. Global warming and 21st
century drying. *Clim. Dyn.* **43**, 2607–2627 (2014).
24. Seager, R. *et al.* Whither the 100th meridian? The once and future physical and
human geography of America's arid–humid divide. Part II: The meridian moves
east. *Earth Interact.* **22**, 1–24 (2018).
25. Knapp, A. K., Briggs, J. M., Blair, J. M. & Turner, C. L. Patterns and controls of
aboveground net primary production in tallgrass prairie. *Grassland dynamics: long-
term ecological research in tallgrass prairie. Oxford University Press, New York*
193–221 (1998).
26. Fred B. Samson, Fritz L. Knopf, and Wayne R. Ostlie. Great Plains ecosystems:
past, present, and future. *Wildlife Society Bulletin* **32**, 6–15 (2004).
27. Ren, Z. *et al.* The role of dominant prairie species ecotypes on plant diversity
patterns of restored grasslands across a rainfall gradient in the US Great Plains.
Appl. Veg. Sci. **26**, (2023).
28. Hoekstra, J. M., Boucher, T. M., Ricketts, T. H. & Roberts, C. Confronting a biome
crisis: global disparities of habitat loss and protection. *Ecol. Lett.* **8**, 23–29 (2005).
29. St. Laurent M. Trypaluk C. Unruh D. Martin D. & Wilhite O., P. S. P. S. *NOAA Atlas
14 Volume 11 Version 2: Precipitation-Frequency Atlas of the United States,
Northeastern States. NOAA* (2020).
30. Combs, S. M. and Nathan, M. V. *Recommended Chemical Soil Test Procedures for
the North Central Region North Central Regional*. vol. 211 (Reised) (University of
Missouri Agricultural Experiment Station, Columbia, MO, 1998).
31. Dakota, S. Recommended Chemical Soil Test Procedures for the North Central
Region. (1998).
32. Watson, M. pH and lime requirement.
33. Mclean, E. O. Soil pH and Lime Requirement. in *Agronomy Monographs* 199–224
(American Society of Agronomy, Soil Science Society of America, Madison, WI,
USA, 2015). doi:10.2134/agronmonogr9.2.2ed.c12.

34. Brown, J. R. & Cisco, J. R. An improved Woodruff buffer for estimation of lime
requirements. *Soil Sci. Soc. Am. J.* **48**, 587–592 (1984).
35. Mehlich, A. Mehlich 3 soil test extractant: A modification of Mehlich 2 extractant.
Commun. Soil Sci. Plant Anal. **15**, 1409–1416 (1984).
36. Murphy, J. & Riley, J. P. A modified single solution method for the determination of
phosphate in natural waters. *Anal. Chim. Acta* **27**, 31–36 (1962).
37. LECO Corporation, St. Joseph, MI. Carbon and Nitrogen in Soil and Sediment.
(2005).
38. Foth, H.D., H.S. Jacobs, and L.V. Withee. Laboratory Manual for Introductory Soil
Science, 3rd Ed. 12–13 (1971).
39. Bouyoucos, G. J. Hydrometer method improved for making particle size analysis of
soils. *Agron. J.* **54**, 464–465 (1962).
40. Omega Bio-tek, Inc, Norcross, GA, USA. E.Z.N.A.® Soil DNA Kit (Revision No: 1.3).
41. Caporaso, J. G. *et al.* Ultra-high-throughput microbial community analysis on the
Illumina HiSeq and MiSeq platforms. *ISME J.* **6**, 1621–1624 (2012).
42. Bolyen, E. *et al.* Reproducible, interactive, scalable and extensible microbiome data
science using QIIME 2. *Nat. Biotechnol.* **37**, 852–857 (2019).
43. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. *EMBnet J.* **17**, 10 (2011).
44. Callahan, B. J. *et al.* DADA2: High-resolution sample inference from Illumina
amplicon data. *Nat. Methods* **13**, 581–583 (2016).
45. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data
processing and web-based tools. *Nucleic Acids Res.* **41**, D590–6 (2013).
46. Cao, P. *et al.* Environmental filtering process has more important roles than
dispersal limitation in shaping large-scale prokaryotic beta diversity patterns of
grassland soils. *Microb. Ecol.* **72**, 221–230 (2016).
47. Eren, A. M. *et al.* Oligotyping: Differentiating between closely related microbial taxa
using 16S rRNA gene data. *Methods Ecol. Evol.* **4**, 1111–1119 (2013).
48. Eren, A. M. *et al.* Minimum entropy decomposition: unsupervised oligotyping for
sensitive partitioning of high-throughput marker gene sequences. *ISME J.* **9**, 968–
979 (2015).
49. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. *Nucleic Acids Res.* **32**, 1792–1797 (2004).
50. RStudio Team. RStudio: Integrated Development for R. RStudio. *PBC* (2024).
51. R Core Team. R: A language and environment for statistical computing. *R*
Foundation for Statistical Computing, Vienna, Austria (2024).
52. J. Fox, S. W. *An R Companion to Applied Regression, 3rd Ed.* (SAGE Publications,
Thousand Oaks, CA, 2019).
53. H. Wickham, R. François, L. Henry, K. Müller. *dplyr: A Grammar of Data*
Manipulation, R package version 1.0.7. (2021).
54. Chang, H. W. J. H. *devtools: Tools to Make Developing R Packages Easier*, R
package version 2.4.3. (2021).
55. Dinno, A. dunn.test: Dunn's Test of Multiple Comparisons Using Rank Sums. R
package version 1.3.5. (2017).
56. Arbizu, F. *pairwiseAdonis: Pairwise Multilevel Comparison Using Adonis*, R
package version 0.4. (2020).

57. Hadley Wickham and Davis Vaughan and Maximilian Girlich. tidyr: Tidy Messy
Data. *R package version 1.3.1*.
58. Wickham et al., H. Welcome to the Tidyverse. *J. Open Source Softw.* **4**, 1686
(2019).
59. Wickham, H. ggplot2: Elegant Graphics for Data Analysis. *Springer-Verlag New*
York (2016).
60. Derek H. Ogle and Jason C. Doll and A. Powell Wheeler and Alexis Dinno. FSA:
Simple Fisheries Stock Assessment Methods. *R package version 0.9.5* (2023).
61. David Robinson and Alex Hayes and Simon Couch. broom: Convert Statistical
Objects into Tidy Tibbles. *R package version 1.0.7* (2024).
62. Kassambara, A. rstatix: Pipe-Friendly Framework for Basic Statistical Tests. *R*
package version 0.7.2 (2023).
63. J. Oksanen, F.G. Blanchet, M. Friendly, et al. vegan: Community Ecology Package
R package version 2.5-7. (2020).
64. Hadley Wickham and Jim Hester and Jennifer Bryan. readr: Read Rectangular Text
Data. *R package version 2.1.5* (2024).
65. Pedersen, T. L. patchwork: The Composer of Plots. *R package version 1.2.0*
(2024).
66. Bretz, W. T. H. Simultaneous Inference in General Parametric Models. *Biometrical*
Journal **50**, 346–363 (2008).
67. R. Graves, F. Piepho, L. Selzer, A. Dorai-Raj. *multcompView: Visualizations of*
Paired Comparisons, R package version 0.1-8. (2019).
68. de Mendiburu, F. *agricolae: Statistical Procedures for Agricultural Research*, R
package version 1.3-5. (2020).
69. Dusa, A. venn: Draw Venn Diagrams. *R package version 1.12* (2024).
70. Martin Maechler and Peter Rousseeuw and Anja Struyf and Mia Hubert and Kurt
Hornik. cluster: Cluster Analysis Basics and Extensions. (2023).
71. Pebesma, E. J. Simple Features for R: Standardized Support for Spatial Vector
Data. *R J.* **10**, 439–446 (2018).
72. Becker, R.A., Wilks, A.R., Brownrigg, R., Minka, T.P., & Deckmyn, A. *maps: Draw*
Geographical Maps. R package version 3.3.0. (2018).
73. Wickham, H. Reshaping Data with the {reshape} Package. *Journal of Statistical*
Software **21**, 1–20 (2007).
74. Kolde, R. *pheatmap: Pretty Heatmaps*, R package version 1.0.12. (2017).
75. Neuwirth, E. RColorBrewer: ColorBrewer Palettes, R package version 1.1-2. (2014).
76. Auguie, B. gridExtra: Miscellaneous Functions for 'Grid' Graphics. R package
version 2.3. (2017).
77. P. J. McMurdie, S. H. Phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. *PLoS One* **8**, e61217 (2013).
78. G. Csardi, T. N. The igraph software package for complex network research.
InterJournal **1695**, (2006).
79. Campitelli, D. *ggnewscale: Multiple Fill and Colour Scales in 'ggplot2'*, R package
version 0.4.5. (2021).
80. Arnold, J. B. ggthemes: Extra Themes, Scales and Geoms for 'ggplot2'. *R package*
version 5.1.0 (2024).
81. Claus O. Wilke and Brenton M. Wiernik. ggtext: Improved Text Rendering Support

for 'ggplot2'. *R package version 0.1.2* (2022).

82. Dusa, C.-H. G. A. ggVennDiagram: A 'ggplot2' Implement of Venn Diagram. *R*
package version 1.5.2 (2024).

83. Ammar, R. randomcoloR: Generate Attractive Random Colors. **R packageersion**
1.1.0.1, (2019).

84. Wickham, H. stringr: Simple, Consistent Wrappers for Common String Operations.
R package version 1.5.1 (2023).

85. Chen, H., & Boutros, P.C. VennDiagram: A package for the generation of highly-
customizable Venn and Euler diagrams in R. *BMC Bioinformatics* **12**, (2011).

86. Yang, C. *et al.* Influence of plant genotype and soil on the cotton rhizosphere
microbiome. *Front. Microbiol.* **13**, 1021064 (2022).

87. Easterling, D. R. *et al.* Precipitation change in the United States. (2017).

88. Seager, R. *et al.* Whither the 100th meridian? The once and future physical and
human geography of America's arid–humid divide. Part I: The story so far. *Earth*
Interact. **22**, 1–22 (2018).

89. Walters, K. E. & Martiny, J. B. H. Alpha-, beta-, and gamma-diversity of bacteria
varies across habitats. *PLoS One* **15**, e0233872 (2020).

90. Zhang, Q. *et al.* Alpha, beta and gamma diversity differ in response to precipitation
in the Inner Mongolia grassland. *PLoS One* **9**, e93518 (2014).

91. Yao, M. *et al.* The differentiation of soil bacterial communities along a precipitation
and temperature gradient in the eastern Inner Mongolia steppe. *Catena* **152**, 47–56
(2017).

92. Chen, P. *et al.* Water management-mediated changes in the rhizosphere and bulk
soil microbial communities alter their utilization of urea-derived carbon.
Microorganisms **12**, 1829 (2024).

93. Vieira, S. *et al.* Drivers of the composition of active rhizosphere bacterial
communities in temperate grasslands. *ISME J.* **14**, 463–475 (2020).

94. Orwin, K. H. *et al.* Linkages of plant traits to soil properties and the functioning of
temperate grassland: Links of plant traits to soil properties. *J. Ecol.* **98**, 1074–1083
(2010).

95. Badri, D. V. & Vivanco, J. M. Regulation and function of root exudates. *Plant Cell*
Environ. **32**, 666–681 (2009).

96. Rengel, Z. Role of pH in availability of ions in soil. *Handbook of plant growth pH as*
the master variable (2002) doi:10.1201/9780203910344-20/role-ph-availability-ions-
soil-zdenko-rengel.

97. Tan, W., Wang, J., Bai, W., Qi, J. & Chen, W. Soil bacterial diversity correlates with
precipitation and soil pH in long-term maize cropping systems. *Sci. Rep.* **10**, 6012
(2020).

98. Zhou, Z., Wang, C. & Luo, Y. Meta-analysis of the impacts of global change factors
on soil microbial diversity and functionality. *Nat. Commun.* **11**, 3072 (2020).

99. Rousk, J. *et al.* Soil bacterial and fungal communities across a pH gradient in an
arable soil. *ISME J.* **4**, 1340–1351 (2010).

100. Li, J., Benti, G., Wang, D., Yang, Z. & Xiao, R. Effect of alteration in precipitation
amount on soil microbial community in a semi-arid grassland. *Front. Microbiol.* **13**,
842446 (2022).

101. Li, X., Yan, Y., Lu, X., Fu, L. & Liu, Y. Responses of soil bacterial communities to

precipitation change in the semi-arid alpine grassland of Northern Tibet. *Front. Plant*
Sci. **13**, 1036369 (2022).

102.   Bhattacharyya, A. *et al.* Rhizosphere plant-microbe interactions under water
stress. *Adv. Appl. Microbiol.* **115**, 65–113 (2021).

103.   Nicol, G. W. & Schleper, C. Ammonia-oxidising Crenarchaeota: important players
in the nitrogen cycle? *Trends Microbiol.* **14**, 207–212 (2006).

104.   Sutherland, J., Bell, T., Trexler, R. V., Carlson, J. E. & Lasky, J. R. Host genomic
influence on bacterial composition in the switchgrass rhizosphere. *Mol. Ecol.* **31**,
3934–3950 (2022).

105.   Galliart, M. *et al.* Local adaptation, genetic divergence, and experimental
selection in a foundation grass across the US Great Plains' climate gradient. *Glob.*
Chang. Biol. **25**, 850–868 (2019).

106.   Lucas, J. A. *et al.* Structural and functional study in the rhizosphere of Oryza
sativa L. plants growing under biotic and abiotic stress. *J. Appl. Microbiol.* **115**, 218–
235 (2013).

107.   Hartman, K. & Tringe, S. G. Interactions between plants and soil shaping the root
microbiome under abiotic stress. *Biochem. J.* **476**, 2705–2724 (2019).

108.   Ginnan, N. A. *et al.* Persistent legacy effects on soil metagenomes facilitate plant
adaptive responses to drought. *bioRxivorg* (2024) doi:10.1101/2024.08.26.609769.

109.   Hannula, S. E. *et al.* Persistence of plant-mediated microbial soil legacy effects in
soil and inside roots. *Nat. Commun.* **12**, 5686 (2021).

110.   Metze, D. *et al.* Microbial growth under drought is confined to distinct taxa and
modified by potential future climate conditions. *Nat. Commun.* **14**, 5895 (2023).

111.   Xie, J. *et al.* Drought stress triggers shifts in the root microbial community and
alters functional categories in the microbial gene pool. *Front. Microbiol.* **12**, 744897
(2021).

112.   Waring, B. & Hawkes, C. V. Ecological mechanisms underlying soil bacterial
responses to rainfall along a steep natural precipitation gradient. *FEMS Microbiol.*
Ecol. **94**, fiy001 (2018).

113.   Richter-Heitmann, T. *et al.* Stochastic dispersal rather than deterministic
selection explains the spatio-temporal distribution of soil bacteria in a temperate
grassland. *Front. Microbiol.* **11**, 1391 (2020).

114.   Evans, S., Martiny, J. B. H. & Allison, S. D. Effects of dispersal and selection on
stochastic assembly in microbial communities. *ISME J.* **11**, 176–185 (2017).

115.   Naylor, D., DeGraaf, S., Purdom, E. & Coleman-Derr, D. Drought and host
selection influence bacterial community dynamics in the grass root microbiome.
ISME J. **11**, 2691–2704 (2017).

116.   Albright, M. B. N. *et al.* Interactions of microhabitat and time control grassland
bacterial and fungal composition. *Front. Ecol. Evol.* **7**, 447321 (2019).

117.   Staley, C. *et al.* Urea amendment decreases microbial diversity and selects for
specific nitrifying strains in eight contrasting agricultural soils. *Front. Microbiol.* **9**,
634 (2018).

118.   Boukhatem, Z. F., Merabet, C. & Tsaki, H. Plant growth promoting
Actinobacteria, the most promising candidates as bioinoculants? *Front. Agron.* **4**,
849911 (2022).

119.   Trujillo, M. E., Riesco, R., Benito, P. & Carro, L. Endophytic Actinobacteria and

the interaction of Micromonospora and nitrogen fixing plants. *Front. Microbiol.* **6**,
1341 (2015).

120.   Sim, J. X. F. *et al.* Pesticide effects on nitrogen cycle related microbial functions
and community composition. *Sci. Total Environ.* **807**, 150734 (2022).

121.   Millar, N. S. & Bennett, A. E. Stressed out symbiotes: hypotheses for the
influence of abiotic stress on arbuscular mycorrhizal fungi. *Oecologia* **182**, 625–641
(2016).

122.   Muhammad, A. *et al.* Exploring plant-microbe interactions in adapting to abiotic
stress under climate change: a review. *Front. Plant Sci.* **15**, 1482739 (2024).

123.   Chen, W.-C., Ko, C.-H., Su, Y.-S., Lai, W.-A. & Shen, F.-T. Metabolic potential
and community structure of bacteria in an organic tea plantation. *Appl. Soil Ecol.*
157, 103762 (2021).

124.   Schmidt, V. T. *et al.* Oligotyping reveals community level habitat selection within
the genus Vibrio. *Front. Microbiol.* **5**, 563 (2014).

125.   Eren, A. M. *et al.* A single genus in the gut microbiome reflects host preference
and specificity. *ISME J.* **9**, 90–100 (2015).

126.   Giongo, A., Arnhold, J., Grunwald, D., Smalla, K. & Braun-Kiewnick, A. Soil
depths and microhabitats shape soil and root-associated bacterial and archaeal
communities more than crop rotation in wheat. *Front. Microbiomes* **3**, 1335791
(2024).

127.   Dastogeer, K. M. G., Tumpa, F. H., Sultana, A., Akter, M. A. & Chakraborty, A.
Plant microbiome–an account of the factors that shape community composition and
diversity. *Curr. Plant Biol.* **23**, 100161 (2020).

128.   Tripathi, B. M. *et al.* Trends in taxonomic and functional composition of soil
microbiome along a precipitation gradient in Israel. *Microb. Ecol.* **74**, 168–176
(2017).

129.   Naidoo, Y., Valverde, A., Pierneef, R. E. & Cowan, D. A. Differences in
precipitation regime shape microbial community composition and functional
potential in Namib Desert soils. *Microb. Ecol.* **83**, 689–701 (2022).

130.   Gray, M. M. *et al.* Ecotypes of an ecologically dominant prairie grass
(Andropogon gerardii) exhibit genetic divergence across the U.S. Midwest
grasslands' environmental gradient. *Mol. Ecol.* **23**, 6011–6028 (2014).

1171
1172
1173

ASV Distribution by pH Category, Type, and Ecotype
% of Total Relative Abundance by Phylum

ASV Distribution by PR Category, Ecotype, and Type
% of Total Relative Abundance by Phylum

Significant Changes in Phylum Relative Abundance Across PR Gradient by Type