Assessing plant phenological changes based on drivers of spring phenology

Yong Jiang[1], Stephen J. Mayor[2], Xiuli Chu[3], Xiaoqi Ye[4], Rongzhou Man[2]*, Jing Tao[5]*, and Qing-Lai Dang[6]

[1]Key Laboratory of Ecology of Rare and Endangered Species and Environmental Protection of Ministry of Education, Guangxi Normal University, China yongjiang226@126.com

[2]Ontario Ministry of Natural Resources and Forestry, Ontario Forest Research Institute, 1235 Queen Street East, Sault Ste. Marie, ON, P6A 2E5 Canada stephen.mayor@ontario.ca; rongzhou.man@ontario.ca

[3]Shanghai Botanical Garden, Shanghai Engineering Research Center of Sustainable Plant Innovation, 1111 Longwu Road, Xuhui District, Shanghai, China 200231 xiulic0207@163.com

[4]Research Institute of Subtropical Forestry, Chinese Academy of Forestry, Hangzhou, China mengxqi@yahoo.com

[5]Jilin Provincial Academy of Forestry Sciences, Changchun, China taojing8116@126.com

[6]Faculty of Natural Resources Management, Lakehead University, Thunder Bay, Canada P7B 5E1 qdang@lakeheadu.ca

*Corresponding authors: rongzhou.man@ontario.ca; taojing8116@126.com

ABSTRACT

Understanding plant phenological responses to climate warming is crucial for predicting changes

in plant communities and ecosystems but difficult with sensitivity analysis that is not linked to

drivers of spring phenology. In this article, we present a new measure *phenological lag* to

quantify the overall effect of phenological constraints including insufficient winter chilling,

photoperiod, and environmental stresses, based on observed response and that expected from

species-specific changes in spring temperatures, i.e., changes in spring forcing (degree days)

from warming and average temperature at time of species events. We applied this new analytical

framework to a global dataset with 980 species and 1527 responses to synthesize observed

changes in spring phenology and investigate the contributions of forcing change, growth

temperature, and phenological lag to differential phenological responses reported previously.

Keywords: spring phenology, phenological lag, forcing change, growth temperature,

phenological response, meta-analysis

Mechanisms of changes in spring phenology

Plant phenology, particularly in spring, is shifting with warming climate (Fitter and Fitter, 2002; Post et al., 2018; Primack and Gallinat, 2016). Numerous observational studies and controlled experiments, conducted over a range of climatic and phenological conditions (Huang et al., 2020; Parmesan, 2007; Prevéy et al., 2019; Root et al., 2003; Wolkovich et al., 2012), have produced varying results, with observed phenological changes differing by research approach (Wolkovich et al., 2012), climatic region (Parmesan, 2007; Post et al., 2018; Prevéy et al., 2017; Zhang et al., 2015), and functional group (Parmesan, 2007; Root et al., 2003; Willis et al., 2010; Wolkovich et al., 2013; Zettlemoyer et al., 2019). For example, observed changes are often smaller with experimental studies (Wolkovich et al., 2012), native species (Wolkovich et al., 2013), and warmer regions (Prevéy et al., 2017). Understanding these differences (also referred to as discrepancies or mismatches) is crucial for predicting changes in plant communities and ecosystems (Fitter and Fitter, 2002; Polgar et al., 2014; Primack and Gallinat, 2016) but difficult with sensitivity analysis that is not linked to drivers of spring phenology.

Plants in the northern hemisphere require an accumulation of cool winter temperatures (winter chilling) to break dormancy and an accumulation of warm spring temperatures (spring forcing, degree days) to initiate species event (e.g., leafing or flowering) (Piao et al., 2019; Way and Montgomery, 2015). This means that changes in spring phenology result primarily from changes in spring forcing prior to species events if chilling requirements are fully met and other phenological constraints (e.g., photoperiod effect and environmental stresses) remain constant. While the direction of phenological changes (advance or delay) is set by the sign of forcing changes (+ or -), the magnitude of changes depends on quantity of forcing changes, as well as level of growth temperatures at time of species event with a warmer climate, i.e., higher growth

temperatures are associated with smaller observed changes and sensitivity and lower growth

temperatures with larger observed changes and sensitivity (Chu et al., 2021, 2023; Prevéy et al.,

2017; Shen et al., 2014). As species often differ in timing of spring events, both forcing change

and growth temperature are species-specific, reflecting changes in spring temperatures that are

relevant to phenological changes of individual species (Chu et al., 2021, 2023). The effects of

photoperiod (day length) and environmental stresses (e.g., drought and spring freezing) are

similar to those of insufficient winter chilling in constraining the advance of spring phenology

(Chen et al., 2011; Huang et al., 2019; Körner and Basler, 2010; Ma et al., 2019; Way and

Montgomery, 2015). The difference between observed phenological change and that expected

from forcing change and growth temperature represents phenological difference due to effects of

phenological constraints and can be thereafter named phenological lag. Separating effects of

different constraints is possible if soil moisture/plant water status is monitored (Huang et al.,

2019; Post et al., 2022), or species-specific chilling and photoperiod effects (Man et al., 2017a,

2021a; Fu et al., 2019) or cold hardiness (Man et al., 2017b, 2021b) are known from previous

research. For example, phenological lag can represent the effect of insufficient chilling when

photoperiod effect and environmental stresses are constant (Chen et al., 2011).

The changes in spring phenology are often quantified with sensitivity, a parameter that

reflects rates of changes but not uneven warming and responses (Beaubien and Hamann, 2011;

Keenan et al., 2020; Post et al., 2018; Rafferty et al., 2020) or magnitude and underlying

mechanisms of changes in spring phenology (Chu et al., 2021, 2023). Here, we present a new

method to partition observed phenological change into forcing change, growth temperature, and

phenological lag, based on changes in spring phenology and temperatures with control (baseline)

and warmer climates. We applied this analytical approach to meta-analyze phenological changes

reported in the literature and examined the contributions of forcing change, growth temperature,

and phenological lag to differential responses reported previously, i.e., research approach

(observational or experimental study), species origin (native or exotic), climatic region (boreal or

temperate), and growth form (tree, shrub, herb, or grass).

Partitioning observed changes

Species-specific forcing change (F_C) can be calculated from the difference between degree days

(> 0 °C, see Man and Lu, 2010) of baseline (or control) ($\sum_1^i T_{Ci}$) and warmer ($\sum_1^i T_{Wi}$)

temperatures at the time of species event (leafing or flowering) with baseline temperatures (O_{Ci})

(see Figure 1).

$$F_C = \sum_1^i T_{Wi} - \sum_1^i T_{Ci}$$

The phenological response expected under the null hypothesis that climate warming does

not induce changes in phenological constraints can be determined from forcing change and

species phenology with baseline temperatures (Figure 1):

$$N_{Ei} = \min \left| \sum_1^{i-n} T_{Wi} - \sum_1^i T_{Ci} \right|$$

where N_E is the difference in the number of days (n) between baseline $\sum_1^i T_{Ci}$ and warmer

$\sum_1^{i-n} T_{Wi}$ temperatures in reaching species forcing needs with baseline temperatures (spring

forcing at O_{Ci}). Under this hypothesis, spring phenology is expected to shift from O_{Ci} with

baseline temperatures to E_{Wi} with warmer temperatures. N_E is typically positive, meaning earlier

leafing and flowering, but can be negative if temperatures decrease over time in observational

studies.

Growth temperature (T_N) is the average temperature or rate of forcing accumulation

within the window of expected phenological response (between O_{Ci} and E_{Wi}) and can be

determined from forcing change and expected response.

115                     $$T_N = F_C/N_E$$

116   The difference between expected (N_E) and observed (N_O) responses is phenological lag

117   (N_C) reflecting phenological difference due to effects of phenological constraints.

118                    $$N_C = N_E - N_O$$

$N_C > 0$ indicates increasing phenological constraints that may result from increasing chilling

deficiency, photoperiod restriction, moisture stress, or risks of spring frosts; $N_C = 0$ indicates no

changes; and $N_C < 0$ indicates decreasing constraints with warming climate.

Meta-analysis of global data

Phenological Data

We followed the guidelines of PRISMA (preferred reporting items for systematic reviews and

meta-analyses) (O'Dea et al., 2021; Page et al., 2020) to build up the dataset for responses of

plant spring phenology to experimental and climatic warming. We used Web of Science and

Google Scholar to search experimental and observational studies on warming-induced changes in

spring phenology. The following criteria were used to select papers: (a) accessible peer-reviewed

articles published in scientific journals for boreal (MAT < 6 °C) and temperate (MAT ≥ 6 °C)

regions in the northern hemisphere, (b) studies reporting spring leafing or flowering before Julian

day 213 (July 31), (c) studies containing different climatic conditions of baseline (lower

temperatures in early period of observations or unwarmed control treatment) and warmer (higher

temperatures in later period of observations or warmer treatment) climates, (d) access to local

temperature data (only one study was excluded due to lack of nearby temperature data within 50

km), and (e) reported phenological changes in unique locations, species, and periods of

observations/warmer treatments. When publications did not contain sufficient phenological

information, we accessed data directly from phenological databases (e.g., China's National Earth

System Science Data Center). When results were reported in graphical format, we extracted the

required data using the distance measuring tool in Adobe Acrobat Reader DC. In experimental

studies with multiple warming treatments, we selected the treatment with the smallest

temperature increases to represent a likely scenario of climate change and to maintain data

independence. We also excluded studies where the timing of phenological events, total

temperature change, or observational time periods were not available.

In total, 66 studies were identified from 87 locations, containing 1,527 phenological

responses for 980 species (Tables S1–S3). For studies reporting multiple phenological stages, we

only used data for the first occurrence of the event. To account for variation in phenological

responses, each study location was characterized by research approach (observation or

experiment), climatic region (boreal or temperate), biogeographic origin of species (exotic or

native to study area), and growth form (tree, shrub, herb, or grass).

Climate Data

Baseline and warmer temperature data for calculating forcing change, expected response, growth

temperature, phenological lag, spring warming (average temperature change from Julian day 1 to

182), and long-term MAT and MAP were obtained from 9 different sources (Table S4). When

temperature data were available from multiple sources, we selected the data set with the fewest

missing values. For each phenological response, the number of weather stations within 50 km of

data collection ranged from 1 for studies at a single location to 14 for observational studies

across large geographic areas (Table S2).

Data analysis

W calculated forcing change, expected response, growth temperature, and phenological lag for

each of the 1527 responses compiled. We used linear mixed-effects models to explore the

differences in observed responses and phenological lags between research approaches

(observations vs. experiments), species origins (native vs. exotic), and climatic regions (boreal

vs. temperate), and among growth forms (trees, shrubs, herbs, and grasses). Separate analysis

was conducted for each of the four fixed effects and two species events (leafing and flowering).

Location and species were treated as random effects.

To assess the influences of climatic, phenological, and biological variables, we used a

stepwise regression with automated combined forward and backward selection by Akaike

information criterion (AIC) (Burnham and Anderson, 2002) to select the best combination of

variables for predicting observed responses (N_O).

$$N_O \sim A_L + L_A + MAT + MAP + D_T + T_N + F_C + S_W + N_C$$

The variables included were altitude (A_L), latitude (L_A), MAT, MAP, spring phenology (D_T),

growth temperature (T_N), forcing change (F_C), spring warming (S_W), and phenological lag (N_C).

Observed changes in spring phenology

Across 66 studies with 980 species and 1527 responses, observed responses (N_O) ranged from

−23.8 to +26.8 days for leafing and −36.3 to +49.9 days for flowering, with an average of +6.0

days for both events. These changes differed from expected responses (N_E) which ranged from

−3 to +27 days for leafing and −3 to +26 days for flowering and averaged +7.8 and +9.5 days, respectively. Observed phenological responses were thus smaller than the expected by 1.8 and 3.5 days, on average, for leafing and flowering, respectively.

In experimental studies, observed flowering response was non-significantly smaller than those from observational studies (1.7 days, p=0.232, Figure 2a), while phenological lag was significantly longer for both leafing (3.0 days, p=0.028) and flowering (2.7 days, p=0.027) (Figure 2b). Observed flowering response in exotic species was 2 days greater than in native species (p=0.016), likely due to differences in phenological lag (1.9 days, p=0.019). Observed leafing and flowering responses in boreal region were 4.3 and 1.9 days smaller (p=0.004, p=0.088), respectively, than in temperate region, likely due to smaller forcing changes (equivalent to 2.8 days in leafing and 1.5 days in flowering based on forcing changes and growth temperatures, Table 1 note a). Observed leafing response in grasses was 3.8 days smaller than in trees and shrubs (p=0.033), likely due to differences in forcing changes (6.0 days, Table 1 note b).

Other than phenological lag, forcing change and growth temperature were the most influential variables affecting observed responses (Table 2). The remaining variables combined (altitude, latitude, MAT, and spring warming) explained <2.5% variations in observed leafing and flowering responses.

Differential phenological responses

Observations vs. experiments

By partitioning observed changes based on drivers of spring phenology, we clarified some of the uncertainty regarding the underlying mechanisms of differences in plant phenological response

to climate warming. By synthesizing global data, we found that observed flowering responses in

experimental studies were non-significantly smaller than that in observational studies, but this

was not the case for leafing responses; these findings differ from those based on temperature

sensitivity (Wolkovich et al., 2012). The greater warming in experimental studies did not

produce greater observed responses, due to longer phenological lags, which may have resulted in

the reduced sensitivity (Wolkovich et al., 2012). First, experimental studies are often conducted

at higher altitude sites with lower MAT, later spring phenology, and therefore longer

accumulation of winter chilling. The longer phenological lags are therefore unlikely the result of

insufficient winter chilling. Second, artificial warming has typically provided higher forcing

changes and growth temperatures than natural climate warming (see Table 1). Greater forcing

changes produce larger expected responses (observed response + phenological lag), whereas

higher growth temperatures reduce expected responses (Chu et al., 2021; Prevéy et al., 2017;

Wolkovich et al., 2021). The high temperatures in the artificial warming structures (Marion et

al., 1997) can also reduce humidity and soil moisture (Ettinger et al., 2019; Huang et al., 2019),

slowing spring development (Ganjurjav et al., 2020; Huang et al., 2019; Moore et al., 2015),

particularly in early spring when temperatures in the surrounding environment are low, thus

restricting soil water movement. While the different forcing changes adequately accounts for the

differences in expected leafing responses (2.5 out of 3.3 days, Table 1 note c), the large

differences in forcing changes but similar expected responses in flowering (8.63 and 9.07 days,

Figure 2) suggests a strong effect of growth temperature (Table 1 note d) (Chu et al., 2023).

Thus, warming-induced moisture stress and high growth temperatures may interact to alter

observed phenological responses and temperature sensitivity in experimental studies (Wolkovich

et al., 2012). Future experiments should consider humidity, soil moisture, and plant water status

to better understand disparities between observational and experimental studies and relate plant

phenological changes from experimental warming to those from natural climate change

(Wolkovich et al., 2012).

Native vs. exotic species

Exotic species are often reported with more noticeable observed responses to warming, a trend

that is generally based on flowering (Calinger et al., 2013; Willis et al., 2010; Wolkovich et al.,

2013). The ecological implications of this response are well described (Polgar et al., 2014;

Zettlemoyer et al., 2019), but the underlying mechanisms are not well understood. Lower growth

temperatures faced by early-start exotics are likely partially the cause (Chu et al., 2021, 2023), as

shown by a reverse trend for late-start exotics (Zohner and Renner, 2014). In this synthesis,

phenological lag did not increase from early leafing to late flowering for exotic species, a trend

that is consistent with dormancy release commonly reported on leaf and flower buds (Campoy et

al., 2013; Gariglio et al., 2006; Hussain et al., 2015; Wall et al., 2008), contrary to that for native

species (Figure 2b). The smaller observed response in flowering for native species is associated

with longer phenological lags and accumulations of winter chilling (Table 1), indicatingg more

stressful environments later in the season or higher stress sensitivity with reproductive events.

Comparatively, plants starting growth early in the growing season may be less restricted by soil

moisture that often recharges from winter precipitation and is depleted with increased moisture

consumption over time (Sherry et al., 2007; Wolkovich et al., 2013; Zettlemoyer et al. 2019),

particularly in dry climates (Man and Greenway, 2013). Moisture stress can progressively

develop over the season (Piao et al. 2019; Wolkovich et al., 2013; Zettlemoyer et al., 2019) and

differentially affect exotic and native species with differing spring phenology (Stuble et al.,

2021; Willis et al., 2010; Wolkovich et al., 2013) The smaller observed responses in flowering of

native species suggests reproductive disadvantage (IPCC, 2014; Zettlemoyer et al., 2019). The

consistent leafing response, however, does not support suggestions that late-start native species

would have relatively shorter active growing seasons and, therefore, a competitive growth

disadvantages relative to early-start exotic species with climate warming (Fitter and Fitter, 2002;

Morin et al., 2009; Primack and Gallinat, 2016). Total thermal benefits from climate warming do

not differ among species with differing spring phenology (Chu et al., 2021). Given the projected

increases in temperature and decreases in precipitation for many parts of the world (IPCC, 2014),

future studies should assess the differences in drought sensitivity between vegetative and

reproductive events and among functional groups and climatic regions.

Boreal vs. temperate region

Spatial variations along altitudinal and latitudinal gradients are often reported but not well

understood (Ge et al., 2015; Morin et al., 2009; Parmesan, 2007; Zhang et al., 2015). Our

analysis indicates smaller observed responses in boreal region (MAT<6 °C) due to less forcing

changes (Tables 1), but not to longer phenological lag or chilling deficiency (Figure 2). The

similar spring warming but less forcing changes suggests that the temperature increases in boreal

region occur most frequently in the winter (Beaubien and Hamann, 2011; Shen et al., 2015)

when temperatures are below freezing and don't contribute to spring forcing (Man and Lu,

2010). The uneven warming has been reported in both natural and experimental settings

(Beaubien and Hamann, 2011; Prevéy et al., 2017; Shen et al.; 2015; Slaney et al., 2007; Yang et

al., 2020) and supports the claim that average temperature changes do not represent forcing

changes (Chu et al., 2021; Keenan et al., 2020; Wolkovich et al.; 2021). When forcing changes

are held constant, however, observed responses increase with altitude and latitude, as shown by

the positive regression coefficients in both leafing and flowering models (Table 2). The possible

mechanisms behind this may be lower growth temperature and less moisture stress at higher

altitude and latitude (Rafferty et al., 2020), as shown by the negative collinearity of altitude and

latitude with growth temperature and phenological lag in both leafing and flowering (data not

shown). Therefore, boreal region, depending on forcing changes, can have smaller (Ge et al.,

2015; Menzel et al., 2006; Rafferty et al., 2020; Shen et al., 2015) or greater (Ge et al., 2015;

Parmesan, 2007; Post et al., 2018; Prevéy et al., 2017) phenological responses, spatial trends that

may be different from those suggested by phenological sensitivity (Liu et al., 2019; Post et al.,

2018). Consequently, climate change may not result in general phenological convergence across

altitudes and latitudes (Rafferty et al., 2020; Shen et al., 2015; Tao et al., 2021), contrary to

suggestions by others for small scale studies (Prevéy et al., 2017; Vitasse et al., 2018; Ziello et

al., 2009).

Trees, shrubs, herbs, and grasses

Smaller observed responses in herb and grass leafing support findings from some early syntheses

(König et al, 2018; Parmesan, 2007), but not those that show a greater response from non-woody

plants (Ge et al., 2015; Post and Stenseth, 1999; Root et al., 2003) or no differences (Stuble et

al., 2021). Herbs and grasses often leaf out earlier than woody shrubs and trees (Badeck et al.,

2004; Heberling et al., 2019) and should have lower growth temperatures and therefore greater

phenological responses to climate warming (Chu et al., 2021; Prevéy et al., 2017; Wolkovich et

al., 2021). However, different growth forms are often studied separately on sites in different

climatic regions, making comparisons challenging. Furthermore, most studies on herb and grass

leafing are conducted in boreal region at high altitudes, low MAT, late spring phenology, low

forcing changes, and low spring warming (Table 1); the extremely small forcing changes resulted

in small observed responses despite cooler growth temperatures (Figure 2a). The observed

responses with herbs and grasses would not be smaller if converted to temperature sensitivity

(see Table 1 for large differences in spring warming for leafing). In contrast to leafing, herb and

grass flowering studies have greater forcing changes and warmer growth temperatures (Table 1),

which lowers both expected and observed responses (Table 1 note d) (Chu et al., 2021; Prevéy et

al., 2017; Wolkovich et al., 2021) and potentially induces moisture stress and increases

phenological lags (Ganjurjav et al., 2020; Huang et al., 2019; Moore et al., 2015; Post et al.,

2022), particularly for grasses (Sherry et al., 2007; Zettlemoyer et al., 2019) that have shallower

root systems (Kulmatiski and Beard, 2013; Schenk and Jackson, 2002).

Mechanistic assessment of changes in spring phenology

Averaged across all observational and experimental studies, observed responses and

phenological lags are both positive, suggesting that climate warming advances spring phenology

but not at rates expected from changes in spring temperatures. The positive lags are likely due to

more stressful environments with warmer and drier climate (Huang et al., 2019; Sherry et al.,

2007; Zettlemoyer et al., 2019). We quantified phenological lags from changes in spring

phenology and temperatures, an approach that does not require species-specific chilling or

forcing needs that are often unavailable (Fitter and Fitter, 2002; Morin et al., 2009) or variable in

methods of estimation across species and studies (Man and Lu, 2010; Ettinger et al., 2020; Zhang

et al., 2018). The relatively early stage of climate warming and association of longer

phenological lags with longer accumulation of winter chilling (Table 1) suggest that climate

warming is not likely to induce a general chilling shortage (Chu et al., 2023; Ettinger et al., 2020;

Tao et al., 2021; Yang et al., 2020), although incidental effects on individual species or in

particular conditions can occur (Fu et al., 2015). Similarly, the variable and minor effects of

photoperiod (Basler and Körner, 2012; Chuine et al., 2010; Ettinger et al., 2020; Zohner et al.,

2016) (insignificant spring phenology; see Table 2) often reported in studies with extreme

warming scenarios (Caffarra and Donnelly, 2011; Fu et al., 2019; Laube et al., 2014) or

compounded with that of growth temperatures (Chu et al., 2021), minor effects by nutrient

availability (Piao et al., 2019), or incidental spring freezing (Man et al., 2009; 2021b) are not

likely to cause the systemic differences in phenological lag between observational and

experimental studies or between native and exotic species. Phenological lag indicates the overall

lag effect and the need to investigate individual constraints.

Our analysis indicates that both quantity (forcing change) and rate (growth temperature)

of spring forcing accumulation strongly influence the magnitude of change in spring phenology,

consistent with the findings by Chu et al. (2021) that smaller phenological response with late

season species are largely due to their higher growth temperatures. In phenological research, the

role of growth temperature is not adequately recognized (Chu et al., 2021, 2023) and can be

confounded with progressive increases of phenological lag with greater chilling requirements or

insufficient chilling (Asse et al., 2018; Morin et al., 2009; Primack and Gallinat, 2016),

photoperiod constraints (Körner and Basler, 2010; Shen et al., 2014; Way and Montgomery,

2015), or moisture stress (Piao et al., 2019; Wolkovich et al., 2013; Zettlemoyer et al., 2019)

suggested for late season species. Compared to average spring temperature change that explained

<0.5% of the variation in observed responses (Table 2), forcing change quantifies climate

warming effect relevant to phenological change of individual species and reduces uneven

warming effects among species with different spring phenology (Chu et al., 2021) and areas in

different climatic regions (Post et al., 2018; Yang et al., 2020). Both forcing change and growth

temperature can be readily extracted from phenological and temperature data, as demonstrated in

this synthesis, but not examined in assessment of plant phenology changes in spring.

Conclusions

Differential phenological responses reported previously are mainly due to different forcing

changes, growth temperatures, and environmental stresses from warmer and drier climates. The

overall effects of insufficient winter chilling and photoperiod restriction are likely limited but

may change with continuous progress of climate warming and advance of spring phenology

(IPCC, 2014; Ettinger et al., 2020; Tao et al., 2021). Our method allows for mechanistic

partitioning of observed changes by drivers of spring phenology and helps to understand changes

in spring phenology and differences in plant phenological responses across different functional

groups or climatic regions.

Acknowledgements

This work was supported by Ontario Ministry of Natural Resources and Forestry (Canada),

Guangxi Normal University (China), Jilin Provincial Academy of Forestry Sciences (China),

Shanghai Botanical Garden (China), Research Institute of Subtropical Forestry (China), and

Lakehead University (Canada). Lisa Buse and Gillian Muir of Ontario Ministry of Natural

Resources and Forestry and two anonymous reviewers provided constructive suggestions for

improving earlier drafts of the manuscript. Phenological and temperature data provided by

China's National Earth System Science Data Center are greatly appreciated.

Author Contributions

R.M. and J.T. conceived the general idea of the method; Y.J., X.C., X.Y., R.M., and J.T.

compiled data; and Y.J., S.J.M., X.C., X.Y., R.M., J.T., and Q.L.D. contributed to the

interpretations of the results and to the writing of the paper.

Supporting Information

Temperature data and R codes for calculating forcing changes, expected responses, growth

temperature, spring warming, and phenological lags are deposited at Dryad

https://datadryad.org/stash/share/SGMComxqEbUCX1N-1BX1eAOMTSmoZFtN5YR_Mx0PPvE

REFERENCES

Asse D, Chuine I, Vitasse Y, Yoccoz NG, Delpierre N, Badeau V, … Randin CF (2018) Warmer
winters reduce the advance of tree spring phenology induced by warmer springs in the
Alps. *Agricultural* and Forest *Meteorology* **252**: 220–230.

Badeck FW, Bondeau A, Böttcher K, Doktor D, Lucht W, Schaber J, Sitch S (2004) Responses
of spring phenology to climate change. *New Phytologist* **162**: 295–309.

Basler D, Körner C (2012) Photoperiod sensitivity of bud burst in 14 temperate forest tree
species. *Agricultural* and Forest *Meteorology* **165**: 73–81.

Beaubien E, Hamann A (2011) Spring flowering response to climate change between 1936 and
2006 in Alberta, Canada. <u>*BioScience*</u> **61**: 514–524.

Burnham KP, Anderson DR (2002) *Model selection and multimodel inference: a practical
information-theoretic approach*, 2nd edn. Springer, New York.

Caffarra A, Donnelly A (2011) The ecological significance of phenology in four different tree
species: effects of light and temperature on bud burst. *International
Journal* of *Biometeorology* **55**: 711–721.

Calinger KM, Queenborough S, Curtis PS (2013) Herbarium specimens reveal the footprint of
climate change on flowering trends across north-central North America *Ecology Letters*
16: 1037–1044.

Campoy JA, Ruiz D, Nortes MD, Egea J (2013) Temperature efficiency for dormancy release in
apricot varies when applied at different amounts of chill accumulation *Plant Biology* **15**:
28–35.

Chen H, Zhu Q, Wu N, Wang Y, Peng CH (2011) Delayed spring phenology on the Tibetan

Plateau may also be attributable to other factors than winter and spring warming.

Proceedings of the National Academy of Sciences USA **108**: E93.

Chu X, Man R, Dang QL (2023) Spring phenology, phenological response, and growing season

length. *Frontiers in Forests and Global Change* **6**: 104136.

Chu X, Man R, Zhang H, Yuan W, Tao J, Dang QL **(**2021) Does climate warming favour early

season species? *Frontiers in Plant Science* **12**: 765351.

Chuine I, Morin X, Bugmann H (2010) Warming, photoperiods, and tree phenology. *Science*

329: 277–278.

Ettinger AK, Chuine I, Cook BI, Dukes JS, Ellison AM, Johnston MR, … Wolkovich EM (2019)

How do climate change experiments alter plot-scale climate? *Ecology Letters* **22**: 748–

763.

Ettinger AK, Chamberlain CJ, Morales-Castilla I, Buonaiuto DM, Flynn DFB, Savas T, …

Wolkovich EM (2020) Winter temperatures predominate in spring phenological

responses to warming. *Nature Climate Change* **10**: 1137–1142.

Fitter AH, Fitter RSR (2002) Rapid changes in flowering time in British plants. *Science* **296**:

1689–1691.

Fu YH, Piao S, Zhou X, Geng X, Hao F, Vitasse Y, Janssens IA (2019) Short photoperiod

reduces the temperature sensitivity of leaf-out in saplings of Fagus sylvatica but not in

horse chestnut. *Global Change Biology* **25**: 1696–1703.

Fu YH. Zhao H, Piao S, Peaucelle M, Peng S, Zhou G, . . . Janssens IA (2015) Declining global

warming effects on the phenology of spring leaf unfolding. *Nature* **526**: 104–107.

Ganjurjav H, Gornish ES, Hu G, Schwartz MW, Wan Y, Li Y, Gao Q (2020) Warming and

precipitation addition interact to affect plant spring phenology in alpine meadows on the

central Qinghai-Tibetan Plateau. *Agricultural and Forest Meteorology* **287**: 107943.

Gariglio N, Gonza´lez Rossia DE, Mendow M, Reig C, Agusti M (2006) Effect of artificial

chilling on the depth of endodormancy and vegetative and flower budbreak of peach and

nectarine cultivars using excised shoots. *Scientia Horticulturae* **108**: 371–377.

Ge Q, Wang H, Rutishauser T, Dai J (2015) Phenological response to climate change in China: a

meta-analysis. *Global Change Biology* 21: 265–274.

Heberling JM, McDonough MacKenzie C, Fridley JD, Kalisz S, Primack RB (2019)

Phenological mismatch with trees reduces wildflower carbon budgets. *Ecology Letters*

22: 616–623.

Huang W, Dai J, Wang W, Li J, Feng C, Du J (2020) Phenological changes in herbaceous plants

in China's grasslands and their responses to climate change: a meta-analysis.

International Journal of Biometeorology **64**: 1865–1876.

Huang W, Ge Q, Wang H, Dai J (2019) Effects of multiple climate change factors on the spring

phenology of herbaceous plants in Inner Mongolia, China: Evidence from ground

observation and controlled experiments. *International Journal* of *Climatology* **39**: 5140–

5153.

Hussain S, Niu Q, Yang F, Hussain N, Teng Y (2015) The possible role of chilling in floral and

vegetative bud dormancy release in *Pyrus pyrifolia*. *Biologia Plantarum* **59***: 726–734.

IPCC (2014) *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and*

III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change

[Core Writing Team, R. K. Pachauri and L. A. Meyer (eds.)]. IPCC, Geneva,

Switzerland, 151 pp.

Keenan TF, Richardson AD, Hufkens K (2020) On quantifying the apparent temperature

sensitivity of plant phenology. *New Phytologist* **225**: 1033–1040.

König P, Tautenhahn S, Cornelissen JHC, Kattge J, Bönisch G, Römermann C (2018) Advances

in flowering phenology across the Northern Hemisphere are explained by functional

traits. *Global Ecology* and *Biogeography* **27**: 310–321.

Körner C, Basler D (2010) Phenology under global warming. *Science* **327**: 1461–1462.

Kulmatiski A, Beard KH (2013) Root niche partitioning among grasses, saplings, and trees

measured using a tracer technique *Oecologia* **171**: 25–37.

Laube J, Sparks TH, Estrella N, Höfler J, Ankerst DP, Menzel A (2014) Chilling outweighs

photoperiod in preventing precocious spring development. *Global Change Biology* **20**:

170–182.

Liu Q, Piao S, Fu YH, Gao M, Peñuelas J, Janssens IA (2019) Climatic warming increases

spatial synchrony in spring vegetation phenology across the Northern Hemisphere.

Geophysical Research Letters **46***: 1641–1650.

Man R, Greenway K J (2013) Effects of soil moisture and species composition on growth and

productivity of trembling aspen and white spruce in planted mixtures: Five-year results

New Forests **44**: 23–38.

Ma Q, Huang JG, Hänninen H, Berninger F (2019) Divergent trends in the risk of spring frost

damage to trees in Europe with recent warming. *Global Change Biology* **25**: 351–360.

Man R, Kayahara GJ, Dang QL, Rice JA (2009) A case of severe frost damage prior to budbreak

in young conifers in Northeastern Ontario: consequence of climate change? *Forestry*

Chronicle **85**: 453–462.

Man R, Lu P (2010) Effects of thermal model and base temperature on estimates of thermal time

to bud break in white spruce seedlings. *Canadian Journal of Forest Research* **40**: 1815–

1820.

Man R, Lu P, Dang QL (2017a) Insufficient chilling effects vary among boreal tree species and

chilling duration. *Frontiers in Plant Science* **8**: 1354.

Man R, Lu P, Dang QL (2017b) Cold hardiness of white spruce, black spruce, jack pine, and

lodgepole pine needles during dehardening. *Canadian Journal of Forest Research* **47**:

1116–1122.

Man R, Lu P, Dang QL (2021a) Effects of insufficient chilling on budburst and growth of six

temperate forest tree species in Ontario. *New Forests* **52**: 303–315.

Man R, Lu P, Dang QL (2021b) Cold tolerance of black spruce, white spruce, jack pine, and

lodgepole pine seedlings at different stages of spring dehardening. *New Forests* **52**: 317–

328.

Marion GM, Henry GHR, Freckman DW, Johnstone J, Jones G, Jones MH, …Virginia RA

(1997) Open-top designs for manipulating field temperature in high latitude ecosystems.

Global Change Biology **3***: 20–32.

Menzel A, Sparks T, Estrella N, Koch E, Aasa A, Ahas R, ... Zust A (2006) European

phenological response to climate change matches the warming pattern. *Global Change*

Biology **12**: 1969–1976.

Moore LM, Lauenroth WK, Bell DM, Schlaepfer DR (2015) Soil water and temperature explain

canopy phenology and onset of spring in a semiarid steppe. *Great Plains Research* **25**:

121–138.

Morin X, Lechowicz MJ, Augspurger C, O'Keefe J, Viner D, Chuine I (2009) Leaf phenology

in 22 American tree species during the 21st century. *Global Change Biology* **15**: 961–

975.

O'Dea RE, Lagisz M, Jennions MD, Koricheva J, Noble DW, Parker TH, Gurevitch J, Page MJ,

Stewart G, Moher D, Nakagawa S (2021) Preferred reporting items for systematic

reviews and meta-analyses in ecology and evolutionary biology: a PRISMA extension.

Biological Reviews **96**: 1695–722.

Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, Shamseer L,

Tetzlaff JM, Akl EA, Brennan SE, Chou R (2021) The PRISMA 2020 statement: an

updated guideline for reporting systematic reviews. *British Medical Journal* **372**: n71.

Parmesan C (2007) Influences of species, latitudes and methodologies on estimates of

phenological response to global warming. *Global Change Biology* **13**: 1860–1872.

Piao S, Liu Q, Chen A, Janssens IA, Fu Y, Dai J, ... Zhu X (2019) Plant phenology and global

climate change: Current progresses and challenges. *Global Change Biology* **25**: 1922–

1940.

Polgar C, Gallinat A, Primack RB (2014) Drivers of leaf-out phenology and their implications

for species invasions: insights from Thoreau's Concord. *New Phytologist* **202**: 106–115.

Post AK, Hufkens K, Richardson AD (2022) Predicting spring green-up across diverse North

American grasslands. *Agricultural* and Forest *Meteorology* **327**: 109204.

Post E, Stenseth NC (1999) Climatic variability, plant phenology, and northern ungulates

Ecology **80**: 1322–1339.

Post E, Steinman BA, Mann ME (2018) Acceleration of phenological advance and warming with

latitude over the past century. *Scientific Reports* **8**: 3927.

Prevéy JS, Rixen C, Rüger N, Høye TT, Bjorkman AD, Myers-Smith IH, …Wipf S (2019)

Warming shortens flowering seasons of tundra plant communities. *Nature Ecology and*

Evolution **3**: 45–52.

Prevéy J, Vellend M, Rüger N, Hollister RD, Bjorkman AD, Myers-Smith IH, … Rixen C

(2017) Greater temperature sensitivity of plant phenology at colder sites: implications for

convergence across northern latitudes. *Global Change Biology* **23**: 2660–2671.

Primack RB, Gallinat AS (2016) Spring budburst in a changing climate. *American Scientist* **104**:

102–109.

Rafferty NE, Diez JM, Bertelsen CD (2020) Changing climate drives divergent and nonlinear

shifts in flowering phenology across elevations. *Current Biology* **30**: 432–441.

Root TL, Price JT, Hall KR, Schneider SH, Rosenzweig C, Pounds JA (2003) Fingerprints of

global warming on wild animals and plants. *Nature* **421**: 57–60.

Schenk HJ, Jackson RB (2002) Rooting depths, lateral root spreads and below-ground/above-

ground allometries of plants in water-limited ecosystems *Journal of Ecology* **90**: 480–

494.

Shen M, Cong N, Cao, R (2015) Temperature sensitivity as an explanation of the latitudinal

pattern of green-up date trend in Northern Hemisphere vegetation during 1982–2008

International Journal of *Climatology* **35**: 3707–3712.

Shen M, Tang Y, Chen J, Yang X, Wang C, Cui X, … Cong N (2014) Earlier-season vegetation

has greater temperature sensitivity of spring phenology in northern hemisphere. *PLOS*

ONE **9**: e88178.

Sherry RA, Zhou X, Gu S, Arnone III JA, Schimel DS, Verburg PS, … Luo Y (2007)

Divergence of reproductive phenology under climate warming. *Proceedings of the*

National Academy of Sciences USA **104**: 198–202.

Slaney M, Wallin G, Medhurst J, Linder S (2007) Impact of elevated carbon dioxide

concentration and temperature on bud burst and shoot growth of boreal Norway spruce.

Tree Physiology **27**: 301–312.

Stuble KL, Bennion LD, Kuebbing SE (2021) Plant phenological responses to experimental

warming—A synthesis. *Global Change Biology* **27**: 4110–4124.

Tao J, Man R, Dang QL (2021) Earlier and more variable spring phenology projected for eastern

Canadian boreal and temperate forests with climate warming. *Trees, Forests and People*

6: 100127.

Vitasse Y, Signarbieux C, Fu YH (2018) Global warming leads to more uniform spring

phenology across elevations. *Proceedings of the National Academy of Sciences USA* **115***:*

1004–1008.

Wall C, Dozier W, Ebel RC, Wilkins B, Woods F, Foshee III W (2008) Vegetative and floral

chilling requirements of four new kiwi cultivars of *Actinidia chinensis* and *A. deliciosa*.

HortScience **43**: 644–647.

Way DA, Montgomery RA (2015) Photoperiod constraints on tree phenology, performance and

migration in a warming world. *Plant, Cell and Environment* **38**: 1725–1736.

Willis CG, Ruhfel BR, Primack RB, Miller-Rushing AJ, Losos JB, Davis CC (2010) Favorable

climate change response explains non-native species' success in Thoreau's woods. *PLOS*

ONE **5**: e8878.

Wolkovich EM, Auerbach J, Chamberlain CJ, Buonaiuto DM, Ettinger AK, Morales-Castilla I,

Gelman A (2021) A simple explanation for declining temperature sensitivity with

warming. *Global Change Biology* **27**: 4947–4949.

Wolkovich EM, Cook BI, Allen JM, Crimmins TM, Betancourt JL, Travers SE, … Cleland EE

(2012) Warming experiments underpredict plant phenological responses to climate

change. *Nature* **485***: 494–497.

Wolkovich EM, Davies TJ, Schaefer H, Cleland EE, Cook BI, Travers SE, … Davis CC (2013)

Temperature-dependent shifts in phenology contribute to the success of exotic species

with climate change. *American Journal of Botany* **100**: 1407–1421.

Yang Y, Wu Z, Guo L, He HS, Ling Y, Wang L, … Li MH (2020) Effects of winter chilling vs.

spring forcing on the spring phenology of trees in a cold region and a warmer reference

region. *Science of the Total Environment* **725**: 138323.

Zettlemoyer MA, Schultheis EH, Lau JA (2019) Phenology in a warming world: differences

between native and non-native plant species. *Ecology Letters* **22**: 1253–1263.

Zhang H, Liu S, Regnier P, Yuan W (2018) New insights on plant phenological response to

temperature revealed from long-term widespread observations in China. *Global Change*

Biology **24**: 2066–2078.

Zhang H, Yuan W, Liu S, Dong W (2015) Divergent responses of leaf phenology to changing

temperature among plant species and geographical regions. *Ecosphere* **6**: 250.

Ziello C, Estrella N, Kostova M, Koch E, Menzel A (2009) Influence of altitude on phenology of

selected plant species in the Alpine region (1971–2000). *Climate Research* **39**: *227*–234.

Zohner CM, Mo L, Renner SS, Svenning JC, Vitasse Y, Benito BM, … Crowther TW (2020)

Late-spring frost risk between 1959 and 2017 decreased in North America but increased

in Europe and Asia. *Proceedings of the National Academy of Sciences USA* **117**: 12192–

12200.

Zohner CM, Renner SS (2014) Common garden comparison of the leaf-out phenology of woody

species from different native climates, combined with herbarium records, forecasts long-

term change *Ecology Letters* **17**: 1016–1025.

Table 1. Mean values of climatic, phenological, and biological variables by species event (leafing and flowering), research approach (observation and experiment), species origin (native and exotic), climatic region (boreal and temperate), and growth form (tree, shrub, herb, and grass). Growth temperature=temperatures at species event with warmer climate, forcing change =warming-induced changes in spring forcing experienced by individual species, spring warming=average spring temperature changes, and phenological lag= phenological difference due to effects of phenological constraints.

Species event	Spring phenology (Julian)	Growth temperature (°C)	Altitude (m)	Latitude (°N)	MAT (°C)	Forcing change (degree days)	Spring warming (°C)	Phenological lag (day)	Sample size (N)
Leafing	111	11.1	687	42.8	7.8	85	1.3	1.8	447
Observation	106	11.0	487	42.6	8.4	79	1.1	1.2	359
Experiment	130	11.2	1500	43.8	5.6	106	1.8	4.1	88
Native	113	11.0	744	43.3	7.4	85	1.3	1.9	379
Exotic	100	11.7	366	40.0	10.2	78	1.2	1.1	68
Boreal	128	10.2	1147	45.7	2.0	65	1.3	2.4	154
Temperate	102	11.5	444	41.3	10.9	94	1.3	1.5	293
Tree	105	11.7	364	41.9	10.1	101	1.4	1.9	246
Shrub	109	10.9	433	43.5	7.8	80	1.4	1.1	107
Herb	131	10.0	1955	43.3	1.8	51	0.8	1.2	54
Grass	123	8.9	1637	45.9	2.1	37	0.7	3.6	40
Flowering	138	13.7	534	45.5	9.0	126	1.3	3.5	1080
Observation	136	13.4	454	46.0	9.2	120	1.2	3.3	956
Experiment	154	16.7	1153	41.8	7.3	174	1.8	4.7	124
Native	142	13.8	617	45.9	8.5	124	1.3	3.7	850
Exotic	125	13.5	229	44.3	10.9	135	1.4	2.4	230
Boreal	167	14.6	1998	42.9	3.3	109	1.4	1.3	175
Temperate	133	13.6	251	46.0	10.1	130	1.3	3.9	905

Tree	114	12.7	289	43.5	10.5	119	1.5	1.4	183
Shrub	125	12.3	446	44.8	8.5	103	1.3	2.9	148
Herb	146	14.2	608	46.1	8.7	129	1.4	3.8	659
Grass	150	15.3	636	46.5	8.9	160	1.4	5.8	90

Note: effects of forcing change and growth temperature illustrated with equation 3 in Materials and Methods

a) the difference in forcing change between boreal and temperate regions for leafing is 29 (94 – 65), equivalent to 2.8 day expected responses under the average growth temperature of boreal region (10.2). Similarly, the difference in flowering is equivalent to 1.5 day expected responses, i.e., 21/14.6.

b) the difference in forcing change between grasses and trees + shrubs for leafing is 53 ((101+80)/2 – 37), equivalent to 6.0 day expected responses under the average growth temperature of grasses (8.9).

c) the difference in forcing change between experimental and observational studies for leafing is 27 (106 –79), equivalent to 2.5 day expected responses under the average growth temperature of observational studies (11.0).

d) different growth temperatures explain 2.6 day difference in flowering between experimental and observational studies, i.e., 174/13.4 – 174/16.7, and 1.8 day difference in flowering between woody (trees and shrubs) and non-woody (herbs and grasses) plants, i.e., (129+160)/(12.7+12.3) – (129+160)/(14.2+15.3).

Table 2. Final stepwise regression coefficients and variable influence on observed phenological responses by Akaike information criterion (AIC).

Species event	Intercept	Growth temperature	Altitude	Latitude	MAT	Forcing change	Spring warming	Phenological lag	R-squared Sample (N)
Leafing	2.6439	-0.6075	0.0003	0.0732	0.1480	0.0736	0.9463	-0.9738	0.9453
AIC (%)	449 (27.7)	340 (20.9)	2 (0.1)	15 (0.9)	11 (0.7)	68 (4.2)	7 (0.4)	732 (45.1)	(447)
Flowering	4.521	-0.5917	0.0003	0.0869	0.1466	0.0521	0.7620	-0.9893	0.9653
AIC (%)	785 (18.1)	247 (5.7)	7 (0.2)	20 (0.5)	32 (0.7)	826 (19.1)	5 (0.1)	2409 (55.6)	(1080)

Note: Growth temperature (°C)=temperatures at species event (leafing or flowering) with warmer climate

MAT (°C)=long-term mean annual temperature

Forcing change = warming-induced changes in spring forcing at species events

Spring warming (°C)=average spring temperature change (Julian day 1 to 182)

Phenological lag (days)= phenological difference due to effects of phenological constraints

Spring phenology (date of species event with baseline climate; Julian days) and long-term mean precipitation (MAP, mm) were excluded from final models due to lack of significance

----

Forcing change, growth temperature, and phenological lag are calculated from phenological and temperature data for individual species, spring phenology and observed responses are provided in the original studies, and spring warming and long-term MAT and MAP are provided in the original studies, calculated from baseline temperatures, or extracted from Google Maps (www.maps.google.com), FreeMapTools (www.freemaptools.com), or WorldClim Database (Fick and Hijmans, 2017)

Figure 1. A diagram showing the relationships among forcing change (F_C, difference in spring forcing (degree days) between baseline $\sum_1^i T_{Ci}$ and warmer $\sum_1^i T_{Wi}$ temperatures at species event with baseline temperatures O_{Ci}), expected response (N_E, difference between climatic conditions of baseline and warmer temperatures in reaching species forcing needs with baseline temperatures, i.e., spring forcing at O_{Ci}), growth temperature (F_C/N_E, average temperature at species event with warmer temperatures), and phenological lag (N_C, difference between expected N_E and observed N_O responses with warmer temperatures).

Figure 2. Observed responses and phenological lags (least square means ± standard errors) in leafing and flowering (left and right set of bars in each pair, respectively) by research approach (observation and experiment), species origin (native and exotic), climatic region (boreal and temperate), and growth form (tree, shrub, herb, and grass) extracted from reported plant phenological changes in spring. Phenological lags are calculated from the differences between observed responses and those expected from species-specific changes in spring temperatures. Different letters indicate means differ significantly ($p<0.05$).

Supplementary data

Table S1 | A list of references for all studies included

Table S2 | A summary of study information

Table S3 | MetaData containing all information used in the analysis

Table S4 | Sources of weather data