Identifying a novel mechanism of L-leucine uptake in *Mycobacterium tuberculosis* using a chemical genomic approach

Nisheeth Agarwal*[1], Himanshu Gogoi[1], Eeba[1], Linus Augustin[1], Md. Younus Khan[1], Yashwant Kumar[1], Sayan Kumar Bhowmick[2] and Bappaditya Dey[2]

[1]Translational Health Science and Technology Institute, NCR Biotech Science Cluster, 3rd Milestone, Faridabad–Gurgaon Expressway, Faridabad- 121001 (Haryana), India

[2]National Institute of Animal Biotechnology (NIAB), Gachibowli, Hyderabad- 500032 (Telangana), India

***Corresponding author:** Nisheeth Agarwal

E-mail: nisheeth@thsti.res.in

Running Title: L-leucine uptake in *Mycobacterium tuberculosis*

Keywords: *Mycobacterium tuberculosis*; Drug-repurposing, Semapimod, Amino-acid uptake, PDIM

Abstract

Amino acid biosynthesis is vital for *Mycobacterium tuberculosis* (Mtb) proliferation and

tuberculosis pathogenesis. However, it is not clear how amino acids are transported in Mtb,

particularly the branched chain amino acids (BCAAs) that contribute to the production of the

cell-wall lipid component precursors such as acetyl-CoA and propionyl-CoA.

While performing the screening of an FDA-approved repurposed library of small molecule

inhibitors against the auxotrophic strain Mtb mc^2 6206, which lacks *leuC-leuD* and *panC-*

panD genes, we identified a molecule namely semapimod, which exclusively inhibits growth

of the auxotrophic strain, whereas no effect is observed against the wild-type Mtb H$_{37}$Rv.

Interestingly, 24 h of exposure of Mtb mc^2 6206 to semapimod causes massive transcriptional

reprogramming with differential expression of >450 genes associated with a myriad of

metabolic activities. By performing a series of experiments, we affirm that semapimod indeed

inhibits the L-leucine uptake in Mtb mc^2 6206 by targeting a protein involved in the cell-wall

lipid biosynthesis pathway. Remarkably, semapimod treatment of mice infected with Mtb

H$_{37}$Rv causes a significant reduction of bacterial load in lungs and spleen, despite showing no

efficacy against the pathogenic strain *in vitro*.

Overall findings of our study reveal that together with an endogenous pathway for L-leucine

biosynthesis, a well-orchestrated machinery for its uptake is functional in Mtb which is

important for intracellular survival of the TB pathogen.

Introduction

Tuberculosis (TB), caused by the pathogen *Mycobacterium tuberculosis* (Mtb) is the leading

cause of death due to microbial infections. As per the recent estimates, approximately 25% of

the global population is asymptomatically infected with Mtb and about 11 million people

developed the disease annually resulting in the loss of ~1.25 million lives. In 2023, eight

countries accounted for more than two thirds of the global total burden, with India leading the

count[1].

TB is curable and the usual treatment of pulmonary TB (PTB) cases involves 4 antibiotics

(isoniazid (H), rifampicin(R), pyrazinamide (Z) and ethambutol (E)) for the first 2 months

followed by four months of treatment with R and H[1, 2, 3, 4]. Unfortunately, due to multiple

reasons such as delayed or mis-diagnosis, inadequate availability of drugs in resource-limited

settings, poor administration, and noncompliance to the prolonged toxic drug regimen, TB

becomes resistant to these drugs[5, 6, 7]. The drug-resistant TB (DR-TB) is classified under five

categories: H-resistant TB[8], R-resistant TB (RR-TB)[9], multidrug-resistant TB (MDR-TB)

which is resistant to both H and R, pre-extensively drug-resistant TB (pre-XDR-TB) which is

resistant to H, R and any fluoroquinolone or any of the second-line injectables such as

amikacin, capreomycin, and kanamycin, and XDR-TB which is resistant to H, R, any

fluoroquinolone, and a second-line injectable or at least one of the recently developed drugs

bedaquiline and linezolid. Treatment of DR-TB involves lengthy treatment with a course of

second-line drugs for 6-18 months, causing huge socioeconomic burden[10]. DR-TB continues

to be a threat to the public health system as number of people developing MDR/RR-TB

remains stable since 2020. Globally, ~400,000 MDR/RR-TB cases were reported in 2023,

with about one fourth of the global MDR-TB cases reported from India alone[1, 11].

Considering the grim situation of everlasting DR-TB cases and emergence of resistance to

newly developed drugs, there is a pressing need to explore novel therapeutic approaches.

With advancements in the medicinal chemistry, large numbers of new chemical entities

(NCEs) have been synthesized over the last three decades that were evaluated for

antimicrobial activity by the target-based and the phenotypic screening approaches[12, 13].

However, only a few of the newly developed antibiotics that got approval for human use

exhibit a novel mechanism of action (MoA), whereas majority are the derivatives of the

existing antibiotic classes[12, 14]. Evaluation of the antibiotic potential of the existing drugs that

are approved for human use by the Federal Drug Agency (FDA) for other diseases, also

known as drug repurposing, has emerged as an alternative approach to expedite treatment of

infectious diseases including TB, as it avoids much of the hurdles at the early phases of drug

development[12, 15].

Herein, we report that a new anti-inflammatory molecule– semapimod, identified from a

screen of the repurposed library of FDA-approved drugs, kills Mtb mc^2 6206 (a leucine and

pantothenate auxotroph) at a reasonably low concentration of 20nM. To our great surprise,

semapimod fails to inhibit other mycobacterial species including pathogenic Mtb H$_{37}$Rv.

These unexpected findings led us to identify a functional L-leucine uptake system in Mtb,

which is blocked by the newly-identified compound. We show that semapimod targets an

enzyme, PpsB, involved in the synthesis of the cell-wall phthiocerol dimycocerosate (PDIM),

and influences Mtb virulence during host infection.

Results

Screening of FDA-approved library of molecules against Mtb mc^2 6206.

In this study we screened a repurposed library (MedChemExpress) consisting of 3614 FDA-approved small molecule inhibitors against Mtb mc^2 6206, which is commonly used as a surrogate strain for pathogenic Mtb H$_{37}$Rv[16]. These molecules primarily target microbial infections (790), GPCR/G Protein (646), apoptosis (371), metabolic enzyme (including protease) (355), membrane transporter (236), autophagy (210), cell cycle/DNA damage (157), immunology/inflammation (109), protein tyrosine kinase/RTK (99) and a large number of molecules targeting other miscellaneous pathways (402) (Supplementary Fig. 1a). The library contains a versatile set of molecules targeting different diseases such as cancer, cardiovascular disease, endocrinology, infection, inflammation/immunology, metabolic disease and neurological disease (Supplementary Fig. 1b). While these are intended for human use, 66% of the molecules have recently been launched, whereas remaining inhibitors are at different phases of trial, as depicted in Supplementary Fig. 1c. Of these, we identified 422 molecules showing complete inhibition of growth at 50μM concentration. As anticipated, majority (158) of the Mtb-inhibitors show anti-infection activity. However, several other compounds acting on host metabolic pathways, such as GPCR/G protein (68), apoptosis (45), autophagy (24), cell cycle/DNA damage (18), metabolic enzyme/protease (18), protein tyrosine kinase/RTK (17), JAK/STAT signaling (14) and membrane transporter (13), substantially inhibit *in vitro* growth of Mtb mc^2 6206 (Fig. 1a).

Next, we determined minimum concentration of newly identified anti-mycobacterial compounds at which ~90% growth is suppressed (MIC$_{90}$), by using plate-based resazurin assay. Our results reveal 300 molecules showing an MIC$_{90}$ of $\geq$12.5 μM, whereas remaining 122 molecules exhibit MIC$_{90}$ values ranging from 6.25 to $\leq$0.025μM (Fig. 1b). Surprisingly, out of the eight molecules which inhibit growth at <0.025μM, we find an anti-inflammatory molecule namely semapimod which demonstrates anti-microbial activity against Mtb mc^2 6206. Notably, semapimod is an inhibitor of proinflammatory cytokine production, which suppresses the production of TNF-α, IL-1β, and IL-6[17, 18, 19, 20], and its anti-mycobacterial activity has never been reported to the best of our knowledge.

Semapimod shows bactericidal activity against Mtb mc^2 6206 under extra- and intracellular growth conditions.

To corroborate the findings from initial screen, we tested the effect of the fresh batch of semapimod against Mtb mc^2 6206, which reveals complete inhibition of mycobacterial

growth at ~15nM (Fig. 1c). The colony forming unit (CFU) estimation further shows a
bactericidal activity of this molecule which causes 88% reduction of bacterial viability on day
2 and >99% reduction after 5 days of incubation (Fig. 1d). In addition to its effect on *in vitro*
growth, we also find a substantial reduction in viable bacterial counts during macrophage
infection. Incubation of Mtb mc^2 6206-infected THP1 macrophages with different doses of
semapimod reveals 60% reduction (p <0.005) in intracellular CFU counts after 6 days of
incubation with 1μM inhibitor (Fig. 1e). Notably, no cytotoxic effect was observed at this
concentration against THP1, thus ruling out the possibility of cell lysis by semapimod.

Global transcriptional profile of Mtb in response to semapimod treatment
To gain mechanistic insights into the effect of semapimod on Mtb mc^2 6206, we analysed the
global transcriptional profile of the bacterium in response to this drug. Bacterial cultures in
the logarithmic growth phase (OD600 of ~0.4) were treated with 50nM semapimod,
equivalent to ~3x MIC, and total RNAs were harvested from semapimod-treated and
untreated bacteria after 24 h of incubation, as described in the Materials and Methods. RNA
samples from three biological replicates were subsequently used in RNA sequencing
(RNASeq), which reveals differential expression of 482 genes by ≥2.0-fold (false discovery
rate (FDR)-adjusted p value of ≤ 0.01). While 251 genes are downregulated by semapimod
treatment, expression of 231 genes is induced (Supplementary Dataset 1 and Fig. 2a).
Remarkably, the effect of semapimod was consistent across three biological replicates (Fig.
2b). Genes, prominently affected by semapimod, primarily belong to intermediary
metabolism and respiration (n=124), cell-wall and cell processes (n=98), lipid metabolism
(n=50) and those encoding for regulatory proteins (n=31) (Fig. 2c). Expression of some of the
representative genes was also validated by qRT-PCR using gene-specific primers, which
shows a similar pattern of expression (Supplementary Fig. 2), thus confirming the specificity
of the RNASeq data.
Our results show that genes involved in oxidative phosphorylation such as *atpF*, *atpH*, *atpG*,
atpD, *cydB*, *cydA*, *Rv1812c* and *qcrC* as well as several *nu*o operon genes (*nuoB*, *nuoC*,
nuoD, *nuoE*, *nuoF*, *nuoG*, *nuoI*, *nuoJ*, *nuoL*, *nuoM* and *nuoN*) are downregulated in the drug-
treated bacteria. Not only expression level of these respiratory genes is perturbed, but also a
significant reduction in the cellular ATP levels is observed upon semapimod treatment of Mtb
mc^2 6206 (Supplementary Fig. 3). Furthermore, we find several genes that encode for
subunits of RNA polymerase (*rpoA* and *rpoB*) and ribosome (*rplJ*, *rplL*, *rplC*, *rplD*, *rplW*,
rplB, *rpmC*, *rpsN1*, *rpsH*, *rplF*, *rplR*, *rpmD*, *rplT* and *rpsD*) exhibiting distinct upregulation

in response to the drug treatment, which indicate perturbation of transcription and translation
machineries in the drug-treated bacteria. Notably, semapimod treatment also interferes with
expression of a number of genes associated with metabolism of amino acids such as
branched-chain amino acids (*ilvC, ilvB1, ilvN, leuA, bkdB, bkdA, Rv2499c, accA1* and
accD1), cysteine (*csd, cysM, cysO, cysE, cysK1* and *sahH*), lysine (*pcd, lat, lysA*), methionine
(*metZ, metK* and *metH*), threonine (*thrB, thrC* and *thrA*), glutamate (*gdh*), glutamine (*glnA1*)
and tryptophan (*trpA*).

Among the cell-wall and cell processes functional category, we find a substantial reduction in
the expression of genes involved in peptidoglycan biosynthesis (*pbpA, ponA1, murI, dacB1*),
cell-wall arabinogalactan linker formation (*wbbL1*), cell-wall arabinan biosynthesis (*embB*
and *aftB*), cell-wall LPS biosynthesis (*Rv0225, gca, gmhA, gmhB* and *hddA*), cell division
(*Rv1708, scpA, ftsZ* and *gid*) and transport across the bacterial cell membrane (*ctpB, glnH,*
ctpE, pstA1, pstA2, narK2, Rv1747, ansP1, fecB and *Rv3200c*).

Another category of genes exhibiting notable change in expression are those involved in the
biosynthesis of lipids such as cell-wall mycolic acid (*accD6, fabD, kasA, kasB, fabG4,*
accD4, umaA, and *cmaA2*), methyl-branched lipids (*ppsB, ppsC, ppsD, mas, fadD26, fadD29,*
tesB1, Rv2953, papA1, pks2, mmpL8, fadD23, pks3, pks4, papA3 and *fadD21*), phospholipids
(*cdh*) and triglycerides (*desA3*). In addition, several genes controlling lipid degradation
(*fadE5, fadE35, fadD12, fadE19, tesB2, fadE21, fadE24, fadB2, fadD36, fadE15* and *fadD9*)
and transport (*lprK, mce1F, mce4F, mce4D, lprN,* and *lucA*) are also modulated in the drug-
treated bacteria. Noteworthy to mention, a marked reduction in expression is observed for
genes involved in synthesis of succinyl-CoA *via* vitamin B12-dependent methylmalonyl
pathway (MMP; *accA3, mutA, mutB, cobB* and *cobO*), whereas those associated with
methylcitrate cycle (MCC; *prpD, prpC, icl1* and *pks11*) exhibit significant upregulation,
together indicating perturbation of acetyl CoA levels in semapimod-treated Mtb mc^2 6206.

Furthermore, we find that semapimod treatment modulates expression of genes encoding a
variety of regulatory proteins such as kinases (*pknA, pknG* and *pknD*), WhiB-family
transcription regulators (*whiB2, whiB3* and *whiB7*), hypoxia regulators (*Rv0081, dosS* and
dosR), regulators of lipid transport (*Rv0302* and *mce1R*), a regulator of MCC genes (*regX3*)
and a leucine-responsive transcription regulator (LrpA).

Semapimod treatment interferes with L-leucine uptake in Mtb
To validate the effect of semapimod on the virulent Mtb, we sought to determine its MIC
against Mtb H$_{37}$Rv. Surprisingly, the drug was found ineffective against virulent Mtb strain.

Similar to virulent Mtb strain, no effect was observed against other mycobacterial species
such as *M. bovis* BCG, *M. abscessus* and *M. smegmatis* (Fig. 3a). These results pointed out
the effect of semapimod on uptake of leucine and/or pantothenate that are indispensable for
growth of the auxotrophic strain. Altered expression of a set of LrpA-regulated genes[21, 22, 23]
by semapimod treatment indicates perturbation of leucine uptake. To test this hypothesis, we
cultured Mtb mc^2 6206 in the absence of pantothenate and L-leucine (PL-), pantothenate (P-)
or L-leucine (L-) for 24 h followed by analysis of the expression of some of the leucine-
responsive genes such as *lat*, *leuA*, *lrpA* and *icl1* as surrogate markers. Interestingly, all of
these genes exhibit marked increase in expression under PL- and L-, but not under P-
conditions (Supplementary Fig. 4). A similar expression profile of these genes is observed in
semapimod-treated bacteria by RNASeq, thus suggesting the interference of L-leucine
uptake. To corroborate these findings, the intracellular leucine concentrations were estimated
in the untreated and semapimod-treated Mtb mc^2 6206 by mass-spectrometry. Our results
show ~45% reduction ($p < 0.05$) in the leucine levels in bacteria exposed to semapimod (Fig.
3b). Next, we examined the effect of semapimod against Mtb mc^2 6206, expressing *leuC-*
leuD and *panC-panD*, respectively, under a constitutive promoter. Remarkably, expression of
leuC-leuD, and not of *panC-panD* genes, alleviates the growth-inhibitory effect of
semapimod (Fig. 3c-e). Mtb mc^2 6206::*leuCD* exhibits comparable growth in the presence
and the absence of semapimod (Fig. 3c). In addition, Mtb mc^2 6206::*leuCD* exhibits
resistance to killing by as high as 3.37 μM (>200-fold excess of MIC) semapimod (Fig. 3d-
e). In contrast, Mtb mc^2 6206::*panCD* exhibits a similar level of inhibition by semapimod, as
observed with the Mtb mc^2 6206. Overall, these results substantiate that semapimod inhibits
L-leucine uptake in Mtb.

Generation and characterization of semapimod resistant strain of Mtb mc^2 6206
Semapimod-mediated inhibition of L-leucine uptake suggests the presence of a committed L-
leucine transport machinery in mycobacteria. While a number of putative L-leucine
transporter genes are found in *M. smegmatis* (Supplementary Fig. 5), none of their
orthologues are present in Mtb.
To gain insights into the potential mechanism of L-leucine transport in Mtb, we generated the
semapimod resistant (Sem^R) strain of Mtb mc^2 6206 by repeated passaging in the presence of
drug (Fig. 4a). *In vitro* growth analysis reveals a significantly increased growth of the two
independent Sem^R strains (Fig. 4b). While, the wild-type bacteria do not grow beyond OD$_{600}$
of 2.0, both the Sem^R strains achieve an OD$_{600}$ of ~4.0 after 10 days of culturing in 7H9-

OADS medium supplemented with L-leucine and pantothenate (Fig. 4b). The semapimod
resistance does not lead to any change in susceptibility to standard TB drugs such as
rifampicin or isoniazid (Supplementary Fig. 6), however, a marked increase in susceptibility
to vancomycin is observed in the Sem^R bacteria. While, vancomycin inhibits Sem^R with the
IC$_{50}$ of ~7μg/ml, no effect is observed for this antibiotic against the wild-type Mtb mc^2 6206
(Fig. 4c). It has been reported earlier that bacteria lacking PDIM grow faster and exhibit more
susceptibility to vancomycin than the wild-type Mtb with intact PDIM which provides barrier
to nutrients as well as large molecules such as vancomycin[24]. Interestingly, whole genome
sequence analysis of the Sem^R strain reveals Ile896Asn mutation in PpsB, involved in PDIM
synthesis (Fig. 4d). In addition, three other genes show altered sequence in Sem^R namely,
mce1D, *Rv3837c* and *ppe60*. While *mce1D* accumulates a nonsense mutation yielding a stop
codon at 225th position, sequence alteration in *Rv3837c* is at the extreme 3' end causing Thr
→ Ile substitution at the 228th position of the 332 amino acid long polypeptide. Furthermore,
multiple mutations were observed exclusively in *ppe60* (Supplementary Fig. 7), despite the
presence of two other paralogues of this gene (*ppe18* and *ppe19*), suggesting random
accumulation of mutations in *ppe60* during passaging. These observations led us to
investigate the association of *ppsB, mce1D* and *ppe60* in semapimod resistance, whereas, the
possibility of involvement of *Rv3837c* is unlikely.

Overexpression of *ppsB* alters susceptibility of semapimod^R strain to drugs
To further identify genes that might contribute to semapimod resistance, we overexpressed
ppsB, mce1D and *ppe60* genes in the wild-type and Sem^R strains of Mtb mc^2 6206 under the
constitutive promote. Notably, susceptibility to vancomycin and semapimod was completely
reversed in Sem^R strain upon overexpression of *ppsB*, whereas no effect of *ppsB*
overexpression is observed in the wild-type bacteria (Fig. 5). Sem^R strain, which is
susceptible to vancomycin, becomes resistant akin to the wild-type strain by overexpression
of *ppsB* (Fig. 5a).   Similarly, susceptibility to semapimod is restored in Sem^R by
overexpressing *ppsB* (Fig. 5b). In contrast, no effect of *mce1D* or *ppe60* overexpression is
observed on bacterial sensitivity to semapimod (Supplementary Figs. 8 and 9).

Semapimod targets PpsB of *M. tuberculosis*
The above results strongly suggest that the cell-wall PDIM biosynthesis protein PpsB is
involved in semapimod-mediated interference of L-leucine uptake in Mtb. To corroborate our
findings, we sought to examine if PpsB of Mtb is directly targeted by semapimod. The

4617bp *ppsB* open reading frame of Mtb was cloned in an *E. coli* expression plasmid, pET28,

for overexpression and purification of 6x His-tagged PpsB of Mtb. Analysis of purified

protein by SDS-PAGE reveals near-homogeneous preparation of the protein migrating at its

predicted molecular mass of ~170kDa (Supplementary Fig. 10). The purified 6x His-PpsB

was then subjected to interaction with different concentrations of semapimod using Biolayer

Interferometry-Octet system (Sartorius), as mentioned in the Materials and Methods. The

results, presented in Fig. 6a, clearly demonstrate a dose-dependent interaction of semapimod

with PpsB, immobilized on AR2G sensor. Further analysis of binding kinetics reveals that

semapimod strongly binds to PpsB with a dissociation constant (K_d) of 110nM. As a control,

interaction of semapimod was also analysed with the purified Ppe60, which fails to exhibit

any binding.

PDIM is compromised in Sem^R strain

As hitherto mentioned, PDIM has been found to act as a barrier for uptake of nutrients and

large-size inhibitors. Moreover, our results clearly establish that semapimod targets PDIM

biosynthesis protein PpsB by direct binding. Based on our findings, we speculate that

semapimod resistance is acquired in Mtb owing to disruption in the cell-wall PDIM, which

alleviates the regulation of L-leucine uptake. To test our hypothesis, we analysed the cell-wall

lipids of the wild-type and Sem^R strains of Mtb mc^2 6206. Both mycolic acids and PDIM

were extracted from these strains and analysed by thin layer chromatography, as described in

the Materials and Methods. While no difference in the profiles of mycolic acid methyl esters

(Supplementary Fig. 11) or the tri-acyl glycerol (TAG) was noticed between the two strains,

the Sem^R exhibits a marked reduction in PDIM (Supplementary Fig. 11 and Fig. 6b).

Intracellular survival of Mtb H$_{37}$Rv pathogen in the organelles of infected BalB/C mice

is decreased upon semapimod treatment

Although, endogenous leucine biosynthesis is extremely essential for bacterial survival and

virulence, there is no study describing underlying mechanism of L-leucine uptake from

extracellular environment and its importance in mycobacterial virulence during animal

infections. To examine if L-leucine uptake is critical for intracellular survival of the

pathogenic Mtb H$_{37}$Rv strain with intact endogenous leucine biosynthesis, we tested the

effect of semapimod treatment on intracellular burden of Mtb H$_{37}$Rv during mouse infection.

Infection was performed via aerosol route as described in the Materials and Methods. After

21 days of infection, a group of 3 mice were treated with 5mg/kg dose of semapimod,

whereas another group of 3 mice were treated with vehicle (1x phosphate buffered saline) by
daily oral gavaging for 4 weeks, typically as shown in the schematic. Bacterial load was
examined in lungs and spleen of animals from both the groups by colony forming unit (CFU)
estimation (Fig. 7a). Examination of the gross lung pathology reveals a large number of
granulomatous nodules in the untreated infected group, whereas semapimod treatment led to
a significant reduction in the disease pathology (Fig. 7b). Remarkably, semapimod treatment
caused ~82% and ~85% reduction in CFU counts in lungs and spleen, respectively, after 4
weeks of semapimod treatment (Fig. 7c and d). Noteworthy to mention, *in vitro* susceptibility
to vancomycin remained unaffected in bacteria retrieved from lungs of the untreated and
semapimod-treated mice (Supplementary Fig. 12), thus indicating comparable levels of cell
wall PDIMs, one of the virulence factors, in both untreated and semapimod-exposed Mtb
H$_{37}$Rv.
Overall, these results provide first evidence of the effect of a repurposed compound
obstructing the *in vitro* L-leucine uptake on Mtb virulence and TB pathogenesis during
infection.

Discussion
Conventional approaches of drug discovery involve screening of NCEs against either the
validated 'drug target' (target-based screen) or the pathogen as a whole (phenotypic screen)[25].
The aim is to identify inhibitor(s) showing a novel MoA. However, any promising NCE faces
a multitude of barriers to reach to market launch from initial development[26]. Repurposing
offers the possibility of identifying new MoA of the existing drugs that are already approved
for human use[27, 28]. Therefore, it bypasses the traditional drug discovery and development
pipeline involving *de novo* synthesis of NCEs. Several non-antibiotic drugs that are currently
approved or under development may inhibit microbial pathogens, and therefore can be
explored as novel antimicrobials, either alone or in combination with the existing
antibiotics[29, 30].
The present study was designed with the aim to identify novel TB inhibitors using the FDA-
approved repurposed library comprising of over 3600 molecules. Remarkably, in addition to
the known anti-TB drugs, we found several novel hit molecules that were developed against
non-communicable diseases such as cancer, cardiovascular, metabolic and neurological
diseases (Fig. 1a-b) that can be explored as novel anti-TB drugs in future. Notably, one of the
top hits, namely semapimod, which inhibited Mtb mc^2 6206 growth at less than 20nM
concentration, has never been reported for antibiotic activity. We show that this drug, not only

kills Mtb mc^2 6206 *in vitro*, but also significantly reduces bacterial growth during
macrophage infection (Fig. 1c-e). Surprisingly, semapimod did not exhibit growth inhibition
of other mycobacteria including the pathogenic Mtb H37Rv (Fig. 3), which was used to
develop the auxotroph by deletion of *leuC-leuD* and *panC-panD* genes. Remarkably, we
demonstrate that growth inhibition by semapimod is due to blockage of L-leucine availability
in Mtb mc^2 6206 (Fig. 3). We rule out the possibility of L-leucine quenching or modification
by semapimod, as both the *leuC-leuD*-expressing and the Sem^R strains of Mtb mc^2 6206
continue to metabolize exogenous L-leucine in the presence of drug. Instead, our results
suggest that semapimod restricts L-leucine uptake in Mtb. As a consequence, this drug is not
effective against other mycobacterial species with the functional endogenous leucine
biosynthesis pathway.
Cell membrane acts as a permeability barrier to foreign substances and selectively filters
molecules that are essential for the cellular physiology. Transport of solutes involves three
different classes of transport mechanisms, viz., passive diffusion, facilitated diffusion and
active transport. Of these three different classes, amino acid transport fulfil the criteria of
facilitated diffusion and active transport[31]. The two best characterized amino acid transport
systems are the high-affinity periplasmic binding protein-(BP) dependent transport system for
leucine, isoleucine, and valine (LIV-I and LIV-II) in *E. coli* and the high-affinity histidine
transport system in *S. typhimurium* [32, 33, 34, 35, 36]. Although, Mtb is able to synthesize all 20
amino acids[37, 38, 39], however, to ensure constant supply of amino acids under a wide variety
of environments, the TB pathogen must be equipped for their transport. Genome analysis of
the fast-growing soil organism, *M. smegmatis* reveals 15 genes spread across three different
loci that are putatively involved in BCAA transport (Supplementary Fig. 5). However,
homologues of none of these are found in Mtb. Although, intracellular Mtb exploits multiple
host-derived amino acids as nitrogen sources including BCAAs during growth in
macrophages[40, 41], thus far, it remains unclear how amino acids are transported across the cell
envelope of Mtb. To the best of our knowledge, this is the first study providing evidence for
the presence of a functional transport system for a BCAA in Mtb. We show that semapimod,
which is an anti-inflammatory drug, targets a polyketide synthase involved in the cell-wall
PDIM biosynthesis, and selectively kills *in vitro* the leucine auxotroph by blocking the L-
leucine uptake, possibly by modulating the PDIM architecture. The cell-wall PDIM regulates
uptake of nutrients and provides barrier to large molecule inhibitors. PDIM tends to shed
during culturing of Mtb in the presence of ionic detergents, thus providing better access to
nutrients and large molecules such as vancomycin. As a consequence, Mtb strains lacking

PDIM not only grow faster, but also become more susceptible to vancomycin[24]. Remarkably,

the Sem[R] strain which accumulates mutation in the PpsB near acetyltransferase domain,

exhibits both the phenotypes, i.e., a dramatic increase in its susceptibility to vancomycin and

faster growth rate compared to the wild-type Mtb, despite culturing in the presence of the

non-ionic detergent. Loss of PDIM in the drug-resistant strain is further confirmed by TLC

analysis of the PDIM, whereas other lipids such as triacyl glycerol (TAG) or mycolic acids

remain unaltered (Fig. 6b and Supplementary Fig. 11). Most importantly, susceptibility to

semapimod is restored in the Sem[R] Mtb mc^2 6206 by overexpression of *ppsB*. Taken together,

these results clearly demonstrate for the first time the involvement of the cell-wall PDIM in

L-leucine uptake in Mtb (Fig. 8). At present, it is not understood how PDIM selectively

facilitates movement of L-leucine, and whether other apolar hydrophobic molecules follow

the same route for their transport across the mycobacterial cell membrane. Since, PDIM is an

important virulence factor[42], understanding its role in the transport of other metabolites

including amino acids will shed important light on virulence strategies employed by Mtb.

Semapimod treatment interferes with the expression of more than 10% of protein-coding

genes of Mtb involved in a variety of functions. A careful analysis of the transcriptional

profile further provides evidence corroborating the deviation of leucine levels in the cell by

semapimod treatment. For instance, *lrpA*, which encodes for a transcriptional regulator of the

leucine-responsive regulatory protein (Lrp) family[43], is upregulated by 9.25-fold. It has been

shown that binding of Mtb LrpA with the target promoters is influenced by amino acids and

vitamins[43]. Not only LrpA, but also some of its regulons such as *lat* (94.01-fold), *whiB2*

(0.33-fold), *pks2* (12.58-fold), and *gid* (0.37-fold) also show significant modulation.

Among other differentially expressed gene sets, the most notable ones are those involved in

energy and lipid metabolism. Since, leucine is catalyzed into acetyl-CoA[44], we hypothesize

that in the presence of semapimod, level of acetyl-CoA is perturbed in the cell. Acetyl-CoA,

being the precursor of energy metabolism, subsequently impacts the overall energy state, as

evidenced by >80% reduction in ATP following drug treatment. In other bacteria such as

Salmonella, L-leucine supplementation stimulates the TCA cycle, causing increased

intracellular NADH and ATP concentration[45], which further substantiates the contribution of

intracellular leucine in energy production in mycobacteria.

Strikingly, semapimod-treated Mtb exhibits significant upregulation of genes that encode for

proteins involved in MCC such as methylcitrate synthase PrpC (19.48-fold), methylcitrate

dehydratase PrpD (92.5-fold), the regulator of *prpC* and *prpD* genes– PrpR (28.55-fold), and

an enzyme, Icl1 (68.69-fold). Icl1 of Mtb exhibits both the isocitrate lyase activity involved

in the glyoxylate shunt, and methylisocitrate lyase activity which is responsible for catalysing

the last reaction of the MCC, resulting in the production of succinate and pyruvate that are

routed into the TCA cycle[46]. Although, MCC is responsible for metabolizing the toxic

intermediates produced from catabolism of odd-chain fatty acids (OCFAs) such as

cholesterol, which is the primary source of nutrient during host infection , it has been

observed that both the glyoxylate shunt and MCC are functional in wild-type Mtb, even in the

presence of glycolytic carbon source[47]. Icl1-deficient Mtb undergoes a progressive depletion

of TCA cycle intermediates, which in-turn impair Mtb's respiratory activity. Although, excess

of leucine in the extracellular environment is lethal, intracellular leucine production is critical

as it provides an alternative pool for acetyl-CoA biosynthesis which feeds into the respiratory

and lipid metabolic pathways[48]. We believe that decline in intracellular leucine together with

reduced expression of genes involved in its catabolism (*accD1*, *bkdA* and *bkdB*) triggers the

induction of alternate pathway such as MCC to maintain the cellular acetyl-CoA pool.

The complex cell-wall lipids of Mtb provide a hydrophobic barrier around the bacterium.

These can be esterified with multiple methyl-branched (MB) long chain fatty acids such as

PDIM, sulfolipid-1, and di-, tri-, and poly-acyl trehalose. These MB lipids are synthesized by

individual polyketide synthase (PKS) complexes requiring malonyl-CoA and methyl

malonyl-CoA (MMCoA), which are derived from acetyl-CoA and propionyl-CoA,

respectively. PDIM is an important virulence factor, and controls the movement of nutrients

and large molecule inhibitors such as vancomycin across the bacterial cell membrane. While

genes involved in synthesis of phthiocerol moiety of PDIM such as *ppsB*, *ppsC* and *ppsD* are

downregulated, those involved in biosynthesis of SL-1 (*papA1* and *pks2*) and poly-acyl

trehalose (*pks3*, *pks4*, *papA3*, and *fadD21*) are highly upregulated in drug-treated bacteria.

Since, both PDIM and SL-1 biosynthetic pathways share a common precursor, MMCoA, it

has been observed that reduction of PDIM synthesis favours SL-1 production due to

increased flux of MMCoA to the SL-1 biosynthetic pathway, and *vice versa*[49]. Intrigued with

these observations, we speculate that semapimod induces remodelling of cell-wall lipids,

restricting L-leucine uptake by an unknown mechanism (Fig. 8). Targeting of PpsB by

semapimod further supports the above hypothesis and warrants future studies to identify

alterations in the cell envelope which impose restriction on L-leucine transport.

Uptake of host-derived nutrients including amino acids supports bacterial viability during

infection[40, 41]. Targeting the amino acid transport system offers a novel strategy to curtail the

supply of host-derived nutrients to the pathogens, which are metabolically quiescent during

intracellular stay[50]. Semapimod, which has been found to inhibit the L-leucine uptake in Mtb,

may not be the preferred choice of L-leucine inhibitor owing to its anti-inflammatory effect.
It inhibits production of proinflammatory cytokine such as TNF-α, IL-1β, and IL-6. Also, it
has been reported that semapimod interferes with the TLR4 signalling[17, 18, 19, 20]. These
cytokines play a critical role in providing protection to the host against active TB infection[51].
Nonetheless, despite its anti-inflammatory activity which may promote bacterial
proliferation, semapimod is able to reduce the intracellular burden of virulent Mtb H37Rv
strain in lungs and spleen of the infected mice (Fig. 7). These findings provide an indirect
evidence of the importance of L-leucine uptake for Mtb virulence during host infection.
Advancements in the genetic manipulation tools such as CRISPR-Cas technology[52] has
significantly contributed to enhancing our understanding of metabolic networks in
mycobacteria. Screening of inhibitors against other Mtb auxotrophic strains is expected to
provide novel insights into transport mechanisms of essential metabolites in Mtb that can be
explored as novel anti-TB drug targets.

Materials and Methods

Strains and culturing of bacteria. For propagation of plasmids, *Escherichia coli* strain
XL1Blue[TM](Agilent) was used, whereas expression and purification of PpsB was performed
in *E. coli* BL21 DE3 (Novagen). Mtb H37Rv was received from Dr. Ramandeep Singh at
THSTI, India, and Mtb mc^2 6206 strain was provided by Dr. William Jacobs at Albert
Einstein College of Medicine, NY, USA. *E. coli* was cultured in the Luria-Bertani medium
(Becton Dickinson), Mtb H37Rv was cultured in Middlebrook 7H9 containing 0.05%
tyloxapol (Merck), 0.5% glycerol (Merck) and 1X OADS (oleic acid-albumin-dextrose-
saline) (7H9-O) or Middlebrook 7H11 containing 0.5% glycerol (Merck) and 1X OADS
(7H11-O). For culturing Mtb mc^2 6206 strain, media were supplemented with 50μg/ml L-
leucine (Merck) and 24μg/ml pantothenate (Merck) (7H9-PLO). Liquid cultures were grown
either in Corning 50 mL centrifuge tubes (Corning) or in 250ml flasks (Corning) with not
more than one-third volume of bacterial cultures, whereas plates were incubated at 37°C.
Wherever required, we used 50μg/ml kanamycin (Merck), 50μg/ml ampicillin (Merck), and
150μg/ml hygromycin (Invitrogen) for *E. coli*, whereas for Mtb 25μg/ml kanamycin and
50μg/ml hygromycin were used.

Screening of inhibitors. Repurposed library comprising of 3614 FDA-approved drugs was
obtained from MedChemExpress (www.medchemexpress.com) in the 96-well plate format.
All compounds were suspended in DMSO (Merck) at 5mM concentration, and stored frozen
at -80C before use. For initial evaluation of compounds, 100μl Mtb mc^2 6206 culture, diluted

to OD_{600} of ~0.01 in 7H9-PLO medium was aliquoted in each well of the round-bottom 96-

well plate containing 1µl of respective drugs, so that final conc. of drug became 50µM. Side

wells were filled with 100µl medium to maintain humidity and to prevent loss of samples

from drug-containing wells. As an untreated (UT) control, 100µl bacterial culture was treated

with 1µl DMSO. Plates were incubated at 37 ˚C for 2-3 weeks until UT wells exhibited

substantial growth. Drugs showing complete inhibition of growth by visual inspection of the

pellet formation at the bottom were considered inhibitors of Mtb mc^2 6206, and used for

determination of drug concentration inhibiting 90% growth (MIC_{90}).

Determination of MIC_{90}. The MIC_{90} of active inhibitors was determined by using the

resazurin assay. Briefly, Mtb mc^2 6206 was incubated at OD_{600} of 0.005 with different

concentrations of drug ranging from 25-0.10µM in the flat bottom 96-well plates. After 2

weeks of incubation at 37 ˚C, cultures were transferred in the black 96-well plates and

incubated with 10% of Alamar Blue^{TM} reagent (Thermo Scientific) for 16hrs 37 ˚C. Viability

was calculated by comparing fluorescence of the drug-treated and DMSO-treated samples at

560nm of excitation and 590nm of emission. The minimum drug concentration at which

~90% growth inhibition is found, was considered as MIC_{90}.

Macrophage infection and intracellular CFU analysis. For macrophage infection, THP1

cells were seeded in 24-well plates at a density of 2×10^5 cells per well and differentiated

using 20nM phorbol 12-myristate 13-acetate (PMA). After 48hr of differentiation, cells were

washed once with incomplete RPMI medium (Himedia). Thereafter cells were infected with

Mtb mc^2 6206 at 1:5 MOI (macrophage: bacteria) using RPMI medium containing 10% FBS

medium, 24µg/ml pantothenate and 50µg/ml L-leucine (RPMI-PL). After 4hr of incubation,

cells were washed with 1x phosphate buffered saline (1x PBS) for three times and

replenished with RPMI-PL. After overnight incubation, cells were treated with 0.1, 0.2 and

1.0µM semapimod, and 2.5µM INH in four wells each and remaining 4 wells were left

untreated. Drug treatment was repeated again after 3 days of incubation. On the 6th day, cells

were lysed in 0.1% Triton X100, and serial dilutions of the lysates were plated on 7H11-

OADS-PL agar plates for CFU enumeration.

Global gene expression analysis by RNA sequencing. Freshly inoculated culture of Mtb

mc^2 6206 was treated with 50nM of semapimod at 0.40 OD, whereas another set was left

untreated. After 24hrs of incubation at 37 ˚C with shaking at 200rpm, both semapimod-

treated and untreated cultures were pelleted down, washed twice with 1x PBS and stored at -

80 ˚C before RNA extraction. Total RNAs were extracted from three biological replicates of

untreated (control) and semapimod treated cultures followed by treatment with DNase I, as
described previously[53]. The DNase-treated RNA samples were supplied to Clevergene
(https://clevergene.in/) for further processing and sequencing, typically as reported earlier[53].
Genes that exhibit absolute log2 fold change ≥ 1, and FDR-adjusted p value of ≤ 0.01 were
considered significant, and their expression profile is presented in the volcano plot and
heatmap.

Quantitative reverse transcription real-time PCR (qRT-PCR). Total RNA was treated
with DNase I (Ambion), and 500ng of DNase-treated RNA samples were subjected to
complementary DNA (cDNA) synthesis using SuperScript IV reverse transcriptase, as
recommended by the manufacturer (Thermo Scientific). qRT-PCR was performed by using
50ng cDNA and gene-specific primers (Supplementary Table 1) with the help of SYBR
Green PCR master Mix (ABI). Amplification was monitored in real-time and quantification
was carried out using ABI 7500 Fast Real-Time PCR System (Applied Biosystems), as
mentioned earlier[53].

Measurement of ATP. Mtb mc2 6206 was cultured in the presence or the absence of 50nM
semapimod for 24 h. For intracellular ATP estimation, 1ml of untreated and treated cultures at
OD_{600} of 0.1 were pelleted by centrifugation, washed twice with 1x PBS and lysed in 200µl
of 1x PBS by boiling at 98 ˚C for 10min. The bacterial lysates were centrifuged at 14,500
rpm for 3min to remove debris, and supernatant was collected for ATP estimation by using
Bac Titer- Glo[TM] assay kit as recommended by the manufacturer (Promega).

Estimation of intracellular leucine: Intracellular leucine was estimated by liquid
chromatography mass spectrometry (LC-MS/MS), as follows:

    a) **Preparation of samples:** Freshly-grown Mtb mc[2] 6206 cultures were diluted to 1.0
       OD_{600} and divided into two portions, one treated with 50ng/ml semapimod and the
       other one left untreated. After 24 hours of incubation at 37 ˚C with shaking at 200rpm,
       cultures equivalent to 5.0 OD_{600} were drawn from both the samples and pelleted by
       centrifugation. After through washing with 1 x PBS, bacterial metabolites were
       extracted using methanol precipitation method, and subjected to LC-MS/MS.

    b) **LC-MS/MS:** QTRAP® 6500+ MS system coupled with Ultra-high-performance
       liquid chromatographic system, (SCIEX ExionLC[TM]) was used for targeted LC-MS
       analysis. The ACQUITY UPLC HSS T3 column (1.6µm, 2.1 x 100mm) was used for
       separation, and the column oven temperature was set at 40 °C. Solvent A (10mM

Ammonium formate with 0.1% formic acid in water) and solvent B (0.1% formic acid

in methanol) were used for LC gradient with changing the concentration of solvent B

as follows: 0 min, 1% B; 1 min, 15% B; 4 min, 35% B; 7 min, 95% B; 9 min, 95% B;

10 min, 1% B, 20 min, 1% B. The flow rate was kept at 300 µl/min and the injection

volume was optimized at 5 µl. Following parameters were used for electrospray

ionization: electrospray voltage, +5500 V; ion source temperature, 500 ˚C; curtain gas

of 24, CAD gas 9, and gas 1 and 2 of 60 and 40 psi, respectively.

c) **Compound parameter:** Following optimized MRM parameters of de-clustering

potential (DP), collision energies (CE), entrance potential (EP), and collision cell exit

potential (CXP) were used: DP-40, EP-10, CE-15 & CXP-22. Commercially procured

ultrapure L-leucine (Merck) was used for selection of Q1 and Q3 masses. Sharp peaks

of pure L-leucine were chosen as references to identify the leucine peaks in samples.

Data analysis for LC-MS/MS was performed using the SCIEX (ANALYST 1.7.2)

software. Analyte was confirmed by comparing the retention time and the ratio of

characteristic transition between the test sample and the reference.

Estimation of percent viability in response to vancomycin or semapimod treatment.

Bacterial cultures were incubated at OD_{600} of 0.005 with different concentrations of

vancomycin or semapimod in the flat bottom 96-well plates. After 2 weeks of incubation at

37 ˚C, growth was estimated by measuring OD_{600} of the cultures using multi-mode plate

reader, which was used for measuring percent viability in the drug-treated cultures with

respect to the untreated control.

Cloning and expression of genes in Mtb. The *leuC-leuD* and *panC-panD* open reading

frames (ORFs) were PCR amplified using Mtb H37Rv genomic DNA with gene-specific

primer pairs (Supplementary Table 1) and GoTaq polymerase, as recommended by the

manufacturer (Promega). PCR amplicons with *Nde* I and *Hind* III overhangs were gel

purified and subjected to restriction digestion with the respective enzymes. The digested

DNA fragments were column-purified and annealed at the same sites in a Hyg[R] replicative *E.*

coli-mycobacterium shuttle plasmid, pHsp60 under a constitutive promoter of *hsp60*[54]. The

resulting recombinant plasmids, pHsp-*leuCD* and pHsp-*panCD* were electroporated in Mtb

mc[2] 6206 after sequence verification by Sanger's sequencing. To ascertain expression of the

respective genes, recombinant clones obtained by transformation with pHsp-*leuCD* were

selected on 7H11-OADS agar plates containing pantothenate and hygromycin, whereas those

obtained with pHsp-*panCD* were selected on 7H11-OADS agar plates containing L-leucine
and hygromycin.

For expression of *ppsB*, the ~4.6kb *ppsB* ORF was PCR amplified using Mtb H37Rv
genomic DNA with Pr. 2396-2428 (PCR-A) and 2429-2397 (PCR-B) (Supplementary Table
1). Both the PCR-A and PCR-B fragments of ~2.2 and ~2.4 kb, respectively, were
subsequently cloned in the pGEMT-easy plasmid by following the TA cloning method, as
suggested by the manufacturer (Promega). Both the recombinant clones were sequence
verified, which revealed the presence of PCR-A in the forward orientation in pGEMT-A
clone and PCR-B in the reverse orientation in pGEMT-B clone. Both the clones were
subjected to restriction digestion with *Nde I* and *Sma I*. PCR-A fragment of ~2.2kb was gel
eluted and cloned in the pGEMT-B plasmid at the same sites, to create full-length ppsB of
~4.6kb. Next, DNA fragment comprising of full-length *ppsB* ORF was obtained from the
positive clone of pGEMT-*ppsB* by *Nde I-Hind III* digestion, and cloned in pHsp60 at the
same sites. To express *mce1D* and *ppe60*, the respective ORFs were PCR amplified from Mtb
H37Rv genomic DNA using Pr. 2277-2278 and Pr. 2281-2282, respectively, (Supplementary
Table 1), restriction digested and cloned in pHsp60 at *Nde I-Hind III* sites, as described
above. The resulting recombinant plasmids– pHsp-*ppsB*, pHsp-*mce1D* and pHsp-*ppe60* were
verified by Sanger's sequencing and electroporated in wild-type or Sem^R Mtb mc^2 6206. The
transformants were selected on 7H11-PLO agar plates containing hygromycin, and positive
clones verified by colony PCR were used in subsequent experiments.

Generation of Sem^R strain. To induce semapimod-resistance, 5ml culture of Mtb mc^2 6206
at OD$_{600}$ of ~2.0 was pelleted by centrifugation followed by suspension in ~500µl 7H9-PLO
medium. Aliquots of 50µl of the culture suspension were spread on multiple 7H11-PLO agar
plates containing 100nM semapimod. After 4-8 weeks of incubation at 37 ˚C, single colonies
appeared on drug-containing plates were cultured in 1ml 7H9-PLO medium. Subsequently,
cultures of the wild-type and putative Sem^R strain of Mtb mc^2 6206 were spotted on 7H11-
PLO agar plates with or without 50nM semapimod to monitor growth.

Genomic DNA extraction from mycobacteria and whole genome sequencing. Mtb
mc^{2}6206 wild type and the Sem^R strains were grown in 7H9-PLO, and genomic DNA
extraction was performed using the conventional cetyltrimethylammonium bromide (CTAB)-
based method. Whole genome sequencing (WGS) of the Mtb strains was conducted using the
PacBio Sequel II long read sequencing platform. The WGS was outsourced to Nucleome
Informatics Private Limited, Hyderabad, India. A detailed sequencing workflow was

employed to ensure the generation of high-quality data. Briefly, DNA quantification was

performed using Qubit fluorometry, while DNA quality was assessed on a 1% agarose gel.

Purity ratios were analyzed using the NanoDrop 2000 spectrophotometer. DNA fragments

between 7–10 kb were prepared for SMRTbell library creation utilizing the Megaruptor 3

system, followed by purification with AMPure PB beads. The Agilent FEMTO Pulse analyzer

was used to verify size distribution, and DNA concentration was measured with the DNA HS

assay on the Qubit 3.0 fluorometer. Libraries were constructed with the Template Prep Kit,

starting with the removal of single-strand overhangs, followed by DNA damage repair and

End-Repair/A-tailing. Adapters were then ligated to the repaired DNA fragments to form

SMRTbell libraries, with hairpin dimers removed using AMPure PB beads. The final pooled

libraries were further purified and analyzed with the Agilent FEMTO Pulse. Primer annealing

and polymerase binding were conducted using the Sequel II binding kit 2.2 before loading the

libraries onto SMRTcells for sequencing in CCS/HiFi mode on the PacBio Sequel II system.

This robust workflow ensured the production of high-quality sequencing data essential for

precise genomic analysis.

Bioinformatic analysis for genome assembly preparation and polymorphism detection.

PacBio Sequel II raw subreads were processed into high-fidelity (HiFi) reads using the 'CCS'

tool (version 6.2.0). The genome assembly was constructed using 'Trycycler' (version 0.5.4)

with its default parameters[55]. Three individual assemblies were initially created with the tools

'Flye' (version 2.9.2), 'Minipolish' (version 0.1.3), and 'wtdbg2' (version 2.5)[56, 57, 58]. These

assemblies were combined to generate a consensus assembly using Trycycler's default

configuration. Genome completeness was evaluated using BUSCO (version 5.4.5), with the

actinobacteria_class_odb10 lineage dataset as the reference[59]. For comparative genomics,

variant calling was performed using Genome Analysis Toolkit (GATK, version 4.5),

FreeBayes (version 1.3.7), and TB-Profiler (version 5.0.1)[60, 61, 62]. Mutations of interest were

carefully selected based on their confidence levels to ensure accuracy.

Cloning, expression and purification of PpsB. Full-length *ppsB* ORF fragment was

obtained from pGEMT-*ppsB* by *Nde I-Hind III* digestion and cloned at the same sites in

pET28b (Novagen) to express the recombinant protein with 6x His tags at the N-terminus *in*

E. coli. For this, pET28b-*ppsB* was transformed in *E. coli* C43(DE3) cells and recombinant

clones were selected on LB agar containing kanamycin. Single colony of *E. coli*::pET28b-

ppsB was inoculated in 10ml of LB broth medium containing kanamycin and culture was

grown to turbidity at 37 ˚C with shaking at 200rpm. Secondary culture was subsequently

inoculated from the seed culture in one liter of LB broth medium with kanamycin. Expression
of 6XHis-PpsB was induced by adding 1mM IPTG in the culture at OD_{600} of 0.80. After
overnight incubation with IPTG at 18 ˚C, culture was pelleted and washed twice with 1x
PBS. Induced culture pellet was suspended in 50ml of lysis buffer (50mM Tris, pH 8.0,
50mM NaCl and 5% glycerol) and cell lysis was performed by using French press. Cell
debris was removed by centrifugation and clarified cell lysate was incubated with Ni-NTA
beads (Qiagen) for 3 h at 4 ˚C. Beads were subsequently transferred into an empty column
and unbound proteins were allowed to pass through. Beads immobilized with 6XHis-PpsB
were washed with 5 bed volumes of the lysis buffer, followed by washing with 5 bed volumes
of wash buffer (lysis buffer containing 20mM imidazole). Immobilized 6XHis-PpsB was
subsequently eluted with elution buffers (lysis buffer containing 100-300mM imidazole). The
purity of eluted protein in different elution fractions was assessed by SDS-PAGE. Fractions
showing >90% purity of PpsB were pooled and dialyzed against lysis buffer. The purified
proteins were stored in small aliquots at -80°C for subsequent use.

Protein-drug interaction by BLI-Octet. Optical interference-based Bio layer interferometry
(BLI) from Octet system by ForteBIO was used to ascertain the interaction of semapimod
with Mtb PpsB. Briefly, purified 6xHis-tagged PpsB protein was dialyzed in 10 mM sodium
acetate pH 3.9, followed by immobilization onto the amine reactive group sensor (2nd-
generation) (AR2G) up to a level of 2.1 nm. Different concentrations of semapimod were
prepared in water and used to acquire differential graded response. Binding constant was
calculated as per the standard steps of baseline (60 s), association (180 s) and dissociation
(180 s).

Extraction of lipids. To obtain the apolar lipids, bacterial cultures equivalent to OD_{600} of
~150 were pelleted, washed with 10ml of water and suspended in 2 mL methanol-0.3% NaCl
(10:1) and 1 mL petroleum ether. After mixing vigorously to emulsify the organic and the
aqueous layers, samples were centrifuged at a speed of 2000 x *g* for 5 minutes to allow
mixture to separate. The upper organic layer was collected in the microcentrifuge tubes, and
the extraction was repeated by adding 1 mL petroleum ether to the lower aqueous phase and
stirred for another 30 min. The organic layer was collected as described above and combined
with the previous extracts. Petroleum ether was subsequently evaporated by vacuum drying at
the room temperature and the resulting apolar lipids were stored at -80 ˚C for analysis by thin
layer chromatography (TLC).

To prepare MAMES, culture pellets equivalent to OD_{600} of ~150 were suspended in 1ml of

20% tetrabutylammonium hydroxide and incubated at 100 ˚C for overnight. After overnight

incubation, 0.4 mL of dichloromethane and 50μl of iodomethane were added to the reaction

mixture and stirred for 1 h at room temperature. The biphasic mixture was separated by

centrifugation at 2000 x g for 5 minutes, and lower organic layer containing MAMES was

collected in the microcentrifuge tubes. The MAMES fraction was completely dried and

suspended in 1 mL of diethyl ether. The supernatant was collected in a new tube and vacuum

dried. The dried fraction was suspended in 0.2ml of toluene and 0.1ml of acetonitrile, and

after complete dissolution 0.2ml more acetonitrile was added. Samples were chilled at -20 ˚C

for overnight and the MAMES were collected by centrifugation at 10000 x g for 20 minutes.

MAMES present in the pellet fraction were stored at -80 ˚C for further analysis by TLC.

Analysis of lipids by TLC. Both the apolar lipids and MAMES were detected on aluminum-

backed silica gel 60F254 plates (Merck). Dried lipid pellets were suspended in

dichloromethane and equal volume of samples from both the strains were spotted on the TLC

plates. Apolar lipids were resolved by two dimensional-TLC (2D-TLC) using petroleum

ether:ethyl acetate solvent (98: 2) for 3 times in the first dimension, and petroleum

ether:acetone mixture (98: 2) for one time in the second dimension. MAMES were resolved

by one dimensional-TLC using petroleum ether:diethyl ether (95:5) solvent for 6 times. After

complete run, TLC plates were air dried and sprayed with 5% phosphomolybdic acid

solution, prepared in ethanol. Lipids were visualized on the TLC plates after charring at 100

˚C for 2-5 minutes.

Mice infection. Mice infection was performed as per the guidelines by the committee

for   the purpose   of   control   and   supervision   of experiments on   animals

(CPCSEA,   India). Animal infection experiment was performed after receiving approval

from the   Institutional Animal Ethics Committee of THSTI. Briefly, 6- to 8-week-old

female BALB/c mice were infected with single cell suspension of mouse-passaged Mtb

$H_{37}Rv$ at ~$5x10^7$ cells/ml via aerosol route in a closed aerosol chamber in the biosafety level

3 laboratory. To check the establishment of infection, bacterial load in lungs were examined

at day 1 post-infection and the disease progression was assessed at day 21 post-infection, by

CFU plating on selective 7H11-OADS agar plates containing 100μg/ml carbenicillin,

22μg/ml trimethoprim, 22μg/ml amphotericin B, 11μg/ml vancomycin, 27.5μg/ml polymyxin

B and 0.88μg/ml cycloheximide. At day 21, mice were divided into two groups; one group

was given semapimod (5mg/kg in 5% sucrose) by daily oral gavaging whereas the other half

was given equal dose of 5% sucrose. CFUs were enumerated at day 28 post-semapimod or

sucrose treatment to examine bacterial burden in lungs, and in spleen for dissemination.

Statistics and reproducibility. Some experiments were only performed with 2 biological

replicates. Statistical analysis was performed with data obtained from 3 or more biological

repeats by determining *p* values with the help of GraphPad Prism version 10.3.1 (464)

software.

Legends to Figures

Figure 1. Screening of FDA-approved library of molecules against Mtb mc^2 6206 reveals

growth-inhibitory effect of semapimod, an anti-inflammatory small molecule. a, Status

of inhibitors showing activity against Mtb mc^2 6206. Bar-graph depicts Mtb mc^2 6206

inhibitors acting on different host metabolic pathways. **b**, MIC$_{90}$ status of Mtb inhibitors. Bar-

graph shows inhibitors with different MIC$_{90}$ against Mtb mc^2 6206, *in vitro*. Molecules with

<0.025µM MIC$_{90}$ (encircled) are tabulated in the inset. **c**, Determination of minimum dose of

semapimod for complete inhibition of Mtb mc^2 6206 growth by visual inspection. Isoniazid

(INH) was used as a control in the 96-well plate-based assay. Drug concentration beyond

which no visual growth is observed, is written in bold. **d**, Effect of semapimod on the *in vitro*

growth of Mtb mc^2 6206. Shown is the CFU enumeration of drug-treated and untreated

(control) bacteria at the indicated time points. **e**, Effect of semapimod on the intracellular

proliferation of Mtb mc^2 6206. Intracellular growth of Mtb mc^2 6206 in the THP1-derived

macrophages was examined after 6 days of infection in the absence (UT) or the presence of

0.10-1.0µM semapimod. Treatment with 2.5µM INH was used as control. A significant

reduction in CFU counts of Mtb mc^2 6206 is observed in the presence of semapimod under *in*

vitro culture conditions (**d**) as well as during intracellular (*p*< 0.005) growth (**e**). Data

represent mean±s.d of at least n=2 replicates in **d** and n=4 replicates in **e**. Data in **c** are

representative of n=2 independent experiments. *p* values in **e** were obtained after comparison

of CFUs between the UT and semapimod-treated samples, as described in Materials and

Methods.

Figure 2. Effect of semapimod treatment on the expression profile of Mtb mc^2 6206

transcripts. a, Volcano plot of differentially expressed genes in semapimod-treated Mtb mc^2

6206. The plot shows distribution of genes that are differentially expressed *via* log2 fold-

change and the –log *p* values. Broken vertical lines represent the cutoff of ≥1.0 log2 fold-

change, and horizontal line represents the cutoff of >2.583 –log *p* values. Genes below the -

log p cutoff are represented by grey dots. Downregulated genes are represented by red dots,

and those showing upregulation in response to semapimod treatment are shown with green

dots. **b**, Status of differentially accumulated transcripts. Heatmap representation of transcripts

showing accumulation (left) or suppression (right) upon exposure to semapimod across three

biological replicates. **c**, Functional categorization of differentially regulated genes. The

butterfly chart shows distribution pattern of differentially regulated genes according to their

function, as classified in the Mycobrowser database (https://mycobrowser.epfl.ch/genes/).

Mean fold-change values from n=3 biological replicates are shown in **a**.

Figure 3. Semapimod treatment perturbs L-leucine uptake in Mtb mc2 6206. a, Effect of

semapimod on the *in vitro* growth of different mycobacterial species. Growth inhibitory

effect of semapimod was assessed against slow-growing Mtb H_{37}Rv and *M. bovis* BCG, and

fast-growing *M. abscessus* and *M. smegmatis*, respectively, by visual inspection using 96-

well plate-based assay. Rifampicin (Rif) and INH were used as control drugs against Mtb

H_{37}Rv. **b,** Estimation of intracellular leucine in Mtb mc^2 6206. Intracellular leucine was

estimated in untreated (UT) and semapimod-treated (Sem) bacteria, after 24 hours of drug

treatment, by mass spectrometry as described in Materials and Methods. **c–e**, Effect of *leuC-*

leuD and *panC-panD* expression in Mtb mc^2 6206 on bacterial killing by semapimod. Time-

(**c**) and dose- (**d–e**) dependent growth kinetics reveals loss of bactericidal effect of

semapimod against Mtb mc^2 6206 by *leuC-leuD*, and not by *panC-panD* expression. Percent

viability in **e** was calculated with respect to untreated (UT) cultures after 2 weeks of exposure

to different drug concentrations, using 96-well plate-based method as described in Materials

and Methods. Data in **a** and **c–d** are representative of n=2 independent experiments.

Mean+s.d values from n=4 replicates are shown in **b.**

Figure 4. Generation and characterization of semapimod-resistant strain of Mtb mc^2

6206. a, Propagation of putative semapimod-resistant (Sem^R) strain, but not the wild-type

(WT) Mtb mc^2 6206 in the presence of 50nM semapimod confirms drug-resistance in Sem^R.

In contrast, both the strains exhibit substantial growth on 7H11-PLO agar plate in the absence

of drug. Shown is the growth pattern of two different colonies– C1 and C2 of the respective

strains. **b**, *In vitro* growth analysis of the WT and the Sem^R strains of Mtb mc^2 6206.

Comparative analysis of OD_{600} at different time points reveals substantially increased growth

of Sem^R strain compared to WT under *in vitro* culture conditions. **c**, *In vitro* susceptibility of

the WT and the Sem^R strains of Mtb mc^2 6206 to vancomycin. Viability of WT and Sem^R Mtb

strains was determined in the presence of vancomycin by 96-well plate-based assay, as
described in Materials and Methods. Percent viability was calculated with respect to the
untreated (UT) cultures after 2 weeks of exposure to different concentrations of antibiotic.
Results show a substantial increase in susceptibility of Sem^R to vancomycin compared to WT.
d, Whole genome map analysis of Sem^R. Shown is the genome map of Sem^R highlighting
positions of genes undergoing substitutions, when compared with WT Mtb mc^2 6206.
Mutations leading to corresponding changes at the amino acid level in the respective protein
are indicated alongside. Suspect genes presumably involved in providing semapimod
resistance, are marked in red fonts. Data are representative of at least n=2 independent
experiments in **a**–**b**. Mean+s.d values from n=3 biological replicates are shown in **c**.

Figure 5. Overexpression of *ppsB* alters susceptibility of Sem^R strain of Mtb mc^2 6206 to
semapimod and vancomycin. a–**b**, *In vitro* susceptibility of different strains of Mtb mc^2
6206 to vancomycin (**a**) and semapimod (**b**). Viability of WT and Sem^R Mtb strains in the
presence of drugs was compared with those constitutively expressing *ppsB* by 96-well plate-
based assay, as described in Materials and Methods. Percent viability was calculated with
respect to UT cultures after 2 weeks of exposure to different concentrations of inhibitors.
Remarkably, response of Sem^R to both semapimod and vancomycin is reversed upon
overexpression of *ppsB*. Contrarily, *ppsB* expression in WT does not affect bacterial
sensitivity to either of these drugs, thus indicating a specific effect of the PDIM biosynthesis
gene in Sem^R. Mean values from n=3 biological replicates are shown in **a**–**b**.

Figure 6. Semapimod targets PDIM biosynthesis protein, PpsB. a, Analysis of
semapimod-PpsB interaction by BLI-Octet. Interaction of semapimod with Mtb PpsB was
examined by optical interference–based biolayer interferometry from the Octet system
(ForteBIO). Briefly, dialyzed 6xHis-PpsB was immobilized onto AR2G sensor up to a level
of 2.1 nm. Binding was observed at the indicated concentrations of ligand (semapimod) to
acquire differential graded response. Binding constant was calculated as per the standard
steps, described in Materials and Methods. **b**, Analysis of cell-wall apolar lipids in the WT
and Sem^R strains of Mtb mc^2 6206. Cell-wall apolar lipids were extracted and analyzed by
two-dimensional TLC as described in Materials and Methods, which reveals significant
reduction in PDIMs of Sem^R when compared with the WT strain. Contrary to PDIMs, no
change is observed in the triacyl glycerol (TAG) levels between the two strains. Position of
PDIMs is marked by broken circle in both the images for clarity.

Figure 7. Effect of semapimod treatment on survival of Mtb H37Rv in the organelles of
infected BALB/c mice. a, Schematic of mouse infection. Infection was performed by aerosol
route with the virulent Mtb H_{37}Rv strain. After 21 days of infection, mice were divided into
two groups: one receiving only 5% sucrose (sham) and others receiving 5mg/kg semapimod
prepared in 5% sucrose. Intracellular bacterial load was determined by CFU plating of lung
homogenates at days 1, 21 and 49, and in spleen homogenates prepared on day 49. **b**, Gross
pathology of lungs. Images of lungs obtained from both the sham- and semapimod-treated
groups of mice after 28 days of treatment (i.e., day 49 post-infection) are presented. Scale bar
is shown for size reference. **c**–**d**, Effect of semapimod treatment on intracellular survival of
Mtb H_{37}Rv. Intracellular survival was determined by estimating the bacterial burden in lungs
at the respective time points (**c**), and in spleen at day 49 post-infection (**d**), by CFU
enumeration. Data represent mean $\pm$ s.d. values from n=3 animals in **c**–**d**. *p* values in **c** and **d**
were obtained after comparison CFUs between the two groups, as described in Materials and
Methods.

Fig. 8. Proposed model describing the effect of semapimod on L-leucine uptake in Mtb
mc^2 6206. a, Uptake of L-leucine is controlled by cell-wall PDIM. **b**, Semapimod targets
PDIM biosynthesis protein (PpsB) causing altered PDIM profile which restricts L-leucine
uptake in the auxotroph by an unknown mechanism (denoted by '?'), leading to death. **c**, L-
leucine is freely accessible to Sem^R strain, owing to perturbation in the cell-wall PDIM. As a
consequence, semapimod is ineffective against this strain. However, overexpression of wild-
type *ppsB* in Sem^R reverts the bacterial susceptibility to semapimod. The illustration was
created in BioRender. (Agarwal, N. (2025) https://BioRender.com/h04h035).

Data availability

Raw data of the RNASeq are available from the GEO database under project accession
number GSE284673. GEO accession GSE284673 can be reviewed by using the token
qvgleqketjwjlaf on https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE284673. Raw
data of the whole genome sequence of Mtb mc^2 6206 can be accessed from the Sequence
Read Archive (SRA) database under the Bio-project PRJNA1118666, which can be reviewed
by using the reviewer's
link:https://dataview.ncbi.nlm.nih.gov/object/PRJNA1118666?reviewer=i6d7semeju6p35ks3
vnp6gvda3.

Acknowledgements

We thank Dr. Ramandeep Singh at the Translational Health Science and Technology Institute,
India, Dr. Rajesh Gokhale at the National Institute of Immunology, India and Dr. William
Jacobs at the Albert Einstein College of Medicine, USA for providing us the mycobacterial
strains. Ms. Shivani Goswami (Lab Technician) and Mr. Manish Bansal (Technical Officer)
are acknowledged for providing assistance in the screening of inhibitors. We acknowledge
technical support by the research staffs at the Infectious Disease Research Facility (IDRF) at
THSTI in the animal studies. Experimental Animal Facility (EAF) at THSTI is acknowledged
for providing animals. This work was supported by the core research funding from THSTI.
Research fellowships to L.A. (Award No. 15/12/2019 (ii) EU-V) and MYK (Award No.
15/12/2019 (ii) EU-V) by Council of Scientific & Industrial Research (CSIR), and to H.G
from Science and Engineering Research Board, Department of Science and Technology,
Govt. of India (Award No. RJF/2022/000040) are acknowledged.

Author information

Author contributions

N.A. designed the research, performed the experiments, analysed the data, wrote the paper
and provided overall supervision of the study. H.G., Eeba, L.A., and M.Y.K. performed the
experiments and provided inputs in manuscript writing. S.K.B. and B.D. analysed genome
sequence data.

Competing interests

The authors declare no competing interests.

References

1.   WHO. Global tuberculosis report 2024.) (2024).
819

820 2.   Rendon A, *et al.* Classification of drugs to treat multidrug-resistant tuberculosis
821   (MDR-TB): evidence and perspectives. *J Thorac Dis* **8**, 2666-2671 (2016).
822

823 3.   Caminero JA, Scardigli A. Classification of antituberculosis drugs: a new proposal
824   based on the most recent evidence. *Eur Respir J* **46**, 887-893 (2015).
825

826 4.   Caminero JA, Sotgiu G, Zumla A, Migliori GB. Best drug treatment for multidrug-
827   resistant and extensively drug-resistant tuberculosis. *Lancet Infect Dis* **10**, 621-629
828   (2010).
829

830 5.   Akalu TY, Clements ACA, Gebreyohannes EA, Xu Z, Bai L, Alene KA. Risk factors
831   for diagnosis and treatment delay among patients with multidrug-resistant
832   tuberculosis in Hunan Province, China. *BMC Infect Dis* **24**, 159 (2024).
833

834 6. Bakshi A, Narasimhan P, Li J, Chernih N, Ray PK, MacIntyre R. mHealth for the
835 control of TB/HIV in developing countries. In: *IEEE 13th International Conference*
836 *on e-Health Networking, Applications and Services*) (2011).
837

838 7. Hapolo E, Ilai J, Francis T, du Cros P, Taune M, Chan G. TB treatment delay
839 associated with drug resistance and admission at Daru General Hospital in Papua New
840 Guinea. *Public Health Action* **9**, S50-S56 (2019).
841

842 8. Jhun BW, Koh WJ. Treatment of Isoniazid-Resistant Pulmonary Tuberculosis. *Tuberc*
843 *Respir Dis (Seoul)* **83**, 20-30 (2020).
844

845 9. Malenfant JH, Brewer TF. Rifampicin Mono-Resistant Tuberculosis-A Review of an
846 Uncommon But Growing Challenge for Global Tuberculosis Control. *Open Forum*
847 *Infect Dis* **8**, ofab018 (2021).
848

849 10. Mase SR, Chorba T. Treatment of Drug-Resistant Tuberculosis. *Clin Chest Med* **40**,
850 775-795 (2019).
851

852 11. Muniyandi M, Ramachandran R. Current and developing therapies for the treatment
853 of multi drug resistant tuberculosis (MDR-TB) in India. *Expert Opin Pharmacother*
854 **18**, 1301-1309 (2017).
855

856 12. Boyd NK, Teng C, Frei CR. Brief Overview of Approaches and Challenges in New
857 Antibiotic Development: A Focus On Drug Repurposing. *Front Cell Infect Microbiol*
858 **11**, 684515 (2021).
859

860 13. Kim S*, et al.* PubChem 2025 update. *Nucleic Acids Res*, (2024).
861

862 14. Butler MS, Henderson IR, Capon RJ, Blaskovich MAT. Antibiotics in the clinical
863 pipeline as of December 2022. *J Antibiot (Tokyo)* **76**, 431-473 (2023).
864

865 15. Beachy SH, Johnson SG, Olson S, Berger AC. Drug Repurposing and Repositioning:
866 Workshop Summary.) (2014).
867

868 16. Jain P*, et al.* Specialized transduction designed for precise high-throughput unmarked
869 deletions in Mycobacterium tuberculosis. *mBio* **5**, e01245-01214 (2014).
870

871 17. Nishimatsu H*, et al.* Blockade of endogenous proinflammatory cytokines ameliorates
872 endothelial dysfunction in obese Zucker rats. *Hypertens Res* **31**, 737-743 (2008).
873

874 18. Wehner S*, et al.* Inhibition of p38 mitogen-activated protein kinase pathway as
875 prophylaxis of postoperative ileus in mice. *Gastroenterology* **136**, 619-629 (2009).
876

877 19. Wang J, Grishin AV, Ford HR. Experimental Anti-Inflammatory Drug Semapimod
878 Inhibits TLR Signaling by Targeting the TLR Chaperone gp96. *J Immunol* **196**, 5130-
879 5137 (2016).
880

881 20. Miller IS*, et al.* Semapimod sensitizes glioblastoma tumors to ionizing radiation by
882 targeting microglia. *PLoS One* **9**, e95885 (2014).
883

884   21.   Cho BK, Barrett CL, Knight EM, Park YS, Palsson BO. Genome-scale reconstruction
885   of the Lrp regulatory network in Escherichia coli. *Proc Natl Acad Sci U S A* **105**,
886   19462-19467 (2008).
887

888   22.   Duan X*, et al.* Mycobacterium Lysine epsilon-aminotransferase is a novel alarmone
889   metabolism related persister gene via dysregulating the intracellular amino acid level.
890   *Sci Rep* **6**, 19695 (2016).
891

892   23.   Reddy MC, Gokulan K, Jacobs WR, Jr., Ioerger TR, Sacchettini JC. Crystal structure
893   of Mycobacterium tuberculosis LrpA, a leucine-responsive global regulator associated
894   with starvation response. *Protein Sci* **17**, 159-170 (2008).
895

896   24.   Mulholland CV*, et al.* Propionate prevents loss of the PDIM virulence lipid in
897   Mycobacterium tuberculosis. *Nat Microbiol* **9**, 1607-1618 (2024).
898

899   25.   Rutten A, Kirchner T, Musiol-Kroll EM. Overview on Strategies and Assays for
900   Antibiotic Discovery. *Pharmaceuticals (Basel)* **15**, (2022).
901

902   26.   Payne DJ, Gwynn MN, Holmes DJ, Pompliano DL. Drugs for bad bugs: confronting
903   the challenges of antibacterial discovery. *Nat Rev Drug Discov* **6**, 29-40 (2007).
904

905   27.   Strittmatter SM. Overcoming Drug Development Bottlenecks With Repurposing: Old
906   drugs learn new tricks. *Nat Med* **20**, 590-591 (2014).
907

908   28.   Tobinick EL. The value of drug repositioning in the current pharmaceutical market.
909   *Drug News Perspect* **22**, 119-125 (2009).
910

911   29.   Koh Jing Jie A, Hussein M, Rao GG, Li J, Velkov T. Drug Repurposing Approaches
912   towards   Defeating   Multidrug-Resistant   Gram-Negative   Pathogens: Novel
913   Polymyxin/Non-Antibiotic Combinations. *Pathogens* **11**, (2022).
914

915   30.   Temel A, Aksoyalp ZS. A Preliminary Study on the Effect of Deferoxamine on the
916   Disruption of Bacterial Biofilms and Antimicrobial Resistance. *Turk J Pharm Sci* **21**,
917   267-273 (2024).
918

919   31.   Quigley R. Amino Acids Transported by the Neutral Amino Acid System B0AT1. In:
920   *Fetal and Neonatal Physiology*). Elsevier (2017).
921

922   32.   Haney SA, Oxender DL. Amino acid transport in bacteria. *Int Rev Cytol* **137**, 37-95
923   (1992).
924

925   33.   Antonucci TK, Landick R, Oxender DL. The leucine binding proteins of Escherichia
926   coli as models for studying the relationships between protein structure and function. *J
927   Cell Biochem* **29**, 209-216 (1985).
928

929   34.   Landick R, Oxender DL. The complete nucleotide sequences of the Escherichia coli
930   LIV-BP and LS-BP genes. Implications for the mechanism of high-affinity branched-
931   chain amino acid transport. *J Biol Chem* **260**, 8257-8261 (1985).
932

935 | 35. | Guardiola J, De Felice M, Klopotowski T, Iaccarino M. Mutations affecting the different transport systems for isoleucine, leucine, and valine in Escherichia coli K-12. *J Bacteriol* **117**, 393-405 (1974).

36. Landick R, Oxender DL, Ferro-Luzzi Ames G. Bacterial Amino Acid Transport Systems. In: *The Enzymes of Biological Membranes* (ed Martonosi AN). Springer, Boston, MA. (1985).

37. Yelamanchi SD, Surolia A. Targeting amino acid metabolism of Mycobacterium tuberculosis for developing inhibitors to curtail its survival. *IUBMB Life* **73**, 643-658 (2021).

38. Prakash P, Pathak N, Hasnain SE. pheA (Rv3838c) of Mycobacterium tuberculosis encodes an allosterically regulated monofunctional prephenate dehydratase that requires both catalytic and regulatory domains for optimum activity. *J Biol Chem* **280**, 20666-20671 (2005).

39. Giffin MM, Modesti L, Raab RW, Wayne LG, Sohaskey CD. ald of Mycobacterium tuberculosis encodes both the alanine dehydrogenase and the putative glycine dehydrogenase. *J Bacteriol* **194**, 1045-1054 (2012).

40. Borah K*, et al.* Intracellular Mycobacterium tuberculosis Exploits Multiple Host Nitrogen Sources during Growth in Human Macrophages. *Cell Rep* **29**, 3580-3591 e3584 (2019).

41. Gouzy A, Poquet Y, Neyrolles O. Nitrogen metabolism in Mycobacterium tuberculosis physiology and virulence. *Nat Rev Microbiol* **12**, 729-737 (2014).

42. Rens C, Chao JD, Sexton DL, Tocheva EI, Av-Gay Y. Roles for phthiocerol dimycocerosate lipids in Mycobacterium tuberculosis pathogenesis. *Microbiology (Reading)* **167**, (2021).

43. Song N, Cui Y, Li Z, Chen L, Liu S. New Targets and Cofactors for the Transcription Factor LrpA from Mycobacterium tuberculosis. *DNA Cell Biol* **35**, 167-176 (2016).

44. Massey LK, Sokatch JR, Conrad RS. Branched-chain amino acid catabolism in bacteria. *Bacteriol Rev* **40**, 42-54 (1976).

45. Yang H, Zhou Y, Luo Q, Zhu C, Fang B. L-leucine increases the sensitivity of drug-resistant Salmonella to sarafloxacin by stimulating central carbon metabolism and increasing intracellular reactive oxygen species level. *Front Microbiol* **14**, 1186841 (2023).

46. Lee W, VanderVen BC, Fahey RJ, Russell DG. Intracellular Mycobacterium tuberculosis exploits host-derived fatty acids to limit metabolic stress. *J Biol Chem* **288**, 6788-6800 (2013).

47. Eoh H, Rhee KY. Methylcitrate cycle defines the bactericidal essentiality of isocitrate lyase for survival of Mycobacterium tuberculosis on fatty acids. *Proc Natl Acad Sci U S A* **111**, 4976-4981 (2014).

984 48. Ehebauer MT, *et al.* Characterization of the mycobacterial acyl-CoA carboxylase holo
985 complexes reveals their functional expansion into amino acid catabolism. *PLoS*
986 *Pathog* **11**, e1004623 (2015).
987

988 49. Jain M, *et al.* Lipidomics reveals control of Mycobacterium tuberculosis virulence
989 lipids via metabolic coupling. *Proc Natl Acad Sci U S A* **104**, 5133-5138 (2007).
990

991 50. Cumming BM, Addicott KW, Adamson JH, Steyn AJ. Mycobacterium tuberculosis
992 induces decelerated bioenergetic metabolism in human macrophages. *Elife* **7**, (2018).
993

994 51. Nosik M, *et al.* Decreased IL-1 beta Secretion as a Potential Predictor of Tuberculosis
995 Recurrence in Individuals Diagnosed with HIV. *Biomedicines* **12**, (2024).
996

997 52. Choudhary E, Thakur P, Pareek M, Agarwal N. Gene silencing by CRISPR
998 interference in mycobacteria. *Nat Commun* **6**, 6267 (2015).
999

1000 53. Pal P, *et al.* ResR/McdR-regulated protein translation machinery contributes to drug
1001 resilience in Mycobacterium tuberculosis. *Commun Biol* **6**, 708 (2023).
1002

1003 54. Steyn AJ, Joseph J, Bloom BR. Interaction of the sensor module of Mycobacterium
1004 tuberculosis H37Rv KdpD with members of the Lpr family. *Mol Microbiol* **47**, 1075-
1005 1089 (2003).
1006

1007 55. Wick RR, *et al.* Trycycler: consensus long-read assemblies for bacterial genomes.
1008 *Genome Biol* **22**, 266 (2021).
1009

1010 56. Khoshnood S, *et al.* Mechanism of Action, Resistance, Synergism, and Clinical
1011 Implications of Delamanid Against Multidrug-Resistant Mycobacterium tuberculosis.
1012 *Front Microbiol* **12**, 717045 (2021).
1013

1014 57. Wick RR, Holt KE. Benchmarking of long-read assemblers for prokaryote whole
1015 genome sequencing. *F1000Res* **8**, 2138 (2019).
1016

1017 58. Ruan J, Li H. Fast and accurate long-read assembly with wtdbg2. *Nat Methods* **17**,
1018 155-158 (2020).
1019

1020 59. Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO:
1021 assessing genome assembly and annotation completeness with single-copy orthologs.
1022 *Bioinformatics* **31**, 3210-3212 (2015).
1023

1024 60. Phelan JE, *et al.* Integrating informatics tools and portable sequencing technology for
1025 rapid detection of resistance to anti-tuberculous drugs. *Genome Med* **11**, 41 (2019).
1026

1027 61. Garrison E, Marth G. Haplotype-based variant detection from short-read sequencing.
1028 *arXiv* **12073907**, (2012).
1029

1030 62. McKenna A, *et al.* The Genome Analysis Toolkit: a MapReduce framework for
1031 analyzing next-generation DNA sequencing data. *Genome Res* **20**, 1297-1303 (2010).
1032

1033

Compound	MIC
Bleomycin (sulfate)	<0.024µM
Rifabutin	<0.024µM
Rifaximin	<0.024µM
Bleomycin (hydrochloride)	<0.025µM
Rifapentine	<0.025µM
BTZ043	<0.02µM
Macozinone	<0.02µM
Semapimod	<0.02µM

a

b

c

d

ppsB: Ile896Asn
ppe60: (Multiple mutations)
Rv3837c: Thr228Ile
mce1D: Gln225*

a

b

WT

Sem^R

a L-leucine

PpsA PpsB PpsC PpsD PpsE

Semapimod

b

?

PpsA PpsB PpsC PpsD PpsE

Successive Passaging

ppsB O/E

c

PpsA PpsB* PpsC PpsD PpsE