Rapid Eye Movement Sleep Displays Distinct Fractal Dynamics Between Phasic and Tonic States

Yiqing Lu[1,2], Liang Wu[1,2], Jingyu Liu[1,2], Yongcheng Li[3], Yaping Huai[1, 2*]

1 Department of Rehabilitation Medicine, Shenzhen Longhua District Central Hospital, Shenzhen, Guangdong, China

2 Shenzhen Longhua District Rehabilitation Medical Equipment Development and Transformation Joint Key Laboratory, Shenzhen, Guangdong, China

3 CAS Key Laboratory of Human-Machine Intelligence-Synergy Systems, Shenzhen Institutes of Advanced Technology (SIAT), Chinese Academy of Sciences (CAS), Shenzhen, Guangdong, China

* Corresponding authors:

Yaping Huai:

Department of Rehabilitation Medicine, Shenzhen Longhua District Central Hospital,

Guanlan Avenue No. 187, Shenzhen, 518110, Guangdong, China

Email: huaiyaping@163.com

Abstract

Rapid eye movement (REM) sleep consists of phasic and tonic microstates with unique neurophysiological properties, yet their fractal characteristics remain underexplored. Using Higuchi's fractal dimension (HFD) analysis of electroencephalographic (EEG) data from healthy adults, this study investigated complexity differences between REM microstates. The results showed that both phasic and tonic REM exhibited significantly lower global HFD values compared to wakefulness, while displaying similar overall complexity levels between microstates. Importantly, phasic REM demonstrated regionally specific reductions in fractal dimensionality, with pronounced decreases observed in frontocentral areas. These localized reductions exhibited a negative association with theta band power, yet remained statistically unrelated to Lempel-Ziv complexity (LZC) measures, indicating that HFD and LZC capture distinct aspects of neural signal organization. The findings reveal that although phasic and tonic REM maintain comparable global complexity, they differ in their spatiotemporal fractal patterns. The association between increased theta power and reduced fractal dimensionality suggests that phasic REM represents a neurophysiological state favoring rhythmic regularity, potentially optimized for internal information processing. These results position HFD as a valuable complementary approach for characterizing REM microstates, with potential applications in understanding sleep disorder pathophysiology.

Keywords: Complexity; EEG; Fractal dimension; Nonlinear dynamics; REM Sleep

1. Introduction

Sleep is categorized into distinct stages, differentiated by neurophysiological signatures such as characteristic electroencephalographic (EEG) patterns and concomitant autonomic activity (Le Bon, 2020). Among these, rapid eye movement (REM) sleep is subdivided into phasic and tonic microstates (phasic REM and tonic REM), each exhibiting unique neural dynamics. Phasic REM is marked by transient events, including ponto-geniculo-occipital (PGO) waves, autonomic fluctuations, and muscle twitches (Dickerson et al., 1993; Fernandez-Mendoza et al., 2009; Gott et al., 2017), whereas tonic REM reflects a stable neurophysiological state with diminished motor activity (Simor et al., 2020). These microstates are not only linked to cognitive functions like memory consolidation (Boyce et al., 2016; Klinzing et al., 2019; van den Berg et al., 2023) and emotional regulation (Leitner et al., 2025; Wassing et al., 2019) but also implicated in disorders such as REM sleep behavior disorder (RBD) and insomnia (Dauvilliers et al., 2018; Feige et al., 2018; Simor et al., 2020; Yang et al., 2024).

Research on REM neurophysiology has employed both linear and nonlinear analytical frameworks. Linear spectral analyses identify distinct EEG power distributions between phasic and tonic REM, demonstrating heightened alpha and beta oscillations during tonic states and increased gamma activity in phasic states (Jouny et al., 2000; Simor et al., 2016). Non-periodic neural activity during REM sleep, distinguished by steeper low-frequency band slopes and flatter high-frequency slopes in phasic states, further provides a novel biomarker for differentiating these states (Rosenblum et al., 2024). Event-related potential (ERP) studies reveal preserved auditory processing during tonic REM in contrast to diminished responses in phasic states (Sallinen et al., 1996; Takahara et al., 2002). Extending these insights, nonlinear approaches such as Lempel-Ziv complexity (LZC) uncover reduced EEG complexity in phasic REM compared to tonic REM, correlating inversely with delta power and positively with alpha activity (Lu et al., 2024). Together, these findings demonstrate distinct information processing mechanisms across REM microstates.

Despite advances in nonlinear EEG characterization, the fractal properties of REM microstates remain underexplored. Fractal analysis, particularly Higuchi's fractal dimension (HFD), offers a robust framework to quantify signal complexity by assessing self-similarity patterns in time-series data, a characteristic inherent to nonlinear systems such as the brain (Lau et al., 2022; Rodriguez-Bermudez and Garcia-Laencina, 2015). Unlike LZC, which evaluates sequence compressibility, HFD captures the irregularity and hierarchical structure of neural activity, providing complementary insights into the dynamical organization of sleep stages (Armonaite et al., 2023; Olejarczyk et al., 2022). This approach proves effective in distinguishing pathological from healthy brain states (Cukic et al., 2020; Gomez et al., 2009; Kawe et al., 2019), and may elucidate how phasic and tonic REM contribute to sleep-related disorders.

This study employs HFD to quantify the fractal properties of phasic and tonic REM sleep EEG signals. Expanding upon our prior LZC analyses, which identified distinct complexity patterns in REM microstates, we systematically assess fractal dynamics to differentiate phasic and tonic REM, thereby refining their nonlinear neurophysiological profiles. By integrating HFD methodology, this investigation advances the understanding of REM sleep dynamics, offering novel insights into their role in neuroregulatory mechanisms and pathological manifestations of sleep disorders.

2. Materials and Methods

2.1. Dataset

We analyzed a publicly available REM EEG dataset (https://osf.io/2vptx) originally collected by Simor et al. (2021). The dataset comprises whole-night polysomnographic recordings from 40 healthy young adults (Study 1: $N = 20$, 10 males and 10 females, 21.7±1.4 years; Study 2: $N = 20$, 6 males and 14 females, 21.6±1.6 years). All participants met strict health criteria and provided written informed consent under local university ethics board approval. For full methodological details, see Simor et al. (2021).

2.2. EEG acquisition and preprocessing

EEG data were acquired using 19 electrodes (10-20 system, Study 1) or 17 electrodes (Fp1/Fp2 were not used, Study 2), with impedances <8 kΩ. Signals were digitized at 4096 Hz and downsampled (Study 1: 1024 Hz; Study 2: 512 Hz). Sleep staging followed the American Academy of Sleep Medicine (AASM) criteria, with REM periods classified as phasic (4-second segments with eye movement bursts meeting amplitude criteria) or tonic (4-second segments without notable bursts). Each Study 1 participant contributed 100 segments per state; Study 2 required ≥6 minutes per condition. For comparisons, we analyzed awake periods (eyes-closed only), including pre-sleep onset wakefulness and post-sleep onset wakefulness when available, applying identical preprocessing pipelines as used for REM segments. Three participants were excluded from Study 2 due to insufficient trials, resulting in final samples of $N = 20$ (Study 1) and $N = 17$ (Study 2). Artifact removal was performed using independent component analyses (ICA) implemented through EEGLAB and FieldTrip toolboxes (Delorme and Makeig, 2004; Oostenveld et al., 2011), where 2-4 independent components associated with eye movements were identified based on topography and waveform characteristics.

2.3. Estimation of HFD analysis

We quantified EEG signal complexity using HFD, a nonlinear measure that estimates the fractal characteristics of time series data. Following Higuchi's algorithm (Higuchi, 1988), the fractal

dimension was derived by calculating the mean curve length of the signal at multiple temporal scales (defined by parameter k), where k represents the number of intervals used to subsample the time series. The slope of the logarithmic relationship between scale (k) and curve length yields the HFD value. For each participant, HFD values were computed across 4-second epochs for each electrode, with the maximum scale parameter k_{max} set to 30 based on empirical optimization. Condition-specific averages (phasic REM, tonic REM, wakefulness) were obtained by averaging HFD values across epochs within each state. This analysis was implemented using custom MATLAB scripts.

2.4. Spectral analysis

We examined REM-related EEG spectral changes across all electrodes using Hanning-tapered Fourier transforms with adaptive frequency smoothing (5-cycle windows, 1 Hz resolution). Power was computed for frequency bands: delta (1-4 Hz), theta (4-7 Hz), alpha (8-12 Hz), beta (13-30 Hz), and gamma (31-40 Hz). Grand averages were calculated per condition after excluding three participants (one in Study 1; two in Study 2) due to technical artifacts, yielding final samples of 19 (Study 1) and 15 (Study 2) participants for spectral and correlation analyses.

2.5. Correlation analysis

We computed condition differences (phasic minus tonic REM) for both HFD and spectral power within significant electrode clusters, then assessed their relationship across participants using Pearson's correlation. Additionally, we examined global HFD-LZC correlations by averaging across all electrodes (phasic minus tonic REM) and performing Pearson's correlation.

2.6. Statistical analysis

HFD values were computed from 4-second EEG epochs using custom MATLAB scripts, averaged per condition (phasic REM, tonic REM, wakefulness) either by electrode or globally. Condition effects were assessed via one-way repeated-measures ANOVA with Greenhouse-Geisser correction when the assumption of sphericity was violated (Mauchly's test). Post-hoc pairwise comparisons were examined using two-tailed paired t-tests, followed by Bonferroni correction. To address the multiple comparisons problem inherent in EEG data analysis, we implemented a cluster-based non-parametric randomization test (Maris and Oostenveld, 2007). The analysis employed a paired-sample design with an alpha level of $\alpha = 0.05$ defining statistical significance. Spatiotemporal clusters were formed by aggregating adjacent electrodes and time points demonstrating effects below the predetermined threshold, with cluster-level statistics derived from the summation of t-values within each contiguous region. For null hypothesis testing, we performed 5,000 Monte Carlo simulations by randomly shuffling condition labels (phasic vs. tonic REM) to construct an empirical distribution of maximum cluster statistics. Final significance determination involved comparing the observed cluster statistics against the 95th percentile of this permuted null distribution, thereby controlling family-wise error rate while maintaining sensitivity to genuine neural effects. All correlation analyses employed false discovery rate (FDR) correction for multiple comparisons

(Benjamini and Hochberg, 1995). Visualizations combined FieldTrip functions with custom scripts.

3. Results

3.1. Reduced fractal dimensionality in REM sleep states

A one-way repeated measures ANOVA was conducted to examine differences in HFD across three states: phasic REM sleep, tonic REM sleep, and wakefulness. The HFD values were calculated as the grand average across all EEG channels, representing the overall complexity of neural activity at the whole-brain level. Mauchly's test indicated violations of the sphericity assumption in both studies (Study 1: $W = 0.091$, $p < 0.001$; Study 2: $W = 0.224$, $p < 0.001$), prompting Greenhouse-Geisser corrections. Both studies revealed strong main effects of state on HFD (Study 1: $F(1.048,19.909) = 12.853$, $p = 0.002$, $\eta^2 = 0.404$; Study 2: $F(1.126,21.394) = 46.083$, $p < 0.001$, $\eta^2 = 0.708$), indicating that neural complexity, as measured by HFD, varied significantly across the different states. Post-hoc analyses employing Bonferroni-corrected pairwise comparisons revealed a highly consistent pattern of results across both studies (Figure 1): no statistically significant differences were observed between phasic and tonic REM sleep states (Study 1: $p = 0.202$; Study 2: $p = 0.067$). However, both phasic and tonic REM sleep showed significantly reduced HFD compared to wakefulness (Study 1: mean differences = -0.041 for phasic REM, -0.037 for tonic REM, both $p < 0.01$; Study 2: mean difference = -0.073 for phasic REM, -0.065 for tonic REM, both $p < 0.001$).

Figure 1. *Distribution of individual HFD (averaged across EEG electrodes) across phasic REM, tonic REM, and wakefulness conditions. (A) Study 1; (B) Study 2. For each condition, the density trace shows the smoothed data distribution. The boxplots display the distribution using median, interquartile range (IQR), and 1.5×IQR whiskers, with outliers plotted as individual circles when present; individual data points are jittered vertically. A one-way*

repeated-measures ANOVA with post-hoc testing was employed to determine statistical significance. Both phasic REM sleep and tonic REM sleep demonstrated significantly lower HFD values compared to wakefulness.

3.2. Spatial distribution of HFD differences between phasic and tonic REM

Following our initial analysis of grand average HFD across all EEG channels, we next examined the topographic distribution of HFD differences between phasic and tonic REM periods. cluster-based non-parametric randomization test revealed significantly lower HFD during phasic REM compared to tonic REM (negative clusters, $p < 0.05$; see Figure 2). In Study 1, this reduction was most pronounced in a frontocentral cluster that included prefrontal (Fp1, Fp2), frontal (F3, Fz, F4), and central (C3, Cz) regions ($t(19)$ = -20.921, p = 0.017). The mean phasic HFD in this cluster (1.128 ± 0.025) was 0.62% lower than during tonic periods (1.135 ± 0.023). Study 2 demonstrated a similar but more extensive pattern of reduced phasic HFD ($t(16)$ = -30.471, p = 0.008), with a significant cluster distributed across frontal, central, and parietal regions (F7/F8, F3/F4, Fz, C3/C4, Cz, P3/P4). Within this cluster, phasic HFD value (1.263 ± 0.037) was 0.86% lower than tonic value (1.274 ± 0.035), further supporting the robustness of this difference.

Figure 2. *Reduced HFD during phasic REM compared to tonic REM. (A, B) Study 1; (C, D) Study 2. A & C: Topographical maps display mean HFD values per electrode (Fp1 and Fp2 were not recorded in study 2), with difference maps (Diff) highlighting phasic-tonic REM contrasts. Black asterisks denote electrodes with significantly lower HFD in phasic REM (p < 0.05, cluster-based non-parametric randomization test). B & D: Plots of HFD (averaged across significant electrodes) for individual participants (dots), with boxplots indicating median, IQR, and 1.5×IQR whiskers. Density traces illustrate data distributions. In both studies, phasic REM consistently exhibited reduced HFD relative to tonic REM.*

3.3. Correlational Analysis of HFD with Spectral and Complexity Measures

Following our examination of grand average HFD and its spatial distribution across phasic and tonic REM periods, we further investigated the functional relationships between HFD and spectral power across multiple frequency bands. Particularly, among all tested bands (delta, theta, alpha, beta, and gamma), only theta power showed a significant negative correlation (FDR corrected) with HFD within the identified electrode clusters. This inverse relationship was consistent across both studies (Study 1: $r = -0.699$, $p = 9.0 \times 10^{-4}$; Study 2: $r = -0.771$, $p = 8.0 \times 10^{-4}$; see Figure 3), suggesting a specific coupling between decreased signal complexity and increased theta-band activity during phasic REM (relative to tonic REM).

Figure 3. *Neurophysiological coupling between HFD and spectral power. (A, B) Study 1 & 2. Scatter plots display condition-dependent changes (phasic minus tonic REM) in HFD versus spectral power amplitude for each participant (dots represent electrode cluster averages). Significant negative correlations between HFD and theta power were observed in both studies (Pearson's r and p-values are presented; *p < 0.01, FDR-corrected), indicating that phasic REM-related HFD decreases co-occurred with theta power increases.*

Given our prior work demonstrating the relevance of LZC in REM sleep dynamics (Lu et al., 2024), we additionally examined whether the phasic-tonic HFD differences were related to corresponding LZC differences (averaged across all electrodes). However, no significant association emerged (Figure 4), indicating that these complexity measures may capture distinct aspects of neural activity.

Figure 4. Absence of significant correlation between HFD and LZC. (A, B) Study 1 & 2. Scatter plots display condition-dependent changes (phasic minus tonic REM) in HFD versus LZC for each participant (dots represent all electrode averages). No significant correlations were observed between HFD and LZC measures in either study (Pearson's r and p-values are presented; all p>0.05).

4. Discussion

The present study provides a comprehensive examination of fractal neural dynamics during REM sleep through HFD analysis, revealing distinct nonlinear EEG characteristics that differentiate phasic and tonic microstates. Our results demonstrate that phasic REM exhibits localized decreases in HFD within frontocentral regions, suggesting microstate-specific modifications in neural signal processing. Both phasic and tonic REM displayed lower fractal dimensions relative to wakefulness, their overall complexity measures did not differ significantly at the global level. Furthermore, we identified an inverse relationship between HFD and theta-band power, independent of LZC measures. These results advance our understanding of REM sleep neurodynamics and highlight the utility of fractal analysis in dissecting neural complexity.

The observed reduction in HFD during REM sleep aligns with prior studies showing diminished nonlinear complexity during sleep compared to wakefulness (Aamodt et al., 2022; Lu et al., 2024; Ma et al., 2018). However, the lack of global HFD differences between phasic and tonic REM diverges from previous LZC results (Lu et al., 2024), suggesting that these metrics assess different neural characteristics. This distinction arises from their underlying computational principles: LZC measures temporal unpredictability through sequence compressibility (Lempel and Ziv, 1976), while HFD evaluates self-similarity patterns across temporal scales (Higuchi, 1988). This divergence implies that REM microstates may share comparable overall complexity but differ in hierarchical signal organization, which is supported by the spatially specific HFD reductions in phasic REM. The frontocentral topography of these differences overlaps with regions implicated in

REM-related cognitive processes, including memory consolidation and emotional regulation (Eichenlaub et al., 2018; Kim et al., 2020), suggesting that localized fractal alterations may reflect state-specific neurocomputational demands. The statistical independence of HFD and LZC measurements highlights the complementary nature of these metrics, as LZC primarily reflects temporal unpredictability whereas HFD captures multiscale self-similarity, with their combined use providing a more comprehensive assessment of neural complexity.

The negative correlation between HFD and theta power within phasic REM clusters (compared to tonic REM) provides mechanistic insights into the neurophysiological processes governing REM neurodynamics. Theta oscillations, known to facilitate hippocampal-neocortical communication during memory processing (Boyce et al., 2016), exhibit increased spectral power in phasic REM (Simor et al., 2016). This increased theta activity may drive reduced fractal dimensionality by introducing rhythmic regularity to otherwise irregular neural activity. The rhythmic regularity imposed by theta dominance appears to facilitate the coordination of large-scale neural assemblies for memory reprocessing while promoting perceptual isolation from external inputs through oscillatory entrainment. This sensory disengagement is supported by evidence indicating that phasic REM reflects a neurobiological state preferentially oriented toward internal information processing (Simor et al., 2020; Wehrle et al., 2007), as demonstrated by both attenuated sensory evoked potentials (Sallinen et al., 1996; Takahara et al., 2002) and increased activation of limbic networks (Corsi-Cabrera et al., 2016; Wassing et al., 2019). The reduction in fractal dimensionality may thus represent an adaptive neural signature of this shifted processing priority, where increased theta synchrony concurrently optimizes memory-related circuit integration and attenuates exteroceptive signaling. These findings reveal how the interaction between fractal properties and oscillatory dynamics modulates both information-processing capacity and phenomenological characteristics during distinct REM microstates.

Clinically, the differential fractal properties of REM microstates may inform sleep disorder pathophysiology. For instance, the frontocentral HFD reductions observed here are consistent with EEG abnormalities reported in RBD (Peng et al., 2021; Valomon et al., 2021), suggesting that these fractal alterations may represent a characteristic neurophysiological signature of RBD pathology. The distinct neurophysiological signatures captured by HFD and LZC provide complementary biomarkers that could enhance differential diagnosis. Specifically, HFD's sensitivity to theta power modulations may prove particularly relevant for disorders involving thalamocortical dysregulation, while LZC's association with alpha activity, as observed from our previous results, could better characterize disorders of cortical integration (Lu et al., 2024). Future studies should investigate whether these fractal measures can predict conversion from prodromal RBD to neurodegenerative syndromes, stratify insomnia subtypes based on their neurophysiological profiles, and monitor treatment response in sleep disorders targeting REM mechanisms. Additionally, the development of standardized HFD protocols across clinical populations will be crucial for translating these research

findings into practical diagnostic tools.

Several methodological constraints should be considered when interpreting these findings. First, while HFD provides valuable insights into the fractal properties of neural activity, its interpretation remains constrained by the lack of spatial specificity inherent in scalp EEG. Combining HFD with multimodal neuroimaging techniques could help localize these fractal dynamics to specific neural circuits, thereby clarifying their functional significance. Second, the restriction to healthy young adults of this study limits the extrapolation of these findings to clinical populations and older individuals, who may exhibit distinct REM sleep characteristics and fractal dynamics. Future studies should include more heterogeneous populations to validate and extend these observations.

In conclusion, this study demonstrates that fractal analysis using HFD provides unique insights into the neurophysiological distinction between phasic and tonic REM sleep. By complementing spectral and complexity metrics, HFD enhances our capacity to characterize the multiscale organization of neural activity during REM sleep, offering new avenues for understanding its role in cognitive processes and sleep-related pathologies.

Author Contributions

Conceptualization, Y.Lu. and Y.H.; methodology, Y.Lu.; software, Y.Lu. and Y.Li.; validation, Y.Lu., L.W. and J.L.; formal analysis, Y.Lu. and L.W.; investigation, Y.Lu., J.L. and Y.Li.; resources, Y.Lu. and Y.H.; data curation, Y.Lu.; writing—original draft preparation, Y.Lu.; writing—review and editing, Y.Lu. and Y.H.; visualization, Y.Lu.; supervision, Y.Lu. and Y.H.; project administration, Y.Lu. and Y.H.; funding acquisition, Y.Lu., L.W. and Y.H. All authors have read and agreed to the published version of the manuscript.

Funding

This research was funded by the Scientific Research Projects of Medical and Health Institutions of Longhua District, Shenzhen (2024001, 2024004), and the Shenzhen Science and Technology Program (JCYJ20210324123414039, JCYJ20240813152959014, KCXFZ20230731093300001).

Data Availability Statement

The dataset was originally released by Simor et al. (2021): Interoception in REM microstates. Accessible at osf.io/2vptx.

Acknowledgments

We gratefully acknowledge Simor et al. (2021) for making publicly available the EEG dataset used in this research, which was accessed via the Open Science Framework (OSF).

Conflicts of Interest

The authors declare no conflicts of interest.

References

Aamodt A, Sevenius Nilsen A, Markhus R, Kusztor A, HasanzadehMoghadam F, Kauppi N, Thurer B, Storm JF, Juel BE, 2022, EEG Lempel-Ziv complexity varies with sleep stage, but does not seem to track dream experience. Front Hum Neurosci 16:987714. doi:10.3389/fnhum.2022.987714

Armonaite K, Nobili L, Paulon L, Balsi M, Conti L, Tecchio F, 2023, Local neurodynamics as a signature of cortical areas: new insights from sleep. Cereb Cortex 33:3284-3292. doi:10.1093/cercor/bhac274

Benjamini Y, Hochberg Y, 1995, Controlling the false discovery rate: a practical and powerful approach to multiple testing. J R Stat Soc B 57:289-300. doi:10.1111/j.2517-6161.1995.tb02031.x

Boyce R, Glasgow SD, Williams S, Adamantidis A, 2016, Causal evidence for the role of REM sleep theta rhythm in contextual memory consolidation. Science 352:812-816. doi:10.1126/science.aad5252

Corsi-Cabrera M, Velasco F, Del Rio-Portilla Y, Armony JL, Trejo-Martinez D, Guevara MA, Velasco AL, 2016, Human amygdala activation during rapid eye movements of rapid eye movement sleep: an intracranial study. J Sleep Res 25:576-582. doi:10.1111/jsr.12415

Cukic M, Stokic M, Radenkovic S, Ljubisavljevic M, Simic S, Savic D, 2020, Nonlinear analysis of EEG complexity in episode and remission phase of recurrent depression. Int J Methods Psychiatr Res 29:e1816. doi:10.1002/mpr.1816

Dauvilliers Y, Schenck CH, Postuma RB, Iranzo A, Luppi PH, Plazzi G, Montplaisir J, Boeve B, 2018, REM sleep behaviour disorder. Nat Rev Dis Primers 4:19. doi:10.1038/s41572-018-0016-5

Delorme A, Makeig S, 2004, EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. J Neurosci Methods 134:9-21. doi:10.1016/j.jneumeth.2003.10.009

Dickerson LW, Huang AH, Thurnher MM, Nearing BD, Verrier RL, 1993, Relationship between coronary hemodynamic changes and the phasic events of rapid eye movement sleep. Sleep 16:550-557. doi:10.1093/sleep/16.6.550

Eichenlaub JB, van Rijn E, Gaskell MG, Lewis PA, Maby E, Malinowski JE, Walker MP, Boy F, Blagrove M, 2018, Incorporation of recent waking-life experiences in dreams correlates with frontal theta activity in REM sleep. Soc Cogn Affect Neurosci 13:637-647. doi:10.1093/scan/nsy041

Feige B, Nanovska S, Baglioni C, Bier B, Cabrera L, Diemers S, Quellmalz M, Siegel M, et al., 2018, Insomnia-perchance a dream? Results from a NREM/REM sleep awakening study in good sleepers

and patients with insomnia. Sleep 41. doi:10.1093/sleep/zsy032

Fernandez-Mendoza J, Lozano B, Seijo F, Santamarta-Liebana E, Ramos-Platon MJ, Vela-Bueno A, Fernandez-Gonzalez F, 2009, Evidence of subthalamic PGO-like waves during REM sleep in humans: a deep brain polysomnographic study. Sleep 32:1117-1126. doi:10.1093/sleep/32.9.1117

Gomez C, Mediavilla A, Hornero R, Abasolo D, Fernandez A, 2009, Use of the Higuchi's fractal dimension for the analysis of MEG recordings from Alzheimer's disease patients. Med Eng Phys 31:306-313. doi:10.1016/j.medengphy.2008.06.010

Gott JA, Liley DT, Hobson JA, 2017, Towards a Functional Understanding of PGO Waves. Front Hum Neurosci 11:89. doi:10.3389/fnhum.2017.00089

Higuchi T, 1988, Approach to an irregular time series on the basis of the fractal theory. Physica D 31:277-283. doi:10.1016/0167-2789(88)90081-4

Jouny C, Chapotot F, Merica H, 2000, EEG spectral activity during paradoxical sleep: further evidence for cognitive processing. Neuroreport 11:3667-3671. doi:10.1097/00001756-200011270-00016

Kawe TNJ, Shadli SM, McNaughton N, 2019, Higuchi's fractal dimension, but not frontal or posterior alpha asymmetry, predicts PID-5 anxiousness more than depressivity. Sci Rep 9:19666. doi:10.1038/s41598-019-56229-w

Kim SY, Kark SM, Daley RT, Alger SE, Reboucas D, Kensinger EA, Payne JD, 2020, Interactive effects of stress reactivity and rapid eye movement sleep theta activity on emotional memory formation. Hippocampus 30:829-841. doi:10.1002/hipo.23138

Klinzing JG, Niethard N, Born J, 2019, Mechanisms of systems memory consolidation during sleep. Nat Neurosci 22:1598-1610. doi:10.1038/s41593-019-0467-3

Lau ZJ, Pham T, Chen SHA, Makowski D, 2022, Brain entropy, fractal dimensions and predictability: A review of complexity measures for EEG in healthy and neuropsychiatric populations. Eur J Neurosci 56:5047-5069. doi:10.1111/ejn.15800

Le Bon O, 2020, Relationships between REM and NREM in the NREM-REM sleep cycle: a review on competing concepts. Sleep Med 70:6-16. doi:10.1016/j.sleep.2020.02.004

Leitner C, Greco V, Casoni F, Lewis PA, Ferini-Strambi L, Galbiati A, 2025, Isolated Rem Sleep Behavior Disorder: A Model to Assess the Overnight Habituation of Emotional Reactivity. Clocks Sleep 7. doi:10.3390/clockssleep7010009

Lempel A, Ziv J, 1976, On the complexity of finite sequences. IEEE Trans Inf Theory 22:75-81. doi:10.1109/TIT.1976.1055501

Lu Y, Yang W, Zhang X, Wu L, Li Y, Wang X, Huai Y, 2024, Unraveling the complexity of rapid eye movement microstates: insights from nonlinear EEG analysis. Sleep 47. doi:10.1093/sleep/zsae105

Ma Y, Shi W, Peng CK, Yang AC, 2018, Nonlinear dynamical analysis of sleep electroencephalography using fractal and entropy approaches. Sleep Med Rev 37:85-93. doi:10.1016/j.smrv.2017.01.003

Maris E, Oostenveld R, 2007, Nonparametric statistical testing of EEG- and MEG-data. J Neurosci Methods 164:177-190. doi:10.1016/j.jneumeth.2007.03.024

Olejarczyk E, Gotman J, Frauscher B, 2022, Region-specific complexity of the intracranial EEG in the sleeping human brain. Sci Rep 12:451. doi:10.1038/s41598-021-04213-8

Oostenveld R, Fries P, Maris E, Schoffelen JM, 2011, FieldTrip: Open source software for advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell Neurosci 2011:156869. doi:10.1155/2011/156869

Peng A, Wang R, Huang J, Wu H, Chen L, 2021, Abnormalities of Resting-State Electroencephalographic Microstate in Rapid Eye Movement Sleep Behavior Disorder. Front Hum Neurosci 15:728405.

doi:10.3389/fnhum.2021.728405

Rodriguez-Bermudez G, Garcia-Laencina PJ, 2015, Analysis of EEG signals using nonlinear dynamics and chaos: a review. Appl Math Inf Sci 9:2309. doi:10.12785/amis/090512

Rosenblum Y, Bogdany T, Nadasy LB, Chen X, Kovacs I, Gombos F, Ujma P, Bodizs R, et al., 2024, Aperiodic neural activity distinguishes between phasic and tonic REM sleep. J Sleep Res:e14439. doi:10.1111/jsr.14439

Sallinen M, Kaartinen J, Lyytinen H, 1996, Processing of auditory stimuli during tonic and phasic periods of REM sleep as revealed by event-related brain potentials. J Sleep Res 5:220-228. doi:10.1111/j.1365-2869.1996.00220.x

Simor P, Bogdany T, Bodizs R, Perakakis P, 2021, Cortical monitoring of cardiac activity during rapid eye movement sleep: the heartbeat evoked potential in phasic and tonic rapid-eye-movement microstates. Sleep 44. doi:10.1093/sleep/zsab100

Simor P, Gombos F, Szakadat S, Sandor P, Bodizs R, 2016, EEG spectral power in phasic and tonic REM sleep: different patterns in young adults and children. J Sleep Res 25:269-277. doi:10.1111/jsr.12376

Simor P, van der Wijk G, Nobili L, Peigneux P, 2020, The microstructure of REM sleep: Why phasic and tonic? Sleep Med Rev 52:101305. doi:10.1016/j.smrv.2020.101305

Takahara M, Nittono H, Hori T, 2002, Comparison of the event-related potentials between tonic and phasic periods of rapid eye movement sleep. Psychiatry Clin Neurosci 56:257-258. doi:10.1046/j.1440-1819.2002.00999.x

Valomon A, Riedner BA, Jones SG, Nakamura KP, Tononi G, Plante DT, Benca RM, Boly M, 2021, A high-density electroencephalography study reveals abnormal sleep homeostasis in patients with rapid eye movement sleep behavior disorder. Sci Rep 11:4758. doi:10.1038/s41598-021-83980-w

van den Berg NH, Gibbings A, Baena D, Pozzobon A, Al-Kuwatli J, Ray LB, Fogel SM, 2023, Eye movements during phasic versus tonic rapid eye movement sleep are biomarkers of dissociable electroencephalogram processes for the consolidation of novel problem-solving skills. Sleep 46. doi:10.1093/sleep/zsad151

Wassing R, Lakbila-Kamal O, Ramautar JR, Stoffers D, Schalkwijk F, Van Someren EJW, 2019, Restless REM Sleep Impedes Overnight Amygdala Adaptation. Curr Biol 29:2351-2358 e2354. doi:10.1016/j.cub.2019.06.034

Wehrle R, Kaufmann C, Wetter TC, Holsboer F, Auer DP, Pollmacher T, Czisch M, 2007, Functional microstates within human REM sleep: first evidence from fMRI of a thalamocortical network specific for phasic REM periods. Eur J Neurosci 25:863-871. doi:10.1111/j.1460-9568.2007.05314.x

Yang S, Liu GX, Zhou YS, Ma ZJ, Xia L, Zhang HL, Li XY, Kong XY, et al., 2024, Relationships between rumination and different types of rapid eye movement sleep in patients with chronic insomnia disorder. Sleep Med 117:162-168. doi:10.1016/j.sleep.2024.03.032