1
2
3   Newly trained navigation and verbal memory skills elicit changes in task-related

4   networks but not brain structure

6   Li Zheng[1,2], Zachary Boogaart[3], Andrew McAvan[1], Joshua Garren[1], Stephanie Doner[1], Bradley

7   J. Wilkes[5], Will Groves[4], Ece Yuksel[4], Lucia Cherep[1], & Arne D. Ekstrom[1,2*], & Steven M.

8   Weisberg[4,6*]

9   [1]Department of Psychology, University of Arizona, 1503 E. University Blvd.,
10   Tucson, AZ 85719
11   [2]Evelyn McKnight Brain Institute, University of Arizona, 1503 E. University Blvd.,
12   Tucson, AZ 85719
13   [3]The University of Miami Miller School of Medicine, University of Miami, 1600 NW 10th Ave
14   #1140, Miami, FL 33136
15   [4]Department of Psychology, University of Florida, 945 Center Dr., Gainesville, FL 32611
16   [5]Department of Applied Physiology & Kinesiology, University of Florida, 945 Center Dr.,
17   Gainesville, FL 32611
18   [6]Evelyn McKnight Brain Institute, University of Florida, Gainesville, FL 32611
19

20   *These two authors shared equally in the work

21   CORRESPONDING AUTHORS:
22   Dr. Steven M. Weisberg, email: stevenweisberg@ufl.edu
23   Dr. Arne Ekstrom, email: adekstrom@arizona.edu
24

25   Funding Sources: NIH/NINDS R21NS120237-01 to SMW and ADE; NIH/NIA K01AG070333-01
26   to SMW
27
CRediT statement: Li Zheng: Data curation, Methodology, Formal Analysis, Software,
Validation, Visualization, Writing – original draft, Writing – review & editing; Zachary Boogaart:
Investigation, Data curation, Methodology, Project administration, Software, Validation; Andrew
McAvan: Data curation, Methodology, Software, Validation; Joshua Garren: Investigation,
Project administration, Validation; Stephanie Doner: Data curation, Formal Analysis,
Investigation, Methodology, Project administration, Validation, Visualization; Will Groves:
Investigation, Project administration, Visualization; Ece Yuksel: Data curation, Formal Analysis,
Software, Validation, Visualization; Lucia Cherep: Investigation, Methodology, Project
administration, Supervision; Bradley Wilkes: Formal Analysis, Investigation, Software,
Visualization; Arne Ekstrom: Conceptualization, Formal Analysis, Funding acquisition,
Investigation, Methodology, Project administration, Resources, Supervision, Writing – original
draft, Writing – review & editing; Steven Weisberg: Conceptualization, Data curation, Formal
Analysis, Funding acquisition, Investigation, Methodology, Project administration, Resources,
Software, Supervision, Validation, Visualization, Writing – original draft, Writing – review &
editing

Abstract

Training cognitive skills, such as remembering a list of words or navigating a new city, has important implications for both health and disease. Yet, understanding what brain changes underlie the acquisition of complex cognitive skills remains unresolved. Here, we developed and validated intensive multiweek interventions in which participants were randomly assigned training in either navigation or verbal memory. Healthy young participants (N=75) underwent structural and functional imaging prior to and following the training. Based on pre-registered and exploratory analyses, we did not find any evidence for changes to hippocampal volume, hippocampal subfields, cortical brain volume, or white matter connectivity due to the training. In contrast, network-based analyses suggested changes in task-related informational connectivity, which occurred primarily between cortical areas and mostly involved putative cognitive control networks. These results suggest that cognitive interventions target more transient configurations in network connectivity rather than more durable structural changes.

Introduction

Improving cognitive skills can contribute positively to multiple aspects of daily life, such as education and professional endeavors[1,2]. Yet, the neural changes underlying cognitive skill acquisition remain an unresolved question in cognitive neuroscience, limiting our ability to target interventions designed to improve these skills. Some past studies have suggested a relationship between cognitive skills and focal gray matter volume[3-9], although many of these studies have involved comparison of experts with non-experts or specialized populations (e.g., older adults with cognitive decline). Therefore, it remains unclear whether gray matter volume differences underlie the behavioral changes that occur as part of new cognitive skill acquisition and whether the same mechanisms apply to healthy younger adults. Other studies have suggested the importance of changes in either the recruitment of specific brain areas (i.e., functional activation) or interactions between brain regions (i.e., functional connectivity) that may not result in macroscopic structural changes[1,10,11]. Resolving the neural mechanism(s) underlying the acquisition of new cognitive skills in healthy populations, with the potential for maximal behavioral improvement, is important so we can determine how to target interventions in future studies.

One possibility is that training different cognitive skills leads to changes in similar underlying neural mechanisms by training a cognitive process or enhancing function in a brain region common to both skills. Past studies suggest such overlap for a brain region thought to be broadly important to memory, the hippocampus[12,13]. People with better navigation and/or verbal memory skills typically have larger hippocampi [3,14,15], although this finding has not been consistently replicated[16,17]. Alternatively, it might be that learning different cognitive skills targets only partially overlapping or even independent brain networks, which in turn might hint at why transfer of newly trained cognitive skills to other cognitive domains is often modest[18]. For example, some studies suggest only partial overlap between navigation and verbal memory[19-21], and it is possible that some of the observed differences between experts and non-experts in past studies arose due to partial overlap between verbal memory and navigation. For example, when comparing expert navigators, street and landmark names would be intertwined with wayfinding skills, and thus it is important to determine whether superior verbal memory or navigation skills underlie greater hippocampal volume.

One issue with comparing experts and non-experts is that they involve between-participant designs, which may be underpowered compared to within-participant approaches[22] and may introduce other uncontrolled variables. One potential way to address this issue is to employ a within-participant manipulation to measure how a brain region such as the hippocampus (and connected brain networks) might change in the same participant as a result of training. Successful cognitive interventions suggest that targeted cognitive training, even for as little as 1-2 weeks, can result in improvements to specific cognitive functions. In some instances, interventions may even generalize to areas not explicitly trained but closely related to the training (termed "near transfer")[23-25]. We developed and pre-registered two cognitive training regimens in which we attempted to separately target navigation and verbal memory skills to allow us to test hypotheses related to the degree of neural overlap between the acquisition of these cognitive skills.

Here, we tested three different cohorts of healthy young adults using two different
cognitive interventions, each consisting of 10 training sessions targeting either
navigation or verbal memory. A third active control group watched educational videos
containing information about navigation and verbal memory but did not receive explicit
training[26,27]. Before and after training, participants underwent high-resolution structural
imaging targeting the hippocampus (T2), diffusion weighted imaging (DWI), whole brain
structural imaging (T1), and task-related functional imaging during a spatial and
temporal associative memory task to understand how the hippocampus and related
brain networks changed. We pre-registered three hypotheses, which are described in
more detail here (https://osf.io/etxvj): (1) targeted training over two weeks will result in
improvements in navigation or verbal memory, (2) improvements will be specific to the
trained task and will transfer to a separate behavioral measure of that skill ("near
transfer"), (3a) if the hippocampus is central to navigation, hippocampal volume should
increase from pre-test to post-test following navigation training but not verbal memory
training or the video control condition, (3b) if the hippocampus is central to verbal
memory, hippocampal volume should increase from pre-test to post-test in the Verbal
Memory group but not in the Navigation or Video Control groups, or (3c) alternatively,
hippocampal volume does not reflect acquisition of new verbal memory nor navigation
skills. As part of planned exploratory analyses, we also examined changes in structural
connectivity (DWI), gray matter volume outside of the medial temporal lobes, functional
changes to individual brain regions (i.e., univariate functional activation), and changes
to brain networks (i.e., task-related informational connectivity).

144   Results

145   Behavioral Results

146   Training-related effects

Participants were randomly assigned to one of three conditions, each of which
consisted of 10 training sessions, separated by at least one day over the course of
approximately 4–8 weeks (4.12 ± 1.87 weeks). Each training session lasted two hours.
In the navigation training condition, we recruited a total of 29 participants, with 27
successfully completing 10 sessions of navigation training in a large city-scale virtual
environment ("Virtual Arida", Fig. 1b left). Briefly, participants were placed at one
building and given the name of another building to find. Participants traveled to and from
that pair of buildings until they were able to take a path not exceeding 120% of the
shortest possible path. Upon reaching the criterion, they navigated between a new pair
of landmarks. After one subsection of Virtual Arida was learned completely, participants
were trained on integrating that subsection with previously learned subsections, with no
more than one subsection learned in a day (see Methods). To quantify the training
effect, we analyzed the average distance error travelled and normalized it by 120% of
the length of the shortest possible path. While all 27 participants learned subsections A
(although data was lost due to technical reasons for one participant), B, and A-B
integration, 26 participants progressed to C, while 23 made it to A-B-C integration. 19
participants made it to subsection D, and 14 participants made it to A-B-C-D integration.
Thus, to determine the training effect, we performed a repeated-measures ANOVA with
subsection as the only factor and threshold-normalized distance error as the dependent
measure.

We observed a significant main effect of subsection $F(2, 42) = 13.46$, $p < .001$, η^2
$= .39$; removing sex and site from the model did not change the effect (Supplemental
Note 2). Participants improved over the course of the training, showing higher error
initially for subsection A ($M = 79.09$, $SD = 33.22$) compared to subsection B ($M = 65.52$,
$SD = 26.89$, Fig. 1b right), which was significantly reduced for subsection C ($M = 47.44$,
$SD = 31.22$). This was also the case for subsections involving integration $F(1,22) =$
5.16, $p = .03$, $\eta^2 = .19$, although this finding did not survive inclusion of sex and site in
the model ($p = .06$). The distance error was reduced from subsections A-B ($M = 22.03$,
$SD = 18.47$) to A-B-C ($M = 15.17$, $SD = 14.47$, Fig. 1b right). In sum, participants
improved both in learning new subsections of our large virtual training environment and
in integrating multiple areas of the training environment together, suggesting that the
navigation training resulted in significant reductions in path error over the course of the
training.
In the verbal memory condition, we recruited a total of 32 participants, with 27
successfully completing 10 sessions of training. Briefly, participants viewed words for
one second (one at a time), completed math problems as a distractor after all words
were presented, and then recalled as many words as they could by typing them in any
order they wished. They were trained on a time-based method of loci ("temporal method
of loci") to improve their verbal memory[28]. During the first five sessions, participants
learned the same list of words, with the list growing in length after each successful trial
(Same Word List, Fig. 1a left). In the last 5 sessions, a new list of words was used for
each trial (New Word List, Fig. 1a left; see Methods). To quantify the training effect, we
conducted a linear regression analysis on the maximum number of correctly recalled
words for each day. The slopes of the regression lines for the first five days and the last
five days were calculated separately (Fig. 1a right), as the training paradigm differed
between these two phases (see Methods). Both slopes were significantly greater than
zero, indicating a significant improvement in memory performance over the training
period: first five days, $t(26) = 16.971$, $p < .001$; last five days, $t(26) = 23.579$, $p < .001$.
This effect remained significant after controlling for sex and site (Supplementary Note
2), suggesting robust improvements in verbal memory performance as a result of the
training.
In the active control condition, we recruited a total of 22 participants, with 21
successfully completing 10 sessions of watching videos. These videos included
interviews, lectures, documentaries, and TV series on topics related to memory and
navigation. After watching the videos for that session, participants responded to multiple
choice questions. We did not expect to find a training effect (i.e., improved performance
on the questions over sessions), but did find that the percentage of correct responses
each day was significantly higher than a 50% guess rate (Fig. 1c right, $ts(20) > 8.28$, ps
< 0.001, mean $= 88.58\%$, SD $= 5.82\%$), indicating that participants were engaged with
the video content.
207

208
Fig. 1 Experimental design. Participants were randomly assigned to one of three training
conditions as follows. (a) In the verbal memory training condition (n = 27), participants
underwent 10 sessions of verbal memory training involving word free recall after a distractor
task, with list complexity increasing over sessions. (b) In the navigation training condition (n =
27), participants trained in a large virtual environment, navigating between buildings until
optimal paths were learned. Subsections of the environment were integrated progressively
across sessions. (c) In the active control condition (n = 21), participants watched videos related
to memory and navigation, answering multiple-choice questions afterward, with accuracy
consistently above 50%, indicating engagement. Training schedules spanned 4–8 weeks, with
each session lasting two hours.
219
Pre-test and post-test behavioral transfer
We administered a set of cognitive tests related to navigation and verbal memory
before and after training to track transfer from the training. This involved learning a large
virtual environment not related to the training (i.e., Navigation Transfer task: Virtual
Silcton[29]) and learning an untrained list of 12 words (i.e., Verbal Memory Transfer task)
until all could be recalled perfectly (or after five trials). When we considered our pre-
registered dependent measures specific to verbal memory (average number of words
recalled, number of trials to criteria, and slope) and navigation (path error of Navigation
Transfer task, pointing error of Navigation Pointing task, and map accuracy of

Navigation Model Building task), we found some evidence of improvements in slope
specific to the verbal memory training group and path error specific to the navigation
training group. The other dependent measures, however, did not show any clear
evidence of a training-specific benefit (see Supplemental Note 3). We derived an
exploratory measure that allowed us to compare between navigation and verbal
memory, termed "*learning rate*." The learning rate attempts to capture how quickly a
participant might learn new information at post-test given that they were trained in
improving their acquisition of this information. For example, participants who underwent
navigation training should show a smaller difference in path distance on the first ten
trials compared to the last ten at post-test compared to pre-test, reflective of faster
learning due to the training (see Methods).
To quantify the learning rate for the Navigation Transfer task, a mixed-design
ANOVA was conducted with condition (Navigation/Verbal Memory/Video Control) as a
between-subjects factor and session (pre/post) as a within-subjects factor. This analysis
revealed a significant main effect of session ($F(1,70) = 37.52$, $p < 0.001$, $\eta^2 = 0.202$) and
a marginally significant main effect of condition ($F(2, 70) = 2.66$, $p = 0.08$, $\eta^2 = 0.028$),
with no significant interaction between condition and session ($F(2, 70) = 1.82$, $p = 0.17$,
$\eta^2 = 0.020$). Paired-sample t-tests indicated that all three groups demonstrated an
increase in learning rate from pre to post, with the Navigation group showing the largest
effect (Fig. 2a; Navigation: $t(26) = 4.43$, $p < 0.001$, Cohen's d = 1.32; Verbal Memory:
$t(26) = 2.31$, $p = 0.03$, Cohen's d = 0.596; Video Control: $t(18) = 3.97$, $p < 0.001$,
Cohen's d = 1.20). Results held when controlling for sex and site (see Supplementary
Note 3). The larger effects size for navigation suggests a greater benefit to the learning
rate for the navigation condition, although this conclusion is tempered by the lack of a
significant interaction effect.
To quantify the learning rate for the Verbal Memory Transfer task, we employed
a mixed-design ANOVA, with condition (Navigation/Verbal Memory/Video Control) as a
between-subjects factor and session (pre/post) as a within-subjects factor. We found a
main effect of session ($F(1, 71) = 3.21$, $p = 0.078$, $\eta^2 = 0.014$) and a significant
interaction between condition and session ($F(2, 71) = 2.9$, $p = 0.025$, $\eta^2 = 0.034$).
Paired-sample t-tests indicated that only the Verbal Memory group showed a significant
increase in learning rate from pre-test to post-test (Fig. 2b; Verbal Memory: $t(25) = 3.32$,
$p = 0.003$, Cohen's d = 0.608; Navigation: $t(26) = 0.488$, $p = 0.63$, Cohen's d = 0.113;
Video Control: $t(20) = -0.55$, $p = 0.588$, Cohen's d = 0.164). This effect remained
significant after controlling for sex and site (see Supplemental Note 3). These findings
suggest that verbal memory training resulted in significant improvements from pre-test
to post-test in the Verbal Memory Transfer task, specific to the Verbal Memory group.
266
267

268
Fig. 2 Changes in learning rates across training conditions from pre-test to post-test. (a)
Learning rate for spatial navigation improvement on the Navigation Transfer task. All three
groups improved from pre-test to post-test, with the Navigation group demonstrating the largest
effect. (b) Learning rate for verbal memory improvement in the Verbal Memory transfer task.
Only the Verbal Memory group significantly improved from pre-test to post-test. Red diamonds
represent the mean value. $*p < 0.05$, $**p < 0.01$, $***p < 0.001$.
275

Brain Structural Results

Medial temporal lobe volume did not change due to the training

As our task provided a separation of navigation and verbal memory skills, with
some evidence of their independence in the transfer tests, we next tested the
hypothesis that navigation training or verbal memory training (or both) would result in
increases in hippocampal volume (as measured with T2 high-resolution imaging, see
Methods). We conducted a mixed-design ANOVA with condition (Navigation/Verbal
Memory/Video Control) as a between-subjects factor and session (pre/post) as a within-
subjects factor. The analysis revealed no significant main effect of session ($F(1, 71) =$
0.199, p = .657, $\eta^2 = 0.0002$, $BF_{10} = 0.151$, moderate evidence against the inclusion of
the main effect of session), no significant main effect of condition ($F(2, 71) = 0.113$, p =
.893, $\eta^2 = 0.003$, $BF_{10} = 0.241$, moderate evidence against the inclusion of the main
effect of condition), and no significant interaction between session and condition ($F(2,$
71) = 1.224, p = .300, $\eta^2 = 0.003$, $BF_{10} = 0.054$, strong evidence against the inclusion of
the interaction effect). Paired t-tests further confirmed that none of the three training
groups exhibited significant changes in total hippocampal volume from pre to post-test
(Fig. 3b, ps > .265). This nonsignificant result persisted even after controlling for sex
and site as covariates (ps > .256).
We further investigated changes in hippocampal volume by conducting a mixed-
design ANOVA, with a particular focus on lateralization. Specifically, we included
regions of interest (ROI: left/right hippocampus) and session (pre/post) as a within-
subjects factors, alongside condition (Navigation/Verbal Memory/Video Control) as a
between-subjects factor. Consistent with the previous analysis, no significant main
effects or interactions were observed (session: $F(1, 71) = 0.199$, p = .657, $\eta^2 = .0002$,
$BF_{10} = 0.131$, moderate evidence against the inclusion of the main effect of session;
condition: $F(2, 71) = 0.113$, p = .893, $\eta^2 = 0.003$, $BF_{10} = 0.189$, moderate evidence
against the inclusion of the main effect of condition; ROI: $F(1, 71) = 0.208$, p = .650, $\eta^2 =$
0.0001, $BF_{10} = 0.115$, moderate evidence against the inclusion of the main effect of
ROI; and session × condition × ROI, $F(2, 71) = 0.072$, p = .930, $\eta^2 < 0.001$, $BF_{10} =$

0.0002, strong evidence against the inclusion of the interaction effect). Paired-sample t-
tests confirmed that none of the three training groups exhibited significant changes in
hippocampal volume from pre-test to post-test in either the left hippocampus (Fig. 3b, ps
> .316) or the right hippocampus (ps > .265). These findings remained consistent after
controlling for sex and site as covariates (left hippocampus: ps > .306; right
hippocampus: ps > .256).
Because some studies have shown long-axis specialization in correlations
between hippocampal volume and behavior[3,30,31], we considered whether anterior and
posterior hippocampus might change differentially due to the training, which we
analyzed based on the T1 whole brain structural image (see Methods). The previously
described mixed-design ANOVA model was expanded to incorporate ROI (anterior and
posterior hippocampus) as a within-subjects factor, thereby allowing for the examination
of volume differences along the hippocampal anterior-posterior axis. Consistent with the
above result, although we found a significant main effect of ROI ($F(1,71) = 22.311$, $p <$
0.001, $\eta^2 = 0.052$, $BF_{10} > 100$, strong evidence for the inclusion of the main effect of
ROI), we found no significant main effect of session ($F(1, 71) = 2.914$, $p = 0.092$, $\eta^2 <$
0.001, $BF_{10} = 0.439$, weak evidence against the inclusion of the main effect of session),
no significant main effect of condition ($F(2, 71) = 1.045$, $p = 0.357$, $\eta^2 < 0.022$, $BF_{10} =$
0.277, moderate evidence against the inclusion of the main effect of session), and no
significant interaction effect between session × condition × ROI ($F(2, 71) = 0.032$, $p =$
$.968$, $\eta^2 < 0.001$, $BF_{10} = 0.005$, strong evidence against the inclusion of the main effect
of condition). Simple main effects confirmed that none of the three training groups
exhibited significant changes in hippocampal volume from pre-test to post-test neither in
the anterior hippocampus (ps > .272) nor the posterior hippocampus (ps > .115). These
findings remained consistent after controlling for sex and site as covariates (anterior
hippocampus: ps > .260; posterior hippocampus: ps > .106).
Next, we divided the medial temporal lobe into six subregions: three hippocampal
subfields (CA1, CA23DG, and SUB) and three adjacent subregions (PRC, ERC, and
PHC). We then conducted a mixed-design ANOVA to examine changes in these
subfields and subregions. Although results revealed a significant main effect of ROI
($F(1.766, 125.390) = 602.143$, $p < 0.001$, $\eta^2 = 0.895$, Greenhouse-Geisser corrected;
$BF_{10} > 100$, strong evidence for the inclusion of the main effect of ROI), we did not find
any significant main effect of session ($F(1,71) = 0.766$, $p = 0.384$, $\eta^2 < 0.001$, $BF_{10} =$
0.047, strong evidence against the inclusion of the main effect of session), condition
($F(2,71) = 0.131$, $p = 0.878$, $\eta^2 < 0.001$, $BF_{10} = 0.002$, strong evidence against the
inclusion of the main effect of session) and no significant interaction effect between
session × condition × ROI ($F(3.21,110.806) = 0.888$, $p = .453$, $\eta^2 < 0.001$, Greenhouse-
Geisser corrected; $BF_{10} < 0.001$, extremely strong evidence against the inclusion of the
main effect of session). Paired-sample t-tests confirmed that none of the three training
groups exhibited significant changes from pre to post-test in any of the six MTL
subregions (Fig. 3a, ps > .588, FDR corrected). These findings remained nonsignificant
after controlling for sex and site as covariates (ps > .559, FDR corrected).
We also analyzed gray matter changes outside of the medial temporal lobe using
FreeSurfer (see Methods) to determine if any cortical or other relevant brain areas might
have been affected by the training. We applied a vertex-wise analysis, again finding no
significant differences across the entire cortex. This finding was further validated using

the Destrieux atlas[32], which included 34 cortical parcellations. Paired t-tests revealed
that none of the ROIs exhibited significant volume changes from pre- to post-test in any
of the three groups (ps > .542, FDR-corrected). These findings suggest that training did
not result in volumetric changes to the cortex.
355
Diffusion weighted imaging (DWI) metrics did not change due to the training
For DWI, we separately evaluated free water (FW) and free water-corrected
fractional anisotropy (fwcFA), as these measures capture different aspects of tissue
microstructure. We also chose to analyze ROIs in gray matter (GM) and white matter
(WM) separately, as these tissue types have dramatically different values for both FW
and fwcFA, which involved eight GM ROIs and seven WM ROIs (see Methods). We
conducted a mixed-design ANOVA with condition (Navigation/Verbal Memory/Video
Control) as a between-subjects factor, with ROI and session (pre/post) as within-
subjects factors. For the WM-fwcFA analysis, we observed a significant main effect of
ROI ($F(4.093, 233.279) = 2340.878$, $p < .001$, Greenhouse-Geisser corrected, $\eta^2 =$
0.628, $BF_{10} > 100$, extremely strong evidence for the inclusion of the main effect of
ROI). Yet, we found no main effect of session ($F(1,57) < 0.001$, $p = .981$, $\eta^2 < 0.001$,
$BF_{10} = 0.090$, strong evidence against the inclusion of the main effect of session), no
significant main effect of condition ($F(2,57) = 0.025$, $p = .975$, $\eta^2 < 0.001$, $BF_{10} = 0.011$,
strong evidence against including the main effect of condition), and no significant
interaction between session × condition × ROI ($F(6.924, 197.328) = 1.954$, $p = .064$, η^2
< 0.001, $BF_{10} < 0.001$, strong evidence against including this interaction). Paired-sample
t-tests confirmed that none of the three training groups exhibited significant changes
from pre- to post-test in any of the seven ROIs (ps > .083, FDR corrected). These
findings remained nonsignificant after controlling for sex and site as covariates (ps >
0.063, FDR corrected). Analyses involving WM-FW, and GM regions (GM-fwcFA and
GM-FW) were also not significant (see Supplementary note 4). These findings suggest
that no DWI measures changed as a result of the training. Together, these findings
suggest that neither navigation nor verbal memory training for 20 hours across 10
sessions was sufficient to induce changes in macroscopic hippocampal volume or
microstructure of relevant gray matter regions and white matter tracts.
382
Improvements in the learning rate of the Verbal Memory Transfer task, but not the
Navigation Transfer task, correlated with both baseline total hippocampal volume
and the baseline volume of the CA23DG subfield
We then performed an exploratory analysis to determine whether individual
differences in hippocampal volume independent of the training might predict how well
they learned from pre-test to post-test based on the different trainings. We conducted
analyses employing more stringent corrections for multiple comparisons to examine the
association between hippocampal volume and the change in learning rate from pre- to
post-test based on the Verbal Memory and Navigation Transfer tasks. This analysis
addresses the idea of whether those with larger hippocampi might improve more from
pre- to post-test based on the training. Because there were no pre-post differences in
hippocampal volume, we averaged the pre- and post-test hippocampal volume. We
consider other behavioral correlations with baseline hippocampal volume in the
Supplemental material; none of these were significant (see Supplementary Note 5).

For the Navigation Transfer task, the learning rate did not correlate with
hippocampal volume, hemispheric volume, anterior and posterior hippocampus, or any
hippocampal subfields, which was also not significant when controlling for sex and site
in all three groups (Supplementary Note 5 and Supplementary table 5). For the Verbal
Memory Transfer task, the changes in learning rate did not correlate with hippocampal
volume ($r(23) = 0.367$, $p = .071$). When accounting for sex and site as covariates,
however, we found a significant positive correlation ($r(23) = 0.605$, $p = .006$, FDR
corrected). This effect was specific to the Verbal Memory group; no significant
correlation was identified for either the Navigation group ($r(25) = 0.028$, $p = .889$) or the
Video Control group ($r(19) = -0.206$, $p = .371$); these findings remained consistent when
controlling for sex and site as covariates (Navigation: $r(25) = 0.038$, $p = .858$; Video
Control: $r(19) = -0.123$, $p = .858$, FDR corrected).
We further examined the correlation between learning rate and hippocampal
subfield volumes after correcting for multiple comparisons (see Methods), specifically
focusing on CA1 and CA23DG. Only the CA23DG subfield showed a positive
correlation with the change in learning rate from pre- to post-test in the Verbal Memory
Group ($r(23) = 0.532$, $p = .036$, FDR corrected, Fig. 3c), and this correlation persisted
after controlling for sex and site as covariates ($r(23) = 0.493$, $p = .017$). No significant
correlations were observed for the CA23DG subfield in the Navigation group ($r(25) = -$
0.083, $p = .953$, FDR corrected) or the Video Control group ($r(19) = -0.12$, $p = .953$,
FDR corrected), regardless of whether sex and site were included as covariates. For
correlations by hemisphere, anterior/posterior hippocampus and other subfields, please
see Supplementary Note 5 and Supplementary Table 4. These findings suggested that
individuals in the Verbal Memory group with larger CA23DG volumes exhibited greater
improvement in verbal memory performance from pre- to post-test.
Fisher's z-tests revealed that the positive correlation in the Verbal Memory group
was significantly stronger than that in the Navigation and Video Control group even after
controlling for sex and site (Navigation: total hippocampus: $Z = 2.48$, $p = 0.018$;
CA23DG: $Z = 1.972$, $p = .036$; Video Control group: total hippocampus: $Z = 2.596$, $p = $
0.015; CA23DG: $Z = 2.091$, $p = .036$; FDR corrected, one-side); there was no significant
difference between the Navigation and Video Control group (total hippocampus: $Z = -$
0.518, $p = 0.698$; CA23DG: $Z = -0.264$, $p = .604$; FDR corrected, one-side). Together,
these findings suggest that baseline hippocampal volume, and CA23DG baseline
subfield volume, correlated with the change in learning rate on the Verbal Memory
Transfer task following verbal memory training, but not after navigation training or video
control. We highlight, however, that these analyses are exploratory and somewhat
underpowered for between-subject comparisons.

434
435
Fig. 3 Relationship Between Medial Temporal Lobe Volumes and Training-Induced
Learning Rate Changes Across Conditions. (a) Normalized MTL subregion volumes across
conditions (Verbal Memory/Navigation/Video Control) and sessions (pre-test/post-test). No
significant changes were detected in CA1, CA23DG, SUB, ERC, PRC, or PHC volumes
between pre-test and post-test. (b) Normalized hippocampal volumes (HIP), including left and
right hippocampus, across conditions and sessions. No significant changes in total, left, or right
hippocampal volume from pre-test to post-test. (c) Correlations between changes in learning
rate (post-test minus pre-test) and average CA23DG volume across groups. A significant
positive correlation was observed in the Verbal Memory group, suggesting that participants with
larger CA23DG volumes exhibited greater improvements in verbal memory performance. No
such correlation was observed in the Navigation group or Video Control group. (d) Correlations
between the changes in learning rate (post-test minus pre-test) and average total hippocampal
volume across groups. A significant positive correlation was observed only in the Verbal
Memory group.
450

Task-related fMRI Results
Task-related informational connectivity analyses suggested network changes
specific to the Navigation and Verbal Memory training groups
Although we did not observe changes in hippocampal volume as a result of
training, it is likely that other brain-related changes might be mediating the
improvements specific to the navigation and verbal memory training. We investigated
this issue by collecting fMRI data both pre- and post-training during a source memory
encoding (Fig. 4c) and retrieval task (Fig. 5a). During the encoding phase, participants
learned objects by placing them within either an autobiographical timeline (temporal) or
a familiar spatial layout (spatial). During the retrieval phase, they saw the objects again
and indicated which of the "sources" (spatial or temporal) they had placed the object in.
Because we trained the Verbal Memory group to use the temporal method of loci but
trained the Navigation group to learn spatial locations, we reasoned that training-
specific strategies might lead to context-specific changes (i.e., depending on whether
the object was encoded in time or space) that would be reflected in network changes
during the source memory tasks. We performed a univariate analysis across the whole

brain and an informational connectivity analysis using 242 pre-defined brain regions
involved in spatial processing and memory (see Methods, Fig. 4b). Informational
connectivity involved correlating multivariate patterns between trials within each ROI
and then correlating again between all 242 ROIs (Fig. 4a). Informational connectivity
could reflect changes in network configurations during source memory encoding and
retrieval based on skills gained from training.
473
Encoding phase: Decreases in informational connectivity specific to the
Navigation group and increases specific to the Verbal Memory group
We first considered both spatial and temporal source memory during the
encoding phase, which could reveal how training affected the encoding of novel object-
source pairings more broadly. For the univariate analyses, we found some changes
from pre-test to post-test, but no interactions between session and training condition
(see Supplementary Note 7, Supplementary Table 16 and Supplementary Fig. 6). When
considering spatial or temporal encoding separately, we also found changes from pre-
test to post-test, but no interaction between session and training group (see
Supplementary Note 7, Supplementary Table 16 and Supplementary Figure 6). All
statistical analyses were corrected for multiple comparisons (see Methods).
Informational connectivity changes between pre-test and post-test were then
compared across the three training groups. For the Verbal Memory group, we observed
significantly increased informational connectivity at post-test compared to pre-test (i.e.,
post > pre), relative to the combined Navigation and Video Control groups (Fig.4).
Specifically, we found enhanced connectivity between the left precuneus and left
angular gyrus (t = 3.94, p < 0.05), the right lateral occipital cortex (LOC) and left
intracalcarine (visual) cortex (t = 3.63, p < 0.05), and the right LOC and left superior
parietal lobule (SPL) (t = 4.16, p < 0.05). In contrast, no significant decreases in
informational connectivity (i.e., post < pre) were observed. Considering spatial and
temporal encoding separately, for the Verbal Memory group during spatial encoding, we
found enhanced informational connectivity from pre-test to post-test (i.e., post > pre)
between the frontal cortex and visual cortex (see Supplementary Note 6 and Fig. 4d).
For temporal encoding, increased connectivity from pre- to post-test (i.e., post > pre)
was observed between the frontal cortex and temporal gyrus (Fig. 4d). No significant
decreases in informational connectivity from pre-test to post-test (i.e., post < pre) were
observed in the Verbal Memory group relative to the combined Navigation and Video
Control groups during either spatial or temporal encoding (Fig. 4d).
Conversely, for the Navigation group compared to the combined Verbal Memory
and Video Control groups, we observed significantly decreased informational
connectivity at post-test compared to pre-test (i.e., post < pre). Specifically, reduced
connectivity was observed between the left paracingulate gyrus and right middle frontal
gyrus (MFG) (t = 3.80, p < 0.05), left superior frontal gyrus (SFG) and right
supramarginal gyrus (SMG) (t = 5.14, p < 0.05), left frontal pole and right MFG (t = 3.85,
p < 0.05), left dorsal LOC and right ventral LOC (t = 3.7, p < 0.05) and right MFG and
left MFG (t = 3.79, p < 0.05). In contrast, no significant increases in connectivity were
observed from pre- to post-test (i.e., post > pre) for the Navigation group relative to the
combined Verbal Memory and Video Control groups (Fig. 4d, bottom-left). When
considering spatial and temporal encoding separately, during spatial encoding, reduced

connectivity was observed from pre-test to post-test (i.e., post < pre) between the
primary visual cortex and higher-order visual areas, while increased connectivity was
identified from pre-test to post-test (i.e., post > pre) between the parietal and temporal
cortices. For temporal encoding, the Navigation group exhibited significant decreases in
connectivity from pre-test to post-test (i.e., post < pre) across multiple regions, including
the frontal and parietal cortices (see Supplementary Note 6), with no significant
increases in connectivity from pre-test to post-test (i.e., post > pre). No significant
changes in task-related informational connectivity to the hippocampus survived after
multiple comparisons correction for either navigation or verbal memory training. Broadly,
these findings suggest changes in informational connectivity specific to each training
group (i.e., decreases in informational connectivity from pre-test to post-test in the
Navigation group and increases in the Verbal Memory group), with no changes specific
to the hippocampus.

**Fig. 4 Task-related informational connectivity changes during encoding as a result of
verbal memory and navigation interventions**. (a) Representational similarity matrices (RSMs)
that illustrate within-context correlations (spatial and temporal) for each of the 242 brain ROIs.
Informational connectivity between 242 ROIs was derived by correlating the RSMs across
regions within different contexts, resulting in three types of informational connectivity matrices
(ICMs): the spatial ICM, the temporal ICM, and the combined ICM for all within-context trials.(b)
Visualization of the 242 predefined ROIs, color-coded by functional networks. (c) Schematic of
the experimental design for the source memory task during the encoding stage, highlighting
tasks related to spatial and temporal contexts. (d) Differences in task-related informational
connectivity during encoding (Verbal Memory > Navigation + Video; Navigation > Verbal
Memory + Video) across training conditions. Results are shown for all trials, as well as
separately for spatial and temporal encoding contexts. Red lines indicate regions with
significantly increased connectivity (post > pre), while blue lines indicate regions with

542 significantly decreased connectivity (post < pre). The top panels display results for the Verbal
543 Memory group, while the bottom panels display results for the Navigation group. Nav:
544 Navigation
545
Retrieval phase: Increases in information connectivity specific to the Navigation
group and decreases specific to the Verbal Memory group
For the univariate analyses for both spatial and temporal source memory
retrieval, no significant changes were observed from pre-test to post-test, nor were
there any interactions between session and training condition (see Supplementary
Table 16). When examining spatial and temporal retrieval separately, we similarly found
no significant changes from pre-test to post-test and no interactions between session
and training group (see Supplementary Table 16).
We next examined informational connectivity changes from pre-test to post-test
during the source memory retrieval stage (Fig. 5a), finding a pattern somewhat opposite
to that seen during encoding. In the Verbal Memory group, we observed significantly
decreased informational connectivity in post-test compared to pre-test (i.e., post < pre)
relative to the combined Navigation and Video Control groups. These reductions were
identified between the right inferior frontal gyrus (IFG) and right frontal pole (t = 4.01, p
< 0.05), left frontal pole and right SFG (t = 4.48, p < 0.05), right medial frontal gyrus and
right posterior cingulate cortex (PCC) (t = 4.42, p < 0.05), right ventral LOC and left
temporo-occipital fusiform cortex (TOFC) (t = 4.30, p < 0.05), and right lingual gyrus and
left SPL (t = 3.82, p < 0.05). No significant increases in connectivity from pre-test to
post-test (i.e., post > pre) were observed in the Verbal Memory group relative to the
combined Navigation and Video Control groups (Fig. 5b, top-left). Considering spatial
and temporal source retrieval separately, spatial retrieval showed decreased
connectivity from pre- to post-test (i.e., post < pre) between frontal regions (e.g., left and
right superior frontal gyri) and between the right frontal pole and PCC, and increased
connectivity from pre to post (i.e., post > pre) between temporal and parietal regions
(e.g., left middle temporal gyrus [LMTG] and right SMG) and between the visual cortex
and right SMG. For temporal retrieval, decreased connectivity from pre-test to post-test
(i.e., post < pre) was prominent across multiple regions, including the frontal pole,
angular gyrus, precuneus, and lateral occipital cortices. One increase specific to the
Verbal Memory group was observed from pre-test to post-test (i.e., post > pre),
however, in the left ventral lateral occipital cortex and left MFG (Fig. 5b, see details in
Supplementary Note 6).
In the Navigation group during retrieval, we found significantly increased
informational connectivity in the post-test compared to the pre-test (i.e., post > pre)
relative to the combined Verbal Memory and Video Control groups. These increases
were found between the right occipital pole and dorsal LOC (t = 4.27, p < 0.05), and the
right SPL and right angular gyrus (AG) (t = 4.09, p < 0.05). Considering spatial and
temporal retrieval separately, spatial retrieval showed enhanced connectivity from pre-
test to post-test (i.e., post > pre) between regions such as the left frontal pole and the
occipital and parahippocampal cortices. Temporal retrieval revealed increases from pre-
to post-test (i.e., post > pre) between the precuneus and SFG, lingual gyrus and
occipital gyrus, and SFG and MTG. No significant decreases in connectivity were
observed from pre-test to post-test (i.e., post < pre) in the Navigation group relative to
the combined Verbal Memory and Video Control groups (Fig. 5b, bottom).

Again, no significant changes in task-related informational connectivity during
source retrieval were present to the hippocampus when controlling for multiple
comparisons. To summarize, verbal memory training resulted in decreased connectivity
during retrieval across frontal, parietal, and occipital regions, especially during temporal
retrieval, with limited increases. In contrast, navigation training consistently increased
connectivity for both spatial and temporal retrieval, particularly between frontal, occipital,
parietal, and temporal regions, with no significant decreases observed. These findings
suggest that verbal memory primarily alters network connectivity during temporal
contextual retrieval, whereas navigation training enhances connectivity more broadly
across both contexts.
599

600
601
Fig. 5 Task related informational connectivity changes during source retrieval as a result
of verbal memory and navigation interventions. (a) Schematic of the source memory task
structure during retrieval. Participants were scanned during source retrieval, in which they
identified whether an item was previously seen and determined its spatial or temporal context.
(b) Differences in task-related informational connectivity during source retrieval (Verbal Memory
> Navigation + Video; Navigation > Verbal Memory + Video) across training conditions. Results
are shown for all trials, as well as separately for spatial and temporal encoding contexts. Red
lines indicate regions with significantly increased connectivity (post > pre), while blue lines
indicate regions with significantly decreased connectivity (post < pre). The top panels display
results for the Verbal Memory group, while the bottom panels display results for the Navigation
group. Nav: Navigation

613
Specific changes in network-wide informational connectivity pattern distance
during retrieval: Enhanced due to verbal memory but not navigation training
The above analysis focused on changes in informational connectivity between
pre-test and post-test for individual connectivity edges between pairs of ROIs. Next, we
examined changes in multivariate informational connectivity patterns, which we defined
as the connectivity between a single ROI, and the remaining 241 ROIs. This analysis
helped address network-wide changes in connectivity rather than those restricted to two
different ROI pairs. Multivariate informational connectivity patterns were extracted
separately for spatial and temporal informational connectivity matrices (ICMs), and the
distance between the two patterns was computed using 1 - r (Pearson's, see Methods).
While no significant changes in pattern distance were observed during the
encoding stage for any of the three conditions, a greater distance between spatial and
temporal patterns in the left SFG network connectivity during source retrieval was found
in the Verbal Memory group at post-test compared to pre-test (t(23) = -4.275, p = 0.043,
FDR corrected; Fig. 6b). This effect was not observed in the Navigation and Video
Control groups (Fig. 6c). A similar result was obtained using the Spearman correlation
to compute the distance, with a greater distance in the left SFG for the Verbal Memory
group at post-test relative to pre-test (t(23) = -4.400, p = 0.050, FDR corrected). No
significant differences were found in the Navigation or Video Control groups. These
findings also suggest some network-wide changes that are specific to verbal memory
training.
635
636

637
Fig. 6 Multivariate informational connectivity pattern distance between pre-test and post-
test. (a) Schematic of multivariate distance analysis. For each ROI, informational connectivity
between the current ROI and the remaining 214 ROIs was extracted from both spatial and
temporal ICMs, and distances were calculated as 1 minus the Pearson or Spearman correlation
coefficient. (b) and (c) Among all 242 ROIs, only the SFG showed a greater distance between
spatial and temporal ICMs in post-test compared to pre-test in the Verbal Memory group, but not
in the Navigation or Video Control groups. ICM: informational connectivity matrix. *p < 0.05,
FDR corrected.
646

Discussion
Acquiring complex cognitive skills is a fundamental human capability, but we
have little knowledge about the neural changes that accompany such new learning. As
suggested in past work, we are capable of remarkable improvements in performance
across a wide range of domains, yet the neural basis for these changes remains

unclear. Here, we build on expert training regimens in verbal memory (e.g., the method
of loci) and spatial navigation (e.g., London taxi driver training) by employing a within-
subject intervention approach to determine the neural correlates of training two related
but distinct complex skills: navigation and verbal memory. Our results show that non-
expert young adults substantially improved in the task they were trained on and
exhibited some limited near transfer (to new lists of words or new environments). Unlike
the neuroimaging literature on experts or impaired populations (like older adults or
individuals with dementia who cannot perform these tasks as well), we found no
evidence that training modified brain structure, as measured by hippocampal volume
and DWI microstructure in relevant GM regions and WM tracts. Rather, analyses of
task-based functional imaging data showed training-specific differences in informational
connectivity, a neural metric that indexes how similar representations are across brain
regions[33]. Our findings have implications for both basic research and clinical translation.
From a basic research perspective, our results suggest that structural changes in gray
and white matter are relatively insensitive to cognitive intervention but are instead
reflected in more transient changes to network-based connectivity patterns. From a
clinical-translational perspective, our results suggest promise in training new complex
cognitive skills such as navigation and verbal memory, although the efficacy of our
specific training regimen in clinical populations remains to be tested.
Our study involved two novel and extensive in-lab training regimens, which
shared key features but allow for dissociable cognitive training. While previous training
studies have examined network configurations before and after participants learned the
method of loci or evaluated navigation training, these experiments often involved expert
groups (memory athletes or London taxi drivers)[3,34], or older adults[9]. In our experiment,
we evaluated healthy young adults while carefully controlling the duration and nature of
the verbal memory and navigation trainings to match the intensity and the degree to
which the skill was scaffolded. In this way, one training serves as a control for the other,
while also allowing us to compare with an active control group who did not gain a
complex skill (video control) but was passively exposed to similar material.
Both training regimens were effective: participants in the Verbal Memory group
were able to learn longer word lists, while participants in the Navigation group were able
to reduce the length of their paths and required fewer trials to learn landmarks in new
subsections of the environment. Yet, it was difficult to ascertain whether either training
generalized beyond immediate transfer. Our near transfer measures exhibited strong
individual differences at baseline, which were difficult to overcome with 10 sessions of
training. Both near transfer measures also likely had practice and ceiling effects, which
dampened our ability to measure the effects of training (e.g., Supplementary Figure 2).
Exploratory analyses, however, revealed training-specific differences in the learning rate
on the transfer tasks from pre- to post-test. In addition to good face validity as a match
for the skills we were training (that is, for example, in the navigation training, participants
learned how to quickly acquire spatial information about the whole environment after
only 10 trials compared to those who did not undergo that training), the change in
learning rate from pre- to post-test was also the only dependent measure we found to
correlate with baseline hippocampal volume. These findings, though exploratory,
suggest that baseline hippocampal volume may enable more receptiveness to training,
aligning with concepts like neural reserve[35]. In addition to replicating these findings,

future research using these training regimens should develop a more comprehensive
near-transfer measure and especially evaluate far transfer to completely untrained
tasks.
Our navigation training is particularly notable in that it partially mimics the training
undertaken by London taxi drivers, who first learn individual subsections of an
environment before connecting these subsections to one another. While other virtual
navigation trainings exist[36,37], their environments are often small, simplified, or non-
hierarchical. Thus, a key contribution of this work is the development of a navigation
training environment that is suitable for training wayfinding skills. We also developed a
novel verbal memory training paradigm that has not been reported previously in the
literature. Using this training, we were able to dramatically increase the number of freely
recalled words of the average participant, with the typical participant showing an
improvement from about 25-30 words freely recalled (repeated) words on day 1 to 60-
65 words after only five days of training.
Despite the efficacy of the training, neither the Navigation nor the Verbal Memory
group showed macroscopic neural structural differences in the hippocampus, which has
been linked to verbal memory and/or navigation function in past studies[3,30,38]. These
results supply causal evidence supporting the notion that hippocampal structural
changes may not impart superior navigation or verbal memory skills. Although previous
work has shown effects of training, to our knowledge, this work has occurred in
specialized populations or involved between-subject comparisons. For example, London
taxi drivers who undergo extensive training to learn the city of London show increased
hippocampal volume from pre- to post-training compared to taxi drivers who fail to
complete the training[3]. Older adults who undergo an exercise-only training or a
navigation + exercise training show reduced hippocampal atrophy (i.e., less volume
loss) compared to a no-contact control[39]. Here, in healthy younger adults, who
nevertheless showed robust individual differences in navigation and verbal memory and
showed strong evidence of improvement across both types of training, there were no
observed structural changes to hippocampal gray or white matter tracts, nor were there
any significant structural changes to other brain regions.
In exploratory analyses, informational connectivity analyses on fMRI data from a
spatial and temporal source memory task revealed unique network reconfiguration
signatures as a result of navigation and verbal memory training. These changes were
based on a strict correction for multiple comparisons and suggest that verbal memory
and navigation training results in robust changes in task-related functional connectivity
patterns. Interestingly, the vast majority of these changes were to areas outside of the
medial temporal lobe, an area often considered central to both navigation and memory.
During encoding, we observed increases in connectivity due to the verbal memory
training primarily in the frontal and occipital lobes. Conversely, we observed decreases
in informational connectivity in some of these same areas as a result of navigation
training. We observed somewhat the opposite pattern during retrieval: increases in
informational connectivity during navigation and decreases during verbal memory
training. Again, many of these connectivity changes were areas within the frontal cortex,
with some longer-range changes to parietal, parahippocampal, and occipital cortex.
Finally, a multivariate approach centered around superior frontal gyrus as a hub
revealed differences in network connectivity patterns for spatial compared to temporal

retrieval, suggesting task-related network-level changes due to the training as well.
Similarly, we note that two studies, which trained participants on an n-back task, also
observed changes in dynamic functional connectivity in frontal and parietal cortices[40,41].
Thus, our results suggest that the networks most malleable during acquisition of
complex cognitive skills largely involve areas related to cognitive control, with one
possibility being that such changes allow modulation of existing networks that include
the medial temporal lobe[42]. At the same time, navigation and verbal memory training
resulted in differential changes within these frontal-parietal networks, suggesting that
such cognitive skills involve at least partially dissociable changes. While it is reasonable
to conjecture that microstructural changes in brain tissue (as measured with DWI) or
even macrostructural gray matter changes may occur after considerable expertise is
developed, such structural reorganization would likely be preceded by network changes
in functional connectivity. In this scenario, functional changes that reflect training effects
could potentially become solidified through long-term maintenance or the development
of domain-specific expertise.
759

Methods

Participants

83 participants between the ages of 18-45 were recruited for enrollment in the study
across two sites (University of Florida and University of Arizona). Participants were
recruited through advertisements and flyers posted on campus. After recruitment,
participants were screened for: 1) being right-handed; 2) having no metal in their bodies
or other MRI contraindications; 3) having no COVID-19 symptoms. Participants also
completed and passed an MRI prescreen before fMRI scans. This study included 75
participants (Supplementary Table 1) who fully completed the entire study: 27 in the
verbal memory condition, 27 in the navigation condition, and 21 in the video control
condition. One verbal memory participant, despite completing training, was excluded
from MRI scanning due to ear piercings, resulting in 26 participants for brain structure
and connectivity analyses. Some participants were excluded from certain tests due to
outlier performance, excessive head movement during scanning, or missing data. For
detailed sample size information for each test, please refer to Supplementary Table 2.
775

Overview

Participants were recruited via flyers or word of mouth. During prescreening, each
participant was randomly assigned to one of three training conditions (Navigation,
Verbal Memory, or Video Control). Participants then completed a prescreening phone
call for MRI eligibility, COVID safety, and availability to determine participation in the
study. They then received instructions for their first session verbally and via email. The
study took place over 12 sessions, each of which occurred on a separate day. These 12
sessions took place over the course of three to four weeks (average duration = 4.116
weeks). The first and last sessions were 3-hour sessions, referred to as pre-test and
post-test, and consisted of behavioral and neuroimaging measures. These sessions
were identical for all participants, regardless of which condition they were assigned to.
Following the pre-test session, participants completed ten 2-hour training sessions,
during which they completed training based on their assigned condition. Following
training, participants completed the post-test session, were debriefed and released.

Participants were paid for their participation. See Supplementary Table 3 for more
details about conditions and scheduling.
792
Training Conditions*: Navigation Training*
Participants assigned to the Navigation group navigated a virtual environment in
Unity 3D called "Virtual Arida" using an Xbox game controller on a desktop computer.
Virtual Arida has six individual subsections (A, B, C, D, E, F), with each containing eight
different target locations. Participants began training on day 1 in subsection A and were
spawned in front of a target location. They were then asked to navigate using the
shortest possible route to a second target location. If the participants did not perform a
sufficiently direct route, they repeated the route; their route was determined to be
sufficiently direct if it did not exceed 120% of the shortest possible path between two
buildings. Once they successfully employed a sufficiently direct route, they moved on to
the next route. Participants learned all 28 possible routes in the subsection. When they
learned all routes successfully, they completed random target location pairings for the
remainder of the 2-hour session. After completing a subsection, participants moved on
to a new subsection. In some cases, the new "subsection" involved navigating between
two previously learned subsections. For example, once individual subsections A and B
were completed, participants navigated between targets in subsection A and subsection
B. This involved "integrating" their previously learned knowledge from prior subsections
(A+B). Participants then learned the next individual subsection (subsection C), and after
that, the next integrated subsection A+B+C. This continued in this manner through the
final integrated subsection A+B+C+D+E+F (for those participants that reached this far).
813
Training Conditions*: Verbal Memory Training*
Participants placed in the verbal memory condition completed free recall training
on a desktop computer using a task built with Unity 3D. Similar to the verbal memory
transfer task during pre/post-test, in this task participants were shown a list of words
from the Toronto word pool one at a time, which they memorized using a trained
strategy (the "temporal" method of loci)[43]. After being shown the list of words, they
completed a distractor task by typing the numeric answer for five simple
addition/subtraction questions. Then, they recalled the words by typing them into the
provided space. The strategy that participants were given to memorize the words was a
modified method of loci termed the "temporal method of loci"[43]. At the beginning of Day
1 of training, participants wrote 30 of their most memorable, vivid, and non-traumatic
memories from their life thus far. They were asked to order these memories into a
mental timeline to help them with the word recall task. Participants were asked to
associate each word they were given with a memory from their mental timeline. Then,
when they reached the recall portion of the task, participants were to use their mental
timeline memories to help them recall the associated words in the current list. The
number of words increased with each successful trial. If there were multiple
unsuccessful trials, the number of words decreased. Once they were successful with
this shorter list, the previous list was repeated. For example, if the list was "red,"
"orange," "yellow," and the participant did not recall all 3 words, the next trial's list would
be "red," "orange." On successful completion, the next trial would be "red", "orange,"
"yellow," "blue."

Session 1-5 of verbal memory training (same words list)

For the first five sessions of training, participants recalled a list that increased in length on each trial when they successfully recalled all the words. This was done using an algorithm based on a power law learning function. For example, they could start with the words: "Jersey," "rifle," "pony." Upon successful completion of this list, the list would grow to "Jersey," "rifle," "pony," "castle." (Fig. 1a).

Session 6-10 *of verbal memory training (new word list)*

For the final 5 sessions of training, once participants successfully recalled a list, they learned a completely new list of words that was increased in length in the same manner as described above. For example, they could start with the words: "sister," "soldier," "beauty." Upon successful recall of this list, the next trial, the next list could be "mother," "genius," "number," "sailor." (Fig. 1b).

Active Control Condition- Video Watching

Participants in this condition watched a wide range of videos on a desktop computer. The types of videos varied, but included interviews, lectures, documentaries, shows, and YouTube videos covering different aspects of memory and navigation. Participants were given a set of questions for each video to ensure that they paid attention. Videos and quiz questions were administered using Qualtrics.

Pre-test overview

In the Pre-test session, participants provided informed consent, completed a demographics questionnaire, and confirmed their MRI prescreen information. Participants then completed pre-test behavioral measures, which included a navigation transfer task (Virtual Silcton), a verbal memory transfer task (a new set of words from the Toronto word pool), and an attention task (Posner cueing task). Next, the experimenter described the source memory task that would take place during the fMRI scan. The participants created the temporal and spatial contexts for this task, then after a short break to prepare for the scanner, they completed the source memory task in the scanner (see details in Supplementary Note 1 and Supplementary Table 3).

Post-test overview

The post-test session was similar to the pre-test session. The only changes in the post-test session were different words for the verbal memory transfer task, a different section of Virtual Silcton, and different images for the source memory task. After reviewing the MRI prescreening form for any changes, participants completed the post-test measures (navigation, verbal memory, and attention tasks). Participants then reviewed the spatial and temporal contexts created in the pre-test session, took a short break, and completed the source memory task in the scanner. Finally, participants completed a debriefing survey and received payment (see details in Supplementary Note 1 and Supplementary Table 2).

Pre-post tests
Navigation Transfer task: Virtual Silcton

The pre-test and post-test navigation transfer task was completed on a laptop
computer with an Xbox game controller using the "Virtual Silcton" environment
recreated in Unity 3D[29] (http://www.virtualsilcton.com/). Participants were instructed to
freely navigate from their starting location to one of the eight possible destination
buildings within the environment. If the participant walked for more than 60 seconds
during a trial, a compass appeared at the bottom of the screen to point them to the
target building. Over the trials, participants learned the spatial environment from the
routes taken between the buildings. Participants completed trials until all routes were
completed, or until 30 minutes elapsed. After the learning phase, participants completed
an on-site pointing task, in which they were placed next to one of the eight buildings and
had to point to each of the other seven in a random order. Next, they completed a
model-building task, in which they dragged and dropped images of the eight buildings
around an on-screen box. These tasks demonstrate configural knowledge of the
environment.
896
897 ### ***Verbal Memory Transfer task***
898   Participants memorized and recalled a list of 12 words from the Toronto Word
899 pool. The structure of the task was the same as that for the verbal memory training task
900 although it was programmed in PsychoPy. All participants had a maximum of five
901 attempts to recall all 12 of the words correctly.
902
Source Memory task
Prior to entering the scanner, participants were asked to generate a *temporal* and
a *spatial* context to use during the source memory task. For their temporal context, they
were asked to create a list of 10-15 salient memories of events from their life. We then
asked them to write these down on a piece of paper and arranged them into a mental
timeline of life events. For their spatial context, participants drew a blueprint of their
living space (e.g., dorm room or apartment) on a second piece of paper. Participants
were given time to review their contexts to ensure they knew them well enough prior to
entering the scanner. They were then instructed on how they would use these contexts
to recall objects. They were provided with an example of how to use each context and
practiced verbally by using these contexts verbally.
Participants underwent functional MRI scanning while performing the source
memory task. This task was comprised of two phases, each with four runs. During the
encoding phase (runs 1-4), participants were presented with images and instructed to
encode them within either a temporal or spatial context, consistent with prior out-of-
scanner practice. During the retrieval phase (runs 5-8), participants were presented with
images and asked to make two judgments: first, whether the image was new, and
second, if recognized, the context in which it had been previously encoded. All
participants viewed the same images, but in counterbalanced orders.
922
Source Memory Encoding
This source memory task included 4 encoding and 4 retrieval runs. In the
encoding task, participants viewed an image and were asked to imagine it in either
"time" or "space", referring to their previously generated temporal or spatial contexts.
Participants were to either visualize the image as a part of their mental timeline if it was

to be imagined in "time" or visualize this object existing in their living space if it was
indicated that it was to be imagined in "space". Following this visualization period,
participants were asked to indicate how vividly they visualized this item on a Likert scale
by pressing 1-4 on the button box (see image below). All trials were jittered. At the end
of the block, participants completed a short odd/even task. In this 60-second task, an
"X" or an "O" would be displayed. Participants pressed "1" if it was an "X" and "2" if it
was an "O". This served as a baseline task for hippocampal activation[44].
935
Source Memory Retrieval
In the retrieval task, participants were shown an image of an object and asked
whether they recognized it from the encoding phase or if it was a new image by
pressing "1" for recognition and "2" for new. They were then asked if this image was
previously viewed in "time", or "space", or if it was new by pressing "1", "2" or "3",
respectively. Four retrieval runs were completed. All trials were jittered. Detailed
descriptions of other pre-post tests are provided in Supplementary Note 1.
943

Behavioral Data Analysis
944
Navigation task (navigation training in Virtual Arida)
945
To quantify the training effect, we analyzed the average distance error travelled,
normalized by the threshold (120% of the length of the shortest possible path). We
removed all trials that occurred after all unique pairs had been learned. Because we
controlled the amount of time per task (2 hours per session), not all participants
progressed to the same extent. Repeated-measures ANOVAs were conducted with
subsection or integration as the within-subjects factors and threshold-normalized
distance error as the dependent measure.
953

Verbal Memory task (verbal memory training with the modified method of loci)
954
955   To quantify the training effect, we performed linear regression analysis on the
956 maximum number of correctly recalled words per day. Regression slopes were
957 calculated separately for the first five and last five days (Fig. 1a right), reflecting the
958 distinct training paradigms implemented during these two phases (see Methods).
959

Learning rate in the Navigation Transfer task (Virtual Silcton) during the pre-/post-
960
test
To assess navigation performance in the Navigation Transfer task, for each trial,
we calculated distance error by subtracting the optimal distance from the participant's
actual navigated distance between their starting location and the target building. Since
distance errors are inherently influenced by the distance between the starting location
and the target building, we normalized these errors by dividing each distance error by
the corresponding optimal distance. The task consisted of 20 trials, during which
participants progressively learned the environment. As expected, normalized distance
errors decreased over time, indicating improved navigation performance in later trials.
To quantify this learning effect, we calculated a learning rate, defined as the negative of
the difference between the normalized distance errors of the first 10 trials and the last
10 trials, divided by their sum (see Eq. 1). A higher/positive learning rate suggests
relatively stable and consistent performance across early and late trials whereas a

lower/negative learning rate reflects a greater difference in performance between early
and late trials.

$$Learning\ Rate = -(\frac{Distance\ errors\ of\ first\ 10\ trials - Distance\ errors\ of\ last\ 10\ trials}{Distance\ errors\ of\ first\ 10\ trials + Distance\ errors\ of\ last\ 10\ trials})\ \text{[Eq. 1]}$$

Learning rate in the Verbal Memory Transfer task during the pre-/post-test

For the Verbal Memory Transfer task, we calculated a learning rate to quantify
memory performance. To determine the learning rate, we first calculated the difference
in the number of words correctly recalled between the first and last trials. This difference
was then divided by the total number of trials to yield the learning rate (see Eq. 2).

$$Learning\ Rate = \frac{Last\ trial - First\ trial}{number\ of\ trials} \qquad\qquad \text{[Eq. 2]}$$

Correlation analysis

To examine the relationship between learning rate of either task and brain
structure (volume/DWI, see below), we collapsed the session factor by averaging values
from pre- and post-training sessions for each ROI, given the absence of significant
session effects or interactions between session and training condition. We conducted
Pearson correlation analyses (parametric) for data that met the normality assumption
based on the Shapiro-Wilk test. If normality was violated, Spearman correlation analysis
(nonparametric) was used instead. Sex and site were included as covariates in the
corresponding partial correlation analyses.

fMRI Data Analysis
MRI data acquisition

Scanning was performed using a 32-channel 3T Siemens "Skyra" scanner at the
University of Arizona and 32-channel 3T Siemens "Prisma" scanner at the University of
Florida. Visual stimuli were presented on a screen positioned behind the scanner and
viewed by participants through a mirror attached to the head coil. Stimuli and responses
were presented and collected using PsychoPy (https://www.psychopy.org) running on a
Windows 10 laptop. High-resolution anatomical images of the hippocampus and
surrounding cortex were acquired with a T2-weighted turbo-spin echo (TSE) anatomical
sequence (FOV = 200 mm × 200 mm, matrix = 448 × 448, TR = 4200.0 ms, TE = 93.0 ms,
flip angle = 139 degree, slice thickness = 1.9 mm, 28 slices, bandwidth = 199 Hz/pixel).
High-resolution structural images of the whole brain were obtained using a 3D, T1-
weighted MPRAGE (1 mm^3 isotropic) sequence (FOV = 256 mm, matrix = 256 × 256,
slice thickness = 1 mm, TR = 2300 ms, TE = 2.41 ms, flip angle = 8 degree,
bandwidth = 330 Hz/pixel). Functional images were acquired using a whole-brain echo
planar imaging (EPI) sequence (TR = 1560 ms, TE = 30 ms, flip angle = 70 degree, field
of view (FOV) = 220 mm, matrix = 88 × 88, slice thickness = 2.5 mm, slices = 48,
bandwidth = 2030 Hz/pixel), involving a voxel resolution of 2.5 × 2.5 × 2.5 mm. Diffusion
Weighted Imaging (DWI) data were acquired using two sequences with opposite phase-
encoding directions to correct for distortion artifacts without signal loss. Acquisition

parameters included: TR = 9200 ms, TE = 86 ms, FOV = 256 mm, 30 diffusion
directions, 60 slices with a thickness of 2.0 mm, and a voxel size of 2.0 × 2.0 × 2.0 mm.
The diffusion-weighted images were obtained using an echo plane sequence with the
following parameters: number of b0 images = 1, b-value = 1000 s/mm^2, number of
directions = 30, TR = 9200 ms, TE = 86 ms, and voxel size: 2 × 2 × 2 mm. High-
resolution resting state images were acquired using another whole-brain EPI
sequence(TR = 3000 ms, TE = 36 ms, flip angle = 90 degree, field of view
(FOV) = 240 mm, matrix = 160 × 160, slice thickness = 2.5 mm, slices = 48, bandwidth =
802 Hz/pixel), involving a voxel resolution of 1.5 × 1.5 × 2.5 mm. Resting state data
were not considered in this manuscript.
1029
DWI data preprocessing
We performed pre-processing of DWI data using a customized pipeline that
combines tools from the fMRIB Software Library 6.0[45], Advanced Normalization Tools[46],
and MRtrix3[47]. First, we removed Gaussian noise present in the DWI data by fitting a
Marchenko-Pastor distribution to the signal matrices to generate a threshold for PCA
denoising[48-50]. We then removed Gibbs-ringing artifacts that can occur at tissue borders
such as the outer surface of the brain and near ventricles[47]. Next, we generated brain
masks for the DWI images using the dwi2mask function from MRtrix3, which uses
information from both diffusion-weighted and non-diffusion weighted (b=0) volumes to
generate an accurate brain mask. We then corrected for Eddy current and movement
related distortions using FSL Eddy[51-53]. Using the root mean square motion output
provided by Eddy, we then applied a motion threshold, excluding participants with > 2
mm absolute displacement or > 0.5 mm relative displacement between diffusion
weighted volumes. Application of this motion threshold at both timepoints resulted in 20
participants in the Verbal Memory group, 22 participants in the Navigation group and 18
participants in the Video control group in final DWI analysis.
Consistent with prior work[54-58], we reconstructed the DWI data using a bi-tensor
model, in which freely diffusing water is modeled by one tensor and anisotropic water
diffusion is modeled with a separate tensor after removing the contribution from
isotropic free water. In this model, the free water (FW) compartment of each voxel is
interpreted primarily as originating from extracellular water diffusion and the second
tensor represents the tissue compartment after removing the contribution from FW. We
performed whole brain estimation of FW and calculated free water corrected tensor
metrics using custom MATLAB scripts (R2023a, The Mathworks, Natick, MA, USA).
Briefly, we calculated FW from single shell diffusion data based on minimization of a
variational regularization framework outlined in Pasternak et al. (2009). Initialization of
the free water estimate in this pipeline used mean diffusivity (MD) maps calculated from
a single tensor fit. Thus, prior to FW estimation we performed conventional single tensor
reconstruction using FSL's DTIFIT. Next, conventional MD maps were for initialization of
the bi-tensor reconstruction in MATLAB. Voxels with MD values greater than 0.8 x d
(*i.e., d* is constant diffusivity) were assumed to be comprised of CSF and omitted from
the fitting process. Fitting a bi-tensor model with DWI data from a single non-zero
diffusion weight (i.e. b-value) has been described as an ill-posed problem with nearly
infinite solutions, thus we employed sensible biological constraints to the minimization
process, as performed in Pasternak et al. (2009). We set the reference MD_t at 0.6

$\mu m^2/ms$. and diffusivities were limited to λ_{max} = 2.5 $\mu m^2/ms$ and λ_{min} = 0.1 $\mu m^2/ms$.
We also assumed isotropic water diffusion at 37° C, corresponding to human body
temperature, as a constant (d = 3.0 x 10^{-3} $\mu m^2/ms$). After initialization, 100 iterations
were used to refine the FW estimates corresponding free water corrected metrics. We
used an automated quality assurance procedure and visually inspected output images
(e.g., FW and fwcFA) to confirm appropriate data quality.
We used ANTs to perform nonlinear registration to MNI standard space by
warping each participant's FA image to the HCP 1065 template[59]. We used participant's
uncorrected FA for the registration process because the HCP 1065 template was
created with FA images from single tensor reconstruction, not fwcFA images. We then
applied the same transformation matrix to align FW and fwcFA images from the same
participant into MNI space. Following registration, we extracted FW and fwcFA values
from ROIs in MNI space. We separately evaluated ROIs from gray matter (GM) regions
and white matter (WM) tracts, selected based on prior literature and relevance to
navigation. Eight bilateral GM ROIs from the Mayo Clinic Adult Lifespan Template[60]
were used: hippocampus, parahippocampal gyrus, entorhinal cortex, inferior temporal
cortex, cuneus, superior parietal cortex, retrosplenial cortex, and caudate nucleus[61].
Seven bilateral WM ROIs from the Johns Hopkins University (JHU) white-matter
tractography atlas were used: three subregions of the corpus callosum (body, genu, and
splenium), fornix, posterior thalamic radiation, cingulum (hippocampus), and the fornix
cres / stria terminalis (these two small tracts are not able to be differentiated at 2mm
voxel resolution).
1087
DWI statistics
Similar to previous work[56], we performed separate statistical analyses of ROIs
from GM and WM regions, as these tissue types have dramatically different values for
DWI metrics. We also separately evaluated FW and fwcFA, as these metrics provide
separate but complementary information about extracellular and intracellular
contributions to tissue microstructure. For each of these four combinations (GM-FW,
GM-fwcFA, WM-FW, WM-fwcFA), we performed mix-designed ANOVAs with condition
(Navigation/Verbal Memory/Video Control) as a between-subjects factor and ROI and
session (pre/post) as within-subjects factors with corresponding post-hoc comparisons.
We also performed correlations (Pearson or Spearman, as appropriate) to evaluate the
relationship between DWI metrics and performance on behavioral measures.
Correlations were corrected for multiple comparisons using the false discovery rate
(FDR) method[62], which was applied across p-values for each ROI, performed separately
for FW and fwcFA analyses.
1102
MTL subfield demarcation
Automatic hippocampal subfield segmentation software (ASHS)[63,64] was used to
segment the subfields of the MTL based on each participant's high-resolution T2-
weighted MRI image. We used ASHS with the ASHS-Princeton-1.0.0-Young-Adult[64].
The MTL was segmented into CA1, CA2/3, DG, and subiculum (SUB), as well as the
perirhinal cortex (PRC), entorhinal cortex (ERC), and parahippocampal cortex (PHC).
We combined the CA2/3 and DG subfields as finer distinctions could not be made at the
acquired resolution[65]. Each participant's subfield segmentations were manually

inspected to ensure accuracy of the segmentation protocol. The hippocampus was
further subdivided into anterior (head) and posterior (body + tail) regions along its
longitudinal axis using the T1-weighted MPRAGE sequence and FreeSurfer 7.4.1
software. To ensure reliable volume estimates, images were processed using
FreeSurfer's longitudinal pipeline [66]. This method creates an unbiased within-subject
template using a robust, inverse-consistent registration approach[67]. To account for
individual differences in brain size, we normalized the volume of each MTL subregion by
dividing each volume by intracranial volume (ICV) and scaling by a factor of 100.
1119
Regions of Interest
To identify brain regions associated with memory and spatial navigation, we
conducted meta-analyses using Neurosynth (https://neurosynth.org/). We first defined
11 maps based on Neurosynth association tests, using the following 11 key terms:
"memory," "episodic memory," "autobiographical memory," "recognition memory,"
"subsequent memory," "memory encoding," "memory retrieval," "memory performance,"
"memory processes," "memory tasks," and "memory test." These maps were then
combined to create a single, comprehensive memory-related brain mask. Then,
following the same steps, we created a single, comprehensive spatial navigation-related
mask using six key terms: "navigation", "spatial", "spatial information", "spatial temporal",
"time task", and "visual spatial". The memory-spatial map was then created by summing
up the memory-related and navigation-related maps. Finally, this combined map was
overlapped with the 400 regions of interest (ROIs) from the "Schaefer 2018 parcellation"
to yield 242 ROIs, each containing a minimum of 30 voxels.
1134
fMRI data preprocessing
We performed fMRI data preprocessing using FEAT (FMRI Expert Analysis
Tool), version 6.00, implemented in FSL (http://www.fmrib.ox.ac.uk/fsl). The EPI images
underwent motion correction, slice-timing correction, and temporal filtering with a
nonlinear high-pass filter (100 s cutoff). Six motion parameters were included as
confounding regressors in the model. Additionally, outlier timepoints which identified
using the FSL's motion outlier detection tool (framewise displacement [FD] > 0.9 mm)
were incorporated as additional confounds in the first-level general linear model (GLM)
analysis. If more than 20% of the volumes in a run had an FD exceeding 0.9 mm, or if
the absolute head movement exceeded half the voxel size (i.e., 1.25 mm), the entire run
was excluded from further analyses. For single-trial estimation, no spatial smoothing
was applied. All functional images were linearly registered to the middle image of the
first run, and all analyses were conducted in MNI standard space.
1148
Univariate Activation Analysis
We examined encoding and retrieval-related neural activity using the general
linear model (GLM) within the FILM module of FSL. During the encoding stage, the
GLM included 5 regressors: (1) the encoding stage of all pictures (i.e., the 6 s of the trial
in which the participants were presented with an image and asked to encode that image
using a temporal or spatial context); (2) rating period (i.e., reaction time, RT) when the
participant made a rating judgment of all trials; (3) the remaining time period after the
participant made a rating judgment of all trials (i.e., 6s minus RT) while the stimulus

stayed on the screen;(4) The duration of inter-trial interval (ITI) fixations; (5) all the
instruction period within the same run. The "X" or "O" judgement baseline period was
not coded and thus was treated as an implicit baseline. Events were modeled at the
time of the stimulus onset and convolved with canonical hemodynamic response
function (double gamma function). The encoding phase consisted of four runs, equally
divided between spatial and temporal encoding runs. For the retrieval phase,
remembered and forgotten pictures were separately modeled for both item memory
retrieval and source memory retrieval. Source memory retrieval was further categorized
into spatial and temporal retrieval trials. The source memory retrieval effect was defined
as the difference in activity between correctly retrieved contextual pictures and the
implicit baseline.
A second-level analysis was conducted to compute cross-run averages for
spatial encoding effects, temporal encoding effects, combined encoding effects (spatial
and temporal concatenated), spatial source memory retrieval effects, temporal source,
and combined source memory retrieval effects (spatial and temporal concatenated)
using a fixed-effects model. These contrasts were then used in a group analysis with a
random-effects model using full FMRIB's Local Analysis of Mixed Effect 1 with
automatic outlier detection[68,69]. Unless otherwise noted, group images were thresholded
using cluster detection statistics, with a height threshold of $z > 3.1$ and a cluster
probability of $P < 0.05$, corrected for whole-brain multiple comparisons using Gaussian
Random Field Theory. To specifically examine univariate activation within the
hippocampus, a key region of interest in our experiment, we conducted small volume
correction using the hippocampal mask derived from the Harvard-Oxford subcortical
atlas in MNI152 space.
1181
Single-trial response estimates
General linear models (GLMs) were conducted separately to estimate the
activation patterns for each of 80 encoding trials and 100 retrieval trials. In each single-
trial model, a Least Square–Separate (LS-S) approach was used, in which the trial of
interest was modeled as one regressor, with all other trials modeled as a separate
regressor[70]. Specifically, during the encoding stage, each single-trial GLM included six
regressors: (1) the trial of interest (i.e., the 6 s of the trial where the participants were
presented with an image and asked to encode that image using a temporal or spatial
context); (2) all other remaining trials within the same run; (3) rating period (i.e. reaction
time, RT) when the participant made a rating judgment of all trials; (4) the remaining
time period after participant made a rating judgment of all trials (i.e., 6s minus RT) while
the stimulus stayed on the screen; (5) The duration of inter-trial interval (ITI) fixations;
(6) the instruction period within the same run. The "X" or "O" judgement period was not
coded and thus was treated as an implicit baseline.
Similarly, for the retrieval stage, each single-trial GLM included six regressors:
(1) the trial of interest (i.e., the RT of the trial in which participants were asked to
retrieve source information for the image); (2) all other remaining source retrieval trials
within the same run; (3) the remaining time period after the participant made a source
judgment (i.e., 6 seconds minus RT) while the stimulus remained on the screen; (4) all
item retrieval periods in which participants judged whether they could recognize the item
from the encoding phase by pressing "Yes" or "No" within the same run; (5) the

remaining time period after the participant made a Yes/No judgment (i.e., 6 seconds
minus RT) while the stimulus stayed on the screen; and (6) all instruction periods within
the same run. The ITI fixation period was not modeled and was treated as an implicit
baseline. Each event was modeled at the time of stimulus onset and convolved with a
canonical hemodynamic response function (double gamma). To control the effects of
head motion, six motion parameters were included in the GLM model as a covariate, as
well as a regressor for each TR that was flagged as having greater framewise
displacement (FD) than 0.9 during preprocessing. The t-map for each trial was used for
multivariate pattern similarity analysis to increase the reliability by normalizing for
noise[71].

Multivariate pattern similarity analysis (MPS)
Multi-voxel pattern similarity (MPS) was used to measure the similarity of
activation patterns by calculating the correlation between trials that were either encoded
or correctly retrieved, within each ROI[72]. Following the approach of Power[73], volumes
with a framewise displacement greater than 0.9 mm were censored, and trials
containing any censored frames within the duration of the modeled GLM response were
excluded. Specifically, within-context pattern similarity (PS) during encoding was
calculated using pairwise Pearson correlation coefficients between trials within the
same context (i.e., spatial or temporal). Within-spatial context was calculated using only
spatial trials, while within temporal context was calculated using only temporal trials. For
source memory retrieval, we performed analogous calculations for within-context PS,
separately measuring within-spatial context PS and within-temporal context PS, using
correct retrieval trials. All MPS analyses were performed across trials from different runs
to avoid temporal autocorrelations that could otherwise inflate or bias results. The
resulting correlation coefficients were then transformed into Fisher's z-scores.

Informational connectivity analysis
We performed informational connectivity among the 242 predefined ROIs by
calculating pairwise correlations between representational similarity matrices (RSMs)[74-
78]. This was done by correlating the within-context PS for each ROI with another ROI. In
contrast to traditional functional connectivity (FC) analyses, which quantify temporal
correlations of BOLD signal fluctuations between brain regions, the informational
connectivity approach examines the similarity of information coding patterns across
regions[75,76,78]. While FC analyses rely on averaged BOLD signals within regions,
representational similarity analysis (RSA) captures the fine-grained spatial patterns of
neural responses to different experimental conditions[77]. Specifically, RSA quantifies
how similarly or differently a brain region responds to different stimuli or conditions by
comparing multivariate activity patterns, creating a RSM that characterizes each
region's representational structure[77]. By correlating these RSMs between regions, we
can assess whether different brain areas encode information in similar ways, providing
insights into information sharing and processing that complement traditional FC
measures.
Three types of within-context RSMs were calculated: one based on all trials, one
based on spatial trials, and one based on temporal trials (Fig. 4a top). All Pearson
correlation coefficients (r values) were Fisher-Z transformed prior to statistical analysis.

Significance thresholds for informational connectivity were determined using 10,000 permutation simulations. The reported results have been corrected for multiple comparisons using the FDR method, with a q-value threshold of less than 0.05.

Multivariate informational connectivity pattern distance analysis

To assess changes in multivariate informational connectivity patterns between pre and post, distances were computed for each ROI by comparing spatial and temporal informational connectivity patterns before and after tranining. Specifically, for each ROI, the informational connectivity with the remaining 241 ROIs was extracted separately from the spatial and temporal informational connectivity matrices (ICM). The distance between the two connectivity patterns was calculated using 1 minus the Pearson or Spearman correlation coefficient (Fig. 6a). Paired t-tests were conducted to evaluate statistical differences between pre and post, with FDR correction applied across the 242 ROIs at a q-value threshold of < 0.05.

References

Kraft, J. N. *et al.* Task-based functional connectivity of the useful field of view (UFOV) FMRI task. *GeroScience* **45**, 293-309 (2023).
Edwards, J. D., Fausto, B. A., Tetlow, A. M., Corona, R. T. & Valdés, E. G. Systematic review and meta-analyses of useful field of view cognitive training. *Neuroscience & Biobehavioral Reviews* **84**, 72-91 (2018).
Maguire, E. A. *et al.* Navigation-related structural change in the hippocampi of taxi drivers. *Proc Natl Acad Sci U S A* **97**, 4398-4403, doi:10.1073/pnas.070039597 (2000).
Draganski, B. *et al.* Changes in grey matter induced by training. *Nature* **427**, 311-312 (2004).
Boyke, J., Driemeyer, J., Gaser, C., Büchel, C. & May, A. Training-Induced Brain Structure Changes in the Elderly. *The Journal of Neuroscience* **28**, 7031-7035, doi:10.1523/jneurosci.0742-08.2008 (2008).
Wenger, E., Papadaki, E., Werner, A., Kühn, S. & Lindenberger, U. Observing Plasticity of the Auditory System: Volumetric Decreases Along with Increased Functional Connectivity in Aspiring Professional Musicians. *Cerebral Cortex Communications* **2**, doi:10.1093/texcom/tgab008 (2021).
Hänggi, J., Koeneke, S., Bezzola, L. & Jäncke, L. Structural neuroplasticity in the sensorimotor network of professional female ballet dancers. *Human Brain Mapping* **31**, 1196-1206, doi:https://doi.org/10.1002/hbm.20928 (2010).
Draganski, B. *et al.* Temporal and Spatial Dynamics of Brain Structure Changes during Extensive Learning. *The Journal of Neuroscience* **26**, 6314-6317, doi:10.1523/jneurosci.4628-05.2006 (2006).
Engvig, A. *et al.* Effects of memory training on cortical thickness in the elderly. *NeuroImage* **52**, 1667-1676, doi:https://doi.org/10.1016/j.neuroimage.2010.05.041 (2010).

1294   10   Dresler, M. *et al.* Mnemonic training reshapes brain networks to support superior
1295   memory. *Neuron* **93**, 1227-1235. e1226 (2017).
1296   11   Braun, U. *et al.* Dynamic reconfiguration of frontal brain networks during
1297   executive cognition in humans. *Proc Natl Acad Sci U S A* **112**, 11678-11683,
1298   doi:10.1073/pnas.1422487112 (2015).
1299   12   Spiers, H. J. *et al.* Unilateral temporal lobectomy patients show lateralized
1300   topographical and episodic memory deficits in a virtual town. *Brain* **124**, 2476-
1301   2489 (2001).
1302   13   Buzsaki, G. & Moser, E. I. Memory, navigation and theta rhythm in the
1303   hippocampal-entorhinal system. *Nature neuroscience* **16**, 130-138,
1304   doi:10.1038/nn.3304 (2013).
1305   14   Brunec, I. K. *et al.* Cognitive mapping style relates to posterior–anterior
1306   hippocampal volume ratio. *Hippocampus* **29**, 748-754 (2019).
1307   15   Poppenk, J., Evensmoen, H. R., Moscovitch, M. & Nadel, L. Long-axis
1308   specialization of the human hippocampus. *Trends in cognitive sciences* **17**, 230-
1309   240 (2013).
1310   16   Clark, I. A. *et al.* Does hippocampal volume explain performance differences on
1311   hippocampal-dependant tasks? *NeuroImage* **221**, 117211 (2020).
1312   17   Weisberg, S. M., Newcombe, N. S. & Chatterjee, A. Everyday taxi drivers: Do
1313   better navigators have larger hippocampi? *Cortex* **115**, 280-293,
1314   doi:10.1016/j.cortex.2018.12.024 (2019).
1315   18   Sala, G. & Gobet, F. Does Far Transfer Exist? Negative Evidence From Chess,
1316   Music, and Working Memory Training. *Current Directions in Psychological
1317   Science* **26**, 515-520, doi:10.1177/0963721417712760 (2017).
1318   19   Ekstrom, A. D. & Hill, P. F. Spatial navigation and memory: A review of the
1319   similarities and differences relevant to brain models and age. *Neuron* **111**, 1037-
1320   1049, doi:10.1016/j.neuron.2023.03.001 (2023).
1321   20   Fan, C. L., Abdi, H. & Levine, B. On the relationship between trait
1322   autobiographical episodic memory and spatial navigation. *Mem Cognition* **49**,
1323   265-275 (2021).
1324   21   Weisberg, S. M. & Newcombe, N. S. How do (some) people make a cognitive
1325   map? Routes, places, and working memory. *J Exp Psychol Learn Mem Cogn* **42**,
1326   768-785, doi:10.1037/xlm0000200 (2016).
1327   22   Marek, S. *et al.* Reproducible brain-wide association studies require thousands of
1328   individuals. *Nature* **603**, 654-660, doi:10.1038/s41586-022-04492-9 (2022).
1329   23   Jaeggi, S. M., Buschkuehl, M., Shah, P. & Jonides, J. The role of individual
1330   differences in cognitive training and transfer. *Memory & cognition* **42**, 464-480
1331   (2014).
1332   24   Schweizer, S., Grahn, J., Hampshire, A., Mobbs, D. & Dalgleish, T. Training the
1333   emotional brain: improving affective control through emotional working memory
1334   training. *J Neurosci* **33**, 5301-5311, doi:10.1523/jneurosci.2593-12.2013 (2013).
1335   25   Karbach, J. & Verhaeghen, P. Making working memory work: a meta-analysis of
1336   executive-control and working memory training in older adults. *Psychol Sci* **25**,
1337   2027-2037, doi:10.1177/0956797614548725 (2014).
1338   26   Simons, D. J. *et al.* Do "brain-training" programs work? *Psychological science in
1339   the public interest* **17**, 103-186 (2016).

Gobet, F. & Sala, G. Cognitive Training: A Field in Search of a Phenomenon. *Perspect Psychol Sci* **18**, 125-141, doi:10.1177/17456916221091830 (2023).

Bouffard, N., Stokes, J., Kramer, H. J. & Ekstrom, A. D. Temporal encoding strategies result in boosts to final free recall performance comparable to spatial ones. *Memory & Cognition* **46**, 17-31, doi:10.3758/s13421-017-0742-z (2018).

Weisberg, S. M., Schinazi, V. R., Newcombe, N. S., Shipley, T. F. & Epstein, R. A. Variations in cognitive maps: understanding individual differences in navigation. *Journal of Experimental Psychology: Learning, Memory, and Cognition* **40**, 669 (2014).

Brunec, I. K. *et al.* Multiple scales of representation along the hippocampal anteroposterior axis in humans. *Current biology* **28**, 2129-2135. e2126 (2018).

Angeli, P. A., DiNicola, L. M., Saadon-Grosman, N., Eldaief, M. C. & Buckner, R. L. Specialization of the human hippocampal long axis revisited. *Proceedings of the National Academy of Sciences* **122**, e2422083122 (2025).

Destrieux, C., Fischl, B., Dale, A. & Halgren, E. Automatic parcellation of human cortical gyri and sulci using standard anatomical nomenclature. *Neuroimage* **53**, 1-15 (2010).

Mooraj, Z. *et al.* Toward a functional future for the cognitive neuroscience of human aging. *Neuron* **113**, 154-183, doi:10.1016/j.neuron.2024.12.008 (2025).

Maguire, E. A., Valentine, E. R., Wilding, J. M. & Kapur, N. Routes to remembering: the brains behind superior memory. *Nat Neurosci* **6**, 90-95, doi:10.1038/nn988 (2003).

Stern, Y. Cognitive reserve in ageing and Alzheimer's disease. *The Lancet Neurology* **11**, 1006-1012 (2012).

Lövdén, M. *et al.* Spatial navigation training protects the hippocampus against age-related changes during early and late adulthood. *Neurobiology of aging* **33**, 620. e629-620. e622 (2012).

McLaren-Gradinaru, M. *et al.*   (2020).

Burgess, N., Maguire, E. A. & O'Keefe, J. The Human Hippocampus and Spatial and Episodic Memory. *Neuron* **35**, 625-641, doi:10.1016/S0896-6273(02)00830-9 (2002).

Lövdén, M. *et al.* Spatial navigation training protects the hippocampus against age-related changes during early and late adulthood. *Neurobiology of Aging* **33**, 620.e629-620.e622, doi:https://doi.org/10.1016/j.neurobiolaging.2011.02.013 (2012).

Bassett, D. S., Yang, M., Wymbs, N. F. & Grafton, S. T. Learning-induced autonomy of sensorimotor systems. *Nature neuroscience* **18**, 744-751 (2015).

Finc, K. *et al.* Dynamic reconfiguration of functional brain networks during working memory training. *Nature communications* **11**, 2435 (2020).

Schedlbauer, A. M. & Ekstrom, A. D. Flexible network community organization during the encoding and retrieval of spatiotemporal episodic memories. *Network Neuroscience* **3**, 1070-1093 (2019).

Bouffard, N., Stokes, J., Kramer, H. J. & Ekstrom, A. D. Temporal encoding strategies result in boosts to final free recall performance comparable to spatial ones. *Mem Cognit* **46**, 17-31, doi:10.3758/s13421-017-0742-z (2018).

Squire, L. R., Stark, C. E. & Clark, R. E. The medial temporal lobe. *Annu. Rev. Neurosci.* **27**, 279-306 (2004).

Jenkinson, M., Beckmann, C. F., Behrens, T. E., Woolrich, M. W. & Smith, S. M. Fsl. *Neuroimage* **62**, 782-790 (2012).

Tustison, N. J. *et al.* The ANTsX ecosystem for quantitative biological and medical imaging. *Scientific reports* **11**, 9068 (2021).

Tournier, J.-D. *et al.* MRtrix3: A fast, flexible and open software framework for medical image processing and visualisation. *Neuroimage* **202**, 116137 (2019).

Cordero-Grande, L., Christiaens, D., Hutter, J., Price, A. N. & Hajnal, J. V. Complex diffusion-weighted image estimation via matrix recovery under general noise models. *Neuroimage* **200**, 391-404 (2019).

Veraart, J., Fieremans, E. & Novikov, D. S. Diffusion MRI noise mapping using random matrix theory. *Magnetic resonance in medicine* **76**, 1582-1593 (2016).

Veraart, J. *et al.* Denoising of diffusion MRI using random matrix theory. *Neuroimage* **142**, 394-406 (2016).

Andersson, J. L., Graham, M. S., Zsoldos, E. & Sotiropoulos, S. N. Incorporating outlier detection and replacement into a non-parametric framework for movement and distortion correction of diffusion MR images. *Neuroimage* **141**, 556-572 (2016).

Bastiani, M. *et al.* Automated quality control for within and between studies diffusion MRI data using a non-parametric framework for movement and distortion correction. *Neuroimage* **184**, 801-812 (2019).

Smith, S. M. *et al.* Advances in functional and structural MR image analysis and implementation as FSL. *Neuroimage* **23**, S208-S219 (2004).

Mitchell, T. *et al.* Advanced diffusion imaging to track progression in Parkinson's disease, multiple system atrophy, and progressive supranuclear palsy. *NeuroImage: Clinical* **34**, 103022 (2022).

Pasternak, O., Sochen, N., Gur, Y., Intrator, N. & Assaf, Y. Free water elimination and mapping from diffusion MRI. *Magnetic Resonance in Medicine: An Official Journal of the International Society for Magnetic Resonance in Medicine* **62**, 717-730 (2009).

Wilkes, B. J. *et al.* Cortico-basal ganglia white matter microstructure is linked to restricted repetitive behavior in autism spectrum disorder. *Molecular Autism* **15**, 6 (2024).

Wilkes, B. J. *et al.* Distinct cortical and subcortical predictors of Purdue Pegboard decline in Parkinson's disease and atypical parkinsonism. *npj Parkinson's Disease* **9**, 85 (2023).

Shin, Y. S. *et al.* Transcallosal white matter and cortical gray matter variations in autistic adults aged 30–73 years. *Molecular Autism* **16**, 16, doi:10.1186/s13229-025-00652-6 (2025).

Yeh, F.-C. Population-based tract-to-region connectome of the human brain and its hierarchical topology. *Nature communications* **13**, 4933 (2022).

Schwarz, C. *et al.* (2018).

Ekstrom, A. D., Huffman, D. J. & Starrett, M. Interacting networks of brain regions underlie human spatial navigation: a review and novel synthesis of the literature. *Journal of neurophysiology* **118**, 3328-3344 (2017).

62   Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)* **57**, 289-300 (1995).

63   Yushkevich, P. A. *et al.* Automated volumetry and regional thickness analysis of hippocampal subfields and medial temporal cortical structures in mild cognitive impairment. *Hum Brain Mapp* **36**, 258-287, doi:10.1002/hbm.22627 (2015).

64   Yushkevich, P. A. *et al.* Quantitative comparison of 21 protocols for labeling hippocampal subfields and parahippocampal subregions in in vivo MRI: towards a harmonized segmentation protocol. *Neuroimage* **111**, 526-541, doi:10.1016/j.neuroimage.2015.01.004 (2015).

65   Zeineh, M. M., Engel, S. A., Thompson, P. M. & Bookheimer, S. Y. Unfolding the human hippocampus with high resolution structural and functional MRI. *The Anatomical Record: An Official Publication of the American Association of Anatomists* **265**, 111-120 (2001).

66   Reuter, M., Schmansky, N. J., Rosas, H. D. & Fischl, B. Within-subject template estimation for unbiased longitudinal image analysis. *Neuroimage* **61**, 1402-1418 (2012).

67   Reuter, M., Rosas, H. D. & Fischl, B. Highly accurate inverse consistent registration: a robust approach. *Neuroimage* **53**, 1181-1196 (2010).

68   Beckmann, C. F., Jenkinson, M. & Smith, S. M. General multilevel linear modeling for group analysis in FMRI. *Neuroimage* **20**, 1052-1063, doi:10.1016/s1053-8119(03)00435-x (2003).

69   Woolrich, M. Robust group analysis using outlier inference. *Neuroimage* **41**, 286-301 (2008).

70   Mumford, J. A., Turner, B. O., Ashby, F. G. & Poldrack, R. A. Deconvolving BOLD activation in event-related designs for multivoxel pattern classification analyses. *Neuroimage* **59**, 2636-2643, doi:10.1016/j.neuroimage.2011.08.076 (2012).

71   Walther, A. *et al.* Reliability of dissimilarity measures for multi-voxel pattern analysis. *Neuroimage* **137**, 188-200 (2016).

72   Kriegeskorte, N., Goebel, R. & Bandettini, P. Information-based functional brain mapping. *Proceedings of the National Academy of Sciences* **103**, 3863-3868 (2006).

73   Power, J. D. *et al.* Methods to detect, characterize, and remove motion artifact in resting state fMRI. *Neuroimage* **84**, 320-341, doi:10.1016/j.neuroimage.2013.08.048 (2014).

74   Kriegeskorte, N., Mur, M. & Bandettini, P. Representational similarity analysis - connecting the branches of systems neuroscience. *Front Syst Neurosci* **2**, 4, doi:10.3389/neuro.06.004.2008 (2008).

75   Gao, Z. *et al.* Context free and context-dependent conceptual representation in the brain. *Cerebral Cortex* **33**, 152-166, doi:10.1093/cercor/bhac058 (2022).

76   Zheng, L. *et al.* Hippocampal contributions to novel spatial learning are both age-related and age-invariant. *Proceedings of the National Academy of Sciences* **120**, e2307884120, doi:doi:10.1073/pnas.2307884120 (2023).

77   Kriegeskorte, N., Mur, M. & Bandettini, P. A. Representational similarity analysis-connecting the branches of systems neuroscience. *Frontiers in systems neuroscience* **2**, 249 (2008).

78   Aly, M. & Turk-Browne, N. B. Attention promotes episodic encoding by stabilizing
1478   hippocampal representations. *Proceedings of the National Academy of Sciences*
1479   **113**, E420-E429, doi:doi:10.1073/pnas.1518931113 (2016).

1480