Engineering ATP Import in Yeast Uncovers a Synthetic Route to Extend Cellular Lifespan

Naci Oz[1,2], Hetian Su[3], Vedat Sari[4], Praveen Patnaik[5], Rohil Hameed[4], Jong Hee Song[6], Derek C. Prosser[1], Vyacheslav M. Labunskyy[5], Vadim N. Gladyshev[7], Nan Hao[3], Alaattin Kaya[1,2,4#]

[1]School of Life Sciences and Sustainability, Virginia Commonwealth University, Richmond, VA, USA

[2]Integrative Life Sciences, Virginia Commonwealth University, Richmond, VA, USA

[3]Department of Molecular Biology, University of California San Diego, La Jolla, CA, USA

[4] Department of Biochemistry and Molecular Genetics, University of Virginia, Charlottesville, VA, USA

[5] Department of Dermatology, Boston University School of Medicine, Boston, USA

[6]Biomolecular Analysis Facility, School of Medicine, University of Virginia, Charlottesville, VA 22903, USA

[7]Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, USA

#Corresponding author: Department of Biochemistry and Molecular Genetics, University of Virginia, 1340 Jefferson Park Ave. Charlottesville, VA, USA. E-mail: qcc6zm@virginia.edu

Keywords: ATP, Mitochondria, Aging, Energy metabolism, Nucleotide transport

Major research organism: *Saccharomyces cerevisiae*

Abstract

Aging results from the gradual accumulation of molecular damage as a result of cellular processes and is characterized by impaired functions, most notably an age-related decline in ATP production. However, the causal relationship between cellular ATP homeostasis and aging has not been established. In this study, we used a novel approach by harnessing a nucleotide transporter from a eukaryotic intracellular parasite to facilitate the direct import of extracellular ATP into budding yeast cells, enabling us to effectively manipulate their intracellular ATP levels. We found that depletion of ATP significantly reduces lifespan, while the supplementation of ATP in the growth medium fully restores it thereby extending lifespan. Moreover, gene expression analysis revealed that elevated ATP levels inhibit catabolic processes, indicating a suppression of glucose metabolism. Overall, our study revealed the direct impact of cellular ATP homeostasis on lifespan regulation that has never been directly tested before. This work offers new insights into the bioenergetic control of aging and positions energy metabolism as a promising target for longevity interventions.

Significance

Cellular energy homeostasis is a crucial factor in determining the health and longevity of organisms. While intracellular ATP levels are tightly regulated, the idea that cells can directly take in extracellular ATP to influence metabolism has not been thoroughly explored. In this study, we engineered yeast cells to import external ATP and demonstrated that this approach significantly alters mitochondrial function, metabolic flow, and aging processes. Our findings show that ATP uptake inhibits catabolic pathways and enhances mitochondrial maintenance, thereby extending cellular lifespan through a novel and non-traditional mechanism. This research reveals an unexpected degree of metabolic flexibility and introduces a synthetic biology-based method to reprogram energy metabolism and longevity. The principles established in this study provide a new framework for understanding the role of cellular bioenergetics in aging, highlighting how the modulation of ATP availability can impact metabolic states and lifespan regulation.

Introduction

Aging, a complex and multifactorial process, is marked by the gradual decline in physiological function, culminating in increased susceptibility to diseases and mortality (1,2). Central to this process is mitochondrial dysfunction, a critical factor in the cellular and systemic manifestations of aging. (3–5). Mitochondria produce ATP through oxidative phosphorylation and regulate various cellular pathways such as calcium homeostasis, redox signaling, and apoptosis (6,7). Over the past decade, a growing body of research has established a strong connection between mitochondrial dysfunction and key characteristics of the aging process. Alterations such as somatic mutations in mitochondrial DNA (mtDNA), impairments in the respiratory chain, and elevated production of mitochondrial reactive oxygen species (ROS) have been implicated. These changes lead to oxidative stress and the accumulation of molecular damage, ultimately disrupting cellular function and promoting senescence and age-related decline. Furthermore, mitochondrial dysfunction, resulting from structural and functional changes, is linked to age-related degenerative diseases (4,8,9). However, recent evidence suggests that mild mitochondrial stress, known as mitohormesis, can promote adaptive responses, enhancing longevity through hormetic mechanisms (5,10–13).

Mitochondrial dynamics, encompassing fusion, fission, and mitophagy, are crucial in maintaining mitochondrial integrity and metabolic and bioenergetic function throughout the lifespan (14–16). As organisms age, disruption in these dynamics can lead to the accumulation of dysfunctional mitochondria. Furthermore, mitochondrial quality control, mediated by mitophagy and fission/fusion processes, declines with age, exacerbating mitochondrial metabolic dysfunction, resulting in an imbalance of energy supply and demand (8,9,14–19). Caloric restriction (CR), a well-established metabolic intervention to extend lifespan, physical activity, and pharmacological interventions targeting conserved longevity pathways improve mitochondrial biogenesis and reduce oxidative damage, indicating that metabolic interventions targeting mitochondria can mitigate age-related imbalance of energy supply and demand (20–23).

The concept of ATP homeostasis, essential for cellular metabolism and signaling, emerges as a pivotal factor in mitochondrial-mediated lifespan regulation (24–26). While several studies have

explored how mitochondrial dysfunction impacts aging, few have directly examined the role of intracellular ATP availability and energy supply and demand balance in modulating longevity (6,9,12,24). For example, in *Caenorhabditis elegans*, a mild reduction in mitochondrial metabolic function during developmental stages confers increased lifespan, highlighting the nuanced relationship between energy production and aging (12). Similarly, research on mild mitochondrial uncoupling in mice demonstrates that metabolic shifts favoring increased energy expenditure can reduce ROS production and extend lifespan (11,27).

In this study, we investigated the impact of intracellular ATP homeostasis on cellular aging. We employed a novel synthetic biology approach, introducing a nucleotide transporter gene from an intracellular parasite (28,29) into budding yeast to enhance the uptake of extracellular ATP. This strategy allowed us to explore whether restoring ATP homeostasis could extend lifespan. Our findings revealed that ATP depletion significantly shortens lifespan, while ATP supplementation restores it. Intriguingly, we also observed that extracellular ATP alone could extend lifespan without cellular uptake, suggesting the existence of ATP-sensing pathways that modulate aging independently of conventional energy metabolism. RNA sequencing data further supported our hypothesis, indicating that increased ATP availability downregulates key catabolic processes, including glucose metabolism, thereby altering energy homeostasis. These findings underscore the intricate relationship between energy metabolism and cellular aging, with potential implications for developing energy-based interventions to promote healthy aging.

Furthermore, our research suggests that mitochondrial bioenergetics, redox balance, and ATP homeostasis are tightly intertwined, with disruptions in these processes playing a central role in cellular lifespan. Overall, our study provides novel insights into the role of ATP homeostasis in lifespan regulation. Our findings highlight the potential of targeting ATP levels as a therapeutic strategy to delay aging and promote longevity. By elucidating the mechanisms underlying ATP-sensing pathways, we open new avenues for research to develop interventions that optimize energy metabolism to enhance healthspan and mitigate the impact of age-associated mitochondrial dysfunction.

Results

Overexpression of *NTT1* mediates ATP import into *S. cerevisiae* cells

To alter intracellular ATP levels through extracellular feeding, we created a yeast model expressing a nucleotide transporter (*NTT1*) from the intracellular fungal parasite *Encephalitozoon cuniculi*. *E. cuniculi* naturally uses this exchanger to import ATP from the cytosol of eukaryotic host cells to sustain its survival (28,29). To examine Ntt1 localization in yeast, we modified its sequence by fusing the Pleckstrin homology (PH) domain of *SLM1* to its N-terminus—a well-established plasma membrane targeting signal. We generated expression cassettes with or without a C-terminal GFP tag and integrated them into the genome of wild-type (*Wt-BY4741*) yeast. *NTT1* expression was driven by the constitutive *TEF1* promoter (**Fig. 1A**). Fluorescence microscopy confirmed plasma membrane localization of the PH-Ntt1 fusion protein (Fig. 1B). This engineered system enables the exchange of extracellular ATP for intracellular ADP via *NTT1* (**Fig. 1C**).

Next, to test whether *NTT1* expression mediates ATP import from the extracellular environment, we created a second yeast cell model by integrating the functional *NTT1*-cassette (without GFP) into the genome of a Wt yeast. We also genome-integrated a fluorescent ATP biosensor (QUEEN), to quantify intracellular ATP levels in single living yeast cells (30,31) (**Fig. 2A**). Based on this reporter strain, our analysis showed that *NTT1* expression in cells severely depleted intracellular ATP levels, compared to Wt control cells (**Fig. 2B**). Supplementing the medium with 5 mM ATP increased ATP significantly (p=7.5E-09) exceeding levels observed in Wt controls, indicating efficient ATP uptake via Ntt1 from the extracellular environment (**Supplementary File 1**). Interestingly, Wt cells lacking *NTT1* exhibited a significant increase in ATP levels (p = 2.22E-16) upon ATP supplementation, though to a lesser extent than NTT1-expressing cells (**Fig. 2B**). Since *S. cerevisiae* lacks known ATP transporters, this suggests that extracellular ATP may activate downstream signaling via unknown nucleotide-sensing receptors, promoting mitochondrial function and ATP synthesis. Alternatively, an unidentified transporter might facilitate ATP uptake.

To validate these findings, we designed an experiment to inhibit endogenous ATP synthesis by treating cells with 2-deoxyglucose (2DG), a glucose analog that competitively inhibits glycolysis. Once phosphorylated by hexokinase, 2DG becomes 2DG-6-phosphate, which

accumulates and inhibits glycolysis, leading to rapid ATP depletion (32). As expected, 30 minutes of 2DG treatment sharply reduced intracellular ATP levels in both wild-type and NTT1-expressing cells (**Fig. 2C**). When NTT1-expressing cells were subsequently treated with 5 mM ATP, intracellular ATP levels increased, confirming ATP import via *NTT1*. In contrast, ATP treatment did not restore ATP levels in wild-type cells, arguing against the presence of an unidentified ATP transporter (**Fig. 2C and Supplementary File 1**).

Finally, we further validated the *NTT1*-mediated ATP import by analyzing cells with LC-MS/MS after ATP treatments. Consistent with the QUEEN measurement of controls cells, our analysis revealed significantly decreased ATP level under non-treated condition in *NTT1*-cells in comparison to the Wt controls (p=0.008) (**Fig. 2D**). Similarly, 30 minutes treatment of 1mM exogenous ATP caused 6-fold increase in intracellular ATP in *NTT1* expressing cells (p=0.035), and similar result was also observed under the condition of 5 mM ATP treatment (p=0.029). For the Wt cells, although we detected an increase trend in ATP level under 1 mM treatment, the same effect was not observed under 5 mM condition (**Fig. 2D**) and analyses revealed no significant alteration under both conditions. The two methods may have different sensitivities and linear ranges, but these data further validate our observation that Wt cells do not import exogenous ATP. The QUEEN reporter provided real-time, dynamic measurements of ATP concentration in living cells under continuous treatment of freshly supplemented medium conditions, mediated by NTT1 import mechanism. This allows it to capture rapid, transient increases or fluxes in ATP that occur immediately after the continuous exogenous treatment. However, since there is ATP import in Wt cells, ATP treatment though continuously supplemented medium might be needed to induce downstream signaling leading to ATP synthesis. Furthermore, LC-MS/MS typically involves sample extraction and processing steps that effectively provide a "snapshot" of the average, steady-state ATP levels across the cell population at a single point in time. The transient slow alteration of ATP abundance detected by QUEEN in Wt cells may be rapidly metabolized or pumped out of the WT cells before the LC-MS/MS sample preparation. Considering the rapid import and increased abundance of ATP in *NTT1* cells, which is detected using both QUEEN and LC-MS/MS, Wt cells might need longer and continuous treatment conditions. Overall, these data further validate modulation of intracellular ATP levels through extracellular feeding in cells harboring functional *NTT1*.

Altered ATP level targets cellular metabolic pathways

To investigate the effect of exogenous ATP treatment on gene expression in Wt and *NTT1*-expressing cells, we performed RNA-seq analysis (**Fig. 3A**). Our goal was to distinguish molecular signatures triggered by extracellular ATP-mediated signaling in wild-type cells from those driven by increased intracellular ATP via Ntt1-mediated uptake in engineered cells. We analyzed differentially expressed genes (DEGs) under ATP-treated and untreated conditions for both strains and compared their expression profiles (**Fig. 3B and C**). In Wt cells, ATP treatment significantly altered the expression of 500 genes: 198 upregulated and 302 downregulated. In contrast, *NTT1*-expressing cells displayed a broader response: 1,034 genes were significantly changed, with 586 upregulated and 448 downregulated. Among these, 146 upregulated and 188 downregulated genes were shared between the two strains, suggesting that these changes result directly from ATP treatment and occur independently of *NTT1* expression (**Fig. 3B and C**).

To further dissect ATP-mediated responses in Wt cells, we performed Gene Ontology (GO) (**Fig. S1A, B**) and KEGG pathway enrichment analyses (**Fig. S1C, D**). ATP treatment led to significant upregulation of genes involved in ribosome biogenesis, ATP biosynthesis, MAP kinase (MAPK) signaling, and oxidative phosphorylation (**Fig. S1A, C**). Conversely, downregulated genes were enriched in carbohydrate metabolism and uptake, protein phosphorylation, autophagy, and meiosis pathways (**Fig. S1B, D**). These findings indicate that extracellular ATP triggers intracellular signaling cascades capable of modulating mitochondrial function, cytosolic translation, and protein phosphorylation (**Supplementary File S2**). Notably, genes in the MAPK cascade were significantly enriched (**Fig. S2**), suggesting this pathway might play a key role in mediating the downstream ATP response in Wt cells.

Next, we applied the same enrichment analyses for the DEGs identified by comparing *NTT1*-expressing cells (treated vs non-treated). We found a complete rewiring of transcriptional regulation of metabolic genes in *NTT1*-expressing cells following ATP treatment. Similar to Wt cells, *NTT1*-expressing cells also showed upregulation of ribosome biogenesis and RNA processing genes (**Fig. 3D and Fig. S3**). However, while ATP treatment upregulated genes associated with mitochondrial respiratory activity, oxidative phosphorylation, and MAPK

signaling in Wt cells (**Fig. S1**), genes from these pathways were downregulated in *NTT1* cells (**Fig. 3D and Fig. S3**).

Further, the downregulated gene set in *NTT1*-expressing cells was enriched in glucose uptake, energy generation from precursor metabolites, carbohydrate phosphorylation, general catabolic processes, and autophagy (**Fig. 3E and Fig. S3**). This repression mirrors *SNF1*-mediated metabolic rewiring. *SNF1*, the yeast homolog of mammalian AMP-activated protein kinase (AMPK), plays a pivotal role in energy homeostasis by modulating metabolic responses and glucose de-repression under varying nutrients and ATP levels (32). Low *SNF1* activity is expected under high ATP concentrations (33).

To confirm this, we examined the expression of *SNF1* downstream targets associated with nutrient stress. We observed significant downregulation of several genes, including *MIG1*, *TDA1*, *HXK1*, *HXK2*, and *HXT3*, which are involved in glucose transport and metabolism (**Fig. S4; Supplementary File S2**). Additionally, *RGT1*, a major transcription factor that represses glucose transporter (*HXT*) genes under high-glucose conditions, was significantly upregulated in *NTT1* cells (**Supplementary File S2**). Together, these results indicate repression of glucose uptake and metabolism, further supporting the presence of a global metabolic rewiring in response to ATP elevation in *NTT1*-expressing cells (34,35).

Collectively, our data revealed a striking biological phenomenon: *NTT1*-mediated ATP import suppresses glucose uptake and catabolism, dampening mitochondrial energy production and reconfiguring cellular metabolism. Given that catabolic pathways are a major source of intracellular damage through the production of reactive by-products, this reduction in metabolic flux may also minimize damage accumulation. Consequently, autophagy, a key mechanism for clearing damaged macromolecules, was downregulated in these cells. These findings highlight two mechanistically distinct but complementary models by which ATP influences cellular physiology and lifespan. In *NTT1*-expressing cells, ATP is directly imported from the extracellular environment, leading to a robust and specific rewiring of cellular metabolism characterized by the suppression of glucose uptake, glycolysis, and mitochondrial respiration. This *NTT1*-dependent ATP influx mimics a high-energy state, repressing energy-generating catabolic pathways and reducing the need for damage-clearing processes such as autophagy,

potentially preserving cellular integrity over time. In contrast, extracellular ATP also exerted effects in Wt cells lacking *NTT1*, suggesting a separate, *NTT1*-independent mechanism possibly mediated through extracellular ATP sensing. Based on our transcriptome data, altered stress-responsive signaling cascades, MAPK might mediate the effect of extracellular ATP, that cells can detect and respond to extracellular nucleotides without direct ATP uptake.

In yeast, multiple MAPK pathways branch off and share components to regulate various cellular responses. These responses include pheromone signaling, filamentous growth (also known as pseudohyphal or starvation-induced growth), the high osmolarity glycerol (HOG) pathway, and other stress response pathways (36–38) The pheromone response and filamentous growth pathways are both initiated by G-protein signaling and share common components such as the MAPKKK Ste11p and MAPKK Ste7p. In contrast, the HOG pathway comprises two parallel branches (Sln1 and Sho1) that converge at a common MAPKK (Pbs2) and MAPK (Hog1). (**Fig. S2**). Our data indicated that the expression pattern of gene components of G-protein signaling cascades might be regulated positively while the Hog1-mediated osmolarity pathway was repressed (**Fig. S2**). To further confirm this, we performed western blot analyses to investigate the expression and phosphorylation level of Hog1 (Hog1^P) protein. We found that ATP treatment decreased the Hog1^P level in both Wt and *NTT1* cells, further validating decreased activity thereby repression of the HOG pathway. We further investigated Hog1^P level in *rho^0* isolates of both cell types (**Fig. 4, Fig. S5**), in which we eliminated mitochondrial DNA (mtDNA). These cells are unable to perform oxidative phosphorylation (OXPHOS) because they lack the necessary components of the electron transport chain encoded by the mitochondrial genome (39,40). The HOG pathway both promotes and regulates mitochondrial function and selective mitochondrial degradation (37,41). Prolonged activation was also associated with mitochondria-mediated increase in reactive oxygen species (ROS) levels (36,41). Although, in Wt *rho^0* cells, both phosphorylated and non-phosphorylated levels decreased, we did not observe overall change in Hog1^P level under ATP treatment conditions. The same observation was also true for *NTT1 rho^0* cells; however, overall abundance of Hog1^P level was lower in comparison to the Wt *rho^0* cells. These data suggest that the presence and absence of respiratory active mitochondria can mediate signaling cascades differentially when cells are treated with extracellular ATP (**Fig. 4, Fig S5**).

Together, these data from two cell types (Wt versus *NTT1*-expressing cells) suggest that ATP can modulate cellular pathways through intracellular energy repletion, extracellular ATP sensing, and activating downstream signaling, revealing a compartment-specific role of ATP. The unexpected divergence of ATP treatment between *NTT1*-expressing cells and Wt underscores the complexity of regulating cellular energy demand, further validating ATP as both a metabolic substrate and a signaling molecule with profound implications in cellular homeostasis.

Altered intracellular ATP level modulates cellular lifespan

We further investigated the consequences of altered ATP abundance on cellular aging. Due to the unstable nature of ATP molecules in the culture medium, we employed a state-of-the-art microfluidic aging chamber (42–46), which allowed cells to be continuously exposed to a medium supplemented with fresh ATP. This system was also integrated with high-resolution fluorescence microscopy, enabling real-time measurement of ATP levels throughout the yeast lifespan at the single-cell level using the fluorescence intensity of the QUEEN-ATP reporter (30,31). Our analysis of ATP abundance throughout the yeast lifespan showed that yeast cells are born with low ATP levels, which gradually increase during their lifespan. Some cells completed their lifespan without any observable reduction in ATP abundance, while others showed a drastic decrease in ATP levels during late life (**Fig. 5A–D, Supplementary File S3**), consistent with previous observations supporting two modes of yeast lifespan, mediated by mitochondrial and/or *SIR2* function (42,46–49).

Consistent with our data presented in Figure 2, we also observed significantly lower ATP abundance in *NTT1*-expressing cells throughout their entire lifespan compared to Wt control cells (**Fig. 5A–C**). Furthermore, these cells displayed significantly reduced mean and maximum replicative lifespan (RLS), directly indicating that intracellular ATP depletion shortens lifespan (**Fig. 5D**). Next, we assessed RLS and age-associated ATP changes under ATP supplementation. We found that exposing *NTT1* cells to medium supplemented with 10 µM ATP restored intracellular ATP levels (**Fig. 5A–C**) and significantly (p = 4.03E-18) and increased both mean and maximum RLS to levels comparable to WT cells (**Fig. 5D**).

In line with our previous observations, our analysis also showed a significant increase in ATP levels and RLS in Wt control cells upon continuous ATP supplementation, although the observed changes in both ATP levels and RLS extension were not as pronounced as those in *NTT1*-expressing cells (**Fig. 5A–D**). These findings are consistent with our transcriptomic data, which suggests that extracellular ATP may activate downstream signaling through as-yet unidentified nucleotide-sensing receptors, thereby increasing mitochondrial function, transient ATP production, and other canonical pathways.

ATP homeostasis dictates aging pathway selection and longevity in yeast

To further understand how ATP availability influences aging dynamics, we examined the two known trajectories of yeast replicative aging, mode-1 and mode-2, under both ATP-treated and untreated conditions (42). These modes are distinguished by their daughter cell morphology: mode-1 cells produce elongated daughters, whereas mode-2 cells give rise to small, round daughters. At the molecular level, mode-1 aging is characterized by rDNA instability and nucleolar enlargement and fragmentation, yet mitochondria remain functional. In contrast, mode-2 cells exhibit impaired mitochondrial function (42,46–49). Consistent with a previous report (40), we found that approximately 60% of Wt cells aged through mode-1, while 40% aged through mode-2 under control (no ATP treatment) conditions. Cells aged through mode-1 exhibited significantly longer lifespans than mode-2 cells (**Fig. 6A**).

We previously showed that mode-1 cells can maintain a relatively constant intracellular ATP level during aging, whereas mode-2 cells show a rapid depletion of ATP due to mitochondrial dysfunction (48). Therefore, Ntt1-mediated ATP depletion may have a more dramatic effect on the lifespan of mode-1 cells than that of ATP-depleted mode-2 cells. As expected, under the condition of no ATP treatment, *NTT1* expression caused a significant reduction in the lifespan of mode-1 cells (Wt mode-1, median RLS = 19 vs. *NTT1* mode-1, median RLS = 14; p = 2.39E-4), but not in mode-2 cells (Wt mode-2, mean RLS = 14 vs. *NTT1* mode-2, mean RLS = 14, p = 0.74) (**Fig. 5A**). We then analyzed the effect of ATP treatment. In both Wt and *NTT1*-expressing cells, ATP treatment increased the lifespans of mode-1 and mode-2 cells (**Fig. 6B**). Furthermore, the treatment also enhanced the proportion of mode-1 cells (**Fig.**

6C), suggesting that ATP treatment can compensate for the age-induced ATP decline in mode-2 cells, biasing the fate decision toward mode-1 aging.

These results indicate that elevating intracellular ATP, whether through *NTT1*-mediated import or via extracellular sensing, can extend cellular lifespan and influence the trajectory of aging. By restoring energy balance, ATP not only delays senescence but also shifts cells toward aging modes associated with greater mitochondrial integrity and extended replicative potential.

NTT1-driven ATP import alters lifespan through mitochondria-dependent mechanisms

To further understand the downstream effect of altered extracellular and intracellular ATP abundance and whether it interferes with mitochondrial function, we designed experiments with rho^0 isolates. RLS of Wt and *NTT1*-expressing cells, as well as their rho^0 isolates, were assayed under the control of higher dose (10 mM) ATP-supplemented medium conditions using a microfluidic aging chamber. Interestingly, under control conditions (without ATP supplementation), eliminating mtDNA prevented the adverse lifespan effect (**Fig. 5D and Fig. 7A, Supplementary File S3**) of *NTT1*-expression, consequently *NTT1* rho^0 cells displayed full lifespan to the level of Wt cells (**Fig. 7B**). It should be noted here that under the no ATP treatment condition, we did not observe any significant lifespan differences between Wt and Wt rho^0 cells (**Fig. 7A, B**), further confirming that eliminating mtDNA was the primary driver of the increased lifespan, specifically to the cells expressing *NTT1*. These data strongly suggest that the toxicity associated with *NTT1*-driven ATP import requires functional mitochondria. In other words, high intracellular ATP levels become deleterious only when mitochondrial respiration is intact.

Next, 10 mM ATP treatment resulted in 39% mean RLS increase (p =4.52E-12) in WT cells (**Fig. 7A**), comparable to the altered Wt lifespan that we observed under 10 µM treatment condition (**Fig. 5D**). However, in comparison to the 10 µM treatment condition, in which *NTT1* cells extended lifespan to the level of treated Wt cells (**Fig. 5D**), the positive effect of ATP treatment on lifespan was lost, indicating the increased ATP dosage causes toxicity and thereby significantly shortened (p= 1.25E-05) the lifespan in *NTT1*-expressing cells but not in Wt cells (**Fig. 7B**). However, we made an interesting observation that the toxicity effect of 10 mM ATP treatment on lifespan of *NTT1*-expressing cells was alleviated in *NTT1* rho^0 isolates. The 10 mM

ATP treatment significantly (p=3.07E-05) increased the lifespan of *NTT1 rho^0* cells (**Fig. 7B**), further supporting the conclusion that mitochondrial function mediates ATP toxicity.

Overall, these experiments allowed us to uncouple mitochondrial respiration from ATP-mediated lifespan effects. Strikingly, under baseline (no ATP treatment) conditions, the lifespan-shortening effect of *NTT1* expression was completely suppressed in *rho□* cells, suggesting that the toxicity of *NTT1*-driven ATP import requires functional mitochondria. In other words, high intracellular ATP becomes toxic only in the presence of intact mitochondrial function. Supporting this, at high extracellular ATP levels (10 mM), *NTT1*-expressing cells showed a shortened lifespan, whereas *NTT1 rho□* isolates were protected and displayed significantly extended lifespan, again implicating mitochondria as mediators of ATP toxicity. Conversely, in wild-type cells lacking *NTT1*, extracellular ATP sensing (without import) robustly extended lifespan in both mtDNA-positive and *rho□* backgrounds, demonstrating that ATP sensing acts independently of mitochondrial function and does not confer toxicity.

These combined observations point to a dual mechanism (**Fig. 8**): (i) intracellular ATP import through *NTT1* affects lifespan in a mitochondria-dependent manner, with potential toxicity at high levels; and (ii) extracellular ATP sensing promotes lifespan extension independently of mitochondrial respiration. Thus, mitochondria serve as a key integrator of energy input via *NTT1*, while also distinguishing between beneficial signaling and detrimental metabolic overload. Additionally, the observation that ATP treatment extended lifespan in both Wt and Wt *rho^0* isolates (**Fig. 7B**), indicating the positive lifespan effect of ATP sensing and downstream signaling cascade, mediated through MAPK is independent of mitochondrial function in Wt cells. In summary, these data further suggest the direct synergic interaction of *NTT1* function, altered ATP abundance on mitochondrial function, and differentiated effects of extracellular ATP treatment on Wt and *NTT1* cells, where uncharacterized extracellular nucleotide-sensing, potentially through G-protein signaling cascades of MAPK, initiating downstream signaling to induce cellular changes that alters lifespan in yeast.

Discussion

Catabolic systems (e.g., glycolysis, TCA cycle, respiration) are known to contribute significantly to cellular damage by generating by-products of cellular metabolism and reactive oxygen species

(ROS) (50–52). A major purpose of catabolism is to generate ATP (51). However, downregulating subsets of these systems has been shown to extend lifespan in various organisms (53–56). Interestingly, as cells age, their ability to produce ATP often declines, leading to impaired cellular function and contributing to the aging process (57–59). The effect of altered ATP abundance in the context of aging remains largely unclear. This question has been difficult to address because ATP generation is essential for all life forms.

However, some eukaryotic intracellular parasites have evolved the ability to obtain ATP from their hosts. For example, *E. cuniculi* naturally uses nucleotide transporters to import ATP from the cytosol of eukaryotic host cells into its own cytosol as well as cellular compartments, such as mitochondria, to sustain its obligate intracellular lifestyle (28,29). We applied xenotopic-based synthetic biology strategy (60) to incorporate a nucleotide transporter from *E. cuniculi* into a budding yeast model and investigated the effects of altered ATP abundance on aging. Specifically, we expressed the *E. cuniculi* nucleotide transporter gene (*NTT1*) in yeast to enable ATP import from the extracellular environment, thereby aiming to downregulate much of the cellular catabolism while preserving energy supplies. This novel approach of directly providing energy to cells in the form of ATP alleviated organelle dysfunction and revealed a complex picture of metabolic and phenotypic alteration resulting in lifespan extension.

We found that overexpression of *NTT1* led to severe depletion of intracellular ATP and reduced lifespan. Follow-up experiments in cells lacking mitochondrial respiratory function revealed that this effect was linked to mitochondrial stress, as rho^0 isolates (lacking mitochondrial DNA and OXPHOS) of *NTT1*-expressing cells restored lifespan to the level of WT controls. In addition, while low-dose ATP treatment significantly increased intracellular ATP levels and extended lifespan, high-dose ATP supplementation was detrimental to lifespan in *NTT1*-expressing cells but not in WT cells. However, *NTT1*- rho^0 isolates did not display decreased lifespan under toxic ATP concentrations.

These observations strongly indicate that *NTT1* overexpression and altered ATP abundance interfere with mitochondrial function. Lifespan mode analysis under both treated and non-treated conditions further confirmed that age-associated decline in mitochondrial function can be compensated for by ATP supplementation, thereby promoting a positive lifespan effect.

Our data showed that the percentage of cells aging through mitochondrial dysfunction (mode-2) was significantly reduced when the medium was supplemented with ATP. The majority of cells then followed a mode-1 aging trajectory, which is associated with a loss of Sir2 function. Interestingly, mode-1 cells also showed increased lifespan following ATP treatment. This suggests that elevated ATP levels are tightly connected to regulating cellular aging. In fact, previous studies that characterized these lifespan modes demonstrated that overexpression of *HAP4* (a transcription factor regulating mitochondrial respiratory function) (61) alone did not extend lifespan. However, combined overexpression of *SIR2* and *HAP4* had a synergistic lifespan-extending effect, where cells lived significantly longer than those overexpressing *SIR2* alone (42). This supports the idea that mitochondrial function and Sir2 activity work together to modulate lifespan.

Perhaps the most striking observation from our transcriptome data was that *NTT1*-expressing cells downregulated glucose uptake and catabolic pathways, including energy generation and mitochondrial respiration, when supplemented with ATP. This observation suggests that when cellular energy demand is supported through a direct import of ATP, cells can adapt by suppressing carbohydrate catabolic processes to balance energy homeostasis. Furthermore, the observed concurrence of decreased expression of autophagy pathway genes and lifespan extension can be attributed to the fact that catabolic processes are a major source of cellular damage through the generation of reactive by-products (e.g., oxidants and secondary metabolites). Reduced catabolic activity may therefore generate fewer damaging agents, leading to decreased activation of damage-clearing pathways such as autophagy.

Remarkably, extracellular ATP sensing independent of Ntt1 also exhibited lifespan-extending effects following ATP treatment in Wt cells. Our transcriptomics data suggests the presence of a previously unknown nucleotide-sensing mechanism that links extracellular ATP to downstream cellular processes in yeast. Considering the altered expression of genes enriched in MAPK and glucose homeostasis pathways in these cells, we hypothesize that the stress responsive MAPK pathway may link extracellular ATP sensing to downstream pathways that promote mitochondrial respiration and increase ATP abundance. In fact, extracellular ADP treatment has also been shown to increase intracellular ATP levels in yeast (31). This suggests the existence of specific nucleotide-sensing mechanisms in yeast, similar to mammalian purinergic

signaling, where purinergic receptors modulate downstream signaling cascades (62,63). A well-conserved *p38/HOG1* MAPK module in yeast is essential for cell survival under various stress conditions, such as osmotic, cell wall, nutrient, and oxidative stress (64–66). In addition, a wide variety of stimuli can elicit MAPK activation; however, it is still unclear how MAPKs regulate the distinct coupling of signals to downstream cellular responses (67,68). It is also known that MAPK is involved in cellular metabolic regulation and mitochondrial function and dynamics (69–71). Hence, our study suggests a previously uncharacterized MAPK–mitochondria signaling axis that regulates mitochondrial function in response to extracellular nucleotide-sensing and provides new insights into the molecular and physiological adaptations of cells to ATP homeostasis. However, the components of the upstream sensing and downstream signaling cascades connecting ATP sensing to mitochondrial function require further investigation. It should also be highlighted that the increased mitochondrial respiration in these cells was not essential for the lifespan extension mediated by ATP supplementation, indicating modulation of other downstream canonical pathways. Our data also suggests that the presence or absence of respiratory active mitochondria might mediate alternative pathways that differentially regulate cellular lifespan following ATP treatment in Wt cells.

In part, this positive lifespan effect may also be associated with increased mitochondrial function and ATP abundance in Wt cells. However, we observed that WT rho^0 isolates also showed significantly increased lifespan following ATP treatment, indicating that mitochondrial function is not essential for ATP signaling-mediated lifespan extension. It should be noted that caloric restriction (CR) also induces stress response pathways, increases respiration, and intracellular ATP abundance in yeast (72,73). However, CR-mediated lifespan extension is also independent of mitochondrial function, since WT rho^0 isolates extend lifespan under CR conditions (74). This suggests that ATP-sensing mechanisms may regulate overlapping CR-mediated pathways to induce lifespan extension and may modulate alternative pathways mediated by mitochondrial functional status. However, the exact mechanism of ATP-mediated lifespan extension in yeast requires further exploration.

Overall, our study directly validates that maintaining energy metabolism modulates longevity. Decline in mitochondrial function is considered one of the key hallmarks of aging, a finding supported by numerous previous studies. However, our study is novel in directly

demonstrating the role of ATP homeostasis in modulating cellular metabolism and aging. This research establishes that cellular energy balance and mitochondrial function are interconnected regulators of aging, and targeting ATP metabolism provides a promising strategy for extending lifespan and improving cellular health.

Materials and Methods

Yeast Strains and Plasmids

The Pleckstrin Homology (PH) domain of the *SLM1* gene was amplified from genomic DNA of the wild-type *S. cerevisiae* (BY4741) strain. Primers *PH-F* and *PH-R* were designed to amplify the PH domain with overlapping sequences compatible with assembly (primer sequences are listed in **Supplementary File S4**). The *NTT1* gene was amplified from a plasmid (28) using primers NTT1-F and NTT1-R, which included overlaps with the PH domain at the 5' end and GFP at the 3' end. The yeast-enhanced green fluorescent protein (yeGFP) sequence was amplified from the pYM25 plasmid (75) using primers GFP-F and GFP-R, incorporating overlaps with the *NTT1* sequence. The purified PH domain, *NTT1*, and yeGFP PCR products were fused using the NEBuilder HiFi DNA Assembly Master Mix (New England Biolabs, NEB #E2621).

The assembled *PH_NTT1_GFP* (**Supplementary File S4**) fusion construct was used as a template for PCR amplification with Gateway-compatible attB primers (*attB1-F* and *attB2-R*), which were designed to add recombination sites for Gateway cloning (primer sequences are provided in **Supplementary File S4**). The purified attB-flanked *PH_NTT1_GFP* fragment was recombined into the *pAG413GPD-ccdB* plasmid (Addgene plasmid #14142) (Alberti et al., 2007). The confirmed plasmid was transformed into BY4741 cells using the lithium acetate method.

A fusion cassette (**Supplementary File S4**) containing the PH sequence and *NTT1* (with a SmaI restriction site between them) was amplified by PCR from the appropriate DNA template using primers *Attb_PH_F* and *Attb_NTT1_R*. The reverse primer included a stop codon at the 3′ end of *NTT1*. The resulting attB-PCR product was purified and transferred into pAG306-GPD-ccdB (Addgene plasmid #41894). The final plasmid construct was integrated into the yeast genome

after NotI digestion (Hughes et al., 2012). The empty pAG306-GPD plasmid (Addgene plasmid #41895) was integrated into the yeast genome as a control strain.

Finally, the yeast strains containing the QUEEN ATP reporter gene were obtained from the National BioResource Project Yeast Genetic Resource Center. The *PH_NTT1* construct (with a stop codon) in pAG306-GPD-ccdB was integrated into the BY29034 strain. The empty pAG306-GPD plasmid was also integrated into the BY29034 genome as a control strain. All sequences were verified by Sanger sequencing after PCR amplification and cloning.

RNA Isolation and RNA-seq Analysis

Cells (three independent cultures for each strain) were cultivated in Complete Synthetic Medium (CSM) at 30°C with constant agitation (200–220 rpm) until they reached an optical density at 600 nm (OD600) of approximately 0.4–0.5. To examine the effect of extracellular ATP, a 50 mM ATP stock solution was prepared in CSM to maintain a consistent medium composition during treatment. Cells were treated with ATP at a final concentration of 5 mM for 1 hour at 30°C under continuous shaking. Untreated control cultures of the same strains were grown under identical conditions but without ATP supplementation.

After treatment, cells were harvested by centrifugation at 4°C. Total RNA was extracted using the Quick-RNA Fungal/Bacterial MiniPrep™ Kit (Zymo Research R2014) according to the manufacturer's instructions. The resulting RNA samples were used for downstream RNA-sequencing analysis.

To prepare RNA-seq libraries, the Illumina TruSeq RNA library preparation kit was used according to the user manual. RNA-seq libraries were loaded onto the Illumina NovaSeq 6000 platform to produce 150 bp paired-end sequences. After quality control and adapter removal, paired-end clean reads were aligned to the yeast reference genome (Ensembl, R64-1-1) using Hisat2 v2.0.5. FeatureCounts v1.5.0-p3 was used to count the number of reads mapped to each gene. Fragments per kilobase of transcript per million mapped reads (FPKM) values were calculated based on the length of the gene and the read count mapped to that gene.

Differential expression analysis was performed using the DESeq2 R package (v1.20.0). The resulting p-values were adjusted using the Benjamini and Hochberg approach for controlling the false discovery rate. Genes with an adjusted p-value ≤ 0.05 were classified as differentially expressed. Gene Ontology (GO) enrichment analysis of differentially expressed genes was implemented using the clusterProfiler R package, which corrects gene length bias. GO terms with corrected p-values less than 0.05 were considered significantly enriched among differentially expressed genes.

Microscopy for NTT1 Localization

The *S. cerevisiae* strain harboring *NTT1* was grown in yeast peptone dextrose (YPD) medium at 30°C with constant agitation (200–220 rpm) until mid-log phase. For visualization of *NTT1* localization, cells were harvested and immediately examined by confocal fluorescence microscopy using a GFP-compatible excitation/emission range (e.g., 488 nm excitation/505–530 nm emission). Images were acquired under identical settings for all samples to enable direct comparisons of fluorescence signals.

Relative ATP Measurement Using QUEEN ATP Reporter and LC-MS/MS

The *S. cerevisiae* strains expressing the QUEEN ATP reporter (NTT1 and WT control) were cultivated in Complete Synthetic Medium (CSM) at 30°C with constant agitation (200–220 rpm) until reaching mid-log phase (OD600 ~ 0.4–0.6). The cells were retained in CSM for one set of experiments and either treated with 5 mM ATP for 30 minutes or left untreated as a control.

In a separate experiment, cells were transferred to CSM supplemented with 20 mM 2DG (Sigma #D-8375) for 30 minutes to modulate intracellular ATP levels. Following the indicated treatments, cells were harvested, washed once in fresh CSM, and resuspended in CSM for microscopy. Cell viability was assessed using propidium iodide (PI) staining.

Images were acquired using a Leica DMi8 inverted fluorescence microscope equipped with 405, 488 and 561 nm lasers, an LED3 fluorescence illumination system, a 100X, 1.47 numerical aperture Plan-Apochromat oil immersion lens, a Flash 4.0 v3 sCMOS camera (Hamamatsu), a W-View Gemini beam splitting device (Hamamatsu), and LAS X v3.7.6.25997 software. All

strains within an experiment were imaged under identical exposure settings. A series of four images were captured for each field: one bright-field image and three fluorescence images corresponding to excitation/emission filters optimized for QUEEN (QWF-T ex405, QWF-T ex488), or for PI (QWF-T ex561).

Digital images were processed using Fiji software. First, the background signal was subtracted from the fluorescence channels after converting images to 32-bit floating-point format. Intracellular ATP was estimated by calculating a QUEEN ratio, defined as the pixel intensity of the ex405 image divided by the corresponding intensity of the ex488 image on a per-pixel basis. The mean QUEEN ratio for each cell was determined by averaging pixel ratios within a manually defined region of interest (ROI) encompassing the cell. Higher QUEEN ratios indicate elevated intracellular ATP levels. PI fluorescence (captured via ex561) was used to identify and exclude nonviable or compromised cells from the analysis. Based on the normality and homogeneity assumptions (Shapiro-Wilk and Levene's tests, respectively), a non-parametric Kruskal-Wallis test was chosen for comparison of groups. Post hoc pairwise comparisons using the Wilcoxon rank sum test with Bonferroni adjustment were used when comparing only two groups.

For the LC-MS/MS analyses, yeast cultures of both the *NTT1* mutant strain and the WT strain were grown in liquid medium at 30 °C with shaking at 210 rpm until mid-logarithmic phase (OD□□□ ≈ 0.4). A 50 mM ATP stock solution was freshly prepared from ATP disodium salt (Cayman Chemical, Cat. No. 21121). Cultures were divided into experimental groups consisting of untreated controls, 1 mM ATP, and 5 mM ATP supplementation. ATP was added to cultures at the indicated concentrations, and cells were incubated for 30 min before harvesting. Cells were collected by centrifugation at 5000 rpm for 10 min, washed three times in PBS, and quenched in 500 µL of 80% methanol (–20 °C). Quenched cells were homogenized by tip sonication for 30 seconds in an ice bath, followed by a 30-second pause, and the cycle was repeated three times. Lysates were transferred to 4 mL glass vials, and the tubes were rinsed with an additional 200 µL of 80% methanol, which was combined with the extract. Subsequently, 1.4 mL of cold chloroform (–20 °C) was added, and the mixture was vortexed for 30 seconds and shaken at 300 rpm for 10 min at 4 °C. After the addition of 700 µL of cold water, samples were shaken vigorously for 1 min and centrifuged at 3000 rpm for 5 min at 4 °C. The upper aqueous phase (1

mL) was collected, dried using a SpeedVac concentrator for approximately 3 h, and stored at –80 °C. Prior to LC-MS/MS measurement, dried extracts were reconstituted in 100 µL of formic acid in water containing isotopically labeled internal standards (Metabolomics QReSS, Cambridge Isotope Laboratories). A pooled quality control (QC) sample was prepared by combining 15 µL from each extract, and 10 µL was injected per run. Orbitrap IDX Tribrid mass spectrometer (Thermo Fisher Scientific) coupled with a Vanquish UHPLC system was used for detecting the ATP levels. Separation was achieved on a Waters BEH C18 column (2.1×150 mm, $1.8 \, \mu$m) maintained at 30 °C with a flow rate of 250 µL/min. Raw LC-MS/MS data were processed in MS-DIAL v4.92 (Tsugawa et al., 2015). Blank samples were included to identify and remove background signals. It is important to note that NTT1 and WT experiments were performed and analyzed on different days. Each strain was processed and analyzed independently, and all subsequent metabolite extraction, LC-MS/MS acquisition, and data processing were carried out separately.

Immunoblotting Bioassays

Hog1 phosphorylation was analyzed using antibodies that recognize the total and phosphorylated forms. Mid-log phase cells were grown on YPD medium, treated or not treated with 5mM ATP. Proteins were extracted using yeast protein extraction buffer YPER (Thermo, Catalog number 78990) supplemented with protease and phosphatase inhibitors mix (Thermo, Catalog number 87785). Whole cell extracts were resuspended in boiling SDS-PAGE sample buffer for 5 min. Following regular SDS-PAGE and transfer to nitrocellulose, the membrane was probed with antibodies to phospho-p38 (Hog1[p]) at 1:1000 (Cell Signaling Technology, Catalog number 9215T), Hog1 at 1:1000 (Santa Cruz, Catalog number sc-165978), Pgk1 at 1:5,000 (from Abcam, 113687). It is important to note that the phospho-p38 antibody Fallowing the treatment of secondary antibodies, membranes were developed using chemiluminescent substrate (BioRAD, Catalog number 1705060) and visualized with BioRAD systems.

Setting up Microfluidics Experiments

The microfluidics device fabrication and experimental setup have been described previously (44,46,76). Yeast cells were inoculated into 2 ml synthetic complete medium (SD, 2% glucose)

and incubated on a 30°C shaker overnight. From this culture, 2 µL of saturated culture was transferred to 20 ml SD and incubated until its OD$_{600}$ reached ~0.6 (early exponential growth phase). A microfluidics chip containing 4 microfluidics channels was placed under vacuum for at least 15 min. In the meanwhile, 50 ml SD with 0.04% Tween-20 (Sigma) was transferred to a 60 ml syringe, to which plastic tubing (TYGON, ID 0.020 IN, OD0.060 IN, wall 0.020 IN) was connected. For samples that have additional ATP in the medium, ATP was added to this 50 ml medium to reach the corresponding concentrations. The syringe was sealed with aluminum foil. 4 such syringes were prepared, 1 for each microfluidics channel. After vacuum, the microfluidics chip was quickly connected to the syringes through plastic tubing from its inlet port to avoid air flowing into the microfluidics channels. The outlets were then connected to plastic tubing filled with ddH$_2$O. The chip with inlets and outlets was then fixed to a microscope stage. For cell loading, cells were transferred into a 60 ml syringe with plastic tubing connected. Connection of the microfluidics inlets to the media supply syringe was then replaced by connection to the cell loading syringe. The flow of medium in the device was maintained by gravity to drive cell loading into the traps. After loading, the media supply tubing was switched back to the inlet. Heights of all supply syringes were then adjusted to have a height difference between inlet and outlet of around 240cm. Tween-20 is a non-ionic surfactant that helps reduce friction during cell loading. We have validated previously that this low concentration of Tween-20 has no significant effect on cellular lifespan or physiology (77).

For assaying rho^0 isolates, yeast cells were cultured overnight in synthetic complete (SC) media supplemented with 2% glucose. The following day, cultures were diluted 1:300 into 4 mL fresh SC media and incubated at 30°C for 3 hours. Prior to loading into the microfluidic device, 0.05% bovine serum albumin (BSA) was added to the culture to reduce nonspecific adhesion. Cells were then introduced into the microfluidic device, and all experiments were conducted using a Nikon Eclipse Ti2 inverted microscope equipped with a 60× 1.4 NA oil immersion objective (Nikon) and maintained at 30°C using a stage-top incubation system (Okolabs). Media was continuously delivered to the device via two 30 mL syringe pumps (total flow rate of 4 µL/min) to ensure nutrient availability and waste removal. Image acquisition was carried out every 5 minutes for up to 72 hours using NIS-Elements software, with the Nikon Perfect Focus System (PFS) engaged throughout all imaging sessions to preserve focal stability. Cell viability and

death events were monitored in real time, and all downstream image segmentation, cell tracking, and data extraction were performed using MATLAB (MathWorks).

Time-lapse Microscopy

Time-lapse Microscopy experiments were conducted using a Nikon Ti-E inverted fluorescence microscope with an EMCCD camera (Andor iXon X3 DU897). The light source is a spectra X LED system. Images were taken using a CFI plan Apochromat Lambda DM 60X oil immersion objective (NA 1.4 WD 0.13mm). In the yeast aging experiments, images were taken for each imaging channel every 15min for a total of 90 hours. The setting of each channel was set as the following and remains consistent across time-lapse imaging aging experiments: Phase, 60ms; GFP, 100ms at 10% lamp intensity with an EM gain of 100.

Quantification of Single-cell Aging Traces

The procedure of image quantification has been described previously (46). Image processing was done using ImageJ and custom Matlab codes. The background of fluorescence channel was subtracted by rolling ball background subtraction. Cells were masked and tracked by thresholding the GFP channel. The correctness of cell tracking was manually checked. The mean intensity of the top 40% pixels was quantified for each mother cell at each time point.

Cell aging mode was determined by daughter cell morphology (76). Cell replicative lifespan was manually quantified. Cells that showed obvious abnormal morphology upon loading were filtered out of RLS analysis.

Acknowledgments

We thank Dr. Robert P. Hirt (Newcastle University) for providing plasmids containing the cDNA-based expression cassette of *NTT1*, and Dr. Satoshi Yoshida (Waseda University, Japan) for providing strains and plasmids harboring the QUEEN ATP reporter vector through the National BioResource Project (NBRP-Japan) Yeast Genetic Resource Center. We also thank the NBRP staff for handling the collection and preparing and sending these strains. Additionally, we acknowledge the support provided by The Cancer Center Shared Resource for Metabolomics & Lipidomics at the University of Virginia for performing LC-MS/MS analyses.

Funding

This work was supported by NIH/NIGMS Grant R35GM150858 and the Impetus Grant (to A.K.) and NIH R01AG038004 (to V.N.G.). N.H. was supported by NIH R01AG086348, and NIH R01GM144595. D.C.P. was supported by an NSF CAREER Award (MCB 1942395).

Author contributions

A.K. and V.N.G. designed research; N.O., H.S., V.S., P.P., R.H., J.H.S., D.C.P., performed research. N.O., H.S., V.S., P.P., R.H., J.H.S., V.M.L., N.H., and A.K. analyzed data; and N.O., H.S., V.S., P.P., V.M.L., D.C.P., V.N.G., N.H., and A.K, wrote the paper.

Competing interests

The authors declare no competing interest

References:

1. Tenchov R, Sasso JM, Wang X, Zhou QA. Aging Hallmarks and Progression and Age-Related Diseases: A Landscape View of Research Advancement. ACS Chem Neurosci. 2024 Jan 3;15(1):1–30.
2. Gladyshev VN. The origin of aging: imperfectness-driven non-random damage defines the aging process and control of lifespan. Trends Genet. 2013 Sep;29(9):506–12.
3. Gonzalez-Freire M, de Cabo R, Bernier M, Sollott SJ, Fabbri E, Navas P, et al. Reconsidering the Role of Mitochondria in Aging. J Gerontol A Biol Sci Med Sci. 2015 Nov;70(11):1334–42.
4. Sun N, Youle RJ, Finkel T. The Mitochondrial Basis of Aging. Mol Cell. 2016 Mar 3;61(5):654–66.
5. Phua CZJ, Zhao X, Turcios-Hernandez L, McKernan M, Abyadeh M, Ma S, et al. Genetic perturbation of mitochondrial function reveals functional role for specific mitonuclear genes, metabolites, and pathways that regulate lifespan. Geroscience. 2023 Apr 22;45(4):2161–78.
6. Monzel AS, Enríquez JA, Picard M. Multifaceted mitochondria: moving mitochondrial science beyond function and dysfunction. Nat Metab. 2023 Apr;5(4):546–62.
7. Picard M, McEwen BS. Psychological Stress and Mitochondria: A Systematic Review. Psychosom Med. 2018;80(2):141–53.
8. Bratic A, Larsson NG. The role of mitochondria in aging. J Clin Invest. 2013 Mar;123(3):951–7.
9. Amorim JA, Coppotelli G, Rolo AP, Palmeira CM, Ross JM, Sinclair DA. Mitochondrial and metabolic dysfunction in ageing and age-related diseases. Nat Rev Endocrinol. 2022 Apr;18(4):243–58.
10. Lima T, Li TY, Mottis A, Auwerx J. Pleiotropic effects of mitochondria in aging. Nat Aging. 2022 Mar;2(3):199–213.
11. Caldeira da Silva CC, Cerqueira FM, Barbosa LF, Medeiros MHG, Kowaltowski AJ. Mild mitochondrial uncoupling in mice affects energy metabolism, redox balance and longevity. Aging Cell. 2008 Aug;7(4):552–60.
12. Maglioni S, Schiavi A, Runci A, Shaik A, Ventura N. Mitochondrial stress extends lifespan in C. elegans through neuronal hormesis. Exp Gerontol. 2014 Aug;56:89–98.
13. Ristow M, Schmeisser K. Mitohormesis: Promoting Health and Lifespan by Increased Levels of Reactive Oxygen Species (ROS). Dose Response. 2014 May;12(2):288–341.
14. Sharma A, Smith HJ, Yao P, Mair WB. Causal roles of mitochondrial dynamics in longevity and healthy aging. EMBO Rep. 2019 Dec 5;20(12):e48395.
15. Sebastián D, Palacín M, Zorzano A. Mitochondrial Dynamics: Coupling Mitochondrial Fitness with Healthy Aging. Trends Mol Med. 2017 Mar;23(3):201–15.
16. Chen W, Zhao H, Li Y. Mitochondrial dynamics in health and disease: mechanisms and potential targets. Signal Transduct Target Ther. 2023 Sep 6;8(1):333.
17. Picca A, Faitg J, Auwerx J, Ferrucci L, D'Amico D. Mitophagy in human health, ageing and disease. Nat Metab. 2023 Dec;5(12):2047–61.
18. Jiménez-Loygorri JI, Villarejo-Zori B, Viedma-Poyatos Á, Zapata-Muñoz J, Benítez-Fernández R, Frutos-Lisón MD, et al. Mitophagy curtails cytosolic mtDNA-dependent activation of cGAS/STING inflammation during aging. Nat Commun. 2024 Jan 27;15(1):830.
19. Berry BJ, Kaeberlein M. An energetics perspective on geroscience: mitochondrial protonmotive force and aging. Geroscience. 2021 Aug;43(4):1591–604.
20. McLean S, Lee M, Liu W, Hameed R, Gujjala VA, Zhou X, et al. Molecular Mechanisms of Genotype-Dependent Lifespan Variation Mediated by Caloric Restriction: Insight from Wild Yeast Isolates. bioRxiv. 2024 May 25;

21. Guarente L. Mitochondria--a nexus for aging, calorie restriction, and sirtuins? Cell. 2008 Jan 25;132(2):171–6.

22. O'Reilly CL, Miller BF, Lewis TL. Exercise and mitochondrial remodeling to prevent age-related neurodegeneration. J Appl Physiol (1985). 2023 Jan 1;134(1):181–9.

23. López-Lluch G, Hunt N, Jones B, Zhu M, Jamieson H, Hilmer S, et al. Calorie restriction induces mitochondrial biogenesis and bioenergetic efficiency. Proc Natl Acad Sci U S A. 2006 Feb 7;103(6):1768–73.

24. Picard M, McEwen BS, Epel ES, Sandi C. An energetic view of stress: Focus on mitochondria. Front Neuroendocrinol. 2018 Apr;49:72–85.

25. Moldakozhayev A, Gladyshev VN. Metabolism, homeostasis, and aging. Trends Endocrinol Metab. 2023 Mar;34(3):158–69.

26. Chaudhari SN, Kipreos ET. The Energy Maintenance Theory of Aging: Maintaining Energy Metabolism to Allow Longevity. Bioessays. 2018 Aug;40(8):e1800005.

27. Rohas LM, St-Pierre J, Uldry M, Jäger S, Handschin C, Spiegelman BM. A fundamental system of cellular energy homeostasis regulated by PGC-1alpha. Proc Natl Acad Sci U S A. 2007 May 8;104(19):7933–8.

28. Tsaousis AD, Kunji ERS, Goldberg A V, Lucocq JM, Hirt RP, Embley TM. A novel route for ATP acquisition by the remnant mitochondria of Encephalitozoon cuniculi. Nature. 2008 May 22;453(7194):553–6.

29. Heinz E, Hacker C, Dean P, Mifsud J, Goldberg A V, Williams TA, et al. Plasma membrane-located purine nucleotide transport proteins are key components for host exploitation by microsporidian intracellular parasites. PLoS Pathog. 2014 Dec;10(12):e1004547.

30. Takaine M, Ueno M, Kitamura K, Imamura H, Yoshida S. Reliable imaging of ATP in living budding and fission yeast. J Cell Sci. 2019 Apr 17;132(8).

31. Takaine M, Imamura H, Yoshida S. High and stable ATP levels prevent aberrant intracellular protein aggregation in yeast. Elife. 2022 Apr 19;11.

32. Laussel C, Albanèse V, García-Rodríguez FJ, Ballin A, Defenouillère Q, Léon S. 2-deoxyglucose transiently inhibits yeast AMPK signaling and triggers glucose transporter endocytosis, potentiating the drug toxicity. PLoS Genet. 2022 Aug;18(8):e1010169.

33. Martinez-Ortiz C, Carrillo-Garmendia A, Correa-Romero BF, Canizal-García M, González-Hernández JC, Regalado-Gonzalez C, et al. SNF1 controls the glycolytic flux and mitochondrial respiration. Yeast. 2019 Aug;36(8):487–94.

34. Müller H, Lesur A, Dittmar G, Gentzel M, Kettner K. Proteomic consequences of TDA1 deficiency in Saccharomyces cerevisiae: Protein kinase Tda1 is essential for Hxk1 and Hxk2 serine 15 phosphorylation. Sci Rep. 2022 Oct 27;12(1):18084.

35. Lubitz T, Welkenhuysen N, Shashkova S, Bendrioua L, Hohmann S, Klipp E, et al. Network reconstruction and validation of the Snf1/AMPK pathway in baker's yeast based on a comprehensive literature review. NPJ Syst Biol Appl. 2015;1:15007.

36. Zeinoun B, Teixeira MT, Barascu A. Hog1 acts in a Mec1-independent manner to counteract oxidative stress following telomerase inactivation in Saccharomyces cerevisiae. Commun Biol. 2024 Jun 22;7(1):761.

37. Pastor MM, Proft M, Pascual-Ahuir A. Mitochondrial function is an inducible determinant of osmotic stress adaptation in yeast. J Biol Chem. 2009 Oct 30;284(44):30307–17.

38. Zhang S, Wang H, Sipko EL, Li S, Daugird TA, Legant WR, et al. Shared and redundant proteins coordinate signal cross-talk between MAPK pathways in yeast. Mol Biol Cell. 2024 Oct 1;35(10):ar126.

39. Woo DK, Poyton RO. The absence of a mitochondrial genome in rho0 yeast cells extends lifespan independently of retrograde regulation. Exp Gerontol. 2009;44(6–7):390–7.

40. Kaya A, Gerashchenko M V, Seim I, Labarre J, Toledano MB, Gladyshev VN. Adaptive aneuploidy protects against thiol peroxidase deficiency by increasing respiration via key

mitochondrial proteins. Proc Natl Acad Sci U S A. 2015 Mar;112(34):10685–90.

41. Mao K, Klionsky DJ. MAPKs regulate mitophagy in Saccharomyces cerevisiae. Autophagy. 2011 Dec;7(12):1564–5.

42. Li Y, Jiang Y, Paxman J, O'Laughlin R, Klepin S, Zhu Y, et al. A programmable fate decision landscape underlies single-cell aging in yeast. Science. 2020 Jul 17;369(6501):325–9.

43. Li Y, Jin M, O'Laughlin R, Bittihn P, Tsimring LS, Pillus L, et al. Multigenerational silencing dynamics control cell aging. Proc Natl Acad Sci U S A. 2017 Oct 17;114(42):11253–8.

44. O'Laughlin R, Jin M, Li Y, Pillus L, Tsimring LS, Hasty J, et al. Advances in quantitative biology methods for studying replicative aging in Saccharomyces cerevisiae. Transl Med Aging. 2020;4:151–60.

45. O'Laughlin R, Forrest E, Hasty J, Hao N. Fabrication of Microfluidic Devices for Continuously Monitoring Yeast Aging. Bio Protoc. 2023;13(15).

46. Zhou Z, Liu Y, Feng Y, Klepin S, Tsimring LS, Pillus L, et al. Engineering longevity-design of a synthetic gene oscillator to slow cellular aging. Science. 2023 Mar;380(6643):376–81.

47. Jin M, Li Y, O'Laughlin R, Bittihn P, Pillus L, Tsimring LS, et al. Divergent Aging of Isogenic Yeast Cells Revealed through Single-Cell Phenotypic Dynamics. Cell Syst. 2019 Mar;8(3):242-253.e3.

48. Liu Y, Zhou Z, Su H, Wu S, Ni G, Zhang A, et al. Enhanced cellular longevity arising from environmental fluctuations. Cell Syst. 2024 Aug;15(8):738-752.e5.

49. Paxman J, Zhou Z, O'Laughlin R, Liu Y, Li Y, Tian W, et al. Age-dependent aggregation of ribosomal RNA-binding proteins links deterioration in chromatin stability with challenges to proteostasis. Elife. 2022 Oct 4;11.

50. Roubenoff R. Catabolism of aging: is it an inflammatory process? Curr Opin Clin Nutr Metab Care. 2003 May;6(3):295–9.

51. Terman A, Brunk UT. Aging as a catabolic malfunction. Int J Biochem Cell Biol. 2004 Dec;36(12):2365–75.

52. Gladyshev VN, Kritchevsky SB, Clarke SG, Cuervo AM, Fiehn O, de Magalhães JP, et al. Molecular Damage in Aging. Nat Aging. 2021 Dec;1(12):1096–106.

53. Kaeberlein M, Kirkland KT, Fields S, Kennedy BK. Genes determining yeast replicative life span in a long-lived genetic background. Mech Ageing Dev. 2005 Apr;126(4):491–504.

54. Parkhitko AA, Ramesh D, Wang L, Leshchiner D, Filine E, Binari R, et al. Downregulation of the tyrosine degradation pathway extends Drosophila lifespan. Elife. 2020 Dec 15;9.

55. Parkhitko AA, Filine E, Mohr SE, Moskalev A, Perrimon N. Targeting metabolic pathways for extension of lifespan and healthspan across multiple species. Ageing Res Rev. 2020 Dec;64:101188.

56. Kondoh H, Lleonart ME, Gil J, Wang J, Degan P, Peters G, et al. Glycolytic enzymes can modulate cellular life span. Cancer Res. 2005 Jan 1;65(1):177–85.

57. Chistiakov DA, Sobenin IA, Revin V V, Orekhov AN, Bobryshev Y V. Mitochondrial aging and age-related dysfunction of mitochondria. Biomed Res Int. 2014;2014:238463.

58. Lanza IR, Befroy DE, Kent-Braun JA. Age-related changes in ATP-producing pathways in human skeletal muscle in vivo. J Appl Physiol (1985). 2005 Nov;99(5):1736–44.

59. Chaudhari SN, Kipreos ET. The Energy Maintenance Theory of Aging: Maintaining Energy Metabolism to Allow Longevity. Bioessays. 2018 Aug;40(8):e1800005.

60. Parkhitko AA, Cracan V. Xenotopic synthetic biology: Prospective tools for delaying aging and age-related diseases. Sci Adv. 2025 Mar 28;11(13):eadu1710.

61. Buschlen S, Amillet JM, Guiard B, Fournier A, Marcireau C, Bolotin-Fukuhara M. The S. Cerevisiae HAP complex, a key regulator of mitochondrial function, coordinates nuclear and mitochondrial gene expression. Comp Funct Genomics. 2003;4(1):37–46.

62. Huang Z, Xie N, Illes P, Di Virgilio F, Ulrich H, Semyanov A, et al. From purines to purinergic signalling: molecular functions and human diseases. Signal Transduct Target Ther. 2021 Apr

28;6(1):162.

63. Ai Y, Wang H, Liu L, Qi Y, Tang S, Tang J, et al. Purine and purinergic receptors in health and disease. MedComm (Beijing). 2023 Oct;4(5):e359.

64. Bermejo C, Rodríguez E, García R, Rodríguez-Peña JM, Rodríguez de la Concepción ML, Rivas C, et al. The sequential activation of the yeast HOG and SLT2 pathways is required for cell survival to cell wall stress. Mol Biol Cell. 2008 Mar;19(3):1113–24.

65. Gustin MC, Albertyn J, Alexander M, Davenport K. MAP kinase pathways in the yeast Saccharomyces cerevisiae. Microbiol Mol Biol Rev. 1998 Dec;62(4):1264–300.

66. Johnson AN, Li G, Jashnsaz H, Thiemicke A, Kesler BK, Rogers DC, et al. A rate threshold mechanism regulates MAPK stress signaling and survival. Proc Natl Acad Sci U S A. 2021 Jan 12;118(2).

67. Schaeffer HJ, Weber MJ. Mitogen-activated protein kinases: specific messages from ubiquitous messengers. Mol Cell Biol. 1999 Apr;19(4):2435–44.

68. Sabbagh W, Flatauer LJ, Bardwell AJ, Bardwell L. Specificity of MAP kinase signaling in yeast differentiation involves transient versus sustained MAPK activation. Mol Cell. 2001 Sep;8(3):683–91.

69. Sarg NH, Zaher DM, Abu Jayab NN, Mostafa SH, Ismail HH, Omar HA. The interplay of p38 MAPK signaling and mitochondrial metabolism, a dynamic target in cancer and pathological contexts. Biochem Pharmacol. 2024 Jul;225:116307.

70. Mishra P, Chan DC. Metabolic regulation of mitochondrial dynamics. J Cell Biol. 2016 Feb 15;212(4):379–87.

71. Gehart H, Kumpf S, Ittner A, Ricci R. MAPK signalling in cellular metabolism: stress or wellness? EMBO Rep. 2010 Nov;11(11):834–40.

72. Ocampo A, Liu J, Schroeder EA, Shadel GS, Barrientos A. Mitochondrial respiratory thresholds regulate yeast chronological life span and its extension by caloric restriction. Cell Metab. 2012 Jul 3;16(1):55–67.

73. Choi KM, Hong SJ, van Deursen JM, Kim S, Kim KH, Lee CK. Caloric Restriction and Rapamycin Differentially Alter Energy Metabolism in Yeast. J Gerontol A Biol Sci Med Sci. 2017 Dec 12;73(1):29–38.

74. Kaeberlein M, Hu D, Kerr EO, Tsuchiya M, Westman EA, Dang N, et al. Increased life span due to calorie restriction in respiratory-deficient yeast. PLoS Genet. 2005 Nov;1(5):e69.

75. Cohen R, de la Rocque F, Boucherat M, Bouhanna A, Lecompte MD, Brami A, et al. Prevention of acute otitis media. Amoxicillin versus glycoproteins from Klebsiella pneumoniae. Study in children under 5 years of age. Presse Med. 1992 Mar 21;21(11):509–14.

76. Li Y, Jiang Y, Paxman J, O'Laughlin R, Klepin S, Zhu Y, et al. A programmable fate decision landscape underlies single-cell aging in yeast. Science. 2020 Mar;369(6501):325–9.

77. Li Y, Jin M, O'Laughlin R, Bittihn P, Tsimring LS, Pillus L, et al. Multigenerational silencing dynamics control cell aging. Proc Natl Acad Sci U S A. 2017 Mar;114(42):11253–8.

Figures:

Figure 1: Strain design: (**A**) Pleckstrin homology (PH) domain of *SLM1* gene fused with *NTT1* and inserted in the vector (*pAG306GPD-ccdB-chr1*). The constructed (*PH-NTT1*) plasmid, digested with *NotI*, was integrated into the yeast genome. (**B**) In a separate construct, *yeGFP* sequence was fused to the 3' end of *PH-NTT1* sequence and expressed in yeast and visualized by confocal microscopy to assess cellular localization. (**C**) Overall yeast strain design with *NTT1* overexpression mediating ATP import from extracellular environment.

Figure 2: Measurement of relative ATP levels (**A**) Yeast cells, expressing fluorescent QUEEN ATP reporters were imaged using fluorescence microscopy. Relative ATP levels were measured in cells grown in media containing (**B**) 2% glucose or (**C**) 20 mM 2DG and are presented as violin plots. Sample sizes for each strain are between 265 and 315 cells. (**D**) LC-MS/MS measurement of intracellular ATP changes in Wt and *NTT1*-expressing cells. A pairwise t-test to compare each cell to their respective control was used to compare the means of two dependent groups and determine if a statistically significant difference exists between them. Statistical significance between groups and raw QUEEN ratio as well as LC-MS/MS data are available in **Supplementary File 1.**

Figure 3: Transcriptomic analysis of Wt and Wt_Ntt1 cells with and without ATP treatment. (**A**) Heatmap showing hierarchical clustering of gene expression profiles from Wt and Wt_Ntt1 cells cultured with ATP (red) and without ATP (green). Expression values are Z-score normalized, with red indicating higher expression and blue indicating lower expression. Three independent cultures for each group were used to isolate total RNA and perform RNA-seq analysis. (**B, C**) Venn diagrams displaying the number of genes uniquely or commonly (**B**) upregulated or (**C**) downregulated in Wt and Wt_Ntt1 cells treated with ATP compared to untreated controls. (**D, E**) Enrichment analysis showing significantly enriched gene ontology (GO) terms associated with genes (**D**) upregulated and (**E**) downregulated in Wt_Ntt1 cells compared to their corresponding untreated controls. Circle size reflects the number of genes in each GO term, and color indicates significance level as represented by -log10(FDR). The normalized counts and the complete list of DEGs with *p* values can be found in **Supplementary File S2**.

Figure 4: ATP treatment represses Hog1 activation through a mitochondria-dependent mechanism. The effect of 5 mM ATP treatment on the decreased Hog1[P] level was lost in *NTT1* rho[0] cells. Both phosphorylated and non-phosphorylated protein abundance of Hog1 was assessed with western blot analyses using specific antibodies raised against each form. Pgk1 was used as an internal loading control. Uncropped membranes are provided as Fig S5.

Figure 5: Quantifications of ATP levels and replicative lifespans of the Wt strain and *NTT1* strain under ATP treatment condition. (**A**) Heatmaps show the fluorescent signal of the QUEEN intracellular ATP reporter throughout cells whole lifespan. Each line is a color-coded (Red=low ATP and Blue = high ATP) time trace of a single yeast cell. Wt control without addition of external ATP showed rapid increase of cellular ATP during the aging process (left), while Wt control with ATP showed much higher level (p = 1.15E-143) and similar trend of cellular ATP (right). (**B**) *NTT1* strain without ATP shows lower level of cellular ATP compared to WT without ATP throughout the aging process whereas addition of ATP restored the cellular ATP level to that similar to Wt control with addition of ATP (right). (**C**) Average of single-cell time traces of the QUEEN reporter signal aligned by percent lifespan. Error bars are standard errors. (**D**) Addition of ATP significantly increased the replicative lifespan of the Wt strain (p = 1.28E-

12) and the *NTT1* strain (p = 4.03E-18). Without ATP, the *NTT1* strain showed significantly shorter lifespan than the Wt strain (p = 0.005), while with ATP, the 2 strains showed similar lifespans (p = 0.669). With addition of ATP, the *NTT1* strain showed a significantly enhanced replicative lifespan as compared to Wt without ATP (p = 1.03E-18). P-values were calculated with Mann-Whitney U test. The AU units and lifespan data can be found in **Supplementary File 3.**

Figure 6: Replicative lifespan of different aging modes and mode distribution under ATP treatment condition. (**A**) Without ATP, the *NTT1* strain showed similar lifespans between the two modes of yeast cell aging (p = 0.74), while Wt strain showed different lifespans (p = 2.18E-4) between 2 modes of aging. While the mode-2 lifespans between the 2 strains showed no significant difference (p = 0.826), *NTT1* strain showed significantly shorter mode-1 lifespan than that of the Wt strain (p = 2.39E-4). (**B**) With ATP, the *NTT1* strain showed significantly increased mode-1 lifespan (p = 1.24E-16), but not mode-2 lifespan (p = 0.15), whereas the Wt strain showed significantly increased mode 1 lifespan (p = 3.17E-7) and also mode-2 lifespan (p = 5.98E-6), which was similar to mode 1 lifespan (p = 0.4). The mode 1 lifespans between the 2 strains were similar (p = 0.849). (**C**) Both strains show increased percentage of mode-1 cells under the effects of ATP. The p-values were calculated using Mann-Whitney U test. The lifespan data can be found in **Supplementary File 3.**

Figure 7: Replicative Lifespan analyses of *rho⁰* Isolates. Replicative lifespan (RLS) phenotypes of (**A**) parental controls and (**B**) their *rho⁰* isolates were measured using a microfluidic chip under untreated and ATP-treated (w/ATP) conditions. Elimination of mtDNA in rho0 isolates alleviate the toxic effect of ATP treatment on Wt_Ntt1 cells. Raw lifespan data and statistical significance are provided in **Supplementary File 3**.

Figure 8: Proposed Models for ATP Treatment Effects in Wt and Wt_Ntt1 Cells. The type-1 effect involves ATP sensing, analogous to mammalian purinergic signaling, which activates a kinase cascade (e.g., MAPK) to modulate downstream pathways and mitochondrial function (**left panel**). In contrast, the type-2 effect is specific to Wt_Ntt1 cells and is regulated through increased ATP abundance imported from the medium via Ntt1. In this model, elevated ATP levels inhibit mitochondrial respiration and catabolic processes, potentially modulating some pathways associated with the type-1 effect (**right panel**). Both type-1 and type-2 effects increase replicative lifespan (RLS), with the type-2 effect producing a stronger lifespan extension.

Figure S1: Gene Ontology and KEGG Pathway Analyses for Wt Cells. Gene ontology (GO) enrichment analysis for (**A**) upregulated and (**B**) downregulated genes, and KEGG pathway analysis for (**C**) upregulated and (**D**) downregulated genes in Wt cells after ATP treatment. GO categories include BP (Biological Process), CC (Cellular Component), and MF (Molecular Function).

Figure S2: Alteration of MAP Kinase Pathway in Wt Cells. Genes uniquely upregulated (green) or downregulated (red) are shown for each component of the MAPK pathway in Wt cells under ATP treatment: (A) Cell Wall Stress, (B) High Osmolarity, (C) Pheromone, and (D) Starvation.

Figure S3: Gene Ontology and KEGG Pathway Analyses for Wt_Ntt1 Cells. Gene ontology (GO) enrichment analysis for (**A**) upregulated and (**B**) downregulated genes, and KEGG pathway analysis for (**C**) upregulated and (**D**) downregulated genes in Wt_Ntt1 cells after ATP treatment compared to untreated controls. GO categories include BP (Biological Process), CC (Cellular Component), and MF (Molecular Function).

Figure S4: SNF1-Mediated Regulatory Gene Network in Wt_Ntt1 Cells. Genes regulated by SNF1 in Wt_Ntt1 cells under ATP treatment are depicted in red (downregulated), green (upregulated), and black (no expression changes). The types of interactions are represented by arrows and lines.

Figure S5: Uncropped western blot membranes for assessing the abundance of active Hog1 protein.