Comparative gene annotation and orthology assignments across 301 species of Drosophilidae

Pankaj Dhakad[1,2]*, Bernard Kim[3], Dmitri Petrov[4,5], Darren J. Obbard[1]

[1]Institute of Ecology and Evolution, University of Edinburgh, Edinburgh, United Kingdom

[2]Evolution and Ecology, University of California, Davis, United States of America

[3]Department of Ecology and Evolutionary Biology, Princeton University, Princeton, United States of America

[4]Department of Biology, Stanford University, Stanford, California, United States of America

[5]CZ Biohub, Investigator, San Francisco, California, United States of America

* Corresponding author
Email: Pankaj.Dhakad@ed.ac.uk

Abstract

High-quality genome annotations are essential if we are to address central questions in comparative genomics, such as the origin of new genes, the drivers of genome size variation, and the evolutionary forces shaping gene content and structure. Here, we present protein-coding gene annotations for 301 species of the family Drosophilidae, generated using the Comparative Annotation Toolkit (CAT) and BRAKER3, and incorporating available RNA-seq and protein evidence. We take a comparative phylogenetic approach to annotation, with the aim of improving consistency and accuracy, and to generate a robust set of gene annotations and orthology assignments. We analyze our annotations using a phylogenetic mixed-model approach and find that gene number and CDS length exhibit moderate phylogenetic heritability (40% and 9.7%, respectively). For comparison, we also present analyses using a subset of the 215 highest quality genomes, although the findings were not markedly different. Our work suggests that while evolutionary history contributes to variation in these traits, species-specific factors—including assembly error—play a substantial role in shaping observed differences. To illustrate the utility of our annotations for comparative analyses, we investigate codon usage bias and amino acid composition across Drosophilidae. We find that codon usage is correlated with overall GC content and evolves slowly, but that it is also strongly shaped by selection—such that, in general, species with the strongest selection on synonymous codon usage show the lowest GC bias in third codon positions. This comparative annotation dataset forms part of an on-going collaborative project to sequence and annotate all species of Drosophilidae, with data and annotations being made rapidly and freely available on an on-going basis. We hope that this effort will serve as a foundation for studies in evolutionary and functional genomics and comparative biology across Drosophilidae.

Introduction

Fundamental questions in comparative genomics include the origin of new genes, the causes of genome size variation, and the factors that shape rates of genome evolution. However, addressing these questions requires not just the genome sequence, but also accurate identification and characterization of genomic features such as coding DNA sequence. High-quality genome annotations are thus essential for the identification of loci that underpin phenomena such as adaptation and speciation [1]. However, the annotation of genomes remains challenging, due to complex gene structures, long non-coding regions, and species-specific features [1-3]. Automated annotation methods, while efficient, often produce artefacts that can mislead interpretations of gene function and evolutionary relationships [4-6]. For example, an early annotation of the *Daphnia pulex* genome may have over-estimated the number of genes and over-predicted paralogs of genes involved in environmental responsiveness, potentially leading to initial misinterpretations of the basis of its adaptive capabilities [7, 8].

Over the past decade, advances in long-read sequencing technologies and scaffolding methods, together with the declining cost of sequencing, have led to a dramatic increase in the number and quality of genome assemblies across the tree of life [9, 10]. This surge is

exemplified by large-scale initiatives such as the Darwin Tree of Life project [11], which aims to
sequence around 70,000 species in the UK and Ireland, the Vertebrate Genome Project (VGP)
[12], which has the goal of sequencing all extant vertebrate species, and the African
BioGenome Project [13], which seeks to sequence more than 105 thousand species in Africa.
However, despite advances in sequencing and assembly, genome annotation remains a major
bottleneck [14]. Most genomes from large-scale sequencing projects are annotated
independently using automated pipelines such as BRAKER, Augustus, and MAKER [15-17].
While these annotations can incorporate evidence from reference protein databases (e.g.,
OrthoDB), they typically do not exploit whole-genome alignments across multiple closely
related species or conserved gene order (synteny) to improve annotation accuracy and
consistency across genomes [18, 19]. As a result, different pipelines can identify slightly
different sets of genes due to variation in prediction algorithms, parameter settings, and
assumptions about gene structure [20]. This inconsistency means that genes missed by one
pipeline might be detected by another, while certain gene models may only partially align
between annotations [14]. Consequently, such independent annotations make it difficult to
achieve a standardized gene set for comparative analyses [7, 8, 20]. Comparative annotation,
on the other hand, aims to address these issues by using alignment with well-annotated
reference genomes to help guide predictions, reducing discrepancies and aligning gene
models more consistently across closely related species [21, 22]. This approach can not only
increase the accuracy of gene predictions, but can also ensure a more robust, comparable
gene set for evolutionary studies. The Comparative Annotation Toolkit (CAT) is one such
method, designed to annotate genomes by projecting known gene models from a reference
genome onto target genomes within a phylogenetic framework [22]. CAT integrates evidence
from such a 'lift-over' with short and long read RNA-seq data, Iso-seq data, and protein
alignments to refine gene model predictions, weighting features that are shared among close
relatives more heavily.
91
For over a century, *Drosophila melanogaster* and its relatives have been at the forefront of
genetics, genomics, and evolutionary research, leading to influential discoveries that have
shaped these fields [23]. High-quality genome assemblies and annotations have been
developed for several key species, beginning with the pioneering sequencing of the *Drosophila*
melanogaster genome, which served as a reference for subsequent genomic studies [24, 25].
The *Drosophila* 12 Genomes Project expanded this foundation, offering comparative insights
across multiple species, while initiatives such as the ModENCODE project further enriched our
knowledge with detailed transcriptomic and epigenomic data [26, 27]. Individual research
groups have continued to sequence target species [28-30, 31, such as *D. miranda*, *D. guanche*
etc], making possible resources such as DrosOMA (https://drosoma.dcsr.unil.ch/), which
provides genus-wide orthology information for 36 Drosophila species [32]. This progress has
now culminated in the on-going community effort to achieve a comprehensive genomic study
of the entire family Drosophilidae [33, 34]. This effort includes the *de novo* sequencing of new
species, scaffolding and improvement of existing genomes, and the generation of new
transcriptomic data [33]. As of April 2024, around 360 different drosophilid species had been
sequenced to varying levels of completeness—some fragmentary, from short-read data alone
[e.g. *Drosophila setifemur* and *Drosophila ironensis*; 31], but many to chromosome-level

assemblies, using long-read data and/or scaffolding information from HiC [e.g. *Chmomyza*
fuscimana; 35].
111
Here we contribute to this continuing community effort by providing a comparative coding-
sequence annotation for 301 drosophilid genomes. We do this using a combination of CAT and
BRAKER3, combining publicly available RNAseq data and previous RefSeq reference
genomes [15, 22]. We use phylogenetic linear mixed models to assess and compare the
annotations, and as an example of the utility of our comprehensive annotation we analyze
codon and amino-acid usage bias across the family. To facilitate its use in both single-gene
and genome-wide studies, the annotation is made freely available in the form of genome
annotation files and also as aligned (and optionally masked) orthology groups, with
annotations linked to gene orthology. This is available as two complementary datasets; the first
is complete set of 301 species selected on the basis of a more permissive quality filter, most
suitable for single-gene studies of sequence evolution (e.g., identifying conserved regions,
protein structure prediction, or site-level evolutionary inference), the second is a more
stringently-filtered subset of 215 species, which may be more robust for studying micro-
synteny, gene family evolution or copy-number variation. In the future, we plan to continue
updating this resource with regular new releases as new genomes become available. We hope
that this will be a key resource, enabling gene-based analyses of evolution within this
important model system.

Methods

Genome assemblies

We selected an initial candidate set of genome assemblies by supplementing those of Kim et
al [33] with all other publicly available drosophilid genomes available as of February 2024. This
included genomes available in the RefSeq database (Release 222) [36], those generated by
the Darwin Tree of Life project [11], and many assemblies generated by individual labs [37-40].
For our primary dataset, we shortlisted 301 genome assemblies that had a scaffold N50
greater than 50 Kbp and a BUSCO (Benchmarking Universal Single-Copy Orthologs)
completeness score of over 90% [41], selecting the assembly with the highest scaffold N50
where multiple assemblies were available. To identify and mask repetitive elements, we used
RepeatMasker v4.1.2 [42] and Dfam release 3.7 repeat library [43]. These soft-masked
genomes were used for all subsequent analyses. To provide a secondary, higher-stringency,
dataset for comparison we selected a subset of those genomes which had BUSCO
completeness ≥97%, contig N50 ≥2 Mbp, and assemblies generated using long-read
sequencing technologies (Oxford Nanopore or PacBio).

RNA-seq and protein data

To annotate the genomes, we used the Comparative Annotation Toolkit [CAT; 22], a pipeline
that leverages external evidence ('hints') combining data such as RNA-seq, Iso-seq, proteins,

and attempted lift-over from aligned references. For each of the 301 species, we first gathered
available RNA-seq data to provide transcript evidence that can help resolve ambiguities in
gene models. We identified suitable RNA-seq datasets using the ENA Portal API
(https://www.ebi.ac.uk/ena/portal/api), selecting up to 10 paired-end RNA-seq datasets and
prioritizing those with Poly-A selection to enrich mRNA [44]. Where available, we included data
from up to 10 tissue types, including whole body, carcass, thorax, brain, testes and ovaries, as
well as different developmental stages and both sexes (selected SRA numbers for each
species are in S1 Table). The chosen RNA-seq reads were then down-sampled and
normalized to 100x coverage using BBNorm of BBMap v38.95 [45]. The normalized RNA-seq
reads were aligned to their respective species genomes using the STAR v2.7.9a with default
parameters [46].
158
To provide protein 'hints', we extracted predicted protein sequences from Arthropoda using the
OrthoDB v10 protein database [https://orthodb.org; 47]. Such protein sequences provide
evolutionary conserved evidence that complements RNA-seq data, particularly for genes that
may be underrepresented or absent in the RNA-seq datasets. We aligned these protein
sequences to the genomes using miniprot v0.12 [48] with parameters "-ut8 --gtf genome_file",
which are optimized for mapping proteins to genomic sequences. The alignment files
generated by miniprot were then converted into hints files using the aln2hints.pl script from the
GALBA toolkit [49].

Reference species and cactus alignment

In addition to RNAseq and protein hints, the CAT pipeline attempts a lift-over of annotations
[22]. This uses genome-scale alignments in the Hierarchical Alignment Format [50], each
comprising a single reference species and several target species. To define reference clades
for annotation, we first generated a preliminary species tree including the 301 drosophilid
species (plus seven outgroup species) using 1824 single-copy BUSCO loci [41]. Nucleotide
sequences from each locus were aligned separately using MAFFT v7.520 [51] and used to
infer a maximum likelihood (ML) gene tree using IQ-TREE v2.2.6 [52] under a GTR+I+G4
substitution model. These ML gene trees were then combined to infer a species tree using
ASTRAL-III v5.15.5 [53], which aims to resolve gene-tree species tree incongruence under a
model of incomplete lineage sorting.
178
We selected 37 'reference' species for lift-over annotations based on the completeness and
quality of their genomes, as indicated by RefSeq annotations [36]. Using the ETE 3 python
package [54], we applied a preorder tree traversal strategy to identify subclades that contained
at least one reference species and included the most distantly related leaf within a predefined
phylogenetic distance (measured as the expected number of substitutions per site). We varied
the phylogenetic distance threshold between 0.005 and 0.35 to ensure each subclade included
3 to 15 species, with lower thresholds for densely populated regions of Drosophilidae tree and
higher thresholds to include more distantly related species. In clades containing multiple
potential RefSeq-annotated species, we selected a single reference species based on the
similarity of its gene count to that of the best-annotated drosophilid model, *Drosophila*

melanogaster. This criterion was used because the *D. melanogaster* annotation is likely to be
the most complete and accurate within the family, having benefited from extensive manual
curation and extensive transcriptomic and functional data [55]. This process resulted in 17
RefSeq-annotated species for use in the attempted lift-over. For species located on very long
branches (i.e. divergence >0.35 from the closest potential reference) and for subclades lacking
any reference-species annotations, we attempted a lift-over directly from *Drosophila*
melanogaster.
196
197 We then used these 'lift-over subclades' as guide trees to generate multiple whole-genome
198 alignments with ProgressiveCactus [56]. This approach ensured that the alignments were
199 computationally feasible, and that closely-related genomes aligned together. Finally, we
200 employed the Comparative Annotation Toolkit [22] to annotate multiple target genomes
201 simultaneously, using a lift-over of the selected reference annotation for each subclade.

Running CAT

To perform the genome annotations, we first prepared the reference annotations and extrinsic
'hints' for use in CAT [22]. RefSeq annotation files were converted using the
"convert_ncbi_gff3" script provided by CAT, and the resulting GFF3 files were validated with
the "validate_gff3" script to ensure compatibility. We then employed three modes of
AUGUSTUS [16] in CAT: two based on transMap projections (AugustusTM/R) that project
annotations from reference genomes onto target genomes, and one using ab-initio and
comparative gene predictions (AugustusCGP) guided by extrinsic hints [57]. We used
Comparative Gene Prediction (CGP) parameters trained on 12 well annotated Drosophila
species from the Drosophila 12 Genomes Project, based on exon and intron scoring
[https://bioinf.uni-greifswald.de/augustus/datasets/; 57].

Complementation with BRAKER3

To complement the comparative annotations generated by CAT, and to reduce any potential
reference bias, we additionally incorporated *de novo* non-comparative CDS predictions made
by BRAKER3 [15]. BRAKER3 is an automated gene prediction pipeline that integrates RNA-
seq data with gene prediction algorithms to generate gene models without reference to a
reference annotation, improving the identification of novel genes and gene duplicates that may
be absent from the reference. However, among the annotations generated by BRAKER3 we
identified a number of transposable elements (TEs), which are not the focus of our study.
Therefore, to simplify the annotation, we removed these using EarlGrey v4.1.0 [58], applying
TEstrainer (https://github.com/jamesdgalbraith/TEstrainer) to ensure that recently duplicated
non-TE genes were retained. Finally, to combine the BRAKER3 annotations with those from
CAT we compared the coding sequences (CDS) of overlapping genes between the two
annotation sets. For one-to-one overlapping genes, we selected the annotation with the
longest CDS. In cases of one-to-many or many-to-one overlaps, we preferred the CDS
annotations from CAT, as these reflect the additional support provided by comparison among

relatives. Additionally, we retained all non-overlapping genes with a CDS length greater than
150 nucleotides.

Annotation quality assessment

To assess the annotation quality, we compared our CAT-BRAKER annotations and
annotations generated using BRAKER3 alone with RefSeq annotations. These comparisons
focused on coding sequence (CDS) concordance, quantified using overlap-based precision
and recall (>90% CDS overlap) and overall Jaccard similarity compared to RefSeq CDSs. In
addition to these direct annotation comparisons, we assessed completeness and gene
repertoire quality using BUSCO and OMArk, which evaluate annotations against evolutionary
expectations of conserved gene content [41, 59]. BUSCO was run in protein mode using the
Diptera OrthoDB dataset to estimate the proportion of conserved single-copy orthologs
recovered in each annotation [60]. For OMArk, we first generated omamer search databases
for each species using the LUCA.h5 orthology database
(https://omabrowser.org/oma/current/). We then ran OMArk with taxon ID 7214
(Drosophilidae), assigning proteins to orthologous groups based on phylogenetically informed
gene family classifications.

Orthogroup assignment and CDS alignment

We identified coding sequence homology across the 301 *Drosophila* species and one outgroup
species (*Musca domestica*) using OrthoFinder v2.5.5 [61]. OrthoFinder first identifies homology
using an all-vs-all blast similarity search and then clusters sequences using a Markov
clustering algorithm (MCL inflation parameter of 1.5). Subsequently, it can then identify
"Hierarchical Orthologous Groups" (HOGs) that comprise the genes descended from a
common ancestral gene at a specific taxonomic level—with HOGs defined for different clades
nested within each other along the species phylogeny. We extracted HOGs at the level of
Drosophilidae (i.e., sets of homologous sequences that have their MRCA at the base of
Drosophilidae, or more recently) and retained for further analysis those HOGs that included at
least two species and contained more than three sequences.
255
We aligned the sequences for each of the chosen HOGs using the aligners MACSE v2 [62]
and MAFFT v7.520 [51], an approach intended to minimize the impact of any frameshifts and
in-frame stop codon errors that might arise from sequencing, assembly, or annotation
problems, while maximizing codon-aligned sequence length. MACSE v2 incorporates several
steps to improve alignment quality, including a prefilter to remove long non-homologous
insertions that may result from incorrect annotations, such as intron inclusions or alternative
splicing. It then uses HMMCleaner to mask residues that appear misaligned and applies post-
processing filters to mask isolated codons and patchy sequences, removing sequences if more
than 80% of the residues are masked [63]. Finally, the alignments were trimmed at both ends
until a nucleotide position represented by at least 70% of the sequences was reached,
ensuring a high-quality alignment for downstream analyses of protein coding sequence. From
the 301 annotated drosophilid genomes, we generated 35,642 HOGs and 22,355 high-quality

alignments. In addition, we provide HOGs and corresponding alignments for a high-confidence
annotation subset comprising 215 species, defined by more stringent assembly and annotation
quality criteria (BUSCO completeness ≥97%, contig N50 ≥2 Mbp, estimated OMArk
contamination ≤5%, and assemblies generated using long-read sequencing technologies).
These HOGs form the basis of subsequent analyses of evolutionary relationships and
functional conservation of genes across family Drosophilidae, and all masked and unmasked
aligned sequences are made available at https://doi.org/10.5281/zenodo.15016917.

Phylogenetic generalized linear mixed model analyses

We used Generalized Linear Mixed Model (GLMM) analyses of gene number and CDS length
to identify clade-to-clade variation and outlier genomes across all 301 species, and separately
across 215 species high-confidence set. Such variation may result from true evolutionary
divergence, or from reference bias and the availability (or otherwise) of RNA-seq data, or from
systematic errors in assembly or annotation quality. The relative impact of such factors is
naturally addressed in a linear mixed model framework, treating species as a random effect
and phylogenetic distance from reference species, genome size, assembly contiguity (contig
N50), read-type (ONT/PacBio vs Illumina), and the availability of RNAseq as fixed-effect
predictors. Because related species exhibit correlated traits [leading to pseudo-replication, if
not accounted for; 64], and because the phylogenetic correlation among related species (e.g.
'phylogenetic inertia' or 'phylogenetic heritability') may be of direct interest—reflecting clade-to-
clade variation in gene content or the efficacy of selection—we employed a Phylogenetic
mixed model approach [65], implemented in the R package MCMCglmm [66]. This
incorporates phylogenetic relationships to model the covariance among species, while
evaluating the influence of fixed predictors such as phylogenetic distance from the lift-over
reference and RNA-seq availability.
292
To do this, we first generated a revised species-tree topology of the 301 drosophilid species
using 251 single-copy HOGs (those HOGs containing ≥ 300 species) employing IQTREE2 and
ASTRAL-III, as for BUSCO genes above. To infer relative branch lengths in approximate time,
we randomly selected 10,000 amino acid sites from the HOGs, and we used BEAST [67] to re-
infer branch lengths on the (fixed) ASTRAL tree topology under a LG+G+I model with 7
gamma categories and an uncorrelated relaxed log-normal clock [68]. For the tree prior, we
used birth-death process model [69], setting the fully-informative prior for the MRCA of the
subgenera *Drosophila* and *Sophophora* to 47 Mya [95% prior density 42-52 mya; 34], a
uniform step prior between 0 and 1 on the birth-death growth rate, and the remaining priors to
their default values. We ran the MCMC chain for 10^6 generations sampling every 1,000 steps,
and stationarity and mixing were assessed from visual inspection of the MCMC chain and
Effective Sample Size (ESS) in Tracer [70]. After discarding 10% of the sampled estimates as
burn-in, we report divergence times as median node height for each of the clades in the
summary tree.
307
To assess the factors influencing gene number and CDS length across drosophilid species, we
fitted a multivariate phylogenetic mixed model using MCMCglmm [66]. Our model included

gene number and mean CDS length as response variables, allowing us to analyze their (co-
)variation with respect to predictors such as status as a lift-over reference, distance from lift-
over reference, RNAseq availability, assembled genome size, assembly contig N50 and read-
type. This phylogenetic approach can help to disentangle the effects of biological and technical
factors on genome annotation metrics, while accounting for phylogenetic relatedness among
species. We inferred statistical 'significance' on the basis of 95% highest posterior density
(HPD) credibility intervals. The syntax for MCMCglmm models and priors is described in S1
File.

Evolution of GC, codon, and amino acid composition across Drosophilidae

Genome-wide GC content, codon usage, and amino-acid composition are shaped by a
combination of mutational bias and natural selection [71, 72]. To illustrate the utility of our
coding sequence annotations and alignments for large-scale evolutionary sequence analyses,
we examined the evolution of these coding-sequence traits across the 301 drosophilid species.
We used the R package 'cubar' [73] to calculate the overall GC content of coding sequences
('GC'), and GC content at third codon positions (GC3). Additionally, we estimated the whole-
genome GC content and the GC content of non-coding sequences using 'geecee' [74]. We
used the 'cusp' tool from EMBOSS [75] to calculate amino acid frequencies in each species,
and the nitrogen to carbon ratio (N/C) for protein sequences was calculated as the weighted
average of the N/C ratios of individual amino acids, with weights corresponding to the
proportion of each amino acid in the sequence. We used a PCA analysis of amino acid
frequencies to reduce the dimensionality.
332
We estimated the strength of selection on codon usage bias (S) using the approach of Dos
Reis and Wernisch [76], which compares codon frequencies in highly expressed versus
reference gene sets. To do this, we ranked *Drosophila melanogaster* genes according to their
overall expression level (S2 Table; expression data obtained from FlyBase: https://flybase.org/)
and analyzed the HOGs that contained these genes, assuming that the globally most highly-
expressed genes in *Drosophila melanogaster* are also highly expressed in other species. As
expected, these genes were dominated by those encoding ribosomal proteins, yolk proteins,
salivary gland secretions and elongation factors whose high expression is likely to be
conserved. HOGs were then ranked in order of *Drosophila melanogaster* expression level and
binned in 20 expression categories of *ca.* 600 genes in each category (S3 Table). Codon
usage bias (S) was then estimated as the log-odds ratio of optimal to non-optimal codon
frequencies for two-fold degenerate codons, where the preferred codon was identified from the
most highly expressed gene category [76, 77]. Note that, to distinguish selection from
mutational bias, this method assumes the reference and highly expressed gene sets have
similar mutational patterns. Finally, we obtained bootstrap confidence intervals for S by
resampling genes within expression categories.
349
350 As above, we used a multivariate Phylogenetic Generalized Linear Mixed Model (PGLMM)
351 implemented in MCMCglmm to assess the among-species phylogenetic (co)variance in GC3,

352 non-coding GC, estimated strength of selection on codon usage bias (*S*), observed
353 frequencies of amino acids and N/C ratio, while accounting for phylogenetic effects.

Data availability

355 All code used for data processing, generating figures, and statistical analyses is publicly
356 available from Zenodo (https://doi.org/10.5281/zenodo.18444670). The Drosophila genome
357 annotations and orthology assignments generated in this study are also publicly available via
358 Zenodo (https://doi.org/10.5281/zenodo.15016917).

Results & Discussion

Gene annotation of 301 species

We selected 17 NCBI RefSeq annotations for use as potential lift-over references and
combined this information with RNAseq data from 91 species, protein hints (from mapping of
OrthoDB proteins) from all species, and *de novo* prediction to annotate the remaining genomes
using CAT and Braker3. On average, we identified 14,549 genes in each genome, with a mean
CDS length of 1.60 Kbp. This is very similar to the gold-standard reference, *Drosophila*
melanogaster, which currently has 13,904 protein-coding genes of mean CDS length 1.54 Kbp
(Genome assembly Release 6.53; GCF_000001215.4). However, there was substantial
variation, reflecting both evolutionary variation among species and potential variation in
genome assembly quality and RNAseq availability (Fig 1).
370
Fig 1. Overview of 301 Drosophila genome annotations.
(A) Time-calibrated phylogenetic tree showing ancestral reconstruction of protein-coding
gene number mapped onto branches. Tip labels are coloured by the reference species used
for comparative annotation within each clade; stars denote reference species, and filled
versus open circles indicate the presence or absence of RNA-seq data, respectively.
Concentric tile layers summarise key assembly and annotation metrics for each species:
from the innermost to the outermost ring, mean CDS length (Kbp), BUSCO completeness
(%), genome assembly size (Mbp), and contig N50. Major drosophilid species groups are
indicated by arcs positioned outside the outermost tile layer. (B) The box plot represents the
range in gene number and mean CDS length across family Drosophilidae. The numerical
data underlying both panels are provided in S4 Table. An alternative version of the tree,
with taxon labels, is provided in S2 File.
383
Most species (274 of 301) were annotated with between 12,500 and 16,000 protein-coding
genes. There were three species that appear to possess more than 20,000 genes; *Drosophila*
vulcana [28], *Drosophila miranda* [78] and *Drosophila punjabiensis* [28] (Fig 1 and S4 Table).
In *D. miranda*, the elevated gene number is consistent with the recent evolutionary history of
its sex chromosomes, namely a very young neo-sex chromosome system formed by fusion of
Muller element C with the ancestral Y chromosome that resulted in homologous gene copies

retained on both the neo-X and neo-Y. In our annotation, we identified 3,944 genes located on
the two largest neo-Y scaffolds and 2,944 genes on neo-X, consistent with extensive retention
of neo-Y homologs reported previously [78]. When these neo-sex homologs are considered,
the inferred gene number of *D. miranda* is closer to the genes in *D. melanogaster*, indicating
that it is not anomalous in terms of gene content. In contrast, the elevated gene counts
observed in *D. vulcana* and *D. punjabiensis* are more likely to reflect technical rather than
biological factors (see below). The mean CDS length across species ranged between 1.40 and
1.74 Kbp. The shortest CDSs were observed in *D. miranda*, consistent with extensive
fragmentation and degeneration of amplified genes on its neo-X and neo-Y chromosomes
reported previously [78]. Note that these latter two species are excluded from our subset of
215 high-confidence genomes.
401
Establishing whether the variation in gene number and mean CDS lengths reflects differences
in annotation or assembly quality, or true evolutionary processes, is necessarily challenging in
the absence of ground truth annotation for most species. We therefore assessed the quality of
our CAT-based annotations using multiple complementary approaches. First, we compared
CAT-BRAKER CDS annotations with available RefSeq annotations for a subset of non-*D.*
melanogaster species. Using CDS overlap (>90%) as a criterion, CAT-BRAKER consistently
showed higher precision relative to RefSeq than BRAKER3 alone, while recall was often
slightly lower (S1 Fig). However—except perhaps in the especially well-studied case of *D.*
melanogaster—RefSeq annotation should not necessarily be treated as a strict ground truth
[79], and it is notable that reduced recall primarily reflects additional gene models detected by
CAT and/or BRAKER that are absent from RefSeq. Consistent with this, the Jaccard similarity
between CAT-BRAKER and RefSeq annotations was generally higher than between
BRAKER3 and RefSeq, indicating greater overall concordance in gene content.
415
Second, we assessed BUSCO completeness at both the genome and annotated-protein
levels. For the majority of species, BUSCO completeness estimates were highly concordant
between genome-based and protein-based assessments, indicating that the annotations
recover most conserved genes detectable at the genome level (S2 Fig). Only *Drosophila*
guttifera and *Drosophila bifasciata* showed differences greater than 5% between genome and
protein BUSCO completeness.
422
Finally, we used OMArk to further assess and compare the quality of protein-coding gene
annotations. OMArk estimates the completeness, consistency, fragmentation and
contamination of gene-repertoire in a species by comparison with conserved orthologous
groups (HOGs). We observed high levels of HOG completeness across most species.
However, *Drosophila recens* and *Drosophila miranda* showed a high number of duplications
(S5 Table). In *D. miranda*, this pattern is explained by retention of homologous gene copies on
the neo-X and neo-Y chromosomes (above). The source of apparent duplication in *Drosophila*
recens remains unclear but is also observed in its close relatives *D. subquinaria* and *D.*
suboccidentalis. (S5 Table). Additionally, OMArk identified a substantial level of contamination
in the genome of *Drosophila vulcana* (5,235 of 23,519 genes identified as contaminant), a
genome that is also flagged as 'contaminated' in NCBI database [28]. Other genomes with

apparently high levels of contamination include *Drosophila punjabiensis* (3,131 of 20,339
genes), and *Drosophila nannoptera* (2,006 of 15454 genes). Importantly, all these genomes,
except *D. miranda*, showing higher duplication or contamination were excluded from the high-
quality subset. Together, these assessments indicate that most annotations are highly
complete at the gene level, while highlighting a small number of genomes where duplication or
contamination likely inflates apparent gene counts.

Orthology inference

We used OrthoFinder [61] to infer CDS orthology across 301 species of Drosophilidae, using
Musca domestica as an outgroup. OrthoFinder assigned 98.6% of predicted proteins to
orthogroups (OGs), with 96% further classified into Hierarchical Orthologous Groups (HOGs).
We identified a total of approximately 35 thousand HOGs across the 301 drosophila species
genomes (Table 1). More than 90% of genes in each species were assigned to HOGs,
although some species—such as *Leucophenga varia*, *Drosophila vulcana*, *Drosophila*
quasianomalipes, and *Drosophila differens*—had lower assignment rates (S3 Fig).
448
449   **Table 1. Summary statistics of HOGs in the full and high-stringency datasets.**

Feature	Full dataset	High stringency subset
Number of species	301	215
Total number of proteins	4379331	3074657
Number of Orthogroups (OGs)	29738	21946
Number of proteins in OGs	4316768 (98.6%)	3032610 (98.6%)
Number of HOGs	35642	26046
Number of proteins in HOGs	4202078 (96%)	2964679 (96.4%)
Number of universal* single-copy HOG	618	1077
Number of HOGs with all species present	1131	3406
Number of universal* HOGs	6276	7732
Number of HOGs containing Dmel genes	12151	12145
Number of ancient$ HOGs	12961	12539

This table summarizes the composition and coverage of orthologous gene groups inferred across
drosophilid genomes using the full dataset of 301 species and a more stringently filtered subset of
215 species. *Universal HOGs are defined as those containing genes from at least 99% of species,
while $ancient HOGs represent groups whose most recent common ancestor dates to ≥50 million
years ago and that include at least 30 species. Differences between datasets primarily reflect the
exclusion of lower-quality assemblies in the high-stringency subset, which increases the number of

universal and complete HOGs while retaining a high proportion of proteins assigned to orthologous
groups.

To interpret patterns of gene conservation and turnover, we classified HOGs into two broad
categories: widely conserved HOGs, which include genes shared across most Drosophilidae
species, and species- or clade-specific HOGs, which appear to be restricted in their
distribution.

The widely conserved HOGs were further divided into 'universal' HOGs, present in nearly all
species (≥99%), and ancient HOGs, shared by species with MRCA of at least 50 mya, but
missing in a substantial minority of species (Table 1; S4 Fig). Universal HOGs likely represent
genes experiencing strong evolutionary constraint, encoding core biological functions
necessary across all species. Ancient HOGs, while also conserved, may include genes that
have been differentially retained or lost in some major lineages. In contrast, species- and
clade-specific HOGs could represent recent gene family expansions or lineage-specific
adaptations. Where restricted HOGs encode functional proteins, they may contribute to
species-specific traits; however, they also likely arise from methodological artifacts, such as
genome annotation errors or orthology misassignment.

In the full dataset, over half of all the predicted protein-coding genes fell into universal HOGs,
reinforcing the idea that most genes are highly conserved. However, a substantial fraction of
the HOGs (~20,000) contained genes from a small number of species (<30), suggesting either
recent evolutionary gains or problems with orthology inference (S4 Fig). Interestingly, some
HOGs classified as ancient were present in only a few species, and it seems probable that
many of these 'sparse' HOGs reflect fragmented assemblies or annotation inconsistencies,
rather than true biological patterns. This highlights the importance of caution when interpreting
or analyzing low-representation HOGs in comparative studies. By classifying HOGs based on
their evolutionary conservation and phylogenetic distribution, we provide a framework for future
studies of gene conservation and turnover in Drosophilidae. Universal and ancient HOGs likely
represent functionally essential genes, while restricted HOGs may point to lineage-specific
innovations or methodological challenges. This classification allows us to distinguish between
broad evolutionary patterns and potential technical noise, improving the reliability of our
comparative analyses.

To obtain a well-supported set of orthogroups, and to mitigate potential errors from sparse
HOGs, we examined in more detail those HOGs that contained at least one *Drosophila*
melanogaster gene. Given that *Drosophila melanogaster* has one of the most complete and
thoroughly verified gene sets among multicellular eukaryotes, its representation in a HOG
provides additional confidence that the group represents a biologically meaningful gene family
rather than an artifact. We identified 12,151 such HOGs, which were broadly shared across the
majority of Drosophilidae species (S4 Fig). This approach provides greater confidence for the
analysis of conserved gene families, and allows us to assess how widely well-characterized
genes are distributed across phylogeny. This work substantially expands the most recently-

available gene-orthology set for Drosophilidae from 36 species [32] to 301 species, offering a
more comprehensive understanding of gene families across the entire clade, and we hope that
this dataset will be valuable for future studies focusing on comparative genomics, evolutionary
biology, and functional genomics within Drosophilidae.

Phylogenetic inference using BUSCO and HOG genes

Comparative analyses generally use an ultrametric phylogenetic tree describing relationships,
and thus the expected covariance in traits, among species [65], as this allows the inferences of
'phylogenetic heritability'. The most comprehensive molecular phylogeny of Drosophilidae to
date encompasses 704 species, but this is based on only 17 reference genes and thus has
many deep branches that are not resolved with high confidence [80]. More recently, genome-
sequencing of 360 species has enabled a BUSCO-gene tree based on one thousand loci—
greatly improving confidence in deeper relationships [33]. However, branch lengths were
inferred for that tree using 4-fold degenerate sites, which will tend to underestimate deep
branch lengths due to substitution saturation. Here we infer the species tree using both
BUSCO and HOG approaches, the first dataset comprising 251 single-copy HOGs, and the
second comprising 1,824 BUSCO genes. We found that the HOG tree and the BUSCO tree
were highly concordant and showed highly supported relationships for all but five species (S4
Fig). For example, in the HOG tree *Drosophila fuyamai* was positioned as a sister to
Drosophila carrolli, *Drosophila rhopaloa*, and *Drosophila prolongata*, whereas the BUSCO tree
also included *Drosophila kurseongensis* in this clade. Other conflicting relationships can be
found in the S3 File. Most internal branches were well supported in both trees, but in some
places, the HOG tree exhibited slightly lower local posterior probabilities, particularly for short
branches (S4 File). These discrepancies likely reflect increased discordance due to incomplete
lineage sorting [34], as resolving such branches requires a larger number of gene trees.

Factors affecting annotated gene number and CDS Length

525 To better understand apparent variation in gene number and CDS length across
526 Drosophilidae, we fitted a phylogenetic generalized linear mixed model [65] to assess the
527 impact of 'reference' genome status, phylogenetic distance from the reference, genome size,
528 assembly quality (Contig N50), read-type (i.e. long-read versus exclusively short-read), and the
529 availability of RNA-seq data on these two traits.
530
Our analysis found no significant differences in gene number or CDS length between our
annotations and the established reference annotations, indicating that our annotations are of
comparable quality (gene number: p=0.8, 95% HPD CI [-487, 494]; CDS length: p=0.67, 95%
HPD CI [-0.05, 0.004]). However, species lost an average of *ca.*16 genes for each extra million
years of divergence from their liftover reference (p<0.001, CI [-23.2, -7.7]), while on average
CDS length increased by just one nucleotide per million years divergence from reference
(p=0.002; CI [0.2, 1.5]; Fig 2; S1 File). This reflects the increased challenges associated with
lift-over between more divergent genomes, but—likely because lift-over was only one source of
data among many—the effect is relatively small. Repeating the analyses on the high-

stringency subset of 215 genomes produced qualitatively identical results, with only minor
changes in effect size (15 fewer genes and one nucleotide difference in mean CDS length; S1
File).
543
Fig 2. Variation in gene number and CDS length across Drosophilidae.
(A) Coding sequence (CDS) length increases with phylogenetic distance from the reference
species used for lift-over annotation. (B) CDS length decreases as genome size increases.
(C) Gene number remains largely unaffected by phylogenetic distance from the reference.
(D) Gene number increases with genome size, suggesting a relationship between genome
expansion and gene content. In all plots the points are colored by genome contiguity (contig
N50). The numerical data underlying all panels are provided in S4 Table. Fitted lines and
95% confidence windows are derived from a non-phylogenetic linear model and are for
illustration only; see main text for phylogenetic mixed model analyses.
553
In the full 301 species dataset, 'Read type' (i.e. long-read versus exclusively short-read)
emerged as the strongest overall predictor of gene content. Assemblies based on short-read
data alone contained *ca.* 1,000 more annotated genes than long-read-based assemblies (p <
0.001; 95% HPD CI [588, 1,460]), but exhibited shorter CDSs by 60 nucleotides on average (p
= 0.004; 95% HPD CI [−106.8, −20.2]). Increased assembly contiguity (measured as contig
N50) was associated with an average of *ca.* 13 *fewer* predicted genes per mega base-pair
increase in contig N50, but this effect was marginal (p = 0.05; CI [-26, -0.6]; Fig 2; S1 File).
CDS length was unaffected by the assembly contiguity. In the high-stringency subset, where
all short-read-only assemblies were excluded, the effect of assembly contiguity on gene
number was substantially reduced and no longer statistically supported (*ca.* 8 fewer genes per
Mbp increase in contig N50; p = 0.07), and CDS length remained unaffected (S1 File). These
patterns suggest that short-read assemblies tend to inflate gene counts by fragmenting genes
into multiple partial models, whereas long-read assemblies and higher contiguity facilitate the
recovery of longer, more complete CDSs and reduce spurious gene inflation.
568
The inclusion of RNA sequencing data unexpectedly *reduced* gene number (by *ca.* 440 genes;
p=0.002; CI [-715, -201])—without affecting CDS length (p=0.3, CI [-37.2, 17.1]). This may
reflect a reduction in the overall false-positive rate, a reduction in annotated pseudogenes, and
the joining of disjunct exons. In the high-stringency subset, the effect of RNA sequencing data
on gene number was no longer detectable (S1 File). This suggests that the apparent reduction
in gene number associated with RNAseq in the full dataset primarily reflects improvements in
annotation accuracy for lower-quality or more fragmented assemblies, rather than a
fundamental dependence on transcriptomic evidence. In high-quality, long-read assemblies,
comparative annotation alone appears sufficient to recover most gene models reliably,
rendering additional RNA-seq data less influential.
579
We found that *ca.* 13 genes were on average gained with each additional 1Mbp of genome
assembly size (p<0.001; CI [10.4, 16.5]), but mean CDS length only decreased by less than 1
bp (p<0.001; CI [-0.9, -0.4]) (Fig 2)—that is, larger genome assemblies contained more genes,
but these genes were slightly shorter on average. In addition, total CDS length increased only

weakly with genome size, by approximately 13 Kbp per Mbp (p < 0.001; CI [9, 17]), indicating
that increases in gene number are not accompanied by proportional expansion of coding
capacity (S1 File).This pattern is consistent with increased fragmentation of gene models in
larger or more repetitive genomes [81], but is also observed in the high-stringency dataset,
suggesting that biological processes such as lineage-specific expansion of small gene families
may contribute alongside technical effects.
590
After accounting for these fixed effects, we found little evidence for major differences in gene
number or CDS length among *Drosophila* clades. Posterior estimates of differences among
internal nodes generally had confidence intervals overlapping zero, suggesting that number of
genes and CDS length have remained relatively stable across lineages (S5 Fig). Nevertheless,
the *obscura* group had a generally higher inferred gene number and shorter CDS lengths
compared to other clades (S5 Fig), which may reflect lineage-specific effects such as
variations in karyotype, including transitions among sex chromosomes [82]. Correspondingly,
phylogenetic heritability was moderate for gene number (40%; CI [20.5, 63.8]) and low for
mean CDS length (9.7%; CI [1.5, 18.7]), indicating that while evolutionary history plays a role,
species-specific factors contribute substantially to variation in these traits. Interestingly, when
restricting the analysis to the high-confidence subset of 215 genomes, point estimates of
phylogenetic heritability increased for gene number (61%; CI [47.4, 77.8]) and slightly
decreased for mean CDS length (6.8%; [0.2, 15.3]), although these differences were not
significant. This shift suggests that lower-quality assemblies may add noise to gene number
estimates, thereby obscuring underlying phylogenetic signal.
606
607 Overall, our results indicate that while gene number and CDS length variation occur at the
608 species level, they do not seem to have strong phylogenetic structuring at deeper evolutionary
609 timescales, perhaps predominantly reflecting differences in assembly and annotation rather
610 than long-term evolutionary trends.

GC composition and Codon Usage Bias in Drosophilidae

To illustrate the potential utility of our new annotations, we analyzed variation in GC content
and its relationship with codon usage bias (CUB) across the full set of 301 drosophilid species
[72, 83]. Genomic GC content ranged from 21% in *Drosophila neohypocausta* to 49% in
Drosophila nannoptera (S6 Table), with coding regions, as expected, showing higher GC
content (range: 41–57% GC) than the genome-wide average. GC content at third codon
positions (GC3) is a widely used proxy for codon bias, and reflects a balance between
mutational pressure and selection acting on synonymous mutations [72, 83]. In our analysis,
GC3 was highly correlated between related species, with an estimated phylogenetic heritability
of 1 (i.e. no residual variance that is not captured by the phylogenetic effect), indicating strong
conservation within clades and little variation among closely related species. GC3 was also
positively correlated with non-coding GC content (Fig 3 and Fig 4; phylogenetic correlation
from the PGLMM 0.52; p<0.001; CI [0.41, 0.59]), suggesting that genome-wide mutational
biases contribute to both coding and non-coding base composition. Nevertheless, we observed
substantial clade-specific deviations; notably the willistoni and saltans groups, along with

subfamily Steganinae, had much lower GC3, whereas the genus *Zaprionus* and the
melanogaster, montium, obscura, ananassae, repleta, and virilis species groups exhibited
elevated GC3 (S5 Fig). These differences mirror a recent analysis of 29 *Drosophila* species, in
which subgenera *Sophophora* and *Drosophila* exhibited distinct codon preferences [72]. Such
lineage-specific differences could reflect factors beyond mutation bias or overall GC
composition, potentially including selection for translational efficiency or efficacy [84].
632
633
Fig 3. Codon usage, amino acid composition and selection on codon usage across
Drosophilidae.
(A) Phylogenetic distribution of GC content at third codon positions (GC3), non-coding GC
content, and strength of selection on codon usage bias (*S*) across Drosophilidae. The tree
is color-coded by clades, and branches are colored according to GC3. Inner ring shows GC
content in non-coding regions and outer ring shows strength of selection on codon usage.
Major drosophilid species groups are indicated by arcs positioned outside the outermost tile
layer. (B) Principal component analysis (PCA) of amino acid usage across species, showing
that closely related species exhibit similar amino acid composition patterns. The numerical
data underlying both panels are provided in S6 Table.
644
645
Fig 4. Correlation matrix of codon usage, genome features, and amino acid
composition.
Pairwise phylogenetic correlations between GC content at third codon positions (GC3), GC
content in non-coding regions, strength of selection on codon usage bias (*S*), genome size,
principal components PC1 and PC2 of amino acid usage, and nitrogen-to-carbon (N/C) ratio
of amino acids. Correlations are derived from the posterior distribution of the phylogenetic
variance–covariance matrix and represent evolutionary covariation among traits after
accounting for shared ancestry. Values shown correspond to posterior modes; uncertainty
is reported as 95% highest posterior density intervals. Strong correlations (positive or
negative) are highlighted in green and orange, respectively, indicating relationships
between nucleotide composition, codon usage, and amino acid preferences. The numerical
data underlying this figure are provided in S7 Table.
658
659
To quantify the role of selection in determining codon usage, we estimated the strength of
selection on two-fold codons [quantified by the 'S' statistic of 76]. Estimates of *S* ranged from
0.24 in *Drosophila pachea* (95% bootstrap interval across genes [0.22, 0.29]) up to 1.08 [0.96,
1.20] in *Drosophila takahashii* (Fig 3). As expected, the melanogaster, montium, and
ananassae groups showed elevated GC3 and *S*, confirming stronger selection in favour of GC-
ending codons in these groups [72, 84, 85]. However, the willistoni and saltans groups—which
display low GC3—also showed relatively high *S*, confirming that the AT-bias seen in
Drosophila willistoni is (at least in part) a result of selection [84, 86, 87]. A similar pattern was
also seen in the subfamily Steganinae and overall GC3 and *S* are negatively correlated across
Drosophilidae—indicating that species with higher GC3 are, on average, actually experiencing

weaker selection on codon usage (Fig 4; phylogenetic correlation from the PGLMM: -0.2;
p<0.001; CI [-0.33, -0.10]). Interestingly, a weak *positive* correlation between *S* and genome
size (Fig 4; phylogenetic correlation from the PGLMM 0.15; p=0.02; CI: [0.03, 0.32]) suggests
that species with larger genomes tend to experience slightly stronger selection on codon
usage—in contrast to what might be expected under relaxed constraint in species with small
effective population size [88].

Amino Acid Composition

To investigate whether the variation in codon usage is associated with variation in amino acid
composition, we analyzed the relative proportions of all 20 amino acids across the annotated
proteins of Drosophilidae. In general, it is thought that amino acid usage is influenced by a
combination of mutational biases, translational selection, and functional constraints—but
genome-wide nucleotide composition has been shown to play a significant role in shaping
amino acid frequencies [83, 89]. Our principal component analysis (PCA) revealed that more
closely related species share more similar amino acid usage patterns (Fig 3). Principal
component 1 primarily separated species based on GC content in the codons, with high-GC3
genomes enriched for GC-rich amino acids (Pro, Gly, Ala, Arg) and low-GC3 genomes
enriched for AT-rich amino acids (Asn, Tyr, Ile; see Fig 5). To assess whether the patterns
were linked to biochemical properties of the amino-acids, such as nitrogen-to-carbon (N/C)
ratio or amino acid essentiality [as measured in Drosophila melanogaster; 90, 91], we
examined the remaining PCA loadings. However, we found no clear patterns, suggesting that
other factors, such as protein structure or functional constraints, may play a small role in
shaping variation in amino acid use among species of Drosophilidae (Fig 5).
692
Fig 5. Principal component analysis (PCA) loadings of amino acid usage.
Bar plots show the loadings of individual amino acids on the first two principal components:
(A) PC1 and (B) PC2. Amino acids are coloured according to their nitrogen to carbon (N:C)
ratio, with higher ratios in red and lower ratios in blue. Essentiality categories (essential,
semi-essential, non-essential) are indicated alongside each bar. Positive and negative
loadings reflect the relative contribution of each amino acid to the corresponding principal
component. The numerical data underlying both panels are provided in S8 Table.

Conclusions

This work, to generate standardized, simultaneous multi-species coding DNA sequence
annotations across 301 species of Drosophilidae, forms part of an ongoing community effort
working toward a comprehensive genomic study of the entire family [33]. We envisage that
these new annotations, orthology assignments, and multiple sequence alignments will provide
a valuable resource for both single-gene and genome-wide evolutionary studies. And, along
with future updates as new genomes are sequenced, this resource will support future research
in studies of adaptation and functional genomics within this key model clade.

Supporting information

S1 Fig. Comparison of coding sequence annotations generated by CAT-BRAKER and BRAKER3 alone relative to RefSeq annotations.
Panels show CDS-level precision (A), recall (B), and Jaccard similarity (C), quantified based on pairwise CDS overlap with RefSeq gene models. The numerical data underlying all panels are provided in S9 Table.

S2 Fig. BUSCO completeness scores for genome assemblies and corresponding annotated protein sets across drosophilid species, illustrating concordance between assembly-level and annotation-level recovery of conserved single-copy orthologs.
(A) Annotation-level BUSCO completeness plotted against gene number, showing a weak negative association between completeness and the number of predicted genes. (B) Parallel-axis dot plot comparing genome-level and annotated protein-level BUSCO completeness for each species. Species with less than 5% difference are colored grey. (C) Relationship between genome-level and annotated protein-level BUSCO completeness, demonstrating a strong positive correlation. Fitted lines and 95% confidence intervals in panels (A) and (C) are derived from non-phylogenetic linear models and are shown for illustrative purposes only. The numerical data underlying all panels are provided in S5 Table.

S3 Fig. Overview of orthology assignments across 301 Drosophilidae species.
The figure shows a time-calibrated phylogenetic tree with ancestral reconstruction of protein-coding gene number mapped onto branches. Tip labels are coloured by the reference species used for comparative annotation within each clade; stars denote reference species, and filled versus open circles indicate the presence or absence of RNA-seq data, respectively. A concentric tile layer indicates the percentage of genes assigned to orthogroups for each species. The outer stacked bar layer shows the number of genes belonging to hierarchical orthologous groups (HOGs) with different levels of species representation (>99%, 75–99%, 50–75%, <50%), as well as species-specific HOGs. The numerical data underlying this figure is provided in S4 Table.

S4 Fig. Classification of hierarchical orthologous groups (HOGs) by phylogenetic depth and species representation.
(A) Number of HOGs plotted against the estimated age of their most recent common ancestor (MRCA, in million years), reflecting the evolutionary depth of gene families. (B) Number of HOGs plotted against the number of species in which they are present. (C) Number of ancient HOGs (defined as having an MRCA ≥50 million years ago) plotted against species representation. (D) Number of HOGs containing at least one *Drosophila melanogaster* gene plotted against the number of species represented. This figure was generated using the HOG summary tables available from the Zenodo repository (https://doi.org/10.5281/zenodo.15016917) and the time-calibrated species phylogeny provided in S5 file.

S5 Fig. Posterior estimates of genomic and annotation features reconstructed at the most recent common ancestors (MRCAs) of major drosophilid species groups. Panels show inferred values for gene number (A), mean CDS length (B), GC3 content (C) and strength of selection (S) on codon usage (D), estimated using phylogenetic mixed models. The R model objects used to generate these estimates are provided in S1 Data.

S1 Table. SRA accession numbers for RNA-seq datasets used in genome annotation.

S2 Table. *Drosophila melanogaster* genes ranked by overall expression level (FPKM).

S3 Table. Expression category assignments for HOGs based on *D. melanogaster* gene expression ranks.

S4 Table. Summary statistics for genome assemblies, genome annotations and orthology assignments across species.

S5 Table. BUSCO and OMArk completeness assessments for genome annotations.

S6 Table. Estimates of codon usage bias metrics (GC3, GC content in non-coding regions and selection strength S) across species.

S7 Table. Posterior modes and 95% HPD intervals for phylogenetic correlations among codon usage bias, nucleotide composition, genome size, and amino acid properties.

S8 Table. Principal component loadings for amino acid usage, including nitrogen-to-carbon ratios and essentiality categories.

S9 Table. CDSs Precision, Recall and Jaccard similarity between CAT-BRAKER and BRAKER only annotations compared to RefSeq annotations.

S1 File. MCMCglmm model summaries and phylogenetic heritability's for full set and high-stringency subset.

S2 File. Ultrametric species phylogeny inferred using 251 hierarchical orthologous groups (HOGs).

S3 File. Dendrograms comparing species phylogenies inferred from BUSCO and HOG-based datasets.

S4 File. HOG and BUSCO phylogenetic trees annotated with posterior probabilities from Bayesian analyses.

S5 File. Newick files of species trees generated in this study.

S1 Data. R MCMCglmm model objects generated and used in this study.

Funding

This work was supported by a Darwin Trust PhD studentship to PD, Biotechnology and Biological Sciences Research Council grant BB/T007516/1 to DJO, and National Institutes of Health (NIH) National Institute of General Medical Sciences (NIGMS) grant R35 GM118165 to DAP. DAP is a CZ Biohub Investigator. For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY) license to any Author Accepted Manuscript version arising from this submission.

Acknowledgements

We wish to thank members of the University of Edinburgh Institute for Ecology and Evolution for the collaborative provision of shared computational resources, James Galbraith for help with EarlGrey and TEstrainer, and Eric Lai and Garima Setia for feedback on missing gene models. We thank the editor and two reviewers for their positive and helpful comments on an earlier version of this manuscript. We would particularly like to acknowledge the contributions of the broader *Drosophila* research community, whose collaborative efforts in genome sequencing have made this work possible.

References

1. Ejigu GF, Jung J. Review on the Computational Genome Annotation of Sequences Obtained by Next-Generation Sequencing. Biology (Basel). 2020;9(9):295. doi: 10.3390/biology9090295.
2. Kwon T, Hanschen ER, Hovde BT. Addressing the pervasive scarcity of structural annotation in eukaryotic algae. Sci Rep. 2023;13(1):1687. doi: 10.1038/s41598-023-27881-0.
3. Vuruputoor VS, Monyak D, Fetter KC, Webster C, Bhattarai A, Shrestha B, et al. Welcome to the big leaves: Best practices for improving genome annotation in non-model plant genomes. Appl Plant Sci. 2023;11(4):e11533. doi: 10.1002/aps3.11533.
4. Mathe C, Dunand C. Automatic Prediction and Annotation: There Are Strong Biases for Multigenic Families. Front Genet. 2021;12:697477. doi: 10.3389/fgene.2021.697477.
5. Promponas VJ, Iliopoulos I, Ouzounis CA. Annotation inconsistencies beyond sequence similarity-based function prediction - phylogeny and genome structure. Stand Genomic Sci. 2015;10(1):108. doi: 10.1186/s40793-015-0101-2.
6. Scalzitti N, Jeannin-Girardon A, Collet P, Poch O, Thompson JD. A benchmark study of ab initio gene prediction methods in diverse eukaryotic organisms. BMC Genomics. 2020;21(1):293. doi: 10.1186/s12864-020-6707-9.
7. Colbourne JK, Pfrender ME, Gilbert D, Thomas WK, Tucker A, Oakley TH, et al. The ecoresponsive genome of Daphnia pulex. Science. 2011;331(6017):555-61. doi: 10.1126/science.1197761.

8.   Mendes MF, Valer FB, Vieira JGA, Blauth ML, Gottschalk MS. Diversity of Drosophilidae
(Insecta, Diptera) in the Restinga forest of southern Brazil. Revista Brasileira De
Entomologia. 2017;61(3):248-56. doi: 10.1016/j.rbe.2017.05.002.
9.   van Dijk EL, Naquin D, Gorrichon K, Jaszczyszyn Y, Ouazahrou R, Thermes C, et al.
Genomics in the long-read sequencing era. Trends Genet. 2023;39(9):649-71. doi:
10.1016/j.tig.2023.04.006.
10. Espinosa E, Bautista R, Larrosa R, Plata O. Advancements in long-read genome
sequencing technologies and algorithms. Genomics. 2024;116(3):110842. doi:
10.1016/j.ygeno.2024.110842.
11. Darwin Tree of Life Project C. Sequence locally, think globally: The Darwin Tree of Life
Project. Proc Natl Acad Sci U S A. 2022;119(4):e2115642118. doi:
10.1073/pnas.2115642118.
12. Rhie A, McCarthy SA, Fedrigo O, Damas J, Formenti G, Koren S, et al. Towards complete
and error-free genome assemblies of all vertebrate species. Nature. 2021;592(7856):737-
46. doi: 10.1038/s41586-021-03451-0.
13. Sharaf A, Ndiribe CC, Omotoriogun TC, Abueg L, Badaoui B, Badiane Markey FJ, et al.
Bridging the gap in African biodiversity genomics and bioinformatics. Nat Biotechnol.
2023;41(9):1348-54. doi: 10.1038/s41587-023-01933-2.
14. Freedman AH, Sackton TB. Building better genome annotations across the tree of life.
Genome Research. 2025;35(5):1261-76. doi: 10.1101/gr.280377.124.
15. Gabriel L, Bruna T, Hoff KJ, Ebel M, Lomsadze A, Borodovsky M, et al. BRAKER3: Fully
automated genome annotation using RNA-seq and protein evidence with GeneMark-ETP,
AUGUSTUS, and TSEBRA. Genome Res. 2024;34(5):769-77. doi: 10.1101/gr.278090.123.
16. Stanke M, Diekhans M, Baertsch R, Haussler D. Using native and syntenically mapped
cDNA alignments to improve de novo gene finding. Bioinformatics. 2008;24(5):637-44. doi:
10.1093/bioinformatics/btn013.
17. Cantarel BL, Korf I, Robb SM, Parra G, Ross E, Moore B, et al. MAKER: an easy-to-use
annotation pipeline designed for emerging model organism genomes. Genome Res.
2008;18(1):188-96. doi: 10.1101/gr.6743907.
18. Venkatraman M, Fleischer RC, Tsuchiya MTN. Comparative Analysis of Annotation
Pipelines Using the First Japanese White-Eye (Zosterops japonicus) Genome. Genome Biol
Evol. 2021;13(5). doi: 10.1093/gbe/evab063.
19. Nachtweide S, Romoth L, Stanke M. Comparative Genome Annotation. Methods in
Molecular Biology: Springer US; 2024. p. 165-87.
20. Weisman CM, Murray AW, Eddy SR. Mixing genome annotation methods in a comparative
analysis inflates the apparent number of lineage-specific genes. Curr Biol.
2022;32(12):2632-9 e2. doi: 10.1016/j.cub.2022.04.085.
21. König S, Romoth L, Stanke M. Comparative Genome Annotation. Methods in Molecular
Biology: Springer New York; 2018. p. 189-212.
22. Fiddes IT, Armstrong J, Diekhans M, Nachtweide S, Kronenberg ZN, Underwood JG, et al.
Comparative Annotation Toolkit (CAT)-simultaneous clade and personal genome
annotation. Genome Res. 2018;28(7):1029-38. doi: 10.1101/gr.233460.117.
23. Beller M, Oliver B. One hundred years of high-throughput Drosophila research.
Chromosome Res. 2006;14(4):349-62. doi: 10.1007/s10577-006-1065-2.
24. Richards S, Liu Y, Bettencourt BR, Hradecky P, Letovsky S, Nielsen R, et al. Comparative
genome sequencing of Drosophila pseudoobscura: chromosomal, gene, and cis-element
evolution. Genome Res. 2005;15(1):1-18. doi: 10.1101/gr.3059305.
25. Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, Amanatides PG, et al. The
genome sequence of Drosophila melanogaster. Science. 2000;287(5461):2185-95. doi:
10.1126/science.287.5461.2185.

26. Drosophila 12 Genomes C, Clark AG, Eisen MB, Smith DR, Bergman CM, Oliver B, et al.
Evolution of genes and genomes on the Drosophila phylogeny. Nature.
2007;450(7167):203-18. doi: 10.1038/nature06341.
27. mod EC, Roy S, Ernst J, Kharchenko PV, Kheradpour P, Negre N, et al. Identification of
functional elements and regulatory circuits by Drosophila modENCODE. Science.
2010;330(6012):1787-97. doi: 10.1126/science.1198374.
28. Bronski MJ, Martinez CC, Weld HA, Eisen MB. Whole Genome Sequences of 23 Species
from the Drosophila montium Species Group (Diptera: Drosophilidae): A Resource for
Testing Evolutionary Hypotheses. G3 (Bethesda). 2020;10(5):1443-55. doi:
10.1534/g3.119.400959.
29. Puerma E, Orengo DJ, Cruz F, Gomez-Garrido J, Librado P, Salguero D, et al. The High-
Quality Genome Sequence of the Oceanic Island Endemic Species Drosophila guanche
Reveals Signals of Adaptive Evolution in Genes Related to Flight and Genome Stability.
Genome Biol Evol. 2018;10(8):1956-69. doi: 10.1093/gbe/evy135.
30. Mahajan S, Wei KH, Nalley MJ, Gibilisco L, Bachtrog D. De novo assembly of a young
Drosophila Y chromosome using single-molecule sequencing and chromatin conformation
capture. PLoS Biol. 2018;16(7):e2006348. doi: 10.1371/journal.pbio.2006348.
31. Li F, Rane RV, Luria V, Xiong Z, Chen J, Li Z, et al. Phylogenomic analyses of the genus
Drosophila reveals genomic signals of climate adaptation. Mol Ecol Resour.
2022;22(4):1559-81. doi: 10.1111/1755-0998.13561.
32. Thiebaut A, Altenhoff AM, Campli G, Glover N, Dessimoz C, Waterhouse RM. DrosOMA:
the Drosophila Orthologous Matrix browser. F1000Res. 2023;12:936. doi:
10.12688/f1000research.135250.2.
33. Kim BY, Gellert HR, Church SH, Suvorov A, Anderson SS, Barmina O, et al. Single-fly
genome assemblies fill major phylogenomic gaps across the Drosophilidae Tree of Life.
PLoS Biol. 2024;22(7):e3002697. doi: 10.1371/journal.pbio.3002697.
34. Suvorov A, Kim BY, Wang J, Armstrong EE, Peede D, D'Agostino ERR, et al. Widespread
introgression across a phylogeny of 155 Drosophila genomes. Curr Biol. 2022;32(1):111-23
e5. doi: 10.1016/j.cub.2021.10.052.
35. Obbard DJ, Wellcome Sanger Institute Tree of Life p, Wellcome Sanger Institute Scientific
Operations DNAPc, Tree of Life Core Informatics c, Darwin Tree of Life C. The genome
sequence of a drosophilid fruit fly, Chymomyza fuscimana (Drosophilidae) (Zetterstedt,
1838). Wellcome Open Res. 2023;8:477. doi: 10.12688/wellcomeopenres.20122.1.
36. O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, et al. Reference
sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional
annotation. Nucleic Acids Res. 2016;44(D1):D733-45. doi: 10.1093/nar/gkv1189.
37. Zhou Q, Bachtrog D. Sex-specific adaptation drives early sex chromosome evolution in
Drosophila. Science. 2012;337(6092):341-5. doi: 10.1126/science.1225385.
38. Renschler G, Richard G, Valsecchi CIK, Toscano S, Arrigoni L, Ramirez F, et al. Hi-C
guided assemblies reveal conserved regulatory topologies on X and autosomes despite
extensive genome shuffling. Genes Dev. 2019;33(21-22):1591-612. doi:
10.1101/gad.328971.119.
39. Sanchez-Flores A, Penaloza F, Carpinteyro-Ponce J, Nazario-Yepiz N, Abreu-Goodger C,
Machado CA, et al. Genome Evolution in Three Species of Cactophilic Drosophila. G3
(Bethesda). 2016;6(10):3097-105. doi: 10.1534/g3.116.033779.
40. Wei KH, Mai D, Chatla K, Bachtrog D. Dynamics and Impacts of Transposable Element
Proliferation in the Drosophila nasuta Species Group Radiation. Mol Biol Evol. 2022;39(5).
doi: 10.1093/molbev/msac080.
41. Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO: assessing
genome assembly and annotation completeness with single-copy orthologs. Bioinformatics.
2015;31(19):3210-2. doi: 10.1093/bioinformatics/btv351.

42. Smit A, Hubley, R & Green, P. RepeatMasker Open-4.0 2013-2015. Available from:
http://www.repeatmasker.org.
43. Storer J, Hubley R, Rosen J, Wheeler TJ, Smit AF. The Dfam community resource of
transposable element families, sequence models, and genome annotations. Mob DNA.
2021;12(1):2. doi: 10.1186/s13100-020-00230-y.
44. Yuan D, Ahamed A, Burgin J, Cummins C, Devraj R, Gueye K, et al. The European
Nucleotide Archive in 2023. Nucleic Acids Res. 2024;52(D1):D92-D7. doi:
10.1093/nar/gkad1067.
45. Bushnell B. BBMap: A Fast, Accurate, Splice-Aware Aligner 2014. Available from:
https://sourceforge.net/projects/bbmap/.
46. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21. doi:
10.1093/bioinformatics/bts635.
47. Kriventseva EV, Kuznetsov D, Tegenfeldt F, Manni M, Dias R, Simao FA, et al. OrthoDB
v10: sampling the diversity of animal, plant, fungal, protist, bacterial and viral genomes for
evolutionary and functional annotations of orthologs. Nucleic Acids Res. 2019;47(D1):D807-
D11. doi: 10.1093/nar/gky1053.
48. Li H. Protein-to-genome alignment with miniprot. Bioinformatics. 2023;39(1). doi:
10.1093/bioinformatics/btad014.
49. Bruna T, Li H, Guhlin J, Honsel D, Herbold S, Stanke M, et al. Galba: genome annotation
with miniprot and AUGUSTUS. BMC Bioinformatics. 2023;24(1):327. doi: 10.1186/s12859-
023-05449-z.
50. Hickey G, Paten B, Earl D, Zerbino D, Haussler D. HAL: a hierarchical format for storing
and analyzing multiple genome alignments. Bioinformatics. 2013;29(10):1341-2. doi:
10.1093/bioinformatics/btt128.
51. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol. 2013;30(4):772-80. doi:
10.1093/molbev/mst010.
52. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, et al.
IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic
Era. Mol Biol Evol. 2020;37(5):1530-4. doi: 10.1093/molbev/msaa015.
53. Zhang C, Rabiee M, Sayyari E, Mirarab S. ASTRAL-III: polynomial time species tree
reconstruction from partially resolved gene trees. BMC Bioinformatics. 2018;19(Suppl
6):153. doi: 10.1186/s12859-018-2129-y.
54. Huerta-Cepas J, Serra F, Bork P. ETE 3: Reconstruction, Analysis, and Visualization of
Phylogenomic Data. Mol Biol Evol. 2016;33(6):1635-8. doi: 10.1093/molbev/msw046.
55. Matthews BB, Dos Santos G, Crosby MA, Emmert DB, St Pierre SE, Gramates LS, et al.
Gene Model Annotations for Drosophila melanogaster: Impact of High-Throughput Data. G3
(Bethesda). 2015;5(8):1721-36. doi: 10.1534/g3.115.018929.
56. Armstrong J, Hickey G, Diekhans M, Fiddes IT, Novak AM, Deran A, et al. Progressive
Cactus is a multiple-genome aligner for the thousand-genome era. Nature.
2020;587(7833):246-51. doi: 10.1038/s41586-020-2871-y.
57. Konig S, Romoth LW, Gerischer L, Stanke M. Simultaneous gene finding in multiple
genomes. Bioinformatics. 2016;32(22):3388-95. doi: 10.1093/bioinformatics/btw494.
58. Baril T, Galbraith J, Hayward A. Earl Grey: A Fully Automated User-Friendly Transposable
Element Annotation and Analysis Pipeline. Mol Biol Evol. 2024;41(4). doi:
10.1093/molbev/msae068.
59. Nevers Y, Warwick Vesztrocy A, Rossier V, Train CM, Altenhoff A, Dessimoz C, et al.
Quality assessment of gene repertoire annotations with OMArk. Nat Biotechnol.
2025;43(1):124-33. doi: 10.1038/s41587-024-02147-w.

60. Kuznetsov D, Tegenfeldt F, Manni M, Seppey M, Berkeley M, Kriventseva EV, et al.
OrthoDB v11: annotation of orthologs in the widest sampling of organismal diversity. Nucleic
Acids Res. 2023;51(D1):D445-D51. doi: 10.1093/nar/gkac998.
61. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biol. 2019;20(1):238. doi: 10.1186/s13059-019-1832-y.
62. Ranwez V, Douzery EJP, Cambon C, Chantret N, Delsuc F. MACSE v2: Toolkit for the
Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. Mol Biol
Evol. 2018;35(10):2582-4. doi: 10.1093/molbev/msy159.
63. Di Franco A, Poujol R, Baurain D, Philippe H. Evaluating the usefulness of alignment
filtering methods to reduce the impact of errors on evolutionary inferences. BMC Evol Biol.
2019;19(1):21. doi: 10.1186/s12862-019-1350-2.
64. Freckleton RP. The seven deadly sins of comparative analysis. J Evol Biol.
2009;22(7):1367-75. doi: 10.1111/j.1420-9101.2009.01757.x.
65. Hadfield JD, Nakagawa S. General quantitative genetic methods for comparative biology:
phylogenies, taxonomies and multi-trait models for continuous and categorical characters. J
Evol Biol. 2010;23(3):494-508. doi: 10.1111/j.1420-9101.2009.01915.x.
66. Hadfield JD. MCMC Methods for Multi-Response Generalized Linear Mixed Models: The
MCMCglmm R Package. Journal of Statistical Software. 2010;33(2):1-22. doi: DOI
10.18637/jss.v033.i02.
67. Suchard MA, Lemey P, Baele G, Ayres DL, Drummond AJ, Rambaut A. Bayesian
phylogenetic and phylodynamic data integration using BEAST 1.10. Virus Evol.
2018;4(1):vey016. doi: 10.1093/ve/vey016.
68. Drummond AJ, Ho SY, Phillips MJ, Rambaut A. Relaxed phylogenetics and dating with
confidence. PLoS Biol. 2006;4(5):e88. doi: 10.1371/journal.pbio.0040088.
69. Gernhard T. The conditioned reconstructed process. J Theor Biol. 2008;253(4):769-78. doi:
10.1016/j.jtbi.2008.04.005.
70. Rambaut A, Drummond AJ, Xie D, Baele G, Suchard MA. Posterior Summarization in
Bayesian Phylogenetics Using Tracer 1.7. Syst Biol. 2018;67(5):901-4. doi:
10.1093/sysbio/syy032.
71. Hershberg R, Petrov DA. Selection on codon bias. Annu Rev Genet. 2008;42(Volume 42,
2008):287-99. doi: 10.1146/annurev.genet.42.110807.091442.
72. Kokate PP, Techtmann SM, Werner T. Codon usage bias and dinucleotide preference in 29
Drosophila species. G3 (Bethesda). 2021;11(8). doi: 10.1093/g3journal/jkab191.
73. Zhang H. cubar: Codon Usage Bias Analysis. R package 2024. Available from:
https://CRAN.R-project.org/package=cubar.
74. Blankenberg D, Taylor J, Schenck I, He J, Zhang Y, Ghent M, et al. A framework for
collaborative analysis of ENCODE data: making large-scale analyses biologist-friendly.
Genome Res. 2007;17(6):960-4. doi: 10.1101/gr.5578007.
75. Rice P, Longden I, Bleasby A. EMBOSS: the European Molecular Biology Open Software
Suite. Trends Genet. 2000;16(6):276-7. doi: 10.1016/s0168-9525(00)02024-2.
76. dos Reis M, Wernisch L. Estimating translational selection in eukaryotic genomes. Mol Biol
Evol. 2009;26(2):451-61. doi: 10.1093/molbev/msn272.
77. Eyre-Walker A, Bulmer M. Synonymous substitution rates in enterobacteria. Genetics.
1995;140(4):1407-12. doi: 10.1093/genetics/140.4.1407.
78. Bachtrog D, Mahajan S, Bracewell R. Massive gene amplification on a recently formed
Drosophila Y chromosome. Nat Ecol Evol. 2019;3(11):1587-97. doi: 10.1038/s41559-019-
1009-9.
79. Yang H, Jaime M, Polihronakis M, Kanegawa K, Markow T, Kaneshiro K, et al. Re-
annotation of eight Drosophila genomes. Life Science Alliance. 2018;1(6):e201800156. doi:
10.26508/lsa.201800156.

80. Finet C, Kassner VA, Carvalho AB, Chung H, Day JP, Day S, et al. DrosoPhyla: Resources
for Drosophilid Phylogeny and Systematics. Genome Biol Evol. 2021;13(8). doi:
10.1093/gbe/evab179.
81. Konstantinidis KT, Tiedje JM. Trends between gene content and genome size in prokaryotic
species with larger genomes. Proc Natl Acad Sci U S A. 2004;101(9):3160-5. doi:
10.1073/pnas.0308653100.
82. Bracewell R, Bachtrog D. Complex Evolutionary History of the Y Chromosome in Flies of
the Drosophila obscura Species Group. Genome Biology and Evolution. 2020;12(5):494-
505. doi: 10.1093/gbe/evaa051.
83. Behura SK, Severson DW. Comparative analysis of codon usage bias and codon context
patterns between dipteran and hymenopteran sequenced genomes. PLoS One.
2012;7(8):e43111. doi: 10.1371/journal.pone.0043111.
84. Heger A, Ponting CP. Variable strength of translational selection among 12 Drosophila
species. Genetics. 2007;177(3):1337-48. doi: 10.1534/genetics.107.070466.
85. Vicario S, Moriyama EN, Powell JR. Codon usage in twelve species of Drosophila. BMC
Evol Biol. 2007;7(1):226. doi: 10.1186/1471-2148-7-226.
86. Singh ND, Arndt PF, Petrov DA. Minor shift in background substitutional patterns in the
Drosophila saltans and willistoni lineages is insufficient to explain GC content of coding
sequences. BMC Biol. 2006;4(1):37. doi: 10.1186/1741-7007-4-37.
87. Powell JR, Sezzi E, Moriyama EN, Gleason JM, Caccone A. Analysis of a shift in codon
usage in Drosophila. J Mol Evol. 2003;57 Suppl 1(0):S214-25. doi: 10.1007/s00239-003-
0030-3.
88. Petit N, Barbadilla A. Selection efficiency and effective population size in Drosophila
species. J Evol Biol. 2009;22(3):515-26. doi: 10.1111/j.1420-9101.2008.01672.x.
89. Williford A, Demuth JP. Gene expression levels are correlated with synonymous codon
usage, amino acid composition, and gene architecture in the red flour beetle, Tribolium
castaneum. Mol Biol Evol. 2012;29(12):3755-66. doi: 10.1093/molbev/mss184.
90. Croset V, Schleyer M, Arguello JR, Gerber B, Benton R. A molecular and neuronal basis for
amino acid sensing in the Drosophila larva. Sci Rep. 2016;6(1):34871. doi:
10.1038/srep34871.
91. Park J, Carlson JR. Physiological responses of the Drosophila labellum to amino acids. J
Neurogenet. 2018;32(1):27-36. doi: 10.1080/01677063.2017.1406934.
1065

A

Species labels
- ✳ Reference species
- ● Species with RNAseq
- ○ Species without RNAseq

Clades
- ▬ ananassae group
- ▬ Hawaiian/Scaptomyza
- ▬ immigrans group
- ▬ *Lordiphosa* genus
- ▬ melanogaster subgroup
- ▬ montium group
- ▬ obscura group
- ▬ quinaria group
- ▬ repleta group
- ▬ saltans group
- ▬ Subfamily Steganinae
- ▬ virilis group
- ▬ willistoni group
- ▬ *Zaprionus* genus

6 / mya

Protein-coding genes
- 21000
- 18000
- 15000

Mean CDS Length (Kbp)
- 1.4
- 1.5
- 1.6
- 1.7

BUSCO (%)
- 94
- 96
- 98

Genome size (Mbp)
- 200
- 300
- 400

Contig N50 (Mbp)
- 10
- 20
- 30

B

Total Number of Genes: 24k, 21k, 18k, 15k, 12k

Mean CDS length (Kbp): 1.6, 1.4, 1.2, 1.0

Genes   CDS length

A

Clades
- ananassae group
- Hawaiian/Scaptomyza
- immigrans group
- *Lordiphosa* genus
- melanogaster subgroup
- montium group
- obscura group
- quinaria group
- repleta group
- saltans group
- Subfamily Steganinae
- virilis group
- willistoni group
- *Zaprionus* genus

6 / mya

GC3 (%)
70
60
50
40

GC non-Coding (%)
40
35
30
25

S
1.0
0.8
0.6
0.4

B

PC2 22.453 %

PC1 45.801 %

Dcolorata Dbusckii

Snigrithor

Lmaculata

Cindagator Lmontana

Lvaria

Acommunis Aminor

Amariae