Real-world objects scaffold visual working memory for features: Increased neural delay activity when colors are remembered as part of meaningful objects

Yong Hoon Chung[1]
Timothy F. Brady[2]
Viola S. Störmer[1]

[1]Dartmouth College, Department of Psychological and Brain Sciences
[2]University of California San Diego, Department of Psychology

Abstract

Visual working memory is a core cognitive function that allows active storage of task-relevant visual information. While previous studies have postulated that the capacity of this system is fixed with respect to a single feature dimension, recent research has shown that working memory performance for a simple visual feature – color – is improved when this feature is encoded as part of a real-world object relative to an unrecognizable scrambled shape. Using EEG (N = 24), we here demonstrate that this performance increase is supported by enhanced neural delay activity during the retention period (indexed by the contralateral-delay-activity of the event-related potential), suggesting that the behavioral benefit is linked to the active maintenance process of working memory. Furthermore, using time-resolved neural similarity measures, we show that the neural activity is not only increased during the delay, but also more stable over time when colors are remembered as part of real-world objects. Finally, we report a novel fronto-central event-related potential that distinguishes between real-world objects and scrambled objects during encoding and maintenance processes. Overall, our results demonstrate that active visual working memory capacity for simple features is not fixed but can expand depending on what context these features are encoded in.

Introduction

Visual working memory supports the active storage of task-relevant visual information (Baddeley & Hitch, 1974; Bays et al. 2024). Its capacity is typically assessed by presenting multiple items composed of simple visual features (color, orientation) and abstract shapes (circles, polygons) for brief durations, as these conditions and stimuli are believed to best isolate visual working memory capacity separate from rehearsal, chunking, and passive memory systems (Cowan, 2001). One major conclusion from studies using such stimuli is that the capacity to maintain visual information is strictly limited, often quantified in terms of the number of objects (Luck & Vogel, 1997; Cowan, 2001; Adam, Vogel, & Awh, 2017), or content-less pointers (Awh & Vogel, 2024), or in terms of a fixed resource pool within each simple visual feature dimension that is distributed among the encoded items (e.g., Bays, Catalao, & Hussain, 2009; Bays, 2014).

However, several findings have demonstrated that how much information can be actively maintained is not fixed, but depends on the meaningfulness of the information: Familiar and meaningful stimuli, such as real-world objects, result in increased neural maintenance activity and higher behavioral performance compared to abstract and non-meaningful shapes (Brady et al., 2016; Thibeault et al., 2024; Torres et al., 2024; Asp et al., 2021). These meaningful stimuli may allow participants to extract more relevant visual and semantic features, thereby boosting capacity (see Chung et al, 2024a for a review); similar to how maintaining both color and orientation allows the storage of more total information than color alone (Luck & Vogel, 2013; Shin & Ma, 2017).

Interestingly, a recent study found that memory performance can be increased even within a single feature dimension – color – when this feature is encoded in a meaningful context (Chung et al., 2023a; Chung et al., 2023b). Specifically, in a series of behavioral experiments, participants showed better color memory performance when colors were presented as parts of real-world objects (e.g., a blue backpack) compared to unrecognizable scrambled shapes, even though the color-object associations were randomly chosen on every trial, and object identity itself was task-irrelevant. These findings challenge fixed-capacity models of working memory which assume that how much information can be maintained is strictly limited within a single feature dimension.

What explains this improvement? The observed behavioral effects could be supported by *increases* in active working memory usage; or they could be due to *more efficient* coding of real-world objects compared to abstract stimuli, potentially freeing up resources to better remember the colors; or, the performance benefits could be merely due to differences at encoding or retrieval, with no changes during maintenance. To disambiguate between these possibilities, we here examine both the amplitude and stability of neural activity during working memory maintenance. Persistent neural activity during the retention period is the hallmark of working memory, and not present when participants rely on other non-active forms of storage for retaining information (Buschman, Siegel, Roy, & Miller, 2011; Vogel & Machizawa, 2004). One particularly strong marker of active working memory is the contralateral-delay activity (CDA), a sustained negativity of the event-related potential (ERP) indexing how much visual information is actively maintained (Vogel & Machizawa, 2004; for a review, see Luria et al., 2016). We capitalize on this well-established neural marker to investigate whether better color memory performance for real-world objects is supported by increased active maintenance, decreased active maintenance, or causes no change in active maintenance. To examine representational stability, we employ time-resolved neural pattern similarity analysis across encoding and

maintenance periods. Overall, we find evidence consistent with increased active maintenance and earlier-arising neural stability for color working memory representations when they are encoded as part of real objects relative to scrambled control stimuli. Finally, we report a novel fronto-central ERP component that closely tracks this real-world object advantage by differentiating between the two object conditions during encoding and maintenance periods.

Materials & Methods

All data and example analysis codes are available on OSF (https://osf.io/bgmuw/). The experiments were approved by the Committee for the Protection of Human Subjects at Dartmouth College.

Participants

  All participants gave written informed consent prior to participating in this experiment. To obtain our goal of twenty-four usable EEG participants, twenty-nine participants were recruited from Dartmouth College using their online recruitment platform. Participants' ages ranged from 18 to 35 years, and all participants had normal or corrected-to-normal vision. Prior to the experiment participants were given the Ishihara color blindness test (Clark, 1924). Following our usual exclusion criteria in this type of working memory task (e.g., Chung et al., 2023a), we decided a priori that participants' behavioral and EEG data would be excluded if their overall behavioral performance was lower than d'< 0.5, and their EEG data would be excluded if more than 40% of trials were excluded due to EEG artifact rejection. No participant was excluded due to behavioral performance. Five participants were excluded from the EEG analysis due to artifact rejection, totaling our target of twenty-four participants. On average, about 17.7% of trials were rejected due to EEG artifacts across the remaining participants. The final sample size of 24 for the EEG analysis was determined following protocols of previous studies investigating CDA measures of meaningful stimuli (e.g., 19 participants Asp et al., 2021; 18 participants in Brady et al., 2016; 22 participants in Thibeault et al., 2024).

Stimuli

  For the intact object condition, 540 real-world object images were sampled from the database of Brady et al. (2013). These objects are generally color-neutral in everyday-life and can be assigned any single arbitrary color (see Fig. 1A). Following Chung et al. (2023a), we rotated the stimuli in hue space along a CIE L*a*b* color wheel, so that any object could occur in any color along the wheel. On a given trial, three colored objects appeared on each side of the screen. Two of the stimuli were centered 4.1° of visual angle away from the fixation along the horizontal meridian and +/-3.5° of visual angle along the vertical meridian, while the other one was centered 5.6° away from fixation along the horizontal meridian, centered on the vertical meridian (see Fig 1B). The exact stimulus size varied slightly across objects, but they were all centrally placed in a white area that extended 5.2° in height and 5.2° in width in visual angle for the two of objects at the top and bottom locations, and 5.7° in height and 5.7° in width for the object presented along the horizontal meridian that was more distant from fixation. The colors of objects on each side of the screen were randomly selected along the 360-degree color wheel with the constraint that they had to be at least 30 degrees apart from each other. For the scrambled object condition, these object images were then morphed using the diffeomorphic scrambling technique (Stojanoski & Cusack, 2014), similar to previous work (e.g., Chung et al., 2023a;

Chung et al., 2023b; Chung et al., 2024b; Thibeault et al., 2024; Brady & Störmer, 2022). This transformation technique is particularly useful as it can make object images unrecognizable while maintaining their basic perceptual features and visual complexity. This is particularly important as some previous studies have shown CDA amplitudes may be sensitive to large perceptual differences among stimuli (e.g., Woodman & Vogel, 2008; Rajsic et al., 2018). Due to the limited number of objects sampled, some objects repeated throughout the experiments. However, none of the objects repeated more than once.

Experimental Procedure

Participants were instructed to remember the colors of three stimuli over a brief delay period. Throughout each trial, a small black fixation cross was presented in the center of a white screen and participants were instructed to maintain fixation. We used a lateralized memory display to allow the measurement of the lateralized CDA component. Thus, the memory encoding display consisted of three stimuli presented on each the left and right sides of the screen along with two black arrow cues at the center of the screen (one above and one below the fixation cross) pointing towards the to-be-remembered side (left or right). On half of the trials, colored real-world objects were presented and on the remaining half of trials, colored scrambled objects were presented. Participants were instructed to remember the colors of the stimuli on the side that the arrows pointed to while maintaining their gaze on the central fixation cross. After a 800-ms encoding period, a 800-ms delay period followed displaying a blank screen with just the central fixation cross, followed by a probe (a small black square, 0.9° height and width; 500ms) that appeared at the center of the screen, prompting participants of the beginning of the response period. Subsequently, a stimulus from the to-be-remembered side appeared in two different colors around the center of the screen (slightly shifted along the vertical meridian), and participants were asked to indicate which of the two colors they had encoded initially by pressing up- and down-arrow keys on the keyboard corresponding to the relative position of the items on the screen (top vs. bottom). Following previous work (Chung et al. 2023a), one of the stimulus choices appeared in the color that matched the encoded color (target) and the other stimulus appeared in a color 180° away from the target color on the color wheel (foil). The foil colors were also at least 30° away from the other stimulus colors on the cued side of the encoding array, ensuring no overlap in colors among stimuli on each trial. After this 2-alternative forced choice (2-AFC) task, participants received feedback in the form of a fixation cross color change (green correct, red incorrect) after which the screen turned fully white. After each trial, participants could start the next trial with a button press. The next trial then began after an inter-trial interval (ITI) of 800ms (first 19 participants) or randomly jittered 600-1,000ms (last 5 participants). Note that throughout the experiment, participants were never asked to recall the identities of the stimuli and were only asked about their colors. Thus, the experiment could be done without any regard for the identity of the stimuli. Participants completed 432 trials in total (216 trials each condition, randomly intermixed), and also went through a practice block of 10 trials prior to the main experiment. See Fig. 1B for an illustration of the trial structure.

Electrophysiological recording

EEG was recorded continuously from 32 Ag/AgCl electrodes (arranged in the 10-20 system) mounted in an elastic cap and amplified by an ActiCHamp amplifier (BrainProducts, GmbH). The horizontal electrooculogram (HEOG) was recorded from two additional electrodes positioned on the external ocular canthi which were grounded with another electrode placed on

the neck of the participant. All scalp electrodes were referenced to the right mastoid online and were digitized at 500 Hz. EEG data was filtered with a bandpass of 0.01–112.5 Hz online.

Figure 1. **A)** Example stimuli: Colored real-world objects (left) and colored scrambled objects (right). **B)** Trial structure. All stimulus sizes and distances are for illustration purposes only. **C)** Behavioral performance results based on the 24 participants used in the main EEG analysis.

Statistical Analysis

Behavioral data analysis

Performance on the color memory task was quantified as d' for a 2-AFC task (($[z$hits] – [$z$false alarms])/√2) for each participant and condition separately (intact object condition vs. scrambled object condition). The d' values were then statistically compared using a paired t-test.

EEG Analysis

All EEG data analyses were performed using EEGLAB (Delorme & Makeig, 2004) and ERPLAB (Lopez-Calderon & Luck, 2014) toolboxes and custom-written scripts. Data were epoched into trials aligned to the onset of the encoding display. Artifacts were detected using an automated pipeline and artifact rejection was performed from 200 ms before the onset of memory stimuli to 1600 ms afterwards (the end of the delay period). Trials with eye blinks (peak-to-peak at channel FP1, threshold at 150 µV), excessive eye movements (step function at HEOG, threshold at 17 µV which corresponds to ~1° of saccade; cf., Luck, 2014; Hillyard &

Galambos, 1970), and large noise (peak-to-peak all channels, threshold at 300 µV) were excluded from the analysis. The average rejection rate of participants included in the final analysis was 17.7%. Artifact-free data were re-referenced to the average of the left and right mastoids, digitally low-pass filtered at 30 Hz (Butterworth filter, 12dB/octave roll-off), and baseline corrected to the 200 ms prestimulus interval.

Parietal-occipital lateralized ERPs. We examined lateralized activity over posterior electrode sites both during the encoding (400-800 ms) and delay (1200-1600 ms) periods. Our main interest was to assess whether remembered colors of real-world objects vs. scrambled objects result in a difference in the delay period (the CDA) as this marks the active maintenance of visual representations (Vogel & Machizawa, 2004; McCollough et al., 2007; Unsworth et al., 2015; Adam et al., 2018). However, based on recent findings indicating that lateralized differences may emerge during the encoding period when using relatively long encoding times and meaningful stimuli (cf., Asp et al., 2021; Thibeault et al., 2024), we included both time intervals in our main analyses. EEG epochs were averaged separately for each object condition (intact vs. scrambled) and hemifield (remember left vs. right) and were then collapsed across to-be-remembered visual hemifield (left/right) and hemisphere of recording (left/right) to obtain waveforms recorded over the hemisphere contralateral and ipsilateral with respect to the to-be-remembered side. To quantify activity that was related to the encoding and memorization of the cued items in particular, the ipsilateral waveform was subtracted from the contralateral waveform. Thus, this difference wave reflects the increase in neural activity that's specific to the to-be-remembered items. The mean amplitude for each participant was calculated by averaging activity across four posterior-occipital electrode pairs (PO3/PO4, PO7/PO8, P7/P8, and O1/O2; Asp et al., 2021) from 400 to 800 ms post stimulus onset for the encoding period and 400 ms to 800 ms after the stimuli offset (1200-1600 ms post encoding display onset). Electrode sites and CDA time windows were chosen a-priori based on previous studies (see Roy & Faubert, 2023 for a review); the encoding period time window was chosen to avoid early visually-evoked potentials (i.e., P1/N1) and match the duration of the CDA analysis. For each time window, mean amplitudes were compared across conditions (intact vs. scrambled) using paired t-tests.

We also examined differences in the topographies across the two conditions during both encoding and delay time periods. Scalp topographies of the lateralized encoding and retention periods are plotted in Figure 2D as the contralateral-minus-ipsilateral difference waveform, projected on the right side of the scalp. To statistically compare these topographies, we first normalized the difference waveform by dividing the amplitude from each electrode site by the square root of the sum of the squared voltages across all electrodes within each condition (McCarthy & Wood, 1985) to ensure differences in overall amplitudes would not drive the differences in results. Then we ran a 2X2 ANOVA with 13 lateralized electrode pairs and two conditions (intact vs. scrambled objects) as within-subjects factors.

Cross-Temporal Pattern Similarity. To evaluate the neural stability of the memory representations for each condition, we adopted a novel analysis method (based on Spaak et al., 2017; see also Liu et al., 2020) comparing the pattern similarity of the lateralized difference waveforms across time. Specifically, separately for each condition, we computed the cosine similarities of the activity pattern of these ERP waveforms over time, smoothed over 10ms. Because this was a visual working memory task, we separated this similarity analysis into two sets of electrodes, one using a set of parietal-occipital electrodes (P7/P8, P3/P4, PO3/PO4,

PO7/PO8, O1/O2), and one using a set of fronto-central electrode (FP1/FP2, F3/F4, FC1/FC2, FC5/FC6, C3/C4). Figure 3A depicts the resulting cross-temporal analysis matrices which show the neural similarity time by time across the entire trial period, separately for posterior (top) and anterior (bottom) scalp sites. Higher cross-temporal similarity reflects more stable representations. The critical question was whether this increase in neural stability arose at different points in time for the intact vs. scrambled object condition. To test this, we collapsed the cross-temporal similarity matrices by averaging the similarity values between a given time point and the average of all later time points (i.e., mean cosine similarities of t_1 vs. t_2-t_n; t_2 vs. t_3-t_n, etc.). We then statistically compared the time courses of the averaged similarity values across conditions using a permutation approach, where all single EEG trials were shuffled randomly across conditions 10,000 times. This generated a null distribution of cross-temporal similarity matrices against which the observed effects were compared to at every time point (every 10ms).

Bilateral fronto-central bilateral ERPs. As an exploratory analysis, we also compared non-lateralized neural activity between the two experimental conditions at fronto-central electrode sites (FC1, FC2, CP1, CP2, Cz) during both the encoding and delay periods using the same time windows as for the lateralized ERPs (400-800 ms and 1200-1600 ms post stimulus onset). The mean ERP amplitudes between the two conditions were compared using a paired-t-test. Note that such global differences in bilateral activity is controlled for in the main CDA analysis as non-lateralized activity is subtracted out.

Results

Behavioral Results
We first analyzed the behavioral data including all the participants who met the behavioral inclusion criteria (all 29 participants). This revealed the expected increase in color working memory performance for the intact objects (mean d' = 1.09) compared to scrambled objects (mean d' = 0.98): $t(28) = 2.25$, $p = 0.03$, Cohen's $d_z = 0.43$, replicating previous work (Chung et al., 2023a; Chung et al., 2023b). When only including the 24 participants that were retained for the EEG analysis after artifact rejection, this effect remained numerically present though was statistically non-significant (mean d' = 1.05 for intact objects vs. 0.98 for scrambled $t(23) = 1.54$, $p = 0.14$, Cohen's $d_z = 0.32$ (see Fig. 1C).

Contralateral Encoding Activity (CEA)
All EEG analyses focused on the 24 participants who met the EEG inclusion criteria. During the encoding period (400-800ms), colors superimposed on intact objects elicited a larger contralateral negativity (mean amplitude = -0.23 µV) relative to colors superimposed on scrambled stimuli (mean amplitude = 0.24 µV): $t(23) = 5.03$, $p < 0.001$, Cohen's $d_z = 0.89$ (see Fig. 2A/2B).
Statistical analysis of the spatial topography resulted in a significant interaction between the memory conditions and electrode sites ($F(12, 276) = 2.42$, $p = 0.005$, $\eta_p^2 = 0.095$), indicating a shift in the pattern of activity across conditions. There was also a significant main effect of condition ($F(1, 23) = 5.18$, $p = 0.03$, $\eta_p^2 = 0.18$) but no main effect of electrode locations ($F(12, 276) = 1.42$, $p = 0.15$, $\eta_p^2 = 0.06$). Visual inspection suggests that the intact object condition showed a more lateralized parietal-occipital focus, whereas the scrambled object condition showed a more central-parietal focus of activity (see Fig. 2D).

Contralateral Delay Activity (CDA)

During the retention period, mean CDA amplitude (1200-1600 ms) was significantly more negative (mean amplitude = -0.90 µV) for the intact object condition than the scrambled object condition (mean amplitude = -0.54 µV): $t(23) = 2.90$, p = 0.008, Cohen's d_z = 0.59 (see Fig. 2A/2B), indicating that active maintenance activity was increased for colors superimposed on real objects at encoding.

The topographical analysis revealed a significant interaction between the memory conditions and electrode sites ($F(12, 276) = 2.003$, p = 0.02, $\eta_p^2 = 0.08$), indicating a reliable difference in the spatial distribution of activity, similar as during the encoding period. There was no main effect of condition ($F(1, 23) = 1.23$, p = 0.28, $\eta_p^2 = 0.05$), but a significant main effect of electrode location ($F(12, 276) = 3.89$, p < 0.001, $\eta_p^2 = 0.15$). Visual inspection of the spatial topographies suggests that the activity pattern shifted more towards parietal-central can temporal electrode sites for the intact object condition compared to the scrambled condition, in addition to increasing in amplitude (see Fig. 2D).

Figure 2. **A)** Contralateral-minus-ipsilateral waveforms averaged over parietal-occipital electrodes separately for intact (light blue) and scrambled (dark red) object conditions. Highlighted areas depict the time windows for the ERP analyses for the contralateral-encoding (CEA) and contralateral-delay activities (CDA). **B)** Mean CEA and CDA amplitudes separately for each condition. Error bars reflect within-subject standard errors of the mean. **C)** HEOG subtraction waveforms. Highlighted areas are the time windows for CEA and CDA analyses. **D)** Spatial topographies of the contralateral - minus - ipsilateral waveforms during the encoding and delay time periods. Mean difference is projected on the right side of the scalp. Note that the scale differs between encoding (top) and delay (bottom) time periods.

Cross-Temporal Pattern Similarity

The cross-temporal matrices for intact vs. scrambled object conditions, separately for parietal-occipital and fronto-central electrode channels, are shown in Figure 3A. Overall, as time evolves, neural stability increases across both conditions. Visual examination of these cross-temporal matrices suggests that the neural pattern became stable earlier for intact vs. scrambled objects, at least over parietal-occipital electrode sites. Our statistical analysis examining the time course of neural stability supported this, resulting in a cluster of significant differences in stability between the two conditions around 800~900 ms (see Fig 3B, top). No reliable differences in neural stability between the two conditions were found for the fronto-central electrode sites.

Figure 3. **A)** Cross-temporal stability matrices depicting cosine similarities of lateralized ERP waveform patterns across time. Matrices on top are parietal-occipital electrode sites while the bottom ones are fronto-central electrode sites. **B)** Cosine similarity values between each time window and all later time windows were averaged to compare the time course of stability across the two conditions. Generally, ERP patterns became more stabilized over time, indicated by upwards trends of the averaged similarity values. For parietal-occipital sites, the neural activity pattern for intact objects became stable earlier compared to scrambled objects, as indicated by the significant difference around 800-900ms.

HEOG

To ensure that the observed lateralized effects (e.g., CEA, CDA) were not attributable to systematic eye movement differences between conditions, we analyzed the HEOG waveforms for each cued side (left/right) across the object conditions (intact/scramble) during both the CEA and CDA time windows. Specifically, we subtracted the HEOG activity of the remember-right condition from the activity of the remember-left condition, separately for each object condition; then, we subtracted the scrambled object condition from the intact object condition. These subtracted HEOG waveforms were then tested against zero using single-sample t-tests. The results indicated that the subtracted HEOG waveforms did not significantly deviate from zero in either time window: (CEA: $t(23) = 1.07$, p = 0.29; CDA: $t(23) = 0.02$, p = 0.98; see Fig. 2C). This suggests that the observed differences in CEA and CDA across the conditions were not driven by differences in systemic eye movements.

Fronto-Central Negativity during Encoding and Delay Period
We observed a robust non-lateralized ERP difference over fronto-central electrode sites across the intact and scrambled object conditions. Specifically, the intact object condition elicited a large negative slow-wave relative to the scrambled object condition during the encoding period ($t(23) = 9.25$, p < 0.001, Cohen's $d_z = 0.69$), which sustained throughout the delay period ($t(23) = 4.06$, p < 0.001, Cohen's $d_z = 0.32$; see Fig. 4A/4B). This suggests that activity over fronto-central electrode sites tracks the meaningfulness of items at encoding and during the delay.

Figure 4. **A)** Non-lateralized waveforms recorded over fronto-central electrode sites separately for intact and scrambled objects. Highlighted areas indicate the time windows for statistical analyses. **B)** Mean ERP amplitudes during the encoding and delay time periods. **C)** Spatial topographies of the intact - minus - scrambled object conditions for each time window.

Discussion

We recently found improved working memory performance for colors when they were encoded as parts of meaningful objects, relative to colors that were superimposed on non-meaningful shapes that were otherwise matched in visual complexity (Chung et al., 2023a, Chung et al., 2023b; see also Loaiza et al., 2024). Here, we demonstrate that this performance benefit is accompanied by increases in neural delay activity and earlier-arising neural stability for intact objects compared to scrambled objects. This finding of increased neural engagement stands in contrast to alternative theories, like efficient coding of color information (which would result in reduced memory engagement), or non-active processes such as more effective retrieval at test from passive memory systems, like activated long-term memory (e.g., Cowan, 1988).

We also observed differences in the spatial topography across conditions: the CDA appeared to be more spread-out throughout parietal, central, and temporal sites in the intact object compared to the scrambled object condition that showed a more occipital focus. Such shifts in CDA topography may indicate recruitment of more task-relevant neural regions in the intact object condition during the encoding and maintenance periods (for similar shifts in CDA from working memory task relevance, see Breitinger et al., 2022). For example, it has been shown that colors of real-world objects activate not only low-level color maps (e.g., V4), but also areas in higher-level visual regions that are tuned to objects and their associated colors (e.g., Bramão et al., 2010; Lafer-Sousa et al., 2016). Thus, simple features like colors may be coded – and maintained – redundantly across the visual hierarchy, creating stronger and more stable representations. Further, associating simple features like color to real-world objects may activate additional semantic knowledge structures of these objects, effectively increasing the dimensionality of the memory representations and thereby reducing interference (Wyble, 2016). This interpretation is consistent with other recent studies showing increased neural maintenance activity and higher performance when participants remember not just simple visual features but entire identities of real-world objects (Brady et al., 2016; Thibeault et al., 2024; Asp et al., 2021). The current results go beyond these previous findings by demonstrating that increases in representational dimensionality due to links to semantic knowledge not only improve memory for the objects themselves but also provide an effective scaffold for memorizing their low-level and abstract visual features (Chung et al., 2023a; b). Overall, this suggests that expanded neural recruitment enables higher-dimensional working memory representations that support the maintenance of even simple features, either through redundant coding and/or by connecting lower-level visual features to semantic meaning in a hierarchically structured way.

Is it possible that the observed increase in CDA reflects memory for the intact objects themselves, and not their colors? Such an account would suggest that this study is unique in finding both a behavioral benefit for a condition and a CDA benefit for the same condition, but those benefits somehow being due to different causes — in direct conflict with repeated findings of a strong relationship between CDA and behavior at both the group and individual level (Vogel & Machizawa, 2004; Störmer et al., 2013, Roy & Faubert, 2022; Luria et al., 2016). This seems unlikely, especially given several previous studies demonstrating that the CDA component represents storage of task-relevant information only (Woodman & Vogel, 2008; Zhu et al., 2022). For instance, Zhu et al. (2022) showed that features that are used for the selection in working memory tasks but are not tested at retrieval are not reflected in the CDA. Similarly, Williams & Drew (2020) reported that processing of task-irrelevant real-world objects was not reflected in the CDA (Exp. 1, Williams & Drew, 2020). Thus, while it seems likely that some aspects of object information beyond color are incidentally represented in memory (Chung et al.,

2024b), the current results seem most compatible with the interpretation that the observed CDA increase is not merely due to active maintenance of objects, but instead reflects a boost in maintenance capacity for the task-relevant colors — consistent with behavioral findings that people are better at remembering colors presented on real objects even if those objects are not presented at test (Exp. 5 in Chung et al., 2023a).

The present study used carefully generated control stimuli that matched the low-level visual features and complexity of real-world objects, to ensure that observed neural differences cannot be attributed to differences in visual complexity. This is important because previous work has shown that the CDA amplitude is sensitive to the stimulus types maintained (Woodman & Vogel, 2008). However, more complex but meaningless stimuli actually resulted in lower CDA amplitudes than simple features, consistent with lower behavioral performance for such stimuli (Luria et al., 2010; Gao et al., 2009); only stimuli that are both complex and meaningful resulted in higher CDA amplitudes compared to abstract stimuli, again consistent with higher behavioral performance for such stimuli (Asp et al., 2021; Thibeault et al., 2024; Brady et al., 2016). This suggests that the CDA may reflect a broader neural marker of general memory strength in visual working memory (i.e., indexing how much information is being held in visual working memory more generally; Salahub et al., 2019; Gao et al., 2009), and is not just an assessment tool for discrete item limits (Ikkai et al., 2010; Fukuda et al., 2010; Balaban et al., 2019; Hakim et al., 2019).

We observed earlier neural stabilization of memories in the real-world object condition. This is consistent with the contralateral negativity also emerging earlier for real-world objects. In particular, prior to the interval classically studied as the CDA, we observed a larger contralateral negativity during the encoding period already. This matches other recent studies that also reported lateralized EEG differences emerging during encoding when comparing working memory for meaningful vs. non-meaningful stimuli (Thibeault et al., 2024; Asp et al., 2021). This could reflect faster and more accurate encoding of the stimuli into working memory, as well as limits in how much information can be extracted and maintained during perception (Tsubomi et al., 2013; Balaban & Luria, 2019; for a review see Emrich, 2022). The finding that the condition differences appeared early also relates to recent behavioral evidence indicating that in contrast to earlier claims in the literature of no effects of encoding time (Luck & Vogel, 1997; Alvarez & Cavnagh, 2004), new work has shown that encoding format and time can play a critical role in working memory processes for simple stimuli (Schurgin et al., 2020; Quirk et al., 2020) and meaningful objects (Brady et al., 2016; Brady & Störmer, 2023; Chung et al, 2023b). The present data provide further support that memory capacity depends on limits that can be observed even when the stimuli are still on the screen – and that real-world objects can provide a useful structure to initially build, and then maintain, robust representations.

Finally, our study also reveals a novel neural signature that appears to track processes related to recognizing an intact vs. scrambled object. Beginning during encoding and continuing throughout the delay period, we observed a large non-lateralized slow-wave over fronto-central electrode sites that was more negative for the intact vs. scrambled condition. This sustained negativity for recognizable vs. completely scrambled objects resembles an ERP finding in the word reading literature: the N400, a well-known marker related to semantic understanding (Kutas & Hillyard, 1980), is more negative for pseudowords compared to words (as well as semantically inconsistent vs. consistent words), but interestingly, has also been reported to be more *positive* for fully scrambled nonwords (Bentin et al., 1999; Ziegler et al., 1997; Coch & Mitra, 2010). This is consistent with our results showing that fully scrambled objects result in a

relative positivity compared to intact objects. This suggests that amplitude changes in the N400 component, traditionally linked to word processing, may reflect broader cognitive processes that distinguish between semantically meaningful versus semantically inconsistent and, critically, also fully scrambled stimuli across domains (verbal and visual). While further research is necessary to fully understand what cognitive processes underlie the observed amplitude differences between intact vs. scrambled objects over fronto-central sites, they appear to robustly distinguish between these stimulus conditions. [1]

In conclusion, our results show that working memory for simple features is shaped by the context these features are encoded in. Meaningful context at encoding – being part of a real-world object – promotes stronger neural engagement and earlier temporal stability of memory representations during encoding and maintenance processes. Broadly, this indicates that working memory processes, and consequently its capacity, are flexible, even with respect to a single feature dimension.

---

[1] A similar pattern of non-lateralized ERP result was observed in a previous study investigating visual working memory of real-world objects and scrambled objects (Thibeault et al., 2024; personal correspondence).

References

Adam, K. C., Robison, M. K., & Vogel, E. K. (2018). Contralateral delay activity tracks fluctuations in working memory performance. Journal of Cognitive Neuroscience, 30(9), 1229-1240.

Adam, K. C., Vogel, E. K., & Awh, E. (2017). Clear evidence for item limits in visual working memory. Cognitive psychology, 97, 79-97.

Alvarez, G. A., & Cavanagh, P. (2004). The capacity of visual short-term memory is set both by visual information load and by number of objects. Psychological science, 15(2), 106-111.

Asp, I. E., Störmer, V. S., & Brady, T. F. (2021). Greater visual working memory capacity for visually matched stimuli when they are perceived as meaningful. Journal of cognitive neuroscience, 33(5), 902-918.

Awh, E., & Vogel, E. K. (2024). Working memory needs pointers. Trends in Cognitive Sciences. Bays, P. M., Catalao, R. F., & Husain, M. (2009). The precision of visual working memory is set by allocation of a shared resource. Journal of vision, 9(10), 7-7.

Baddeley, A. D., & Hitch, G. J. (1974). Working memory. In G. H. Bower (Ed.), Psychology of learning and motivation, (8), 47-89.

Balaban, H., Drew, T., & Luria, R. (2019). Neural evidence for an object-based pointer system underlying working memory. Cortex, 119, 362-372.

Balaban, H., & Luria, R. (2019). Using the contralateral delay activity to study online processing of items still within view. In Spatial learning and attention guidance (pp. 107-128). New York, NY: Springer US.

Bays, P. M., Schneegans, S., Ma, W. J., & Brady, T. F. (2024). Representation and computation in visual working memory. Nature Human Behaviour, 8(6), 1016-1034.

Bays, P. M. (2014). Noise in neural populations accounts for errors in working memory. Journal of Neuroscience, 34(10), 3632-3645.

Bentin, S., Mouchetant-Rostaing, Y., Giard, M. H., Echallier, J. F., & Pernier, J. (1999). ERP manifestations of processing printed words at different psycholinguistic levels: time course and scalp distribution. *Journal of cognitive neuroscience*, *11*(3), 235-260.

Brady, T. F., & Störmer, V. S. (2022). The role of meaning in visual working memory: Real-world objects, but not simple features, benefit from deeper processing. Journal of Experimental Psychology: Learning, Memory, and Cognition, 48(7), 942.

Brady, T. F., & Störmer, V. S. (2023). Comparing memory capacity across stimuli requires maximally dissimilar foils: Using deep convolutional neural networks to understand visual working memory capacity for real-world objects. Memory & Cognition, 52(3), 595-609.

Brady, T. F., Störmer, V. S., & Alvarez, G. A. (2016). Working memory is not fixed-capacity: More active storage capacity for real-world objects than for simple stimuli. Proceedings of the National Academy of Sciences, 113(27), 7459-7464.

Bramão, I., Faísca, L., Forkstam, C., Reis, A., & Petersson, K. M. (2010). Cortical brain regions associated with color processing: An FMRI study. The open neuroimaging journal, 4, 164.

Breitinger, E., Pokorny, L., Biermann, L., Jarczok, T. A., Dundon, N. M., Roessner, V., & Bender, S. (2022). What makes somatosensory short-term memory maintenance effective? An EEG study comparing contralateral delay activity between sighted participants and participants who are blind. NeuroImage, 259, 119407.

Buschman, T. J., Siegel, M., Roy, J. E., & Miller, E. K. (2011). Neural substrates of cognitive capacity limitations. Proceedings of the National Academy of Sciences, 108(27), 11252-11255.

Chung, Y. H., Brady, T. F., & Störmer, V. S. (2023a). No fixed limit for storing simple visual features: Realistic objects provide an efficient scaffold for holding features in mind. Psychological Science, 34(7), 784-793.

Chung, Y. H., Brady, T. F., & Störmer, V. S. (2023b). Sequential encoding aids working memory for meaningful objects' identities but not for their colors. Memory & Cognition, 52(8), 2119-2131.

Chung, Y. H., Brady, T. F., & Störmer, V. S. (2024a). Meaningfulness and familiarity expand visual working memory capacity. Current Directions in Psychological Science, 33(5), 275-282.

Chung, Y. H., Tam, J., Wyble, B., & Störmer, V. S. (2024b). Conceptual information of meaningful objects is stored incidentally. Journal of Experimental Psychology: Learning, Memory, and Cognition.

Clark, J. H. (1924). The Ishihara test for color blindness. American Journal of Physiological Optics.

Coch, D., & Mitra, P. (2010). Word and pseudoword superiority effects reflected in the ERP waveform. Brain research, 1329, 159-174.

Cowan, N. (2001). The magical number 4 in short-term memory: A reconsideration of mental storage capacity. Behavioral and Brain Sciences, 24(1), 87–114. doi:10.1017/S0140525X01003922

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. Journal of neuroscience methods, 134(1), 9-21.

Druzgal, T. J., & D'Esposito, M. (2001). Activity in fusiform face area modulated as a function of working memory load. Cognitive Brain Research, 10(3), 355-364.

Druzgal, T. J., & D'esposito, M. (2003). Dissecting contributions of prefrontal cortex and fusiform face area to face working memory. Journal of cognitive neuroscience, 15(6), 771-784.

Emrich, S. M., Salahub, C., & Katus, T. (2022). Sensory delay activity: more than an electrophysiological index of working memory load. Journal of Cognitive Neuroscience, 35(1), 135-148.

Fukuda, K., Awh, E., & Vogel, E. K. (2010). Discrete capacity limits in visual working memory. Current opinion in neurobiology, 20(2), 177-182.

Gao, Z., Li, J., Liang, J., Chen, H., Yin, J., & Shen, M. (2009). Storing fine detailed information in visual working memory—Evidence from event-related potentials. Journal of vision, 9(7), 17-17.

Hakim, N., Adam, K. C., Gunseli, E., Awh, E., & Vogel, E. K. (2019). Dissecting the neural focus of attention reveals distinct processes for spatial attention and object-based storage in visual working memory. Psychological science, 30(4), 526-540.

Hillyard, S. A., & Galambos, R. (1970). Eye movement artifact in the CNV. Electroencephalography and clinical neurophysiology, 28(2), 173-182.

Ikkai, A., McCollough, A. W., & Vogel, E. K. (2010). Contralateral delay activity provides a neural measure of the number of representations in visual working memory. Journal of neurophysiology, 103(4), 1963-1968.

Kriegeskorte, N., Mur, M., & Bandettini, P. A. (2008). Representational similarity analysis-connecting the branches of systems neuroscience. Frontiers in systems neuroscience, 2, 249.

Kutas, M., & Hillyard, S. A. (1980). Reading senseless sentences: Brain potentials reflect semantic incongruity. Science, 207(4427), 203-205.

Lafer-Sousa, R., Conway, B. R., & Kanwisher, N. G. (2016). Color-biased regions of the ventral visual pathway lie between face-and place-selective regions in humans, as in macaques. Journal of Neuroscience, 36(5), 1682-1697.

Liu, J., Zhang, H., Yu, T., Ni, D., Ren, L., Yang, Q., ... & Xue, G. (2020). Stable maintenance of multiple representational formats in human visual short-term memory. Proceedings of the National Academy of Sciences, 117(51), 32329-32339.

Loaiza, V. M., Cheung, H. W., & Goldenhaus-Manning, D. T. (2024). What you don't know can't hurt you: Retro-cues benefit working memory regardless of prior knowledge in long-term memory. Psychonomic Bulletin & Review, 31(3), 1-12.

Lopez-Calderon, J., & Luck, S. J. (2014). ERPLAB: an open-source toolbox for the analysis of event-related potentials. Frontiers in human neuroscience, 8, 213.

Luck, S. J. (2014). An introduction to the event-related potential technique. MIT press.

Luck, S. J., & Vogel, E. K. (1997). The capacity of visual working memory for features and conjunctions. Nature, 390(6657), 279-281.

Luck, S. J., & Vogel, E. K. (2013). Visual working memory capacity: from psychophysics and neurobiology to individual differences. Trends in cognitive sciences, 17(8), 391-400.

Luria, R., Balaban, H., Awh, E., & Vogel, E. K. (2016). The contralateral delay activity as a neural measure of visual working memory. Neuroscience & Biobehavioral Reviews, 62, 100-108.

Luria, R., Sessa, P., Gotler, A., Jolicœur, P., & Dell'Acqua, R. (2010). Visual short-term memory capacity for simple and complex objects. Journal of cognitive neuroscience, 22(3), 496-512.

McCollough, A. W., Machizawa, M. G., & Vogel, E. K. (2007). Electrophysiological measures of maintaining representations in visual working memory. Cortex, 43(1), 77-94.

McCarthy, G., & Wood, C. C. (1985). Scalp distributions of event-related potentials: an ambiguity associated with analysis of variance models. Electroencephalography and Clinical Neurophysiology/Evoked Potentials Section, 62(3), 203-208.

Postle, B. R., Druzgal, T. J., & D'Esposito, M. (2003). Seeking the neural substrates of visual working memory storage. Cortex, 39(4-5), 927-946.

Quirk, C., Adam, K. C., & Vogel, E. K. (2020). No evidence for an object working memory capacity benefit with extended viewing time. Eneuro, 7(5).

Rajsic, J., Burton, J. A., & Woodman, G. F. (2019). Contralateral delay activity tracks the storage of visually presented letters and words. Psychophysiology, 56(1), e13282.

Roy, Y., & Faubert, J. (2023). Is the Contralateral Delay Activity (CDA) a robust neural correlate for Visual Working Memory (VWM) tasks? A reproducibility study. Psychophysiology, 60(2), e14180.

Salahub, C., Lockhart, H. A., Dube, B., Al-Aidroos, N., & Emrich, S. M. (2019). Electrophysiological correlates of the flexible allocation of visual working memory resources. Scientific Reports, 9(1), 19428.

Schurgin, M. W., Wixted, J. T., & Brady, T. F. (2020). Psychophysical scaling reveals a unified theory of visual memory strength. Nature human behaviour, 4(11), 1156-1172.

Shin, H., & Ma, W. J. (2017). Visual short-term memory for oriented, colored objects. Journal of Vision, 17(9), 12-12.

Störmer, V. S., Li, S. C., Heekeren, H. R., & Lindenberger, U. (2013). Normative shifts of cortical mechanisms of encoding contribute to adult age differences in visual–spatial working memory. Neuroimage, 73, 167-175.

Spaak, E., Watanabe, K., Funahashi, S., & Stokes, M. G. (2017). Stable and dynamic coding for working memory in primate prefrontal cortex. Journal of neuroscience, 37(27), 6503-6516.

Thibeault, A. M., Stojanoski, B., & Emrich, S. M. (2024). Investigating the effects of perceptual complexity versus conceptual meaning on the object benefit in visual working memory. Cognitive, Affective, & Behavioral Neuroscience, 24(3), 453-468.

Torres, R. E., Duprey, M. S., Campbell, K. L., & Emrich, S. M. (2024). Not all objects are created equal: The object benefit in visual working memory is supported by greater recollection-like memory, but only for memorable objects. Memory & Cognition, 1-13.

Tsubomi, H., Fukuda, K., Watanabe, K., & Vogel, E. K. (2013). Neural limits to representing objects still within view. Journal of Neuroscience, 33(19), 8257-8263.

Unsworth, N., Fukuda, K., Awh, E., & Vogel, E. K. (2015). Working memory delay activity predicts individual differences in cognitive abilities. Journal of Cognitive Neuroscience, 27(5), 853-865.

Vogel, E. K., & Machizawa, M. G. (2004). Neural activity predicts individual differences in visual working memory capacity. Nature, 428(6984), 748-751.

Williams, L. H., & Drew, T. (2021). Maintaining rejected distractors in working memory during visual search depends on search stimuli: Evidence from contralateral delay activity. Attention, Perception, & Psychophysics, 83, 67-84.

Woodman, G. F., & Vogel, E. K. (2008). Selective storage and maintenance of an object's features in visual working memory. Psychonomic bulletin & review, 15(1), 223-229.

Wyble, B., Swan, G., & Callahan-Flintoft, C. (2016). Measuring visual memory in its native. *Trends in Cognitive Sciences*, *20*(11), 790-791.

Zhu, P., Fu, Y., Wyble, B., Shen, M., & Chen, H. (2022). A new aspect of cognitive selectivity: Working memory reselection for attended information. PsyArxiv.

Ziegler, J. C., Besson, M., Jacobs, A. M., Nazir, T. A., & Carr, T. H. (1997). Word, pseudoword, and nonword processing: A multitask comparison using event-related brain potentials. *Journal of Cognitive Neuroscience*, *9*(6), 758-775.

Acknowledgements

Special thanks to Sadye Law, our research assistant who helped with initial data collection and pilot experiments, and Alison Sasaki and Sam (Seho) Jung who helped finish data collection.

Supplemental Material

Analyses including all 29 participants
For completeness, we have run the same main analysis for CDA amplitudes including all 29 participants. The 5 participants who were excluded for main analyses had on average 54.12% of trials excluded due to EEG artifacts. Same pattern of results emerged with all 29 participants: the CDA amplitude was larger for the intact object condition than scrambled object condition ($t(28) = 2.64$, $d_z = 0.45$, $p = 0.01$).