Whole-brain EEG dynamics depending on the stimulus modality and task

requirements in oddball tasks

Shiho Kashihara[1], Tomohisa Asai[1], & Hiroshi Imamizu[1,2]

[1] Department of Cognitive Neuroscience, Cognitive Mechanisms Laboratories, Advanced

Telecommunications Research Institute International (ATR), Keihanna Science City, Kyoto

619-0288, Japan.

[2] Department of Psychology, Graduate School of Humanities and Sociology and Faculty of

Letters, The University of Tokyo, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8654, Japan.

* Corresponding author: Shiho Kashihara, Department of Cognitive Neuroscience, Cognitive

Mechanisms Laboratories, Advanced Telecommunications Research Institute International

(ATR), Keihanna Science City, Kyoto 619-0288, Japan. TEL: +81-774-95-2685, FAX: +81-

774-95-1236, E-mail: skashihara@atr.jp

ORCID: Shiho Kashihara (0000-0002-5443-9495), Tomohisa Asai, Hiroshi Imamizu

19   **<u>Abstract</u>**

Electroencephalography (EEG) microstates constitute temporal map configurations

that reflect the whole brain electrical state. The dynamics of EEG microstates may serve as

an effective discretization method for capturing spatiotemporally continuous neural dynamics

with high temporal resolution. In this study, we employed polarity-sensitive microstate

analysis to investigate whole-brain state transitions during audiovisual oddball tasks.

Moreover, we examined how sensory modality and its coupling, types of response to target

stimuli, and the physical presence or absence of target stimuli affected EEG dynamics. The

results demonstrated that the abovementioned factors affected both behavioral indices and

the event-related potential (ERP) components, particularly the P300. Importantly, when

considering the topographical polarity of map configuration, transitions to microstate E-,

which originates within 300 to 600 ms after stimulus onset and coincides with the typical

latency of the P300 component potentially reflect attentional and conscious processes that

are associated with the P300. These novel insights into the dynamic transitions of whole

brain states during cognitive processes complement the results of traditional ERP analyses.

Keywords: EEG microstates, state transition, whole-brain dynamics, oddball task, stimulus

modality

Statements and Declarations

Funding: SK and TA were supported by Innovative Science and Technology Initiative for Security Grant Number JPJ004596, ATLA, Japan. SK, TA and HI were supported by JST [Moonshot R&D][Grant Number JPMJMS2291].

Conflict of interest: The authors declare that they have no conflict of interest.

Ethics approval: The study was approved by the Ethical Committee of ATR (approval numbers: 21-144 and 21-143).

Availability of data and material: Data are available from the authors upon reasonable request.

Acknowledgments: We are grateful to Yuki Inoue, Yoshiko Umemoto, and Ayako Oura for their assistance with data collection.

Authors' contributions: SK: Conceptualization, Methodology, Data curation, Investigation, Formal analysis, Visualization, Writing – original draft; TA: Conceptualization, Methodology, Validation, Writing – review and editing; HI: Project Administration, Supervision, Writing – review and editing. All authors read and approved the final manuscript.

1. <u>Introduction</u>

Cognitive functions, such as recognizing and responding to external stimuli, are

mediated by neural activity in the brain. Traditional approaches in cognitive neuroscience,

known as the "Sherringtonian view," attributed specific cognitive functions to localized brain

areas and circuits (Barack & Krakauer, 2021). However, the "Hopfieldian view" or "population

doctrine" constitutes a paradigm shift wherein whole-brain dynamics and neural state-space

transformations explain cognition (Barack & Krakauer, 2021; Ebitz & Hayden, 2021). Studies

using functional magnetic resonance imaging (fMRI) and intracranial electrodes described

cognition in terms of neural state-space transitions (Iyer et al., 2022; Mau et al., 2020;

Pessoa, 2019; Shine et al., 2019); however, noninvasive methods with high temporal

resolution, such as scalp EEG, offer promising alternative approaches for investigating the

neural dynamics that underlie cognition.

EEG microstate analysis provides a promising method for capturing whole-brain

neural activity with high temporal resolution (e.g., Khanna et al., 2015; D. Lehmann et al.,

1987; Michel & Koenig, 2018; Mishra et al., 2020). EEG microstates are quasi-stable electric

field patterns in scalp EEG that reflect the globally synchronized activity of large neural

networks (D. Lehmann et al., 1987; Michel & Koenig, 2018). Frequently referred to as the

"atoms of thought" (Dietrich Lehmann, 1990), EEG microstates have been extensively

studied in relation to psychiatric disorders, consciousness, and cognition.

To explore the functional role of EEG microstates, studies that used simultaneous

EEG-fMRI and EEG source localization linked specific microstates to distinct brain networks

(Al Zoubi et al., 2022; Britz et al., 2010; Musso et al., 2010; Yuan et al., 2012). Furthermore,

researchers have investigated whether particular microstates emerge during cognitive tasks

and how task characteristics influence microstate dynamics (Bréchet et al., 2019; D'Croz-

Baron et al., 2021; Kim et al., 2021; Seitzman et al., 2017). For instance, Milz et al. (2016)

found that microstate A and B (msA and msB) parameters increased during visualization and

verbalization tasks, respectively. More recently, D'Croz-Baron et al. (2021) analyzed EEG

topographical maps during auditory and visual encoding and recognition tasks, and reported

the increased occurrence of msB following visual stimulus encoding. However, the

relationship between specific microstates and cognitive functions remains inconsistent and is

sometimes contradictory, which leaves their precise role unclear.

Previous studies have primarily focused on describing conditional differences in the

discrete, static properties of EEG microstates, such as their frequency of occurrence and

duration. However, recent research has challenged the discrete nature of microstates and

emphasized the importance of considering the continuous dynamics that underlie EEG

microstate sequences (Asai et al., 2023; Kashihara et al., 2024; Michel & Koenig, 2018;

Mishra et al., 2020; Shaw et al., 2019; Sikka et al., 2020; Van de Ville et al., 2010). Despite

the growing body of knowledge on discrete microstate properties and the recognition of their

continuous dynamics, the dynamic patterns of EEG microstate transition that emerge in

response to external stimuli remain poorly understood. In contrast, Tamano et al. (2022)

reported microstate dynamics during an N-back task and identified the relationship of certain

transition probabilities comprising polarity with performance on the N-back task. However,

these authors did not consider topographical polarity a better discretization method,

assuming the continuity of EEG dynamics, and the significance of polarity consideration was

limited. In this study, we address the knowledge gap indicated by this issue by using a

polarity-sensitive microstate analysis and focusing on their transitions, while considering the

continuous trajectory of the essentially continuous EEG dynamics in the state space

(Kashihara et al., 2024).

This study aimed to investigate EEG dynamics during cognitive tasks from the

perspective of whole-brain state transitions while considering the spatiotemporal continuity of

EEG activity. Instead of the traditional Sherringtonian approach of event-related potentials

(ERPs), we sought to elucidate the neural basis of cognition through whole-brain state

transition dynamics in the neural state space. To achieve this, we introduced a polarity-

sensitive microstate analysis (Kashihara et al., 2024) to descriptively analyze the stimulus-

locked whole-brain state dynamics. Specifically, we extended the conventional ERP

microstate approach (Kindler et al., 2011; Kondakor et al., 1997; D. Lehmann et al., 1994;

Pizzagalli et al., 2000; Ruggeri et al., 2019) by incorporating topographical polarity, and

thereby enabled a more precise characterization of state transitions in EEG activity. Previous

studies on ERP microstates have examined how pre-stimulus topographical configurations

influenced responses in oddball tasks (Kondakor et al., 1997; D. Lehmann et al., 1994).

Research conducted in recent years has elucidated EEG microstate topography sequences

that reflect the conflict-monitoring processes during the Stroop task (Ruggeri et al., 2019)

and EEG microstate dynamics that predict working memory performance during the N-back

tasks (Tamano et al., 2022). By incorporating topographical polarity, our approach enables

the appropriate discretization of stimulus-locked, inherently continuous spatiotemporal EEG

transitions while preserving trajectory alignment in the neural state-space based on

topographical dissimilarity. This refinement enhances the ability to more accurately capture

the dynamics of overlapping cognitive processes.

Therefore, we decided to use an oddball task as a cognitive task. The oddball

paradigm, in which two or more stimulus events are presented at different frequencies, is a

well-known procedure in ERP studies because it allows the measurement of neural

processes related to stimulus perception and internal processing. In particular, among the

salient ERP components that are elicited by this task, P300 – a peak that appears in the

positive direction approximately 300 ms after target-stimulus presentation – has been used

to retrieve endogenous mental activity. Furthermore, P300 may represent brain activity

during the modification of a model of the environment, that is, during a contextual update

(Donchin & Coles, 1988). Thus, by using the oddball task, wherein the association of ERP

responses and cognitive processes has been well examined, clues to examine the functional

role of EEG microstate dynamics may be found.

Taken together, this study aims to clarify the functional role of continuous whole-brain

state dynamics during task performance by designing the task features of the oddball tasks

in accordance with the findings accumulated in previous ERP studies. Therefore, we

conducted experiments wherein several factors that influence the P300 component were

manipulated to examine differences in microstate responses. Specifically, the following three

points were investigated: the influence of the sensory modality of the stimulus (Experiment

1), the influence of the response format to the target (Experiment 2), and the influence of the

presence or absence of physical stimuli on the microstate dynamics of the target

(Experiment 3).

First, as a factor that affects the P300, the sensory modality of the stimulus (auditory

or visual) affects the amplitude and latency of P300 (Polich & Heine, 1996). Furthermore,

when the standard and target stimuli are intermodal (e.g., visual standard and auditory

target), the late ERP component, including P3, increased, as compared to that in the

unimodal case (Brown et al., 2006, 2007). Considering the possibility that sensory modality-

specific processing may be associated with specific classes of EEG microstates (e.g., Britz

et al., 2010), it is possible that the effects of the stimulus modality itself and its coupling with

stimulus conditions can also be ascertained in the emerged EEG microstate classes or

dynamics even when examining continuous whole-brain state transitions. Therefore, we

prepared oddball tasks with different stimulus modalities and coupled them with conditions

(unimodal auditory task, i.e., auditory target–auditory standard; unimodal visual task, i.e.,

visual target–visual standard; inter-modal auditory-target task, that is, the auditory target–

visual standard; intermodal visual-target task, that is, visual target–auditory standard). First,

we examined the effects of the modalities and their coupling on EEG microstate dynamics

(in Experiment 1).

Second, we examined the effect of the response format required for the target

stimulus. In the context of ERP studies, the P3 amplitude or latency may differ depending on

the response format, such as whether the target stimulus requires a key press response or

mental count. The findings pertaining to the effect of the response mode on ERPs are

confusing. Some studies reported no difference between key-press and count for the target

stimuli (Starr et al., 1995), whereas others reported that the P3 in the key-press task was

greater than that in the counting task (Brázdil et al., 2003; Kotchoubey, 2014); moreover,

there were even contradictory results (Barrett et al., 1987; Salisbury et al., 2001). As some

findings suggest that motor behavior affects the discrete parameters of the EEG microstate

(Croce et al., 2022), we examined the effects of the response format on EEG microstate

dynamics by establishing a count-response group that can isolate the influence of motor

behavior compared to key presses (Experiment 1 vs. Experiment 2).

Third, we examined the effect of the presence/absence of physical target stimuli. P3

potentially reflects an endogenous component and appears even when the target stimulus is

not physically present (Ragazzoni et al., 2019). By setting up a task that requires a key press

when a regularly presented physical stimulus is missing (the omitted-target oddball task), we

can isolate the influence of the physical properties of the target stimulus and examine how

purely endogenous processes of the target can be involved in continuous whole-brain state

transitions (Experiment 1 vs. Experiment 3).

178   **2. Methods**

2-1. Participants

Three experiments involved 66 participants, and there was no duplication of

participants. All experiments were approved by the ethics committee of the Advanced

Telecommunications Research Institute International, Kyoto, Japan (approval numbers 21-

144 and 21-143). All participants provided written informed consent before the experiment.

2-1-1. Experiment 1

Twenty participants (12 females, 8 males; age, mean [SD]: 30.7 [6.9] years)

participated in Experiment 1 and declared that they were right-handed. As no previous study

has ascertained the EEG microstate transition indices during the oddball task, we

determined the number of participants under an expected moderate effect size (Cohen's d =

0.45) based on the robustness of the neural activity elicited by the oddball task (e.g., P300

ERP component). Using PANGEA v0.2 (https://jakewestfall.shinyapps.io/pangea/)(Power

ANalysis for GEneral Anova designs, Westfall, 2016), we calculated the sample size

required for detecting the interaction between stimulus modality and condition when

assuming the experimental design of 2 (modality coupling: unimodal, inter-modal) ×2 (target

stimulus modality: auditory, visual) ×2 (stimulus condition: target, standard) with the factor of participants as a random effect. We considered the repetition of stimulus presentations 80 times (in the target condition). Accordingly, PANGEA suggested 15 participants for power (1-β) > 0.80 with Cohen's d = 0.45. We recruited 20 participants, considering the possibility of missing data.

2-1-2. Experiment 2

The number of participants in Experiment 2 was determined according to the sample size in Experiment 1. Twenty-one participants (11 females and 10 males; age, mean [SD]: 24.8 [5.3] years) participated in Experiment 2. All participants were right-handed and had never participated in Experiments 1 and 3. One participant was excluded because of refusal to participate; therefore, data from 20 participants (10 females and 10 males; age, mean [SD]: 25.0 [5.3] years) were included in the analysis.

2-1-3. Experiment 3

In total, 21 participants (11 males and 10 females; age, mean [SD]: 22.6 [4.2] years) were included in Experiment 3, were all right-handed, and had never participated in Experiments 1 and 3. Two participants were excluded because one refused to participate and the other experienced recording failure. Therefore, data from 19 participants (nine males and 10 females; age, mean [SD]: 22.9 [4.4] years) were used in the subsequent analyses.

2-2. Apparatus

Stimuli were presented on a 24-inch color LCD monitor in all three experiments (FlexScan EV 2456, EIZO, Japan), and responses were recorded via a computer keyboard in Experiments 1 and 3 (Realforce91U, Topre Corporation, Japan). The monitor (resolution: 1920 × 1200) was situated approximately 55 cm in front of the seated participants. The size of visual stimuli was approximately 480x480 pixels. Stimulus presentation and response collection were controlled by PsychoPy v2021.1.4 (J. Peirce et al., 2019; J. W. Peirce, 2007, 2008).

2-3. Tasks and materials

We conducted audiovisual oddball tasks in the three experiments with different task

formats. In the task, two types of stimuli with different frequencies of the presentation were

shown sequentially (high-frequency: standard, low-frequency: target). Stimuli were

presented randomly, with the restriction that the target stimuli were not presented

consecutively. Participants were asked to respond to a low-frequency stimulus (target) as

quickly and accurately as possible; that is, they were to press the space key when the target

is presented (Experiment 1), keep counting the number of target presentations in their mind

(Experiment 2), and press the space key when the stimulus was absent (Experiment 3). In

Experiment 3, only the standard stimulus was perceptible, and the target stimulus was a

"missing target" that was physically absent. In general, a total of 400 trials were presented in

each task. The rate of the presentation of the standard and target stimuli was 8:2. Therefore,

the number of standard and target trials were 320 and 80, respectively. Each stimulus was

presented for 200 ms, with a 1000-ms inter-stimulus interval.

2-3-1. Experiments 1 and 2

In Experiments 1 and 2, there were four tasks based on the target stimulus modality

(auditory or visual) and the modality coupling between target and standard stimulus

(unimodal or inter-modal) as follows: unimodal-auditory oddball task, unimodal-visual oddball

task, inter-modal auditory-target oddball task, and inter-modal visual-target oddball task

(Figure 1). The order of tasks was partially counterbalanced; two unimodal tasks were

conducted, followed by the inter-modal tasks, and the modality of the target stimulus was

counterbalanced. Participants were asked to either press the space key (Experiment 1) or

count the number of target stimuli in their minds (Experiment 2) when the target stimulus

was presented.

The experimental materials were 1000- and 2000-Hz pure tone for auditory stimuli,

as well as a black circle (●) and a black star (□) for visual stimuli. The assignment of

materials to stimulus conditions (i.e., standard or target condition) was fixed in Experiment 1

(auditory target: 2000 Hz, auditory standard: 1000 Hz, visual target: black star, visual

standard: black circle). To offset the influence of materials, we counterbalanced the

assignment of materials to conditions across participants in Experiment 2: for half of the

participants (N = 10), the auditory target was a 2000-Hz tone, and the visual target was a

black star; for the other half (N = 10), the auditory target was a 1000-Hz tone, and the visual

target was a black circle.

2-3-2. Experiment 3

The tasks in Experiment 3 were almost the same as those in Experiments 1 and 2.

However, because the target stimulus was not physically present in Experiment 3 ("missing

target"), there were only two unimodal tasks according to the standard stimulus modality

(auditory or visual), and participants perceived only standard stimuli. The order of tasks was

counterbalanced. In the tasks, auditory standard (1000 Hz) or visual standard stimuli (black

circle) were presented at regular intervals, sometimes with missing stimuli (target; 20% of all

trials). Participants were asked to press a space key when they noticed the missing target.

Figure 1. The overview of trials of each of the four types of oddball tasks.

In each experiment, four different unimodal/inter-modal, auditory/visual tasks were performed with a combination of stimulus modalities (auditory or visual) for each stimulus with varying presentation frequencies (standard or target). In Experiment 3, only two unimodal tasks were performed because no physical stimulus was presented in the target trials. For each task, a total of 400 trials (320 standard and 80 target trials) of two blocks (200 trials, including 160 standard and 40 target trials) were presented. After finishing the first block, we asked the participants to take a break for an arbitrary time. Each stimulus was presented for 200 ms. The inter-stimulus interval was 1000ms. The order of stimulus presentation was determined pseudo-randomly for each participant under the condition that the target stimulus never appeared twice in a row. Participants were asked to press the space key as quickly and accurately as possible when the target stimulus was presented (Experiment 1), to count the number of times the target stimulus was presented (Experiment 2), or to press the space key as quickly and accurately as possible when they noticed a target trial where the stimulus was not presented (Experiment 3). We recorded the reaction time and correct/incorrect responses.

2-4. Procedure

The general procedure was common to all three experiments, except for what

participants were to do during the oddball tasks. Participants wore an EEG cap, and the

resting-state spontaneous EEG was recorded, for 5 minutes each, with eyes open and eyes

closed. The participants sat in a comfortable chair and were asked not to think too deeply

about certain things and to stay at rest—the order in which the eye-open and eye-closed

resting recordings were counterbalanced. In eye-open resting EEG recording, the fixation

cross appeared at the center of the monitor, and participants were asked to gaze at it. In the

eyes-closed rest condition, participants were asked to keep their eyes closed throughout the

recording and not to fall asleep.

Next, participants performed four oddball tasks; auditory or visual unimodal oddball

tasks and inter-modal oddball tasks with auditory or visual targets. Inter-modal tasks were

conducted after unimodal tasks, and the modalities were counterbalanced for both task

presentation types. In one task, 400 trials were presented, with a break every 200 trials. The

experiment lasted approximately 2 hours, from start to finish (Experiment 3 had two fewer

tasks, but the time required for the experiment was the same as that for Experiments 1 and 2

because other tasks were performed that were not directly related to this study's purpose).

Participants were given different instructions regarding task performance in each of the three

experiments. In Experiment 1, participants were asked to press a space key as quickly and

accurately as possible with either the index or middle finger of their right (i.e., dominant)

hand when the target stimulus was presented. In Experiment 2, participants were asked to

mentally count the number of target presentations and instructed not to speak aloud or use

their fingers or bodies during counting. To increase engagement with the task, they were

asked to answer the number of target presentations for that block every 200 trials. In

Experiment 3, participants were given the same instruction as for Experiment 1; however,

the target stimulus was not physically presented ("missing target"), and participants were

asked to press the key when they noticed that stimuli presented at regular intervals in the

target trials were missing.

2-5. EEG recording and preprocessing

EEGs were recorded in all three experiments using 32ch R-net (Brain Products

GmbH, Gilching, Germany), silver–silver chloride electrodes attached to a saltwater sponge-

based cap. A total of 32 electrodes were placed on the scalp based on the international 10-

10 systems: that is, at Fp1, Fp2, Fz, F3, F4, F7, F8, F9, F10, FC1, FC2, FC5, FC6, Cz, C3,

C4, T7, T8, CP1, CP2, CP5, CP6, Pz, P3, P4, P7, P8, P9, P10, Oz, O1, and O2. The ground

and reference electrodes were placed at Fpz and FCz, respectively, and were off-line re-

referenced to the average potential. The impedances were maintained under 50 kΩ. EEG

signals were recorded at 500 Hz (Experiment 1) or 5000 Hz (Experiments 2 and 3) using an

EEG amplifier (BrainAmp MR, Brain Products GmbH, Gilching, Germany) with an online

bandpass of 0.016–250 Hz.

EEG data were preprocessed using an EEGLAB toolbox (EEGLAB v2019.1; Delorme

& Makeig, 2004) on MATLAB R2019a (MathWorks). The raw EEG data were downsampled

to 250 Hz and filtered using a finite impulse response (FIR) filter from 1 to 45 Hz. Next, we

applied an EEGLAB plugin of *clean_rawdata* (v2.3) to the data to automatically identify bad

channels and reconstruct bad segments using an artifact subspace reconstruction method

(ASR; Mullen et al., 2015). The data was then run through adaptive mixture independent

component analysis (AMICA) using an EEGLAB plugin of *AMICA* (v1.5.1) and also applied

an EEGLAB plugin of *ICLabel* (v1.3; Pion-Tonachini et al., 2019) to identify artifact

components. More specific information about the preprocess after filtering are as follow:

First, the channel was rejected as a bad channel when it had a flat line of at least 5 s, had

more than 4 *SD* line noise relative to its total channel signal, or correlated at less than 0.85

with adjacent channels. Second, continuous data were cleaned and reconstructed if their

power deviated too much from the calibration data when evaluated by the ASR method. As

the experimental booth was unshielded and the data were relatively noisy, we chose a 10-

SD burst-detection criteria threshold as a relatively strict criterion (Chang et al., 2020). If the

data that were decomposed into short segments had an RMS value greater than 10 *SD*

compared to the clean-calibration data portion (the concatenated data was automatically

determined based on the distribution of signal variance), the data in that time window were

reconstructed based on the remaining components as containing artifacts. Third, we ran an

adaptive mixture ICA (Hsu et al., 2018; J. A. Palmer et al., 2008; Jason A. Palmer et al.,

2006) using an EEGLAB plugin of *AMICA*, and thereafter labeled each ICs using ICLabel

(Pion-Tonachini et al., 2019) and conducted the dipole estimation using the EEGLAB plugin

of *DIPFIT*. Finally, using this information, we kept the "signal IC"s that meet the following

three criteria: (a) IC, which was labeled as "brain" by ICLabel; (b) IC in which the residual

variance of the dipole was less than 15% (i.e., the IC with RV >15% is assumed to be mostly

an artifact component); and (c) the dipole was presumed to be in the brain area. In addition,

ICs that still contained data with amplitudes greater than 150 μV after these preprocess were

rejected as likely to be artifactual components.

2-6. EEG analysis

2-6-1. ERP analysis

Preprocessed EEG data were epoched from −200 to 1000 ms, with 0 ms at the

stimulus onset. The trials that received incorrect responses were excluded from later

analysis. Using these epochs, individual ERPs were computed for the two conditions:

averaging target trials (M = 78.89, *SD* = 1.77; range: 68–80, for all four tasks) and averaging

standard trials (M = 319.33, *SD* = 2.16; range: 303–320 for all of four tasks). Then, the

individual ERPs were averaged into two grand-average ERPs for each condition. For the

ERP analysis, the voltage was corrected using the segment from −200 to 0 ms as a

baseline.

2-6-2. EEG microstate analysis

We performed EEG microstate analysis on the preprocessed data using some of the

functions of the EEGLAB plugin "*MST1.0*" (Poulsen et al., 2018). In the microstate analysis,

the canonical microstate template is obtained through clustering topographies extracted from

EEG or ERP signals (Khanna et al., 2014; D. Lehmann et al., 1998; Michel & Koenig, 2018;

Poulsen et al., 2018). In general, in terms of the electric field stability (i.e., excellent signal-

to-noise ratio), the points that receive clustering seeds are selected based on the peak of

global field power (GFP; Khanna et al., 2014; D. Lehmann et al., 1998). After the templates

are obtained, the topographic distribution at each timepoint in each participant's EEG time

series is labeled according to which class of template it most resembles (Khanna et al.,

2014; D. Lehmann et al., 1998; Michel & Koenig, 2018; Poulsen et al., 2018).

In this study, we decided to use stable templates that were created in other large-

scale EEG datasets (Kashihara et al., 2024) as the templates for this study. This is because

we considered microstates to be frequently traversed ("high traffic") points that were defined

in the standard MDS state space with embedded EEG topography (Asai et al., 2023;

Kashihara et al., 2024). Using this 10-state template that considers topographical polarity

(Figure 2), we applied the template-fitting procedure to identify the label of the microstate

class at each timepoint. The microstate template was fitted to individual trial-by-trial ERP

epochs in the target and standard condition of each task.

Figure 2. Stable EEG microstate templates considering topographical polarity from

large-scale datasets.

Microstate templates used for microstate fitting to EEG during oddball tasks in our study.

These templates were originally created in a separate study by Kashihara et al. (2024) *using*

a large-scale eyes-closed resting-state EEG dataset (N = 190, 8-min recordings). In their

study, five microstate maps were initially extracted using the modified k-means method. To

enable polarity-sensitive fitting, polarity-inverted patterns were then generated by reversing

the sign of the voltage values at each electrode. Each state represents a stable

topographical configuration of brain electrical activity. Considering topographical polarity

facilitates an appropriate discretization of essentially spatiotemporally continuous EEG

dynamics within the neural manifold space, where topographies are embedded using a

multidimensional scaling method (Asai et al., 2023; Kashihara et al., 2024).

<div align="center">

3. Results

</div>

3-1. Behavioral results

3-1-1. Experiment 1: Examining the effects of stimulus modality and coupling with conditions

We checked the mean correct rate for the key press of each task and found very high performance by the participants: 99.7% (SD = 0.3) for the unimodal-auditory task, 99.5% (SD = 0.9) for the unimodal-visual task, 99.8% (SD = 0.3) for the inter-modal auditory task, and 99.4% (SD = 1.6) for the inter-modal visual task. We conducted two-way ANOVA (target stimulus modality × modality coupling with stimulus condition) using the R packages of "anovakun (v4.8.9)" under R v4.4.2. Thus, there was neither significant main effects of the modality coupling with stimulus condition (F(1, 19) = 0.09, Greenhouse-Geisser's ε = 0.39, p = .77, η_p^2 = .005) and the target stimulus modality (F(1, 19) = 1.40, Greenhouse-Geisser's ε = 0.39, p = .25, η_p^2 = .07) nor an interaction effect for them (F(1, 19) = 1.42, Greenhouse-Geisser's ε = 0.39, p = .25, η_p^2 = .07; Figure 3a).

Furthermore, we checked the mean reaction time of each task in Experiment 1. The reaction times of all tasks were sufficiently fast: 311 ms (SD = 41) for the unimodal-auditory task, 360 ms (SD = 33) for the unimodal-visual task, 310 ms (SD = 45) for the inter-modal auditory task, and 308 ms (SD = 33) for the inter-modal visual task. We found significant main effects of the modality coupling with stimulus condition (unimodal > inter-modal: F(1,19) = 23.13, p < .001, η_p^2 = .55), the target stimulus modality (auditory < visual: F(1,19) = 19.24, p < .001, η_p^2 = .50), and their significant interaction effect (F(1,19) = 33.22, p < .001, η_p^2 = .64). The post analyses revealed significant differences between two visual-target tasks (unimodal visual > inter-modal visual-target: F(1,19) = 64.51, p < .001, η_p^2 = .77) and between two unimodal tasks (unimodal visual > unimodal auditory: F(1,19) = 39.82, p < .001, η_p^2 = .68; Figure 3b).

3-1-2. Experiment 2: Examining the effects of the response format

We confirmed how correctly participants counted the targets for each task in

Experiment 2. We calculated the counting accuracy for each task by absolutizing the

magnitude of the discrepancy between the number of counts reported by participants (the

total number of targets in two blocks of 200 trials) and the correct answer (80 targets) by

dividing them by the total number. Even when the response format was changed to counting

task, we found that the participants had very high performance: 97.7% (SD = 3.9) for the

unimodal-auditory task, 98.3% (SD = 2.7) for the unimodal-visual task, 97.2% (SD = 4.0) for

the inter-modal auditory task, and 98.3% (SD = 2.9) for the inter-modal visual task. A two-

way ANOVA (target stimulus modality × modality coupling with stimulus condition) revealed

that there was no significant effect of the modality coupling (F(1, 19) = 0.16, p = .69, η_p^2 =

.009) and the target stimulus modality (F(1, 19) = 1.96, p = .18, η_p^2 = .09) and interaction

between them (F(1, 19) = 0.16, p = .70, η_p^2 = .008) as in Experiment 1.

Next, we verified the effect of the response format on the behavioral index, that is, a

comparison of the percentage of correct responses between Experiments 1 and 2. We

conducted three-way mixed ANOVA (response mode × target stimulus modality × modality

coupling with stimulus condition). Thus, there was a significant main effect of response mode

(key press > count: F(1, 38) = 9.02, Greenhouse-Geisser's ε = 0.92, p = .005, η_p^2 = .19).

Furthermore, the primary interaction between the response mode and target stimulus

modality was significant, which indicated that regardless of the modality coupling, when the

target stimulus was visual and auditory, accuracy was higher in the key-press task than in

the counting task, respectively (visual: F(1, 38) = 8.84, p = .005, η_p^2 = .19; auditory: F(1, 38)

= 4.77, p = .04, η_p^2 = .11; Figure 3c). The main effects of other factors and their interactions

were not significant (ps > .40).

3-1-3. Experiment 3: Examining the effect of the presence or absence of

physical target stimuli

The mean correct rates for the omitted targets were still high: 99.1% (SD = 1.1) for

the auditory task and 97.6% (SD = 2.2) for the visual task. As shown with a paired t-test

using the R function of *t.test* and package of "effsize (v0.8.1)," there was a significant

difference between the two tasks (auditory > visual: t(18) = 3.64, p = .002, d = 0.80). In

addition, the mean reaction time was 442 ms (*SD* = 55) in the auditory task and 567 ms (*SD*

= 95) in the visual task. Moreover, we found a significant difference between the two tasks;

participants had slower responses in the visual task than that in the auditory task (t(18) =

6.64, p < .001, d = 1.51).

In addition, we examined the effect of the physical presence of the target stimulus by

comparing Experiments 1 and 3. Because Experiment 3 only included two types of tasks,

one with auditory standard stimuli and one with visual standard stimuli, we decided to focus

on the unimodal tasks in Experiment 1 and not consider the modality coupling of stimulus

conditions here.

We conducted two-way mixed ANOVA (presence or absence of physical target

stimulus × target stimulus modality) with accuracy as the dependent variable. Thus, there

was a significant main effect of the presence or absence of physical target stimulus (existed

target > omitted target: F(1, 37) = 12.10, Greenhouse-Geisser's ε = 1.00, p = .001, η_p^2 = .25).

In addition, there was a significant main effect of target stimulus modality (auditory > visual:

F(1, 37) = 14.48, Greenhouse-Geisser's ε = 1.00, p < .001, η_p^2 = .28). Furthermore, the

interaction between these two factors was significant (F(1, 37) = 9.80, Greenhouse-Geisser's

ε = 1.00, p = .003, η_p^2 = .21). From the post analyses, there were significant differences

between auditory tasks (existed target > omitted target: F(1,37) = 4.23, p = .047, η_p^2 = .10)

and visual tasks (existed target > omitted target: F(1,37) = 13.27, p < .001, η_p^2 = .26).

Moreover, in omitted target tasks, there were significant differences between standard

stimulus modalities (auditory > visual: F(1,18) = 13.23, p = .002, η_p^2 = .42; Figure 3d).

Figure 3. Behavioral results pertaining to the accuracy and reaction time in the three

experiments in this study.

(a) Accuracy differences between target modality and modality coupling in Experiment 1. (b)

Reaction-time differences between target modality and modality coupling in Experiment 1.

(c) Accuracy differences between the target modality, modality coupling, and response

format in Experiment 2. (d) Accuracy differences between stimulus modality and existence or

absence of physical target stimuli in Experiment 3.

3-2. ERPs

ERPs were calculated for each task and condition using 1200-ms epochs (–200 ms

prestimulus to 1000-ms poststimulus). In Experiments 1 and 3, only correct trials (i.e.,

participants responded to the target stimulus and did not respond to the standard stimulus)

were included in the later analysis.

3-2-1. Confirming the P3 in each experiment: Comparison of standard and

target trials in each task of each experiment

Figure 4 presents the grand average ERPs at a Pz electrode of each condition and

task in Experiments 1–3. The results of the permutation test comparing the average ERPs

for target trials and standard trials for each experiment and task showed that, in Experiment

1, the positive potential in the 300–400 ms range for target trials (magenta line) was

significantly larger than that for standard trials (cyan) for each task, even after FDR

correction (the following periods were significant at $p < .05$ after FDR correction for each

task: unimodal-auditory, 264–484, 532–540, and 928–936 ms; unimodal-visual, 92–112,

148–200, 304–516, 720–796, 804, 920–932, and 952–992 ms; inter-modal auditory-target,

180, 312–380 ms; inter-modal visual-target, 100–120, 164–200, 224–608, 684–696, and

988–1000 ms). This trend was almost same in Experiment 2 except for the inter-modal

auditory-target task (unimodal-auditory: 332–404, 436–484, 508–580, 588–616, 628–660,

and 736–740 ms; unimodal-visual: 336–444 and 476–488 ms; inter-modal visual-target: -56,

24–28, 48–56, 92–96, 108–116, 172–196, 256–664, and 1000 ms; inter-modal visual-target:

-60- -56, 24-28, 48-56, 88-96, 108-116, 168-200, 252-620, 636-644, 1000 ms). Even in

Experiment 3, where the target stimulus was not physically present, there was a significant

difference between the target and standard trials in the auditory missing-target oddball task

(–156 to –140, –28 to –8, 100–116, 148–164, 272–616, 632–640, and 800–828 ms) as well

as in the visual missing-target oddball task (136–188, 412–414, and 540–688 ms; ps < .05).

Figure 4. ERPs at Pz and voltage heatmaps of all 32 channel electrodes of the target vs. standard trials for each task in three experiments.

The magenta (target trial) and cyan (standard trial) lines show the grand average ERP time course on the Pz electrode. Gray areas in each ERP panel mean a significant difference between the target and standard trial in t-tests with FDR correction. The heatmaps show the voltage level of all 32 electrodes. Of the two heatmaps in each task, the upper one depicts the grand average potential of the target trial, and the lower one is the standard trial. Red means positive potential whereas blue means negative potential (see the color bar). In each heatmap, the vertical axis shows the array of 32 electrodes, and the horizontal axis shows the time course. The order of the electrodes for all heatmaps is shown in the lower right corner.

3-2-2. Experiment 1: Examining the effects of stimulus modality and coupling

with stimulus conditions

Since the emergence of the P300 after the target stimulus was nearly confirmed, the

peak amplitude at the Pz electrode between 300 and 600 ms in the trial average waveform

of each participant was treated as the P300 amplitude in the subsequent ERP analysis.

First, to examine the effect of stimulus modality and modality coupling with stimulus

condition, we performed a 2 (target modality: auditory, visual) × 2 (modality coupling:

unimodal, intermodal) two-factor within-participant analysis of variance (ANOVA) for the

mean P300 amplitude. Results showed that only the main effect of target modality was

significant, with visual targets having larger P300 amplitudes than auditory targets ($F(1,19)$ =

43.32, $p < .001$, $\eta_p^2 = 0.55$; Figure 5a). The main effect of modality coupling ($F(1,19) = 2.43$,

$p = .14$, $\eta_p^2 = .11$) and interaction ($F(1,19) = .07$, $p = .80$, $\eta_p^2 = .003$) were not significant.

3-2-3. Experiment 2: Examining the effects of the response format

Next, to investigate the effect of response format on P300 amplitude, a 2 (target

modality: auditory, visual) × 2 (modality coupling: unimodal, intermodal) × 2 (response

format: key press, count) mixed-factors ANOVA was conducted.

The results showed that the main effect of modality coupling ($F(1,38) = 5.18$, $p = .03$,

$\eta_p^2 = 0.12$) and the main effect of target modality ($F(1,38) = 55.87$, $p < .001$, $\eta_p^2 = 0.60$) were

significant as was the main effect of the response type ($F(1,38) = 3.51$, $p = .07$, $\eta_p^2 = 0.08$;

Figure 5b). The P300 amplitude was larger when the intermodal condition was used rather

than when the unimodal condition was used; the P300 amplitude was larger when the visual

target was used than when the auditory target was used; and the P300 amplitude was larger

when the participant responded to the target by pressing a key rather than when it was

counted. None of the interactions were significant (ps > .57).

3-2-4. Experiment 3: Examining the effects of the presence or absence of

physical target stimuli

Finally, to investigate the effect of the presence or absence of a physical target

stimulus on P300 amplitude, a 2 (stimulus modality: auditory, visual) x 2 (presence or

absence of physical target: present, absent) mixed-factors ANOVA was conducted by

comparing the unimodal auditory and visual data from Experiment 1 and the data from

Experiment 3.

The results showed that the main effect of stimulus modality ($F(1,37) = 13.57$, $p <$

.001, $\eta_p^2 = 0.27$) and the main effect of the presence or absence of a physical target ($F(1,37)$

$= 6.04$, $p = .02$, $\eta_p^2 = 0.14$) were significant. In each case, the P300 amplitude was larger

when the visual target was presented than when the auditory target was presented, and the

P300 amplitude was larger when the physical target stimulus was presented than when it

was not presented. Furthermore, the interaction was significant ($F(1,37) = 12.50$, $p = .001$,

$\eta_p^2 = 0.25$). Post-hoc tests showed that the P300 amplitude was larger when there was a

physical target stimulus than when there was no physical target stimulus, both in the auditory

task ($F(1,37) = 5.08$, $p = .03$, $\eta_p^2 = 0.12$) and in the visual task ($F(1,37) = 16.38$, $p < .001$, η_p^2

$= 0.31$), and the P300 amplitude was larger for the visual task than for the auditory task only

when there was a physical stimulus ($F(1,19) = 11.22$, $p = .003$, $\eta_p^2 = 0.37$; Figure 5c).

Figure 5. ERP results in the three experiments of this study.

(a) P300 amplitude at the case of target modality and modality coupling in Experiment 1. (b)

P300 differences between target modality, modality coupling, and response format in

Experiment 2. (c) P300 differences between stimulus modality and the existence or absence

of physical target stimuli in Experiment 3.

3-3. EEG microstate transition probabilities

Using the 10-state template with topographical polarity consideration (Figure 2) elucidated by Kashihara et al. (2024), we performed template fitting for the EEG epoch of each trial, each participant, each task, and each experiment to obtain the microstate label sequence. To focus on the whole-brain state transitions during the same period as the P300 component was identified in the ERP described in the previous section, template fitting was performed after sub-epoching from 300 to 600 ms following stimulus presentation.

The grand average transition probabilities (TPs) of the target (top row of each experiment) and standard trials (second row of each experiment) for each of the four tasks in all three experiments are shown in Figures 6—8. The TPs were calculated for 90 transitions, excluding self-recurrence, with the entire label sequence for a given epoch set to 1. Thereafter, the transition probabilities for each transition were averaged for each target trial (for a maximum of 80 trials) and standard trial (maximum of 320 trials) for each task within each participant and then averaged for each participant for each task to obtain the grand average TP for each task.

To further confirm the difference between the TPs of the target and standard trials, we subtracted the grand average TP of the standard trials from that of the target trials to obtain the difference TP and standardized difference between the two (Figures 6—8, bottom two rows of each experiment). For the target and standard trials, similar trends were observed in all experiments and for all task conditions: compared to the standard trials, the target trials had more transitions that converged on E- between 300 and 600 ms after stimulus onset, and conversely, fewer transitions that related to the opposite pole, E+.

600   **Figure 6. EEG microstate transition probabilities in Experiment 1 of this study.**

604   **Figure 7. EEG microstate transition probabilities in Experiment 2 of this study.**

607   **Figure 8. EEG microstate transition probabilities in Experiment 3 of this study.**

4. Discussion

In this study, we clarified the polarity-sensitive microstate transition dynamics in

oddball tasks by conducting multiple oddball tasks with manipulated experimental conditions

that are known to affect the P300 component. In Experiment 1, we examined the effects of

the target stimulus modality and the coupling of the standard and target stimulus modalities

on the EEG. Furthermore, to isolate the motor effects and the effects of the presence or

absence of physical stimuli, we asked the participants to count without any motor behavior

toward the target (Experiment 2), and we used a target without any physical stimulus

(Experiment 3) and then compared these to Experiment 1.

In this study, very high response accuracy was obtained across all experiments and

task conditions. In Experiments 1 and 2, there was no significant difference in the response

accuracy based on the target stimulus modality and modality coupling. However, response

times tended to be higher in the unimodal visual task. Furthermore, in Experiment 3,

response times were higher and accuracy was lower in the visual task. These findings

suggest that responding to visual stimuli may be more challenging than responding to

auditory stimuli. Regarding response format, regardless of the stimulus condition coupling,

accuracy tended to be higher when participants were required to press a key in response to

the target, whether it was visual or auditory, rather than counting the occurrences. The

difference between keypress responses, which involve processing the presented stimulus in

real-time, and counting responses, which require the maintenance of a record of past

occurrences while processing the current stimulus, may reflect differences in task goals and

motivation among participants. The differences in response accuracy observed in this study

may merely reflect the variations in task difficulty.

In this study, P300 ERP responses were observed in almost all oddball tasks. The

amplitude of these P300 components was influenced by the target stimulus modality.

Compared to when the target was an auditory stimulus, a target that comprised a visual

stimulus elicited a higher amplitude P300. This finding is consistent with the results reported

by Polich & Heine (1996), who showed that P300 from auditory stimuli was smaller than that

638 from visual stimuli. Considering this finding alongside the behavioral results that suggest

639 differences in task difficulty generates the inference that visual targets require a greater

640 allocation of attentional resources than auditory targets, which are processed more

641 passively. This increased attentional demand is reflected in the longer response times for

642 visual stimuli and the corresponding P300 amplitude.

643   This hypothesis is supported by the results of Experiment 3, which investigated the

644 effects of the presence or absence of a physical stimulus on the target. In Experiment 3,

645 which focused on an unimodal task, the P300 amplitude was larger for the visual task than

646 for the auditory task, but this was only observed when the physical target stimulus was

647 presented. P300 could reflect brain activity during the modification of an environmental

648 model, that is, during the updating of context (Donchin & Coles, 1988). Therefore, the effect

649 of stimulus modality on P300 amplitude is an event that occurs depending on the

650 characteristics of the physical target stimulus.

651   Regarding the effect of modality coupling between stimulus conditions on P300,

652 Experiment 2 confirmed that, compared to the unimodal condition, the intermodal condition

653 elicited higher P300 amplitudes. Although no significant difference was observed in

654 Experiment 1, the descriptive trend was consistent. This finding aligns with previous studies

655 (Brown et al., 2006, 2007) and extends prior research examining the influence of auditory

656 and visual standard stimuli on late ERP components evoked by auditory targets. Brown et al.

657 (2006, 2007) reported that late components, such as P300, are larger in audiovisual oddball

658 tasks, which suggests the occurrence of distinct intermodal processes at later stages. Our

659 study confirmed the occurrence of this late-stage context-dependent processing and

660 suggests that it applies not only to auditory processing but also to visual processing.

661   Importantly, the target modality and the modality coupling between the target and

662 standard stimuli each independently affected the P300 amplitude. This suggests that the

663 meaning and context of each stimulus are important, regardless of whether the stimulus is

664 visual or auditory or whether the frequent stimulus and rare target are the same or different

665 modalities.

Furthermore, regarding the response format, the P300 amplitudes were greater in the

keypress task than in the counting task. Prior research on oddball tasks has yielded

inconsistent findings regarding the influence of response modality. However, our results

support the notion that late ERP responses are stronger in keypress tasks.

We found that the ERP components were influenced by target stimulus modality,

modality coupling, response format, and the presence or absence of physical stimuli.

Previous ERP studies have suggested that these variables affect the configuration of EEG

topography (e.g., Brown et al., 2006, 2007; Polich & Heine, 1996). To examine the spatial

patterns of whole-brain states at a given moment and further track their temporal dynamics,

polarity-informed microstate analysis was conducted to determine the transitional

probabilities during the period corresponding to P300 generation. The results revealed that

during the P300-related period, target trials exhibited increased cohesion toward a state

referred to as E-. In contrast, the transitions associated with E+ tended to be less frequent

than those in the standard trials. This finding suggests the process of transitioning to a

specific state of a specific polarity rather than indication of an increase in a certain electrical

field configuration pattern, and it also emphasizes the importance of considering

topographical polarity in microstate analysis.

These transition patterns were consistently observed regardless of task conditions,

response formats, or the presence or absence of physical stimuli. This suggests that the

increased transition to E- may reflect a shift in whole-brain processing networks

corresponding to the P300 component in target trials. The microstate E has been associated

with interoceptive sensory and emotional information processing; moreover, its involvement

in the salience network has been reported (Tarailis et al., 2024). The differentiation of the

topographical polarity of microstates is a relatively recent development (Kashihara et al.,

2024), and little evidence has been accumulated regarding the functional distinctions

between E+ and E-. However, fMRI studies on oddball tasks have suggested that key

regions of the salience network, including the anterior cingulate cortex and insular cortex,

are involved in this process (Harsay et al., 2012; Horn et al., 2003), indicating that E- may

reflect the functions governed by the salience network. Specifically, the activation of the

insular cortex, particularly the anterior insular cortex, potentially serves as an interface that

mediates dynamic interactions between large-scale brain networks that are involved in

externally directed attention and internally directed cognition (Menon & Uddin, 2010). That is,

the salience network likely plays a critical role in detecting target stimuli, marking them for

additional processing, and initiating appropriate control signals—such as keypress or

counting responses. The spatiotemporal dynamics of EEG aggregation toward E- may

reflect this process.

Finally, the limitations of this study should be acknowledged. First, this study

manipulated experimental conditions to examine the effects of stimulus modality condition,

response format, and the presence or absence of physical stimuli on relatively late neural

dynamics following stimulus onset, which manifest as P300 and microstate transitions.

However, the factors examined in this study represent only a subset of potential influences

and other cognitive factors, such as attentional load and working memory demands, that

may contribute to the formation of ERP components and microstate dynamics. Furthermore,

investigating whole-brain state transitions using ERP markers in earlier processing stages

may provide valuable insights. Future research should explore how these additional factors

interact with the effects observed in the present study.

Moreover, through polarity-sensitive microstate analysis, we identified a potential role

of transitions to E- in target stimulus detection and switching to appropriate processing

states. However, the interpretation of polarity-sensitive microstate transitions remains an

emerging area of research, and further studies are necessary to elucidate the distinct

functional roles of E+ and E- within broader neural network dynamics. Although EEG

microstate analysis offers high temporal resolution, its spatial resolution is inherently limited.

Future research could integrate EEG with fMRI to more precisely localize the neural sources

that mediate microstate transitions, particularly those associated with E-.

In conclusion, this study shows that EEG microstate transitions, in particular the

increase in cohesiveness toward E-, correspond to the P300 component and reflect the

dynamics of large-scale neural networks involved in target processing. This finding suggests

that transitions to a whole-brain state pattern called E- play an important role in the detection

and marking of target stimuli, facilitating the initiation of an appropriate response. Through

the integration of microstate analysis with conventional ERP measurements, this study

provided new insights into the spatiotemporal dynamics of neural network reconfiguration

during the oddball task. These findings further the understanding of the neural mechanisms

of stimulus-driven attention and highlight the importance of considering topographical

polarity-sensitive microstate transitions in future research.

5. <u>References</u>

Al Zoubi, O., Mayeli, A., Misaki, M., Tsuchiyagaito, A., Zotev, V., Refai, H., Paulus, M. &

Bodurka, J. (2022). Canonical EEG microstates transitions reflect switching among

BOLD resting state networks and predict fMRI signal. *Journal of Neural*

Engineering, 18(6), 066051. https://doi.org/10.1088/1741-2552/ac4595

Asai, T., Kashihara, S., Chiyohara, S., Hiromitsu, K., & Imamizu, H. (2023). Spatio-temporal

"global" neurodynamics of the human brain in continuous and discrete picture: Simple

statistics meet on-manifold microstates as multi-level cortical attractors. bioRxiv.

https://doi.org/10.1101/2023.07.13.548951

Barack, D. L. & Krakauer, J. W. (2021). Two views on the cognitive brain. *Nature Reviews.*

Neuroscience, 22(6), 359-371. https://doi.org/10.1038/s41583-021-00448-6

Barrett, G., Neshige, R. & Shibasaki, H. (1987). Human auditory and somatosensory event-

related potentials: effects of response condition and age. *Electroencephalography and*

Clinical Neurophysiology, 66(4), 409-419. https://doi.org/10.1016/0013-4694(87)90210-

0

Brázdil, M., Roman, R., Daniel, P. & Rektor, I. (2003). Intracerebral somatosensory event-

related potentials: effect of response type (button pressing versus mental counting) on

P3-like potentials within the human brain. *Clinical Neurophysiology, 114*(8), 1489-1496.

https://doi.org/10.1016/s1388-2457(03)00135-4

Bréchet, L., Brunet, D., Birot, G., Gruetter, R., Michel, C. M. & Jorge, J. (2019). Capturing

the spatiotemporal dynamics of self-generated, task-initiated thoughts with EEG and

fMRI. *NeuroImage, 194*, 82-92. https://doi.org/10.1016/j.neuroimage.2019.03.029

Britz, J., Van De Ville, D. & Michel, C. M. (2010). BOLD correlates of EEG topography reveal

rapid resting-state network dynamics. *NeuroImage, 52*(4), 1162-1170.

https://doi.org/10.1016/j.neuroimage.2010.02.052

Brown, C. R., Clarke, A. R. & Barry, R. J. (2006). Inter-modal attention: ERPs to auditory

targets in an inter-modal oddball task. *International Journal of Psychophysiology, 62*(1),

77-86. https://doi.org/10.1016/j.ijpsycho.2006.02.001

Brown, C. R., Clarke, A. R. & Barry, R. J. (2007). Auditory processing in an inter-modal

oddball task: effects of a combined auditory/visual standard on auditory target

ERPs. *International Journal of Psychophysiology, 65*(2), 122-131.

https://doi.org/10.1016/j.ijpsycho.2007.04.002

Chang, C. Y., Hsu, S. H., Pion-Tonachini, L. & Jung, T. P. (2018). Evaluation of artifact

subspace reconstruction for automatic EEG artifact removal. *Annual International*

Conference of the IEEE Engineering in Medicine and Biology Society. IEEE Engineering

in Medicine and Biology Society. Annual International Conference, 2018, 1242-1245.

https://doi.org/10.1109/EMBC.2018.8512547

Croce, P., Tecchio, F., Tamburro, G., Fiedler, P., Comani, S. & Zappasodi, F. (2022). Brain

electrical microstate features as biomarkers of a stable motor output. *Journal of Neural*

Engineering, 19(5), 056042. https://doi.org/10.1088/1741-2552/ac975b

D'Croz-Baron, D. F., Bréchet, L., Baker, M. & Karp, T. (2021). Auditory and visual tasks

influence the temporal dynamics of EEG microstates during post-encoding rest. *Brain*

Topography, 34(1), 19-28. https://doi.org/10.1007/s10548-020-00802-4

Delorme, A. & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-

trial EEG dynamics including independent component analysis. *Journal of Neuroscience*

Methods, 134(1), 9-21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Donchin, E., & Coles, M. G. H. (1988). Is the P300 component a manifestation of context

updating? *The Behavioral and Brain Sciences, 11*(3), 357-374.

https://doi.org/10.1017/S0140525X00058027

Ebitz, R. B. & Hayden, B. Y. (2021). The population doctrine in cognitive

neuroscience. *Neuron, 109*(19), 3055-3068.

https://doi.org/10.1016/j.neuron.2021.07.011

Harsay, H. A., Spaan, M., Wijnen, J. G. & Ridderinkhof, K. R. (2012). Error awareness and

salience processing in the oddball task: shared neural mechanisms. *Frontiers in Human*

Neuroscience, 6, 246. https://doi.org/10.3389/fnhum.2012.00246

Horn, H., Syed, N., Lanfermann, H., Maurer, K. & Dierks, T. (2003). Cerebral networks linked

to the event-related potential P300. *European Archives of Psychiatry and Clinical*

Neuroscience, 253(3), 154-159. https://doi.org/10.1007/s00406-003-0419-4

Hsu, S. H., Pion-Tonachini, L., Palmer, J., Miyakoshi, M., Makeig, S. & Jung, T. P. (2018).

Modeling brain dynamic state changes with adaptive mixture independent component

analysis. *NeuroImage, 183*, 47-61. https://doi.org/10.1016/j.neuroimage.2018.08.001

Iyer, K. K., Hwang, K., Hearne, L. J., Muller, E., D'Esposito, M., Shine, J. M. & Cocchi, L.

(2022). Focal neural perturbations reshape low-dimensional trajectories of brain activity

supporting cognitive performance. *Nature Communications, 13*(1), 4.

https://doi.org/10.1038/s41467-021-26978-2

Kashihara, S., Asai, T., & Imamizu, H. (2024). Topographical polarity reveals continuous

EEG microstate transitions and electric field direction in healthy aging. *bioRxiv*

2024.03.24.586427. https://doi.org/10.1101/2024.03.24.586427

Khanna, A., Pascual-Leone, A. & Farzan, F. (2014). Reliability of resting-state microstate

features in electroencephalography. *PloS One, 9(12)*, e114163.

https://doi.org/10.1371/journal.pone.0114163

Khanna, A., Pascual-Leone, A., Michel, C. M. & Farzan, F. (2015). Microstates in resting-

state EEG: current status and future directions. *Neuroscience and Biobehavioral*

Reviews, 49, 105-113. https://doi.org/10.1016/j.neubiorev.2014.12.010

Kim, K., Duc, N. T., Choi, M. & Lee, B. (2021). EEG microstate features according to

performance on a mental arithmetic task. *Scientific Reports, 11*(1), 343.

https://doi.org/10.1038/s41598-020-79423-7

Kindler, J., Hubl, D., Strik, W. K., Dierks, T. & Koenig, T. (2011). Resting-state EEG in

schizophrenia: auditory verbal hallucinations are related to shortening of specific

microstates. *Clinical Neurophysiology, 122*(6), 1179-1182.

https://doi.org/10.1016/j.clinph.2010.10.042

Kondakor, I., Lehmann, D., Michel, C. M., Brandeis, D., Kochi, K., & Koenig, T. (1997).

Prestimulus EEG microstates influence visual event-related potential microstates in field

maps with 47 channels. *Journal of Neural Transmission*, *104*(2-3), 161–173.

https://doi.org/10.1007/BF01273178

Kotchoubey, B. (2014). First love does not die: a sustaining primacy effect on ERP

components in an oddball paradigm. *Brain Research, 1556*, 38-45.

https://doi.org/10.1016/j.brainres.2014.02.012

Lehmann, D., Michel, C. M., Pal, I. & Pascual-Marqui, R. D. (1994). Event-related potential

maps depend on prestimulus brain electric microstate map. *The International Journal of*

Neuroscience, 74(1-4), 239-248. https://doi.org/10.3109/00207459408987242

Lehmann, D., Ozaki, H. & Pal, I. (1987). EEG alpha map series: brain micro-states by space-

oriented adaptive segmentation. *Electroencephalography and clinical*

neurophysiology, 67(3), 271-288. https://doi.org/10.1016/0013-4694(87)90025-3

Lehmann, D., Strik, W. K., Henggeler, B., Koenig, T. & Koukkou, M. (1998). Brain electric

microstates and momentary conscious mind states as building blocks of spontaneous

thinking: I. Visual imagery and abstract thoughts. *International Journal of*

Psychophysiology, 29(1), 1-11. https://doi.org/10.1016/s0167-8760(97)00098-6

Mau, W., Hasselmo, M. E. & Cai, D. J. (2020). The brain in motion: How ensemble fluidity

drives memory-updating and flexibility. *eLife, 9*, e63550.

https://doi.org/10.7554/eLife.63550

Menon, V. & Uddin, L. Q. (2010). Saliency, switching, attention and control: a network model

of insula function. *Brain structure & function, 214(5-6)*, 655-667.

https://doi.org/10.1007/s00429-010-0262-0

Michel, C. M. & Koenig, T. (2018). EEG microstates as a tool for studying the temporal

dynamics of whole-brain neuronal networks: A review. *NeuroImage, 180(Pt B)*, 577-593.

https://doi.org/10.1016/j.neuroimage.2017.11.062

Milz, P., Faber, P. L., Lehmann, D., Koenig, T., Kochi, K. & Pascual-Marqui, R. D. (2016).

The functional significance of EEG microstates--Associations with modalities of

thinking. *NeuroImage, 125*, 643-656. https://doi.org/10.1016/j.neuroimage.2015.08.023

Mishra, A., Englitz, B. & Cohen, M. X. (2020). EEG microstates as a continuous

phenomenon. *NeuroImage, 208*, 116454.

https://doi.org/10.1016/j.neuroimage.2019.116454

Mullen, T. R., Kothe, C. A., Chi, Y. M., Ojeda, A., Kerth, T., Makeig, S., Jung, T. P. &

Cauwenberghs, G. (2015). Real-time neuroimaging and cognitive monitoring using

wearable dry EEG. *IEEE Transactions on Bio-medical Engineering, 62*(11), 2553-2567.

https://doi.org/10.1109/TBME.2015.2481482

Musso, F., Brinkmeyer, J., Mobascher, A., Warbrick, T. & Winterer, G. (2010). Spontaneous

brain activity and EEG microstates. A novel EEG/fMRI analysis approach to explore

resting-state networks. *NeuroImage, 52*(4), 1149-1161.

https://doi.org/10.1016/j.neuroimage.2010.01.093

Peirce, J. W. (2007). PsychoPy--Psychophysics software in Python. *Journal of Neuroscience*

Methods, 162(1-2), 8-13. https://doi.org/10.1016/j.jneumeth.2006.11.017

Peirce, J. W. (2008). Generating stimuli for neuroscience using PsychoPy. *Frontiers in*

Neuroinformatics, 2, 10. https://doi.org/10.3389/neuro.11.010.2008

Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman, E.

& Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior*

Research Methods, 51(1), 195-203. https://doi.org/10.3758/s13428-018-01193-y

Pessoa, L. (2019). Neural dynamics of emotion and cognition: From trajectories to

underlying neural geometry. *Neural Networks, 120*, 158-166.

https://doi.org/10.1016/j.neunet.2019.08.007

Pion-Tonachini, L., Kreutz-Delgado, K. & Makeig, S. (2019). ICLabel: An automated

electroencephalographic independent component classifier, dataset, and

website. *NeuroImage, 198*, 181-197. https://doi.org/10.1016/j.neuroimage.2019.05.026

Pizzagalli, D., Lehmann, D., Koenig, T., Regard, M. & Pascual-Marqui, R. D. (2000). Face-

elicited ERPs and affective attitude: brain electric microstate and tomography

analyses. *Clinical Neurophysiology, 111*(3), 521-531. https://doi.org/10.1016/s1388-

2457(99)00252-7

Polich, J. & Heine, M. R. (1996). P300 topography and modality effects from a single-

stimulus paradigm. *Psychophysiology, 33*(6), 747-752. https://doi.org/10.1111/j.1469-

8986.1996.tb02371.x

Poulsen, A. T., Pedroni, A., Langer, N., & Hansen, L. K. (2018). Microstate EEGlab toolbox:

An introductory guide. bioRxiv (p. 289850). https://doi.org/10.1101/289850

Ragazzoni, A., Di Russo, F., Fabbri, S., Pesaresi, I., Di Rollo, A., Perri, R. L., Barloscio, D.,

Bocci, T., Cosottini, M. & Sartucci, F. (2019). "Hit the missing stimulus". A simultaneous

EEG-fMRI study to localize the generators of endogenous ERPs in an omitted target

paradigm. *Scientific Reports, 9*(1), 3684. https://doi.org/10.1038/s41598-019-39812-z

Ruggeri, P., Meziane, H. B., Koenig, T. & Brandner, C. (2019). A fine-grained time course

investigation of brain dynamics during conflict monitoring. *Scientific Reports, 9*(1), 3667.

https://doi.org/10.1038/s41598-019-40277-3

Salisbury, D. F., Rutherford, B., Shenton, M. E. & McCarley, R. W. (2001). Button-pressing

affects P300 amplitude and scalp topography. *Clinical Neurophysiology, 112*(9), 1676-

1684. https://doi.org/10.1016/s1388-2457(01)00607-1

Seitzman, B. A., Abell, M., Bartley, S. C., Erickson, M. A., Bolbecker, A. R. & Hetrick, W. P.

(2017). Cognitive manipulation of brain electric microstates. *NeuroImage, 146*, 533-543.

https://doi.org/10.1016/j.neuroimage.2016.10.002

Shaw, S. B., Dhindsa, K., Reilly, J. P. & Becker, S. (2019). Capturing the forest but missing

the trees: microstates inadequate for characterizing shorter-scale EEG

dynamics. *Neural Computation, 31*(11), 2177-2211.

https://doi.org/10.1162/neco_a_01229

Shine, J. M., Breakspear, M., Bell, P. T., Ehgoetz Martens, K. A., Shine, R., Koyejo, O.,

Sporns, O. & Poldrack, R. A. (2019). Human cognition involves the dynamic integration

of neural activity and neuromodulatory systems. *Nature Neuroscience, 22*(2), 289-296.

https://doi.org/10.1038/s41593-018-0312-0

Sikka, A., Jamalabadi, H., Krylova, M., Alizadeh, S., van der Meer, J. N., Danyeli, L.,

Deliano, M., Vicheva, P., Hahn, T., Koenig, T., Bathula, D. R. & Walter, M. (2020).

Investigating the temporal dynamics of electroencephalogram (EEG) microstates using

recurrent neural networks. *Human Brain Mapping, 41*(9), 2334-2346.

https://doi.org/10.1002/hbm.24949

Starr, A., Sandroni, P. & Michalewski, H. J. (1995). Readiness to respond in a target

detection task: pre- and post-stimulus event-related potentials in normal

subjects. *Electroencephalography and Clinical Neurophysiology, 96*(1), 76-92.

https://doi.org/10.1016/0013-4694(94)00162-e

Tamano, R., Ogawa, T., Katagiri, A., Cai, C., Asai, T. & Kawanabe, M. (2022). Event-related

microstate dynamics represents working memory performance. *NeuroImage, 263*,

119669. https://doi.org/10.1016/j.neuroimage.2022.119669

Tarailis, P., Koenig, T., Michel, C. M. & Griškova-Bulanova, I. (2024). The Functional

Aspects of Resting EEG Microstates: A Systematic Review. *Brain topography, 37*(2),

181-217. https://doi.org/10.1007/s10548-023-00958-9

Van de Ville, D., Britz, J. & Michel, C. M. (2010). EEG microstate sequences in healthy

humans at rest reveal scale-free dynamics. *Proceedings of the National Academy of*

Sciences of the United States of America, 107(42), 18179-18184.

https://doi.org/10.1073/pnas.1007841107

Westfall, J. (2016). PANGEA: Power ANalysis for GEneral Anova designs. University of

Texas at Austin. http://jakewestfall.org/publications/pangea.pdf

916