Anaesthesia selectively decouples signal propagation from responsivity in the fronto-parietal loop

Abhilash Dwarakanath[1,2,3,4*], Majid Khalili-Ardali[1,5*], Marion Gay[1], Raghuram HV[1,6], Maxime Roustan[1], Bechir Jarraya[1,7,8], Theofanis I Panagiotaropoulos[1,9,10]

[1] Cognitive Neuroimaging Unit, INSERM, CEA, Université Paris-Saclay, NeuroSpin center, Gif/Yvette, France

[2] Neural Circuits and Cognition, European Institute of Neuroscience, Göttingen 37077, Germany

[3] Cognitive Neuroscience Laboratory, Perception & Plasticity Group, Deutsches Primatenzentrum, Göttingen 37077

[4] SFB 1528 Cognition of Interaction, 37077, Göttingen

[5] Donders Center for Neuroscience, Radboud University, Nijmegen, Netherlands

[6] Department of Psychology, University of Wisconsin-Madison, Madison, USA

16   [7] University of Versailles Saint-Quentin-en-Yvelines, Université Paris-Saclay,

17   Versailles, France

18   [8] Neuromodulation Unit, Foch Hospital, Suresnes, France

19   [9] Department of Psychology, National & Kapodistrian University of Athens,

20   Athens, Greece

21   [10] Centre for Basic Research, Biomedical Research Foundation of the

22   Academy of Athens (BRFAA), Athens, Greece

23 Corresponding authors - Abhilash Dwarakanath (ADwarakanath@dpz.eu);

24   Theofanis Panagiotaropoulos (fpanag@psych.uoa.gr)

Summary

Loss of consciousness under anaesthesia is accompanied by widespread silencing of neurons, and disruption of cortical dynamics. Yet, how this affects mesoscale signal propagation within higher-order associative areas, crucially implicated in theories of consciousness, remains poorly understood. Here we combined intracortical microstimulation, and simultaneous multielectrode recordings in the ventrolateral prefrontal cortex (vlPFC) and the posterior parietal cortex (PPC) of macaques across wakefulness, and graded depths of anaesthesia. Spiking responses revealed distinct regimes: in the PFC, higher-amplitude stimulation elicited a delayed single rebound after sustained inhibition, whereas in the PPC, a faster and double-rebound profile emerged. Despite enhanced local spiking and LFP responsivity under anaesthesia, we found a strong and selective suppression of lateral signal propagation in the PFC - a breakdown strikingly absent in the PPC. This dissociation suggests that anaesthesia disrupts consciousness not merely by silencing cortical populations, but by impairing mesoscale integrative processes critical for neuronal dynamics

47 at multiple scales. Our findings demonstrate in a causal and spatially-

48 resolved manner, that lateral signal spread within higher-order cortex is a

49 key mechanism underlying conscious awareness, and its loss under

50 anaesthesia.

Introduction

Consciousness, a fundamental aspect of human experience, encompasses a spectrum ranging from full awareness and altered states of consciousness to different levels of unconsciousness seen in anaesthesia and consciousness disorders[1–4]. Understanding the neural correlates of such unconscious states, and the disruptive effects of these states on normal cortical functions, has significant implications for elucidating the cortical mechanisms underlying conscious awareness and for the treatment of consciousness disorders[5–7].

Multiple theoretical frameworks have been developed to address how consciousness arises in the brain[8,9]. Among theories that ascribe a central role to higher cortical and association areas[10–14], the Global Neuronal Workspace Theory (GNWT) proposes that conscious experience arises from a higher-order, cortex-mediated ignition, amplification and widespread dissemination of sensory information across the brain, with the prefrontal cortex (PFC) playing a key role in this global broadcasting process[15–18]. In contrast, Integrated Information Theory (IIT) posits that consciousness

corresponds to the amount and structure of integrated information (*phi -* Φ) generated by a system, emphasising locally maximised, irreducible cause-effect structures within highly interconnected neuronal ensembles, often associated with a posterior cortical "hot zone" [1,19–21], rather than with any specific role for these higher order associational cortical areas. Higher-order thought (HOT) [22,23] and related theories likewise emphasise the contribution of higher cortical regions to conscious access, albeit with distinct commitments to representation, state, access, hierarchy, and mechanism/dynamics, while recurrent processing and posterior hot zone accounts argue that posterior cortices [24–26] may be sufficient for conscious experience, with PFC contributions confined to report, metacognition or task demands [27].

Studies in humans using non-invasive whole-brain recording techniques such as BOLD-fMRI have implicated the PFC during conscious, accessible perception of a stimulus[28–30]. More specifically, direct electrophysiological recordings in the PFC show that prefrontal neuronal populations represent the contents of consciousness, while ongoing low-frequency fluctuations in the prefrontal state are involved in the stability and update of these

contents[31–35], even as some authors remain sceptical about a necessary role

of PFC for all forms of conscious experience [36,37], (but see [11].

In terms of known cytoarchitecture at the micro- and mesoscales, the PFC

is of particular interest due to its dense reciprocal connections with other

cortical areas, especially the parietal cortex, forming a fronto-parietal loop

(FP-loop)[38–41], specifically mediated *via* long-range, bidirectional,

monosynaptic projections from layer 5 pyramidal (L5p) neurons. Crucially,

beyond conscious perception *per se*, this fronto-parietal network is widely

believed to be central for sensory awareness and goal-directed access,

which are either suppressed, or abolished under anaesthesia and similar

states[15,42,43]. Work exploring posterior thalamocortical interactions *via* deep

brain stimulation (DBS), and pharmacological manipulation of

associational regions – across species – has both highlighted that the precise

contribution of fronto-parietal circuitry to consciousness [44], and muddied

the contribution of posterior cortical regions, which are supported by

studies employing gross perturbational and complexity metrics [45,46].

Therefore, to critically address the role of these areas, in particular the

ventrolateral prefrontal cortex (vlPFC) and the posterior parietal cortex

(PPC), in the genesis and loss of conscious awareness, direct causal

perturbational studies at the micro- and mesoscale are necessary[10,36].

Studying these regions under anaesthetics, which cause reversible

unconsciousness, can help reveal mechanisms of consciousness loss and

recovery[47–51], and can thus inform competing accounts by probing how

state-dependent changes in effective connectivity and mesoscale

propagation differ across key cortical nodes. Different anaesthetics have

been used to understand local and global neural dynamics during

conscious and unconscious states[47,52]. For GABAergic anaesthetics, cortical

states are dominated by anteriorised slow rhythms[53–57] in the delta-theta

range (1-4 and 5-8 Hz)[58,59], exhibiting high intra-areal but low inter-areal

coherence[55,58,60]. Sevoflurane, a commonly used veterinary inhalant

anaesthetic, potentiates inhibitory neurotransmission by primarily

enhancing $GABA_A$ receptor function[48,61]. At the cellular and microcircuit

level, general anaesthesia can decouple cortical pyramidal neurons [62] and

alter dendritic integration in a way that profoundly reshapes how inputs

are transformed into output activity, as shown for example in the

somatosensory cortex under ketamine-xylazine [63]. Furthermore,

sevoflurane reconfigures cortical neuronal discharge activity from diffuse

and decorrelated to distinct UP/DOWN states, with increased local neuronal

correlations[48,64] while impairing functional connectivity[65–67]. Crucially,

sevoflurane disrupts fronto-parietal communication[65,66] while also leading

to a reduction in large-scale signal complexity (as measured by the

Perturbational Complexity Index), a signature of the level of

consciousness[47,68]. Related work combining recordings and deep brain

stimulation has shown that loss and recovery of consciousness are

accompanied by characteristic changes in large-scale dynamics, including

within fronto-parietal circuits and thalamus [44,55,69,70]. However, the exact

nature of this functional reorganisation at the neuronal population level,

in particular at the mesoscale, remains uncharacterised[71,72]. While studies

have measured the complexity of large-scale brain activity and distinct

states elicited by transcranial magnetic stimulation (TMS) perturbation

using scalp EEG[45,73], investigations specifically targeting nodes of the fronto-

parietal loop to characterise the effects of loss of consciousness on

mesoscale signal propagation are lacking.

This study aims to address three important questions at the mesoscale

cortical level: (1) How does the response profile of prefrontal and parietal

populations to electrical stimulation differ across awake and anaesthetised

states? (2) How do these states affect signal propagation in time and space?

(3) Are there differences in the thus inferred network properties between

these two subunits of the FP-loop? To answer these questions, we employed

a simultaneous electrical microstimulation-recording protocol during

quiet wakefulness (QW), light (LS) and deep anaesthesia (DS), all performed

in the same experiment and recording from the same population, to

maintain sample consistency. The vlPFC and PPC were recorded in

different animals and sessions, a constraint imposed by experimental

feasibility that we address in detail in the Discussion. Apart from quasi-

similar population response profiles in the two cortical regions, our results

reveal a specific signature of anaesthesia in the PFC involving suppression

of lateral electrical signal propagation that is absent in the PPC. In contrast,

the amplitude of responses following electrical microstimulation did not

correlate monotonically with the level of anaesthesia, since deep

anaesthesia resulted in responses that were stronger compared to quiet

wakefulness.

Results

We recorded and stimulated from the ventrolateral prefrontal cortex of two male macaques during quiet wakefulness and anaesthesia, in the same experimental sessions, using a 10 x 10 Iridium Oxide coated Utah Array. Specialised adapters allowed us to stimulate and record members of the same population, simultaneously (Fig. 1A).

After recording and electrical microstimulation periods during quiet wakefulness, we induced anaesthesia with a single injection of ketamine and dexmedetomidine. After waiting for ~90 minutes for the ketamine to metabolise, light anaesthesia was induced by inhalation, and maintained between 1-1.5% of sevoflurane, a GABAergic anaesthetic. After recording and microstimulation during light anaesthesia, we increased the sevoflurane concentration to 3%, inducing deep anaesthesia (Fig. 1B). While most intracortical microstimulation (ICMS) that intend to characterise population response properties employ single-pulse stimulation, we devised our stimulation to mimic those used in studies that

seek to, for e.g., augment stimulus detection, or influence behavioural

outcome[74–77]. This stimulation, designed as a biphasic 50-pulse train at 250

Hz, was delivered to the two electrodes in the centre of the array, each with

opposing polarity, at varying strengths, viz. 1 to 100μA (Fig. 1B). Each

stimulation was spaced 10-20s apart to allow sufficient time for cortical

activity to return to baseline activity post-stimulation. Resting state periods,

both during wakefulness and anaesthesia, were recorded in pre- and post-

stimulation periods (Fig. 1C).

189   State characterisation in wakefulness and different anaesthesia

190   depths

How does the activity of the same cortical population vary between quiet

wakefulness and different levels of anaesthesia?

To address this, in each session from each recorded area in each animal,

we selected a population of neurons based on their firing rate during QW

(firing rate >= 0.5 spks/s), and used the same population for all downstream

analyses. In Monkey A in the PFC, while the firing rates during LS and DS

were significantly lower than during QW ($p_{Comp} < 0.001$), they were not

different from each other ($p_{Comp} > 0.05$). In Monkey C in the PPC however,

there were significant differences in the firing rate between QW and DS,

and LS and DS ($p_{Comp} < 0.001$), but not between QW and LS ($p_{Comp} > 0.07$). In

general, the firing was higher during QW, followed by LS and then DS,

subject to statistical tests as above. In Monkey B in the PFC, due to very

sparse firing, no significant differences were found in the firing rates

between any of the 3 conditions (all $p_{Comp} > 0.5$). The firing rate comparisons

are summarised in SIFig. 1. Because each session contained the same

number of channels recorded, for this statistical comparison, the Student's

t-test was employed.

Qualitatively, quiet wakefulness is typified by neuronal discharge activity

which is diffuse and uncorrelated between channels (Fig. 2A (PFC) and Fig.

2E (PPC) - top row), accompanied by low amplitude high frequency field

fluctuations (gamma) (Fig. 2C (PFC) and 2G - top row). The induction of

anaesthesia leads to a transition from this decoherent state to periods of

coherent discharge interspersed by silence, known as UP/DOWN states,

possibly leading to burst suppression in DS (Fig. 2A (PFC) and Fig. 2E (PPC)

- middle row (LS), and bottom row (DS))[78,79]. This is reflected in significantly

lower delta-theta power, and significantly higher beta and gamma power

during wakefulness (p_{Comp} < 0.0001; SIFig. 2), whereas both states of

anaesthesia are dominated by low-frequency field activity (Figs. 2D (PFC),

2H - PPC). In both recorded areas, the power in the low-frequencies

increased with increasing depth of anaesthesia (LS vs DS; p_{Comp} < 0.0001;

SIFig. 2).

We quantified the observed pattern of activity during these resting state

periods (in both wakefulness and two levels of anaesthesia) using a two-

state Hidden Markov Model (HMM) fit to the population firing vector. From

the learned state sequence, the durations of bursting (UP) and silent

(DOWN) periods were computed. While the model did not fit wakefulness

well (Fig. 2A and 2E top row - purple line P(UP)), indicating the absence of

such states during this period, it detected UP/DOWN states reliably in both

LS and DS, in both PFC and PPC (Fig 2A and 2E, middle (LS) and bottom (DS)

rows). While in Monkey A (PFC) and Monkey C (PPC), DS was typically

characterised by long silent periods up to ~16s (PFC, Fig. 2I Column 1) and

~100s (PFC, Fig. 2I column 3), silent periods in both LS and DS were

restricted to ~2.6s in Monkey B (PFC, SIFig. 3)

Finally, a synchrony index (SI) computed from the positive half of the

autocorrelation function (ACF) excluding the peak suggests that while

synchrony of population firing is relatively high during both LS and DS

periods, QW displays low synchrony (Fig. 2B (PFC) and 2C (PPC); SIFig. 1),

in both the PFC and the PPC. The difference in synchrony between each

state was evaluated statistically, and all differences were significant with

all p_{Comp} < 0.001. We attribute this to the coherent UP/DOWN states observed

under anaesthesia.

State dependence of cortical responsivity to electrical

microstimulation

Following the characterisation of prefrontal and parietal states

during wakefulness and different depths of sevoflurane anaesthesia, we

examined the pattern of cortical response to electrical microstimulation

administered at the centre of the implanted Utah array during these

different states. Trial and channel averaged peri-stimulus time histograms

(PSTHs) for spiking activity (Figs. 3A - PFC & 3C - PPC), and the same for

LFPs (0.1 to 250Hz) (Fig. 4A - PFC & 4C - PPC) were computed aligned at

stimulation offset.

For lower microstimulation amplitudes viz. 1, 5 and 8 µA, the baseline-

normalised spiking activity displayed a suppression during the

microstimulation period, followed by a sharp excitatory transient

immediately after the end of microstimulation for all consciousness states

(QW, LS and DS), in both PFC and PPC. Perhaps counterintuitively, the

excitatory transient was stronger in amplitude for DS compared to QW and

LS, in both recorded regions. However, for higher amplitudes (15-25 µA),

we observed a different pattern present again in all consciousness states.

In the PFC, spiking activity increased during the microstimulation period,

which was then followed by a period of long-lasting inhibition (~1-

1.5seconds), finally bookended by a later rebound. This rebound was the

lowest during light anaesthesia, and highest during deep anaesthesia in the

PFC (Fig. 3A). Interestingly, while only one rebound occurred in the PFC,

two rebounds manifested in the PPC, with the inhibition and rebound

occurring faster and lasting for a shorter time (Fig. 3C). Furthermore, both

the transient and the rebounds in the PPC followed a gradient according to

the state, i.e. DS > LS > QW.

The above-mentioned characteristics are summarised as violin

distributions (Fig. 3B - PFC, and Fig. 3D - PPC), estimated using an n-peak

detection algorithm. In the PFC, for the lower stimulation amplitudes, the

response during QW was the fastest, followed by during DS, with the

slowest response manifesting during LS (Fig. 3B). However, in the PPC, the

response occurred earlier in LS than during QW, with DS being the fastest.

(Fig. 3D). Furthermore, the strength of the responses was highest during

DS, which can potentially be attributed to increased cortical membrane

sensitivity during burst suppression periods[80,81]. While in the PFC, the

response and rebound strengths were weakest during LS, with DS being the

highest, in the PPC, again, the response strength showed an increasing

trend with anaesthesia. The mean occurrence times, their strengths, and

their dispersal, under QW, LS and DS, for both Monkey A (PFC) and Monkey

C (PPC) are catalogued in SI Table 1 (PFC) and SI Table 2 (PPC), along with

confidence intervals (CIs) for the difference in means (If a CI includes 0,

then no significant differences between the two states are seen).

The LFPs also displayed stimulation-dependent heterogeneity in the

response pattern and strength, both in the PFC (Fig. 4A) and the PPC (Fig.

4C). For lower amplitudes (1, 5 and 8 µA), following stimulation offset, a

negative deflection is seen, often followed by a positive rebound. However,

these responses are noisy (Fig. 4C insets), and highly variable. For higher

amplitudes (15-25 µA), following stimulation offset, a strong positive

deflection is elicited followed by a delayed negative dip. These responses

are highly stereotypical and consistent (Fig. 4C insets). While the first

transient happened earliest during QW, and latest during DS, the reversal

occurred latest during QW and earliest during DS, in both the PFC and the

PPC. Furthermore, the initial response transient was sustained for the

longest time (peak width duration) during QW, while it resolved itself the

fastest during LS, pointing to its role in modulating membrane potentials

of the constituent neurons of the population in the slow rhythm range (~0.5-

2Hz; data not shown). The mean occurrence times, their strengths, and

their durations, along with their dispersal, under QW, LS and DS, for both

Monkey A (PFC) and Monkey C (PPC) are catalogued in SI Table 3 (PFC) and

SI Table 4 (PPC) respectively, along with confidence intervals (CIs) for the

difference in means (If a CI includes 0, then no significant differences

between the two states are seen).

In summary, these results show that causal perturbation via targeted

electrical stimulation elicits state, stimulation strength, and space

dependent heterogeneity in the response activity of the fronto-parietal

network.

Spatiotemporal modulation of field activity in different

consciousness states

To study whether the state of consciousness has an effect on lateral

signal propagation in the two cortical regions, the electrodes on the Utah

array were binned according to their distance from the stimulated

electrodes in 0.4mm bins, from the nearest (0.4 to 0.8mm) to the farthest (2-

2.4mm in PFC and 2.4-2.8mm in PPC). Responses to ICMS were pooled for

all stimulation amplitudes in Monkey A and Monkey C, while in Monkey B,

only the response to 15 µA stimulation was analysed (only 2 amplitudes

were tested in Monkey B viz.15 and 100 µA. 100uA displayed complete

suppression in Monkey B, and thus was discarded from the analysis). The

field response profiles by distance to the stimulated electrode are displayed

in Fig. 5A (PFC) and 6A (PPC) respectively, with 2 stimulation amplitudes

being represented, viz. 8 and 15 μA.

Because of the distinct response patterns to the low and high amplitude

regimes elicited in the spiking activity, viz. a simple transient excitation

versus a transient followed by inhibition, and rebound, in both the PFC and

PPC, we restricted the analysis of the spatial modulation of population

activity to the LFPs.

LFP signal energy modulation over distance was computed as the

normalised pre/post change in activity in a 4s bin following the end of

stimulation. The signal response was modulated strongest during QW,

displaying non-monotonicity over distance, while they were modulated the

least during DS, and decreased sharply over distance (Fig. 5B). Neuronal

activity in LS displayed modulation strength in between QW and DS, and

also decayed linearly over distance, albeit slower than DS (Fig. 5B). This

points to a maintenance of the strength of modulation at longer distances

during QW, with proximal and distal populations being modulated equally,

while signal propagation is restricted with increase in distance, in both

states of anaesthesia. These are confirmed by the analysis of the sum of the

rate of change over distance, combined with a Bayesian bootstrapped

comparison-of-means. The spatiotemporal characterisation for Monkey B

is presented in SI Fig. 4.

In the PPC however, modulation decreased with distance in all three states

pointing to no non-monotonic recovery of signal propagation with distance

(Fig. 6B). However, as opposed to the PFC where in the LFPs, the strongest

modulation was observed during QW (Fig. 5B), in the PPC, the strongest

modulation was seen in the DS.

A 0.5s bin by bin analysis of this modulation in time in the PFC, starting

immediately after stimulation offset, displays the preservation of this non-

monotonicity during wakefulness (Fig. 5C; Left Column), even up to 6s after

stimulation offset, in the LFPs. However, during both levels of anaesthesia,

the drop off of signal energy modulation over distance remained in all time

bins (Fig. 5C Middle and Left Columns). In the PPC, the decay in time was

even faster, strong excitatory modulation extinguishing already after ~2.5s

(Fig. 6C).

Taken together, our results suggest that the prefrontal cortex not only

displays distinct differences in responses in spiking activity and LFPs to electrical microstimulation, it also displays differential responsivity to low and high amplitudes of stimulation, under different states of (un)consciousness. Most importantly, our data show that while field signal propagation in the PFC is starkly reduced under anaesthesia, as compared to wakefulness, only the strength of modulation displays significant differences between the states in the PPC.

Discussion

The PFC exhibits a rich diversification of structure[82–84], and consequently, function. It serves as an integrative hub for a multitude of cognitive processes such as inhibitory control, decision making, cognitive control, working memory, and the integration of sensory information with planned actions[85–91]. Under anaesthesia, both rodent and human studies have revealed notable alterations in PFC activity. Specifically, anaesthesia has been shown to induce a transition from desynchronised to highly

synchronised oscillatory activity in the PFC of rodents[72,92,93]. In humans,

anaesthetic agents have been observed to disrupt functional connectivity

within the PFC, which correlates with a decline in conscious awareness and

cognitive function[43,94,95]. Pressingly, what can we learn from characterising

macaque PFC population dynamics by monitoring and perturbing

neuronal populations during these (un)conscious states?

Altered mesoscale cortical dynamics under anaesthesia

Previous studies have monitored cortical populations across the dorsal and

ventral pathways in different animal models during rest. Furthermore,

global dynamics have been studied using gross-scale techniques in

humans. Perturbational studies in humans and monkeys under different

(un)conscious states have been performed with both DBS and TMS[73,96–99].

However, to date, consistent, simultaneously-recorded mesoscale cortical

populations in different states have not been monitored, and perturbed.

We show that chaotic and dispersed spiking activity in the same population

during the awake state reorganises into coherent UP/DOWN states during

anaesthesia, both during light and deep anaesthetic states (Fig. 2). The

distinguishing feature between LS and DS both in the PFC and PPC seems

to be a long tail of the duration distributions of the down states during DS

(Fig. 2), thus signifying a quieter, i.e. less active cortex. However, a higher

population synchrony in DS agrees well with the significantly larger low-

frequency power (Fig. 2, delta and theta), and therefore coherence (data

not shown). These long periods of cortical silence, termed burst

suppression in neurological EEG literature, could reset cortical

responsivity during anaesthesia at the mesoscale.

Response characteristics of the PFC and PPC

In the past, a majority of causal perturbational studies have been used

either to bias perception and behaviour by stimulating stimulus-selective

neurons in sensory cortices[74,100–104] or object recognition areas[105] , or for

brain computer interfaces (BCIs)[106], and therapeutic purposes[107–109]. While

studies that electrically and pharmacologically perturb the PFC in

humans[36,110] and animal models[10,111] exist, they are sparse and not

systematised, neither in the behaviour or sub-behavioural range, nor

under different arousal states. Our study addresses this *lacuna* specifically

with respect to intracortical microstimulation by assaying the primate

higher associational cortices. Crucially, no studies exist that simultaneously

monitor and perturb the same consistent population of prefrontal and

parietal neurons under different states of consciousness, thus targeting the

fronto-parietal loop.

Our results (Figs. 3A and 3B) suggest the existence of two membrane-

responsiveness regimes with respect to electrical challenge in these cortical

populations. We show that low amplitude stimulation seems to display an

initial inhibition followed by excitation (Fig. 4A - Left). However, large

current injections elicit first a strong transient, followed by a long-lasting

inhibition, after which the spiking activity rebounds with a vengeance (Fig.

4A - Right). Inhibitory neuronal populations, though comprising only 15-

30% of the cortex[112,113], tightly regulate microcircuit function[114,115]. Post-

inhibitory rebound and its relationship to recurrent inhibition is widely-

known in a variety of intrinsically and extrinsically-driven contexts[116].

Large current insults could reflect a disruption of this inhibitory control,

and its subsequent release, pointing to an inhibitory population led

network reorganisation[117,118]. This heterogeneity, also preserved in the LFPs

(Figs. 4A and 4B), could therefore point to downstream perisynaptic

modulation, allowing the cortical membrane potential to display

differential activity as a consequence.

In the PPC, in addition to the transient-inhibition-rebound, we observe a

second rebound that happens very close to the first (Fig. 3C). Because the

firing rate before this rebound does not return to baseline, and instead

manifests as a "second peak", this extra rebound could potentially arise

from recurrent excitation, and could be crucial in self-generated and

sustained neural activity, that can be broadcasted to organise on-line

circuit control[119,120].

The responsivity paradox

Our data also indicate that fronto-parietal responsivity is highest during

deep anaesthesia. What seems at first paradoxical, lends itself to a

remarkable interpretation in terms of equilibrium dynamics. During deep

anaesthesia, long periods of iso-electric voltage lines (in the EEG)

interspersed with brief, large fluctuations exist (Fig. 2A), which is termed

"burst suppression"[121]. We speculate that during these extended periods of

cortical silence, the neuronal membrane is hyper-sensitive[80], thereby

allowing it to return to equilibrium, during which, any reasonably strong

perturbation could potentially push the system into an accelerated,

hyperbolic trajectory, manifesting as fast and strong responses. Such

empirically-determined cortical activity patterns under perturbation, rest

and different states are imperative for constraining models of brain

function at multiple scales.

Contrasting resting state activity profiles with the observed response

dynamics underscores the potential limitations in inferring the exact state

of consciousness (for e.g., depth of anaesthesia) from temporal patterns in

population activity. Despite the varying distributions of quiescent intervals

across sessions and subjects, the pattern of increased responsivity coupled

with limited spatial propagation during deep sedation (DS) was a consistent

finding. This observation is bolstered by the analysis of proximity to

criticality (see SI. Fig 1), which showed uniform spatial and temporal

response characteristics across sessions and subjects, regardless of any

significant shifts in criticality between light sedation (LS) and DS.

Moreover, the hypothesis that direct stimulation may induce prolonged

suppression, culminating in a synchronised and markedly amplified

rebound, suggests that during DS, complexity indices - evaluated through

methods like perturbational complexity index (PCI) and pattern analysis -

would be reduced due to this enforced, homogenous, and intensified

rebound effect. Consequently, beyond a certain depth of anaesthesia,

where neuronal populations reach a threshold of inhibitory saturation, a

progressive decline in criticality or complexity metrics may not be evident.

In effect, these metrics potentially only reflect this specific inhibitory

population led dynamics, rather than a true reconfiguration of

information, and thus complexity. Future investigations could explore

these dynamics through BOLD-fMRI under varied anaesthetic conditions

and DBS protocols, coupled with advanced PCI evaluations using complex

TMS protocols, including pulse-train stimulation, to thoroughly examine

and correlate activities across different scales.

Signal communication is maintained during wakefulness in PFC but not in PPC

Anaesthesia is speculated to induce unconsciousness in many ways[47,60,122], a

primary one among which is by suppressing information processing and

communication. This is evidenced by global inhibition, i.e. the reduction in

neuronal spiking activity[123,124]. Additionally, local synchrony increases but

global synchrony has been observed to be disrupted under anaesthesia[125],

while during wakefulness, although locally neurons remain uncorrelated

in their firing, large-scale synchrony is observed across brain regions[126–128],

specifically with the persistence of the Default Mode Network (DMN)[129].

This change in synchrony patterns under anaesthesia suggests a disruption

in normal information processing in the brain, which at the prefrontal

cortical level is confirmed by our results.

Crucially, the dorsal and ventral PFC have been shown to contain features

that distinguish them from lower cortical regions[130]. For example, instead

of distinct organisation into cortical columns, the PFC instead shows

greater lateral connectivity among distributed pools of functionally distinct

neuronal cell types[83,131–133]. Layer 5 pyramidal neurons project out of the

cortex and receive extra-cortical and subcortical efferents to establish

overarching connectivity[134,135.136]. Long-range fronto-parietal connections

exist which form the fronto-parietal loop, a unit central to the GNW[14,16,137].

Most importantly, the PFC displays not only abstract tuning such as task-

phase[138], but also stimulus features[139]. These various modes of

differentiation and integration[70] are ideal for complex information

processing, such as those required for conscious awareness[13,140]. A unique

and computationally significant property of neuronal networks is

functional connectivity, which can be measured using noise correlations at

the mesoscale[141,142]. The structure of this functional connectivity in lower

cortical areas is a linear decrease with lateral distance, both under

wakefulness and anaesthesia, whereas in the PFC it is non-monotonic,

showing that proximal and distal populations are equally connected[143,144].

This is also evidenced by patchy anatomical connectivity between pools of

excitatory and inhibitory populations[83,131,133,145]. This unique pattern of

connectivity in the PFC suggests that it can be used for complex integrative

functions[87,146–148] that lower order areas cannot perform. Interestingly,

primary sensory areas, with the exception of the somatosensory cortex,

aren't much perturbed during anaesthesia, even at deep levels[149,150]. These

observations lend support to the hypothesis that loss of consciousness is

triggered by a lack of higher-order integration in the cortex[66,151]. Lower

areas don't need long-range functional connectivity because they process

low-level stimulus features independently and don't need to integrate.

However, feature-binding (Gestalt) is necessary for conscious perception[152]

[−156], the PFC being a prime candidate[157–159].

From the standpoint of GNW, although the PPC is indeed part of the fronto-

parietal loop[43], and distinct patterns of directionality, strength and

functional organisation are observed across states of consciousness in each

individual module of the loop[42,43], it is the PFC that is supposed to amplify,

broadcast and coordinate brain-wide activity to allow for consciousness to

emerge 15[15,17]. Therefore, we posit that anaesthesia must necessarily

disturb the PFC, not that it is only sufficient to dampen PFC or PPC activity.

If anaesthesia causes an information bottle-neck, long-range coordination

of neuronal activity by an orchestrating node would become weak or

impossible. Our results (Fig. 5) show – that although fast spatial spread of

global activity in V1 is seen immediately post ignition – that strong

horizontal connectivity and lateral information flow, only in the PFC, is a

signature of conscious wakefulness and indeed breaks down under

anaesthetic unconsciousness, suggesting that such complex functional cytoarchitecture in frontal regions is necessary for integrative processes crucial to conscious awareness, which subsequently implicates its disruption in loss of consciousness.

Causal perturbation of the fronto-parietal circuit and its systemic implications

In the ongoing discussion surrounding the role of the PFC in consciousness and DoCs, studies employing intracranial electrical stimulation (iES) to causally perturb networks of interest have recently begun to not only provide unique, unexpected and contradictory insights[110], but also elicit spirited debate[36]. Most importantly, studying cortical response properties under external stimulation can help elucidate the mechanistic underpinnings of consciousness, as opposed to purely correlative studies. Causal perturbational studies are crucial as they allow for the manipulation of specific neural populations, thereby providing direct evidence of their role in consciousness. To even begin to design appropriate experiments addressing this large gamut of questions, it is vital that a

foundation is securely built on thorough electrical and optical

characterisation of these mesoscale networks. Furthermore, insights

gained from such studies can be used to constrain larger neuronal models

of cortical activity under different states of consciousness.

Studies employing iES or ICMS in animal models have also provided

indirect support for a central role of the PFC in conscious awareness. For

example, it has been shown that when the primary visual cortex (V1) is

stimulated, correct detection trials consistently display sustained PFC

activity as opposed to misses[75,76]. In terms of the state of consciousness, a

landmark study in rodents demonstrated robust RoC after carbachol

injection to the PFC, but crucially, not the PPC[10]. Interestingly however,

when the thalamus is stimulated using deep brain stimulation (DBS) to

arouse the experimental subject, both the PFC and the PPC are strongly

activated, indicating the engagement of fronto-parietal populations during

RoC[44,160,161]. In task contexts for instance, global deviants in the local-global

task[162], which are a signature of conscious access, can then be decoded from

the PFC when other behavioural signs of arousal re-manifest[44]. DBS has also

shown promise in "enhancing" conscious access in patients with minimally

conscious state (MCS), marking a significant advancement in DoC

intervention[96]. These findings trace a functional link between the thalamic

relay and higher associational populations in conscious access, potentially

engaging a broader thalamo-cortical circuit, thereby restoring

consciousness in patients with DOCs, while also providing a more nuanced

approach in identifying potential therapeutic targets, such as cautiously

employing direct cortical stimulation - rather than strongly invasive and

high-strength DBS - could. However, systematic protocols for characterising

ICMS effects during wakefulness or anaesthesia (as a proxy for DoCs) are

lacking. Our study attempts to fill this gap by exploring myriad stimulation

protocols and parameters (Fig. 1).

Limitations

In our study, we recorded stable prefrontal and parietal populations during

all states of (un)consciousness. However, it is prudent to note that the PFC

and PPC data come from two different animals, due to the unfortunate loss

of the PFC and the PPC array in each animal respectively (COVID-19 *etc*; in

the other animal, no spiking activity could be recorded from the PPC array).

This inherently limits the direct comparison of the two areas forming the

frontoparietal loop. However, it is our firm belief that the consistency of

responses and functional modulation over space and time seen in our data

would predict the same across simultaneous recordings too.

Furthermore, since we used stimulation regimes that have previously been

shown to elicit behavioural responses[75,76,163–165], we did not probe the cortex

classically with single-pulse stimulation, as our goal was to also investigate

response characteristics and signal spread under these un(conscious)

states.

Conclusion

Taken together, our results reveal that the PFC and the PPC exhibit unique

and intriguing responses when subjected to electrical stimulation,

specifically shedding light on the putative functional role of the PFC in

regulating states of consciousness. Most importantly, we show the

restriction of lateral signal propagation under anaesthesia in the PFC,

which could result in break-down of crucial integrative processes that are

necessary for consciousness and cognition. These results can further

significantly inform and constrain cortical models of brain function in

health and disease.

Further investigation into the effects of elevated electrical stimulation

intensities on these higher order association areas offers a promising

avenue for future research. By delving into the consequences of

stimulation whose parameters are in the range to elicit behavioural effects,

we have the opportunity to elucidate the nuanced relationship between

PFC activity and specific behavioural metrics, particularly in arousal and

its loss. This line of enquiry holds substantial clinical implications,

especially for individuals afflicted with DoCs. Given that our current

stimulation design did not demonstrate behavioural signs of arousal (this

is demonstrated with a different stimulation regime, in forthcoming work)

or long-range cortical excitability, we hypothesise that higher stimulation

intensities or alternative stimulation patterns, as guided by dynamic brain

activity[100,166,167], could potentially increase cortical excitability and thus

awaken subjects under anaesthesia. To this end, we are initiating further

studies aimed at assessing the feasibility of reversing anaesthesia-induced

unconsciousness through targeted electrical stimulation of the

frontoparietal loop.

618
619
620
621

Figure Captions

Figure 1 | Experimental design and ROI

A. Multielectrode array in the inferior convexity of the PFC, thus targeting

the fronto-parietal loop, placed anterior to the bank of the arcuate sulcus

and below the principal sulcus, thus being localised to the left ventrolateral

prefrontal cortex

B. Cortical population activity was recorded in four experimental phases,

viz. Quiet wakefulness (QW), transition to anaesthesia and preparation,

light anaesthesia (LS - 1-1.5% Sevoflurane), and deep anaesthesia (DS - 3%

Sevoflurane), along with the delivery of intracortical microstimulation. The

stimulation was devised as a train of 50 biphasic pulses lasting 200ms, with

varying amplitudes, presented randomly with an inter-stimulation interval

between 10-20s.

C. At the start of each experimental epoch, 15-20 mins of resting state

activity was recorded, followed by the three stimulation regimes, viz.

Centre-array stimulation, full-array stimulation, and full-array high-

frequency, high-amplitude stimulation, finally followed by 15-20 mins of

post-stimulation resting state.

Figure 2 | PFC and PPC display distinct neuronal population

activity patterns under different conscious states

A. PFC - Reorganisation of spiking activity into periods of UP and DOWN

states in anaesthesia (LS - Green, DS - Blue) as compared to desynchronised

activity during QW (Red). DS also displays extended periods of cortical

silence (seconds to minutes) termed "burst suppression".

B. PFC - Autocorrelograms of the population firing vector computed in

chunks. While LS shows a wider peak, the strength of the non-shifted

correlation is highest during DS and lowest during QW.

C. PFC - z-scored time-frequency wavelet decomposition of the

corresponding LFPs. While QW is identified by its characteristic ongoing

high-frequency power which follows spiking, this is suppressed during

both types of anaesthesia, where slower rhythms dominate. The purple line

indicates the learned UP state path from the 2-state HMM. *n.b.* - Shown is

one example period from Monkey A.

D. PFC - z-scored spectrum for the example epochs. While all frequencies

are well-represented during QW, especially beta and gamma, during LS

and DS, the spectral activity is concentrated in the 0.1-8 Hz range.

E, F G and H depict the same as above, but for the PPC.

I. Col 1 & Col 2 - Duration of DOWN and UP states in the PFC. Col 3 & Col 4 -

Duration of DOWN and UP states in the PPC. Green is LS, Blue is DS. While

during both LS and DS, the durations of the up states stay below 3s, the

durations of the silent periods extend to 15s (PFC) and 100+s (PPC) during

DS.

Figure 3 | Spiking characteristics

A. Peri-stimulus time histograms (PSTHs) computed for all stimulation amplitudes (1, 5, 8, 15, 20, 25μA) during QW, LS and DS, in the PFC. The firing rates were averaged over channels and trials, and aligned to stimulation offset. Insets are zoomed in on the response shape.

B. This panel characterises the above activity as violin distributions, estimated using an n-peak detection algorithm. It represents the times at which the peak transients and rebound occurred. While the cortical population response to low amplitude stimulation is temporal fast, transient, and exhibits an initial inhibition followed by excitation, higher amplitudes elicit an initial transient followed by long lasting inhibition, after which the discharge activity rebounds.

C. Same as A, for PPC.

D. While the cortical population response to low amplitude stimulation is temporal fast, transient, and exhibits an initial inhibition followed by excitation, higher amplitudes elicit an initial transient followed by short inhibition, after which the discharge activity rebounds, followed by another rebound. This does not occur in the PFC.

Figure 4 | LFP characteristics

A. LFP activity for all stimulation amplitudes (1, 5, 8, 15, 20, 25μA) during

QW, LS and DS, in the PFC. Insets are zoomed in on the response shape.

B. This panel characterises the above activity in terms of the timing of the

induced up or down state post stimulation. While the cortical population

response to low amplitude stimulation first elicits a negative deflection

followed by a positive overshoot, this pattern is reversed for higher

amplitudes.

C. Same as A for PPC

D. Same as B for PPC

(*n.b.* - In both the LFPs and neuronal discharges, the heterogeneous

response patterns can be grouped into two distinct groups, viz. those in

response to "low" amplitudes (1,5,8 μA) and those in response to "high"

amplitudes (15, 20, 25 μA).)

Figure 5 | Modulation of cortical information flow in different

states - PFC

A. Shows the average LFP response aligned to stimulation offset in the three

states, viz. quiet wakefulness, light and deep anaesthesia, for two example

amplitudes from each regime, i.e. "low" and "high" stimulation, sorted by

distance, from nearest (0.4mm) to farthest (2.6mm) nodes from the centre.

As noted previously, stimulation was delivered to the two centre

electrodes.

B. Whereas during QW, proximal and distal populations show similar

modulation of field activity, this modulation falls off over space during both

types of anaesthesia, with the decrease being stronger during DS (Insets

show the sum of the first derivative for the proximal and distal

populations). Data points show the mean and SEM.

C. Top - Shows the evolution of field signal energy modulation in 0.5s bins

starting immediately after stimulation offset, until 7s after stimulation.

While the non-monotonicity is preserved in almost all time bins during

QW, it does not manifest during LS or DS at all, showing a lack of spatial

spread of information.

Figure 6 | Modulation of cortical information flow in different

states - PPC

A. Shows the average LFP response aligned to stimulation offset in the three states, viz. quiet wakefulness, light and deep anaesthesia, for two example amplitudes from each regime, i.e. "low" and "high" stimulation, sorted by distance, from nearest (0.4mm) to farthest (2.8mm) nodes from the centre. As noted previously, stimulation was delivered to the two centre electrodes.

B. Field signal energy modulation in the PPC shows that no non-monotonic structure is observed at all in the LFPs in the PPC, as opposed to the PPC. Furthermore, As opposed to the PFC, in the PPC, DS modulation remains higher than QW and LS.

C. While the strength of modulation persists for longer after stimulation in the PPC during QW, it extinguishes fastest during LS, and then DS. No non-monotonic structure is observed in any time-bin in any state after stimulation.

Acknowledgments

732   We thank Dr. Lynn Uhrig for her assistance with anaesthesia. We are

733   thankful to Jeremy Bernard for assistance with the custom-made setup for

734   the experiment. We would also like to thank Prof. Dr. B S Dwarakanath for

735   helpful discussions and comments, and proof-reading assistance.

736   Author contributions

Conceptualisation: AD, TIP ; Data curation: AD; Formal analysis: AD;

Funding acquisition: BJ, SD, TIP ; Investigation: AD, MKA, MG, MR, RHV, BJ,

TIP; Methodology: AD, MKA, RHV, TIP; Project administration: TIP;

Resources: MG, MR, BJ, SD, TIP (lead); Software: AD; Supervision: BJ, TIP

(lead); Visualisation: AD (lead), RHV (supporting); Writing - original draft:

AD; Writing - review & editing: AD, MKA, MG, MR, RHV, BJ, TIP (lead).

743   Funding

This work received funding from Inserm (BJ), CEA (TIP), Collège de France

(SD), the European Union's Horizon 2020 Framework Programme for

Research and Innovation under the Specific Grant Agreement No. 945539

(Human Brain Project SGA3, T2.10; TIP), a grant from Templeton World

Charity Foundation, Inc (TWCF0562) to TIP, and a grant from the

Bettencourt-Schueller Foundation to BJ.

750 Declaration of interests

751 The authors declare no competing interests.

753 STAR Methods

754 1.   Resource availability

755 Information and requests for data and specific code should be directed to

756 and will be fulfilled by the lead contact or the first author –

757 Theofanis I Panagiotaropoulos - theofanis.panagiotaropoulos@cea.fr (lead

758 contact); Abhilash Dwarakanath - abhilash.dwarakanath@cea.fr

759 2. Materials and availability –

760 Not applicable

761 3. Data and code availability –

Electrophysiological recordings used in this paper will be used in the future for other studies. Therefore, the data is available upon reasonable request; please contact the lead contact. Custom stimulation, data acquisition and extraction, preprocessing, and analysis scripts written in MATLAB are available on the first author's GitHub - (https://github.com/abhilashdwarakanath/IntracorticalMicrostimulation, https://github.com/abhilashdwarakanath/-Un-ConsciousStates, https://github.com/abhilashdwarakanath/EphysPreprocessing).

The data used in this manuscript will be made available through the Human Brain Project's EBRAINS Knowledge Graph, as a contribution from SGA3's Task 2.10.

4. Experimental model and subject details –

Two adult, male rhesus macaques (*Macaca mulatta)*, weighing between 10-12kg were recruited for the experiments. The subjects, denoted as Monkey "A" and Monkey "J", were pair-housed in an enriched animal facility. They were trained to be chaired and head-posted in the recording chamber.

5. Methods details -

Electrophysiological recordings

We performed extracellular electrophysiological recordings in the inferior convexity of the lateral PFC of 2 awake and anaesthetised monkeys using chronically implanted Utah microelectrode arrays[168] (Blackrock Microsystems, Salt Lake City, Utah USA). We implanted the arrays 1 - 2 millimetres anterior to the bank of the arcuate sulcus and below the ventral bank of the principal sulcus, thus covering a large part of the inferior convexity in the ventrolateral PFC, which forms one node of the fronto-parietal loop. The arrays were 4x4mm wide, with a 10 by 10 electrode configuration and inter-electrode distance of 400μm. Electrodes were 1mm long therefore recording from the middle cortical layers. The monkeys were implanted with form-specific titanium head posts on the cranium after modelling the skull based on CT scans. All experiments were approved by the local authorities' ethical committee (Protocol # - A18_028), and were in full compliance with the guidelines of the European Community (EUVD 86/609/EEC) for the care and use of laboratory animals.

Data acquisition and stimulation

Broadband neural signals (0.1–30 kHz) were recorded using Neural Signal

Processors (NSPs) from Blackrock Microsystems Inc. Signals from the Utah

array were digitised, amplified, and then routed to the NSPs for acquisition.

The digitisation occurred on a CerePlex M Headstage which, coupled with

Omnetics Y adapters, allowed us to simultaneously record and stimulate

from the same electrodes. A CereStim 96 microstimulator from Blackrock

Microsystems driven by MATLAB (2022a, MathWorks Ltd) code interfaced

with their proprietary API was used to provide biphasic pulses between 1

to 210μA.

For the offline detection of action potentials, broadband data were filtered

between 0.6 and 3 kHz using a second-order Butterworth filter (the filter

was chosen such that it allowed a flat response in the passband while

contributing the least phase distortion due to its low order, yet having an

acceptable attenuation in the stop band, i.e. a roll-off starting at -20dB). The

amplitude for spike detection was set to five times the median absolute

deviation (MAD)[169]. Spikes were rejected if they occurred within 0.5 ms of

each other, (multi-unit refractory period temporal threshold) or if they

were larger than 50 times the MAD. Finally, all spikes that occurred at the

same time point (within 3 samples at 30kHz) in a minimum of 60 channels

(out of 96) were also discarded to exclude stimulation artefacts. All

collected spikes were aligned to their minimum. For the analysis of LFP

activity, the broadband signal was decimated to 500 Hz sampling rate using

a Type I Chebyshev Filter, reliably preserving frequency components up to

200 Hz.

Experimental design

The animal was chaired and head-posted in the recording chamber, after

which the eye-recording camera was aligned and centre-calibrated. The

CereStim module was connected to the CerePlex M headstage for

stimulation delivery, which was subsequently connected to the pre-

amplifiers, which fed the recorded and referenced signals to the Neural

Signal Processor (NSP) via fibre-optic cables.

The experiment consisted of four distinct phases viz. Quiet

wakefulness (QW), induction and transition to anaesthesia, light

anaesthesia (LS) and deep anaesthesia (DS). Each of the QW, LS and DS

phases started with an initial recording of 15-20 mins of spontaneous

activity. Next, multiple repeats (10 to 25) of a pulse train consisting of 50

pulses of amplitudes varying between 1 to 100µA were delivered to the two

central electrodes, randomised, separated by 10-20s of dead-time. Each

pulse was biphasic, being either anodal first, or cathodal first. Each phase

lasted for 50µS with the inter-phase interval being 53µS. Therefore, each

stimulation pulse lasted for ~0.1995s. Following certain other stimulation

protocols, a post-stimulation spontaneous phase of 15-20 mins was again

recorded.

Transition to anaesthesia from wakefulness was achieved first by

sedating the animal using a combination of Ketamine 1000 (3 mg/kg

intramuscular) and Dexmedetomidine (15 µg/kg intramuscular). The

animal was then released from head-fixation and carefully extracted from

the experimental chair. The chair was then removed and a custom-built

experimental bed was inserted and fixed. This bed was covered with a

blanket and an inflatable heating pad was inserted. An injection of

Antisedan (75 μg/kg intramuscular) was then given, as an antagonist of the

sedative effect of Dexmedetomidine. In one experiment, the animal was

laid in a prone position inside a stereotactic frame, while in the rest of the

sessions, the animal was laid on its side. Next, the animal was intubated

and an IV line was laid. Inhalation anaesthesia was initiated using 2% of

Sevoflurane. The ventilator was switched on, and on confirmation of stable

vitals, the door of the faraday cage was closed. After ~80-90 minutes from

the first injection, the LS experimental epoch was started by changing to

and maintaining the anaesthesia at a low level of 1-1.5%. In one session, a

paralytic agent, namely Curare, was used. After the end of the LS epoch, the

inhalant anaesthesia was increased to 3% (DS). Crucially, we continued to

record during this entire process thereby making sure that we were

monitoring the same population of neurons throughout the 4 experimental

phases.

At the end of the DS epoch, the inhalant anaesthesia was gradually reduced

to 0. Upon awakening, the monkey was extubated, the IV line and rectal

thermometer was removed, and disconnected from the ventilator and

vitals monitor. Next, the animal was placed in the recovery box and

returned to its cage. Finally, the animal was monitored for full recovery for

the next 24 hours.

Evaluation of resting state activity

To statistically characterise the effects of the levels of anaesthesia,

we fit a 2-state Hidden Markov Model (HMM) to the data. A firing rate

threshold of 0.05 spikes/s (Monkey 1) or 2.5 spikes/s (Monkey 2) was set to

exclude non-firing channels in the resting state periods, after which spikes

were binned in 25ms (Monkey 2) or 50ms (Monkey 1) bins, and then

summed and normalised to obtain the population firing vector. To this

population vector for each state, k-means clustering was applied to obtain

2 distinct firing rate regimes. These cluster-centres were normalised and

used as the initial emission probabilities ($p(EM_{init})$). Initial transition

probabilities ($p(TR_{init})$) were drawn from a positive-normal distribution. An

initial state sequence guess was obtained by setting all data above the mean

to 2, and those below to 1. The HMM was trained on this data to obtain the

learned $p(EM_{final})$ and $p(TR_{final})$. These learned probabilities were then used

to generate the learned state sequence using the Viterbi algorithm, from

which the durations of the UP and DOWN states were finally computed.

This training and testing were performed using the HMM toolbox in

MATLAB.

State time-frequency information was computed via a continuous wavelet

transform using a generalised morse wavelet with 7 cycles and a time-

bandwidth product of 3 x 30 at a sampling rate of 500Hz.

The measure of synchrony (Synchrony Index) was estimated as the sum of

the positive-lag auto-correlation function (computed to a 0.5s lag) divided

by the number of lags, after which it was subtracted from 1. We consider

only the positive lags after 0, therefore we subtract the mean auto-

correlation from the peak-value (i.e. 1 at lag-0 when the ACF is

normalised).

Computing stimulation-related measures

To obtain the onset of each stimulation, the recorded pulse onset-offset

analogue signal was differentiated, and zero crossings were detected. These

onsets and offsets were sorted into their respective amplitude trials using

the trial structure log file corresponding to each epoch. LFPs and spikes

were collected for each trial and channel, and aligned to the stimulation

offset. Resting state epochs were segmented from the recording using their

respective epoch tags.

Mean responses to each stimulation amplitude were computed by

concatenating the channel and trial LFPs and PSTHs and then averaging

them. The errors were thus the standard deviation across these collapsed

samples normalised by the square root of the number of channels, and

finally multiplied by 1.96 to obtain the 95% CI.

Response characteristics of the neuronal discharge patterns, namely the

first transient and subsequent rebound(s), and the LFP response, were

characterised using MATLAB's in-built *findpeaks* function. The dispersal

was calculated as the distance to the cluster centre(s), which roughly

corresponds to the variance of a normal distribution. For multi-peak

rebound, k-means with n clusters was run to fit the underlying normal

distributions.

LFP and spiking response characteristics comparisons were performed

pairwise across state combinations (QW vs LS, QW vs DS, LS vs DS), using

bootstrapped resampling. The differences were evaluated as the 95%

confidence intervals (CI) on 10000 resampling runs.

LFP modulation -

where x(t) represents the LFP signal, k represents the trial index, and n

represents the channel index.

The existence and strength of non-monotonicity in the spatial spread of

field activity modulation was evaluated using a Bayesian bootstrap on 4

hypotheses, viz. flat, increasing, decreasing, and non-monotonic. Because

the distances increase in even number pitch increments, the data and

function were divided into "nearby" and " far away" populations. The sum

of the first differential was computed on these chunks. A 0 or positive index

indicates non-monotonicity, whereas the magnitude of the negative sum

denotes the steepness of the monotonic decrease. After 10000

bootstrapping runs, the hypothesis with the highest probability (above

0.25) was chosen as the most probable curve shape.

The MATLAB statistics and machine learning toolbox was used to

implement all statistical analyses, with custom-written scripts.

References

1.   Koch, C., Massimini, M., Boly, M. & Tononi, G. Neural correlates of consciousness: progress and problems. *Nat. Rev. Neurosci.* **17**, 307–321 (2016).

2.   Boly, M. *et al.* Are the neural correlates of consciousness in the front or in the back of the cerebral cortex? clinical and neuroimaging evidence. *J. Neurosci.* **37**, 9603–9613 (2017).

3.   Laureys, S. The neural correlate of (un)awareness: lessons from the vegetative state. *Trends Cogn Sci (Regul Ed)* **9**, 556–559 (2005).

4.   Boly, M. *et al.* Consciousness in humans and non-human animals: recent advances and future directions. *Front. Psychol.* **4**, 625 (2013).

5.   Edlow, B. L., Claassen, J., Schiff, N. D. & Greer, D. M. Recovery from disorders of consciousness: mechanisms, prognosis and emerging therapies. *Nat. Rev. Neurol.* **17**, 135–156 (2021).

6.   Gosseries, O., Di, H., Laureys, S. & Boly, M. Measuring consciousness in severely damaged brains. *Annu. Rev. Neurosci.* **37**, 457–478 (2014).

7.   Giacino, J. T., Fins, J. J., Laureys, S. & Schiff, N. D. Disorders of consciousness after acquired brain injury: the state of the science. *Nat. Rev. Neurol.* **10**, 99–114 (2014).

8.   Seth, A. K. & Bayne, T. Theories of consciousness. *Nat. Rev. Neurosci.* **23**, 439

–452 (2022).

9.   Signorelli, C. M., Szczotka, J. & Prentner, R. Explanatory profiles of models

of consciousness - towards a systematic classification. *Neurosci. Conscious.* **2021**,

niab021 (2021).

10.   Pal, D. *et al.* Differential role of prefrontal and parietal cortices in

controlling level of consciousness. *Curr. Biol.* **28**, 2145-2152.e5 (2018).

11.   Odegaard, B., Knight, R. T. & Lau, H. Should a few null findings falsify

prefrontal theories of conscious perception? *J. Neurosci.* **37**, 9593–9602 (2017).

12.   Michel, M. & Morales, J. Minority reports: Consciousness and the

prefrontal cortex. *Mind & Language* (2019) doi:10.1111/mila.12264.

13.   Bartels, A. Consciousness: What is the role of prefrontal cortex? *Curr. Biol.*

31, R853–R856 (2021).

14.   Bor, D. & Seth, A. K. Consciousness and the prefrontal parietal network:

insights from attention, working memory, and chunking. *Front. Psychol.* **3**, 63

(2012).

15.   Dehaene, S. & Changeux, J.-P. Experimental and theoretical approaches to

conscious processing. *Neuron* **70**, 200–227 (2011).

16.   Mashour, G. A., Roelfsema, P., Changeux, J.-P. & Dehaene, S. Conscious

processing and the global neuronal workspace hypothesis. *Neuron* **105**, 776–798

(2020).

17.   Dehaene, S. & Naccache, L. Towards a cognitive neuroscience of

consciousness: basic evidence and a workspace framework. *Cognition* **79**, 1–37

(2001).

18.   Dehaene, S., Kerszberg, M. & Changeux, J. P. A neuronal model of a global

workspace in effortful cognitive tasks. *Proc Natl Acad Sci USA* **95**, 14529–14534

(1998).

19. Tononi, G. Integrated information theory of consciousness: an updated

account. *Arch Ital Biol* **150**, 56–90 (2012).

20. Albantakis, L. *et al.* Integrated information theory (IIT) 4.0: Formulating the

properties of phenomenal existence in physical terms. *PLoS Comput. Biol.* **19**,

e1011465 (2023).

21. Lau, H. Where is the 'posterior hot zone'? Open Review of Ferrante et al

(2023): "An Adversarial Collaboration to Critically Evaluate Theories of

Consciousness" (by the ARC-Cogitate Consortium). (2023)

doi:10.31234/osf.io/93ufe.

22. Lau, H. C. A higher order Bayesian decision theory of consciousness. *Prog.*

Brain Res. **168**, 35–48 (2008).

23. Brown, R., Lau, H. & LeDoux, J. E. Understanding the Higher-Order

Approach to Consciousness. *Trends Cogn Sci (Regul Ed)* **23**, 754–768 (2019).

24. Lamme, V. A. F. Visual functions generating conscious seeing. *Front.*

Psychol. **11**, 83 (2020).

25. Lamme, V. A. F. Separate neural definitions of visual consciousness and

visual attention; a case for phenomenal awareness. *Neural Netw.* **17**, 861–872

(2004).

26. Lamme, V. A. & Roelfsema, P. R. The distinct modes of vision offered by

feedforward and recurrent processing. *Trends Neurosci.* **23**, 571–579 (2000).

27. Zaretskaya, N. & Narinyan, M. Introspection, attention or awareness? The

role of the frontal lobe in binocular rivalry. *Front. Hum. Neurosci.* **8**, 527 (2014).

28. Knapen, T., Brascamp, J., Pearson, J., van Ee, R. & Blake, R. The role of

frontal and parietal brain areas in bistable perception. *J. Neurosci.* **31**, 10293–

10301 (2011).

29. Sterzer, P., Kleinschmidt, A. & Rees, G. The neural bases of multistable

perception. *Trends Cogn Sci (Regul Ed)* **13**, 310–318 (2009).

30.   Weilnhammer, V. A., Ludwig, K., Hesselmann, G. & Sterzer, P. Frontoparietal cortex mediates perceptual transitions in bistable perception. *J. Neurosci.* **33**, 16009–16015 (2013).

31.   Panagiotaropoulos, T. I., Deco, G., Kapoor, V. & Logothetis, N. K. Neuronal discharges and gamma oscillations explicitly reflect visual consciousness in the lateral prefrontal cortex. *Neuron* **74**, 924–935 (2012).

32.   Kapoor, V. *et al.* Decoding internally generated transitions of conscious contents in the prefrontal cortex without subjective reports. *Nat. Commun.* **13**, 1535 (2022).

33.   Dwarakanath, A. *et al.* Bistability of prefrontal states gates access to consciousness. *Neuron* **111**, 1666-1683.e4 (2023).

34.   Bellet, J. *et al.* Decoding rapidly presented visual stimuli from prefrontal ensembles without report nor post-perceptual processing. *Neurosci. Conscious.* **2022**, niac005 (2022).

35.   Bellet, M. E. *et al.* Prefrontal neural ensembles encode an internal model of visual sequences and their violations. *BioRxiv* (2021) doi:10.1101/2021.10.04.463064.

36.   Raccah, O., Block, N. & Fox, K. C. R. Does the Prefrontal Cortex Play an Essential Role in Consciousness? Insights from Intracranial Electrical Stimulation of the Human Brain. *J. Neurosci.* **41**, 2076–2087 (2021).

37.   Whyte, C. J., Hohwy, J. & Smith, R. An active inference model of conscious access: How cognitive action selection reconciles the results of report and no-report paradigms. *Current Research in Neurobiology* **3**, 100036 (2022).

38.   Petrides, M. & Pandya, D. N. Distinct parietal and temporal pathways to the homologues of Broca's area in the monkey. *PLoS Biol.* **7**, e1000170 (2009).

39.   Petrides, M. & Pandya, D. N. Projections to the frontal cortex from the

posterior parietal region in the rhesus monkey. *J. Comp. Neurol.* **228**, 105–116

(1984).

40.   Markov, N. T. *et al.* Cortical high-density counterstream architectures.

Science **342**, 1238406 (2013).

41.   Yeterian, E. H., Pandya, D. N., Tomaiuolo, F. & Petrides, M. The cortical

connectivity of the prefrontal cortex in the monkey brain. *Cortex* **48**, 58–81

(2012).

42.   Martín-Signes, M., Paz-Alonso, P. M. & Chica, A. B. Connectivity of

frontoparietal regions reveals executive attention and consciousness interactions.

Cereb. Cortex **29**, 4539–4550 (2019).

43.   Lee, U. *et al.* The directionality and functional organization of

frontoparietal connectivity during consciousness and anesthesia in humans.

Conscious. Cogn. **18**, 1069–1078 (2009).

44.   Tasserie, J. *et al.* Deep brain stimulation of the thalamus restores signatures

of consciousness in a nonhuman primate model. *Sci. Adv.* **8**, eabl5547 (2022).

45.   Casali, A. G. *et al.* A theoretically based index of consciousness independent

of sensory processing and behavior. *Sci. Transl. Med.* **5**, 198ra105 (2013).

46.   Dasilva, M. *et al.* Modulation of cortical slow oscillations and complexity

across anesthesia levels. *Neuroimage* **224**, 117415 (2021).

47.   Alkire, M. T., Hudetz, A. G. & Tononi, G. Consciousness and anesthesia.

Science **322**, 876–880 (2008).

48.   Brown, E. N., Lydic, R. & Schiff, N. D. General anesthesia, sleep, and coma.

N. Engl. J. Med. **363**, 2638–2650 (2010).

49.   MacDonald, A. A., Naci, L., MacDonald, P. A. & Owen, A. M. Anesthesia and

neuroimaging: investigating the neural correlates of unconsciousness. *Trends*

Cogn Sci (Regul Ed) **19**, 100–107 (2015).

50.   Bonhomme, V. *et al.* General anesthesia: A probe to explore consciousness. *Front. Syst. Neurosci.* **13**, 36 (2019).

51.   Purdon, P. L. *et al.* Electroencephalogram signatures of loss and recovery of consciousness from propofol. *Proc Natl Acad Sci USA* **110**, E1142-51 (2013).

52.   Uhrig, L. *et al.* Resting-state Dynamics as a Cortical Signature of Anesthesia in Monkeys. *Anesthesiology* **129**, 942–958 (2018).

53.   Tinker, J. H., Sharbrough, F. W. & Michenfelder, J. D. Anterior shift of the dominant EEG rhytham during anesthesia in the Java monkey: correlation with anesthetic potency. *Anesthesiology* **46**, 252–259 (1977).

54.   Vijayan, S., Ching, S., Purdon, P. L., Brown, E. N. & Kopell, N. J. Thalamocortical mechanisms for the anteriorization of α rhythms during propofol-induced unconsciousness. *J. Neurosci.* **33**, 11070–11075 (2013).

55.   Bastos, A. M. *et al.* Neural effects of propofol-induced unconsciousness and its reversal using thalamic stimulation. *eLife* **10**, (2021).

56.   Vincent, J. L. *et al.* Intrinsic functional architecture in the anaesthetized monkey brain. *Nature* **447**, 83–86 (2007).

57.   Schrouff, J. *et al.* Brain functional integration decreases during propofol-induced loss of consciousness. *Neuroimage* **57**, 198–205 (2011).

58.   Akeju, O. *et al.* Effects of sevoflurane and propofol on frontal electroencephalogram power and coherence. *Anesthesiology* **121**, 990–998 (2014).

59.   Akeju, O. & Brown, E. N. Neural oscillations demonstrate that general anesthesia and sedative states are neurophysiologically distinct from sleep. *Curr. Opin. Neurobiol.* **44**, 178–185 (2017).

60.   John, E. R. & Prichep, L. S. The anesthetic cascade: a theory of how anesthesia suppresses consciousness. *Anesthesiology* **102**, 447–471 (2005).

61.   Franks, N. P. General anaesthesia: from molecular targets to neuronal

pathways of sleep and arousal. *Nat. Rev. Neurosci.* **9**, 370–386 (2008).

62.   Suzuki, M. & Larkum, M. E. General anesthesia decouples cortical

pyramidal neurons. *Cell* **180**, 666-676.e13 (2020).

63.   Luczak, A. & Barthó, P. Consistent sequential activity across diverse forms

of UP states under ketamine anesthesia. *Eur. J. Neurosci.* **36**, 2830–2838 (2012).

64.   Uhrig, L., Dehaene, S. & Jarraya, B. Cerebral mechanisms of general

anesthesia. *Ann Fr Anesth Reanim* **33**, 72–82 (2014).

65.   Lee, U. *et al.* Disruption of frontal-parietal communication by ketamine,

propofol, and sevoflurane. *Anesthesiology* **118**, 1264–1275 (2013).

66.   Hudetz, A. G. & Mashour, G. A. Disconnecting consciousness: is there a

common anesthetic end point? *Anesth. Analg.* **123**, 1228–1240 (2016).

67.   Sellers, K. K., Bennett, D. V., Hutt, A., Williams, J. H. & Fröhlich, F. Awake

vs. anesthetized: layer-specific sensory processing in visual cortex and functional

connectivity between cortical areas. *J. Neurophysiol.* **113**, 3798–3815 (2015).

68.   Sarasso, S. *et al.* Consciousness and Complexity during Unresponsiveness

Induced by Propofol, Xenon, and Ketamine. *Curr. Biol.* **25**, 3099–3105 (2015).

69.   Poulet, J. F. A., Fernandez, L. M. J., Crochet, S. & Petersen, C. C. H. Thalamic

control of cortical states. *Nat. Neurosci.* **15**, 370–372 (2012).

70.   Afrasiabi, M. *et al.* Consciousness depends on integration between parietal

cortex, striatum, and thalamus. *Cell Syst.* **12**, 363-373.e11 (2021).

71.   Deco, G., Jirsa, V. K., Robinson, P. A., Breakspear, M. & Friston, K. The

dynamic brain: from spiking neurons to neural masses and cortical fields. *PLoS

Comput. Biol.* **4**, e1000092 (2008).

72.   Bettinardi, R. G., Tort-Colet, N., Ruiz-Mejias, M., Sanchez-Vives, M. V. &

Deco, G. Gradual emergence of spontaneous correlated brain activity during

fading of general anesthesia in rats: Evidences from fMRI and local field

potentials. *Neuroimage* **114**, 185–198 (2015).

73.   Sarasso, S. *et al.* Quantifying cortical EEG responses to TMS in

(un)consciousness. *Clin. EEG Neurosci.* **45**, 40–49 (2014).

74.   Fetsch, C. R., Kiani, R., Newsome, W. T. & Shadlen, M. N. Effects of cortical

microstimulation on confidence in a perceptual decision. *Neuron* **83**, 797–804

(2014).

75.   Chen, X., Wang, F., Fernandez, E. & Roelfsema, P. R. Shape perception via a

high-channel-count neuroprosthesis in monkey visual cortex. *Science* **370**, 1191–

1196 (2020).

76.   van Vugt, B. *et al.* The threshold for conscious report: Signal loss and

response bias in visual and frontal cortex. *Science* **360**, 537–542 (2018).

77.   Hanks, T. D., Ditterich, J. & Shadlen, M. N. Microstimulation of macaque

area LIP affects decision-making in a motion discrimination task. *Nat. Neurosci.* **9**,

682–689 (2006).

78.   Steriade, M., Nuñez, A. & Amzica, F. A novel slow (< 1 Hz) oscillation of

neocortical neurons in vivo: depolarizing and hyperpolarizing components. *J.*

Neurosci. **13**, 3252–3265 (1993).

79.   Stephen, E. P. *et al.* Broadband slow-wave modulation in posterior and

anterior cortex tracks distinct states of propofol-induced unconsciousness. *Sci.*

Rep. **10**, 13701 (2020).

80.   Kroeger, D. & Amzica, F. Hypersensitivity of the anesthesia-induced

comatose brain. *J. Neurosci.* **27**, 10597–10607 (2007).

81.   Ferron, J.-F., Kroeger, D., Chever, O. & Amzica, F. Cortical inhibition during

burst suppression induced with isoflurane anesthesia. *J. Neurosci.* **29**, 9850–9860

(2009).

82.   Petrides, M., Tomaiuolo, F., Yeterian, E. H. & Pandya, D. N. The prefrontal

cortex: comparative architectonic organization in the human and the macaque

monkey brains. *Cortex* **48**, 46–57 (2012).

83.   Kritzer, M. F. & Goldman-Rakic, P. S. Intrinsic circuit organization of the

major layers and sublayers of the dorsolateral prefrontal cortex in the rhesus

monkey. *J. Comp. Neurol.* **359**, 131–143 (1995).

84.   Petrides, M. Lateral prefrontal cortex: architectonic and functional

organization. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* **360**, 781–795 (2005).

85.   Rigotti, M. *et al.* The importance of mixed selectivity in complex cognitive

tasks. *Nature* **497**, 585–590 (2013).

86.   Mante, V., Sussillo, D., Shenoy, K. V. & Newsome, W. T. Context-dependent

computation by recurrent dynamics in prefrontal cortex. *Nature* **503**, 78–84

(2013).

87.   Miller, E. K. & Cohen, J. D. An integrative theory of prefrontal cortex

function. *Annu. Rev. Neurosci.* **24**, 167–202 (2001).

88.   Miller, E. K., Lundqvist, M. & Bastos, A. M. Working Memory 2.0. *Neuron*

100, 463–475 (2018).

89.   Friedman, N. P. & Robbins, T. W. The role of prefrontal cortex in cognitive

control and executive function. *Neuropsychopharmacology* **47**, 72–89 (2022).

90.   Stuss, D. T. Functions of the frontal lobes: relation to executive functions. *J.

Int. Neuropsychol. Soc.* **17**, 759–765 (2011).

91.   Munakata, Y. *et al.* A unified framework for inhibitory control. *Trends Cogn

Sci (Regul Ed)* **15**, 453–459 (2011).

92.   Jercog, D. *et al.* UP-DOWN cortical dynamics reflect state transitions in a

bistable network. *eLife* **6**, (2017).

93.   Torao-Angosto, M., Manasanch, A., Mattia, M. & Sanchez-Vives, M. V. Up

and Down States During Slow Oscillations in Slow-Wave Sleep and Different

Levels of Anesthesia. *Front. Syst. Neurosci.* **15**, 609645 (2021).

94. Norton, L. *et al.* Disruptions of functional connectivity in the default mode

network of comatose patients. *Neurology* **78**, 175–181 (2012).

95. Bodien, Y. G., Chatelle, C. & Edlow, B. L. Functional networks in disorders

of consciousness. *Semin. Neurol.* **37**, 485–502 (2017).

96. Yang, Y. *et al.* Long-term functional outcomes improved with deep brain

stimulation in patients with disorders of consciousness. *Stroke Vasc. Neurol.* **8**, 368

–378 (2023).

97. Quraishi, I. H., Benjamin, C. F., Spencer, D. D., Blumenfeld, H. & Alkawadri,

R. Impairment of consciousness induced by bilateral electrical stimulation of the

frontal convexity. *Epilepsy Behav. Case Rep.* **8**, 117–122 (2017).

98. West, T. O. *et al.* Stimulating at the right time to recover network states in a

model of the cortico-basal ganglia-thalamic circuit. *PLoS Comput. Biol.* **18**,

e1009887 (2022).

99. Koubeissi, M. Z., Bartolomei, F., Beltagy, A. & Picard, F. Electrical

stimulation of a small brain area reversibly disrupts consciousness. *Epilepsy*

Behav. **37**, 32–35 (2014).

100. Peles, O., Werner-Reiss, U., Bergman, H., Israel, Z. & Vaadia, E. Phase-

Specific Microstimulation Differentially Modulates Beta Oscillations and Affects

Behavior. *Cell Rep.* **30**, 2555-2566.e3 (2020).

101. Salzman, C. D., Britten, K. H. & Newsome, W. T. Cortical microstimulation

influences perceptual judgements of motion direction. *Nature* **346**, 174–177

(1990).

102. Houweling, A. R. & Brecht, M. Behavioural report of single neuron

stimulation in somatosensory cortex. *Nature* **451**, 65–68 (2008).

103.   Logothetis, N. K. *et al.* The effects of electrical microstimulation on cortical

signal propagation. *Nat. Neurosci.* **13**, 1283–1291 (2010).

104.   Tehovnik, E. J., Tolias, A. S., Sultan, F., Slocum, W. M. & Logothetis, N. K.

Direct and indirect activation of cortical neurons by electrical microstimulation. *J.*

Neurophysiol. **96**, 512–521 (2006).

105.   Afraz, S.-R., Kiani, R. & Esteky, H. Microstimulation of inferotemporal

cortex influences face categorization. *Nature* **442**, 692–695 (2006).

106.   Caldwell, D. J., Ojemann, J. G. & Rao, R. P. N. Direct Electrical Stimulation

in Electrocorticographic Brain-Computer Interfaces: Enabling Technologies for

Input to Cortex. *Front. Neurosci.* **13**, 804 (2019).

107.   de Lafuente, V. Regaining the senses of touch and movement. *eLife* **7**,

(2018).

108.   Flesher, S. N. *et al.* Intracortical microstimulation of human somatosensory

cortex. *Sci. Transl. Med.* **8**, 361ra141 (2016).

109.   Najarpour Foroushani, A., Pack, C. C. & Sawan, M. Cortical visual

prostheses: from microstimulation to functional percept. *J. Neural Eng.* **15**, 021005

(2018).

110.   Fox, K. C. R. *et al.* Changes in subjective experience elicited by direct

stimulation of the human orbitofrontal cortex. *Neurology* **91**, e1519–e1527 (2018).

111.   Moore, T. & Armstrong, K. M. Selective gating of visual signals by

microstimulation of frontal cortex. *Nature* **421**, 370–373 (2003).

112.   Torres-Gomez, S. *et al.* Changes in the Proportion of Inhibitory Interneuron

Types from Sensory to Executive Areas of the Primate Neocortex: Implications for

the Origins of Working Memory Representations. *Cereb. Cortex* **30**, 4544–4562

(2020).

113.   Tremblay, R., Lee, S. & Rudy, B. Gabaergic interneurons in the neocortex:

from cellular properties to circuits. *Neuron* **91**, 260–292 (2016).

114. Ferrer, C. & De Marco García, N. V. The role of inhibitory interneurons in
circuit assembly and refinement across sensory cortices. *Front. Neural Circuits* **16**,
866999 (2022).

115. Wu, Y. K., Miehl, C. & Gjorgjieva, J. Regulation of circuit organization and
function through inhibitory synaptic plasticity. *Trends Neurosci.* **45**, 884–898
(2022).

116. Goyal, N. & Chad, D. A. Motor Control, Peripheral. in *Encyclopedia of the*
neurological sciences 100–106 (Elsevier, 2014). doi:10.1016/B978-0-12-385157-
4.00618-7.

117. Herstel, L. J. & Wierenga, C. J. Network control through coordinated
inhibition. *Curr. Opin. Neurobiol.* **67**, 34–41 (2021).

118. Swanson, O. K. & Maffei, A. From hiring to firing: activation of inhibitory
neurons and their recruitment in behavior. *Front. Mol. Neurosci.* **12**, 168 (2019).

119. Santos, B. A., Gomes, R. M. & Husbands, P. The role of rebound spikes in
the maintenance of self-sustained neural spiking activity. *Nonlinear Dyn.* **105**, 767–
784 (2021).

120. Terman, D., Ahn, S., Wang, X. & Just, W. Reducing neuronal networks to
discrete dynamics. *Physica D* **237**, 324–338 (2008).

121. Steriade, M., Amzica, F. & Contreras, D. Cortical and thalamic cellular
correlates of electroencephalographic burst-suppression. *Electroencephalogr.*
Clin. Neurophysiol. **90**, 1–16 (1994).

122. Mashour, G. A. Top-down mechanisms of anesthetic-induced
unconsciousness. *Front. Syst. Neurosci.* **8**, 115 (2014).

123. Hao, X. *et al.* The effects of general anesthetics on synaptic transmission.
Curr. Neuropharmacol. **18**, 936–965 (2020).

124. Antkowiak, B. Different actions of general anesthetics on the firing

patterns of neocortical neurons mediated by the GABA(A) receptor.

Anesthesiology **91**, 500–511 (1999).

125. Ferrarelli, F. *et al.* Breakdown in cortical effective connectivity during

midazolam-induced loss of consciousness. *Proc Natl Acad Sci USA* **107**, 2681–2686

(2010).

126. Arnulfo, G. *et al.* Long-range phase synchronization of high-frequency

oscillations in human cortex. *Nat. Commun.* **11**, 5363 (2020).

127. Poulet, J. F. A. & Crochet, S. The cortical states of wakefulness. *Front. Syst.*

Neurosci. **12**, 64 (2018).

128. Harris, A. Z. & Gordon, J. A. Long-range neural synchrony in behavior.

Annu. Rev. Neurosci. **38**, 171–194 (2015).

129. Greicius, M. D. *et al.* Persistent default-mode network connectivity during

light sedation. *Hum. Brain Mapp.* **29**, 839–847 (2008).

130. *Handbook of brain microcircuits.* (Oxford University Press, 2017).

doi:10.1093/med/9780190636111.001.0001.

131. Rao, S. G., Williams, G. V. & Goldman-Rakic, P. S. Isodirectional tuning of

adjacent interneurons and pyramidal cells during working memory: evidence for

microcolumnar organization in PFC. *J. Neurophysiol.* **81**, 1903–1916 (1999).

132. Constantinidis, C., Williams, G. V. & Goldman-Rakic, P. S. A role for

inhibition in shaping the temporal flow of information in prefrontal cortex. *Nat.*

Neurosci. **5**, 175–180 (2002).

133. Schwartz, M. L., Zheng, D. S. & Goldman-Rakic, P. S. Periodicity of GABA-

containing cells in primate prefrontal cortex. *J. Neurosci.* **8**, 1962–1970 (1988).

134. Lee, A. T. *et al.* Pyramidal neurons in prefrontal cortex receive subtype-

specific forms of excitation and inhibition. *Neuron* **81**, 61–68 (2014).

135. Baker, A. *et al.* Specialized Subpopulations of Deep-Layer Pyramidal

Neurons in the Neocortex: Bridging Cellular Properties to Functional

Consequences. *J. Neurosci.* **38**, 5441–5455 (2018).

136. Murayama, M. & Larkum, M. E. Enhanced dendritic activity in awake rats.

Proc Natl Acad Sci USA **106**, 20482–20486 (2009).

137. Gaillard, R. & Del Cul, A. [A scientific model of consciousness: implications

for neuropsychiatic diseases]. *Psychol. Neuropsychiatr. Vieil.* **5**, 261–267 (2007).

138. Kapoor, V., Besserve, M., Logothetis, N. K. & Panagiotaropoulos, T. I.

Parallel and functionally segregated processing of task phase and conscious

content in the prefrontal cortex. *Commun. Biol.* **1**, 215 (2018).

139. Constantinidis, C. & Qi, X.-L. Representation of spatial and feature

information in the monkey dorsal and ventral prefrontal cortex. *Front. Integr.*

Neurosci. **12**, 31 (2018).

140. Zhao, T. *et al.* Consciousness: new concepts and neural networks. *Front.*

Cell. Neurosci. **13**, 302 (2019).

141. Franke, F. *et al.* Structures of neural correlation and how they favor coding.

Neuron **89**, 409–422 (2016).

142. Stevenson, I. H. *et al.* Functional connectivity and tuning curves in

populations of simultaneously recorded neurons. *PLoS Comput. Biol.* **8**, e1002775

(2012).

143. Safavi, S. *et al.* Nonmonotonic spatial structure of interneuronal

correlations in prefrontal microcircuits. *Proc Natl Acad Sci USA* **115**, E3539–E3548

(2018).

144. Rosenbaum, R., Smith, M. A., Kohn, A., Rubin, J. E. & Doiron, B. The spatial

structure of correlated neuronal variability. *Nat. Neurosci.* **20**, 107–114 (2017).

145. Wang, Y. *et al.* Heterogeneity in the pyramidal network of the medial

prefrontal cortex. *Nat. Neurosci.* **9**, 534–542 (2006).

146.   Sugihara, T., Diltz, M. D., Averbeck, B. B. & Romanski, L. M. Integration of

auditory and visual communication information in the primate ventrolateral

prefrontal cortex. *J. Neurosci.* **26**, 11138–11147 (2006).

147.   O'Reilly, R. C. The What and How of prefrontal cortical organization.

Trends Neurosci. **33**, 355–361 (2010).

148.   Pipa, G. & Munk, M. H. J. Higher Order Spike Synchrony in Prefrontal

Cortex during Visual Memory. *Front. Comput. Neurosci.* **5**, 23 (2011).

149.   Ramani, R. & Wardhan, R. Understanding anesthesia through functional

imaging. *Curr. Opin. Anaesthesiol.* **21**, 530–536 (2008).

150.   Ramani, R., Qiu, M. & Constable, R. T. Sevoflurane 0.25 MAC preferentially

affects higher order association areas: a functional magnetic resonance imaging

study in volunteers. *Anesth. Analg.* **105**, 648–655 (2007).

151.   Vizuete, J. A., Pillay, S., Diba, K., Ropella, K. M. & Hudetz, A. G.

Monosynaptic functional connectivity in cerebral cortex during wakefulness and

under graded levels of anesthesia. *Front. Integr. Neurosci.* **6**, 90 (2012).

152.   Mashour, G. A. Integrating the science of consciousness and anesthesia.

Anesth. Analg. **103**, 975–982 (2006).

153.   Treccani, B. The neuropsychology of feature binding and conscious

perception. *Front. Psychol.* **9**, 2606 (2018).

154.   Treisman, A. Consciousness and perceptual binding. in *The unity of*

consciousnessbinding, integration, and dissociation 95–113 (Oxford University

Press, 2003). doi:10.1093/acprof:oso/9780198508571.003.0005.

155.   Huang, L. Characterizing the nature of visual conscious access: the

distinction between features and locations. *J. Vis.* **10**, 24 (2010).

156.   Pennartz, C. M. A. Neural mechanisms for conscious representations. in

The brain's representational power: on consciousness and the integration of
modalities   233–264   (The   MIT   Press,   2015).
doi:10.7551/mitpress/9780262029315.003.0009.

157.   Parto Dezfouli, M. *et al.* A neural correlate of visual feature binding in
primate lateral prefrontal cortex. *Neuroimage* **229**, 117757 (2021).

158.   Bocincova, A., Buschman, T. J., Stokes, M. G. & Manohar, S. G. Neural
signature of flexible coding in prefrontal cortex. *Proc Natl Acad Sci USA* **119**,
e2200400119 (2022).

159.   Isbister, J. B. *et al.* A new approach to solving the feature-binding problem
in primate vision. *Interface Focus* **8**, 20180021 (2018).

160.   Pirot, S., Glowinski, J. & Thierry, A. M. Excitatory responses evoked in
prefrontal cortex by mediodorsal thalamic nucleus stimulation: influence of
anaesthesia. *Eur. J. Pharmacol.* **285**, 45–54 (1995).

161.   Cruikshank, S. J. *et al.* Thalamic control of layer 1 circuits in prefrontal
cortex. *J. Neurosci.* **32**, 17813–17823 (2012).

162.   Bekinschtein, T. A. *et al.* Neural signature of the conscious processing of
auditory regularities. *Proc Natl Acad Sci USA* **106**, 1672–1677 (2009).

163.   Sombeck, J. T. *et al.* Characterizing the short-latency evoked response to
intracortical microstimulation across a multi-electrode array. *J. Neural Eng.* **19**,
(2022).

164.   Callier, T., Brantly, N. W., Caravelli, A. & Bensmaia, S. J. The frequency of
cortical microstimulation shapes artificial touch. *Proc Natl Acad Sci USA* **117**, 1191
–1200 (2020).

165.   Romo, R., Hernández, A., Zainos, A. & Salinas, E. Somatosensory
discrimination based on cortical microstimulation. *Nature* **392**, 387–390 (1998).

166.   Klink, P. C., Dagnino, B., Gariel-Mathis, M.-A. & Roelfsema, P. R. Distinct

feedforward and feedback effects of microstimulation in visual cortex reveal

neural mechanisms of texture segregation. *Neuron* **95**, 209-220.e3 (2017).

167.   Voigt, M. B., Hubka, P. & Kral, A. Intracortical microstimulation

differentially activates cortical layers based on stimulation depth. *Brain*

Stimulat. **10**, 684–694 (2017).

168.   Maynard, E. M., Nordhausen, C. T. & Normann, R. A. The Utah intracortical

Electrode Array: a recording structure for potential brain-computer interfaces.

Electroencephalogr. Clin. Neurophysiol. **102**, 228–239 (1997).

169.   Quiroga, R. Q., Nadasdy, Z. & Ben-Shaul, Y. Unsupervised spike detection

and sorting with wavelets and superparamagnetic clustering. *Neural Comput.* **16**,

1661–1687 (2004).

1360

a

b

Induction of anaesthesia
using
Ketamine + Dexmedetomidine

Quiet Wakefulness

Light Anaesthesia
(1-1.5% Sevoflurane)

Deep Anaesthesia
(3% Sevoflurane)

50μs

53μs

50μs

50μs

50 pulses per
stimulation train

25μA

25μA

20μA

20μA

15μA

15μA

8μA

8μA

5μA

5μA

1μA

1μA

Experiment flow

c

Pre-stimulation
resting state

Centre-array
stimulation

Full array
stimulation

High amplitude
high Frequency
full array stimulation

Post-stimulation
resting state

a

Quiet wakefulness

8µA — amplitude [µV]
- 0.4mm
- 0.8mm
- 1.2mm
- 1.6mm
- 2mm
- 2.4mm

15µA

1-1,5% (light) Sevoflurane

8µA

15µA

time from stimulation [s]

3% (deep) Sevoflurane

8µA

15µA

b

1, 5, 8, 15, 20, 25 µA pooled

Spatial spread - LFP

signal energy modulation [%]

summed Δ (prox/dist)

distance [mm]

c

Quiet Wakefulness   Light Anaesthesia   Deep Anaesthesia

1, 5, 8, 15, 20, 25 µA pooled

Spatiotemporal modulation of field activity

time from stimulation off [s]

signal energy modulation [%]

distance from stimulated electrode pair [mm]

a

Quiet wakefulness   **1-1,5% (light) Sevoflurane**   **3% (deep) Sevoflurane**

8µA   15µA   8µA   15µA   8µA   15µA

amplitude [µV]

0.4mm
0.8mm
1.2mm
1.6mm
2mm
2.4-2.8mm

time from stimulation [s]

b

1, 5, 8, 15, 20, 25 µA pooled

Spatial spread - LFP

signal energy modulation [%]

distance [mm]

summed Δ (prox/dist)

c   **Quiet Wakefulness**   **Light Anaesthesia**   **Deep Anaesthesia**

1, 5, 8, 15, 20, 25 µA pooled

Spatiotemporal modulation of field activity

time from stimulation off [s]

signal energy modulation [%]

distance from stimulated electrode pair [mm]

SI Figure 1 | Population characteristics under different states

Top row - Col 1 - A comparison of firing rates of the fixed population (based on the firing rates during QW, fr >= 0.5) during different states of (un) consciousness in Monkey A. Col 2 - Metrics of nearness-to-criticality and critical slowing, in 10s bins during different states of (un)consciousness. Col 3 - A comparison of population synchrony of the fixed population of neurons, during different states of (un)consciousness (PFC)

Middle Row - Same as above, for Monkey B (PFC)

Bottom Row - Same as above, for Monkey C (PPC)

SI Figure 2 | Examples of QW, LS and DS

More examples of QW, LS and DS periods, from PFC and PPC

Top rows in each subplot correspond to QW, middle to LS and lower to DS.

Sideplots show the comparison of the collapsed power spectral density (z-scored). The two peaks at 50 and 100Hz are non-physiological, and come from strong line noise during this particular epoch of this particular recording session.

SI Figure 3 | Resting state characteristics for Monkey B - PFC

The raster plot of the fixed population displays sparse and desynchronised spiking during QW (top row), while it displays the characteristic UP/DOWN states during both states of anaesthesia.

The time-frequency decomposition and the the power spectral density both show significantly stronger low-frequency activity during LS and DS, but higher beta and gamma activity during QW.

Although DS was maintained at 3%, there was no significant lengthening of the silent periods as compared to LS (1%). Additionally, there were no significant differences observed between the durations of the UP and DOWN states, pointing to no difference in anaesthesia-induced cortical entrainment.

SI Figure 4 | Spatial and temporal patterns of modulation for Monkey B - PFC

A - Please refer to Fig. 5 in the main text for a description

B - While a conclusive non-monotonic pattern is observed in the modulation in the LFPs in QW, the strength of modulation falls of linearly during both LS and DS in LFPs . Although the sum of rate of change in modulation in all three states do display a weak distal increase, it is not established to be conclusive statistically.

C - Spatio-temporal modulation of field activity.

1   Table for Amplitude: 1 µA

State	Transient Timing [s]	Dispersal	Transient Strength [corrSpks/s]	Dispersal	Rebound Timing [s]	Dispersal	Rebound Strength [corrSpks/s]	Dispersal
QW	0.8438 ± 0.0039	0.6562	93.3845 ± 0.3070	3974.9238	n/a	n/a	n/a	n/a
LS	1.3086 ± 0.0047	1.3967	26.7084 ± 0.0812	419.2467	n/a	n/a	n/a	n/a
DS	0.7064 ± 0.0043	0.5723	206.2554 ± 0.7104	15490.6126	n/a	n/a	n/a	n/a

4   CIs for Amplitude: 1 µA

	QW vs LS	LS vs DS	QW vs DS
Time	CIs:0.0003;0.0162	CIs:0.0003;0.0161	CIs:0.0003;0.0156
Amplitude	CIs:0.0076;0.6921	CIs:0.0151;1.9451	CIs:0.0407;1.7713

6   Table for Amplitude: 5 µA

State	Transient Timing [s]	Dispersal	Transient Strength [corrSpks/s]	Dispersal	Rebound Timing [s]	Dispersal	Rebound Strength [corrSpks/s]	Dispersal
QW	0.7572 ± 0.0026	0.4634	103.1004 ± 0.2183	3266.2309	n/a	n/a	n/a	n/a
LS	1.1009 ± 0.0032	1.0049	39.4883 ± 0.0889	756.4541	n/a	n/a	n/a	n/a
DS	0.5687 ± 0.0024	0.3310	262.2674 ± 0.6476	23749.6579	n/a	n/a	n/a	n/a

8   CIs for Amplitude: 5 µA

	QW vs LS	LS vs DS	QW vs DS
Time	CIs:0.0001;0.0091	CIs:0.0001;0.0113	CIs:0.0001;0.0067

| Amplitude | CIs:0.0193;0.4975 | CIs:0.046;1.5001 | CIs:0.012;1.6153 |

11   Table for Amplitude: 8 µA

State	Transient Timing [s]	Dispersal	Transient Strength [corrSpks/s]	Dispersal	Rebound Timing [s]	Dispersal	Rebound Strength [corrSpks/s]	Dispersal
QW	0.7023 ± 0.0022	0.4075	102.6141 ± 0.2063	3492.3023	n/a	n/a	n/a	n/a
LS	1.1668 ± 0.0028	0.7758	32.5500 ± 0.0682	471.1069	n/a	n/a	n/a	n/a
DS	0.5629 ± 0.0019	0.2651	178.6599 ± 0.3636	10156.9429	n/a	n/a	n/a	n/a

13   CIs for Amplitude: 8 µA

	QW vs LS	LS vs DS	QW vs DS
Time	CIs:0.0002;0.0071	CIs:0.0001;0.0086	CIs:0.0002;0.006
Amplitude	CIs:0.0117;0.4993	CIs:0.0152;0.976	CIs:0.0074;0.8898

15   Table for Amplitude: 15 µA

State	Transient Timing [s]	Dispersal	Transient Strength [corrSpks/s]	Dispersal	Rebound Timing [s]	Dispersal	Rebound Strength [corrSpks/s]	Dispersal
QW	-0.0739 ± 0.0015	0.0234	64.3858 ± 0.2950	967.5470	1.8880 ± 0.0032	0.3152	26.6120 ± 0.1254	473.0090
LS	0.1379 ± 0.0025	0.1121	25.1672 ± 0.1277	281.6930	2.0960 ± 0.0033	0.6020	17.4788 ± 0.0598	192.9809
DS	-0.1092 ± 0.0005	0.0028	57.5013 ± 0.2630	755.9151	1.5646 ± 0.0038	0.4476	41.5794 ± 0.2023	1260.3483

17   CIs for Amplitude: 15 µA

	QW vs LS	LS vs DS	QW vs DS	Rebound QW vs LS	Rebound LS vs DS	Rebound QW vs DS
Time	CIs:0.0001; 0.0077	CIs:0.0002; 0.0076	CIs:0;0.0035	CIs:0.0002; 0.0111	CIs:0.0002; 0.0126	CIs:0.0002; 0.0125

Ampl itude	CIs:0.0126; 0.8382	CIs:0.0097; 0.7864	CIs:0.0107; 0.9128	CIs:0.0044; 0.2845	CIs:0.0068; 0.4393	CIs:0.0073; 0.5404

20   Table for Amplitude: 20 µA

State	Transient Timing [s]	Dispersal	Transient Strength [corrSpks/s]	Dispersal	Rebound Timing [s]	Dispersal	Rebound Strength [corrSpks/s]	Dispersal
QW	0.0784 ± 0.0019	0.0450	43.3549 ± 0.3318	1336.7909	2.1949 ± 0.0031	0.2705	22.3349 ± 0.1030	299.4019
LS	0.1974 ± 0.0017	0.0852	31.8037 ± 0.1786	963.2913	2.4295 ± 0.0030	0.3984	13.3161 ± 0.0519	116.8168
DS	1.3361 ± 0.0050	0.7478	68.4755 ± 0.2659	2098.2474	3.0582 ± 0.0096	0.3002	95.1274 ± 1.4681	7091.3451

22   CIs for Amplitude: 20 µA

	QW vs LS	LS vs DS	QW vs DS	Rebound QW vs LS	Rebound LS vs DS	Rebound QW vs DS
Time	CIs:0.0001; 0.007	CIs:0.0002;0.016	CIs:0.0003; 0.0217	CIs:0.0002; 0.0104	CIs:0.0004; 0.0263	CIs:0.0002; 0.0224
Amplitude	CIs:0.0108; 0.8304	CIs:0.0113;0.7856	CIs:0.0136; 1.0292	CIs:0.004;0.2554	CIs:0.0154; 1.3314	CIs:0.0277; 1.5633

24   Table for Amplitude: 25 µA

State	Transient Timing [s]	Dispersal	Transient Strength [corrSpks/s]	Dispersal	Rebound Timing [s]	Dispersal	Rebound Strength [corrSpks/s]	Dispersal
QW	0.0803 ± 0.0015	0.0323	31.4526 ± 0.2416	799.9227	2.1955 ± 0.0031	0.2691	21.2145 ± 0.1012	285.4046
LS	0.2491 ± 0.0018	0.0959	34.4576 ± 0.1941	1162.2963	2.5425 ± 0.0030	0.3828	15.7265 ± 0.0624	161.8122
DS	0.0693 ± 0.0014	0.0183	49.5116 ± 0.4471	1889.1332	2.1147 ± 0.0024	0.1778	67.1164 ± 0.2642	2212.6116

26   CIs for Amplitude: 25 µA

	QW vs LS	LS vs DS	QW vs DS	Rebound	Rebound	Rebound

				QW vs LS	LS vs DS	QW vs DS
Time	CIs:0.0001; 0.0062	CIs:0.0001; 0.0076	CIs:0.0001; 0.0049	CIs:0.0002; 0.0107	CIs:0.0002; 0.0098	CIs:0.0001; 0.0081
Ampl itude	CIs:0.0107; 0.7699	CIs:0.012;0 .9709	CIs:0.0118; 1.0996	CIs:0.0041; 0.2523	CIs:0.012;0. 6456	CIs:0.0069; 0.8019

1 Table for Amplitude: 1 μA

State	Transient Timing [s]	Dispersal	Transient Strength [corr Spks/s]	Dispersal	1st Rebound Timing [s]	Dispersal	1st Rebound Strength [corr Spks/s]	Dispersal	2nd Rebound Timing [s]	Dispersal	2nd Rebound Strength [corr Spks/s]	Dispersal
QW	0.3209 ± 0.00031	0.5749	28.6237 ± 0.0654	263.2935	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
LS	0.4754 ± 0.00035	0.9590	39.7702 ± 0.1068	909.5022	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
DS	0.2987 ± 0.00029	0.4695	161.8230 ± 0.4905	13495.9911	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a

4 CIs for Amplitude: 1 μA

	QW vs LS	LS vs DS	QW vs DS
Time	CIs:0.0001;0.0109	CIs:0.0002;0.0099	CIs:0.0001;0.0085
Amplitude	CIs:0.0049;0.3375	CIs:0.0107;1.5639	CIs:0.0272;1.6068

6 Table for Amplitude: 5 μA

State	Transient Timing [s]	Dispersal	Transient Strength [corr Spks/s]	Dispersal	1st Rebound Timing [s]	Dispersal	1st Rebound Strength [corr Spks/s]	Dispersal	2nd Rebound Timing [s]	Dispersal	2nd Rebound Strength [corr Spks/s]	Dispersal
QW	0.3463 ± 0.00025	0.4557	28.7935 ± 0.0581	250.4984	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
L	0.44	0.75	56.84	1140	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a

State													
S	70 ± 0.0026	74	67 ± 0.0990	.8940									
DS	0.3390 ± 0.0023	0.4040	133.3902 ± 0.2812	5951.7137	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a

7 CIs for Amplitude: 5 µA

	QW vs LS	LS vs DS	QW vs DS
Time	CIs:0.0002;0.0069	CIs:0.0001;0.0092	CIs:0.0001;0.0074
Amplitude	CIs:0.0085;0.3093	CIs:0.0114;0.6751	CIs:0.0014;0.7452

9 Table for Amplitude: 8 µA

State	Transient Timing [s]	Dispersal	Transient Strength [corr Spks/s]	Dispersal	1st Rebound Timing [s]	Dispersal	1st Rebound Strength [corr Spks/s]	Dispersal	2nd Rebound Timing [s]	Dispersal	2nd Rebound Strength [corr Spks/s]	Dispersal
QW	0.3998 ± 0.0021	0.4138	29.7751 ± 0.0502	229.5956	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
LS	0.4013 ± 0.0021	0.5842	59.5486 ± 0.0943	1133.7954	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a
DS	0.2490 ± 0.0017	0.2754	160.6436 ± 0.2948	7942.8891	n/a	n/a	n/a	n/a	n/a	n/a	n/a	n/a

11 CIs for Amplitude: 8 µA

	QW vs LS	LS vs DS	QW vs DS
Time	CIs:0;0.0072	CIs:0.0002;0.0069	CIs:0;0.0054
Amplitude	CIs:0.0007;0.2662	CIs:0.0121;0.6474	CIs:0.0061;1

13 Table for Amplitude: 15 µA

St	Tra	Dis	Trans	Disp	1st	Dis	1st	Disp	2nd	Dis	2nd	Disp

at e	nsie nt Tim ing [s]	per sal	ient Stren gth [corr Spks/ s]	ersa l	Reb oun d Tim ing [s]	per sal	Rebo und Stren gth [corr Spks/ s]	ersa l	Reb oun d Tim ing [s]	per sal	Rebo und Stren gth [corr Spks/ s]	ersa l
Q W	0.24 97 ± 0.000 07	0.02 34	25.81 01 ± 0.060 8	169. 1815	0.81 55 ± 0.00 09	0.02 54	16.13 93 ± 0.038 9	43.7 187	1.44 64 ± 0.00 14	0.05 29	18.22 87 ± 0.053 2	77.6 375
L S	0.27 03 ± 0.00 08	0.03 58	28.53 94 ± 0.059 8	226. 5878	0.94 06 ± 0.00 08	0.04 72	34.45 57 ± 0.083 2	452. 4760	2.98 94 ± 0.01 01	0.38 89	15.64 93 ± 0.246 1	230. 050 2
D S	0.27 83 ± 0.00 07	0.03 37	69.21 29 ± 0.137 8	1403 .365 9	0.91 16 ± 0.00 09	0.04 82	90.03 69 ± 0.257 4	3853 .478 9	2.94 89 ± 0.01 75	0.35 25	16.84 73 ± 0.663 4	508. 391 3

15   CIs for Amplitude: 15 µA

	QW vs LS	LS vs DS	QW vs DS	1st Reboun d QW vs LS	1st Rebo und LS vs DS	1st Rebo und QW vs DS	2nd Reboun d QW vs LS	2nd Reboun d LS vs DS	2nd Reboun d QW vs DS
Ti m e	CIs:0; 0.002 5	CIs:0; 0.002 3	CIs:0; 0.002 4	CIs:0.00 01;0.003 3	CIs:0; 0.002 8	CIs:0; 0.003 2	CIs:0.00 03;0.023 7	CIs:0.00 04;0.044 2	CIs:0.00 03;0.027 3

16   Table for Amplitude: 20 µA

St at e	Tra nsie nt Tim ing [s]	Dis per sal	Trans ient Stren gth [corr Spks/ s]	Disp ersa l	1st Reb oun d Tim ing [s]	Dis per sal	1st Rebo und Stren gth [corr Spks/ s]	Disp ersa l	2nd Reb oun d Tim ing [s]	Dis per sal	2nd Rebo und Stren gth [corr Spks/ s]	Disp ersa l
Q W	0.14 38 ± 0.000 07	0.01 93	19.11 15 ± 0.049 7	89.0 804	0.59 68 ± 0.00 08	0.02 29	19.46 54 ± 0.048 9	76.8 025	1.19 99 ± 0.00 14	0.08 69	20.00 33 ± 0.045 7	90.6 792
L S	0.37 06 ±	0.05 69	28.73 60 ±	265. 3904	1.10 12 ±	0.05 43	29.95 02 ±	255. 1862	3.22 62 ±	0.32 51	11.16 67 ±	125. 6028

State	Transient Timing [s]	Dispersal	Transient Strength [corr Spks/s]	Dispersal	1st Rebound Timing [s]	Dispersal	1st Rebound Strength [corr Spks/s]	Dispersal	2nd Rebound Timing [s]	Dispersal	2nd Rebound Strength [corr Spks/s]	Dispersal
	0.0009		0.0607		0.0009		0.0621		0.0150		0.2946	
DS	0.1972 ± 0.0006	0.0198	74.4049 ± 0.2012	1966.7066	0.7317 ± 0.0006	0.0215	79.3698 ± 0.1550	1524.3680	1.2736 ± 0.0016	0.1147	81.1950 ± 0.2306	2251.3290

20 CIs for Amplitude: 20 µA

	QW vs LS	LS vs DS	QW vs DS	1st Rebound QW vs LS	1st Rebound LS vs DS	1st Rebound QW vs DS	2nd Rebound QW vs LS	2nd Rebound LS vs DS	2nd Rebound QW vs DS
Time	CIs:0;0.0035	CIs:0;0.0003	CIs:0;0.0024	CIs:0.0001;0.0035	CIs:0.0001;0.0035	CIs:0;0.0025	CIs:0.0004;0.0279	CIs:0.0005;0.0296	CIs:0.0001;0.0047

21 Table for Amplitude: 25 µA

State	Transient Timing [s]	Dispersal	Transient Strength [corr Spks/s]	Dispersal	1st Rebound Timing [s]	Dispersal	1st Rebound Strength [corr Spks/s]	Dispersal	2nd Rebound Timing [s]	Dispersal	2nd Rebound Strength [corr Spks/s]	Dispersal
QW	0.1025 ± 0.00007	0.0300	17.5088 ± 0.0353	74.9733	0.7871 ± 0.0009	0.0444	20.3232 ± 0.0486	137.2494	2.8982 ± 0.0172	0.4077	2.2317 ± 0.0337	1.5632
LS	0.2810 ± 0.0008	0.0340	34.5436 ± 0.0897	481.3793	1.1489 ± 0.0011	0.0729	24.8837 ± 0.0560	174.0582	2.9593 ± 0.0189	0.4159	8.5372 ± 0.2326	63.0340
DS	0.2249 ± 0.0006	0.0212	116.4452 ± 0.2909	5173.3219	0.9078 ± 0.0007	0.0219	76.4889 ± 0.1747	1424.4939	1.3986 ± 0.0011	0.0413	98.3370 ± 0.2917	3078.5002

CIs for Amplitude: 25 µA

	QW vs LS	LS vs DS	QW vs DS	1st Rebound QW vs LS	1st Rebound LS vs DS	1st Rebound QW vs DS	2nd Rebound QW vs LS	2nd Rebound LS vs DS	2nd Rebound QW vs DS
Time	CIs:0;0.0028	CIs:0;0.0022	CIs:0;0.0022	CIs:0.00001;0.00039	CIs:0.00001;0.00036	CIs:0;0.0026	CIs:0.00006;0.0544	CIs:0.00003;0.0253	CIs:0.00003;0.023

Amplitude	CIs:0.004;0.2312	CIs:0.0093;0.8458	CIs:0.0103;0.8835	CIs:0.0033;0.1723	CIs:0.0076;0.4941	CIs:0.0081;0.57	CIs:0.0062;0.5549	CIs:0.0404;3.8546	CIs:0.0541;3.5159

Amplitude	CIs:0.0037;0.2129	CIs:0.0067;0.484	CIs:0.0095;0.6395	CIs:0.0033;0.2272	CIs:0.0059;0.4724	CIs:0.0148;0.6429	CIs:0.0114;0.606	CIs:0.0329;2.7489	CIs:0.0102;0.7123

Am plit ude	CIs:0.0 027;0.1 992	CIs:0.0 053;0. 421	CIs:0.0 074;0.5 301	CIs:0.0 039;0. 295	CIs:0.0 128;0. 6555	CIs:0.0 125;0. 9783	CIs:0.0 04;0.3 774	CIs:0.0 153;1. 3043	CIs:0.0 118;0. 6378

First Peak Table for Amplitude: 1 μA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [μV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	-0.1875 ± 0.0008	0.0025	-146.4080 ± 3.4030	50025.9140	0.0028 ± 0.0001	0.0001
LS	-0.1198 ± 0.0014	0.0087	-292.6658 ± 6.7398	196234.1829	0.0216 ± 0.0010	0.0044
DS	-0.1537 ± 0.0017	0.0068	-111.0850 ± 4.6182	51187.4571	0.0067 ± 0.0006	0.0010

First Peak CIs for Amplitude: 1 μA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0	CIs:0;0.014	CIs:0;0
Amplitudes	CIs:0.0927;6.1429	CIs:0.2752;7.1138	CIs:0.0393;6.115
Widths	CIs:0;0.0001	CIs:0;0.0001	CIs:0;0.0001

5   Second Peak Table for Amplitude: 1 μA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [μV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	-0.0848 ± 0.0019	0.0155	154.6919 ± 7.4938	242600.2345	0.0177 ± 0.0006	0.0016
LS	-0.1421 ± 0.0021	0.0197	591.5271 ± 12.2441	647498.3670	0.0267 ± 0.0006	0.0014
DS	-0.1806 ± 0.0007	0.0013	154.0726 ± 9.4404	213890.9140	0.0197 ± 0.0006	0.0009

Second Peak CIs for Amplitude: 1 μA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0	CIs:0;0	CIs:0;0
Amplitudes	CIs:0.0091;3.9276	CIs:0.0367;2.0518	CIs:0.078;2.7287
Widths	CIs:0;0.0005	CIs:0;0.001	CIs:0;0.0001

First Peak Table for Amplitude: 5 μA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [μV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	-0.1488 ± 0.0019	0.0164	-465.3241 ± 5.2960	121165.5494	0.0170 ± 0.0008	0.0029
LS	-0.0995 ±	0.0102	-475.0833	224492.553	0.0362 ±	0.0055

| | 0.0015 | | ± 7.2087 | 7 | 0.0011 | |
| DS | -0.1418 ± 0.0026 | 0.0161 | -103.1981 ± 3.3370 | 26725.9674 | 0.0072 ± 0.0005 | 0.0005 |

10 First Peak CIs for Amplitude: 5 µA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0.002	CIs:0;0.002	CIs:0;0
Amplitudes	CIs:0.2263;17.2322	CIs:0.0432;15.2422	CIs:0.0237;16.9537
Widths	CIs:0;0.0001	CIs:0;0.0001	CIs:0;0.0001

Second Peak Table for Amplitude: 5 µA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [µV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	0.2394 ± 0.0018	0.0144	357.7109 ± 10.4443	470480.1109	0.0084 ± 0.0007	0.0018
LS	-0.0443 ± 0.0028	0.0335	766.8163 ± 13.6181	800784.5479	0.0147 ± 0.0006	0.0013
DS	-0.1551 ± 0.0018	0.0076	150.5371 ± 10.3800	257184.6153	0.0285 ± 0.0010	0.0025

Second Peak CIs for Amplitude: 5 µA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0.008	CIs:0;0	CIs:0;0.002
Amplitudes	CIs:0.0101;2.4354	CIs:0.0925;7.1158	CIs:0.0367;4.2866
Widths	CIs:0;0.0002	CIs:0;0.0011	CIs:0;0.0003

First Peak Table for Amplitude: 8 µA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [µV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	-0.1386 ± 0.0022	0.0211	-1887.1052 ± 6.9743	210126.3987	0.0691 ± 0.0014	0.0080
LS	-0.1338 ± 0.0016	0.0109	-1066.7797 ± 11.0712	529510.8262	0.0606 ± 0.0011	0.0057
DS	-0.1566 ± 0.0023	0.0131	-466.3530 ± 8.5582	175783.1932	0.0329 ± 0.0013	0.0041

First Peak CIs for Amplitude: 8 µA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0.002	CIs:0;0	CIs:0;0
Amplitudes	CIs:1.0582;30.9494	CIs:0.063;6.9568	CIs:0.445;27.0213
Widths	CIs:0;0.0006	CIs:0;0.0005	CIs:0;0.0001

20   Second Peak Table for Amplitude: 8 μA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [μV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	0.2981 ± 0.0024	0.0256	809.7980 ± 11.1874	540183.9485	0.0118 ± 0.0010	0.0041
LS	0.0434 ± 0.0030	0.0397	798.5892 ± 11.1248	534524.8845	0.0117 ± 0.0007	0.0023
DS	0.1803 ± 0.0019	0.0090	355.2900 ± 9.1696	201794.9599	0.0092 ± 0.0011	0.0029

Second Peak CIs for Amplitude: 8 μA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0.018	CIs:0;0.014	CIs:0;0.01
Amplitudes	CIs:0.2665;6.49	CIs:0.7425;24.6546	CIs:0.1082;7.8635
Widths	CIs:0;0.0002	CIs:0;0.0002	CIs:0;0.0001

First Peak Table for Amplitude: 15 μA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [μV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	0.0756 ± 0.0051	0.0912	4338.7388 ± 17.0864	1026481.6333	0.2256 ± 0.0021	0.0160
LS	0.0662 ± 0.0023	0.0206	3702.9401 ± 19.1846	1477343.5728	0.1518 ± 0.0017	0.0120
DS	0.1732 ± 0.0196	0.2091	3532.5513 ± 66.2855	2381425.0298	0.1056 ± 0.0041	0.0090

First Peak CIs for Amplitude: 15 μA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0.002	CIs:0;0.004	CIs:0;0.002
Amplitudes	CIs:0.5037;25.8877	CIs:1.3405;144.9011	CIs:0.1287;34.5856
Widths	CIs:0.0001;0.0078	CIs:0.0007;0.0138	CIs:0.0003;0.0136

Second Peak Table for Amplitude: 15 μA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [µV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	1.0106 ± 0.0040	0.0686	-1264.0495 ± 6.0337	157271.3427	0.0052 ± 0.0006	0.0016
LS	0.5677 ± 0.0043	0.0796	-853.2645 ± 9.0130	350935.0585	0.0080 ± 0.0008	0.0027
DS	0.6284 ± 0.0024	0.0134	-757.0959 ± 5.2953	67295.8247	0.0041 ± 0.0005	0.0005

30   Second Peak CIs for Amplitude: 15 µA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0.025	CIs:0;0	CIs:0.003;0.03
Amplitudes	CIs:0.5275;32.6493	CIs:0.3925;12.9734	CIs:0.7362;41.1881
Widths	CIs:0;0	CIs:0;0	CIs:0;0

First Peak Table for Amplitude: 20 µA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [µV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	0.0263 ± 0.0034	0.0436	2911.3335 ± 16.6949	1048263.9708	0.0215 ± 0.0011	0.0045
LS	0.0597 ± 0.0053	0.1129	2509.6113 ± 18.1050	1320670.3337	0.0146 ± 0.0007	0.0020
DS	0.0106 ± 0.0008	0.0011	3842.4830 ± 20.5825	811694.7062	0.0241 ± 0.0011	0.0023

First Peak CIs for Amplitude: 20 µA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0	CIs:0;0	CIs:0;0
Amplitudes	CIs:0.9025;82.1748	CIs:1.6208;107.6018	CIs:1.88;92.7815
Widths	CIs:0;0.0002	CIs:0;0.0005	CIs:0;0.0002

Second Peak Table for Amplitude: 20 µA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [µV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	1.1649 ± 0.0051	0.1137	-1737.4405 ± 8.4427	306785.2714	0.0024 ± 0.0001	0.0001
LS	0.6401 ±	0.0793	-921.0162	316656.8692	0.0027 ±	0.0001

	0.0043		± 8.5616		0.0001	
DS	0.8102 ± 0.0043	0.0446	-1426.0558 ± 10.2227	250807.5666	0.0019 ± 0.0001	0.0000

Second Peak CIs for Amplitude: 20 µA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0.038	CIs:0;0.034	CIs:0.001;0.04
Amplitudes	CIs:1.2806;43.1607	CIs:0.6816;34.1415	CIs:0.8829;43.5542
Widths	CIs:0;0	CIs:0;0	CIs:0;0

40   First Peak Table for Amplitude: 25 µA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [µV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	0.0234 ± 0.0008	0.0029	1828.8005 ± 9.0299	342462.3094	0.0132 ± 0.0008	0.0029
LS	0.0106 ± 0.0006	0.0017	2025.3054 ± 11.4609	565729.6258	0.0038 ± 0.0002	0.0002
DS	0.0117 ± 0.0004	0.0004	2352.1052 ± 16.9931	691590.1789	0.0042 ± 0.0008	0.0014

First Peak CIs for Amplitude: 25 µA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0	CIs:0;0	CIs:0;0
Amplitudes	CIs:0.3267;31.0582	CIs:0.1746;39.2103	CIs:0.4628;29.7482
Widths	CIs:0;0	CIs:0;0	CIs:0;0

Second Peak Table for Amplitude: 25 µA

State	Timing Mean [s]	Timing Dispersal	Strength Mean [µV]	Strength Dispersal	Duration Mean [s]	Duration Dispersal
QW	1.0814 ± 0.0056	0.1374	-1590.3188 ± 9.5349	392748.5438	0.0022 ± 0.0001	0.0001
LS	0.7576 ± 0.0033	0.0477	-1135.6794 ± 5.5682	133940.4151	0.0019 ± 0.0002	0.0002
DS	0.8808 ± 0.0048	0.0542	-1629.2031 ± 11.7075	328959.2837	0.0157 ± 0.0025	0.0148

Second Peak CIs for Amplitude: 25 μA

Metric	QW vs LS	LS vs DS	QW vs DS
Times	CIs:0;0.026	CIs:0.001;0.018	CIs:0;0.006
Amplitudes	CIs:0.2473;30.1235	CIs:0.2597;36.3063	CIs:0.9107;28.0431
Widths	CIs:0;0	CIs:0;0	CIs:0;0