1

Genomic and functional insights on *Priestia megaterium* MOD5IV: Enhancing Metal

Phytoremediation Potential in Arid Environments

Luis Pouchucq[1*], Andrés E. Marcoleta[2], Cristian Becerra[1], Carola Bahamondes[1], Pablo Lobos-Ruiz[3]

1. Laboratorio de Biotecnología Vegetal y Ambiental, Departamento de Biotecnología,   Universidad Tecnológica Metropolitana.
2. Grupo de Microbiología Integrativa, Facultad de Ciencias, Universidad de Chile.
3. División de Innovación, Duna Soluciones Biotecnológicas SpA.

* lpouchucq@utem.cl

19   Runnig head: **Metal-Resistant *P. megaterium* for Phytoremediation**

ABSTRACT.

Aims

Metal contamination poses a global threat due to its widespread occurrence and the high toxicity of these elements. Phytoremediation has emerged as a preferred approach for the bioremediation of metal-contaminated soils. The search for microorganisms facilitating phytoremediation, especially plant-growth-promoting bacteria (PGPB), has become critical to advance ecosystem remediation efforts. This research aimed to characterize in-depth a *Priestia megaterium* strain isolated from multimetal contaminated soils located at the Atacama Desert, showing potential for bacteria-assisted phytoremediation .

Methods and Results

The strain MOD5IV exhibited notable PGPB features: phosphate and potassium solubilization, nitrogen fixation, phytohormone production, and growth promotion of *Arabidopsis thaliana*. Genomic analysis revealed a 5,254,635 bp chromosome plus nine plasmids, hosting over 6,000 genes. Functional annotation identified genes associated with resistance to copper, cadmium, lead, mercury, zinc, and cobalt. Also, genes linked to PGPB capabilities as siderophore-production, nutrient-solubilization, IAA-synthesis, and nitrogen-fixation. Accordingly, MOD5IV exhibited robust tolerance to multiple metals and enhanced the phytoremediation potential of *Caesalpinia Spinosa* (Mol.) in laboratory trials.

Conclusions

MOD5IV proved to have promising traits for microbe-assisted phytoremediation of metal-contaminated soils.

Impact Statement

This study contributed to the characterization of new native multi-metal-resistant PGPR bacteria for phytoremediation of metal-contaminated soils. Increasing the evidence of the Atacama Desert as a source of microbiological solutions for climate adaptation and environmental remediation.

KEYWORDS. Plant growth promotion; Arsenic resistance; Copper resistance; Contaminated soils, Indole-acetic acid production.

INTRODUCTION

Soil contamination by heavy metals is a significant concern for governments worldwide. Various regulations establish concentration limits for several heavy metals based on land use (Sánchez-Castro 2023, Parviainen 2022). Additionally, regulations require polluting companies, such as those in the mining industry, to take actions to prevent pollution or to remediate contaminated sites. Governments are making concerted efforts to address soil contamination by heavy metals and mitigate its effects on human health and ecosystems (Liu 2021). There are various strategies for the remediation of these contaminated soils, most of which are based on physical or mechanical methods such as excavation and landfilling, surface capping, chemical solubilization and stabilization, and soil washing, among others. Furthermore, biological strategies are generally based on microorganism-mediated or plant-mediated bioremediation methodologies (Souza 2020, Azhar 2022). The best efficiencies at the field scale have been reported combining physical and biological strategies (Sánchez-Castro 2023).

Among the biological strategies, plant-mediated bioremediation, called phytoremediation, is a widely accepted approach due to its economic feasibility and eco-friendliness (Jacob 2018). However, many factors, such as high metal concentrations, nutrient deficiency, soil structure, and physicochemical factors like pH and salinity, limit biomass production and, therefore, the phytoremediation efficiency of plants (Alves 2022). Improving the phytoremediation efficiency of plants at higher metal concentrations is a promising approach, that may be achieved by utilizing plant growth-promoting bacteria (PGPB) (Manoj 2020; Simmer 2022).

PGPB comprise species from a broad number of genera, including free-living bacteria, those that establish specific symbiotic relationships with plants, bacterial endophytes that colonize plant interior tissues, and cyanobacteria (blue-green algae) (Alves 2022). These bacteria can fix nitrogen, solubilize phosphate and potassium, and produce phytohormones, significantly reducing the need for chemical fertilizers and modulating the effects of environmental stress and pathogen control (Glick 2012, Romero-Estonllo 2023, Asad 2018, Sijilmassi 2020, Khoshru 2020, Backer 2018, Nazari 2020, Myo 2019, Hernández-Soberano 2020, Naqqash 2020). Several bacterial strains isolated from the plant rhizosphere have been reported to improve the productivity and health of plants under metal contamination stress, making them a significant component in phytoremediation processes (Kamaruzzaman 2020, Srivastava 2013, Xu 2018, Wu 2019, Liu 2022).

Among them, *Priestia megaterium* (formerly *Bacillus megaterium*) strains have demonstrated notable

plant growth-promoting (PGP) properties, including potassium solubilization and nitrate reduction (Wu

2023, Wang 2020). These strains also exhibit bioremediation capabilities for environments

contaminated with heavy metals like zinc (Zn) and cadmium (Cd) (Bhatt 2020, Wu 2019) and can

induce plant resistance to environmental stresses or pathogens (Cui 2023, Wang 2020, Chakraborty

2006, Zhou 2016).

In this study, we report on the functional and genomic characterization of the PGP and multi-metal-

resistant strain *P. megaterium* MOD5IV, isolated from the rhizosphere of *Caesalpinia spinosa* (Mol.)

plants thriving in a former mine tailing deposit in the semiarid edge of Atacama Desert. Additionally,

we investigated its capability to promote growth and enhance the phytoremediation performance.

MATERIALS AND METHODS

Strain isolation. Strains were isolated from the rhizosphere of *Caesalpinia spinosa* plants growing on

former tailings from artisanal copper mining near Andacollo, a traditional mining city located in Elqui

Province in northern Chile (30°15.381'S 71°04.551'W). Soil samples (1 g) collected from the mine

tailings were added to 10 mL of sterile water and vortexed vigorously for 1 min. Serial tenfold dilutions

were prepared and then spread on Soy Trypticase Agar (TSA) medium (Difco®). Plates were incubated

at 28°C for 24 h.

Basic qualitative PGPB analysis. Nitrogen fixation was determined by growth on Ashby-agar

medium (0.2 g/L KH_2PO_4, 0.2 g/L $MgSO_4$, 0.2 g/L NaCl, 0.2 g/L $CaSO_4$, 5.0 g/L $CaCO_3$, 5.0 g/L

sucrose, 5.0 g/L glucose, and 15 g/L agar) (Becking 2006). Phosphate and potassium solubilization

index were determined cording Sijilmassi (2020), using Pikovskaya agar medium (10.0 g/L glucose,

0.2 g/L NaCl, 0.002 g/L $FeSO_4$, 0.002 g/L $MnSO_4$, 0.2 g/L KCl, 0.1 g/L $MgSO_4$, 0.5 g/L $(NH_4)_2SO_4$,

0.5 g/L yeast extract, 5.0 g/L $Ca_3(PO4)_2$, and 15 g/L agar) and modified Pikovskaya agar medium (10.0

g/L glucose, 0.2 g/L NaCl, 0.002 g/L $FeSO_4$, 0.002 g/L $MnSO_4$, 0.2 g/L KCl, 0.1 g/L $MgSO_4$, 0.5 g/L

$(NH_4)_2SO_4$, 0.5 g/L yeast extract, 5.0 g/L KNO_3, and 15 g/L agar), respectively. Siderophore production

was tested using the chrome azurol S method in solid medium (CAS-agar), as described by Alexander

and Zuberer (Alexander 1991). The bacterial cultures were incubated at 28°C for 24 h. Orange halos

around the colonies indicated siderophore production.

IAA production was determined colorimetrically, according to Glickmann (1995). The isolates were

inoculated in Soy Trypticase Broth (TSB) medium with and without tryptophan (500 µg/ml) and

incubated at 28°C for 4 days with shaking at 250 rpm. A 1 mL aliquot was removed from each tube and

centrifuged at 10,000 rpm for 10 min. 200 µL of the supernatant from each sample was transferred to a

96-well microplate, to which 50 µL of Salkowski reagent (12 g/L $FeCl_3$ and 7.9 M H_2SO_4) were added.

The mixture was incubated at room temperature for 30 min, and the absorbance was measured at 530

nm using a microplate spectrophotometer (EPOCH-II BioTek). The IAA concentration in the culture

was determined using a calibration curve of pure IAA (Supelco®) as a standard, following linear

regression analysis.

Phosphate uptake quantification. PO_4 was quantitatively determined using the Mo-Blue Method

(Watanabe 1965, Nautiyal 1999). A pre-culture was incubated in 10 mL of TSA medium overnight at

29°C with agitation (150 rpm). The following day, the culture was adjusted to 10^8 CFU/mL, and 100

µL were inoculated into 10 mL of PKV medium. The cultures in PKV medium were incubated at 29°C

with constant agitation (150 rpm) for 72 h. One milliliter of the culture was clarified by centrifugation

(12,000 x g), and 250 µL of the supernatant were placed in a microcentrifuge tube and digested by

adding concentrated sulfuric acid in a 1:1 v/v ratio. 200 µL of the digested samples were placed in

triplicate in a 96-well microplate. 100 µL of Mo-Blue reagent was added to each sample and mixed

gently by pipetting. Absorbance was recorded using a microplate spectrophotometer (EPOCH-II

BioTek) set at 820 nm. A calibration curve was constructed with a PO_4 standard (KH_2PO_4, Certipur®,

Supelco).

Arabidopsis thaliana **growth promotion.** *A. thaliana* (Col-0) seeds were treated with 2% Tween 20

for 10 min with gentle agitation. The seeds were then rinsed thrice with sterile distilled water.

Afterward, the seeds were soaked in 70% ethanol for 30 s and rinsed three times with sterile distilled

water. Subsequently, the seeds were dipped in 1% sodium hypochlorite for 10 min and finally rinsed

with sterile distilled water five times. Seeding was carried out in polycarbonate Petri plates with

Murashige & Skoog (MS) medium supplemented with sucrose (10 g/L) and agar (8 g/L) at pH 5.7.

Three seeds were placed into each plate at 2 cm from each other and incubated at 4°C for 48 h for

stratification. After the stratification period, the plates were placed vertically (at a 65° angle) in an

incubator with a photoperiod of 16:8 h (light:dark) at 24°C for 6 days. On the sixth day of culture, 20

μL of MOD5IV at 10^8 CFU/mL were inoculated 2 cm from the seedling roots. The seedlings were

incubated for an additional 8 days under the same conditions. Finally, the length of the primary root,

number of secondary roots, rosette diameter, leaf area, and number of true leaves were measured. This

assay was carried out in triplicate, with a total of 9 plants for each isolate.

IAA quantification by HPLC. MOD5IV (0.3 mL of 10^8 CFU/mL culture) was inoculated into 30

mL of TSB medium, with and without tryptophan (500 μg/mL), in 50 mL centrifuge tubes, and

incubated at 28°C for 72 h with shaking at 250 rpm. After incubation, the culture was centrifuged at

7,000 rpm for 10 min for cell clarification. 25 mL of the supernatant were adjusted to pH 2.5 with HCl.

IAA was extracted with ethyl acetate in a 1:3 ratio, vortexing for 1 min. The ethyl acetate was

evaporated at 40°C overnight, and the resultant powder was dissolved in 3 mL of methanol. The

sample was analyzed by HPLC (Ultimate 3000 Thermo®) using a pre-equilibrated reverse-phase

column (C18 LiChrosphere 5 μm, Merck®) under isocratic conditions with a mobile phase of

water:acetonitrile: acetic acid (85:15:1). The chromatograms were run at a flow rate of 1 mL/min at

20°C. Detection was performed by UV light absorption at 252 nm. A calibration curve was constructed

using an HPLC-grade standard of IAA (Supelco®) (Shao 2015, Tien 1979).

Quantification of ammonia production. Nitrogen fixation was determined based on ammonia

production using the Nessler's method (Kamaruzzaman 2020, Wu 2019). MOD5IV preinoculum was

prepared in TSB and incubated overnight at 29°C with shaking at 150 rpm. The next day, CFUs were

determined using the microdroplet technique. Inocula were prepared in triplicate tubes with 20 mL of

peptone water (20 g/L peptone and 30 g/L NaCl). The tubes were incubated at 29°C for 5 days with

shaking at 150 rpm. After incubation, CFU was determined using the microdroplet technique. Bacterial

cells were eliminated by centrifugation at 12,000 rpm for 10 min. Then, 200 μL of the culture

supernatant was transferred to new microcentrifuge tubes. Subsequently, 1 mL of Nessler's reagent was

added. The optical density was measured at 450 nm using a sterile 96-well microplate. A calibration

curve was generated using an ammonium standard (NH_4Cl Certipur® Supelco) on the same plate.

Metal exposure and MIC determination. The strain was inoculated from a previous culture adjusted

at 10^8 CFU/mL in a proportion of 1:100 in TSB medium supplemented with previously diluted

gradient of metal concentrations: copper ($CuSO_4$), mercury ($HgCl_2$), chromium ($K_2Cr_2O_7$), lead

($Pb(C_2H_3O_2)_2$), cadmium ($CdCl_2$) and arsenic (As_2O_3) (ReagentPlus®, Sigma-Aldrich). pH was adjusted

to 7.5. The cultures were placed in triplicate into a sterile 96-well microplate and incubated at 28°C
with agitation of 200 rpm into an Epoch-II (BioTeck) microplate reader. Cell proliferation was
determined by absorbance at 600 nm, measuring every 0.5 h for 24 h. The minimum inhibitory
concentration (MIC) was determined for each metal as the concentration needed to avoid culture
growth in the conditions previously mentioned.

Promotion of *C. spinosa* growth in metal-contaminated substrate. *C. spinosa* seeds were treated
with concentrated sulfuric acid for 45 min. The solution was then neutralized with sodium bicarbonate.
The seeds were rinsed with distilled water and placed into plastic containers containing 80 g of
contaminated soil from the same mine tailing where the MOD5IV strain was collected, amended in a
50:50 ratio with sterilized plant substrate based on Sphagnum sp. H2–H4 von Post (Kekkila®
Professional substrate). The MOD5IV inoculum was grown overnight in TSB medium, and the cell
density was adjusted to 10^8 CFU/mL. One milliliter of this inoculum was added to each substrate
container. Control containers received TSB medium without bacteria. One seed was placed in each
container, and the containers were incubated at 28°C for 60 days with a 16:8 light-dark photoperiod.
Five replicates were prepared for each condition.

After the culture period, seedlings were carefully extracted and rinsed with distilled water. Growth
parameters, including root length and dry mass, were recorded. The dried tissues were used to
determine copper (Cu) concentration. The plant material was calcined in a muffle furnace until white
ash was obtained. The white ash was suspended in nitric acid and diluted with distilled water (1:10).
The solutions were clarified by filtering, and Cu concentrations were quantified by atomic absorption
spectroscopy using an AAnalyst™ 700, PerkinElmer instrument. Metal concentrations in the substrate
were quantified by X-ray fluorescence (XRF). A soil (tailing) sample was collected during bacterial
isolation, from a 1 m × 1 m area to a depth of 15 cm using a shovel. The amount of soil was then
reduced using the quartering method (ASTM International 2018) to a representative 1 kg sample. The
tailing material, both alone and amended, was dried in an oven at 60°C until no mass loss was
observed. The samples were analyzed using X-ray fluorescence (XRF) with an S1 Titan 600 Handheld
XRF Analyzer (Bruker).

Genomic sequencing, assembly and annotation. Genomic DNA was purified from an overnight
culture of the MOD5IV strain grown in TSA, using the genomic extraction kit Wizard (Promega®)

following the manufacturer's recommendations. Genomic DNA was sequenced by Illumina® and

Nanopore® platforms. Illumina sequencing (100-pb paired-end reads) was performed through the

TruSeq Nano DNA Kit for library preparation and run in a XLEAP-SBS machine (Illumina SBS

technology). The obtained reads (15,374,518) were quality-checked by the FastQC 0.11.9 software

(FastQC, 2025). The Nanopore sequencing library was prepared using the Ligation Sequencing kit

LSK-110, following the manufacturer's guidelines. 5–50 fmol of the library were loaded into a FLO-

MIN106D flow cell and sequenced using a MinION device (Nanopore Technologies Ltd., Oxford,

UK). The quality and length of the Nanopore reads were assessed by Nanoplot from the Nanopack

toolkit (Oxford Nanopore). Unicycler 0.4.9b (Wick 2017) was used in hybrid mode for genome

assembly. Gene annotation was performed using the Rapid Annotation program by the Subsystem

Technology (RAST) server (Overbeek 2014). Additional functional annotation was carried out using

the KEGG (https://www.kegg.jp/kegg/kegg2.html) database. The Resistance Gene Identifier (RGI) and

the CARD database (https://card.mcmaster.ca/analyze/rgi) (Alcock 2020) were used for resistome

prediction.

RESULTS

Isolation and initial characterization. To screen for the presence of metal-resistant bacteria with

high-efficiency plant growth promotion (PGP) properties, the rhizosphere soil of *Caesalpinia spinosa*

was selected because it was thriving on a former mine tailing deposit with high concentrations of

metals (Supplementary Table 1). The isolation of pure strains was performed on copiotrophic TSA

medium agar plates. A series of bacterial strains were obtained and screened for PGP properties,

including nitrogen fixation, phosphate and potassium solubilization, IAA production, and promotion of

Arabidopsis thaliana growth. The bacterial isolates that most effectively induced the growth of *A.*

thaliana are shown in Figure 1A. Of these, the MOD5IV isolate exhibited the greatest induction of

primary root growth.

The MOD5IV isolate formed white colonies with lobulated margins on TSA medium and showed a

bacillar Gram-positive morphology under microscopic analysis (Figure 1B), with a cell length of 10.5 ±

2.4 µm. To confirm the taxonomic position of MOD5IV, we sequenced the 16S rDNA (Accession

number MN744240.1). BLAST analysis indicated the highest similarity with *Priestia megaterium*

SR01-1 and *Priestia megaterium* DK2 (100% identity for both). The species distribution map of the

16S rDNA gene compared to the 16S rRNA database is shown in Figure 1C, where more than 80% of
the hits obtained corresponded to sequences annotated as belonging to the taxa *Priestia megaterium*
(NCBI: txid1404). This evidence, combined with the comparison results of 16S rDNA gene sequence-
based and whole-genome sequence-based phylogenetic trees elaborated with TYGS (Meier-Kolthoff
2022) (Supplementary Figure 1), allowed us to conclude that MOD5IV belongs to the *Priestia*
megaterium species. This represents the first reported complete genome sequence of the taxa (*P.*
megaterium) for the arid zones of South America (Adeniji 2025).

MOD5IV plant growth promotion properties. The basic plant-growth-promoting properties of
MOD5IV were examined (Figure 1D). MOD5IV was able to solubilize phosphate and potassium in
solid PKV agar medium. The solubilization indexes for both nutrients were 8.6 and 35.3, respectively,
after 72 h of culture. PO_4 solubilization was also quantified using the Mo-Blue method, determining
that MOD5IV was capable of incorporating 69 ± 8 µg(PO_4)/mL directly from inorganic $Ca_3(PO_4)_2$
present in the PKV medium during 72 h of culture. Similarly, MOD5IV was able to grow in nitrogen-
deficient Ashby medium, indicating the presence of a nitrogen fixation mechanism. Quantification of
NH_4 production through the Nessler reaction indicated that MOD5IV produced 100.6 ± 3.6 µgNH4/mL
in a 5-day culture. Siderophore production was evidenced by the formation of a characteristic halo on
CAS-agar medium. The production of the phytohormone IAA was assessed colorimetrically using the
Salkowski method and also by HPLC analysis. In a 72-h culture, the MOD5IV strain produced 40.9
µg/mL of IAA, according to the Salkowski method, and 0.3 µg/mL of IAA using HPLC analysis. This
difference of two orders of magnitude between the methods is due to the Salkowski method reacts
positively with various auxins and their precursors. In contrast, the HPLC method can differentiate
between different types of auxins and precursors as indole-3-lactic acid (ILA), tryptamine, indole-
pyruvic acid, among others (Guardado-Fierros 2024, Tien 1979).

The plant growth-promoting properties (PGPR) of MOD5IV were tested using a co-culture method
with *Arabidopsis thaliana* in solid plant MS medium. Results shown in Figure 2 indicate that the
presence of MOD5IV induced a significant increase in all of the growth parameters tested: length of the
primary root (Figure 2A), number of secondary roots (Figure 2B), and rosette diameter (Figure 2C).
The increase was even more pronounced in the parameters related to the root. Representative images
are shown in Figure 2D.

MOD5IV genomic features. To gain deeper insights regarding the genetic properties of the MOD5IV

strain, we performed complete genome sequencing using Illumina and Nanopore sequencing platforms.

2,321,552,218 bp of raw Illumina data were obtained, corresponding to 15,374,518 100-bp reads. No

quality filtering was required since the FasQC analysis indicated no overrepresented sequences and a

quality Phred score above 30 throughout all the reads. The Nanopore platform generated 575,784,902

bp (56,256 reads, N50=32,731 bp). The complete genome was assembled using a hybrid approach,

taking advantage of the higher-accuracy short reads from the Illumina platform and the significantly

longer Nanopore reads.

The MOD5IV genome showed a total size of 5.69 Mb with an average GC content of 37.73%,

consisting of a 5.25 Mbp circular chromosome and 9 plasmids designated pMOD5IV1 to pMOD5IV9

(Table 1, Figure 3A). Taxonomic classification according to the Genome Taxonomy Database

confirmed this strain belonging to the *Priestia megaterium* species. A schematic representation of the

replicons generated with the CG View server (Wang 2020; Stothard 2005) is provided in Figure 3A.

Genome annotation showed 6,286 protein-coding genes, 75 repeat regions, and 189 RNA genes divided

into 143 tRNAs and 46 rRNA (Table 1). The general features of the *P. megaterium* MOD5IV genome

were compared with three genomes of *P. megaterium* strains with different numbers of plasmids (NTC-

2, DMS319, KF18 and NBRC15305 = ATCC14581) and one *P. flexa* strain genome (KLBMP 4941)

(Table 1). Homologous genome alignment drafts (Figure 3B) (Alikhan 2011) showed that the MOD5IV

chromosome shared the general *Priestia megaterium* genome topology, except for two regions centered

at 3 Mbp and 4.6 Mbp, which apparently correspond to insertions due to differences in the GC content.

These insertions carried some open reading frames related to DNA repair and protein translation, such

as Vsr-endonuclease and Aspartate-tRNA-synthetase. Also, the regions between 1.6 and 1.9 Mbp,

around 3 Mbp, and between 3.6 and 3.8 Mbp showed a marked anomaly in the GC content,

corresponding to divergent regions among the four strains compared. In these regions, it was possible

to find genes related to sporulation, metal resistance, multidrug resistance, transcriptional regulators,

chaperones, and DNA repair. Phage genes were detected only in the region around the coordinate 3.0

Mbp.

MOD5IV genome functional annotation. To obtain better results in genomic annotation, the analysis

was complemented using two different databases, RAST and KEGG (Shrestha 2022). The RAST

platform annotated and functionally categorized 2,068 from a total of 5,722 coding sequences present

in the MOD5IV genome. Of these, 2,025 were found in the chromosome, with the remainder

distributed among the different plasmids. A total of 1,361 chromosomal genes (25%) were assigned to

334 subsystems by RAST. The top five categories in strain MOD5IV were "Amino Acids and

Derivatives" (414), "Carbohydrates" (318), "Cofactors, Vitamins, Prosthetic Groups, Pigments" (159),

"Protein Metabolism" (243), and "Nucleosides and Nucleotides" (108) (Figure 4A). No matches were

found for the "Photosynthesis" and "Phages, Prophages, Transposable Elements, Plasmids" categories.

Meanwhile, the KEGG platform identified 6,022 genes, with 2,915 genes annotated and classified into

KEGG categories. The top five categories were "Protein Families: Genetic Information Processing"

(381), "Carbohydrate Metabolism" (343), "Protein Families: Signaling and Cellular Processes" (324),

"Environmental Information Processing" (227), and "Unclassified: Metabolism" (199) (Figure 4B).

The proteome of MOD5IV was extracted from the RAST server and submitted to the FusionBD server

(https://services.bromberglab.org/fusiondb/) to search for similarities with related strains. The

MOD5IV proteome contained 5449 proteins, which were mapped to 3780 functions. Five hundred

seventy-three proteins could not be mapped to any function in the BD database. The most similar

proteome found in the database was *P. megaterium* DSM319 (TAXID 592022) with 81% similarity

and 3511 shared functions.

Core and accessory genomes were determined using the Spine/AGEnt/ClustAGE tool

(http://spineagent.fsm.northwestern.edu/index_age.html) and 21 *Priestia megaterium* reference strains

(DSM319, YC4-R4, KF18, 207, NCT-2, KNU-01, 5-3, S188, S2, FDU301, BIM B-1314D,

NBRC15305 = ATCC14581, BHS1, IGA-FME-1, CDC 2008724142, H2, 2020WEIHUA_L, BP01R2,

CK7, F20, and KF18) available in BioProject database with an assembly level of complete genome.

Approximately 25.5% of the MOD5IV genome was accessory. The core genome (4,239,675 bp)

consisted of 4,581 coding sequences with a GC content of 38.7%. Meanwhile, the accessory genome

had 1,783 coding sequences and a GC content of 34.9%. The differences in GC content between the

core and accessory genomes could indicate horizontal gene transfer. RAST annotation of the accessory

genome revealed that the principal subsystem categories corresponded to "Amino Acids and

Derivatives," "Stress Response," and "Cell Wall and Capsule." This differed from the core genome,

where the principal categories were "Amino Acids and Derivatives," "Carbohydrates," and "Protein

Metabolism" (Figure 5). The presence of genes classified into the "Virulence, Disease, and Defense"

subsystem also increased considerably in the accessory genome of MOD5IV, including genes related to

copper homeostasis, cobalt-zinc-cadmium resistance, and antibiotic resistance. Meanwhile, in the
"Stress Response" subsystem, the accessory genome of MOD5IV contained genes related to osmotic
stress (choline and betaine uptake), cold shock, and bacterial hemoglobins. A total of 429 coding
sequences were present in the nine plasmids of MOD5IV (Table 2). In the best case, RAST annotation
covered no more than 14% of the sequences. Among the annotated sequences, we found genes related
to isoprenoid biosynthesis, sugar ABC transporters, oxidative stress (catalase), osmotic stress
(glycine/betaine transporters or choline-binding protein A), and phage capsid genes.

Plant growth promotion and resistance genes. As expected, the MOD5IV genome contained several
genes related to both direct and indirect mechanisms of plant growth promotion (Backer 2018).
Regarding direct mechanisms, MOD5IV included ten genes related to PO_4 solubilization and uptake,
such as *phoU, phoP, phoR, phoB,* and *phoH*; two genes related to potassium uptake, *kefA* and the large-
conductance mechanosensitive K channel; 21 genes related to siderophore production and Fe
assimilation; and 12 genes related to nitrogen assimilation, including *norD, norQ,* nitrate reductase, and
glutamine synthase, among others. Concerning indirect plant growth-promoting mechanisms, MOD5IV
housed 10 genes related to auxin production (indole-acetic acid, IAA), primarily belonging to the
indole-3-pyruvic acid and tryptamine synthesis pathways (Supplementary Table 2).

The MOD5IV genome showed a variety of metal resistance genes distributed principally into the
chromosome. These include arsenic resistance genes grouped into two putative operons: one included
arsC, arsB, and *arsR*, while the other contained only *arsR* and *arsB*. Additionally, copper resistance
genes were grouped into three putative operons: one operon encoded a copper-translocating P-type
ATPase, *copD,* and *ycnl*; another contained *copC, copD,* and *cutC*; and the third encoded a copper-
translocating P-type ATPase, *copZ,* and *csoR*. Cadmium resistance genes *cadA* and *cadC* were grouped
into one putative operon. Three mercury-sensitive transcriptional regulator genes (*merR)* not organized
into any *mer* operon were present in the genome. Resistance genes for other metals such as lead, cobalt,
and zinc were also present and organized in an operon containing the *P-type Cd^{2+}/Zn^{2+} transporter,* the
transcriptional repressor *czrA,* and the cobalt/zinc/cadmium resistance protein *czcD* (Supplementary
Table 2).

Using the Resistance Gene Identifier (RGI) tool (Alcock 2023), we further characterized the MOD5IV
resistome, identifying 12 probable antibiotic resistance genes, which include six genes for glycopeptide

antibiotic resistance (*vanY/I/B/W/G/T*), one for tetracycline resistance (*tetQ*), one for cephalosporin resistance *(bcIII)*, one for fosfomycin resistance (*fosB*) and three small multidrug resistance efflux pump (*qacG/J*).

MOD5IV metal resistance and phytoremediation potential. Metal exposure experiments were performed to establish the resistance range of MOD5IV. The strain was exposed to increasing concentrations of common contaminating metals diluted in TSB medium, and growth kinetics were monitored by measuring the absorbance at 600 nm. Figure 6 summarizes the resistance level of the MOD5IV strain, which exhibited high tolerance to multiple metals (and metalloids), including arsenic, chromium, lead, and copper. Conversely, mercury and cadmium induced toxic effects at low concentrations.

MOD5IV was initially isolated from the rhizosphere of a *C. spinosa* specimen. Accordingly, the same plant species was used to test the capabilities of MOD5IV to promote plant growth and improve phytoremediation potential in conditions of high metal concentrations in the substrate. A co-culture experiment was conducted in which *C. spinosa* seedlings were grown for 60 days in tailings material amended with organic substrate and treated with MOD5IV. After the culture period, the seedlings were extracted, and growth parameters were measured. As could be seen, seedlings inoculated with MOD5IV showed a significant increase in growth parameters in comparison with non-inoculated seedlings, including the length of roots and aerial parts (shoots + leaves) (Figure 7A) and dry weight (Figure 7B) parameters. Additionally, an increase in Cu uptake was recorded (Figure 7C), corresponding to an increase in Cu concentration of approximately 63% in the aerial parts and 29% in the roots. Although *C. spinosa* appears to be very tolerant to conditions of high metal concentrations in the substrate, it is not a hyperaccumulating plant, as the Cu concentration in the roots and aerial parts showed an accumulation of only 3% and 0.6%, respectively, of the copper present in the substrate, which registers 1,974 mg(Cu)/kg.

DISCUSSION

Phytoremediation efforts of metal-contaminated soils largely benefit from bacterial strains promoting the growth of plants performing the bioaccumulation process. These strains must have two concomitant features that do not always come together: the capacity to resist the highly toxic metals present, and the metabolic capacities leading to (ideally several) plant growth-promoting effects. As stated above, the

MOD5IV genome contains a complex array of metal resistance genes and operons, conferring resistance to multiple metals, as has been previously reported for other *P. megaterium* strains. For example, the strain HgT21 contains two complete type II *mer* operons in its genome and tolerates about 6.8 mg/L of Hg (Guzmán-Moreno 2022). MOD5IV contains the transcriptional regulator *merR* gene repeated three times in its genome but lacks the rest of the *mer* operon elements. Despite this, MOD5IV could resist 2 mg/L of Hg^{2+} in solution. The HgT21 strain also contains the *ars* operon and tolerates high concentrations of arsenic (5,697.3 mg/L As(V)) in the form of Na_3AsO_4, whereas MOD5IV tolerated over 1,000 mg/L of arsenite (As(III)) (as As_2O_3). This high tolerance correlates with the presence of two *ars* operons in its genome. Higher concentrations of As(III) could not be tested due to its low solubility in the culture medium. MOD5IV also resisted the presence of a high concentration of copper in the culture medium (MIC 125 mg(Cu)/L) at a similar range as the *P. megaterium* HgT21 strain (190 mg(Cu)/L). This resistance could be linked to the presence of three putative copper operons in its genome, including one *csoR-copZA* system that has been described to confer resistance to high levels of copper (Efe 2020, Smaldone 2007).

Regarding cadmium resistance, MOD5IV contained one *cad* operon and other operons conferring resistance to lead, zinc, cadmium, and cobalt, resulting in a resistance of 8 mg/L of Cd in solution, which is slightly greater than the previously reported Cd-resistant *P. megaterium* BM18 strain that has demonstrated resistance of over 4.5 mg/L of Cd (Wu 2019). MOD5IV also showed high resistance to Pb with a MIC of 500 mg/L, considerably more than other *P. megaterium* isolates, which have shown a resistance of 20.7 mg/L of Pb in solution (Roane 1999). However, strains of the genus *Bacillus* have shown resistance to similar amounts of Pb (Abdelkrim 2018). Finally, MOD5IV showed a MIC of 250 mg/L of Cr, despite the absence of specific Cr resistance genes, such as the *chr* operon. Other *P. megaterium* isolates have resisted 150 mg/L of Cr(VI) (Srinath 2002). Notably, MOD5IV presents resistance to a greater number of different metals than other reported *P. megaterium* strains.

Several studies have demonstrated the ability of metal-resistant PGPB strains to increase plant growth and phytoremediation potential in metal-contaminated soils (Bhatt 2020, Chakraborty 2006, Tirry 2021, Tang 2020, Ustiatik 2021, Ullah 2022, Kong 2015, Gao 2010, Ma 2011), including some belonging to *P. megaterium* taxa as the strain HgT21 (Guzmán-Moreno 2022) and M18-2 (Wu 2019). To our knowledge, MOD5IV is the first multi-metal resistant and PGPB *P. megaterium* strain reported with the capacity to increase the phytoremediation potential in experiments using copper mining waste

in the substrate. Plant growth-promoting bacteria can act by reducing the toxic effects of metals or by
producing phytohormones (Wu 2019). Depending on the PGP traits, different bacteria play a key role
in promoting plant development besides reducing the toxicity or damage to plants exposed to stress
from different heavy metals. Additionally, PGPB strains produce various metabolic compounds such as
organic acids, siderophores, EPS, and biosurfactants, which can alter the redox state, chelation,
precipitation, and immobilization of the metals in the rhizospheric environment, thereby enhancing the
plant phytoremediation efficiency (Manoj 2020, Alves 2022).

Biosurfactants, as the lipopeptide surfactin, have shown to play an important role in metal
bioremediation (Mishra 2021) and copper bioleaching (Rozas 2019). Upon searching the MOD5IV
genome, we found no genetic determinants for the surfactin synthesis pathway, despite of its synthesis
has been reported for another *P. megaterium* strain (Cui 2023). Considering the above, it is likely that
copper resistance genes may be crucial in conferring phytoremediation efficiency to *C. spinosa* plants.
Specifically, the *copD* and *copC* genes, which are involved in the import of copper into the cytoplasm
(Lawton 2016), and the cytoplasmic copper-binding protein CutC (Latorre 2011) reduce copper
toxicity and facilitate its safe transport within the cell (Rensing 2003). Additionally, the phosphate
solubilization ability of the tested bacterial strain likely plays an important role in enhancing plant
uptake of soil minerals and increasing the uptake of metals in contaminated soils through the secretion
of organic acids (Manoj 2020, Ma 2011). Bacteria such as MOD5IV and other metal-tolerant
rhizobacteria selected by plants, are crucial for metal detoxification in metalliferous soils. They help
plants endure metal toxicity and improve metal uptake. Thus, the efficiency of phytoextraction can be
enhanced by promoting both plant growth and metal bioavailability in the soil (Ashraf 2019).

Summarizing, The strain *P. megaterium* MOD5IV exhibited outstanding genetic and phenotypic
features, which related plant growth promotion and metal resistance with the increase in the
phytoremediation potential of the desert woody plant *C. spinosa*. These results provide valuable
genomic and experimental resources for further studies and applications of *P. megaterium* MOD5IV in
bioremediation processes, particularly in heavy metal-contaminated soils. Finally, the identification of
the *P. megaterium* MOD5IV strain isolated from metal-contaminated soils in the arid regions of South
America, increases the evidence of the Atacama desert as a source of microbiological solutions for

climate adaptation and environmental remediation (Castro-Severyn 2024, Astorga-Eló 2021, Maza 2019).

ACKNOWLEDGMENTS. We thank Duna Soluciones Biotecnológicas SpA. for providing materials and samples used in this study.

FUNDING. This work was supported by "Funding By "Concurso Interno de Fomento a la I+D+I o Creación" UTEM, LPR21-05, and by the grant FONDECYT 1221193 from Agencia Nacional de Investigación y Desarrollo (ANID, Chile).

REFERENCES

Abdelkrim S, et al. Heavy metal accumulation in *Lathyrus sativus* growing in contaminated soils and identification of symbiotic resistant bacteria. *Arch Microbiol 2019;*201:107–21.

Adeniji A, et al. Unveiling novel features and phylogenomic assessment of indigenous *Priestia megaterium* AB-S79 using comparative genomics. Microbiology Spectrum 2025;13:e01466-24.

Alcock BP, et al. CARD 2023: Expanded Curation, Support for Machine Learning, and Resistome Prediction at the Comprehensive Antibiotic Resistance Database. *Nucleic Acids Research* 2023;*51*:D690-9.

Alcock BP, et al. CARD 2020: Antibiotic resistome surveillance with the comprehensive antibiotic resistance database. *Nucleic Acids Research* 2020;*48*(D1):D517-25.

Alexander DB, Zuberer DA. Use of chrome azurol S reagents to evaluate siderophore production by rhizosphere bacteria. *Biology and Fertility of Soils* 1991;12(1):39–45.

Alikhan NF, et al. BLAST Ring Image Generator (BRIG): simple prokaryote genome comparisons. *BMC Genomics* 2011;12:402.

Alves LS, et al. Plant growth-promoting bacteria in phytoremediation of metal-polluted soils: Current knowledge and future directions. *Science of the Total Environment* 2022*;838*:156435.

Ashraf S, et al. Assessment of phytoremediation potential of native plants growing on Pb mine waste. *Ecotoxicology and Environmental Safety* 2019;174:714-27.

Asad SA, et al. Integrated phytobial heavy metals remediation strategies for sustainable clean environment - A review . *Chemosphere* 2018;*217*:925-941.

ASTM International 2018. Standard Practice for Reducing Samples of Aggregate to Testing Size, C702/C702M- 18. https://store.astm.org/c0702_c0702m-18.html (27 June 2025, date last accessed).

Astorga-Eló M, et al. Rhizobacteria from 'flowering desert' events contribute to the mitigation of water scarcity stress during tomato seedling germination and growth. *Sci Rep*. 2021;11(1): 13745.

Azhar U, et al. Remediation techniques for elimination of heavy metal pollutants from soil: A review. *Environmental Research* 2022;214:113918.

Bhatt K, Maheshwari DK. Zinc solubilizing bacteria (*Bacillus megaterium*) with multifarious plant growth promoting activities alleviates growth in *Capsicum annuum* L. *3 Biotech* 2020;10:36.

14 Backer R, et al. Plant growth-promoting rhizobacteria: Context, mechanisms of action, and
roadmap to commercialization of biostimulants for sustainable agriculture. *Frontiers in Plant*
Science 2018;9;1473.

15 Becking JH. The Family Azotobacteraceae. In: Dworkin M, et al. (ed.) The Prokaryotes. New
York: Springer, 2006, pp 759–783.

16 Castro-Severyn J. et al. Rhizospheric bacteria from the Atacama Desert hyper-arid core:
cultured community dynamics and plant growth promotion. *Microbiol Spectr*
2024;12(6):e0005624.

17 Chakraborty U, Chakraborty B, Basnet M. Heavy metal resistance in microorganisms:
Physiology and molecular mechanisms. *Journal of Basic Microbiology* 2006;*46(3)*:186-95.

18 Cui Z, et al. Isolation and characterization of *Priestia megaterium* KD7 for the biological
control of pear fire blight. *Front Microbiol* 2023;14:1099664.

19 Efe D. Potential Plant Growth-Promoting Bacteria with Heavy Metal Resistance. *Curr*
Microbiol 2020;77: 3861–8.

20 FastQC: A Quality Control Tool for High Throughput Sequence Data. Retrieved from
http://www.bioinformatics.babraham.ac.uk/projects/fastqc (27 June 2025, date last accessed).

21 Gao Y, et al. Improvement of phytoextraction and antioxidative defense in *Solanum nigrum* L.
under cadmium stress by application of cadmium-resistant strain and citric acid. *Journal of*
Hazardous Materials 2010;181:771–7.

22 Glick BR. Plant Growth-Promoting Bacteria: Mechanisms and Applications. *Scientifica*
2012;*2012:963401*.

23 Glickmann E, Dessaux Y. A critical examination of the specificity of the Salkowski reagent for
indolic compounds produced by phytopathogenic bacteria. Appl Environ Microbiol
1995;61(2):793-6.

24 Guardado-Fierros B, et al. Comparative study between Salkowski reagent and chromatographic
method for auxins quantification from bacterial production. *Front. Plant Sci.* 2024;15:1378079.

25 Guzmán-Moreno J, et al. *Bacillus megaterium* HgT21: a Promising Metal Multiresistant Plant
Growth-Promoting Bacteria for Soil Biorestoration. *Microbiol Spectr* 2022;10:e00656-22.

26 Hernández-Soberano C, Ruíz-Herrera LF, Valencia-Cantero E. Endophytic bacteria
Arthrobacter agilis UMCV2 and *Bacillus methylotrophicus* M4-96 stimulate achene

germination, *in vitro* growth, and greenhouse yield of strawberry (*Fragaria × ananassa*)
Scientia Horticulturae 2020;261:109005.

27 Jacob JM, et al. Biological approaches to tackle heavy metal pollution: A survey of literature .
Journal of Environmental Management 2018;217:56-70.

28 Kamaruzzaman MA, et al. Characterisation of Pb-resistant plant growth-promoting
rhizobacteria (PGPR) from Scirpus grossus. *Biocatalysis and Agricultural Biotechnology*
2020*;23*:101456.

29 Khoshru B, et al. Current scenario and future prospects of plant growth-promoting
rhizobacteria: an economic valuable resource for the agriculture revival under stressful
conditions. *Journal of Plant Nutrition* 2020*;43*(20):3062–92.

30 Kong Z, et al. Rhizobial symbiosis effect on the growth, metal uptake, and antioxidant
responses of *Medicago lupulina* under copper stress. *Environ Sci Pollut Res* 2015;22:12479–89.

31 Latorre M, et al. CutC is induced late during copper exposure and can modify intracellular
copper content in *Enterococcus faecalis. Biochemical and Biophysical Research*
Communications 2011;406(4):633-7.

32 Lawton TJ, et al. The CopC Family: Structural and Bioinformatic Insights into a Diverse Group
of Periplasmic Copper Binding Proteins. *Biochemistry* 2016;55(15):2278-90.

33 Liu S, et al. Status and environmental management of soil mercury pollution in China: A
review. *Journal of Environmental Management* 2021;277:111442.

34 Liu A, et al. Improvement of the Cd and Zn phytoremediation efficiency of rice (*Oryza sativa*)
through the inoculation of a metal-resistant PGPR strain. *Chemosphere* 2022*;302*:134900.

35 Ma Y, et al. Plant growth promoting rhizobacteria and endophytes accelerate phytoremediation
of metalliferous soils. *Biotechnology Advances* 2011;29(2):248-58.

36 Manoj SR, et al. Understanding the molecular mechanisms for the enhanced phytoremediation
of heavy metals through plant growth promoting rhizobacteria: A review. *Journal of*
Environmental Management 2020;254:109779.

37 Maza F, et al. Soil Bacterial Communities From the Chilean Andean Highlands: Taxonomic
Composition and Culturability. *Front Bioeng Biotechnol.* 2019;7:10.

38 Meier-Kolthoff JP, et al. TYGS and LPSN: a database tandem for fast and reliable genome-
based classification and nomenclature of prokaryotes. *Nucleic Acids Research*, 2022;50;801-7.

39   Mishra S, et al. Biosurfactant is a powerful tool for the bioremediation of heavy metals from
contaminated soils. *Journal of Hazardous Materials* 2021;418:126253.

40   Myo EM, et al. Indole-3-acetic acid production by *Streptomyces fradiae* NKZ-259 and its
formulation to enhance plant growth. *BMC Microbiol* 2019;19:155.

41   Naqqash T, et al. First report of diazotrophic *Brevundimonas* spp. as growth enhancer and root
colonizer of potato. *Sci Rep* 2020;10:12893.

42   Nautiyal CS. An efficient microbiological growth medium for screening phosphate solubilizing
microorganisms. *FEMS Microbiology Letters* 1999;170(1):265-70.

43   Nazari M, Smith DL. A PGPR-Produced Bacteriocin for Sustainable Agriculture: A Review of
Thuricin 17 Characteristics and Applications. *Frontiers in Plant Science* 2020;11:916.

44   Overbeek R, et al. The SEED and the Rapid Annotation of microbial genomes using
Subsystems Technology (RAST). *Nucleic Acids Research* 2014;42:D206-14.

45   Parviainen A, et al. Human health risks associated with urban soils in mining areas.
Environmental Research 2022;206:112514.

46   Rensing C, Grass G. Escherichia coli mechanisms of copper homeostasis in a changing
environment. *FEMS Microbiology Reviews* 2003;27(2-3):197-213.

47 Roane TM. Lead resistance in two bacterial isolates from heavy metal–contaminated soils.
Microb Ecol 1999;37:218–24.

48   Romero-Estonllo M, et al. Soil amendment and rhizobacterial inoculation improved Cu
phytostabilization, plant growth and microbial activity in a bench-scale experiment. *Front.*
Microbiol. 2023;14:1184070.

49   Rozas EE, et al. Self-assembly of supramolecular structure based on copper-lipopeptides
isolated from e-waste bioleaching liquor. *Journal of Hazardous Materials* 2019;368:63–71.

50   Sánchez-Castro I, et al. Past, present and future trends in the remediation of heavy-metal
contaminated soil - Remediation techniques applied in real soil-contamination events. *Heliyon*
2023;9(6):e16692.

51   Shao J, et al. Analysis and cloning of the synthetic pathway of the phytohormone indole-3-
acetic acid in the plant-beneficial *Bacillus amyloliquefaciens* SQR9. *Microb Cell Fact*
2015;14:130

52   Sijilmassi B, et al. In-vitro plant growth promotion of *Rhizobium* strains isolated from lentil
root nodules under abiotic stresses. *Agronomy* 2020;10(7):1006.

53 Simmer RA, Schnoor JL. Phytoremediation, Bioaugmentation, and the Plant Microbiomes. *Environmental Science & Technology* 2022;56(23):2958-70.

54 Smaldone G, Helmann JD. CsoR regulates the copper efflux operon copZA in *Bacillus subtilis. Microbiology* 2007;153:4123-8.

55 Souza LR, Pomarolli LC, da Veiga M. From classic methodologies to application of nanomaterials for soil remediation: an integrated view of methods for decontamination of toxic metal(oid)s. *Environ Sci Pollut Res* 2020;27:10205–27 .

56 Shrestha P, et al. Prediction of trehalose-metabolic pathway and comparative analysis of KEGG, MetaCyc, and RAST databases based on complete genome of *Variovorax* sp. PAMC28711. *BMC Genomic Data*. 2022;23:4.

57 Srinath T, Verma T, Ramteke PW. Chromium(VI) biosorption and bioaccumulation by chromate resistant bacteria. *Chemosphere* 2002;48:427-35.

58 Srivastava S, et al. Influence of inoculation of arsenic-resistant *Staphylococcus arlettae* on growth and arsenic uptake in *Brassica juncea* (L.) Czern. Var. R-46. *Journal of Hazardous Materials* 2013;262:1039-47.

59 Stothard P, Wishart DS. Circular genome visualization and exploration using CGView. *Bioinformatics* 2005;21(4):537–9.

60 Tang L, et al. *Fava* bean intercropping with *Sedum alfredii* inoculated with endophytes enhances phytoremediation of cadmium and lead co-contaminated field. *Environmental Pollution* 2020;265:114861.

61 Tien TM, Gaskins MH, Hubbell DH. Plant growth substances produced by *Azospirillum brasilense* and their effect on the growth of Pearl millet (*Pennisetum americanum* L.). Applied and Environmental Microbiology 1979;37(5):1016-24.

62 Tirry N, et al. Improved chromium tolerance of *Medicago sativa* by plant growth-promoting rhizobacteria (PGPR). *Journal of Genetic Engineering and Biotechnology* 2021;19:149-62.

63 Ullah A, et al. Phytoremediation of heavy metals assisted by plant growth promoting (PGP) bacteria: A review. *Environmental and Experimental Botany* 2022;117:28-40.

64 Ustiatik R, et al. Mercury resistance and plant growth promoting traits of endophytic bacteria isolated from mercury-contaminated soil. *Bioremediation Journal* 2021;26(3):208–27.

65   Wang B, et al. Genomic Analysis of *Bacillus megaterium* NCT-2 Reveals Its Genetic Basis for
the Bioremediation of Secondary Salinization Soil, *International Journal of Genomics*,
2020;4109186.

66   Watanabe FS, Olsen SR. Test of an ascorbic acid method for determining phosphorus in water
and NaHCO3 extracts from soils. *Soil Science Society of America Journal, 1965;29*:677-67.

67   Wick RR, et al. Unicycler: Resolving bacterial genome assemblies from short and long
sequencing reads. *PLOS Computational Biology* 2017;13(6):e1005595.

68   Wu J, et al. Endophytic *Bacillus megaterium* BM18-2 mutated for cadmium accumulation and
improving plant growth in Hybrid *Pennisetum*. *Biotechnology Reports 2019;23*:e00374.

69   Wu X, et al. Analysis of the Potassium-Solubilizing *Priestia megaterium* Strain NK851 and Its
Potassium Feldspar-Binding Proteins. *Int J Mol Sci.* 2023;24(18):14226.

70   Xu X, et al. Complete Genome Sequence of Cd(II)-Resistant *Arthrobacter* sp. PGP41, a Plant
Growth-Promoting Bacterium with Potential in Microbe-Assisted Phytoremediation. *Curr*
Microbiol 2018;75:1231–9.

71   Zhou C, et al. Rhizobacterial Strain *Bacillus megaterium* BOFC15 Induces Cellular Polyamine
Changes that Improve Plant Growth and Drought Resistance. *Int J Mol Sci.* 2016;17(6):976.

Table 1

Table1. General genome features of *P. megaterium* MOD5IV compared with other five *Priestia*
strains.

Strains	P. megaterium MOD5IV	P. megaterium NCT-2	P. megaterium DSM 319	P. megaterium KF18	P. megaterium NBRC 15308	P. flexa KLBMP 4941
Genome size (Mb)	5.7	5.9	5.1	6.0	5.7	4.1
Chromosome size (Mb)	5.3	5.2	5.1	5.2	5.3	3.96
G+C content (%)	37.7	37.8	38.1	37.6	38.0	38.0
Chromosomal G+C content (%)	38.0	38.2	38.1	38.1	38.0	38.1
Gene number	6,211	6,134	5,337	6,294	6,003	4,332
Coding sequence number	6,022	5,805	4,994	5,979	5,754	3,971
RNA gene number	189	203	153	217	163	143
rRNA genes	46	53	33	46	41	36
tRNA gene number	243	142	114	163	122	102
Plasmid number	9	10	0	9	6	2

647
648

Table 2

663 **Table 2. General features of the chromosome and plasmids present in *P. megaterium* MOD5IV**

664 **genom**

	Chromosome	pMOD 5IV 1	pMOD 5IV 2	pMOD 5IV 3	pMOD 5IV 4	pMOD 5IV 5	pMOD 5IV 6	pMOD 5IV 7	pMOD 5IV 8	pMOD 5IV 9
Size (pb)	5,254,635	125,337	112,536	69,915	47,78	42,938	10,013	9,405	8,643	8,518
GC Content	38.0	34.1	34.5	31.7	32.0	37.3	34.8	34.4	33.8	34.5
Number of RAST Subsystems	334	9	6	6	0	0	0	0	0	1
Number of Coding Sequences	5.554	140	62	84	53	42	14	10	13	11
Number of RNAs	168	0	0	0	0	20	0	0	0	0

Figure 1. Selection of Isolates, Plant Growth Promotion Capacity, and General Characterization of Strain MOD5IV. A) Arabidopsis thaliana primary root growth promotion by the isolated strains. The red column represents the strain MOD5IV, which had the highest plant growth promotion capacity. a, b and c indicate significant differences estimated using Two-way ANOVA ($p < 0.05$). Parametricity was estimated using the Kolmogorov-Smirnov Test of Normality. N = 9. B) MOD5IV morphology as revealed by bright-field microscopy and Gram staining (scale bar indicates 25 µm). C) Species distribution map of MOD5IV 16S rDNA compared to the NR database (numbers represent the percentage of representation of the 100 best hits in BLAST analysis). D) Summary of the morphological characteristics and plant growth promotion properties of strain MOD5IV.

699 <div align="center">**Figure 2**</div>

Figure 2. *P. megaterium* MOD5IV Promotes Arabidopsis thaliana Growth In Vitro. A) Promotion

of *A. thaliana* primary root growth by the MOD5IV strain. B) Promotion of *A. thaliana* lateral root

development induced by the MOD5IV strain. C) Growth of *A. thaliana* rosette induced by MOD5IV. a

and b indicate significant differences estimated by Two-way ANOVA p < 0.05. Parametricity was

estimated by Kolmogorov-Smirnov Test of Normality. N = 9. D) Representative images of *A. thaliana*

plants exposed to MOD5IV and not exposed (Control) in a 15-day co-culture experiment.

Figure 3

CDS
tRNA
misc_feature
GC Content
GC Skew+
GC Skew-

P. megaterium DSM319
P. megaterium K18
P. megaterium NBRC15308
P. megaterium NTC2
GC Content
GC Skew+
GC Skew-

Figure 3. Genome map of *P. megaterium* MOD5IV and comparison with other P. megaterium genomes. A) Chromosome map prepared using Proksee software. Circles from outside to inside show the position of protein-coding sequences (blue), tRNA genes (pink), and rRNA genes (cyan) on the positive (circle 1) and negative (circle 2) strands. Circles 3 and 4 show plots of GC content and GC skew as deviations from the average for the entire sequence. B) BRIG output image. The draft of the P. megaterium MOD5IV genome aligned against other P. megaterium genomes (DM319, K18, NBRC15305 = ATCC 14581, NTC2). GC content and skew are also shown.

Figure 4

Figure 4. Predicted functions encoded in the *P. megaterium* MOD5IV genome. A) Statistical chart of the RAST subsystem category distribution. The y-axis represents the classification of different functional genes in the RAST subsystem classification. B) Statistical chart of the KEGG annotation classes. The y-axis represents the classification of different functional genes in the KEGG database.

Figure 5

Figure 5. ***P. megaterium*** **MOD5IV Accessory Genome Analysis.** RAST subsystem classification of the MOD5IV core and accessory genome determined by Spine/AGEnt. Data is presented as a percentage of the total genes in the core and accessory genome.

Figure 6

Figure 6. *P. megaterium* **MOD5IV Metal Resistance.** Maximum growth of MOD5IV in TSB medium supplemented with increasing concentrations of metals (as indicated in each figure), measured by absorbance at 600 nm. Error bars indicate the standard deviation of the average of three independent experiments. MIC: As = >1000 mg/L; Hg = 2 mg/L; Pb = >500 mg/L; Cr = 500 mg/L; Cd = 8 mg/L; Cu = 120 mg/L.

805   **Figure 7**

Figure 7. *P. megaterium* MOD5IV Increases Growth and Copper Uptake in Caesalpinia spinosa.
Two-month-old C. spinosa plants were grown in mine tailings amended with organic substrate, with or
without inoculation with P. megaterium MOD5IV. A) aerial (shoot + leafs) and root length of C.
spinosa. B) Dry weight of C. spinosa aerial organs and roots. C) Copper accumulation in aerial organs
and roots of C. spinosa. a and b indicate significant differences estimated using Two-way ANOVA (p <
0.05). Parametricity was estimated by the Kolmogorov-Smirnov Test of Normality.