Bulk-based hypothesis weighing to increase power in single-cell differential expression analysis

Pierre-Luc Germain[1,2,3] and Mark D. Robinson[2,3]

Abstract

Due to the costs of single-cell sequencing, sample sizes are often relatively limited, sometimes leading to poorly reproducible results. In many contexts, however, larger bulk RNAseq data is available for the same conditions or experimental paradigm, which can be used as additional evidence of a generalizable differential expression pattern. Here, we show how such data can be used, via bulk-based hypothesis weighing (bbhw), to increase the power and robustness of single-cell differential state analysis. We find that all methods improve performance, with the best results obtained by applying a grouped Benjamini-Hochberg procedure on bins based on proportion-adjusted significance (PAS). These methods are implemented in the *muscat* package, and should be applicable to a broader range of scenarios.

Keywords: Single-cell, differential analysis, multiple testing correction, false discovery rate, hypothesis weighing, power

[1]D-HEST Institute for Neuroscience, ETH Zurich, Switzerland,   [2]Department of Molecular Life Sciences, University of Zurich, Zurich, Switzerland,   [3]SIB Swiss Institute of Bioinformatics, Switzerland

Correspondence
pierre-luc.germain@hest.ethz.ch

Introduction

The development of single-cell technologies has enabled the analysis of complex tissues at the resolution of individual cell types. However, due to the costs of these technologies, sample sizes are often relatively limited, sometimes leading to poorly reproducible results, especially with clinical data (e.g. Maleki et al., 2019; Nakatsuka et al., 2025). In many contexts, however, larger bulk RNAseq datasets are often available for the same conditions or experimental paradigm, which could be used as additional evidence of a differential expression pattern. Of course, many changes observed in specific cell types – especially rare ones – might not be visible in bulk data, and such a prior should not reduce our ability to detect novel changes. Here, we explore some simple methods to use bulk RNAseq data to increase the power and robustness of (multi-sample, multi-condition) single-cell differential expression analysis.

Consider as an example the recent data from Waag et al., 2025 on the response to stress in the hippocampus, where a large bulk RNAseq data (195 samples) is complemented by a smaller single-cell RNA-seq dataset. For each cell type, pseudobulk aggregation was performed to test for differential expression upon stress (Crowell et al., 2020), and the distribution of p-values (here for all neuronal cell types) is shown in Figure 1A. The p-value distribution shows a small excess of low p-values in comparison to the flat distribution that would be expected in the absence of genuine differences. While this is suggestive of a biological effect, the excess is very small, and consequently only a small fraction of genes can be reported as significant after False Discovery Rate (FDR) correction. However, we can partition the genes into quantile bins based on their significance in the much larger bulk dataset using the bulk RNA-seq data (Figure 1B). If we then look again at the distribution of single-cell p-values, but splitting genes according to this prior evidence (i.e. according to the significance bin in the bulk dataset), we can appreciate that some bins (e.g. bin 1) show a much larger excess of small p-values, while others (e.g. bin 3) are rather depleted of small p-values (Figure 1C). As a consequence of this greater (local) excess, the probability that any of those p-values is due to chance becomes lower, and thus a notable gain in power could be obtained by applying FDR to each bin separately.

Figure 1 – Example p-values of differential expression upon stress, from both a single-cell dataset with modest sample size (**A** and **C**) and a much larger bulk dataset (**B**) Waag et al., 2025. **C**: Splitting the single-cell p-values by significance of the respective gene in the bulk data leads to higher local excess of p-values.

Several methods have been proposed to harness such grouping of the hypotheses during FDR correction (Korthauer et al., 2019), in particular the Grouped Benjamini-Hochberg (gBH)

procedures (Hu et al., 2010) and Independent Hypothesis Weighing (IHW) (Ignatiadis et al., 2016). gBH procedures work by estimating the proportion of true null hypotheses in each bin, which can be done through either the Least-Slope (LSL, Benjamini and Hochberg, 2000) or Two-Stage (TST, Benjamini et al., 2006) methods, and weighing the p-values accordingly (Hu et al., 2010). IHW instead uses a cross-validation strategy to optimize weights that lead to the highest rejections of the null hypothesis (see Ignatiadis et al., 2016).

In this brief report, we investigate the application of such adjusted hypothesis testing regimes, and additionally show that taking into consideration the local informativeness of the prior, i.e. the extent to which the cell type contributes to the bulk expression for each gene, further improves both precision and recall.

Methods

Example stress data

The example stress data used in Figures 1 and 3A was taken from the neuronal populations of (Waag et al., 2025). Specifically, we used the differential expression analysis results of dropping all acute stress related covariates from the model, i.e. Sex+CRS*TimePoint vs Sex+CRS (for the bulk, without Sex for the single-cell data that was only on females). For Figure 3A, we used a single cell type: ExN.CA1-do.

Toy p-value simulations

To explore the behavior of adjustment methods, we generated distributions of p-values (Figure 3B-D). For null simulations, we simulated 80k p-values using a uniform distribution. For simulations with a signal, we generated 74k uniform p-values (true null hypotheses), and complemented them with the absolute of normally-distributed values centered at 0, using 1000 with a standard deviation of 0.001, 3000 with 0.01, and 2000 with 0.025. Uninformative bins were generated by randomly sampling the desired number of bins with equal probability. Every scenario was simulated with 10 random seeds.

To generate informative bins, we first split the true null hypotheses (i.e. p-values generated with a uniform distribution) into 5 equally-size bins, and grouped the other p-values (i.e. with false null hypothesis) according to the p-value distribution from which they were sampled. We created noise by randomly assigning one third of the true null hypotheses to any of the bins with signal.

scRNA-seq simulation

To evaluate different approaches, we first simulated single-cell RNAseq data using muscat, using as a reference the dataset published with the package (Crowell et al., 2020). Specifically, we simulated two different groups (with 8 samples each) and six cell types (Figure 2): three abundant cell types, only one of which differs between groups, and three rare, of which two differ between groups. The two affected rare cell types differ chiefly in that smallAffected1 has an expression profile similar to one of the large unaffected clusters, while smallAffected2 is more distinct but has smaller differential expression between groups (10% of the genes with an average log2FC of 0.3, versus 20% of the genes with an average log2FC of 0.4 for other populations).

For the evaluation, the 16 samples were summed to generate artificial bulk data, on which a differential expression analysis was performed using edgeR (Robinson et al., 2010), and 12

Figure 2 – Simulated dataset with six cell types across two groups of 8 samples each.

random sets of 2 vs 2 samples were used for pseudo-bulk differential state analysis using muscat-edgeR (Crowell et al., 2020).

Downsampled real data

To have a complementary evaluation based on real data, we used the large single-nuclei dataset of human multiple sclerosis (MS) from Macnair et al., 2025, specifically restricting ourselves to white matter samples. We performed a pseudo-bulk differential analysis between the 13 control samples and the 51 diseased samples. As ground truth, we took the hypotheses with a (locally) adjusted p-value below 0.1 as true positives, and those above 0.5 as true negatives – the hypotheses in-between were judged unclear and discarded for the purpose of evaluation. Only cell types with more than 2 true positives were considered for the evaluation. We then tested the different methods on 12 random subsets of 4 control vs 5 diseased samples.

The MS data has strong condition-associated changes in cell type abundance, which would dominate bulk data. Therefore, to generate artificial bulk data, we first equalized the cell type abundance across samples, before summing the counts.

Tested methods

We tested different ways of grouping hypotheses based on the bulk data, as well as different ways of using this grouping for p-value adjustment, all of which are now implemented in the `bbhw` function of the *muscat* package.

Creation of the evidence bins. We tested the following ways of grouping hypotheses based on the bulk data:

p: bins are based on quantiles of the bulk p-values. Missing (i.e. NA) p-values are put in their own bin if there are at least 50 of them, and otherwise set to 0.5.

p+sign: for each hypothesis (i.e. gene in a given cell type), if the direction of the change is different between bulk and the cell type, the bulk covariate is set to 0.7. Bins are then based on this modified covariate as for 'p'.

combined: : first, significance-based bins are created in the same fashion as in 'p+sign'. Each significance bin is then further split into genes for which the cell type contributes much to the bulk, and genes for which the cell type contributes little.

PAS (Proportion-Adjusted Significance): : for each hypothesis, the bulk significance is first adjusted for discordant directions as described for previous variants. Then the covariate is adjusted based on the cell type contribution to the bulk of that gene using $inv.logit(logit(p)*$

sqrt(*c*)), where *p* and *c* are respectively the bulk p-value and the proportion of bulk reads
contributed by the cell type). Bins are then based on the quantiles of this modified co-
variate.

PA-logFC (Proportion-Adjusted logFC): : same as for *PAS*, except that shrunk bulk logFCs
(from the `edgeR::prefFC` function) are used instead of the significance. The logFCs are
converted to signed rank and normalized between -1 and 1, so that the same procedure
as PAS can be applied.

asNA: : for hypotheses for which the cell type contributes little to the bulk profile, the
covariate (i.e. bulk p-value) is set to NA, resulting in the hypothesis being placed in a
separate bin. Non-NA values are then split into quantile bins.

In all cases, the number of bins is determined in relation to the number of hypotheses, ensur-
ing a minimum of 1000 hypotheses per bin (except for the NA bin), and using a maximum of 12
bins except where indicated otherwise.

p-value adjustment. Once the bins are created, the following options were tested to make use
of them:

binwise: : the Benjamini-Hochberg (BH) procedure is applied separately for each bin. Doing
this can lead to an increase in false positives if the number of bins is large, and to correct
for this the resulting adjusted p-values are multiplied by $\min(1, \text{nbins}/\text{rank}(p))$, which
results in proper FDR control even across a large number of bins.

IHW: : The Independent Hypothesis Weighing (IHW) method by Ignatiadis et al., 2016 is
applied, using the *IHW* R package with the default alpha of 0.1. If the 25% quantile of the
bin sizes was above 1000, 5-fold validation was performed. If it was above 400, 4-fold,
and otherwise 3-fold.

gBH.LSL: : Grouped Benjamini-Hochberg (gBH) is used (Hu et al., 2010) with the Least-
Slope (LSL) estimation of the proportion of true null hypotheses (Benjamini and Hochberg,
2000).

gBH.TST: : Grouped Benjamini-Hochberg (gBH) is used (Hu et al., 2010) with the Two-Stage
(TST) estimation of the proportion of true null hypotheses. The Two-Stage approach
amounts to using BH within each group, and use the rejection rate at a given nominal
FDR threshold (here we used 0.05) to estimate the proportion of true null hypotheses
within the group.

Each method was run in two flavors: a local one, which is applied for each cell type separately,
and a global one, which is applied once across all cell types.

Results

General behavior of the adjustment methods

We first tested the different methods that leverage hypothesis grouping using the Waag et al.
(2025) stress single-cell data. While there is no ground truth in this data, we tested the expecta-
tion that using uninformative (i.e. random) bins should not lead to systematically higher reported
significance. As shown in Figure 3A, computing adjusted p-values (using the FDR method) sep-
arately for the different bins (FDRsplit) leads to a systematic decrease in the reported adjusted
p-values, although the effect was very mild unless the number of bins was very high. This could
be easily corrected by multiplying the adjusted p-values by $\max(1, n/\text{rank}(padj))$, where *n* is the

Figure 3 – **A:** Adjusted p-values obtained by the different methods on the stress dataset with increasing numbers of uninformative (i.e. random) bins. **B:** Example toy simulations of p-value distributions with and without signal. **C:** Precision and recall (at nominal p.adj<0.05) of the adjustment methods in simulation with signals, using 8 informative bins. The dots indicate the average, and the segment the range, across 10 random seeds. (Note that the FDRsplit and FDRsplit-adjusted methods overlap here). **D:** Number of false positives reported by the methods in the null simulations with increasing number of uninformative (i.e. random) bins.

number of bins and rank($padj$) is the rank of the (pooled) adjusted p-values. The results are
shown as 'FDRsplit-adjusted'. The Two-Stage grouped BH method (TST) had a similar trend, but
much milder, while the other methods did not lead to a systematic FDR underestimation.

All methods leveraging bulk-based evidence bins improve power

We next turned to our specific case of using bulk data to improve power of per-cell-type
differential expression across condition in single-cell data. To this end, we used simulated data, as
well as downsampled real data with the full-sized data serving as ground truth (see Methods). All
methods led to an improvement in the precision-recall curves, mostly through an improvement in
recall, except for the adjusted bin-wise method that instead led to increased precision (Figure 4A-
B). The two grouped BH variants (LSL and TST) showed the strongest improvement.

Further improvements by considering direction and cell type contribution to the bulk

The prior evidence, i.e. the bulk data, is not equally informative for all genes and cell types.
It is more informative for abundant cell types than rare cell types, because these influence the
bulk more, and it is also more informative for a gene that is chiefly expressed in the cell type
of interest, versus a gene that is mostly expressed in other cell types (which would dilute, at

Figure 4 – **A:** TPR-FDR curves of the different adjustment methods, using bins based on quantiles of raw bulk significance. The dashed lines indicate an effective FDR of 0.05, 0.1 and 0.25. **B:** Precision, recall and F1 score of the same methods at a nominal adjusted p-value threshold of 0.05. **C:** Precision, recall and F1 score of the top method (gBH-LSL) using different ways to group the hypotheses (i.e. variants of the bin preparation). 'p' stands for defining bins based on the bulk p-value only. 'p+sign' stands for discounting opposite directions. The other three methods additionally take the proportion of contribution into account (see Methods).

the bulk level, the effects in the cell type of interest). We therefore computed, based on the pseudobulk profiles, the proportion of the bulk reads expected to be attributable to each cell type for each gene, and tested ways to include that in the creation of the evidence bins (see Methods). In addition, we discounted (i.e. increased) the bulk p-values when the bulk logFC was in the opposite direction to that of the cell type of interest. Taking the direction into account

resulted in a small, but systematic improvement. To test these changes, we first focused on the
top adjustment method (gBH-LSL) (Figure 4C). Taking the direction of change into account led to
a small but systematic improvement. Instead, not all bin construction methods using the propor-
tion of contribution into account had positive impact, but the Proportion-Adjusted Significance
(PAS) approach generally brought further slight improvements (Figure 4C). Repeating this for all
combinations of binning and adjustment methods (Figure 5A) confirmed the robustness of this
approach across adjustment methods.

Effect of the number of bins used

We tested the impact of the number of bins used on the results (Figure 5B). Both IHW and
gBH were fairly robust to the number of bins. There was an initial gain in recall for all methods
when increasing the number of bins from 5 to 8 and 12, but beyond this IHW did not show
any improvement. This is presumably because the cross-validation strategy, which requires a
certain number of hypotheses per bin to work properly. Instead, gBH sometimes showed further
increase in recall with larger numbers of bins, although the optimum differed across cell types.

Similar results using shrunk logFC instead of significance

It has often been argued that p-values are not the best way of ranking genes, and that fold-
changes can be more appropriate, although involving its own risks (Xiao et al., 2012). As the
accuracy of fold-change estimates in sequencing data is notoriously dependent on that of the
quantification, we tested whether using bulk-level, shrunk logFC estimates (Phipson, 2013) could
provide a better grouping of the hypotheses. While both approaches were superior in some
contexts, overall they appeared to provide a very similar increase in power (Figure 6).

Discussion

We have shown how simple strategies based on grouped FDR correction can leverage bulk
data in order to increase the precision and/or sensitivity of single-cell differential state analy-
sis. The methods discussed here are now all implemented in the *muscat* bioconductor package.
Because they operate at the level of multiple correction, they can however be used in combina-
tion with any differential analysis package. The methods could also be directly applied to other
problems, such as eQTL discovery in single-cell data (Bryois et al., 2022).
While the methods had good FDR control in the simulations, FDR was generally underesti-
mated in the real data. We believe that there are two main reasons for this. First, in order to
establish a ground truth, we removed from the evaluation a lot of hypotheses that were consid-
ered ambiguous (i.e. those which had an adjusted p-value between 0.5 and 0.1 in the full data),
which means that we are not computing precision on the full set of rejected hypotheses. This is a
limitation of this study, for which the accompanying simulations offers some mitigation. Second,
the ground truth established from the larger cohort effectively tests the generalizability of the
findings, and given patient heterogeneity, it is very likely that many differences are true of the
smaller comparison (i.e. would be reproduced if other samples of the same individuals had been
taken) but not generalizable.
In the case of the MS data, we had to standardize the cell type abundance of the 'artificial'
bulk data in order for the bbhw-methods to genuinely improve performance, otherwise the bulk
profiles would be dominated by differences in cell type proportions. Obviously, this cannot be
done with real bulk data. Therefore, a limitation of bulk-based hypothesis weighing is that it is

Figure 5 – **A:** Precision, recall, F1 score, and number of false positives (in unaffected cell/types) reported by each combination of bin- and adjustment-method. **B:** Precision and recall when using different numbers of bins. All values are at a nominal adjusted p-value threshold of 0.05.

207   unlikely to yield much improvement when the experimental condition studied involves major
208   expansion or depletion of some cell types. However, in such cases, we also do not expect the
209   method to deteriorate the results, making it safe for application.
210   Finally, an interesting finding of the study is that grouped Benjamini-Hochberg (Hu et al.,
211   2010) (in particular using the Least-Slope estimator) outperformed Independent Hypothesis

Figure 6 – TPR-FDR curves of the top method based either on bulk significance or shrunk logFC, compared to standard (no-prior) FDR.

Weighing (IHW) (Ignatiadis et al., 2016), although the difference was small. This is in contrast with the results obtained by Ignatiadis et al. (2016), who found gBH 'slightly anti-conservative'. If gBH underestimates FDR, it appears in our setting to be by very little. gBH also has the benefit of being deterministic and very fast to compute (we provide an implementation in the *muscat* package). While this might not generalize to other applications, both methods would be worth further study.

Funding

This work was supported by Swiss National Science Foundation (SNSF) project grant 310030_204869 to MDR. MDR acknowledges support from the University Research Priority Program Evolution in Action at the University of Zurich.

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Data, script, code, and supplementary information availability

The implementation of the methods can be found in the 'bulkIhw' branch of the *muscat* package (and will be in the next Bioconductor release). The scripts to reproduce the analyses and figures can be found at https://github.com/plger/bbhw.

References

Benjamini Y, Hochberg Y (2000). On the Adaptive Control of the False Discovery Rate in Mul-
tiple Testing With Independent Statistics. *Journal of Educational and Behavioral Statistics* **25**.
Publisher: American Educational Research Association, 60–83. https://doi.org/10.3102/
10769986025001060. URL: https://doi.org/10.3102/10769986025001060 (visited on
03/26/2025).

Benjamini Y, Krieger AM, Yekutieli D (2006). Adaptive linear step-up procedures that control the
false discovery rate. *Biometrika* **93**, 491–507. https://doi.org/10.1093/biomet/93.3.
491. URL: https://doi.org/10.1093/biomet/93.3.491 (visited on 03/26/2025).

Bryois J, Calini D, Macnair W, Foo L, Urich E, Ortmann W, Iglesias VA, Selvaraj S, Nutma E, Marzin
M, Amor S, Williams A, Castelo-Branco G, Menon V, De Jager P, Malhotra D (2022). Cell-type-
specific cis-eQTLs in eight human brain cell types identify novel risk genes for psychiatric and
neurological disorders. *Nature Neuroscience* **25**. Publisher: Nature Publishing Group, 1104–
1112. https://doi.org/10.1038/s41593-022-01128-z. URL: https://www.nature.com/
articles/s41593-022-01128-z (visited on 03/11/2024).

Crowell HL, Soneson C, Germain PL, Calini D, Collin L, Raposo C, Malhotra D, Robinson MD
(2020). muscat detects subpopulation-specific state transitions from multi-sample multi-condition
single-cell transcriptomics data. *Nature Communications* **11**. Number: 1 Publisher: Nature Pub-
lishing Group, 6077. https://doi.org/10.1038/s41467-020-19894-4. URL: https:
//www.nature.com/articles/s41467-020-19894-4 (visited on 11/30/2020).

Hu JX, Zhao H, Zhou HH (2010). False Discovery Rate Control With Groups. *Journal of the Amer-
ican Statistical Association* **105**, 1215–1227. https://doi.org/10.1198/jasa.2010.
tm09329. URL: https://www.tandfonline.com/doi/full/10.1198/jasa.2010.tm09329
(visited on 03/26/2025).

Ignatiadis N, Klaus B, Zaugg JB, Huber W (2016). Data-driven hypothesis weighting increases
detection power in genome-scale multiple testing. *Nature Methods* **13**. Publisher: Nature Pub-
lishing Group, 577–580. https://doi.org/10.1038/nmeth.3885. URL: https://www.
nature.com/articles/nmeth.3885 (visited on 06/28/2024).

Korthauer K, Kimes PK, Duvallet C, Reyes A, Subramanian A, Teng M, Shukla C, Alm EJ, Hicks SC
(2019). A practical guide to methods controlling false discoveries in computational biology.
Genome Biology **20**, 118. https://doi.org/10.1186/s13059-019-1716-1. URL: https:
//doi.org/10.1186/s13059-019-1716-1 (visited on 06/02/2022).

Macnair W, Calini D, Agirre E, Bryois J, Jäkel S, Smith RS, Kukanja P, Stokar-Regenscheit N, Ott
V, Foo LC, Collin L, Schippling S, Urich E, Nutma E, Marzin M, Ansaloni F, Amor S, Magliozzi
R, Heidari E, Robinson MD, et al. (2025). snRNA-seq stratifies multiple sclerosis patients into
distinct white matter glial responses. *Neuron* **113**. Publisher: Elsevier, 396–410.e9. https:
//doi.org/10.1016/j.neuron.2024.11.016. URL: https://www.cell.com/neuron/
abstract/S0896-6273(24)00873-0 (visited on 02/18/2025).

Maleki F, Ovens K, McQuillan I, Kusalik AJ (2019). Size matters: how sample size affects the
reproducibility and specificity of gene set analysis. *Human Genomics* **13**, 42. https://doi.
org/10.1186/s40246-019-0226-2. URL: https://doi.org/10.1186/s40246-019-0226-2
(visited on 04/07/2025).

Nakatsuka N, Adler D, Jiang L, Hartman A, Cheng E, Klann E, Satija R (2025). A Reproducibility Focused Meta-Analysis Method for Single-Cell Transcriptomic Case-Control Studies Uncovers Robust Differentially Expressed Genes. *bioRxiv*. https://doi.org/10.1101/2024.10.15.618577. eprint: https://www.biorxiv.org/content/early/2025/02/03/2024.10.15.618577.full.pdf. URL: https://www.biorxiv.org/content/early/2025/02/03/2024.10.15.618577.

Phipson B (2013). Empirical Bayes modelling of expression profiles and their associations. PhD thesis. University of Melbourne, Department of Mathematics and Statistics.

Robinson MD, McCarthy DJ, Smyth GK (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140. https://doi.org/10.1093/bioinformatics/btp616. URL: https://doi.org/10.1093/bioinformatics/btp616 (visited on 04/24/2023).

Waag R, Ziegler Lv, Sturman O, Sonder E, Leonardi J, Frei S, Gapp K, Germain PL, Bohacek J (2025). Distinct molecular mechanisms of stress habituation in the mouse hippocampus. Pages: 2025.03.04.641433 Section: New Results. https://doi.org/10.1101/2025.03.04.641433. URL: https://www.biorxiv.org/content/10.1101/2025.03.04.641433v1 (visited on 03/05/2025).

Xiao Y, Hsiao TH, Suresh U, Chen HIH, Wu X, Wolf SE, Chen Y (2012). A novel significance score for gene selection and ranking. *Bioinformatics* **30**, 801–807. https://doi.org/10.1093/bioinformatics/btr671. eprint: https://academic.oup.com/bioinformatics/article-pdf/30/6/801/48921601/bioinformatics_30_6_801.pdf. URL: https://doi.org/10.1093/bioinformatics/btr671.