Title Page

Title

Linking neuron-axon-synapse architecture to white matter vasculature using high-resolution

multimodal MRI in primate brain

Authors

Ikko Kimura[1, 2], Takuya Hayashi[1, 3, *], Joonas A. Autio[1, *]

Affiliations

[1]Laboratory for Brain Connectomics Imaging, RIKEN Center for Biosystems Dynamics

Research, Kobe, Japan

[2]Neurophysics Group, Danish Research Centre for Magnetic Resonance, Copenhagen

University Hospital - Hvidovre and Amager, Copenhagen, Denmark

[3]Department of Brain Connectomics, Kyoto University Graduate School of Medicine, Kyoto,

Japan

*Corresponding authors

Joonas A. Autio, Ph.D.; Takuya Hayashi, M.D., Ph.D.

Laboratory for Brain Connectomics Imaging

RIKEN Center for Biosystems Dynamics Research

6-7-3 MI R&D Center 3F, Minatojima-minamimachi

Chuo-ku, Kobe 650-0047, Japan

E-mail: joonas.autio@riken.jp, takuya.hayashi@riken.jp

Abstract

Blood vessels and axons align outside the brain due to shared growth factors. However, this

neuron-axon-synapse and vessel relationship within the brain white matter remains unclear,

primarily due to the technical challenges of charting the complex trajectories of fiber tracts

and the dense network of arteries. Consequently, the organizational logic and neurometabolic

factors shaping white matter vasculature remain poorly understood. Here, we address these

questions using high-resolution multimodal MRI, in vitro neuron density, and receptor

autoradiography in macaque monkeys. In superficial white matter, vascularity exhibited

parallel alignment with the cortical surface. This vascularity showed negligible dependence

on overlying gray matter neuron density ($R^2 = 0.01$), minimal dependence on white matter

myelination ($R^2 = 0.10$), and moderate correlation with receptor density ($R^2 = 0.27$). These

suggest an association of vascularity with energy demands and axonal branching. In deep

white matter, axon geometry, density, and proximity to the cortical surface predict vascular

volume with high precision ($R^2 = 0.62$). Overall, these findings establish a relation between

neuron-axon-synapse architecture and white matter vasculature in the primate brain, offering

advances in understanding the organization and pathophysiology of white matter.

1. Introduction

White matter vasculature plays a crucial role in brain function and aging-related diseases. Nevertheless, its organization and function remain underexplored, despite supplying a substantial proportion of brain's blood flow (Wu et al., 2013). A deeper understanding of white matter vascularity is essential for elucidating its contributions to neural function (Autio and Roberts, 2014; Huang et al., 2023, 2020; Schilling et al., 2023; Wang et al., 2023; Weber, 2002), plasticity (Kimura et al., 2024; Sampaio-Baptista and Johansen-Berg, 2017), and pathology (Chen et al., 2013; Chojdak-Łukasiewicz et al., 2021). Early post-mortem studies mapped the arterial supply of white matter from pial vessels (Duvernoy et al., 1981), and research outside the brain showed that blood vessels and axons align due to shared growth factors (Carmeliet and Tessier-Lavigne, 2005). However, the fundamental principles governing the brain white matter vascular organization remain poorly understood (Smirnov et al., 2021), limiting insights into its role in brain health and disease.

The neurometabolic factors influencing white matter vasculature also remain uncertain. Energy demand, reflected by mitochondria density, varies among tracts (Kageyama and Wong-Riley, 1986, 1984; Perge et al., 2012), and regions with high energy demands tend to have denser vascularization (Borowsky and Collins, 1989; Perge et al., 2012). Energy consumption in white matter stems from several biophysical processes, including protein and lipid synthesis, axonal transport of molecules and organelles (Bernstein and Bamburg, 2003) and, to a lesser extent, axonal action potentials (Harris and Attwell, 2012). Although myelin is generally thought to conserve energy through saltatory conduction, simulation studies suggest that the energetic costs of myelin synthesis and oligodendrocyte maintenance may offset these savings (Harris and Attwell, 2012). Variations in gray matter neural and receptor density may further impact white matter energy demands, as axonal transport is essential for

moving synaptic vesicles between the soma and axon terminals (Bernstein and Bamburg, 2003; Goldstein et al., 2008). However, empirical evidence linking neurometabolic factors to white matter vascularization remains limited.

Advancements in high-resolution ferumoxytol-weighted MRI have significantly improved the quantification of brain vascularity (Autio et al., 2024a; Bernier et al., 2019). Our recent work demonstrated that cortical layer vasculature was related to regional variations in neuron, and synaptic or myelin densities (Collins et al., 2010; Froudist-Walsh et al., 2023) in macaque monkeys (Autio et al., 2024a). Additionally, diffusion MRI (dMRI) provides crucial insights into the organization and microstructure of white matter tracts, including neurite density, axon orientation and its dispersion (Basser et al., 1994; Zhang et al., 2012), enabling direct comparisons between fiber architecture and vascular organization.

In this study, we applied multimodal MRI to investigate the neurometabolic factors influencing white matter vascular organization in macaques. By estimating vascular volume in both superficial and deep white matter, we identified substantial inter-areal variability, comparable to that seen in cortical gray matter. To understand the anatomical basis of this variability, we examined the relationship between white matter vasculature and microstructure, as well as the neuronal and synaptic architecture of overlying cortical areas. As a preview, we found that: 1) myelinated regions exhibited lower vascular density, suggesting reduced energy demands, though this explains only a small portion of the variability, consistent with the result of simulation study (Harris and Attwell, 2012); 2) white matter vascularity correlated more strongly with cortical receptor density than with neuron density; and 3) the geometrical arrangement, density, and distance of axons to the cortical surface predict white matter vasculature with high precision. These findings provide new insights into the metabolic and structural determinants of white matter vascular organization.

2. Methods

2.1 Animals

MRI data from four macaques (Macaca mulatta, age: 4 – 6 years, weight: 7.4 – 8.4 kg) (Autio et al., 2024a) were used to investigate the vascularity and microstructure of the white matter in the brain. All experiments were approved by the Animal Care and Use Committee of the Kobe Institute of RIKEN (MA2008-03-11) and conducted in accordance with the institutional guidelines for animal experiments, the Basic Policies for the Conduct of Animals Experiments in Research Institution (MEXT, Tokyo, Japan), and the Guidelines for the Care and Use of Laboratory Animals (ILAR, Washington DC, USA).

2.2 Image acquisition

Structural, ferumoxytol-weighted multi-echo gradient-echo and dMRI data were sourced from our prior studies (Autio et al., 2024a, 2024b, 2020) using the same macaques across modalities (n = 4). All MRI scans were acquired using a Siemens Prisma 3T scanner (Siemens, Erlangen, Germany) equipped with a 24-channel multi-array RF coil for non-human primate brains (Rogue Research, Montreal, Canada / Takashima Seisakusho KK, Tokyo, Japan) (Autio et al., 2020). The animals were scanned under anesthesia and physiological conditions were continuously monitored throughout the experiments. Detailed experimental procedures and imaging protocols can be found in Autio et al., (2024b) for structural MRI, in Autio et al., (2024a) for multi-echo gradient-echo MRI, and in Autio et al., (2020) for dMRI.

In brief, multi-echo gradient-echo MRI data with 10 equidistant echo-times (TE1 = 3.4 ms, TE interval = 2.4 ms) were acquired before and after an intravascular injection of 12

mg/kg ferumoxytol to investigate brain vascularity (0.32 mm isotropic resolution). A subset of these macaques was rescanned after ferumoxytol injection using three equidistant TEs (TE1 = 5.0 ms, interval = 4.0 ms) with spatial resolution adjusted to match the critical (spatial) sampling frequency of intra-cortical vessels (0.23 mm isotropic resolution). Structural T1-weighted (T1w) MPRAGE and T2-weighted (T2w) SPACE-FLAIR images were acquired at 0.32 mm isotropic resolution to accurately reconstruct white matter and pial surfaces. Additionally, T1w/T2w-FLAIR was used to investigate myeloarchitecture across cortical layers and within the adjacent superficial white matter, suppressing Gibbs' ringing artifact. The Human Connectome Project (HCP)-style diffusion MRI data acquisition protocol was applied to study white matter tract orientation and microstructure (Autio et al., 2020; Glasser et al., 2013). These images were obtained at 0.9 mm isotropic resolution with b-values of 0, 1000, 2000, and 3000 mm/s^2, and with 256 diffusion orientations.

2.3 Image Analysis

Structural and dMRI data were preprocessed using a non-human primate version of the preprocessing pipelines developed by the HCP (https://github.com/Washington-University/NHPPipelines) (Autio et al., 2020; Donahue et al., 2016; Glasser et al., 2013; Hayashi et al., 2021). These pipelines utilize FreeSurfer (version 6.0.1), wb_command (version 1.5.0), and FMRIB Software library (FSL; version 6.0.4). Detailed preprocessing procedures for structural and diffusion MRI can be found in Autio et al., (2020).

Briefly, T1w and T2w-FLAIR images were aligned to an anterior-posterior commissural (AC-PC) space, corrected for B_1-bias field, and nonlinearly registered to the

Mac30BS standard space (Hayashi et al., 2021). The pial and white surfaces and subcortical structures were delineated using FreeSurfer. These surfaces were registered to the Mac30BS standard space based on sulcus landmarks using MSMSulc, and created a mid-thickness surface. All surfaces were visually inspected for errors. When required, manual corrections were made on a wm.edit.mgz or subtle manual repositioning was performed to a surface using FreeSurfer 7.1 (by JAA).

Using the corrected white matter and pial surfaces as references, we generated a layer for superficial white matter positioned 0.32 mm beneath the white matter surface(Kirilina et al., 2020). We additionally created 12 equivolumetric layers (ELs) (Autio et al., 2024a, 2024b; Bok, 1929; Van Essen and Maunsell, 1980). These ELs were positioned within the cerebral cortex (i.e., EL1a, EL1b, EL2a, ... EL6a, and EL6b), with EL1a situated just beneath the pial surface and EL6b situated just above the white matter surface. This nomenclature was intended to facilitate comparison with anatomical cortical layers (e.g., Ia, Ib, Ic, ... VIb).

Vascularity was evaluated using ferumoxytol-weighted multi-echo gradient-echo imaging (Autio et al., 2024a). Quantitative R_2* fitting was performed on the multi-echo gradient-echo images before and after ferumoxytol injection using hMRI toolbox (Tabelow et al., 2019). The pre-ferumoxytol R_2* maps were subtracted from post-ferumoxytol maps to generate ΔR_2* maps, providing indirect measures of vascular volume (Boxerman et al., 1995; Kim et al., 2013). These ΔR_2* volume maps were then mapped onto native cortical layers using ribbon-constrained volume-to-surface-mapping in wb_command and resampled to 32k and 164k meshes. Subsequently, these ΔR_2* maps were corrected for the orientation of the cerebral cortex relative to the direction of the static magnetic field (B_0) (Autio et al., 2024a). The median value of each area in M132 Lyon (Markov et al., 2014) or Julich (Froudist-Walsh

et al., 2023) macaque brain atlas was calculated and averaged across the macaques (n = 4). In the M132 Lyon atlas (n = 91 per hemisphere), areas 24a, 24b, 29_30, and V2 were excluded from further analysis (resulting in n = 87 per hemisphere), and, in the Julich atlas (n = 109, left hemisphere), areas a24ab, p24ab, 24ab, 29/30, V2v, and V2d were excluded (resulting in n = 103) due to minor yet significant errors in white surface estimation. Additionally, post-ferumoxytol images with higher spatial resolution (0.23 mm isotropic) were averaged across runs and TEs to improve signal-to-noise ratio. Hessian-based Frangi 'vesselness' filter (https://jp.mathworks.com/matlabcentral/fileexchange/24409-hessian-based-frangi-vesselness -filter) was applied to detect vessels in the 3D volumetric images. To enable surface visualization of vessels, 164k surface mesh was resampled to ultra-high resolution 656k mesh, resulting in an average 0.14 mm spacing between vertices in the superficial white matter. In cortical gray matter, this spacing is smaller than the average distance between the penetrating vessels (Autio et al., 2024a; Weber et al., 2008). To visualize the macrovessels, spatial gradients on each layer were calculated using wb_command -metric-gradient.

dMRI data were intensity-normalized, distortion corrected, and corrected for eddy current and subject motion. The corrected images were co-registered to the native AC-PC aligned T1w image. To investigate the relationship between vascularity and major fiber bundles, tractography was performed to define the location of commissural fibers connecting between bilateral identical areas. We selected these bundles for analysis because they can be robustly delineated and consistently traverse from the cerebral cortex to the deep white matter. Tractography was performed using MRtrix3 (version 3.0.4) (Tournier et al., 2019). First, the basis functions to deconvolve dMRI data to create the fiber orientation density (FOD) were calculated with the dhollander algorithm (i.e., basis function was estimated for gray matter, white matter, and cerebrospinal fluid and for each b value). These basis functions were used

to calculate FOD on each voxel with multi-shell multi-tissue constrained spherical deconvolution. Tissue boundary between the white and cortical gray matter for the seed of tractography was created with preprocessed structural MRI data. One hundred million streamlines were generated with anatomically constrained tractography (maximum length: 200 mm, cut off for FOD amplitude: 0.06) setting the boundary between the white and gray matter as seeds. These streamlines were refined from 100,000,000 streamlines to 10,000,000 streamlines with Spherical-deconvolution Informed Filtering of Tractograms to alleviate the overestimation of the number of streamlines. Commissural fibers connecting between each identical bilateral area defined by M132 Lyon macaque brain atlas (Markov et al., 2014) were extracted. Finally, automated fiber-tract quantification (https://github.com/yeatmanlab/AFQ) (Yeatman et al., 2012) was used to exclude outliers, whose distance from the core of the tract is more than three standard deviations (SD), within each bundle and allocated 20 equidistant points on each tract. The bundles with more than five streamlines for each macaque were utilized for further analysis. ΔR_2^* and the other microstructural property on each point was calculated to obtain the median value for each point of each bundle and was averaged across macaques.

To investigate the relationship between vascularity and white matter microstructure, we estimated microstructural properties from dMRI using Neurite Orientation Dispersion and Density Imaging (NODDI) toolbox (version 1.0.3; www.nitrc.org/projects/noddi_toolbox) (Zhang et al., 2012). The NODDI models three compartments (intra-neurite, extra-neurite, and cerebrospinal fluid), using multi-shell high-angular resolution dMRI data (Zhang et al., 2012). Two indices were used for further analysis: the neurite density index (NDI), representing the density of neurite compartments, and the orientation dispersion index (ODI), reflecting the dispersion of neurite orientations. FSL's *dtifit* was additionally applied in each

voxel to estimate the diffusion orientation of major fibers in each voxel.

2.4 Statistical Analysis

The group-average ΔR_2^* values between the gray and superficial white matter were

compared using a two-tailed paired two-sample t-test. ΔR_2^* values from EL4a were used to

represent the gray matter. To assess how inter-areal variability in ΔR_2^* of the superficial

white matter is influenced by the overlying gray matter, we calculated the Pearson's

correlation coefficient between ΔR_2^* values of the gray matter and superficial white matter.

To corroborate the relations between ΔR_2^* and microstructural properties within each area,

Pearson's correlation coefficients were calculated between ΔR_2^* and T1w/T2w-FLAIR

(myelin map). We also calculated the coefficients with neuron density and receptor density in

the overlying gray matter of each area to evaluate how neuron and synapse affect the

vasculature in the superficial white matter. The myelin map data was taken from our previous

study (Autio et al., 2024b), while neuron and receptor densities were from the literature

(Froudist-Walsh et al., 2023). Consistent with our previous study (Autio et al., 2024a), we

focused on the total receptor density rather than individual receptor densities, as these were

highly correlated with each other. To test whether these Pearson's correlation coefficients

differed between the gray matter and superficial white matter, we applied two-tailed bootstrap

testing to each of these coefficients.

To examine vessel orientation characteristics in the superficial and deep white matter,

we investigated the relationship between B_0 orientation bias and the alignment of the cortex.

The vessel B_0 orientation bias is defined by the following equation (Ogawa et al., 1993):

$\Delta R_2^* \propto cos^2\theta_{vessel, B_0}$ (1)

where, θ_{vessel, B_0} represents the angle between the primary orientation of the vessels within

a voxel and the B_0 direction in the individual's original MRI space. In the superficial white

matter, vessel alignment relative to the normal of cortex was indirectly assessed by

calculating Pearson's correlation coefficients between ΔR_2^* and $cos^2\theta_{normal\ of\ cortex, B_0}$.

To model changes in ΔR_2^*, NDI, and ODI from the superficial to deep white matter,

we fitted the group-average values of each metric against the relative distance from the

superficial white matter to the midline (i.e., 0% distance at the white matter surface and 100%

at the midline [corpus callosum]). The modeling was performed using the following equation:

$\Delta R_2^* = -a\ e^{-b\ \times\ relative\ distance} + c$ (2)

using the *fitnlm* function in Matlab R2016b (MathWorks). We evaluated the relationship

between ΔR_2^* and NDI, ODI, or path length across the white matter bundles using the

Pearson's correlation coefficient.

To determine whether ΔR_2^* at each location could be predicted by fiber density,

tract dispersion, and distance from superficial white matter to the midline (e.g., arteries

branching or weakening toward deeper white matter), we used a forward stepwise nonlinear

regression model. Here, we assumed a nonlinear relationship between the relative distance

from the superficial white matter to the midline and ΔR_2^*, while treating the other properties

as linear. Model selection was guided by the Bayesian information criterion. We evaluated the

prediction accuracy using a leave-one-dataset-out (i.e., one white matter bundle out)

cross-validation method. The significance of each Pearson's correlation coefficient was

assessed using two-tailed bootstrap testing. All statistical analyses were conducted using the

Statistics and Machine Learning Toolbox in Matlab R2016b (MathWorks), with a

significance threshold of $P < 0.05$. All P-values were Bonferroni-corrected for multiple

comparisons.

3. Results

3.1 Charting Superficial White Matter Vascularity using Ferumoxytol-weighted MRI

The gradient map of mean ferumoxytol-weighted gradient-echo images provided clear visualization of macrovessels in superficial white matter (Fig. 1). Specifically, the ferumoxytol-weighted images enabled robust identification of macrovessels projecting from the pial surface and traversing along the superficial white matter (Fig. 1). These vessels likely correspond to those categorized as Durney's Group 5 or 6 arteries as these vessels did not exhibit noticeable contrast in the pre-ferumoxytol images (Duvernoy et al., 1981).

Comparison of vessel orientation in cortical gray matter and superficial white matter is illustrated in Figure 2 using spatial gradients of the mean ferumoxytol-weighted image. In the gray matter, the mean-intensity gradients exhibited ring-shaped patterns (Fig. 2A), indicative of vessels traversing perpendicular to the surface of the cerebral cortex (Autio et al., 2024a). Conversely, in the superficial white matter, the gradients exhibited rod-shaped patterns (Fig. 2C) indicative of vessels traversing parallel to the white matter surface. These distinct vessel orientations are in good agreement with brain vessel anatomy (Duvernoy et al., 1981), and support Viessmann and colleagues' indirect MRI observations of vessel orientation differences between gray and white matter (Viessmann et al., 2022). The sharp contrast was observed in orientation of vessels between equidistant superficial white matter (Fig. 2C) and the innermost EL of cortical gray matter (EL6b; Fig. 2B), supporting that the superficial white matter boundary was precisely delineated.

We also observed areal differences in vascularization patterns. For instance, in V1 we observed sparse distribution of macrovessels in superficial white matter, contrasting the more uniform and dense distribution in the gray matter. In the prefrontal cortex, superficial

white matter vessels exhibited orthogonal orientations whereas less distinct and more subtler

variations in vessel orientation were observed near the temporal pole (Supp. Fig. 1).

Figure 1. Charting macrovessels in the superficial white matter using ferumoxytol-weighted vessel-density informed MRI. Circular gradients represent vessels oriented perpendicular to the superficial white matter, while rod-like gradients denote vessels traversing along the superficial white matter. Gradients were notably stronger in the dorsal regions of the superficial white matter compared to the ventral regions. The black box indicates the location of the magnified view.

Figure 2. Distinct macrovessel arrangement in gray and white matter. Spatial gradients of the ferumoxytol-weighted image on the equivolumetric layer (EL) **(A)** EL4a, located proximal to cortical midthickness, and **(B)** EL6b, located adjacent to the white matter surface. **(C)** Equidistant superficial white matter surface located beneath the cortical gray matter. Note the striking contrast in the gradient pattern between the gray matter (circular structures) and superficial white matter (rod-like structures): circular gradients indicate the presence of vessels oriented perpendicular to the orientation of the cerebral cortex whereas rod-like gradients indicate vessels traversing along the white and gray matter interface The red and green boxes indicate the locations of zoomed views.

Quantitative analysis revealed significant differences in vessel orientation bias

between gray matter and superficial white matter. In the gray matter, correlation between

ΔR_2^* and cerebral cortex orientation ($\cos^2\theta$ in Eq. (1)) were negative (r = -0.32 – -0.26, t_3 =

-20.25, P = 0.0003), indicating that larger vessels predominantly align perpendicularly to the

cortical surface. Conversely, the superficial white matter showed positive correlations (r =

0.05 – 0.11; t_3 = 6.69, P = 0.0068), indicating parallel orientation of vessels relative to the

cortical surface (Supp. Fig. 2).

We quantified vascularity in the superficial white matter by measuring R_2^* before

and after ferumoxytol contrast agent injection, using ΔR_2^* as a well-established proxy for

vascular volume (Kim et al., 2013). To mitigate the influence of macrovessels on ΔR_2^* (Fig.

1), we ascribed median ΔR_2^* to each M132-atlas parcel (Fig. 3A, upper panel) (Autio et al.,

2024a). While occipital and temporal grey matter exhibited a decrease in ΔR_2^* from lower- to

higher-order areas, no such trend was found in the superficial white matter. Interestingly,

ΔR_2^* increased gradually from posterior to anterior regions within the frontal superficial

white matter (Fig. 3A, bottom panel), a pattern being absent in gray matter. Superficial white

matter vascularization was independent of gray matter vascularization (r = -0.10, P = 0.34;

Fig. 3C), consistent with distinct energy demands between these regions (Harris and Attwell,

2012). Although ΔR_2^* was systematically lower in superficial white matter compared to gray

matter (Fig. 3B; paired one-sample t-test, t_{87} = 22.2, P < 0.001), variability in superficial

white matter vascularization was comparable to gray matter (gray matter: SD = 3.56,

interquartile range [IQR] = 4.00; superficial white matter: SD = 3.72, IQR = 4.68; $F_{87,87}$ =

0.92, P = 0.69).

Figure 3. Comparison of vasculature in cortical gray matter and superficial white matter. (A) Vascular volume in cortical gray matter (upper panel) and superficial white matter (lower panel), estimated using ferumoxytol-induced change in transverse relaxation rate (ΔR_2^*). To reduce the influence of macrovessels, data was parcellated using the M132 areal atlas (Markov et al., 2014), and median ΔR_2^* value was assigned to each cortical area. **(B)** Box plot showing comparison of ΔR_2^* values between cortical gray matter and superficial white matter. Red crosses indicate outliers: in gray matter V1 and entorhinal cortex, and in superficial white matter areas 9, 11 and 9/46d. *** $P < 0.001$. **(C)** Scatter plot between vascular volume in superficial white matter and cortical gray matter (r = -0.01, P = 0.34).

3.2 Comparison between Superficial White Matter Vasculature and Overlying Cortical Neuron-Synapse Architecture

Given the large inter-areal heterogeneity in superficial white matter vasculature, we sought to investigate how neuroanatomical factors in the white matter and overlying cortical gray matter might influence superficial white matter energy consumption. To address this, we examined correlations between white matter ΔR_2^* and microstructural metrics, including neuron, myelin, and synaptic densities (Fig. 4) (Autio et al., 2024a; Collins et al., 2010; Froudist-Walsh et al., 2023). In the cortical gray matter, ΔR_2^* demonstrated significant positive correlations with neuron density (r = 0.53, *P* < 0.001) and myelin density (r = 0.54, *P* < 0.001) and a negative correlation with receptor density (r = -0.42, *P* < 0.001) (Autio et al., 2024a) (Fig. 4A–D, G). Conversely, within the superficial white matter, ΔR_2^* was positively correlated with receptor density (r = 0.50, *P* < 0.001) and negatively correlated with myelin density (r = -0.31, *P* < 0.001) (Fig. 4A, C, E–G), with no significant correlation observed for neuron density (r = -0.10, *P* = 0.36). Bootstrap testing confirmed significant differences between these correlations in the cortical gray matter and superficial white matter across all metrics (all *P* < 0.001). The M132-atlas parcellated ΔR_2^* also showed a significant positive correlation with myelin (r = 0.67, *P* < 0.001) within the cortical gray matter and a significant negative correlation with myelin (r = -0.28, *P* = 0.0014) within the superficial white matter (Supp. Fig. 3).

Figure 4. Neuroanatomical correlates of superficial white matter vascularity. Parcellated maps of cortical **(A)** neuron density (Collins et al., 2010), **(B)** T1w/T2w-FLAIR myelin (Autio et al., 2024a), **(C)** receptor density (Froudist-Walsh et al., 2023), **(D)** vascular volume assessed using ferumoxytol-induced change in transverse relaxation rate (ΔR_2*) (Autio et al., 2024b). Superficial white matter **(E)** T1w/T2w-FLAIR myelin and **(F)** vascular volume. **(G)** Correlation between vascular volume and neuroanatomical factors. *** $P <$ 0.001 (Bonferroni-corrected).

3.3 Comparison between White Matter Density, Orientation and Vasculature

Finally, we explored the vascularity deep in the white matter using vessel-density informed ferumoxytol-weighted MRI. To delineate the vessels in the volume-space, we applied a Hessian-based Frangi 'vesselness' filter to the TE-averaged mean intensity images (Frangi et al., 1998). This analysis revealed a large number of vessels traversing from the pia mater to white matter ($2800 - 4220$ mm^3, $13–15$ % of white matter voxels; Fig. 5A). Visual inspection of these vessels revealed that several vessels projected toward lateral ventricles, likely corresponding to the large draining veins. Several vessels appeared to traverse along the major tracts (e.g., commissural fibers connecting brain hemispheres), raising the question whether the feeding arterial vessels are organized along the major tract bundles (Smirnov et al., 2021).

To corroborate this question, we determined vessel orientation using a Frangi-filter and assessed fiber tract orientation using diffusion tensor analysis (Basser et al., 1994). Our analysis revealed that, in most cases, the vessels were aligned with the primary orientation of axons within the commissural fiber bundles (Fig. 5A; white arrows). However, the spatial resolution of dMRI (0.9 mm isotropic; 0.73 mm^3) was 60-fold lower than that of vessel imaging (0.23 mm isotropic; 0.012 mm^3), making it challenging to accurately attribute individual white matter bundles with their corresponding vessels, particularly smaller ones.

To overcome the spatial resolution limitation, we compared vascular density with NODDI, a model that estimates neurite density and orientation dispersion within each white matter voxel (Zhang et al., 2012). Specifically, we focused on commissural fibers connecting identical bilateral regions, as these long bundles can be robustly delineated (Fig. 5B). Within each bundle, as the relative distance from the white surface to the midline increased, ΔR_2* and ODI exhibited exponential decay (ΔR_2*: 11.64 e$^{-0.05 \times \text{relative distance}} +19.13$; ODI: 0.30 e$^{-0.02}$

$\times$ relative distance $+ 0.07$), whereas NDI showed exponential growth (-0.32 e$^{-0.01 \times \text{relative distance}} +$

0.87) (Fig. 5C-E; for the relationship with raw axon path length, see Supp. Fig. 4). Across all

bundles, ΔR_2^* was significantly negatively correlated with NDI (r = -0.51, $P < 0.001$) and

positively correlated with ODI (r = 0.67, $P < 0.001$), while it was not significantly correlated

with path length (r = -0.27, $P = 0.32$).

Given the systematic variation between microstructure, brain geometry and

vasculature, we additionally asked whether the white matter microstructure may predict the

variation in white matter vasculature. To address this question, we performed a forward

stepwise nonlinear regression model selected by Bayesian information criterion (see

methods). This analysis revealed that ΔR_2^* along each bundle can be predicted by the NDI,

ODI, and the relative distance from the white matter surface to the midline (corpus callosum)

with high precision (r = 0.79, $P < 0.001$; Fig. 5G).

Figure 5. Modeling white matter vascularity using tract density, dispersion and distance from cortical surface. (**A**) Diffusion tensor orientation (red, blue, and green color indicate the orientation of major fibers are mainly in right-left, superior-inferior, and anterior-posterior, respectively) overlaid with maximum intensity projection image of ferumoxytol-weighted and frangi-filtered vessel maps (red-yellow). (**B**) Diffusion tractography. (**C**) ferumoxytol-induced change in transverse relaxation rate (ΔR_2^*), (**D**) neurite density index (NDI), and (**E**) orientation dispersion index (ODI) values calculated from white matter surface (0%) toward the cortical midline (100%). (**F**) Pearson's correlation coefficients between ΔR_2^* and NDI, ODI, or path length across tracts. (**G**) Relationship between the ΔR_2^* within the white matter and the value predicted by the microstructural factors. Solid red line

1　indicates the fit by the linear model, while dashed lines indicate 95% confidence interval. ***

2　$P < 0.001$ (Bonferroni-corrected).

4. Discussion

In this study, we explored the relationship between neuron-axon-synapse architecture and the organization of white matter vasculature using high-resolution multimodal MRI and quantitative anatomical data in a non-human primate brain. Our novel surface presentation of superficial white matter vasculature revealed that the spatial arrangement of cortical synapses and myelins, rather than neurons, significantly influences the white matter vascularity. In deep white matter, we found that the geometric arrangement of white matter bundles, neurite density, and the distance from the cortical surface are key determinants of white matter vascular organization. These findings represent both a methodological and conceptual advancement in white matter vasculature MRI, offering potential clinical applications in understanding aging-related white matter pathologies.

4.1 Heterogeneity of Superficial White Matter Vasculature

The superficial white matter macrovasculature is largely unexplored area with limited studies (Duvernoy et al., 1981; Hase et al., 2019; Smirnov et al., 2021). It displays a complex and heterogeneous organization with several notable organizational features. In regions where white matter blades are exceptionally thin (< 0.5 mm), such as underneath V1, spatial constraints confine vessel trajectories to the white matter surface. Certain vessels take the most direct route to the lateral ventricles, possibly reflecting preferential venous drainage patterns (Smirnov et al., 2021). Additionally, some of the macrovessels exhibit orthogonal orientations, appearing to cross each other (Supp. Fig. 1; arrows). Additionally, some of the macrovessels exhibit orthogonal orientations, appearing to cross each other (Supp. Fig. 1; arrows). These vessels may be the arteries that deliver oxygen and nutrients to cortical layer

VI (Duvernoy et al., 1981). Hypoperfusion or ischemia in the superficial white matter

vasculature may selectively disrupt energy supply to cortical layer VI, potentially impairing

corticothalamic and corticoclaustral circuits (Wiser and Callaway, 1996).

One surprising discovery of the study was the substantial two-fold variability in the

superficial white matter vasculature (Fig. 3A, B). This heterogeneity may reflect the diverse

functional properties of different white matter regions. For instance, axonal conduction

velocities vary three-fold (3.1 – 9.4 m/s) (Innocenti et al., 2014), and the frequency bands

involved in inter-areal communication range from theta to gamma band (Shu et al., 2003;

Vezoli et al., 2021). Such functional diversity is enabled by differences in axon diameter and

myelination, influencing local energy requirements and vascular densities (Perge et al., 2012).

However, we found negligible correlation between the vascular volume in gray matter and

superficial white matter (Fig. 3C), implying that the metabolic demands are driven by distinct

biophysical mechanisms. This provides experimental validation to the previous modeling

study (Harris and Attwell, 2012). That study suggested that, while energy demand in gray

matter is primarily influenced by the frequency of electrical activity (e.g., Na^+-K^+ ATPase of

neuronal cell membrane potential), that in white matter is primarily determined by distinct

biophysical processes related to the maintenance of axons and their surroundings.

Another unexpected finding was that myelin density, as assessed using

T1w/T2w-FLAIR, accounted for only about 10% of the variance in the superficial white

matter vasculature (Fig. 4G). This result is surprising given that myelin is generally

considered a key factor in reducing the energetic demands of axonal conduction (Saab et al.,

2013), yet consistent with a simulation study suggesting that myelination reduces energy

consumption by only about 5% (Harris and Attwell, 2012). Meanwhile, we acknowledge that

the utility of T1w/T2w-FLAIR as a biomarker of myelin may be more limited in the white

matter than cortical gray matter (Glasser et al., 2014; Sandrone et al., 2023). Thus, further histological investigations are essential to corroborate how myelin, and its supporting cells, such as oligodendrocytes, influence white matter vascularization (Biswas et al., 2020; Gao et al., 2023).

4.2 Cortical Synaptic Architecture Shapes Superficial White Matter Vasculature

Given that axons originate from neuronal cell bodies, and growth factors guide the alignment of axons and arteries outside the brain (Carmeliet and Tessier-Lavigne, 2005), we initially hypothesized that white matter vascularization would correlate with cortical neuron density. However, we found that the vascular density in the superficial white matter was not significantly associated with neuron density (axon origin) (Fig. 4G). Instead, it showed a positive correlation with receptor density (axon terminals) in the overlying cortical gray matter (Froudist-Walsh et al., 2023). This relationship suggests that the metabolic demand in white matter may be influenced more by the axonal branches that are rich in mitochondrial density (Tao et al., 2014), rather than by the density of axons from soma.

Additionally, axonal transport may contribute to the relationship between cortical receptor and superficial white matter vascular densities (Fig. 4G). Transportation of proteins and nutrients between axon terminals and cell bodies occurs in both anterograde (from cell body toward synapse) and retrograde (from synapse toward cell body) directions. The movement of organelles and particularly the maintenance of the actin cytoskeleton are thought to significantly contribute to white matter energy consumption (Fath and Lasek, 1988; Harris and Attwell, 2012). Considering that receptors are directly associated with retrograde transport, the greater number of axon terminals may facilitate large-scale

retrograde transport, and, thus, increase in energy consumption and vessel density. Meanwhile, the proportion of energy used for anterograde and retrograde transport remains debated, with estimates ranging from 5% to 50% of total white matter energy consumption (Bernstein and Bamburg, 2003; Daniel et al., 1986; Engl and Attwell, 2015; Harris and Attwell, 2012), and more advanced research designs are needed to clarify the energy dynamics of these processes.

Another potential factor is white matter plasticity (Kimura et al., 2024; Sampaio-Baptista and Johansen-Berg, 2017). Increased neural synchronization (between cortical regions) may induce microstructural changes within white matter (Lazari et al., 2022; Sampaio-Baptista et al., 2021), such as myelination or oligodendrogenesis (Gibson et al., 2014). Given that synchronization also increases receptor densities through spike-timing-dependent plasticity (Caporale and Dan, 2008; Lamsa et al., 2010), high receptor density may indicate frequent plasticity in the underlying white matter, leading to increased energy demand.

4.3 White Matter Vasculature Reflects Fiber Architecture

Our study also revealed that macrovessels tend to align with the major orientation of axons. This supports previous human MRI studies that showed B_0 orientation bias in ΔR_2* (Hernández-Torres et al., 2017) and T_2*-weighted signal-intensity (Viessmann et al., 2022). This alignment may reflect shared growth factors like neuropilin-plexin signaling, which regulate the development of both axons and vessels (Carmeliet and Tessier-Lavigne, 2005).

The alignment of feeding arteries with fiber tracts may in part explain why blood volume is higher in regions where multiple white matter tracts intersect, such as beneath

multimodal association areas that form a dense interconnected network (Fig. 5B) (Markov et al., 2014). This axonal-vascular relationship is influenced by anatomical constraints and may also relate to variations in axon diameter (Fig. 5F). Axons with larger diameters transmit signals faster and require more energy to maintain their function, which could result in increased vessel density to support the higher energy demands (Harris and Attwell, 2012; Perge et al., 2012). In contrast, smaller diameter axons, which constitute the majority of axons in the central nervous system, conduct signals more slowly but are more energy-efficient (Perge et al., 2012). Given that high NDI may reflect small axon diameter in that region (Alexander et al., 2010), the negative relationships between vessel density and NDI may be explained by the energy demand defined by axon diameter.

Overall, these findings highlight the complex interplay between anatomical structures that influence the vascularization of white matter. Indeed, a forward stepwise nonlinear regression model selected by Bayesian information criterion predicts white matter vascularity with high precision ($R^2 = 0.62$) (Fig. 5G). This information could be valuable for modeling brain blood circulation, understanding white matter pathology, and guiding neurosurgical operations. However, it will be fundamental to augment vascular models with classification of arteries and veins, which is more feasible at ultra-high magnetic field strengths.

5. Conclusions

This study examined the relationship between neuron-axon-synapse architecture and white matter vasculature using high-resolution multimodal MRI and quantitative anatomy in primates. We found that white matter vasculature was correlated to the synaptic density of the

overlying gray matter, while myelinated tracts were characterized by sparser vascular networks. By integrating axon geometry, density, and distance from the cortical surface, a predictive model was developed to map vascularity, providing new insights into the structural and metabolic determinants of white matter vasculature.

Data and Code Availability

The data for the figures will be made available at https://balsa.wustl.edu/study/show/LPDP.

These datasets require Connectome Workbench v2.0 for visualization.

The NHP-HCP pipelines are available at https://github.com/Washington-University/NHPPipelines.

CRediT authorship contribution statement

Ikko Kimura: Conceptualization, Software, Formal analysis, Writing - original draft, Writing - review & editing, Visualization. **Takuya Hayashi:** Resources, Funding acquisition. **Joonas Autio:** Conceptualization, Methodology, Formal analysis, Writing - original draft, Writing - review & editing, Visualization, Resources, Funding acquisition.

Funding

This research is partially supported by JSPS KAKENHI Grant Number (JP20K15945, J.A.A), by the program for Brain/MINDS and Brain/MINDS-beyond from Japan Agency for Medical Research and development, AMED (JP18dm0307006, JP21dm0525006, JP19dm0307004,

JP23wm0625001, JP24wm0625205, JP24wm0625122, T.H.).

Declaration of Competing Interests

The authors declare that they have no competing interests.

Acknowledgements

References

Alexander, D.C., Hubbard, P.L., Hall, M.G., Moore, E.A., Ptito, M., Parker, G.J.M., Dyrby, T.B., 2010. Orientationally invariant indices of axon diameter and density from diffusion MRI. Neuroimage 52, 1374–1389. https://doi.org/10.1016/j.neuroimage.2010.05.043

Autio, J.A., Glasser, M.F., Ose, T., Donahue, C.J., Bastiani, M., Ohno, M., Kawabata, Y., Urushibata, Y., Murata, K., Nishigori, K., Yamaguchi, M., Hori, Y., Yoshida, A., Go, Y., Coalson, T.S., Jbabdi, S., Sotiropoulos, S.N., Kennedy, H., Smith, S., Van Essen, D.C., Hayashi, T., 2020. Towards HCP-Style macaque connectomes: 24-Channel 3T multi-array coil, MRI sequences and preprocessing. Neuroimage 215, 116800. https://doi.org/10.1016/j.neuroimage.2020.116800

Autio, J.A., Kimura, I., Ose, T., Matsumoto, Y., Ohno, M., Urushibata, Y., Ikeda, T., Glasser, M.F., Van Essen, D.C., Hayashi, T., 2024a. Mapping vascular network architecture in primate brain using ferumoxytol-weighted laminar MRI. https://doi.org/10.7554/elife.99940.1

Autio, J.A., Roberts, R.E., 2014. Interpreting functional diffusion tensor imaging. Front. Neurosci. 8, 68. https://doi.org/10.3389/fnins.2014.00068

Autio, J.A., Uematsu, A., Ikeda, T., Ose, T., Hou, Y., Magrou, L., Kimura, I., Ohno, M., Murata, K., Coalson, T., Kennedy, H., Glasser, M.F., Van Essen, D.C., Hayashi, T., 2024b. Charting cortical-layer specific area boundaries using Gibbs ringing attenuated T1w/T2w-FLAIR myelin MRI. bioRxiv. https://doi.org/10.1101/2024.09.27.615294

Basser, P.J., Mattiello, J., LeBihan, D., 1994. MR diffusion tensor spectroscopy and imaging. Biophys. J. 66, 259–267. https://doi.org/10.1016/S0006-3495(94)80775-1

Bernier, M., Chen, J.E., Ohringer, N., Fultz, N.E., Leaf, R.K., Viessmann, O., Lewis, L.D., Wald, L.L., Polimeni, J.R., 2019. Prospects for improving neuronal specificity of fMRI with ferumoxytol: an evaluation of vascular segmentation and cortical depth-dependent analysis, in: Proc Intl Soc Mag Reson Med. archive.ismrm.org, p. 537.

Bernstein, B.W., Bamburg, J.R., 2003. Actin-ATP hydrolysis is a major energy drain for neurons. J. Neurosci. 23, 1–6. https://doi.org/10.1523/JNEUROSCI.23-01-00002.2003

Biswas, S., Cottarelli, A., Agalliu, D., 2020. Neuronal and glial regulation of CNS angiogenesis and barriergenesis. Development 147. https://doi.org/10.1242/dev.182279

Bok, S.T., 1929. Der Einflu¥ der in den Furchen und Windungen auftretenden Krümmungen der Gro¥hirnrinde auf die Rindenarchitektur. Arch. Psychiatr. Nervenkr. Z. Gesamte Neurol. Psychiatr. 121, 682–750. https://doi.org/10.1007/BF02864437

Borowsky, I.W., Collins, R.C., 1989. Metabolic anatomy of brain: a comparison of regional capillary density, glucose metabolism, and enzyme activities. J. Comp. Neurol. 288, 401–413. https://doi.org/10.1002/cne.902880304

Boxerman, J.L., Hamberg, L.M., Rosen, B.R., Weisskoff, R.M., 1995. MR contrast due to intravascular magnetic susceptibility perturbations. Magn. Reson. Med. 34, 555–566. https://doi.org/10.1002/mrm.1910340412

Caporale, N., Dan, Y., 2008. Spike Timing–Dependent Plasticity: A Hebbian Learning Rule. https://doi.org/10.1146/annurev.neuro.31.060407.125639

Carmeliet, P., Tessier-Lavigne, M., 2005. Common mechanisms of nerve and blood vessel wiring. Nature 436, 193–200. https://doi.org/10.1038/nature03875

Chen, J.J., Rosas, H.D., Salat, D.H., 2013. The relationship between cortical blood flow and sub-cortical white-matter health across the adult age span. PLoS One 8, e56733.

https://doi.org/10.1371/journal.pone.0056733

Chojdak-Łukasiewicz, J., Dziadkowiak, E., Zimny, A., Paradowski, B., 2021. Cerebral small vessel disease: A review. Adv. Clin. Exp. Med. 30, 349–356. https://doi.org/10.17219/acem/131216

Collins, C.E., Airey, D.C., Young, N.A., Leitch, D.B., Kaas, J.H., 2010. Neuron densities vary across and within cortical areas in primates. Proceedings of the National Academy of Sciences 107, 15927–15932. https://doi.org/10.1073/pnas.1010356107

Daniel, J.L., Molish, I.R., Robkin, L., Holmsen, H., 1986. Nucleotide exchange between cytosolic ATP and F-actin-bound ADP may be a major energy-utilizing process in unstimulated platelets. Eur. J. Biochem. 156, 677–684. https://doi.org/10.1111/j.1432-1033.1986.tb09631.x

Donahue, C.J., Sotiropoulos, S.N., Jbabdi, S., Hernandez-Fernandez, M., Behrens, T.E., Dyrby, T.B., Coalson, T., Kennedy, H., Knoblauch, K., Van Essen, D.C., Glasser, M.F., 2016. Using Diffusion Tractography to Predict Cortical Connection Strength and Distance: A Quantitative Comparison with Tracers in the Monkey. J. Neurosci. 36, 6758–6770. https://doi.org/10.1523/JNEUROSCI.0493-16.2016

Duvernoy, H.M., Delon, S., Vannson, J.L., 1981. Cortical blood vessels of the human brain. Brain Res. Bull. 7, 519–579. https://doi.org/10.1016/0361-9230(81)90007-1

Engl, E., Attwell, D., 2015. Non-signalling energy use in the brain: Non-signalling energy use in the brain. J. Physiol. 593, 3417–3429. https://doi.org/10.1113/jphysiol.2014.282517

Fath, K.R., Lasek, R.J., 1988. Two classes of actin microfilaments are associated with the inner cytoskeleton of axons. J. Cell Biol. 107, 613–621. https://doi.org/10.1083/jcb.107.2.613

Frangi, A.F., Niessen, W.J., Vincken, K.L., Viergever, M.A., 1998. Multiscale vessel enhancement filtering, in: Medical Image Computing and Computer-Assisted Intervention — MICCAI'98, Lecture Notes in Computer Science. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 130–137. https://doi.org/10.1007/bfb0056195

Froudist-Walsh, S., Xu, T., Niu, M., Rapan, L., Zhao, L., Margulies, D.S., Zilles, K., Wang, X.-J., Palomero-Gallagher, N., 2023. Gradients of neurotransmitter receptor expression in the macaque cortex. Nat. Neurosci. 26, 1281–1294. https://doi.org/10.1038/s41593-023-01351-2

Gao, L., Pan, X., Zhang, J.H., Xia, Y., 2023. Glial cells: an important switch for the vascular

function of the central nervous system. Front. Cell. Neurosci. 17, 1166770. https://doi.org/10.3389/fncel.2023.1166770

Gibson, E.M., Purger, D., Mount, C.W., Goldstein, A.K., Lin, G.L., Wood, L.S., Inema, I., Miller, S.E., Bieri, G., Zuchero, J.B., Barres, B.A., Woo, P.J., Vogel, H., Monje, M., 2014. Neuronal activity promotes oligodendrogenesis and adaptive myelination in the mammalian brain. Science 344, 1252304. https://doi.org/10.1126/science.1252304

Glasser, M.F., Goyal, M.S., Preuss, T.M., Raichle, M.E., Van Essen, D.C., 2014. Trends and properties of human cerebral cortex: correlations with cortical myelin content. Neuroimage 93 Pt 2, 165–175. https://doi.org/10.1016/j.neuroimage.2013.03.060

Glasser, M.F., Sotiropoulos, S.N., Wilson, J.A., Coalson, T.S., Fischl, B., Andersson, J.L., Xu, J., Jbabdi, S., Webster, M., Polimeni, J.R., Van Essen, D.C., Jenkinson, M., WU-Minn HCP Consortium, 2013. The minimal preprocessing pipelines for the Human Connectome Project. Neuroimage 80, 105–124. https://doi.org/10.1016/j.neuroimage.2013.04.127

Goldstein, A.Y.N., Wang, X., Schwarz, T.L., 2008. Axonal transport and the delivery of pre-synaptic components. Curr. Opin. Neurobiol. 18, 495–503. https://doi.org/10.1016/j.conb.2008.10.003

Harris, J.J., Attwell, D., 2012. The energetics of CNS white matter. J. Neurosci. 32, 356–371. https://doi.org/10.1523/JNEUROSCI.3430-11.2012

Hase, Y., Ding, R., Harrison, G., Hawthorne, E., King, A., Gettings, S., Platten, C., Stevenson, W., Craggs, L.J.L., Kalaria, R.N., 2019. White matter capillaries in vascular and neurodegenerative dementias. Acta Neuropathol. Commun. 7, 16. https://doi.org/10.1186/s40478-019-0666-x

Hayashi, T., Hou, Y., Glasser, M.F., Autio, J.A., Knoblauch, K., Inoue-Murayama, M., Coalson, T., Yacoub, E., Smith, S., Kennedy, H., Van Essen, D.C., 2021. The nonhuman primate neuroimaging and neuroanatomy project. Neuroimage 229, 117726. https://doi.org/10.1016/j.neuroimage.2021.117726

Hernández-Torres, E., Kassner, N., Forkert, N.D., Wei, L., Wiggermann, V., Daemen, M., Machan, L., Traboulsee, A., Li, D., Rauscher, A., 2017. Anisotropic cerebral vascular architecture causes orientation dependency in cerebral blood flow and volume measured with dynamic susceptibility contrast magnetic resonance imaging. J. Cereb. Blood Flow Metab. 37, 1108–1119. https://doi.org/10.1177/0271678X16653134

Huang, Y., Wei, P.-H., Xu, L., Chen, D., Yang, Y., Song, W., Yi, Y., Jia, X., Wu, G., Fan, Q., Cui, Z., Zhao, G., 2023. Intracranial electrophysiological and structural basis of BOLD functional connectivity in human brain white matter. Nat. Commun. 14, 3414. https://doi.org/10.1038/s41467-023-39067-3

Huang, Y., Yang, Y., Hao, L., Hu, X., Wang, P., Ding, Z., Gao, J.-H., Gore, J.C., 2020. Detection of functional networks within white matter using independent component analysis. Neuroimage 222, 117278. https://doi.org/10.1016/j.neuroimage.2020.117278

Innocenti, G.M., Vercelli, A., Caminiti, R., 2014. The diameter of cortical axons depends both on the area of origin and target. Cereb. Cortex 24, 2178–2188. https://doi.org/10.1093/cercor/bht070

Kageyama, G.H., Wong-Riley, M., 1986. Laminar and cellular localization of cytochrome oxidase in the cat striate cortex. J. Comp. Neurol. 245, 137–159. https://doi.org/10.1002/cne.902450202

Kageyama, G.H., Wong-Riley, M.T., 1984. The histochemical localization of cytochrome oxidase in the retina and lateral geniculate nucleus of the ferret, cat, and monkey, with particular reference to retinal mosaics and ON/OFF-center visual channels. J. Neurosci. 4, 2445–2459. https://doi.org/10.1523/jneurosci.04-10-02445.1984

Kim, S.-G., Harel, N., Jin, T., Kim, T., Lee, P., Zhao, F., 2013. Cerebral blood volume MRI with intravascular superparamagnetic iron oxide nanoparticles. NMR Biomed. 26, 949–962. https://doi.org/10.1002/nbm.2885

Kimura, I., Hayashi, M.J., Amano, K., 2024. Immediate effect of quadri-pulse stimulation on human brain microstructures and functions. Imaging Neuroscience 2, 1–15. https://doi.org/10.1162/imag_a_00264

Kirilina, E., Helbling, S., Morawski, M., Pine, K., Reimann, K., Jankuhn, S., Dinse, J., Deistung, A., Reichenbach, J.R., Trampel, R., Geyer, S., Müller, L., Jakubowski, N., Arendt, T., Bazin, P.-L., Weiskopf, N., 2020. Superficial white matter imaging: Contrast mechanisms and whole-brain in vivo mapping. Sci Adv 6. https://doi.org/10.1126/sciadv.aaz9281

Lamsa, K.P., Kullmann, D.M., Woodin, M.A., 2010. Spike-timing dependent plasticity in inhibitory circuits. Front. Synaptic Neurosci. 2, 8. https://doi.org/10.3389/fnsyn.2010.00008

Lazari, A., Salvan, P., Cottaar, M., Papp, D., Rushworth, M.F.S., Johansen-Berg, H., 2022.

Hebbian activity-dependent plasticity in white matter. Cell Rep. 39, 110951. https://doi.org/10.1016/j.celrep.2022.110951

Markov, N.T., Ercsey-Ravasz, M.M., Ribeiro Gomes, A.R., Lamy, C., Magrou, L., Vezoli, J., Misery, P., Falchier, A., Quilodran, R., Gariel, M.A., Sallet, J., Gamanut, R., Huissoud, C., Clavagnier, S., Giroud, P., Sappey-Marinier, D., Barone, P., Dehay, C., Toroczkai, Z., Knoblauch, K., Van Essen, D.C., Kennedy, H., 2014. A weighted and directed interareal connectivity matrix for macaque cerebral cortex. Cereb. Cortex 24, 17–36. https://doi.org/10.1093/cercor/bhs270

Ogawa, S., Menon, R.S., Tank, D.W., Kim, S.G., Merkle, H., Ellermann, J.M., Ugurbil, K., 1993. Functional brain mapping by blood oxygenation level-dependent contrast magnetic resonance imaging. A comparison of signal characteristics with a biophysical model. Biophys. J. 64, 803–812. https://doi.org/10.1016/S0006-3495(93)81441-3

Perge, J.A., Niven, J.E., Mugnaini, E., Balasubramanian, V., Sterling, P., 2012. Why do axons differ in caliber? J. Neurosci. 32, 626–638. https://doi.org/10.1523/JNEUROSCI.4254-11.2012

Saab, A.S., Tzvetanova, I.D., Nave, K.-A., 2013. The role of myelin and oligodendrocytes in axonal energy metabolism. Curr. Opin. Neurobiol. 23, 1065–1072. https://doi.org/10.1016/j.conb.2013.09.008

Sampaio-Baptista, C., Johansen-Berg, H., 2017. White Matter Plasticity in the Adult Brain. Neuron 96, 1239–1251. https://doi.org/10.1016/j.neuron.2017.11.026

Sampaio-Baptista, C., Neyedli, H.F., Sanders, Z.-B., Diosi, K., Havard, D., Huang, Y., Andersson, J.L.R., Lühr, M., Goebel, R., Johansen-Berg, H., 2021. fMRI neurofeedback in the motor system elicits bidirectional changes in activity and in white matter structure in the adult human brain. Cell Rep. 37, 109890. https://doi.org/10.1016/j.celrep.2021.109890

Sandrone, S., Aiello, M., Cavaliere, C., Thiebaut de Schotten, M., Reimann, K., Troakes, C., Bodi, I., Lacerda, L., Monti, S., Murphy, D., Geyer, S., Catani, M., Dell'Acqua, F., 2023. Mapping myelin in white matter with T1-weighted/T2-weighted maps: discrepancy with histology and other myelin MRI measures. Brain Struct. Funct. 228, 525–535. https://doi.org/10.1007/s00429-022-02600-z

Schilling, K.G., Li, M., Rheault, F., Gao, Y., Cai, L., Zhao, Y., Xu, L., Ding, Z., Anderson,

A.W., Landman, B.A., Gore, J.C., 2023. Whole-brain, gray, and white matter time-locked functional signal changes with simple tasks and model-free analysis. Proc. Natl. Acad. Sci. U. S. A. 120, e2219666120. https://doi.org/10.1073/pnas.2219666120

Shu, Y., Hasenstaub, A., McCormick, D.A., 2003. Turning on and off recurrent balanced cortical activity. Nature 423, 288–293. https://doi.org/10.1038/nature01616

Smirnov, M., Destrieux, C., Maldonado, I.L., 2021. Cerebral white matter vasculature: still uncharted? Brain 144, 3561–3575. https://doi.org/10.1093/brain/awab273

Tabelow, K., Balteau, E., Ashburner, J., Callaghan, M.F., Draganski, B., Helms, G., Kherif, F., Leutritz, T., Lutti, A., Phillips, C., Reimer, E., Ruthotto, L., Seif, M., Weiskopf, N., Ziegler, G., Mohammadi, S., 2019. hMRI - A toolbox for quantitative MRI in neuroscience and clinical research. Neuroimage 194, 191–210. https://doi.org/10.1016/j.neuroimage.2019.01.029

Tao, K., Matsuki, N., Koyama, R., 2014. AMP-activated protein kinase mediates activity-dependent axon branching by recruiting mitochondria to axon: Role of Mitochondria in Axonal Morphogenesis. Dev. Neurobiol. 74, 557–573. https://doi.org/10.1002/dneu.22149

Tournier, J.-D., Smith, R., Raffelt, D., Tabbara, R., Dhollander, T., Pietsch, M., Christiaens, D., Jeurissen, B., Yeh, C.-H., Connelly, A., 2019. MRtrix3: A fast, flexible and open software framework for medical image processing and visualisation. Neuroimage 202, 116137. https://doi.org/10.1016/j.neuroimage.2019.116137

Van Essen, D.C., Maunsell, J.H., 1980. Two-dimensional maps of the cerebral cortex. J. Comp. Neurol. 191, 255–281. https://doi.org/10.1002/cne.901910208

Vezoli, J., Vinck, M., Bosman, C.A., Bastos, A.M., Lewis, C.M., Kennedy, H., Fries, P., 2021. Brain rhythms define distinct interaction networks with differential dependence on anatomy. Neuron 109, 3862-3878.e5. https://doi.org/10.1016/j.neuron.2021.09.052

Viessmann, O., Tian, Q., Bernier, M., Polimeni, J.R., 2022. Static and dynamic BOLD fMRI components along white matter fibre tracts and their dependence on the orientation of the local diffusion tensor axis relative to the B0-field. J. Cereb. Blood Flow Metab. 42, 1905–1919. https://doi.org/10.1177/0271678X221106277

Wang, H., Wang, X., Wang, Y., Zhang, D., Yang, Y., Zhou, Y., Qiu, B., Zhang, P., 2023. White matter BOLD signals at 7 Tesla reveal visual field maps in optic radiation and vertical occipital fasciculus. Neuroimage 269, 119916.

https://doi.org/10.1016/j.neuroimage.2023.119916

Weber, B., 2002. White matter glucose metabolism during intracortical electrostimulation: A quantitative [18F]fluorodeoxyglucose autoradiography study in the rat. Neuroimage 16, 993–998. https://doi.org/10.1006/nimg.2002.1104

Weber, B., Keller, A.L., Reichold, J., Logothetis, N.K., 2008. The microvascular system of the striate and extrastriate visual cortex of the macaque. Cereb. Cortex 18, 2318–2330. https://doi.org/10.1093/cercor/bhm259

Wiser, A.K., Callaway, E.M., 1996. Contributions of individual layer 6 pyramidal neurons to local circuitry in macaque primary visual cortex. J. Neurosci. 16, 2724–2739. https://doi.org/10.1523/jneurosci.16-08-02724.1996

Wu, W.-C., Lin, S.-C., Wang, D.J., Chen, K.-L., Li, Y.-D., 2013. Measurement of cerebral white matter perfusion using pseudocontinuous arterial spin labeling 3T magnetic resonance imaging--an experimental and theoretical investigation of feasibility. PLoS One 8, e82679. https://doi.org/10.1371/journal.pone.0082679

Yeatman, J.D., Dougherty, R.F., Myall, N.J., Wandell, B.A., Feldman, H.M., 2012. Tract profiles of white matter properties: automating fiber-tract quantification. PLoS One 7, e49790. https://doi.org/10.1371/journal.pone.0049790

Zhang, H., Schneider, T., Wheeler-Kingshott, C.A., Alexander, D.C., 2012. NODDI: Practical in vivo neurite orientation dispersion and density imaging of the human brain. Neuroimage 61, 1000–1016. https://doi.org/10.1016/j.neuroimage.2012.03.072