Integrating ethnolinguistic and archaeobotanical data to uncover the

origin and dispersal of cultivated sorghum in Africa: a genomic

perspective

Aude Gilabert[1,2*], Monique Deu[1,2#], Louis Champion[3#], Philippe Cubry[3#], Armel Donkpegan[1,2‡],

Jean-François Rami[1,2], David Pot[1,2], Yves Vigouroux[3], Christian Leclerc[1,2]

[1] CIRAD, UMR AGAP Institut, F-34398 Montpellier, France

[2] UMR AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, Montpellier, France

[3] UMR DIADE, Univ Montpellier, IRD, CIRAD, Montpellier, France

[#] Those authors contributed equally to the work

[‡] Present address: SYSAAF-Centre INRAE Val de Loire, UMR BOA, 37380 Nouzilly, France.

*Corresponding author: aude.gilabert@cirad.fr

**ABSTRACT**

Archaeobotanical evidence suggests that the beginning of cultivation and emergence of

domesticated sorghum was located in eastern Sudan during the fourth millennium BCE. Here, we

used a genomic approach, together with archaeobotanical and ethnolinguistic data, to refine the

spatial and temporal origin and the spread of cultivated sorghum in Africa.

We built a probability map of the origin of sorghum domestication in Eastern Africa using genomic

data and spatial Bayesian models. The origin was located in Eastern Sudan and Western Ethiopia,

in perfect concordance with recent archaeobotanical evidence. Calibrated on archaeological

remains, our genomic-based model suggests that the beginning of the expansion of sorghum

agriculture took place around 4,600 years ago.

Spread of sorghum cultivation led to a sorghum population structure fitting ethnolinguistic groups

at different scales, suggesting that human social groups and sorghum populations co-diffused.

Consequently, ethnolinguistic barriers and social preferences, as well as adaptation to specific

climate zones, have contributed to structuring domesticated sorghum diversity during its diffusion.

**Keywords:** Domestication, sorghum origin and diffusion, population genomics, archaeobotany,

ethnolinguistics.

**Introduction**

Sorghum is one of the most climate-resilient and drought-tolerant cereals. It is the fifth most widely

cultivated cereal globally (FAO, 2023) and is the mainstay for millions of inhabitants in the drier

areas of Africa and Asia. Sorghum has a multitude of diversified end uses, including food, feed,

fodder, fiber, and fuel in over 100 countries in a variety of environments.

Anthropological evidence suggests that hunter-gatherers were familiar with wild forms of sorghum

as early as 8,000 BCE (Smith and Frederiksen, 2000). The beginning of cultivation and emergence

of sorghum domesticated characteristics was situated during the fourth millennium BCE in eastern

Sudan (Winchell et al., 2017), but no genetic evidence has supported that hypothesis so far, and

there are still gaps in our understanding of its early diffusion.

Anthropological and human genetics studies suggest that many cultural and linguistic traits,

including agricultural expansion (Diamond and Bellwood, 2003), followed demic diffusion models,

i.e. the spread occurred through the movements of people (Hewlett et al., 2002), an hypothesis

known for the agricultural diffusion as the "farming/language dispersal hypothesis". Assuming that

the initial sorghum populations were propagated under a demic diffusion model through the

movements of people from the crop's area of origin, and were subsequently maintained through

time, it can be hypothesized that sorghum genetic groups should be associated with fine linguistic

clusters at their destination.

Many anthropological studies suggest that seed exchanges and seed circulation are related to

farmers' linguistic identity (Leclerc and Coppens d'Eeckenbrugge, 2012; Westengen et al., 2014).

As noted by Harlan & Stemler (1976), the correspondence between the distribution of the basic

morphological groups of sorghum and the major linguistic groups in Africa may not be fortuitous.

For example, guinea sorghum is associated with Western African farmers, durra sorghum is

confined to the Muslim and Arabic farmers along the fringes of the Sahara and Ethiopia (Kimber,

2000), and kafir, which is widely grown in Southern Africa, is reputed to be associated with the

Bantu-speaking peoples of Eastern and Southern Africa (de Wet and Huckabay, 1967). If such

social barriers were at work in the past and maintained, at least partially, up to present times, they

can explain why, after thousands of years, sorghum genetic groups still remain associated today

with fine linguistic clusters.

The aim of this paper is to shed light on the origin and spread of the cultivated sorghum in Africa

by integrating ethnolinguistic, archaeobotanical and genomic data. Based on a multidisciplinary

approach and on spatially explicit modeling, we seek to identify the geographic origin of cultivated

sorghum, and enlighten its expansion history and dynamics in Africa. The origin and spread of

cultivated sorghum were considered to result from both biological and social processes, with two

leading questions: firstly, how human social groups could have contributed to the dissemination of

initial sorghum populations, and secondly, how linguistic barriers could have subsequently

contributed to the isolation of sorghum populations from each other and to the building of its current

genetic structure.

To examine the origin and dissemination of African sorghum, we initially investigated the

relationships between sorghum populations and ethnolinguistic groups at two classification levels.

Subsequently, we assessed the congruence between our genomic-based estimation of the

geographic origin of cultivated sorghum and the most recent archaeobotanical findings. By

integrating genomic and archaeobotanical data, we finally inferred the timing and rate of the

westward and southward expansions from its origin.

## Materials and methods

**Plant material**

We used a set of cultivated sorghum accessions [*Sorghum bicolor* (L.) Moench] included in several

genotype-by-sequencing (GBS) datasets previously published in Brenton et al. (2016), Lasky et al.

(2015) and Yu et al. (2016). From these datasets combining over 2,000 landraces, we selected 210

cultivated varieties (Supplementary Dataset 1 available on the CIRAD Dataverse repository; see

the section "Data and code availability") with the aim of 1) keeping only African geo-referenced

accessions, 2) covering the African area of distribution of sorghum and 3) limiting over-

representation of any geographic, ethnolinguistic and genetic group. To proceed to that selection,

we first excluded the non-African landraces from the complete datasets. Then, following the

procedure described in the following section "Genomic dataset", we produced a VCF file containing

the African landraces from the Brenton, Morris-Lasky and Yu collections and performed an

individual-based clustering analysis implemented in STRUCTURE (Pritchard et al., 2000) and a

Neighbor Joining tree based on that African dataset, to remove nearest neighbor accessions while

maximizing geographic distance and ethnolinguistic data (results not shown). Latitude and

longitude coordinates associated with each selected accession were checked using passport data

and place names from different databases (genesis, available at https://www.genesys-

pgr.org/a/overview, grin global available at https://www.grin-

global.org/help_pw2/Content/Searching%20Overview.htm, and geonames available at

https://www.geonames.org/). This sampling included representation of all five known cultivated

sorghum botanical groups of cultivated sorghum (*bicolor, caudatum, durra, guinea, and kafir*) and

their intermediate races.

**Genomic dataset**

The GBS data of cultivated sorghum from Brenton, Morris-Lasky and Yu collections were

downloaded from NCBI Sequence Read Archive (Leinonen et al., 2011) (SRA) (see Supplementary

Dataset 1). The SRA files were converted to FASTQ file using fastq-dump of SRA Toolkit

(http://www.ncbi.nlm.nih.gov/Traces/sra/sra.cgi?view=toolkit_doc&f=fastq-dump).

Low quality bases, Illumina adapters, primers and the end of read were removed with

TRIMMOMATIC version 0.33 (Bolger et al., 2014) with the following options: ILLUMINACLIP

2:30:10, LEADING 3, TRAILING 3, SLIDINGWINDOW 4:15, MINLEN 36.

All single-end reads were aligned to the BTx623 sorghum reference genome (McCormick et al.,

2018; release 3.1 of the Phytozome database) with the Burrow-Wheeler Aligner (BWA) software

v0.7.15 using the standard MEM option (Li and Durbin, 2009). The unmapped, not properly mapped

and duplicated reads were filtered with the SAMtools v0.1.9 software (Li et al., 2009). The resulting

BAM files (binary format of sequence data) were used as input for Genome Analysis Toolkit (GATK)

v4.1.4.0 (DePristo et al., 2011). The GATK HaplotypeCaller algorithm was used for joint variant

discovery and genotyping. Single-sample gVCFs were created using the GATK HaplotypeCaller

standard parameters according to GATK Best Practices recommendations (DePristo et al., 2011).

Finally, single-sample gVCF files were merged into one VCF file using the GenomicsDBImport tool

from GATK followed by GenotypeGVCFs for joint variant calling.

Raw VCF was filtered with VCFtools v0.1.13 (Danecek et al., 2011) to obtain high-quality SNPs

according to the following criteria: only bi-allelic sites, missing data < 40%, minQ (sites with Quality

value above) > 60. After applying all filters, we obtained a VCF file containing 95,362 SNPs for the

210 accessions analyzed.

**Data analysis**

*Population structure and diversity*

To identify population structure, we used three distinct and complementary approaches: (i) a

maximum likelihood estimation of individual ancestries approach based on a population genetic

model as implemented in the software ADMIXTURE (Alexander et al., 2009) (ii) a principal

component analysis (PCA) approach; and (iii) an unweighted neighbour-joining (NJ) approach

based on pairwise genetic distances. These analyses were run on the dataset of 210 accessions

and 95,362 SNPs. ADMIXTURE v.1.22 software (Alexander et al., 2009) was used to estimate the

proportion of ancestry of each individual sample in each of the $K$ considered genetic clusters, $K$

ranging from 1 to 20. Ten repetitions of the algorithm were made for each $K$. We used the cross-

validation procedure implemented in ADMIXTURE (5-fold cross-validation) to choose the value of

$K$ that best represents our sampling. We selected as the optimal number of genetic clusters the

value of $K$ that gave the lowest cross-validation errors averaged over the 10 replicates. We

performed the PCA using PLINK version 1.9 (Chang et al., 2015; Purcell et al., 2007). Diversity tree

was constructed by using the unweighted neighbour-joining (NJ) algorithm (Saitou and Nei, 1987)

based on a dissimilarity matrix computed using GBS-div software (available at

https://darwin.cirad.fr/gbsdiv.php): dissimilarity indices $d_{ij}$ were estimated by simple matching, that

is the number of unmatching alleles between two accessions standardized by the number of loci

(Perrier and Jacquemoud-Collet, 2019).

We estimated admixture graphs of geographically defined genetic groups of cultivated sorghum

resulting from Admixture analysis using TreeMix software v1.13 (Pickrell and Pritchard, 2012)

TreeMix builds a maximum likelihood tree for the relationships among populations and allows the

addition of migration events to improve model fit. It allows inference of contemporary and historical

patterns of gene flow among populations. We used the script provided by Fitak (2021) to run

TreeMix for a number of migration events, $m$. We tested $m$ ranging from 0 to 10 with 30 iterations

for each value of $m$. We also considered different sizes of SNP-blocks ranging from 50 to 500 with

an incrementation of 50, to avoid converging to the same likelihood across all runs for each iteration

of $m$, as suggested (Fitak, 2021). The optimal number of migration events was then identified using

the Evanno's method (Evanno et al., 2005) implemented in the R package optM (Fitak, 2021) The

tree inferred with the optimal number of migration events was visualized using the R function

provided with the Treemix software (Pickrell and Pritchard, 2012). Because of its divergence and

its likely independent origin (Mace et al., 2013; Sagnard et al., 2011), we excluded the cluster cl5c

for the Treemix analysis.

In order to characterize the level of diversity within cultivated sorghum, we computed the observed

heterozygosity ($H_O$) and gene diversity ($H_S$) per genetic cluster using the R package hierfstat

(Goudet, 2005). We calculated the number of singletons and private doubletons per accession

using VCFtools 0.1.16 (Danecek et al., 2011) and the number of private alleles ($N$) using an in-

house script (available on the CIRAD gitlab platform; see the section "Data and code availability").

The genetic differentiation between the genetic clusters was estimated by the means of pairwise

Weir & Cockerham $F_{ST}$ using the R package hierfstat (Goudet, 2005). All those statistics were

computed using the whole dataset of 210 accessions. For all the subsequent analyses that are

based on the genetic clusters, we only kept accessions that were unambiguously assigned to a

genetic cluster using the same ADMIXTURE ancestry threshold of 0.7.

To analyze the distribution of the genetic clusters according to the climate and to the language

families, we used the Köppen-Geiger climate classification, which defines climate zones with

different vegetation growth according to the temperature and/or the dryness (Beck et al., 2018),

and the language classification referenced in the database Ethnologue Version 16 (Lewis, 2009).

Regarding the climate, our sampling is distributed in ten of the sixteen zones present in Africa, with

six climate zones being represented by more than five accessions. For the ethnolinguistic analysis,

we considered the three main language families present in Africa: the Afro-Asiatic one, distributed

in Africa in the western and eastern parts of the Sahelian belt and in the Horn of Africa, the Nilo-

Saharan language, whose distribution mostly extends from the central part of the Sahara and

Sahelian belt to northern limit of the Great Lakes region, and the Niger-Congo language family,

widely distributed from the Sahelian belt to South Africa. We further subdivided the Niger-Congo

family into two classes, the Mande distributed in the western part of western Africa and the Atlantic-

Congo languages, which include the Bantoid languages notably, since this level of classification

has been shown to be relevant, at least for the fonio in Western Africa (Diop et al., 2023). We

further subdivided the Nilo-Saharan family into three classes (Central and Eastern Sudanic and

Fur), the Afro-Asiatic family into four classes (the Chadic, the Cushitic, the Omotic and the Semitic)

and the Atlantic-Congo class into 16 classes to consider the Guthrie classification of the Bantoid

languages (Guthrie, 1971, 1948; Maho, 2009).

The association between genetic clusters and botanical races, climate zones or ethnolinguistic

groups were tested using Fisher's exact tests with the R function fisher.test. The associations were

further analyzed by visualizing the Peason residuals to identify the specific groups contributing to

the observed associations. Pearson residuals were retrieved from the R function chisq.test. Cells

with residuals greater than |2| were considered as deviating from the expected values. Kruskal-

Wallis rank sum tests (using the R function kruskal.test) were used to test whether the altitude or

the number of singletons per accession differed between the genetic clusters. Multiple groups

comparisons were performed using Dunn's tests from the R package dunn.test (Dinno, 2024) to

identify the genetic clusters that are differing according to the altitude or the number of singletons

per accession. All R analyses were performed using R Statistical Software version 4.4.0 (R Core

Team, 2024).

***Inference of the geographic origin of the cultivated sorghum***

We used SPLATCHE2 (Currat et al., 2004) to perform spatially-explicit simulations within an

approximate Bayesian framework in order to infer the geographic origin of the diffusion of the

cultivated sorghum in Africa.

SPLATCHE2 enables users to model range expansions using non-equilibrium stepping-stone

models. In those models, the geographical cells are colonized from their neighbors. We considered

here a single origin model. We performed 1,000,000 range expansion simulations from randomly

picked origins based on prior distributions for latitudinal and longitudinal coordinates, considering

a heterogeneous environment and modeling the African continent using an array of 87 by 83

demes. Prior distributions allowed the geographic coordinates of the origin of expansion to vary

over the Sub-saharan region, within an area ranging between -17° and 50° of longitude and

between -35° and 20° of latitude. Despite the importance of gene flow and introgression in the

evolutionary history of sorghum, we decided to consider simple demographic models, to limit the

model complexity and the number of parameters in the model. Because domestication can be

accompanied by a genetic bottleneck, followed by a period of stable low population size before

expansion due to a period of weak cultivation, we considered two different simple demographic

models, one with a constant size of population before the expansion and one with a bottleneck

before the expansion (*Supplementary Information*, Fig. S1). Uninformative prior distributions were

considered for the longitude, latitude, migration rate, the growth rate, the expansion length

(Exp_length), the effective population size before expansion (BtSize, corresponding to population

size during the bottleneck), the population size before the bottleneck (AncSize) and the duration of

the bottleneck in generations (BtDuration) (*Supplementary Information,* Table S1). Simulated

genetic variation was surveyed at 210 geographic sites corresponding to the exact sampling

locations of cultivated sorghum accessions. A total of 10,000 haploid loci were simulated for each

genotype, and an effective mutation rate of $10^{-6}$ per base pair per generation was considered.

Missing data were randomly included in the simulated dataset at the same frequency as the

frequency of missing data in the observed dataset (approx. 0.20). The posterior distributions of the

geographic and demographic parameters were inferred using an approximate Bayesian

computation (ABC) approach relying on a neural network algorithm for posteriors estimation (R

package abc (Csilléry et al.*,* 2012)), considering a tolerance rate of 0.5% and implementing 500

neural networks. Parameter estimations were thus performed accepting 5,000 simulations. The

summary statistics used for the ABC analysis included a statistic related to the rare variants and

the expected heterozygosity in each genetic cluster identified using the ADMIXTURE analysis, as

well as eight classes of the Site Frequency Spectrum (SFS), as described in Cubry et al. (2018).

The statistic based on the rare variants corresponds to the mean number of singletons per

accession within each genetic cluster, relative to the entire dataset. As for the SFS, the eight

classes were defined according to the number of times a SNP was detected in the dataset: one

time (singleton), two times, three times, four times, five to six times, seven to 12 times, 13 to 28

times and more than 28 times. Two analyses were performed, one considering the 210 cultivated

accessions (model including all cultivated genetic clusters) and one without cultivated accessions

that likely originated from an independent event and shows signals of introgression with wild

sorghum (Gilabert et al., 2023; Mace et al., 2013; Sagnard et al., 2011) (cluster cl5c; see

*Supplementary Information*, SI Text section 1.2). We considered for the calculation of the summary

statistics only the accessions assigned to a genetic group with an ancestry > 0.7 according to

ADMIXTURE analysis. We evaluated the adjustment of our models using the goodness-of-fit

approach described in Lemaire et al. (2016) which evaluates the distance between the simulated

summary statistics and the observed ones. We used the gfit function of the abc R package (Csilléry

et al., 2012) and the mean as the goodness-of-fit statistic to perform the prior predictive checks.

We simulated 1,000 additional datasets, sampling the parameters from the posterior distributions

obtained from the previous simulations, to compute the 95% credible intervals of the summary

statistics and perform the posterior predictive checks.

***Estimating the timing of the expansion onset***

The availability of archaeological data for sorghum allowed us to calibrate temporally our diffusion

model following Burgarella et al. (2018) and estimate the timing of the onset of the expansion of

the cultivated sorghum in Africa. From the bibliography, we identified African archaeological sites

where archaeobotanical remains of cultivated (or pre-domesticated) sorghum could be identified

(*Supplementary Information,* Table S2-A), using for each site the oldest estimated dates to get an

estimation of the youngest bound for the beginning of the diffusion of cultivated sorghum in Africa.

We conducted 5,000 new spatial demographic simulations by randomly sampling the parameters

from the posterior distributions estimated in the previous ABC analysis, and monitored the

simulated time for the arrival of the expansion wave at each cell where cultivated sorghum were

identified by archaeobotanical data. A linear regression was fitted between the observed dates from

archaeological sites and the simulated dates at the same geographical coordinates. The intercept

of the real time axis was then estimated as the timing of the onset of the expansion. The regression

analysis was performed for each simulation, which allowed us to obtain a distribution of the timing

of the expansion's inception. Because domesticated sorghum was suggested to have dispersed

through two movements of diffusion (Fuller and Stevens, 2018), a North-South and an East-West

one, we performed two independent regression analyses. The first one considered the sites

distributed along an Eastern-Western transect from Sudan to Mali (westward diffusion), while the

second considered the sites distributed along the North-South axis (southward diffusion, see

*Supplementary Information,* Table S2-A). In addition, because the expansion from the origin toward

the North may correspond to a different movement of expansion than the one toward the South,

and since we only have one archaeological site north of the inferred point of origin, we estimated

the onset of the southward expansion removing the archaeological site from Egypt from the

latitudinal analysis (see *Supplementary Information*, Table S2-A). Similarly, we also estimated the

westward expansion by removing from the longitudinal analysis the archaeological site from

Ethiopia, the only site east of the inferred origin (see *Supplementary Information*, Table S2-A).

***Estimation of the dispersal front speed***

We followed Cobo et al. (2019) to estimate the speed of diffusion of the cultivated sorghum in Africa

by the means of time-space regressions. We used the same list of archaeological sites considered

for the estimation of the beginning of the dispersal of the cultivated sorghum (*Supplementary*

*Information*, Table S2-A). Although based on a low number of archaeological sites, we performed

two estimations, for the longitudinal and latitudinal diffusions, similarly to the estimation of the timing

of the onset of sorghum expansion. Indeed, one might expect that the front speeds may be different

for the two diffusions as the environments and climate do not present the same variability

longitudinally and latitudinally.

We fitted two linear regressions between the oldest estimated dates of the archaeological sites and

their distance to the estimated point of origin of the expansion of the cultivated sorghum. For the

point of origin of the expansion of sorghum, we used the median of the posterior distributions of the

longitude and latitude inferred using the ABC approach. Following Cobo et al. (2019), we used the

Haversine equation to compute the great-circle distance (the shortest distance between two points

on a sphere) between archaeological sites and the point of origin. We estimated the speed as the

inverse of the slope of the time-space regression and its standard error (SE) as being equal to the

standard error of the slope divided by the slope squared (Cobo et al., 2019; Fort et al., 2004;

Jerardino et al., 2014). Because they were based on a low number of sites, we computed for each

of the two regressions the 80% confidence-level interval for the speed, similarly to Jerardino et al.

(2014): slope $\pm$ $t_{1-\alpha/2,\ N-2}$ * $SE_{slope}$, with $t$ being the Student's $t$ value for the confidence level $\alpha$ (0.8

here) and $N$ the number of sites considered in the regression.

***Identification of geographic barriers***

We estimated effective migration surfaces using the software FEEMS v1.0.0 (Marcus et al., 2021)

(available at https://github.com/NovembreLab/feems) to infer geographic barriers and corridors for

the cultivated sorghum. FEEMS assumes an isolation-by-distance pattern (IBD), which stipulates

that dispersal occurs at a local scale and that two organisms will be more likely genetically close if

they are geographically close. FEEMS identifies areas where the genetic distance between two

individuals is either higher or lower than expected given the geographic distance separating them

and assuming an IBD. FEEMS was run following the pipeline provided with the publication. As for

the SPLATCHE analysis, FEEMS was run using the panel of 210 cultivated accessions from which

we removed the nine cultivated accessions of the cluster cl5c, as introgression may bias the results.

**Results**

**Sorghum population structure and diversity**

A panel of 210 georeferenced accessions representative of the African genetic diversity of

cultivated sorghum for which GBS data were available was used to study its domestication history

as a biological and social process. A total of 95,362 Single Nucleotide Polymorphisms (SNPs) was

identified after bioinformatic treatment of the raw genomic data. The genetic structure analysis of

population ancestries of our sampling revealed the existence of nine genetic clusters

(*Supplementary Information*, SI Text section 1.1, Fig. S2 and S3). The clusters, associated with the

known cultivated sorghum botanical races (Fisher's exact test *p*-value < 0.005; *Supplementary*

*Information*, Fig. S4), were geographically structured (Fig. 1A), with clusters associated with both

Köppen-Geiger agroclimatic zones (Beck et al., 2018) and language families (Lewis, 2009) (see

below for the detailed results on the association with the agro-climatic zones and language

families). One genetic cluster, associated with the caudatum botanical race, was widely distributed

(cl1), partially overlapping the other eight genetic clusters. Three genetic clusters were restricted

to Western Africa (cl5a, b and c) and were associated with the guinea botanical race. One cluster

(cl2b), associated with the kafir race, was restricted to Southern Africa. Two other clusters (cl4 and

cl6), respectively associated with the bicolor/durra-bicolor and the caudatum botanical races, were

distributed in Eastern Africa. Lastly, two clusters present in Eastern Africa were also found in

Southern (cl2a, associated with the guinea race) and Western Africa (cl3, associated with the durra

race) (Fig. 1-A, *Supplementary Information*, Fig. S4).

The genetic clusters considered from the structure analysis (*Supplementary Information*, SI Text

section 1.1, Fig. S2 and S3), from the PCA (*Supplementary Information*, SI Text section 1.2, Fig.

S5) and from the neighbor joining tree (*Supplementary Information*, Fig. S6) corresponded. The

first two axes of the PCA distinguished two distinct groups of accessions that belonged to the

Western African cluster cl5c and the Eastern African cluster cl4 (*Supplementary Information*, SI

Text section 1.2, Fig. S5-A and C) respectively. A third group, comprising accessions from the

Western African clusters cl5a and cl5b, stood apart on the third axis of the PCA (*Supplementary*

*Information*, SI Text section 1.2, Fig. S5-B and C). In the neighbor joining tree (*Supplementary*

*Information*, Fig. S6), the Westernmost cultivated sorghum from the cluster cl5c stood out as a

highly divergent and diverse clade, which likely corresponds to accessions belonging to the

margaritiferum subrace (see *Supplementary Information*, SI Text section SI Text section 1.2).

The maximum likelihood tree inferred using the Treemix analysis (Pickrell and Pritchard, 2012) and

excluding the highly divergent margaritiferum cultivated cluster cl5c, was consistent with the NJ

tree. The analysis suggested two gene flow events, one between the clusters cl1 and cl6 and one

between the clusters cl2b and cl3 (*Supplementary Information*, Fig. S7).

Two hotspots of sorghum genetic diversity were localized in Western and Eastern Africa, as

illustrated by the distribution of the singletons per accession (*Supplementary Information*, Fig. S8-

A). The margaritiferum cluster cl5c in Western Africa harbored the highest levels of genetic diversity

compared to all other clusters, as evidenced with the number of singletons per accession (global

Kruskal-Wallis rank sum test $X^2$ = 82.25, df = 9, p < 0.001, Dunn's tests *p*–values < 0.02 for every

pairwise comparison including the cluster cl5c; *Supplementary Information*, Fig. S8-A & B), and the

number of private alleles (*Supplementary Information*, Table S3). Besides the cluster cl5c, the

Eastern African cluster cl4 and the Eastern and Western African cluster cl3, which harbored similar

levels of genetic diversity (Dunn's test *p*–value for the pairwise comparison cl4-cl3 = 0.41;

*Supplementary Information*, Fig. S8-A & B; *Supplementary Information*, Table S3), contributed to

the two genetic diversity hotspots.

**A) Geographic distribution of the cultivated genetic groups**

**B) Classification of the three main African family languages and sorghum genetic clusters**

**C) Classification of the subclasses of the three main African family languages and sorghum genetic clusters**

 **Figure 1. Genetic diversity of the African cultivated sorghum.** A) Geographic distribution
of the cultivated genetic groups defined using ADMIXTURE (Alexander et al., 2009). Each
point corresponds to an accession. B) and C) Chi-squared residuals between the genetic and
the farmers language groups, considering (B) four linguistics groups (the three main language
families present in Africa, the Niger-Congo family being subdivided into two classes, the
Mande and the Atlantic-Congo) or (C) a finer linguistic classification. The size and color of the
circles are proportional to the value of the chi-squared residuals between the language
families and the genetic clusters.

Sorghum genetic clusters were significantly associated with the Köppen-Geiger agroclimatic zones

(Beck et al., 2018) (Fisher's exact test *p*-value < 0.001; *Supplementary Information*, SI Text section

1.3, Fig. S9-A and B, Table S4-A). The chi-squared residuals of the association between the

genetic clusters and the agroclimatic zones indicated that the Eastern African cluster cl4 was

overrepresented in temperate climates with warm summers and a dry season (Csb and Cwb zones)

and cluster cl3 in hot, arid desert climates (Bwh zone), while Western African clusters cl5c and cl5a

were over-represented in the tropical monsoon (Am) and savannah (Aw) climates respectively

(*Supplementary Information*, Fig. S9-B, Table S4-A). Conversely, the Eastern African cluster cl2b

was under-represented in the tropical savanna climate (Aw) but over-represented in semi-arid (Bsh

and Bsk) and Monsoon-influenced humid subtropical climates (Cwa) and cluster cl6 under-

represented in hot semi-arid climate (Bsh) but over-represented in humid tropical climate (Af zone)

(*Supplementary Information*, Fig. S9-B, Table S4-A). Beside the influence of the agroclimatic

zones, the genetic clusters were significantly associated with the elevation (Kruskal-Wallis rank

sum test, Kruskal-Wallis $X^2$ = 120.33, df = 9, p < 0.001). The Eastern African cluster cl4 was found

at the highest elevations while the three Western African clusters (cl5b, cl5c) were found at the

lowest (*Supplementary Information*, Fig. S9-C).

Association between sorghum genetic clusters and the human language families (Lewis, 2009) was

statistically significant (Fisher's exact test *p*-value < 0.001; Fig. 1-B, *Supplementary Information*, SI

Text section 1.3, Fig. S9-D, Table S4-B). The chi-squared residuals showed a clear association

between clusters cl2a and cl2b with Atlantic-Congo languages, and between clusters cl5b and cl5c

with Mande languages. Clusters cl3 and cl4 were associated with Afro-asiatic languages, and

clusters cl1 and cl6 with Nilo-saharan languages (Fig. 1-B). A more precise picture, by considering

a finer linguistic classification, revealed that cluster cl2b was strongly (Chi$^2$ residual over 9)

associated with the Narrow Bantu Guthrie linguistic zone S, which is located in South-Eastern

Africa, while the cluster cl2a was associated with four geographically close Bantu Guthrie linguistic

zones (zones F, G, N and P). The cluster cl6, still associated with Nilo-saharan languages, was

strongly associated with the Narrow Bantu Guthrie linguistic zone J located in the Great Lakes area

(*Supplementary Information*, Table S4-B; Fig. 1-C; and map in Fig. S9-D). Finally, the westernmost

clusters cl5b and cl5c were associated with the Atlantic-Congo and the Mande, the cluster cl5b

being also associated with the Volta-Congo linguistic group. The cluster cl5a was associated with

multiple Niger-Congo, non-Bantu, groups and with the Afro-Asiatic Chadic group.

**Geographic origin and spread of the cultivated sorghum**

The geographic origin of cultivated sorghum was inferred including or not the cluster cl5c. The

cluster cl5c is associated with the margaritiferum subrace (see *Supplementary Information*, SI Text

section 1.2), which most probably emerged from a secondary independent domestication event

and is likely introgressed with wild sorghum (Gilabert et al., 2023; Mace et al., 2013; Sagnard et

al., 2011). Thus, inferring the geographic origin of cultivated sorghum with this cluster could

introduce a bias that we wanted to account for (see details in the *Supplementary Information*, SI

Text section 2.1 and Fig. S10).

The posterior predictive checks indicate that the model with a constant ancestral population size

simulated well and better the observed data than a model with a bottleneck of the ancestral

population (*Supplementary Information*, Fig. S10-IA and Fig.S10-IIA). The simplest model with a

constant ancestral population size was therefore considered. Graphic representations of the

summary statistics (PCA and their distribution) of this model revealed that the posterior distributions

were close to the observed dataset, and most of the summary statistics (15 statistics over the 24)

were found in the 95% credible intervals of their posterior predictive distributions. Predictive checks

are only slightly significant for this no bottleneck model, and more strongly for the other three

models (*Supplementary Information*, Table S5). This result was obtained without integrating gene

flow between the genetic groups in the models due to methodological limitations of Splatche.

Our retained spatially explicit model (i.e. with constant population effective size before

domestication and excluding likely introgressed cluster cl5c) located the geographic origin of

cultivated sorghum in a region situated in eastern Sudan and western Ethiopia (Fig. 2-A,

*Supplementary Information*, Fig. S10-IB), in line with archaeobotanical evidence and most recent

hypotheses (Fuller and Stevens, 2018; Winchell et al., 2017).

Using archaeobotanical remains to calibrate our model of spatial expansion (*Supplementary*

*Information*, Table S2-B) led to an estimation for the onset of both the westward and southward

diffusions of the African sorghum to have occurred around 4,600 ya (4,550 ya, 95% CI 4,340-4,979

for the southward diffusion and 4,600 ya, 95% CI 4,435-5,238 for the westward one; Fig. 2-B;

*Supplementary Information*, SI Text section 2.2; Fig. S11).

The speed of the fronts of diffusion was estimated to be similar for the westward and southward

diffusions, with a spread rate of $1.26 \pm 0.4$ km/year (80% CL: 0.67-1.85 km/year and r = 0.82.) and

$1.14 \pm 0.25$ km/year (80% CL: 0.77-1.5 km/year, r = 0.88), respectively (*Supplementary Information*,

Fig. S12-A).

431
**Figure 2. Geographic origin and diffusion of the African cultivated sorghum.** A) Inferred
geographic origin of the cultivated sorghum in Africa. The posterior distribution of the latitude
and longitude of the origin in the spatial model was represented, the darker, the higher. The
blue symbols correspond to the localizations of the archaeological sites in the inferred area of
origin where the oldest remains of sorghum were found (from Table S4-B). B) Distribution of
the timing of the beginning of the westward (blue) and southward (green) movements of
diffusion of the African cultivated sorghum, removing the Ethiopian and the Egyptian
archaeological sites from the analyses. C) Effective migration surfaces representing dispersal
barriers (orange) and corridors (blue) without the cultivated accessions from the cluster cl5c.
The geographic distribution of the eight genetic clusters considered in the analysis are
illustrated on the FEEMS results map using standard deviational ellipses computed using the
CrimeStat method implemented in the Standard Deviational Ellipse plugin for QGIS.

Effective migration surfaces analysis using FEEMS (Marcus et al., 2021) mirrored the spatial

organization of sorghum genetic diversity at different scales. Low migration rates were observed in

Eastern Sudan and Western Ethiopia, the area of origin of the cultivated sorghum (Fig. 2-C). In this

area, genetic differentiation between two cultivated accessions was higher than expected given

their geographic distance. This suggests the existence of isolated populations (mainly clusters cl3

and cl4) between which gene flow was limited, while high diversity was observed within those two

clusters. High genetic differentiation combined with high intra population diversity could result from

low effective population sizes, and geographic and/or linguistic barriers to diffusion. Effective

migration surfaces revealed other isolated populations in the African Great Lakes area of nilotic

speakers (cluster cl6), in the South-Eastern Africa area of Guthrie S group speakers (cluster cl2b),

and in the Nigeria area, where no Bantoid Benue-Congo speakers are located (mainly cluster cl5a).

## Discussion

In this study, we combined genomic, archaeobotanical and ethnolinguistic data to investigate the

origin and spread of the cultivated sorghum in Africa. The origin of cultivated sorghum and the

beginning of its expansion, as estimated by genomic data, converge with the result deduced from

archaeological remains. The multidisciplinary approach adopted here indeed confirmed the East

African origin of cultivated sorghum. The southward and westward expansions of sorghum were

estimated to have begun at the same period, with a similar spread rate, by 4,600 BP, which is close

to the period estimated from archaeological remains. The complex sorghum population genetic

structure shows how its evolution dynamics were linked to both environmental and ethnolinguistic

factors: effective migration surfaces analysis, mirroring the spatial organization of sorghum genetic

diversity at different scales, confirmed that sorghum diffusion was constrained by geographic and/or

ethnolinguistic barriers. Low migration rates, and high level of genetic diversity between and within

genetic clusters in the area of origin, shed new light on the early period of sorghum cultivation.

470  Considering close associations between genetic clusters with present day ethnolinguistic groups

471  in the area of origin and beyond, on one hand, and social barriers as anthropological characteristics

472  on the other hand, allows upscaling our analysis over time and space.

**Geographic origin and expansion onset of African cultivated sorghum**

Our results localized the onset of the expansion of the African cultivated sorghum in Eastern Sudan,

specifically in the Southern Atbai region (the Eastern desert of Sudan), where the oldest

domesticated sorghum remains are known (Fuller and Stevens, 2018). Besides the probabilistic

map, one of the two hotspots of cultivated genetic diversity was localized in that same area

(*Supplementary Information*, Fig. S-8), in line with the expectation of a higher diversity in the

primary domestication center.

The second hotspot was observed in Western Africa and was mainly due to the accessions from

the Western African cluster cl5c (guinea margaritiferum). The margaritiferum subrace is highly

specific, both genetically (Burgarella et al., 2021) and morphologically, showing close relationships

to wild sorghum with evidence of gene flow (Deu et al., 2006, 1995; Gilabert et al., 2023; Mace et

al., 2013), and has been suggested to have originated from a secondary independent

domestication event (Morris et al., 2013; Sagnard et al., 2011). This western hotspot could thus

result from an independent sorghum domestication, cluster cl5c emerging from local Western

African wild sorghum and/or from introgression between the wild and cultivated sorghum refueling

the cultivated diversity with endemic wild one. More extended genomic study of this guinea

margaritiferum group will certainly shed more light on its precise origin.

Calibrating with archaeobotanical data our genomic based spatial expansion model, we estimated

that the westward and southward diffusions of cultivated sorghum from its center of origin in Eastern

Africa occurred approximately 4,600 years ago (~2,600 BCE). Our estimations are concomitant to

the estimation for the cultivated pearl millet diffusion date (Burgarella et al., 2018), and correspond

to a more ancient diffusion than the diffusion of the African rice and the African yam (Cubry et al.,

2018; Scarcelli et al., 2019).The expansion of sorghum and pearl millet coincides with the

intensification of the aridification of the "green Sahara", and more specifically with the most severe

drought episode the Sahara desert has experienced during the Holocene (Van der Meeren et al.,

2022). The harsher conditions and reduced availability of Poaceae in the environment (Kröpelin et

al., 2008) may have encouraged the cultivation of drought-resistant cereals such as sorghum and

pearl millet, and the migration of people towards regions with more favorable conditions (Fuller et

al., 2019).

It is worth noting that, with our approach, we estimated the onset of the expansion of cultivated

sorghum, rather than the timing of the domestication process itself, which may largely predate the

expansion. Even our estimation of the onset of expansion reflects a lower bound, given that the

calibration is made on archaeobotanical remains that could be found only in the case of a rather

important presence of the taxon, suggesting that reports of the presence of the crop post-date its

actual arrival. The process of crop domestication is now understood as being protracted with the

existence of an initial period of weak cultivation and/or management, rather than a rapid process

accompanied by a domestication bottleneck (Allaby, 2010; Allaby et al., 2022, 2019; Fuller et al.,

2014). The ABC results of our study do not support the hypothesis of a sharp population decline in

the population of cultivated sorghum. Our finding is consistent with a study of modern and ancient

sorghum genomes from Egyptian Nubia, which suggested that the loss of diversity observed in

cultivated sorghum occurred gradually over time and was due to an accumulation of the mutation

load rather than to a domestication bottleneck (Smith et al., 2019).

In line with these findings, sorghum archaeobotanical data argue in favor of the existence of a

protracted period for sorghum in Africa. Charred remains of wild sorghum from huts and storage

pits were identified in Egypt sites dating from approx. 8,000 BCE, and from potteries found in sites

in Central Sudan active during the fifth and fourth millennium, suggesting that wild sorghum were

likely collected and used in the Mid-Holocene (Fuller and Stevens, 2018; Lucarini and Radini, 2020;

Wasylikowa and Dahlberg, 1999). The first evidence of cultivated sorghum was found in

archaeological sites from Northern Sudan dating back to the 4[th] millennium (Winchell et al., 2017)

indicating that the domestication of sorghum was already ongoing at that time.

**Drivers of the genetic diversity of the African cultivated sorghum**

The genetic diversity and structure of African sorghum have been suggested to be shaped by a

combination of environmental, morphological (Billot et al., 2013; Deu et al., 2006; Lasky et al., 2015;

Morris et al., 2013), and ethnolinguistic factors (Labeyrie et al., 2014; Westengen et al., 2014).

Morris et al. (2013) suggested that the diffusion of the cultivated sorghum has been driven by both

geographic isolation and agroclimatic constraints. Similarly, we observed that some genetic

clusters appeared to be preferentially distributed along a specific agroclimatic zone or altitudinal

range, substantiating the role of climatic and environmental conditions in shaping sorghum

diversity.

Analyzing a distinct minicore collection of African cultivated sorghum with simple sequence repeat

markers, Westengen et al. (2014) identified three major sorghum groups that were co-distributed

with the three main language families in Africa, in agreement with the farming/language dispersal

hypothesis (Diamond and Bellwood, 2003). Our study, based on SNP markers, more finely divided

sorghum diversity in nine groups and confirmed this previous finding of structuration of sorghum

diversity at the language family scale, suggesting that sorghum genetic diversity is related to

ethnolinguistic factors. An advantage of a fine description of sorghum genetic diversity is that we

can associate sorghum diversity not only to a wide linguistic family level (Westengen et al., 2014)

but also to a finer level and local linguistic groups, freeing us from potential correlations that could

exist between language families and environmental variables at the higher level. This allows taking

into consideration the ethnolinguistic groups that might have contributed to sorghum diversity

diffusion or structure through the selection of sorghum locally-adapted to environmental conditions

or usage.

The key role of social factors in structuring crop genetic diversity has been suggested to result from

seed exchanges that were historically more oriented within than between sorghum populations

(Westengen et al., 2014). Globally, this hypothesis, which was made explicit by Leclerc & Coppens

d'Eeckenbrugge (2012), proposes that seed circulation is not random, and that speaking the same

or a common language makes such interactions more likely. Linguistic organization of crop genetics

diversity was observed for many crops, such maize (Orozco-Ramírez et al., 2016; Perales et al.,

2005), pearl millet (Naino Jika et al., 2017) or fonio (Diop et al., 2023). In the case of fonio, farmers'

migrations are proposed as the mechanism leading to the fonio spatial structure in Senegal, which

fits language diversity (Diop et al., 2023).

At a finer scale, local populations were detected in the area of origin of sorghum. Interestingly,

those populations (clusters cl3 and cl4) are differentiated whereas they harbored a high level of

intra-population diversity. This suggests that those populations were characterized by a low

effective population size, and by limited gene flow between them, probably because of the

existence of social barriers or adaptation to the environment or elevation. Action of social barriers

on sorghum genetic diversity was well documented at the local scales (Labeyrie et al., 2016;

Westengen et al., 2014). Notably, ethnolinguistic differentiation was also detected in the area of

origin of cultivated sorghum in Northern Ethiopia in three sites with similar agro-ecological and

topographic profiles, yet with admixture between the studied populations (Wendmu et al., 2023).

We identified three genetic clusters in the area of distribution of the Bantoid languages. The cluster

cl6 was associated to the Guthrie zone J in the Great Lakes region, cluster cl2b to the zone S in

South-Eastern Africa. These associations suggest that those sorghum populations moved with

people, and since stayed associated with them. The cluster cl2a was associated with four Guthrie

zones (F, G, N, P), which are located from north to south along a South-Eastern African corridor

that reaches the area of distribution of the cluster cl2b, the cluster presenting the lowest level of

genetic diversity. This could suggest a North to South migration route of sorghum (Fuller and

Stevens, 2018). This is consistent with the two areas in Eastern and South-Eastern Africa with high

effective migration rates. Using a similar analysis of effective migration rates of Bantu-speaking

populations, congruent southeastern corridors along the coast from Tanzania to South Africa and

the area of the Great Lakes were also detected for those human populations (Fortes-Lima et al.,

2023).

Our estimations of the two fronts of dispersal of the cultivated African sorghum, at a rate of approx.

1 km/year, are consistent with previous estimations of the spread rate of agriculture in multiple

areas of the world (reviewed in Fort (2021)). One striking result regarding the timing and the spread

of the two movements of diffusion is that they appear to have occurred simultaneously. Dispersals

along an east-west direction are expected to have happened more rapidly than those along a north-

south axis (Diamond and Bellwood, 2003). Indeed, climates and environments show a higher

variability latitudinally than longitudinally due to the fact that daylength and seasonality depends on

the latitude only. Interestingly, a relatively constant movement rate through time was also

suggested for Bantu-speaking populations, despite the variety of environments and interactions

they had to have encountered during their migration (Fortes-Lima et al., 2023). The results obtained

from Treemix suggest gene flow between cluster cl3, which is split between Eastern and Western

Africa, and cluster cl2b in South-Eastern Africa. The interconnection between Eastern accessions,

which are located in the sorghum area of origin, and cluster cl2b in South-Eastern Africa, suggests

a route of diffusion that corresponds well to the one proposed by Fuller and Stevens (2018). The

gene flow between cluster cl6 (Great Lakes region) and the widely distributed cluster cl1 could

suggest a key role played by Bantu speakers of Guthrie zone J in sorghum diffusion from the area

of origin and the Great Lakes region. The wide distribution of the cluster cl1 could suggest an

independent diffusion, with secondary contact leading to signature of gene flow. The complexity of

sorghum structuration could reflect the fact that African populations underwent multiple waves of

migration during their history, known as the spread-over-spread model (Fortes-Lima et al., 2023;

Seidensticker et al., 2021).

599                                     **Conclusion**

Combining genomics and archaeobotanical data, this study corroborates the Eastern African origin

of the cultivated sorghum. It also proposes two simultaneous and synchronous fronts of diffusions

from the area of origin towards the West and the South. Overall, our results also confirm the

complex geographic-based genetic structuring of the cultivated sorghum, whose genetic structure

at the African continental scale appears to be shaped by a combination of multiple factors,

environmental and ethnolinguistic ones, acting at different temporal and spatial scales. One major

outstanding question on the evolutionary history of cultivated plants and the history of their

domestication relies on their origin and how they dispersed from there: have crops dispersed

through a demic or cultural diffusion, or a combination of the two (Hewlett et al., 2002; Leclerc and

Coppens d'Eeckenbrugge, 2012)? Under a scenario of a demic diffusion, one would expect crops

to be structured by social factors as they move together with the farmers while a cultural diffusion

model would lead to patterns of isolation by distance since it depends on interactions with

neighboring societies. If our study helps the understanding of the beginning of sorghum

domestication, a comparative study of population genetics of ancient human and crop populations

(see for example Tao et al. (2023)) and/or additional archaeobotanical data (see for example Cobo

et al. (2019)) would be necessary to give insight into the main mechanism, a demic and/or a cultural

diffusion, responsible for the diffusion of agriculture, and thus help us to better understand their

evolutionary history.

619                          **Data and code availability**

The raw GBS data were available in the NCBI Sequence Read Archive from references Brenton et

al., 2016; Lasky et al., 2015; Yu et al., 2016. The vcf file containing the filtered SNPs and the

passport data of the accessions analyzed in the study are publicly available on the CIRAD

Dataverse repository, under the doi identifier https://doi.org/10.18167/DVN1/SAAMUT. The R

scripts for the analyses performed in the study are available on the CIRAD gitlab plateform

(https://gitlab.cirad.fr/agap/sorgho/africrop_sorghum).

627                          **Conflict of interest disclosure**

628   The authors declare that they have no conflict of interest in relation to the content of this article.

630                          **Acknowledgments**

This work was supported by an ANR grant (AfriCrop project, ANR-13-BSV7-0017). We

acknowledge the following bioinformatic facilities for providing HPC resources and support: the

Core Cluster of the Institut Français de Bioinformatique, the MESO@LR-Platform at the University

of Montpellier and the INRAE MIGALE bioinformatics facility (MIGALE, INRAE, 2020. Migale

bioinformatics Facility, doi: 10.15454/1.5572390655343293E12).

**References**

Alexander, D.H., Novembre, J., Lange, K., 2009. Fast model-based estimation of ancestry in
unrelated individuals. Genome Res. 19, 1655–1664.
https://doi.org/10.1101/gr.094052.109

Allaby, R., 2010. Integrating the processes in the evolutionary system of domestication. J. Exp. Bot.
61, 935–944. https://doi.org/10.1093/jxb/erp382

Allaby, R.G., Stevens, C.J., Kistler, L., Fuller, D.Q., 2022. Emerging evidence of plant
domestication as a landscape-level process. Trends Ecol. Evol. 37, 268–279.
https://doi.org/10.1016/j.tree.2021.11.002

Allaby, R.G., Ware, R.L., Kistler, L., 2019. A re-evaluation of the domestication bottleneck from
archaeogenomic evidence. Evol. Appl. 12, 29–37. https://doi.org/10.1111/eva.12680

Beck, H.E., Zimmermann, N.E., McVicar, T.R., Vergopolan, N., Berg, A., Wood, E.F., 2018. Present
and future Köppen-Geiger climate classification maps at 1-km resolution. Sci. Data 5,
180214. https://doi.org/10.1038/sdata.2018.214

Billot, C., Ramu, P., Bouchet, S., Chantereau, J., Deu, M., Gardes, L., Noyer, J.-L., Rami, J.-F.,
Rivallan, R., Li, Y., Lu, P., Wang, T., Folkertsma, R.T., Arnaud, E., Upadhyaya, H.D.,
Glaszmann, J.-C., Hash, C.T., 2013. Massive sorghum collection genotyped with SSR
markers to enhance use of global genetic resources. PLOS ONE 8, e59714.
https://doi.org/10.1371/journal.pone.0059714

Bolger, A.M., Lohse, M., Usadel, B., 2014. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics 30, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Brenton, Z.W., Cooper, E.A., Myers, M.T., Boyles, R.E., Shakoor, N., Zielinski, K.J., Rauh, B.L.,
Bridges, W.C., Morris, G.P., Kresovich, S., 2016. A genomic resource for the development,
improvement, and exploitation of sorghum for bioenergy. Genetics 204, 21–33.
https://doi.org/10.1534/genetics.115.183947

Burgarella, C., Berger, A., Glémin, S., David, J., Terrier, N., Deu, M., Pot, D., 2021. The road to
sorghum domestication: evidence from nucleotide diversity and gene expression patterns.
Front. Plant Sci. 12. https://doi.org/10.3389/fpls.2021.666075.

Burgarella, C., Cubry, P., Kane, N.A., Varshney, R.K., Mariac, C., Liu, X., Shi, C., Thudi, M.,
Couderc, M., Xu, X., Chitikineni, A., Scarcelli, N., Barnaud, A., Rhoné, B., Dupuy, C.,
François, O., Berthouly-Salazar, C., Vigouroux, Y., 2018. A western Sahara centre of

domestication inferred from pearl millet genomes. Nat. Ecol. Evol. 2, 1377–1380. https://doi.org/10.1038/s41559-018-0643-y

Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M., Lee, J.J., 2015. Second-generation PLINK: rising to the challenge of larger and richer datasets. GigaScience 4, s13742-015-0047–8. https://doi.org/10.1186/s13742-015-0047-8

Cobo, J.M., Fort, J., Isern, N., 2019. The spread of domesticated rice in eastern and southeastern Asia was mainly demic. J. Archaeol. Sci. 101, 123–130. https://doi.org/10.1016/j.jas.2018.12.001

Csilléry, K., François, O., Blum, M.G.B., 2012. abc: an R package for approximate Bayesian computation (ABC). Methods Ecol. Evol. 3, 475–479. https://doi.org/10.1111/j.2041-210X.2011.00179.x

Cubry, P., Tranchant-Dubreuil, C., Thuillet, A.-C., Monat, C., Ndjiondjop, M.-N., Labadie, K., Cruaud, C., Engelen, S., Scarcelli, N., Rhoné, B., Burgarella, C., Dupuy, C., Larmande, P., Wincker, P., François, O., Sabot, F., Vigouroux, Y., 2018. The rise and fall of African rice cultivation revealed by analysis of 246 new genomes. Curr. Biol. 28, 2274-2282.e6. https://doi.org/10.1016/j.cub.2018.05.066

Currat, M., Ray, N., Excoffier, L., 2004. splatche: a program to simulate genetic diversity taking into account environmental heterogeneity. Mol. Ecol. Notes 4, 139–142. https://doi.org/10.1046/j.1471-8286.2003.00582.x

Danecek, P., Auton, A., Abecasis, G., Albers, C.A., Banks, E., DePristo, M.A., Handsaker, R.E., Lunter, G., Marth, G.T., Sherry, S.T., McVean, G., Durbin, R., 1000 Genomes Project Analysis Group, 2011. The variant call format and VCFtools. Bioinformatics 27, 2156–2158. https://doi.org/10.1093/bioinformatics/btr330

de Wet, J.M.J., Huckabay, J.P., 1967. The origin of *Sorghum bicolor*. II. Distribution and domestication. Evolution 21, 787–802. https://doi.org/10.2307/2406774

DePristo, M.A., Banks, E., Poplin, R., Garimella, K.V., Maguire, J.R., Hartl, C., Philippakis, A.A., del Angel, G., Rivas, M.A., Hanna, M., McKenna, A., Fennell, T.J., Kernytsky, A.M., Sivachenko, A.Y., Cibulskis, K., Gabriel, S.B., Altshuler, D., Daly, M.J., 2011. A framework for variation discovery and genotyping using next-generation DNA sequencing data. Nat. Genet. 43, 491–498. https://doi.org/10.1038/ng.806

Deu, M., Hamon, P., Dufour, P., D'Hont, A., Lanaud, C., Chantereau, J., 1995. Mitochondrial DNA diversity in wild and cultivated sorghum. Genome 38, 635–645. https://doi.org/10.1139/g95-081

Deu, M., Rattunde, F., Chantereau, J., 2006. A global view of genetic diversity in cultivated
sorghums using a core collection. Genome 49, 168–180. https://doi.org/10.1139/g05-092

Diamond, J., Bellwood, P., 2003. Farmers and their languages: the first expansions. Science 300,
597–603. https://doi.org/10.1126/science.1078208

Dinno A (2024). dunn.test: Dunn's Test of Multiple Comparisons Using Rank Sums. R package
version 1.3.6. Available at https://cran.r-project.org/web/packages/dunn.test/

Diop, B.M., Guèye, M.C., Leclerc, C., Deu, M., Zekraoui, L., Calatayud, C., Rivallan, R., Kaly, J.R.,
Cissé, M., Piquet, M., Diack, O., Ngom, A., Berger, A., Ndoye, I., Ndir, K., Vigouroux, Y.,
Kane, N.A., Barnaud, A., Billot, C., 2023. Ethnolinguistic and genetic diversity of fonio
(*Digitaria exilis*) in Senegal. Plants People Planet. https://doi.org/10.1002/ppp3.10428

Evanno, G., Regnaut, S., Goudet, J., 2005. Detecting the number of clusters of individuals using
the software structure: a simulation study. Mol. Ecol. 14, 2611–2620.
https://doi.org/10.1111/j.1365-294X.2005.02553.x

FAO, 2023. Agricultural production statistics 2000–2022.
https://openknowledge.fao.org/handle/20.500.14283/cc9205en

Fitak, R.R., 2021. OptM: estimating the optimal number of migration edges on population trees
using Treemix. Biol. Methods Protoc. 6, bpab017.
https://doi.org/10.1093/biomethods/bpab017

Fort, J., 2021. The spread of agriculture: quantitative laws in Prehistory?, in: Pardo-Gordó, S.,
Bergin, S. (Eds.), Simulating Transitions to Agriculture in Prehistory, Computational Social
Sciences. Springer International Publishing, Cham, pp. 17–28. https://doi.org/10.1007/978-
3-030-83643-6_2

Fort, J., Pujol, T., Cavalli-Sforza, L.L., 2004. Palaeolithic populations and waves of advance. Camb.
Archaeol. J. 14, 53–61. https://doi.org/10.1017/S0959774304000046

Fortes-Lima, C.A., Burgarella, C., Hammarén, R., Eriksson, A., Vicente, M., Jolly, C., Semo, A.,
Gunnink, H., Pacchiarotti, S., Mundeke, L., Matonda, I., Muluwa, J.K., Coutros, P.,
Nyambe, T.S., Cikomola, J.C., Coetzee, V., de Castro, M., Ebbesen, P., Delanghe, J.,
Stoneking, M., Barham, L., Lombard, M., Meyer, A., Steyn, M., Malmström, H., Rocha, J.,
Soodyall, H., Pakendorf, B., Bostoen, K., Schlebusch, C.M., 2023. The genetic legacy of
the expansion of Bantu-speaking peoples in Africa. Nature 1–8.
https://doi.org/10.1038/s41586-023-06770-6

Fuller, D., Champion, L., Stevens, C., 2019. Comparing the tempo of cereal dispersal and the
agricultural transition: two African and one West Asian trajectory, in: Eichhorn, B., Hohn,
A. (Eds.), In: Eichhorn, B and Hohn, A, (Eds.) Trees, Grasses and Crops – People and

735    Plants in Sub-Saharan Africa and Beyond.  Verlag Dr. Rudolf Habelt GmbH: Bonn,
736    Germany. (2019). Verlag Dr. Rudolf Habelt GmbH, Bonn, Germany.

737 Fuller, D.Q., Denham, T., Arroyo-Kalin, M., Lucas, L., Stevens, C.J., Qin, L., Allaby, R.G.,
738    Purugganan, M.D., 2014. Convergent evolution and parallelism in plant domestication
739    revealed by an expanding archaeological record. Proc. Natl. Acad. Sci. 111, 6147–6152.
740    https://doi.org/10.1073/pnas.1308937110

741 Fuller, D.Q., Stevens, C.J., 2018. Sorghum domestication and diversification: a current
742    archaeobotanical perspective, in: Mercuri, A.M., D'Andrea, A.C., Fornaciari, R., Höhn, A.
743    (Eds.), Plants and People in the African Past: Progress in African Archaeobotany. Springer
744    International Publishing, Cham, pp. 427–452. https://doi.org/10.1007/978-3-319-89839-
745    1_19

746 Gilabert, A., Burgarella, C., Calatayud, C., Berger, A., Rami, J.-F., Pot, D., Deu, M., 2023. Shedding
747    light on the evolutionary history of wild and cultivated african sorghum: the guinea
748    margaritiferum case. Poster presented at: Sorghum in the 21st century: Resiliency and
749    sustainability in the face of climate change; June 5-9 2023; Montpellier, France.

750 Goudet, J., 2005. hierfstat, a package for R to compute and test hierarchical $F$-statistics. Mol. Ecol.
751    Notes 5, 184–186. https://doi.org/10.1111/j.1471-8286.2004.00828.x

752 Guthrie, M., 1971. Comparative Bantu: An Introduction to the Comparative Linguistics and
753    Prehistory of the Bantu Languages. Gregg, Famborough, UK.

754 Guthrie, M., 1948. The Classification of the Bantu languages. Oxford University Press.

755 Harlan, J.R., Stemler, A.B.L., 1976. The races of sorghum in Africa, in: Harlan, J.R., Wet, J.M.J.
756    de, Stemler, A.B.L. (Eds.), Origins of African Plant Domestication. De Gruyter Mouton,
757    Berlin, New York, pp. 465–478. https://doi.org/doi:10.1515/9783110806373.465

758 Hewlett, B.S., De Silvestri, A., Guglielmino, C.R., 2002. Semes and genes in Africa. Curr.
759    Anthropol. 43, 313–321. https://doi.org/10.1086/339379

760 Jerardino, A., Fort, J., Isern, N., Rondelli, B., 2014. Cultural diffusion was the main driving
761    mechanism of the Neolithic transition in Southern Africa. PLOS ONE 9, e113672.
762    https://doi.org/10.1371/journal.pone.0113672

763 Kimber, C.T., 2000. Origins of domesticated sorghum and its early diffusion to India and China, in:
764    Sorghum: Origin, History, Technology, and Production. John Wiley & Sons, New York, pp.
765    3–98.

766 Kröpelin, S., Verschuren, D., Lézine, A.-M., Eggermont, H., Cocquyt, C., Francus, P., Cazet, J.-P.,
767    Fagot, M., Rumes, B., Russell, J.M., Darius, F., Conley, D.J., Schuster, M., von
768    Suchodoletz, H., Engstrom, D.R., 2008. Climate-driven ecosystem succession in the

Sahara: The past 6000 Years. Science 320, 765–768.
https://doi.org/10.1126/science.1154913

Labeyrie, V., Deu, M., Barnaud, A., Calatayud, C., Buiron, M., Wambugu, P., Manel, S.,
Glaszmann, J.-C., Leclerc, C., 2014. Influence of ethnolinguistic diversity on the sorghum
genetic patterns in subsistence farming systems in Eastern Kenya. PLOS ONE 9, e92178.
https://doi.org/10.1371/journal.pone.0092178

Labeyrie, V., Thomas, M., Muthamia, Z.K., Leclerc, C., 2016. Seed exchange networks, ethnicity,
and sorghum diversity. Proc. Natl. Acad. Sci. 113, 98–103.
https://doi.org/10.1073/pnas.1513238112

Lasky, J.R., Upadhyaya, H.D., Ramu, P., Deshpande, S., Hash, C.T., Bonnette, J., Juenger, T.E.,
Hyma, K., Acharya, C., Mitchell, S.E., Buckler, E.S., Brenton, Z., Kresovich, S., Morris,
G.P., 2015. Genome-environment associations in sorghum landraces predict adaptive
traits. Sci. Adv. 1, e1400218. https://doi.org/10.1126/sciadv.1400218

Leclerc, C., Coppens d'Eeckenbrugge, G., 2012. Social organization of crop genetic diversity. The
G × E × S interaction model. Diversity 4, 1–32. https://doi.org/10.3390/d4010001

Leinonen, R., Sugawara, H., Shumway, M., on behalf of the International Nucleotide Sequence
Database Collaboration, 2011. The Sequence Read Archive. Nucleic Acids Res. 39, D19–
D21. https://doi.org/10.1093/nar/gkq1019

Lemaire, L., Jay, F., Lee, I.-H., Csilléry, K., Blum, M.G.B., 2016. Goodness-of-fit statistics for
approximate Bayesian computation. https://doi.org/10.48550/arXiv.1601.04096

Lewis, M., 2009. Ethnologue: languages of the World, 16th edition. SIL International.

Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics 25, 1754–1760. https://doi.org/10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,
Durbin, R., 1000 Genome Project Data Processing Subgroup, 2009. The Sequence
Alignment/Map format and SAMtools. Bioinformatics 25, 2078–2079.
https://doi.org/10.1093/bioinformatics/btp352

Lucarini, G., Radini, A., 2020. First direct evidence of wild plant grinding process from the Holocene
Sahara: use-wear and plant micro-residue analysis on ground stone tools from the Farafra
Oasis, Egypt. Quat. Int., Africa under the microscope: What's new in techno-functional
analyses in archaeology? 555, 66–84. https://doi.org/10.1016/j.quaint.2019.07.028

Mace, E.S., Tai, S., Gilding, E.K., Li, Y., Prentis, P.J., Bian, L., Campbell, B.C., Hu, W., Innes, D.J.,
Han, X., Cruickshank, A., Dai, C., Frère, C., Zhang, H., Hunt, C.H., Wang, X., Shatte, T.,
Wang, M., Su, Z., Li, J., Lin, X., Godwin, I.D., Jordan, D.R., Wang, J., 2013. Whole-genome

sequencing reveals untapped genetic potential in Africa's indigenous cereal crop sorghum.
Nat. Commun. 4, 2320. https://doi.org/10.1038/ncomms3320

Maho, J.F., 2009. NUGL Online : the online version of the New Updated Guthrie List, a referential
classification of the Bantu languages.

Marcus, J., Ha, W., Barber, R.F., Novembre, J., 2021. Fast and flexible estimation of effective
migration surfaces. eLife 10, e61927. https://doi.org/10.7554/eLife.61927

McCormick, R.F., Truong, S.K., Sreedasyam, A., Jenkins, J., Shu, S., Sims, D., Kennedy, M.,
Amirebrahimi, M., Weers, B.D., McKinley, B., Mattison, A., Morishige, D.T., Grimwood, J.,
Schmutz, J., Mullet, J.E., 2018. The *Sorghum bicolor* reference genome: improved
assembly, gene annotations, a transcriptome atlas, and signatures of genome
organization. Plant J. 93, 338–354. https://doi.org/10.1111/tpj.13781

Morris, G.P., Ramu, P., Deshpande, S.P., Hash, C.T., Shah, T., Upadhyaya, H.D., Riera-Lizarazu,
O., Brown, P.J., Acharya, C.B., Mitchell, S.E., Harriman, J., Glaubitz, J.C., Buckler, E.S.,
Kresovich, S., 2013. Population genomic and genome-wide association studies of
agroclimatic traits in sorghum. Proc. Natl. Acad. Sci. 110, 453–458.
https://doi.org/10.1073/pnas.1215985110

Naino Jika, A.K., Dussert, Y., Raimond, C., Garine, E., Luxereau, A., Takvorian, N., Djermakoye,
R.S., Adam, T., Robert, T., 2017. Unexpected pattern of pearl millet genetic diversity
among ethno-linguistic groups in the Lake Chad Basin. Heredity 118, 491–502.
https://doi.org/10.1038/hdy.2016.128

Orozco-Ramírez, Q., Ross-Ibarra, J., Santacruz-Varela, A., Brush, S., 2016. Maize diversity
associated with social origin and environmental variation in Southern Mexico. Heredity 116,
477–484. https://doi.org/10.1038/hdy.2016.10

Perales, H.R., Benz, B.F., Brush, S.B., 2005. Maize diversity and ethnolinguistic diversity in
Chiapas, Mexico. Proc. Natl. Acad. Sci. 102, 949–954.
https://doi.org/10.1073/pnas.0408701102

Perrier, X., Jacquemoud-Collet, J.-P., 2019. DARwin - Dissimilarity Analysis and Representation
for Windows. Available at: https://darwin.cirad.fr/

Pickrell, J.K., Pritchard, J.K., 2012. Inference of population splits and mixtures from genome-wide
allele frequency data. PLOS Genet. 8, e1002967.
https://doi.org/10.1371/journal.pgen.1002967

Pritchard, J.K., Stephens, M., Donnelly, P., 2000. Inference of population structure using multilocus
genotype data. Genetics 155, 945–959. https://doi.org/10.1093/genetics/155.2.945

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M.A.R., Bender, D., Maller, J., Sklar,
P., Bakker, P.I.W. de, Daly, M.J., Sham, P.C., 2007. PLINK: a tool set for whole-genome

838  association and population-based linkage analyses. Am. J. Hum. Genet. 81, 559–575.
839    https://doi.org/10.1086/519795

840 R Core Team, 2024. R: A Language and Environment for Statistical Computing. R Foundation for
841    Statistical Computing, Vienna, Austria. URL https://www.R-project.org/

842 Sagnard, F., Deu, M., Dembélé, D., Leblois, R., Touré, L., Diakité, M., Calatayud, C., Vaksmann,
843    M., Bouchet, S., Mallé, Y., Togola, S., Traoré, P.C.S., 2011. Genetic diversity, structure,
844    gene flow and evolutionary relationships within the *Sorghum bicolor* wild–weedy–crop
845    complex in a western African region. Theor. Appl. Genet. 123, 1231–1246.
846    https://doi.org/10.1007/s00122-011-1662-0

847 Saitou, N., Nei, M., 1987. The neighbor-joining method: a new method for reconstructing
848    phylogenetic trees. Mol. Biol. Evol. 4, 406–425.
849    https://doi.org/10.1093/oxfordjournals.molbev.a040454

850 Scarcelli, N., Cubry, P., Akakpo, R., Thuillet, A.-C., Obidiegwu, J., Baco, M.N., Otoo, E., Sonké, B.,
851    Dansi, A., Djedatin, G., Mariac, C., Couderc, M., Causse, S., Alix, K., Chaïr, H., François,
852    O., Vigouroux, Y., 2019. Yam genomics supports West Africa as a major cradle of crop
853    domestication. Sci. Adv. 5, eaaw1947. https://doi.org/10.1126/sciadv.aaw1947

854 Seidensticker, D., Hubau, W., Verschuren, D., Fortes-Lima, C., de Maret, P., Schlebusch, C.M.,
855    Bostoen, K., 2021. Population collapse in Congo rainforest from 400 CE urges
856    reassessment of the Bantu expansion. Sci. Adv. 7, eabd8352.
857    https://doi.org/10.1126/sciadv.abd8352

858 Smith, C.W., Frederiksen, R.A., 2000. Sorghum: origin, history, technology, and production, Wiley
859    Series in Crop Science. Wiley.

860 Smith, O., Nicholson, W.V., Kistler, L., Mace, E., Clapham, A., Rose, P., Stevens, C., Ware, R.,
861    Samavedam, S., Barker, G., Jordan, D., Fuller, D.Q., Allaby, R.G., 2019. A domestication
862    history of dynamic adaptation and genomic deterioration in Sorghum. Nat. Plants 5, 369–
863    379. https://doi.org/10.1038/s41477-019-0397-9

864 Tao, L., Yuan, H., Zhu, K., Liu, X., Guo, J., Min, R., He, H., Cao, D., Yang, X., Zhou, Z., Wang, R.,
865    Zhao, D., Ma, H., Chen, J., Zhao, J., Li, Y., He, Y., Suo, D., Zhang, R., Li, S., Li, L., Yang,
866    F., Li, H., Zhang, L., Jin, L., Wang, C.-C., 2023. Ancient genomes reveal millet farming-
867    related demic diffusion from the Yellow River into southwest China. Curr. Biol. 0.
868    https://doi.org/10.1016/j.cub.2023.09.055

869 Van der Meeren, T., Verschuren, D., Sylvestre, F., Nassour, Y.A., Naudts, E.L., Aguilar Ortiz, L.E.,
870    Deschamps, P., Tachikawa, K., Bard, E., Schuster, M., Abderamane, M., 2022. A

predominantly tropical influence on late Holocene hydroclimate variation in the hyperarid
central Sahara. Sci. Adv. 8, eabk1261. https://doi.org/10.1126/sciadv.abk1261

Wasylikowa, K., Dahlberg, J., 1999. Sorghum in the economy of the early Neolithic nomadic tribes
at Nabta Playa, Southern Egypt, in: van der Veen, M. (Ed.), The Exploitation of Plant
Resources in Ancient Africa. Springer US, Boston, MA, pp. 11–31.
https://doi.org/10.1007/978-1-4757-6730-8_2

Wendmu, T.A., Gebrelibanos, T.S., Kovi, M.R., Ring, K.H., de Boer, H.J., Abera, F.A., Westengen,
O.T., 2023. "People gathered by sorghum": cultural practices and sorghum diversity in
Northern Ethiopia. Hum. Ecol. 51, 923–935. https://doi.org/10.1007/s10745-023-00442-9

Westengen, O.T., Okongo, M.A., Onek, L., Berg, T., Upadhyaya, H., Birkeland, S., Kaur Khalsa,
S.D., Ring, K.H., Stenseth, N.C., Brysting, A.K., 2014. Ethnolinguistic structuring of
sorghum genetic diversity in Africa and the role of local seed systems. Proc. Natl. Acad.
Sci. 111, 14100–14105. https://doi.org/10.1073/pnas.1401646111

Winchell, F., Stevens, C.J., Murphy, C., Champion, L., Fuller, D.Q., 2017. Evidence for sorghum
domestication in fourth millennium BC Eastern Sudan: spikelet morphology from ceramic
impressions of the Butana group. Curr. Anthropol. 58, 673–683.
https://doi.org/10.1086/693898

Yu, X., Li, X., Guo, T., Zhu, C., Wu, Y., Mitchell, S.E., Roozeboom, K.L., Wang, D., Wang, M.L.,
Pederson, G.A., Tesso, T.T., Schnable, P.S., Bernardo, R., Yu, J., 2016. Genomic
prediction contributing to a promising global strategy to turbocharge gene banks. Nat.
Plants 2, 1–7. https://doi.org/10.1038/nplants.2016.150