Genomic and Phenotypic Characterization of Mupirocin Resistant *Staphylococcus aureus* Clinical Isolates

Ariana M. Virgillio[1,2*]; Emily A. Felton[2,3*]; Jessica K. Jackson[3], Sarah J. Kennedy[3], Deanna N. Becker[4]; Amorce Lima[4]; Kimberly Atrubin[4]; Eleonora Cella[5]; Taj Azarian[5]; Suzane Silbert[4]; Lindsey N. Shaw[2,3,6#]; Kami Kim[1,2,4,6#]

[1]

[1]Division of Infectious Disease and International Medicine, University of South Florida, Tampa, Florida, USA

[2] Center for Antimicrobial Resistance, University of South Florida, Tampa, FL, USA

[3] Department of Molecular Biosciences, University of South Florida, Tampa, FL, USA

[4]Tampa General Hospital, Tampa, Florida, USA

[5]Burnett School of Biomedical Sciences, University of Central Florida, Orlando, Florida, USA

[6]Center for Global Health Interdisciplinary Research, University of South Florida, Tampa, Florida, USA

*Equal contribution

#Address correspondence to:

Lindsey Shaw, PhD

shaw@usf.edu

Department of Molecular Biosciences

University of South Florida

4202 East Fowler Avenue, ISA2015

Tampa, FL 33620-5150

USA

OR

Kami Kim, MD

kamikim@usf.edu

Division of Infectious Disease and International Medicine

University of South Florida

Interdisciplinary Research Building (IDRB)

3720 Spectrum Boulevard, Suite 304

Tampa, FL 33612-9415

USA

Key points (40 words):

• In a survey of clinical isolates in Florida, 34.4% of MRSA strains were mupirocin

resistant.

• Mupirocin resistance correlated with mutations in *ileS* or carriage of *mupA*.

• We found evidence for emergence of MRSA mupirocin-resistant strains that were

sequence type ST3390.

Word count: 3499

Author contributions: AMV and EAF performed all experiments, analyzed the data and wrote

the manuscript; JKJ and SJK performed genomic analysis of ST8 sequence types. DNB, AL and

SS provided strains and guided MIC studies/interpretation; KA provided prior mupirocin

surveillance data and advised on current infection prevention practices; EC and TA sequenced

the first 80 strains; LNS and KK supervised the project and edited the final manuscript. All

authors provided input into the manuscript content and interpretation of data.

Abstract

Background: Colonization with *Staphylococcus aureus* is a risk factor for subsequent infection.

Decolonization with the topical antibiotic mupirocin is effective and reduces the risk of subsequent

S. aureus infection for both methicillin-sensitive (MSSA) and methicillin-resistant (MRSA) strains

but may select for mupirocin-resistant isolates.

Methods: We characterized oxacillin and mupirocin susceptibility amongst 384 *S. aureus* strains

isolated from clinical samples isolated 2017-2023 in Tampa, Florida, spanning strains collected

before and after the onset of the COVID-19 pandemic. Whole genome sequencing of bacterial

isolates was conducted in parallel and correlated with drug susceptibility profiles.

Results: Mupirocin resistance (MupR) was nearly exclusively present in MRSA strains (103/106

97.1% of MupR; 103/299 34.4% of MRSA). Although our hospital protocol for decolonization

shifted to povidone iodine in the Post-COVID period, the overall prevalence of MupR did not

change in Pre-COVID and Post-COVID samples (28.9% vs 26%). Genotype correlated with

antibiotic susceptibility with low level MupR (MupLR), linked to mutations in *ileS* and high level

MupR (MupHR), linked to the presence of *mupA*. Genome analysis revealed that most MupR

strains fell into three sequence types (ST) falling into two major clonal complexes (CC): CC8 ST8

(including Community-Associated MRSA strains USA300 and USA500), CC5 ST5 (associated

with Healthcare-Associated MRSA such as USA100), and CC5 ST3390. ST3390 isolates had the

highest prevalence of MupR (30/36 83%; MupHR 20/36 55.6%; MupLR 10/36 27.8%).

Conclusions: Mupirocin resistance was prevalent in our hospital MRSA strains. We also found

evidence for emergence and persistence of ST3390 MRSA-MupR strains in Florida.

Keywords: Staphylococcus aureus, MRSA, mupirocin, antimicrobial resistance, decolonization

Abbreviations: methicillin resistant *Staphylococcus aureus* (MRSA); methicillin sensitive

Staphylococcus aureus (MSSA); Mupirocin Resistant (MupR); clonal complexes (CC); sequence

type (ST)

Introduction

Infections with *Staphylococcus aureus* are major causes of skin and soft tissue infections, as well

as invasive infections such as endocarditis, pneumonia, and osteomyelitis. Colonization of skin

and mucosal surfaces with *S. aureus* is a risk factor for subsequent clinical infections [1].

Decolonization is an effective method for reducing risk of subsequent *S.aureus* infections,

including those caused by methicillin resistant *S. aureus* (MRSA) [2]. Universal decolonization

with mupirocin and chlorhexidine body washes has been shown in randomized control trials as

being more effective than targeted decolonization [3]. Decolonization of *S. aureus* prevents post-

surgical infections, ventilator associated pneumonias and blood stream infections [4].

Mupirocin inhibits bacterial protein synthesis by competitively binding to the isoleucine specific

site of the isoleucyl-tRNA synthetase preventing the conversion of isoleucine and tRNA to

isoleucyl-tRNA. As this is an essential process, this inhibition leads to bacterial cell death. There

have been reports of increasing prevalence of mupirocin resistance, particularly amongst MRSA

strains, hypothesized due to increased usage of decolonization protocols designed to prevent

MRSA infections [5].

Mupirocin resistance (MupR) can be categorized as mupirocin susceptible (MupS, ≤4 μg/mL),

low-level resistant (MupLR, 8-64 μg/mL), and high-level resistant (MupHR, ≥ 512 μg/mL) [6] [7]

[8] [9] [10] [11]. Isolates demonstrating MupLR typically have a chromosomal mutation altering the

Rossman fold of the native isoleucyl-tRNA synthetase (IleS) protein sequence [12]. The most

prevalent mutations are V588F and V631F, which result in bulkier residue within the enzymatic

binding site, blocking association with mupirocin. The major determining factor behind high-level

resistance is the gene *mupA*, although exceptions have been noted, including *mupA*-negative

MupHR-isolates that possess a different gene, *mupB*, which is 65% similar to *mupA* [13]. Whilst

mupirocin binds discriminately to the substrate binding sites of bacterial isoleucyl-tRNA

synthetases, it has low affinity for eukaryotic isoleucyl-tRNA synthetases. Both MupA and MupB

have altered sequences that are closer to that of eukaryotes, whilst still functional in prokaryotes.

Importantly, both genes can be found on multi-resistant plasmids, facilitating the spread of

resistance.

In this study we apply a combination of Whole Genome Sequencing (WGS) and phenotypic

characterization to investigate changes in the molecular epidemiology of *S. aureus* mupirocin

resistance within a large metropolitan hospital in Florida. Our study period includes isolates before

and after the COVID-19 pandemic, which coincided with a switch in hospital protocols from

mupirocin to povidone iodine for decolonization.

Methods

SA Strain Sampling: A random convenience sample of clinical *S. aureus* isolates was collected

from discarded residual clinical samples at Tampa General Hospital (TGH) between 2017-2023.

TGH is a 1040 bed not-for-profit academic medical center serving a population exceeding six

million across western Florida. The TGH research lab archives more MRSA than MSSA isolates

and randomly archives samples from different body sites.

Patient Consent Statement: Collection of samples for this study was reviewed by our local IRB

and was deemed not human subjects research and did not require patient consent.

MIC Testing: Samples were struck to TSA for isolation overnight and transferred to TSB for

overnight growth at 37°C in a shaking incubator. MIC testing was performed using a top agar

method. Clinical susceptibility data were verified using E-tests MupR was designated as

sensitive (MIC <4 µg/mL), low-level or intermediate resistant (MupLR, MIC = 8-256 µg/mL), or

high level resistant (MupHR, MIC >512 µg/mL). Bacteria were suspended to 0.5 McFarland in

Mueller Hinton II agar for mupirocin testing and Muller Hinton II Agar + 2% NaCl for oxacillin

testing. Inoculated agar (10mL) was poured over premade uninoculated plates with their

corresponding agar. Test strips from Liofilchem were placed on dried agar corresponding to the

antibiotic being tested. Muller Hinton II agar with mupirocin test strips were incubated inverted at

37°C for 18 hours. Muller Hinton II +2% NaCl agar with oxacillin test strips were incubated

inverted at 37°C for 24 hours. Mupirocin plates were read with the guidance of "MIC Test Strip

Photographic Guide" with interpretation as in current literature. Oxacillin plates were read using

CLSI breakpoints provided by Liofilchem. Analysis for statistical differences between groups

was performed with SPSS using χ^2 for significance (p=0.05).

Whole Genome Sequencing (WGS): Overnight cultures were grown in TSB at 37ºC with

shaking. Genomic DNA was collected using the DNeasy Blood and Tissue kit (Qiagen) following

the gram-positive specific pretreatment method, with the addition of 20mg/ml lysostaphin. For 80

strains (Run 1), quality and quantity was assessed using a Qubit 4 Fluorometer, and sequencing

occurred on an Illumina MiSeq instrument with a Nextera Flex library preparation kit and V3 flow

cell chemistry. For the remaining 304 strains, quality and quantity was assessed using a Quant-it

fluorescent assay kit (Invitrogen) and a Biotek Citation5 microplate reader. Libraries were

prepared using a Hackflex [14] adapted version of the Illumina DNA-seq library preparation kit.

Sequencing of 70 of the Pre-COVID samples (Run 2) was performed using a Nextseq550 Mid-

Output v2.5 Kit (300 cycles) and 234 Post-COVID samples were sequenced using a Nextseq2000

with P1 flow cell chemistry (Run 3). Runs 2 and 3 were performed at the USF Genomics Center

Core facilities. Strains are listed in Supplementary Table 1.

For long read sequencing of TGH462, DNA was extracted using the QIAGEN genomic tip 100/G

and genomic DNA buffer kits. Genomic DNA integrity and concentration were accessed via qubit

4 (Invitrogen) and Genomic DNA Tapestation (Agilent) QC checks. Samples were then treated

with the short fragment eliminator (Nanopore) and prepared with the Rapid Barcoding library prep

kit (SQK-RBK114.24). Libraries were sequenced on an R10.4.1 flowcell using a MinION Mk1B

with default settings of 72 hour run length and 200 bp minimum read length.

Genomic assembly and molecular characterization: The raw reads for all samples were

trimmed via Trimmomatic v0.39 [15] with SLIDINGWINDOW:10:20 MINLEN:31 TRAILING:2

settings [16]. Trimming quality checks were performed with FastQC v0.12.0. The reads were then

de novo assembled with SPAdes v3.15.5 using default settings [17], and assemblies were

checked for quality via Quast v5.2.0 [18]. Genomes were then sequence-typed with pyMLST [19],

annotated with prokka, and antimicrobial resistance (AMR) genes identified using AMRFinderPlus

v3.10.45 [20]. The presence of DNA and translated Protein sequences for AMR genes of interest

were confirmed and aligned using BLASTncbi-BLAST v2.13.0+ [21]. The staphtyper workflow as

part of Bactopia v3.0.1 was used to further characterize isolates based on Agr group [22],

SCCmec type [23], and Spa type [24].

Phylogenetic analysis: To explore relatedness of the cluster of ST8-t008 that harbor the novel

R112Q *ileS* mutation, Snippy was used to generate an initial core genome single-nucleotide

polymorphism alignment. Following this, Gubbins was used to create a recombination masked

alignment. From this alignment RAxML was used to infer a maximum-likelihood mid-point rooted

tree. The subsequent tree was visualized and annotated using iTOL webserver.

<div align="center">**Results**</div>

Mupirocin Resistance (MupR) is Common Amongst Tampa MRSA isolates: To explore the

prevalence of MupR in our hospital system, we curated and characterized a collection of 384 *S.*

aureus strains. The collection was isolated between February 2017 and September 2023, and

comprised of 299 MRSA isolates (77.9%) and 85 MSSA strains (22.1%) (**Figure 1**). Many infection

control practices changed during the COVID-19 pandemic, and there have been reports of

increased *S. aureus* incidence during the pandemic [25, 26] [27]. Therefore, we designated the

collection as being collected Pre-COVID (**Table 1;** 2017-2020, PreC = 211 strains, 54.9%) and

Post-COVID (2021-2023, PostC = 173 strains, 45.1%). There were 161 Blood samples making

up 41.9% of the sample set (**Table 1)**. Non-Blood strains (223 strains, 58.1%) were isolated from

infections of: bone (4 samples, 1%), eye (12 samples, 3.1%), fluid (29 samples, 7.6%), respiratory

(58 samples, 15.1%), urine (37 samples, 9.6%), and wound (83 samples, 21.6%) (**Table 1**).

The Pre-COVID and Post-COVID periods coincided with switch from a mupirocin-based

decolonization protocol to povidone-based protocol in January 2020. Our hospital had found

>40% prevalence of mupirocin resistance in 2019 in amongst MRSA isolates from surveillance

nasal swabs and a random sampling of MRSA blood cultures (46% of MRSA nares n=127; 43%

of MRSA blood cultures n=46). The first case of COVID-19 in the Tampa Bay area was March 1,

2020 with Post-COVID cases collected more than a year after change in decolonization protocol.

We determined the prevalence of mupirocin resistance amongst both MRSA and MSSA isolates

and whether specific genetic or phenotypic features correlated with resistance over time.

All strains were screened for oxacillin and mupirocin resistance using Etest strips. From this, we

determined that 278 of 384 strains (72.4%; **Figure 1A, Table 1**) were susceptible to mupirocin.

Of the 106 (27.6%) resistant strains, 34 (8.9%) had low-level resistance, whilst 72 (18.8%) had

high level resistance.

Mupirocin Resistance Prevalence was Stable: We observed significant correlation between

oxacillin resistance and mupirocin resistance (**Supplemental Table 2 and 3**). Most MupR strains

were MRSA (**Figure 1A**). From Pre-COVID samples, 2.1% of MSSA samples were also MupR,

whilst 36.6% of MRSA samples were MupR (p<0.001). 30.7% of MupS samples were oxacillin

susceptible, whilst 98.4% of MupR samples were oxacillin resistant. (p<0.001). Post-COVID strain

susceptibilities were similar. For example, 5.3% of MSSA samples were MupR and 31.9% of

MRSA samples were MupR (p<0.001).

When comparing levels of MupR between the Pre- and Post-COVID time points, there was no

significant difference between the two strain sets (p=0.460). In the Pre-COVID population, 71.1%

of strains were susceptible, whilst 28.9% displayed resistance (MupLR = 8.1%, MupHR = 20.9%).

In the Post-COVID strains, 74.0% were susceptible, whilst 26% displayed resistance (MupLR =

9.8%, MupHR = 16.2%).

Pre-COVID Non-Blood Strains Had Higher Levels of MupR: We found a significant difference

for strains isolated from blood versus non-blood. 21.7% of blood isolates displayed some level of

MupR, compared with 31.8% of non-blood samples (p=0.029; **Supplemental Table 4 and 5**).

This was driven by the Pre-COVID population, where 20.7% of blood samples displayed resistant

versus 34.1% of non-blood samples (p=0.037), whilst Post-COVID had no significant correlation

between isolate site and resistance (22.8% of blood samples were resistant versus 28.7% of non-

blood samples, p=0.375). When the non-blood strains were parsed more deeply, 46.2% of the

Pre-COVID respiratory isolates were MupR (p=0.016), which accounts for 29.5% of all Pre-COVID

MupR strains. Amongst our Post-COVID cohort, 26.3% respiratory strains were MupR, making

up 11.1% of MupR samples in the Post-COVID era (p=0.651).

Clonal Clusters and Sequence Types Associated with MupR: All strains from our collection

were subject to WGS and initially parsed into CC and then ST (**Figure 1B**). Strains that share six

of the seven conserved sequences used for multilocus sequence typing (MLST) are grouped into

clonal complexes (CC). MupR was most common in three sequence types (**Figure 2**). CC8 ST8

was the most abundant, with 173 strains (PreC = 96, 45.5% vs. PostC = 77, 44.5%), followed by

61 CC5 ST5 strains (PreC = 35, 16.6% vs. PostC = 26, 15%) and 36 CC5 ST3390 strains (PreC

= 24, 11.4% vs. PostC = 12, 7.5%). CC8 ST8 encompass USA300 and USA500, the most

common CA-MRSA in the US. There were 92 USA300 (ST8 t008) isolates (PreC = 51, 55.4% vs.

PostC = 41, 44.6%), and 31 USA500 (ST8 t064) strains (PreC = 19, 61.3% vs. PostC = 12,

38.7%). CC5 ST5 includes USA100, the CC/ST type classically associated with HA-MRSA) in the

US [28]. ST5, ST3390 and ST105 are closely related strains that are members of CC5.

When we explored MupR as a function of population structure, the three most common STs ST8,

ST5, and ST3390, made up the majority of the MupHR (88.9%) and MupLR (79.4%) strains

(**Figure 3**). MupHR strains comprised eight different STs, with the highest proportion being ST8

(43.1%) with 31 strains (PreC = 19, 43.2% vs. PostC = 12, 42.9%), then ST3390 (27.8%) with 20

strains (PreC = 13, 29.5% vs. PostC = 7, 25%) and 13 ST5 strains (PreC = 8,18.2% vs. PostC =

17.9%). ST3390 made up the highest group of MupLR overall with 10 strains (PreC = 6, 35.3%

vs. PostC = 4, 23.5%), followed by ST8, the majority of which are PostC (PreC = 2, 11.8% vs.

PostC = 7, 41.2%), and then finally, ST5 with most being PreC (PreC = 7, 41.2% vs. PostC = 1,

5.9%). Other MRSA-MupHR were ST105, ST1, and ST59. Of the three MSSA-MupR, two were

ST121 (one MupHR, one MupLR) and one ST1157 (MupHR).

High Level Mupirocin Resistant Strains Carry the *mupA* **Gene:** From our collection, 70 of 72

MupHR isolates contained the *mupA* gene, whilst no isolates were found to be *mupB* positive

(**Figure 3A**). MupHR strains were most commonly classified in the top three STs within two CCs

CC8 (ST8, n=31, 44.3%) and CC5 (ST5, n=13, 18.6% and ST3390, n=20, 28.6%). There was an

overall decrease of *mupA* carriage over time (20.4% PreC vs. 15.6% PostC) that was not

statistically significant (p=0.228). Within the pre-COVID strain collection (**Figure 3B**), we see a

higher number of *mupA*$^+$ isolates (43 strains) from a less diverse mix of the population (four STs

and two untypeable strains in total), whereas in the Post-COVID collection (**Figure 3B**) we see

27 strains distributed across six STs. The percentage of *mupA*$^+$ ST8 (PreC = 19, 44.2% vs. PostC

= 12, 44.4%) and ST5 isolates (PreC = 8, 18.6% vs. PostC = 5, 18.5%) remained largely stable,

however there was a slight decrease in the percentage of ST3390 isolates (PreC = 13, 30.2% vs.

PostC = 7, 25.9%, p-value = 0.264). The remaining *mupA*$^+$ isolates made up a relatively small

proportion of each population (PreC = 3, 6.9% vs. PostC = 3, 11.8%) and did not have any ST in

common. The rates of *mupA* prevalence across the Pre and Post pandemic era were not found

to be significantly different.

In general, the genotype and antibiotic susceptibility phenotype of our *mupA*$^+$ strains conformed,

e.g. *mupA*$^+$ strains were MupHR and *mupA*$^-$ strains were MupS (**Figure 3C**). However, there

were exceptions. Isolate TGH406, was *mupA+* but MupS, and no variation found in *mupA*

sequence to indicate it had become a pseudogene. We also detected a cluster of five *mupA*$^+$

isolates that were phenotypically MupLR (**Figure 3C**). We know from the literature that this is

likely due to *mupA* integration onto the chromosome [8]. There were six samples that were found

to have mutations within *mupA*, leading to varying degrees of protein truncations. Two truncations

were found that resulted in 957/1024 AA (n=2) and 997/1024 AA (n=1) variants that did not have

an impact on the function of *mupA* resistance measured phenotypically. Conversely, three isolates

had larger truncations, all of 137 AA, resulting in a shorter protein sequence of 887 AA. This

variant completely eliminated MupA function, resulting in a MupS phenotype.

Mutations in *IleS* Correlate with Low Level Mupirocin Resistance: Low level mupirocin

resistance is associated with a mutation (V588F) in the native isoleucyl-tRNA synthetase, *ileS*.

This mutation was identified in 27/34 MupLR isolates (PreC = 14, 6.6% vs. PostC = 13, 7.5%).

Three strains had the V588F mutation in *ileS* and also carried *mupA* and were MupHR as a result.

There was also a single MupLR isolate (TGH365) that had a novel *ileS* mutation (W436G) that to

our knowledge has never been reported. We also had four MupLR samples that did not contain

any mutations in the *ileS* sequence.

From a population perspective, there was a notable switch in strains bearing the V588F mutation

Pre-COVID and Post-COVID. Pre-COVID, there are no ST8 strains with MupLR that bear *ileS*

mutations; instead, the most prevalent groups are ST5 and ST3390, represented by six isolates,

each making up 42.9% of the population (**Figure 4A)**. In the Post-COVID population, ST8

represents 46.2% of the observed MupLR population, driven by the acquisition of the *ileS* (V588F)

mutation (**Figure 4B)**. There are no ST8 samples with this mutation prior to this shift in the Pre-

COVID population. Afterwards, Post-COVID ST8 made up 61.5% of all the strains with the

(V588F) mutation. The discrepancy between the percentage of Post-COVID MupLR and Post-

COVID (V588F) mutation ST8 strains is due to two ST8 strains in which the chromosomal

mutation effect is masked by the presence of *mupA,* resulting in a MupHR phenotype. Thus, we

observed CA-MRSA strains acquiring mutations in *ileS* and the *mupA* plasmid associated with

MupHR.

In addition, we identified another conserved novel MupLR *ileS* mutation (R112Q) that has never

been reported in the literature. This only appears within 7 of the 41 PostC USA300 strains,

indicating that this cluster may represents an outbreak within the ST8 clone (**Figure 4C**). Over

time the ST3390 population showed an increase in the presence of MupHR (+25.5% PostC), with

decrease in samples with *ileS* mutations (-16.7% PostC), suggesting evolution to a more highly

resistant population.

282   **Discussion**

Prior studies have shown that the rate of MupR varies by geographic region and clinical

characteristics of the patient populations sampled. The increasing use of decolonization protocols

designed to reduce the incidence of MRSA transmission creates selection pressure for

emergence of MupR. MupR has been reported in both MRSA and MSSA strains [9] [10], and

genome sequencing has implicated transmission of strains within healthcare settings [9] [10].

Amongst our strain collection, prevalence of MupR was higher in MRSA isolates than MSSA

isolates, as was also reported in strains from Stamford, CT [9]. For MRSA MupR strains, nasal

povidone-iodine protocols for decolonization should be effective [29], although recent clinical trials

have shown that nasal povidone iodine is inferior to nasal mupirocin [30]. Thus, optimal

management of MRSA colonization is not clear and will be influenced by local molecular

epidemiology of *S. aureus* and the rate of emergence of MupR.

The mechanism of MupHR is transmission of plasmid mediated *mupA* whilst MupLR is associated

with accumulation of chromosomal point mutations. Amongst the risk factors reported for MupR

are 1) prior exposure to mupirocin in the past year 2) *Pseudomonas aeruginosa* infection in the

year prior to admission and 3) cefepime use in the year prior to culture [31]. Ceasing mupirocin

exposure is reported to result in less MupR [8], although we did not see significant differences

between rates of MupR Pre-COVID and Post-COVID. The Pre-COVID strain collection was less

diverse than the Post-COVID collection so balancing selective pressures may be present. A

limitation of our retrospective study is that we did not have a method to determine mupirocin

exposure outside the inpatient setting. Other selective pressures that may be operative include

exposure to other antibiotics that favor maintenance of the resistance plasmid. Interestingly, many

of the isolates that carry *mupA* (58 out of 70), were also found to have a gene encoding for

resistance to aminoglycoside antibiotics, ANT(4')-Ia. Therefore, changing our decolonization

protocol to povidone iodine may have slowed emergence of MupR, but this trend may have been

masked by other factors affecting the population structure of MRSA in our hospital system.

The isolates examined were convenience samples, and our laboratory archives a random subset

of clinical isolates. We noted differences MupR in some of the subpopulations of isolates tested,

including respiratory samples. This phenotypic change in the Non-Blood category isolates may

be due to a population shift over time from being predominantly CC5 (HA-MRSA) to CC8 (CA-

MRSA), however, we cannot rule out that these differences were due to our small sample size. A

more in-depth characterization of samples from different clinical sites is needed to determine if

the changes we observed represent changes in population structure of clinical isolates, outbreaks

or selection of specific clones.

There were two isolates (Blood/Respiratory) from the same patient that had MupHR with no

identifiable *mupA* or *mupB* present. Both samples are ST105 SCC*mec* type II *Spa*-type t002, a

relative of the New York/Japan clone (USA100/ST5). European ST105-II-t002 has been linked to

skin and soft tissue infections, while the more recent South American ST105-II-t002 has been

heavily associated with blood stream infections [32, 33].

A major theme in *S. aureus* pathogenesis has been the emergence of waves of virulent clones

carrying genetic determinants conferring resistance to antibiotics [34]. In the 1990's strains were

either HA-MRSA or CA-MRSA, but gradually virulent CA-MRSA strains such as USA300 became

dominant [34]. We show that MupR in western Florida is associated with MRSA strains and

common US MRSA types such as ST8 (USA300 and USA500) and ST5 (USA100) are highly

represented within our clinical strains. Our collection also revealed the emergence of an unusual

ST3390 group with unexpected high prevalence of MupR (83%). Our ST3390 strains show

greater toxicity to neutrophils and are more virulent in mice than reference ST5 strain (Felton, *et*

al submitted). The genomes of ST3390 strains show evidence of horizontal exchange of genetic

traits that could promote maintenance in the population, increased virulence, and multi-drug

resistance (Felton, *et al* submitted).

Decolonization protocols for *S. aureus* utilizing antibiotics and antiseptics such as mupirocin or

povidone-iodine with chlorhexidine have shown to decrease surgical site infections and invasive

infections, but only result in temporary decolonization. These agents could promote the selection

of resistant organisms through selective pressure or broad disruption of the microbiome. The

incidence of mupirocin resistance and the molecular epidemiology of resistant strains in local

hospital populations is an important factor in selecting the appropriate agents used for

decolonization.

Acknowledgements:

This study was supported by grants AI124458 and AI157506 (both to L.N.S.) from the National

Institute of Allergy and Infectious Diseases and a USF Provost's CREATE award (LNS and KK).

We thank Emily Coughlin of the MCOM RISE program with advice about statistical analysis, the

USF Genomics facility, including John Adams and Min Zhang, for expert advice and assistance,

and Kartik Cherabuddi for review of the manuscript.

Data availability: Whole genome sequence data for all 384 isolates can be accessed at SRA

with accession numbers PRJNA785927 (Run1 80 isolates) and PRJNA1125331 (Run 2 and 3;

304 isolates). Sequenced strains are listed in Supplementary Table 1.

Conflicts of Interest: KK is a member of the Editorial Board of the Sanford Guide, has consulted

for Shionogi and Regeneron, and has received clinical research grants from Regeneron, Abbott,

Astra Zeneca, Pfizer, Romark, and the NIH. The other authors report no conflicts of interest.

References

1. von Eiff C, Becker K, Machka K, Stammer H, Peters G. Nasal carriage as a source of

Staphylococcus aureus bacteremia. Study Group. N Engl J Med **2001**; 344:11-6.

2. Huang SS, Singh R, McKinnell JA, et al. Decolonization to Reduce Postdischarge Infection

Risk among MRSA Carriers. N Engl J Med **2019**; 380:638-50.

3. Huang SS, Septimus E, Kleinman K, et al. Targeted versus universal decolonization to prevent

ICU infection. N Engl J Med **2013**; 368:2255-65.

4. Sharara SL, Maragakis LL, Cosgrove SE. Decolonization of Staphylococcus aureus. Infect Dis

Clin North Am **2021**; 35:107-33.

5. Prakash R, Garg A, Arya R, Kumawat RK. Chronicity of high and low level mupirocin resistance

in Staphylococcus aureus from 30 Indian hospitals. Sci Rep **2023**; 13:10171.

6. Tacconelli E. Methicillin-resistant Staphylococcus aureus: source control and surveillance

organization. Clin Microbiol Infect **2009**; 15 Suppl 7:31-8.

7. AMR NGHRUoGSo. Whole-genome sequencing as part of national and international

surveillance programmes for antimicrobial resistance: a roadmap. BMJ Glob Health **2020**; 5.

8. Patel JB, Gorwitz RJ, Jernigan JA. Mupirocin resistance. Clin Infect Dis **2009**; 49:935-41.

9. Labrecque S, Shah S, Fergus D, Parry MF. Mupirocin susceptibility of staphylococci 2022: Is it

time for a change in MRSA decolonization protocols? Am J Infect Control **2023**; 51:725-8.

10. Annavajhala MK, Kelly NE, Geng W, et al. Genomic and Epidemiological Features of Two

Dominant Methicillin-Susceptible Staphylococcus aureus Clones from a Neonatal Intensive Care

Unit Surveillance Effort. mSphere **2022**; 7:e0040922.

11. Hetem DJ, Bonten MJ. Clinical relevance of mupirocin resistance in Staphylococcus aureus.

J Hosp Infect **2013**; 85:249-56.

12. Antonio M, McFerran N, Pallen MJ. Mutations affecting the Rossman fold of isoleucyl-tRNA

synthetase are correlated with low-level mupirocin resistance in Staphylococcus aureus.

Antimicrob Agents Chemother **2002**; 46:438-42.

13. Seah C, Alexander DC, Louie L, et al. MupB, a new high-level mupirocin resistance

mechanism in Staphylococcus aureus. Antimicrob Agents Chemother **2012**; 56:1916-20.

14. Gaio D, Anantanawat K, To J, Liu M, Monahan L, Darling AE. Hackflex: low-cost, high-

throughput, Illumina Nextera Flex library construction. Microb Genom **2022**; 8.

15. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.

Bioinformatics **2014**; 30:2114-20.

16. Chen C, Khaleel SS, Huang H, Wu CH. Software for pre-processing Illumina next-generation

sequencing short read sequences. Source Code Biol Med **2014**; 9:8.

17. Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. Using SPAdes De Novo

Assembler. Curr Protoc Bioinformatics **2020**; 70:e102.

18. Mikheenko A, Prjibelski A, Saveliev V, Antipov D, Gurevich A. Versatile genome assembly

evaluation with QUAST-LG. Bioinformatics **2018**; 34:i142-i50.

19. Biguenet A, Bordy A, Atchon A, Hocquet D, Valot B. Introduction and benchmarking of

pyMLST: open-source software for assessing bacterial clonality using core genome MLST. Microb

Genom **2023**; 9.

20. Feldgarden M, Brover V, Gonzalez-Escalona N, et al. AMRFinderPlus and the Reference

Gene Catalog facilitate examination of the genomic links among antimicrobial resistance, stress

response, and virulence. Sci Rep **2021**; 11:12728.

21. Camacho C, Coulouris G, Avagyan V, et al. BLAST+: architecture and applications. BMC

Bioinformatics **2009**; 10:421.

22. Raghuram V, Alexander AM, Loo HQ, Petit RA, 3rd, Goldberg JB, Read TD. Species-Wide

Phylogenomics of the Staphylococcus aureus Agr Operon Revealed Convergent Evolution of

Frameshift Mutations. Microbiol Spectr **2022**; 10:e0133421.

23. Petit RA, 3rd, Read TD. Staphylococcus aureus viewed from the perspective of 40,000+

genomes. PeerJ **2018**; 6:e5261.

24. Sanchez-Herrero J, Sullivan M. spaTyper: Staphylococcal protein A (spa) characterization

pipeline.

25. Sanchez-Osuna M, Pedrosa M, Bierge P, et al. Genomic analysis of Staphylococcus aureus

isolates from bacteremia reveals genetic features associated with the COVID-19 pandemic.

iScience **2024**; 27:110402.

26. Baker MA, Sands KE, Huang SS, et al. The Impact of Coronavirus Disease 2019 (COVID-19)

on Healthcare-Associated Infections. Clin Infect Dis **2022**; 74:1748-54.

27. Gu J, Shen S, Xiong M, et al. ST7 Becomes One of the Most Common Staphylococcus aureus

Clones After the COVID-19 Epidemic in the City of Wuhan, China. Infect Drug Resist **2023**;

16:843-52.

28. Pardos de la Gandara M, Curry M, Berger J, et al. MRSA Causing Infections in Hospitals in

Greater Metropolitan New York: Major Shift in the Dominant Clonal Type between 1996 and 2014.

PLoS One **2016**; 11:e0156924.

29. Miller LG, McKinnell JA, Singh RD, et al. Decolonization in Nursing Homes to Prevent Infection

and Hospitalization. N Engl J Med **2023**; 389:1766-77.

30. Huang SS, Septimus EJ, Kleinman K, et al. Nasal Iodophor Antiseptic vs Nasal Mupirocin

Antibiotic in the Setting of Chlorhexidine Bathing to Prevent Infections in Adult ICUs: A

Randomized Clinical Trial. JAMA **2023**; 330:1337-47.

31. Caffrey AR, Quilliam BJ, LaPlante KL. Risk factors associated with mupirocin resistance in

meticillin-resistant Staphylococcus aureus. J Hosp Infect **2010**; 76:206-10.

32. Viana AS, Totola L, Figueiredo AMS. ST105 Lineage of MRSA: An Emerging Implication for

Bloodstream Infection in the American and European Continents. Antibiotics (Basel) **2024**; 13.

33. Viana AS, Botelho AMN, Feder A, et al. High frequency of increased triclosan MIC among

CC5 MRSA and risk of misclassification of the SCCmec into types. J Antimicrob Chemother **2022**;

77:3340-8.

34. Chambers HF, Fowler VG, Jr. Intertwining clonality and resistance: Staphylococcus aureus in

the antibiotic era. J Clin Invest **2024**; 134.

Table 1

Samples

	Strains	Ox R(MRSA)	Mup R
Pre-COVID	211	164	61
Post-COVID	173	135	45
Blood	161	100	35
Non-Blood	223	199	71
Total	384	299	106

435 **Strain Source**

	PreC			*PostC*			*Total*
	MupHR	MupLR	MupS	MupHR	MupLR	MupS	No (%)
Total #	44	17	150	28	17	128	384 (100)
Blood	12	5	65	9	9	61	161 (41.9)
Non-Blood	32	12	85	19	8	67	223 (58)
Bone	1	0	2	0	0	1	4 (1)
Eye	1	0	1	6	0	4	12 (3.1)
Fluid	3	1	12	1	0	12	29 (7.6)
Respiratory	13	5	21	2	3	14	58 (15.1)
Urine	5	1	12	4	2	13	37 (9.6)
Wound	9	5	37	6	3	23	83 (21.6)

S. aureus **strains used in this study.** Numbers collected Pre-COVID (2017-2020) and Post-

COVID (2021-2023) are indicated, as are the numbers of strains in each group with oxacillin

resistance and mupirocin resistance as defined in the methods. Sites of collection of the strains

are also indicated. All strains underwent whole genome sequencing (details in supplementary

table 2)

Figures:

Figure 1. Mupirocin Resistance and Sequence Typing of Tampa *S. aureus* isolates. A. 384

S.aureus clinical strains from 2017-2023 were characterized. Oxacillin sensitivity (MSSA, n=85),

mupirocin sensitivity (MupS, n=278) and the distribution of MRSA and MSSA strains amongst

mupirocin resistant (MupR n=106) strains is shown. B. Results of classification of strains into

sequence types (ST) are shown with numbers of Pre-COVID isolates (2017-2020) and Post-

COVID isolates (2021-2023) within each ST. Numbers of USA300 and USA500 within ST8 are

shown. ST5 includes USA100 strains. "UT" signifies strains that were untypeable using the

methods deployed herein. ST was determined by analysis of whole genome sequence of each

strain.

Figure 2. Mupirocin Resistance as a Function of Sequence Type. Shown are the sequence

types for high-level mupirocin resistance (MupHR) and low/intermediate-level mupirocin

resistance (MupLR) of the 106 MupR strains as determined by analysis of whole genome

sequence. Mupirocin susceptibility was determined by Etests. Two MupHR and four MupLR

strains were untypeable.

Figure 3. Distribution of *mupA* across Sequence Types and Correlation with Mupirocin Resistance. Summary of the distribution of *mupA*[+] isolates and ST across: (A) the entire 384 strain collection, (B) the Pre-COVID strains and the Post-COVID strains. Shown are the percentages and number of strains for each ST. C. The distribution of clinical strains with and without *mupA* and its phenotypic impact are shown. Most *mupA*[+] strains were MupHR as

determined by eTest. Strains are arrayed by isolation site, isolation period, the presence-absence

of *mupA*, and its phenotypic impact on mupirocin resistance/sensitivity. Strains with mutations in

ileS that are phenotypically MupLR are not included in this depiction.

Figure 4. Correlation of MupLR and *ileS* (V588F) as a Function of Sequence Type and

Isolation Site. Summary of the distribution of *ileS* (V588F) positive isolates **(A)** Pre-COVID and

(B) Post-COVID. **(C)** Shown is a recombination masked core SNP alignment maximum-likelihood

tree of Post-COVID USA300 isolates (ST8). The clade of 7 mupirocin resistant USA300 isolates

that possess the R112Q *IleS* mutation are designated in red. The reference USA300 strain

FPR_3757 used for SNP alignment is shown in blue.

Supplemental Table 2

Mupirocin Resistance Comparisons

Pre-COVID vs Post-COVID * Mup _Resistance

			Mup _Resistance			
			Susceptible	Intermediate	Resistant	Total
Group	Pre-COVID	Count	150	17	44	211
		%	71.1%	8.1%	20.9%	100.0%
		% of Total	39.1%	4.4%	11.5%	54.9%
	Post-COVID	Count	128	17	28	173
		%	74.0%	9.8%	16.2%	100.0%
		% of Total	33.3%	4.4%	7.3%	45.1%
Total		Count	278	34	72	384
		% of Total	72.4%	8.9%	18.8%	100.0%

Chi-Square Tests

	Value	df	Asymptotic Significance (2-sided)
Pearson Chi-Square	1.551[a]	2	0.460

483 **Supplemental Table 3**

Ox_Resistance vs Mup_Resistance

Group			Mup _Resistance Susceptible	Resistant	Total
Pre-COVID	MSSA	Count	46	1	47
		% of MSSA group	97.9%	2.1%	100.0%
		% within Mup _Resistance Group	30.7%	1.6%	22.3%
		% of Total	21.8%	0.5%	22.3%
	MRSA	Count	104	60	164
		% within MRSA group	63.4%	36.6%	100.0%
		% within Mup _Resistance Group	69.3%	98.4%	77.7%
		% of Total	49.3%	28.4%	77.7%
	Total	Count	150	61	211
		% within Ox_Resistance	71.1%	28.9%	100.0%
		% within Mup_Resistance Group	100.0%	100.0%	100.0%
		% of Total	71.1%	28.9%	100.0%
Post-COVID	MSSA	Count	36	2	38
		% within MSSA	94.7%	5.3%	100.0%
		% within Mup _Resistance Group	28.1%	4.4%	22.0%
		% of Total	20.8%	1.2%	22.0%
	MRSA	Count	92	43	135
		% within MRSA	68.1%	31.9%	100.0%
		% within Mup _Resistance Group	71.9%	95.6%	78.0%
		% of Total	53.2%	24.9%	78.0%
	Total	Count	128	45	173
		% within Ox_Resistance	74.0%	26.0%	100.0%

	% within Mup _Resistance Group	100.0%	100.0%	100.0%
	% of Total	74.0%	26.0%	100.0%

Chi-Square Tests

Date Code		Value	df	Asymptotic Significance (2-sided)
Pre-COVID	Pearson Chi-Square	21.105[a]	1	0.000
Post-COVID	Pearson Chi-Square	10.893[c]	1	0.001

487 **Supplemental Table 4**

Blood v. Non-blood * Mup _Resistance

| Date Code | | | Mup _Resistance | | |
			Susceptible	Resistant	Total
Pre-COVID	Blood	Count	65	17	82
		% within Blood vs. Non-blood	79.3%	20.7%	100.0%
		% of Total	30.8%	8.1%	38.9%
	Non-Blood	Count	85	44	129
		% within Blood v. Non-blood	65.9%	34.1%	100.0%
		% of Total	40.3%	20.9%	61.1%
	Total	Count	150	61	211
		% within Blood v. Non-blood	71.1%	28.9%	100.0%
		% of Total	71.1%	28.9%	100.0%
Post-COVID	Blood	Count	61	18	79
		% within Blood v. Non-blood	77.2%	22.8%	100.0%
		% of Total	35.3%	10.4%	45.7%
	Non-Blood	Count	67	27	94
		% within Blood v. Non-blood	71.3%	28.7%	100.0%
		% of Total	38.7%	15.6%	54.3%
	Total	Count	128	45	173
		% within Blood v. Non-blood	74.0%	26.0%	100.0%
		% of Total	74.0%	26.0%	100.0%

Chi-Square Tests

Date Code		Value	df	Asymptotic Significance (2-sided)
Pre-COVID	Pearson Chi-Square	4.365[a]	1	0.037
Post-COVID	Pearson Chi-Square	.787[c]	1	0.375

490   **Supplemental Table 5**

491   **Blood v. Non-blood * Mup _Resistance, Respiratory broken out**

Group			Mup_Resistance Susceptible	Resistant	Total
Pre-COVID	Blood	Count	65	17	82
		% of group	79.3%	20.7%	100.0%
		% of Total Mup_Resistant	43.3%	27.9%	38.9%
		% of Total	30.8%	8.1%	38.9%
	Respiratory	Count	21	18	39
		% of group	53.8%	46.2%	100.0%
		% of Total Mup_Resistant	14.0%	29.5%	18.5%
		% of Total	10.0%	8.5%	18.5%
	Non-Blood	Count	64	26	90
		% of group	71.1%	28.9%	100.0%
		% of Mup_Resistant	42.7%	42.6%	42.7%
		% of Total	30.3%	12.3%	42.7%
	Total	Count	150	61	211
		% of group	71.1%	28.9%	100.0%
		% of Mup_Resistant	100.0%	100.0%	100.0%
		% of Total	71.1%	28.9%	100.0%
Post-COVID	Blood	Count	61	18	79
		% within Source Code	77.2%	22.8%	100.0%
		% within Mup_Resistance	47.7%	40.0%	45.7%
		% of Total	35.3%	10.4%	45.7%
	Respiratory	Count	14	5	19
		% within Source Code	73.7%	26.3%	100.0%
		% within Mup_Resistance	10.9%	11.1%	11.0%

		% of Total	8.1%	2.9%	11.0%
Non-Blood		Count	53	22	75
		% within Source Code	70.7%	29.3%	100.0%
		% within Mup_Resistance	41.4%	48.9%	43.4%
		% of Total	30.6%	12.7%	43.4%
Total		Count	128	45	173
		% within Source Code	74.0%	26.0%	100.0%
		% within Mup_Resistance	100.0%	100.0%	100.0%
		% of Total	74.0%	26.0%	100.0%

Chi-Square Tests

Group		Value	df	Asymptotic Significance (2-sided)
Pre-COVID	Pearson Chi-Square	8.311[a]	2	0.016
Post-COVID	Pearson Chi-Square	.858[b]	2	0.651